HIV-1 Vpr drives a tissue residency-like phenotype during selective infection of resting memory T cells

Ann-Kathrin Reuschl[1], Maitreyi Shivkumar[1,2], Dejan Mesner[1], Laura J. Pallett[1], José Afonso Guerra-Assunção[1], Rajhmun Madansein[,3,4], Kaylesh J Dullabh[3], Alex Sigal[,5,6,7], John P. Thornhill[8,9], Carolina Herrera[9], Sarah Fidler[9,10], Mahdad Noursadeghi[1], Mala K. Maini[1] and Clare Jolly[1*].

[1] Division of Infection and Immunity, University College London, London, UK
[2] Present address: Leicester School of Pharmacy, De Montfort University, Leicester, UK
[3] Department of Cardiothoracic Surgery, University of KwaZulu-Natal, Durban, South Africa
[4] Centre for the AIDS Programme of Research in South Africa, Durban, South Africa
[5] Africa Health Research Institute, Durban, South Africa
[6] School of Laboratory Medicine and Medical Sciences, University of KwaZulu-Natal, Durban, South Africa
[7] Max Planck Institute for Infection Biology, Berlin, Germany
[8] Peter Medawar Building for Pathogen Research, Nuffield Department of Medicine, University of Oxford, Oxford, UK.
[9] Department of Infectious Diseases, Faculty of Medicine, Imperial College, London, UK.
[10] Imperial College NIHR Biomedical Research Centre, London, UK.

Human immunodeficiency virus type 1 (HIV-1) replicates in CD4+ T cells leading to profound

T cell loss, immunological dysfunction and AIDS. Determining how HIV-1 shapes the

immunological niche in which it resides to create a permissive environment is central to

informing efforts to limit pathogenesis, disturb viral reservoirs and achieve a cure. A key

roadblock in understanding HIV-T cell interactions is the requirement to activate CD4+ T cells

***in vitro* in order to make them permissive to infection. This dramatically alters T cell biology,**

obscuring native virus-host interactions. Here we show that HIV-1 cell-to-cell spread permits

efficient and productive infection of resting CD4+ T cells without the need for prior activation.

Infection is preferential for resting memory T cells, is observed with both CXCR4-tropic virus

and CCR5-tropic transmitter-founder viruses and results in virus production and onward

spreading infection. Strikingly, we find that HIV-1 infection of resting memory CD4+ T cells

primes for induction of a tissue-resident memory (T$_{RM}$)-like phenotype evidenced by

upregulation of T$_{RM}$ markers CD69/CXCR6 alongside co-expression of CD49a, PD-1, CD101

as well as transcription factor Blimp-1. Furthermore, we reveal that HIV-1 initiates a

transcriptional program that overlaps with the core T$_{RM}$ transcriptional signature. This

reprograming depends on the HIV-1 accessory protein Vpr. We propose that HIV-1 infection

drives a CD4+ T$_{RM}$-phenotype potentially sequestering infected cells within tissues to support

viral replication and persistence.

Introduction

Resting primary CD4+ T cells cannot be efficiently infected by cell-free HIV-1 virions *in vitro* and

require robust mitogenic stimulation to support viral replication[1–3]. This has led to the notion that T

cell activation is necessary for HIV-1 replication and that resting T cells are not permissive for HIV-

1 infection. However, mitogenic T cell activation *in vitro* results in wide-spread phenotypic and

functional reprogramming which dominates changes to gene and protein expression[4–6], concealing

and altering authentic virus-host interactions. This presents a significant challenge for understanding

the cellular response to HIV-1 infection and the consequences of the virus-host interaction for HIV-

1 replication and persistence. While it is clear HIV-1 efficiently infects and replicates in activated T

cells, the outcomes of the virus-host interaction with resting T cells has been reported to be cell

death[7] or latency[8]. However, previous data demonstrating that HIV-1 cell-to-cell spread is highly-

efficient and drives widespread changes in protein phosphorylation status in both infected and target

cells[9–13] suggested that cell-to-cell spread may overcome the barrier to productive infection of resting

T cells. Here, we comprehensively show for the first time that HIV-1 exploits cell-to-cell spread to

efficiently infect resting memory CD4+ T cells and have used this to uncover a hithertho unknown

consequence of HIV-1 infection for T cell reprogramming driven by the accessory protein Vpr.

Results

To test whether cell-to-cell spread allows for productive infection of resting T cells, HIV-1 infected

Jurkat T cells (Fig. 1a) or primary CD4+ T cells (Fig. 1b) were co-cultured with uninfected resting

CD4+ T cells (Extended data Fig. 1a-e). We confirmed that CD4+ T cells isolated from peripheral

blood display a resting phenotype by staining for Ki67, CD69, HLA-DR and MCM2. (Extended data

Fig. 1c-e). Infection of resting target cells in the absence of mitogenic or cytokine activation was

measured. Direct co-culture of infected and uninfected cells resulted in significant levels of HIV-1

infection of resting CD4+ target T cells (Gag+) measured by intracellular flow cytometry staining (Fig. 1a, b and Extended data Fig. 1f). By contrast, resting CD4+T cells were not infected (<1%) when cell-cell contact was prevented by separating the two cell populations by a transwell (Fig. 1a,b), a condition that only allows for cell-free infection. As expected, mitogenic activation of target T cells made them more permissive to cell-free HIV-1 but, as previously shown, infection was still substantially boosted by direct co-culture allowing for additional cell-to-cell spread (Extended data Fig.1f-h) [9,11,12]. Infection of resting CD4+ target T cells was preferentially detected in CD45RA-ve resting memory T cell populations rather than CD45RA+ naïve T cells, which are both abundant in peripheral blood (Fig. 1c-e, Extended Fig. 1i). Co-staining for CD62L confirmed that the infected CD45RA+ve cells were mainly naïve rather than T_{EMRA} (Extended data Fig. 1j,k)[14]. The preferential infection of CD45RA-ve memory T cell populations rather than the CD45RA+ve naïve population, is in agreement with HIV-1 being predominantly detected in memory CD4+ T cells *in vivo* [15–17]. This was not due to competition between naïve and memory cells because the same effect was observed when CD45RA+ve and CD45RA-ve resting CD4+ target T cells were separated prior to cell-to-cell infection (Fig. 1f and Extended data Fig. 2a). Cell-to-cell infection of resting memory T cells was observed with CXCR4-tropic strain NL4.3, and CCR5-tropic viruses NL4.3 BaL and two transmitter-founder (T/F) primary isolates (CH040 and CH077) (Fig. 1g-i) demonstrating that selective permissivity was not unique to a particular virus or receptor tropism. Preventing viral entry with fusion inhibitor (T20), or blocking reverse transcription (Efavirenz), inhibited the appearance of Gag+ve or GFP+ve cells (Fig. 1j-l and Extended data Fig. 2b,c) demonstrating that this signal reflects productive infection and not simply virus capture[9,11–13]. Consistent with infection, we also observed downregulation of CD4 expression on target cells (Fig. 1m and Extended data Fig. 2d) that was most pronounced in the resting memory T cell population. Importantly, flow-sorting and culturing HIV-1 infected resting target T cells showed that these cells support viral replication by producing new virus and transmitting infection to new target cells (Fig. 1n-p and Extended data Fig. 2e), further demonstrating productive infection.

We confirmed that HIV-1+ target T cells infected by cell-to-cell spread maintained their resting phenotype and did not upregulate Ki67 or MCM2, two markers of cell-cycle progression (Extended

data Fig. 3a,b), showing these cells were not simply being activated and driven into cell cycle, either

by infection or bystander effects during co-culture. Intriguingly, expression of CD69 on HIV-1 infected

resting memory target T cells was significantly increased compared to mock-treated target T cells

(Fig. 2a,b). This was not due to preferential infection of a minor pre-existing population of CD69+ve

CD4+ T cells in blood (Extended data Fig. 1c) because flow cytometry sorting of CD69-ve CD4+ T

cells (prior to co-culture with HIV-1 infected donor T cells) showed *de novo* upregulation of CD69 on

the newly-infected resting memory target cells (Extended data Fig.3c,d). While CD69 is classically

thought of as a marker of early T cell activation, expression can occur independently of cell cycle

progression and T cell activation[18–20]. Consistent with this we did not detect T cell activation

concomitant with CD69 upregulation in HIV-1 infected resting CD4+ memory T cells since CD69+ve

cells remained HLA-DR-negative (Extended data Fig.3e).

Functionally, CD69 is crucial for T cell retention in tissues by interfering with S1P receptor

mediated egress and has been identified as a hallmark of tissue-resident memory (T_{RM}) T cells[21].

Recently it has been demonstrated that although T_{RM} cells are largely absent in peripheral blood

(Extended data Fig. 3f), precursor cells poised to adopt a T_{RM} phenotype are present in

circulation[22,23]. Interestingly, while HIV-1 infection alone was associated with upregulation of CD69,

exposure of HIV-1 infected resting memory T cells to the homeostatic T cell cytokine IL7 further

boosted CD69 upregulation 4-fold, as compared to HIV-1 alone or IL7 alone (Fig. 2c and Extended

data Fig. 3j). IL7 secreted by stromal cells is required for long-term maintenance of CD4+ T_{RM} cells[24–

26]. Although IL7 can enhance HIV-1 infection of T cells[27] (Extended data Fig. 3g), infection of resting

memory T cells mediated by cell-to-cell spread does not require IL7 (Fig.1) and IL7 increased CD69

expression on infected cells even when added 48h post-infection (Extended data Fig. 3j). HIV-1

induced CD69 upregulation was completely abrogated by suppressing infection with the fusion

inhibitor T20 (Fig. 2c) or by treating cells with Ruxolitinib that blocks IL7-mediated JAK signalling

(Fig. 2d) demonstrating that the enhanced CD69 induction requires both infection and cytokine

signalling. Similar infection-driven enhancement of CD69 expression was observed in response to

γ_c-chain cytokine IL15 (Fig. 2e,f) but not IL12 or TGF-β (Extended data Fig.3h,i). Consistent with the

hypothesis that HIV-1 infection induces a CD4+ T_{RM}-like phenotypic signature, resting memory T

cells also upregulated the T_{RM} marker CXCR6[21] during HIV-1 infection, (Fig. 2g,h and Extended data

Fig. 3k,n), as well as increasing the population of CD69+ve cells co-expressing CD49a, PD-1 and

CD101 that are also associated with the core T_{RM} signature[21] (Fig. 2i, j and Extended data Fig. 3l).

As expected for T_{RM}s, we saw no upregulation of CX3CR1 expression (Extended data Fig. 3o) and

no transcriptional upregulation of *S1PR1* or *KLF2* mRNA (Extended data Fig. 5i). Critically, induction

of T_{RM}-associated markers did not occur in uninfected Gag-ve bystander cells (Fig. 2k and Extended

data Fig.3m,n). By contrast to CD8+ T_{RM} cells, CD103 was barely detectable and not upregulated

(Extended data Fig.3p,q), consistent with the observation of limited CD103 expression on CD4+ T

cells[21]. Induction of CD69 expression was also concomitant with upregulation of the T_{RM}-associated

transcription factor Blimp-1[28–30] (Fig. 2l,m). Similar upregulation of T_{RM}-associated markers were also

observed when unstimulated CD4+ T cells from tonsil (Fig. 2n,o) or mediastinal lymph nodes (Fig.

2p,q) were infected with HIV-1 via cell-to-cell spread and exposed to IL7 (Fig. 2o,q), demonstrating

that induction of TRM-like phenotype also occurs in tissue-derived T cells following HIV-1 infection.

Taken together, these data suggest that HIV-1 infection of resting memory CD4+ T cells

reprogrammes cells by upregulating expression of T_{RM}-associated marker proteins, and thus induces

a phenotype characteristic of tissue residency.

HIV-1 expresses four accessory proteins, Vif, Vpu, Vpr and Nef which directly and indirectly

manipulate host cell factors to facilitate efficient viral replication *in vivo* and drive pathogenesis[31]. Co-

culture of resting target T cells with donor T cells infected with HIV-1 accessory protein mutants

showed that deletion of Vpr (HIV-1 ΔVpr) resulted in a complete abrogation of the T_{RM}-like phenotype

as evidenced by a lack of CD69 upregulation, and no detection of the CD69+/CXCR6+/CD49a+

triple-positive T_{RM}-like memory population (Fig. 3a,b,d, Extended data Fig.4 and 5a-e). By contrast,

deletion of Vpu or Nef did not affect HIV-1 induction of T_{RM}-markers on infected resting T cells (Fig.

3a,b, Extended data Fig. 4 and Fig. 5a-e). HIV-1 ΔVif could not be tested because Vif is required to

antagonise APOBEC3-mediated viral restriction and allow infection[32]. HIV-1 Vpr is not required for

infection of T cells *in vitro*[33,34] and concordantly, loss of T_{RM}-marker protein induction was not due to

lack of infection of target cells by HIV-1 ΔVpr virus, nor reduced Gag expression (Fig. 3c and

Extended data Fig. 5f-h). Critically, Vpr was also required for induction of CD69 expression observed

at the mRNA level, as well as induction of *CXCR6* and Blimp1 (*PRDM1*) mRNA (Fig. 3e). As

expected, there was no upregulation of *S1PR1* or *KLF2* mRNA by either HIV-1 WT or ΔVpr virus

(Extended data Fig.5i), consistent with their suppression under conditions of T_{RM} induction[21]. HIV-1

dependent upregulation of CD69 surface expression was inhibited by Ruxolitinib treatment (Fig. 2d)

in a Vpr-dependent manner (Extended data Fig. 5j). Furthermore, HIV-1 infection downregulated the

IL7 receptor alpha-subunit (CD127) from the cell surface (Extended data Fig 5k) and transcriptionally

(fold change=0.693, adjusted p-value=5.89E-08, Supplementary table 1), indicative of activation of

this pathway[35] by HIV-1. This was accompanied by a significant increase in the intracellular levels of

phosphorylated transcription factor STAT5 (Extended data Fig. 5j and k). These data are consistent

with Vpr manipulating cellular signalling pathways to drive induction of a T_{RM}-like phenotype in

response to IL7. Vpr-dependent T_{RM}-like induction was also accompanied by spontaneous

production of IFNγ by infected CD4+ memory T cells (Fig. 3f), in line with the increased capacity of

T_{RM} to produce this cytokine[21,36]. Notably, Vpr-mediated induction of a T_{RM}-like phenotype was not

inhibited by the integrase inhibitor Raltegravir (Fig. 3g, h) that potently suppressed HIV-1 integration

into resting memory CD4+ T cells infected by cell-to-cell spread (Fig. 3i), demonstrating that the T_{RM}-

like phenotype does not require integration. Of note, the presence of integrated provirus in resting

CD4+ T cells (less than or equal to 1 provirus per cell Fig. 3i) further supports the observation that

cell-to-cell spread results in resting CD4+ T cell infection (Fig. 1). Collectively, these data identify

Vpr as the viral determinant required for upregulation of T_{RM}-associated proteins on HIV-1 infected

resting CD4+ T cells.

Vpr is a multifunctional protein that is packaged into viral particles and is present during the

early stages of infection where it plays an important, but as yet poorly-defined role in HIV-1

pathogenesis. Amongst the best-defined functions of Vpr are its ability to a) bind the Cul4A-DDB1

(DCAF1) complex (leading to an interaction with the ubiquitinylation and proteasomal machinery); b)

induce G2/M cell-cycle arrest and; c) drive apoptosis in infected cells[37–41]. We abrogated these

functions individually by introducing the mutations Q65R, S79A or R80A into Vpr in the context of

virus and confirmed that each Vpr mutant is packaged into virions (Fig. 3j). Co-culture of resting

memory target T cells with HIV-1+ T cells infected with different Vpr mutants revealed that the Vpr

mutants S79A or R80A (inhibits cell cycle arrest) behaved similarly to WT virus and did not induce CD69 and CXCR6 upregulation (Fig. 3k-m). By contrast, Q65R (that is most closely associated with loss of DCAF1 binding) was unable to induce a T_{RM}-like phenotype following infection of target cells, behaving like ΔVpr virus in these experiments (Fig. 3k-m).

Next we performed transcriptional profiling by RNA-Seq analysis of flow-sorted cells infected with HIV-1 WT or ΔVpr virus by cell-to-cell spread. Figure 4 shows that HIV-1 infection induced widespread changes to gene expression in resting memory T cells when compared to uninfected cells (232 differentially expressed genes (DEG), fold change >1.2, adjusted p-value<0.01) (Fig. 4). Hierarchial clustering and principal component analysis revealed that the gene expression patterns in response to HIV-1 WT infection were clearly distinct from those induced following infection with ΔVpr virus (Fig. 4 a-c, e,f) demonstrating that *Vpr*-deletion suppresses the global transcriptional response to HIV-1 infection. Specifically, infection with ΔVpr virus resulted in only 13 genes showing statistically significantly changes compared to uninfected cells, by contrast to 232 genes for HIV-1 WT virus (Supplementary table 1 and 3, Fig. 4e). In fact, much of the transcriptional response to HIV-1 infection was regulated by Vpr, as evidenced by changes in DEG in the presence and absence of Vpr (Fig. 4f). The requirement for Vpr in driving many of the changes to DEG was also observed in response to IL7 (Fig. 4d, Supplementary table 2 and 4, and Extended data Fig. 6a). Consistent with Vpr manipulating the T cell response to HIV-1 infection (Supplementary table 5 and 6), gene-set enrichment analysis (GSEA) (Fig. 4g) and Ingenuity Pathway Analysis (IPA) (Fig. 4h and i) revealed enrichment of numerous cellular signalling pathways following HIV-1 infection that appeared Vpr-dependent, most notably pathways associated with cytokine and inflammatory responses as well as immune signalling (Fig. 4g, Extended data Fig. 6 and Supplementary table 7 and 8). This was further evidenced by Upstream Regulator analysis that showed significant enrichment for genes associated with cytokine signalling and transcriptional regulators that were again largely Vpr-dependent (Fig. 4j and k, and Extended data Fig. 6). Taken together, these data reveal that HIV-1 induces dramatic reprogramming during infection of resting memory CD4+ T cells that is driven largely by Vpr.

Tissue residency of T cells has been associated with a 31-gene core transcriptional

signature[21]. We took advantage of this dataset and our RNAseq analysis to determine whether this

core T_{RM} signature was enriched in our transcriptome of HIV-1 infected resting memory T cells.

Hierarchical clustering of our data, compared to the core signature of T_{RM} cells (CD69+ve T cells

isolated from human lung and spleen)[21] from Kumar et al, showed that HIV-1 infected memory T

cells exposed to IL7 grouped distinctly and clustered with CD69+ve T_{RM} cells (Fig. 5a,

Supplementary table 9) distinct from non-T_{RM} T cells (CD69-ve T cells isolated from tissue and blood

from Kumar et al). We further corroborated this finding by calculating a T_{RM} enrichment score based

on the published core transcriptional signature, which showed that HIV-1 infected resting memory T

cells (both +/-IL7) harbour a higher T_{RM} signature score, significantly different from non-T_{RM} T cells,

approximating more closely to that of *bona fide* T_{RM} cells (Fig. 5b). Critically, this was Vpr-dependent,

with mock and ΔVpr infected cells showing an enrichment score that was not statistically different to

non-T_{RM}. Taken together, these data demonstrate that HIV-1 induces both a phenotypic and

transcriptional T_{RM}-like signature in resting T cells via the accessory protein Vpr.

Discussion

Our discovery that resting CD4+ T cells can be productively infected by cell-to-cell spread, allowing

for viral integration, replication and dissemination, transforms our ability to determine how T cells

respond to and suppport HIV-1 replication without confounding activation-induced changes.

Recently, it has been reported that cell-to-cell spread can also facilitate latent infection of resting T

cells without productive infection[8]. Together, these two studies highlight the distinct advantages of

co-culture models (that do not require mitogenic, experimental stimulation of T cells to drive infection)

to study native HIV-1-host cell interactions and the cellular response to infection.

Here we have employed our co-culture model to show that HIV-1 infection of resting memory

T cells induces a T_{RM}-like phenotype evidenced by upregulation and co-expression of T_{RM}-associated

markers on infected cells and induction of a core T_{RM} transcriptional signature. HIV-1 establishes

cellular and tissue reservoirs (both active and latent) that ultimately prevent cure with antiretroviral

therapy. Importantly, T_{RM} cells are long-lived and are thought to be largely confined to tissue[42] providing an alternate model for a tissue-associated reservoir driven by the virus itself. Our results suggest that HIV-1 persistence and the establishment of tissue reservoirs may be driven, in part, through direct viral induction of a T_{RM}-like phenotype via transcriptional reprogramming. Recently, TRM in cervical tissue were found to be preferentially infected by HIV-1 and can harbour an HIV-1 reservoir *in vivo*[43]. The relative contribution of pre-existing versus HIV-1 induced T_{RM} cells to viral reservoirs and their relative abundance in different anatomical compartments *in vivo* remains to be quantified, but we expect T_{RM} cells harbouring virus to be important contributors to viral persistence. In light of these findings it is possible that HIV-1 infected cells circulating in peripheral blood may in fact represent cells that have failed to become part of the tissue reservoir, leading to an underestimation of the true viral burden. Having shown that HIV-1 infection of resting T cells by cell-to-cell spread results in productive infection, we hypothesise that induction of a T_{RM}-like phenotype in infected cells may also play additional roles in establishing and maintaining viral reservoirs by sequestering infected cells in tissue sites where susceptible target T cells are in abundance, thus supporting localised viral replication. Indeed we have shown that infected resting memory T cells support spreading infection to disseminate virus. Given the importance of T_{RM} cells as a population that are increasingly recognised to be critical in providing localised immunity and immunosurveillance[20,30], future work should focus on understanding the contribution of HIV-1 induced T_{RM}-like cells in pathogenesis and persistence.

It is now emerging that committed T_{RM} precursors, imprinted with the capacity to become mature T_{RM} pre-exist in blood and that when exposed to the appropriate cues in tissues or *ex vivo* can become tissue-resident[22,23]. Thus the ontogeny, derivation and maintenance of T_{RM} cells appears more complex that initially appreciated. Our discovery that HIV-1 induces a T_{RM}-like phenotype in CD4+ T cells provides an opportunity to gain new understanding of mechanisms behind CD4+ T_{RM} induction and maintainance.

We found that HIV-1 infection of resting memory T cells was associated with striking transcriptional reprograming that was licenced by Vpr, thus identifting a novel function for this enigmatic HIV accessory protein. Notably, HIV-1 induction of a T_{RM}-like phenotype via Vpr was

accompanied by induction of a T_{RM} transcriptional signature that aligned closely with a published

core T_{RM} signature[21]. Vpr-deletion not only abolished induction of this T_{RM} signature, but also many

HIV-1 induced changes to gene expression following infection of resting T cells. This is in keeping

with wide-spread proteome remodelling by Vpr in activated T cells[44], but suggests that these

changes may be driven in part by a hitherto unappreciated role for Vpr in modulating the host cell

gene expression profile. Whether this reflects wide-spread epigenetic changes mediated by Vpr or

manipulation of key upstream regulators remains to be determined. Our data show HIV-1 Vpr

modulates T cell responsiveness to external stimuli by manipulation of immune signalling pathways,

including innate and inflammatory responses. This is particularly intriguing and suggests HIV-1

manipulates immune signalling pathways to benefit the virus, in this case by priming resting memory

T cells for T_{RM}-like induction. Vpr-mediated induction of T_{RM}-like phenotype was dependent on

residue Q65 and Vpr is reported to drive wide-spread remodelling of the cellular proteome via its

recruitment of DCAF1 through Q65[44] . Whether this requirement for Q65 in induction of a T_{RM}-like

phenotype is DCAF1-dependent or not remains unclear because DCAF1 knockdown in primary T

cells made cells hyper-responsive to HIV-1 induced cell death (Extended data Fig. 7). Thus we

cannot at present formally exclude other functions of Vpr Q65 in the process of T_{RM}-induction.

Notably, a rare case of laboratory-derived infection with Vpr-defective HIV-1 was

characterised by markedly delayed seroconversion, suppressed viremia and normal CD4+ T cell

counts[45], consistent with reduced pathogenesis and failure to establish and maintain a significantly

large tissue reservoir. We envisage therapeutic targeting of Vpr to manipulate persistence and

pathogenesis. In order to achieve an HIV-1 cure it is essential to understand the nature and

establishment of HIV-1 reservoirs and how to manipulate them. By demonstrating that HIV-1

infection drives a T_{RM}-like phenotype during infection of resting memory T cells we have taken a

significant step towards this to help accelerate the quest for an HIV-1 cure.

References

1. Swiggard, W. J. *et al.* Human Immunodeficiency Virus Type 1 Can Establish Latent Infection

in Resting CD4 + T Cells in the Absence of Activating Stimuli. *J. Virol.* **79**, 14179–14188

(2005).

2. Stevenson, M., Stanwick, T. L., Dempsey, M. P. & Lamonica, C. A. HIV-1 replication is

controlled at the level of T cell activation and proviral integration. *EMBO J.* **9**, 1551–60

(1990).

3. Zack, J. A. *et al.* HIV-1 entry into quiescent primary lymphocytes: molecular analysis reveals

a labile, latent viral structure. *Cell* **61**, 213–22 (1990).

4. Wolf, T. *et al.* Dynamics in protein translation sustaining T cell preparedness. *Nat. Immunol.*

21, 927–937 (2020).

5. Szabo, P. A. *et al.* Single-cell transcriptomics of human T cells reveals tissue and activation

signatures in health and disease. *Nat. Commun.* **10**, 4706 (2019).

6. Howden, A. J. M. *et al.* Quantitative analysis of T cell proteomes and environmental sensors

during T cell differentiation. *Nat. Immunol.* **20**, 1542–1554 (2019).

7. Doitsh, G. *et al.* Abortive HIV infection mediates CD4 T cell depletion and inflammation in

human lymphoid tissue. *Cell* **143**, 789–801 (2010).

8. Agosto, L. M., Herring, M. B., Mothes, W. & Henderson, A. J. HIV-1-Infected CD4+ T Cells

Facilitate Latent Infection of Resting CD4+ T Cells through Cell-Cell Contact. *Cell Rep.* **24**,

2088–2100 (2018).

9. Jolly, C., Kashefi, K., Hollinshead, M. & Sattentau, Q. J. HIV-1 cell to cell transfer across an

Env-induced, actin-dependent synapse. *J. Exp. Med.* **199**, 283–93 (2004).

10. Sattentau, Q. Avoiding the void: cell-to-cell spread of human viruses. *Nat. Rev. Microbiol.* **6**,

815–826 (2008).

11. Sourisseau, M., Sol-Foulon, N., Porrot, F., Blanchet, F. & Schwartz, O. Inefficient Human

Immunodeficiency Virus Replication in Mobile Lymphocytes. *J. Virol.* **81**, 1000–1012 (2007).

12. Hübner, W. *et al.* Quantitative 3D video microscopy of HIV transfer across T cell virological

synapses. *Science.* **323**, 1743–7 (2009).

13. Len, A. C. L., Starling, S., Shivkumar, M. & Jolly, C. HIV-1 Activates T Cell Signaling

Independently of Antigen to Drive Viral Spread. *Cell Rep.* **18**, 1062–1074 (2017).

14. Sallusto, F., Lenig, D., Förster, R., Lipp, M. & Lanzavecchia, A. Pillars Article : Two Subsets

of Memory T Lymphocytes with Distinct Homing Potentials and Effector Functions . Nature .

Nature **401**, 708–712 (1999).

15. Shan, L. *et al.* Transcriptional Reprogramming during Effector-to-Memory Transition

Renders CD4 + T Cells Permissive for Latent HIV-1 Infection. *Immunity* **47**, 766-775.e3

(2017).

16. Brenchley, J. M. *et al.* T-cell subsets that harbor human immunodeficiency virus (HIV) in

vivo: implications for HIV pathogenesis. *J. Virol.* **78**, 1160–8 (2004).

17. Chomont, N. *et al.* HIV reservoir size and persistence are driven by T cell survival and

homeostatic proliferation. *Nat. Med.* **15**, 893–900 (2009).

18. Corneau, A. *et al.* Comprehensive Mass Cytometry Analysis of Cell Cycle, Activation, and

Coinhibitory Receptors Expression in CD4 T Cells from Healthy and HIV-Infected

Individuals. *Cytom. Part B - Clin. Cytom.* **92**, 21–32 (2017).

19. Lea, N. C. *et al.* Commitment Point during G 0 → G 1 That Controls Entry into the Cell Cycle

Commitment Point during G 0 3 G 1 That Controls Entry into the Cell Cycle. *Mol. Cell. Biol.*

23, 2351–2361 (2003).

20. Szabo, P. A., Miron, M. & Farber, D. L. Location, location, location: Tissue resident memory

T cells in mice and humans. *Sci. Immunol.* **4**, eaas9673 (2019).

21. Kumar, B. V *et al.* Human Tissue-Resident Memory T Cells Are Defined by Core

Transcriptional and Functional Signatures in Lymphoid and Mucosal Sites. *Cell Rep.* **20**,

2921–2934 (2017).

22.   Kok, L. *et al.* A committed tissue-resident memory T cell precursor within the circulating

CD8+ effector T cell pool. *J. Exp. Med.* **217**, (2020).

23.   Fonseca, R. *et al.* Developmental plasticity allows outside-in immune responses by resident

memory T cells. *Nat. Immunol.* **21**, 412–421 (2020).

24.   Yeon, S. M. *et al.* IL-7 plays a critical role for the homeostasis of allergen-specific memory

CD4 T cells in the lung and airways. *Sci. Rep.* **7**, 1–9 (2017).

25.   Adachi, T. *et al.* Hair follicle-derived IL-7 and IL-15 mediate skin-resident memory T cell

homeostasis and lymphoma. *Nat. Med.* **21**, 1272–1279 (2015).

26.   Amezcua Vesely, M. C. *et al.* Effector TH17 Cells Give Rise to Long-Lived TRM Cells that

Are Essential for an Immediate Response against Bacterial Infection. *Cell* **178**, 1176-

1188.e15 (2019).

27.   Coiras, M. *et al.* IL-7 Induces SAMHD1 Phosphorylation in CD4+ T Lymphocytes, Improving

Early Steps of HIV-1 Life Cycle. *Cell Rep.* **14**, 2100–2107 (2016).

28.   Hombrink, P. *et al.* Programs for the persistence, vigilance and control of human CD8 +

lung-resident memory T cells. *Nat. Immunol.* **17**, 1467–1478 (2016).

29.   Mackay, L. K. *et al.* Hobit and Blimp1 instruct a universal transcriptional program of tissue

residency in lymphocytes. **352**, 459–464 (2016).

30.   Pallett, L. J. *et al.* IL-2(high) tissue-resident T cells in the human liver: Sentinels for

hepatotropic infection. *J. Exp. Med.* **214**, 1567–1580 (2017).

31.   Malim, M. H. & Emerman, M. HIV-1 Accessory Proteins-Ensuring Viral Survival in a Hostile

Environment. *Cell Host Microbe* **3**, 388–398 (2008).

32.   Sheehy, A. M., Gaddis, N. C. & Malim, M. H. The antiretroviral enzyme APOBEC3G is

degraded by the proteasome in response to HIV-1 Vif. *Nat. Med.* **9**, 1404–1407 (2003).

33.   Balliet, J. W. *et al.* Distinct Effects in Primary Macrophages and Lymphocytes of the Human

Immunodeficiency Virus Type 1 Accessory Genes vpr, vpu, and nef: Mutational Analysis of

a Primary HIV-1 Isolate. *Virology* **200**, 623–631 (1994).

34.   Rogel, M. E., Wu, L. I. & Emerman, M. The human immunodeficiency virus type 1 vpr gene

prevents cell proliferation during chronic infection. *J. Virol.* **69**, 882–8 (1995).

35.   Park, J. H. *et al.* Suppression of IL7Rα transcription by IL-7 and other prosurvival cytokines:

A novel mechanism for maximizing IL-7-dependent T cell survival. *Immunity* **21**, 289–302

(2004).

36.   Kumar, B. V. *et al.* Functional heterogeneity of human tissue-resident memory T cells based

on dye efflux capacities. *JCI insight* **3**, (2018).

37.   Jowett, J. B. *et al.* The human immunodeficiency virus type 1 vpr gene arrests infected T

cells in the G2 + M phase of the cell cycle. *J. Virol.* **69**, 6304–13 (1995).

38.   Wu, Y. *et al.* The DDB1-DCAF1-Vpr-UNG2 crystal structure reveals how HIV-1 Vpr steers

human UNG2 toward destruction. *Nat. Struct. Mol. Biol.* **23**, 933–939 (2016).

39.   Laguette, N. *et al.* Premature activation of the slx4 complex by vpr promotes g2/m arrest

and escape from innate immune sensing. *Cell* **156**, 134–145 (2014).

40.   Schröfelbauer, B., Hakata, Y. & Landau, N. R. HIV-1 Vpr function is mediated by interaction

with the damage-specific DNA-binding protein DDB1. *Proc. Natl. Acad. Sci. U. S. A.* **104**,

4130–4135 (2007).

41.   Wen, X., Duus, K. M., Friedrich, T. D. & De Noronha, C. M. C. The HIV1 protein Vpr acts to

promote G2 cell cycle arrest by engaging a DDB1 and cullin4A-containing ubiquitin ligase

complex using VprBP/DCAF1 as an adaptor. *J. Biol. Chem.* **282**, 27046–27051 (2007).

42.   Kumar, B. V., Connors, T. J. & Farber, D. L. Human T Cell Development, Localization, and

Function throughout Life. *Immunity* **48**, 202–213 (2018).

43.   Cantero-Pérez, J. *et al.* Resident memory T cells are a cellular reservoir for HIV in the

cervical mucosa. *Nat. Commun.* **10**, (2019).

44.   Greenwood, E. J. D. *et al.* Promiscuous Targeting of Cellular Proteins by Vpr Drives

Systems-Level Proteomic Remodeling in HIV-1 Infection. *Cell Rep.* **27**, 1579-1596.e7

(2019).

45.   Ali, A. *et al.* Highly attenuated infection with a VPR-deleted molecular clone of human

immunodeficiency virus-1. *J. Infect. Dis.* **218**, 1447–1452 (2018).

Methods

Cells

Peripheral blood mononuclear cells (PBMC) were isolated from buffy coats from healthy donors (UK NHS Blood and Transplant Service) by density centrifugation using FicollPaque Plus (GE Life Sciences) and cryopreserved in 10% DMSO (Sigma-Aldrich) in 90% FBS (LabTech). Resting CD4+ T cells were isolated from total PBMCs by negative selection using the MojoSort Human CD4+ T Cell Isolation kit (Biolegend) according to the manufacturer's instructions. CD45RA+ve naïve and CD45RA-ve memory populations were further separated after CD4+ T cell isolation with CD45RA MicroBeads (Biolegend). For activated CD4+ T cells, PBMCs were treated with 5µg/ml PHA (Sigma) and 10IU/ml IL2 (Centre For AIDS Reagents, National Institute of Biological Standards and Control, UK [CFAR]) in RPMI1640 with 20% FBS for 72h prior to CD4+ T cell isolation. Once purified, CD4+ T cells were cultured in RPMI supplemented with 20% FBS and 10IU/ml IL2. Jurkat T cell lines (Clone E6-1; ATCC TIB-152) were cultured in RPMI with 10% FBS and 100U/ml penicillin/streptomycin. HEK 293T/17 cells (ATCC, CRL-11268) were cultured in DMEM with 10% FBS and 100U/ml penicillin/streptomycin. Tonsil tissue was obtained from an individual with primary HIV infection who underwent routine tonsillectomy (2 months after commencement of ART). As previously described[46],the tonsillar tissue from elective tonsillectomy was dissected and mechanically digested, prior to cryopreservation of the cellular suspension. This was collected under the Imperial College Infectious Diseases Biobank (REC: 15/SC/0089). Lymph nodes were obtained from the field of surgery of participants undergoing surgery for diagnostic purposes and/or complications of inflammatory lung disease. Informed consent was obtained from each participant,

and the study protocol approved by the University of KwaZulu-Natal Institutional Review Board (approval BE024/09).

Plasmids and virus production

The HIV-1 clone pNL4.3 was obtained from the CFAR, NIBSC (cat# 2006). HIV-1 NL4.3 ΔNef and pNL4.3 ΔVpr were provided by R. Sloan (University of Edinburgh, UK)[47]. NL4.3 ΔVpu was provided by S. Neil (King's College London, UK)[48]. NLENG1-IRES was provided by D. Levy (NYU, USA)[49]. NL4.3 bearing the CCR5-tropic BaL *Env* was provided by G. Towers (UCL, UK)[50]. CCR5 tropic transmitter/founder virus plasmids CH044 and CH077 were provided by G. Towers (UCL, UK) and were originally obtained through the NIH AIDS Reagent Program [NIHARP], Division of AIDS, NIAID, NIH: pCH040.c/2625 (cat# 11740) and pCH077.t/2627 (cat# 11742) from Dr. John Kappes and Dr. Christina Ochsenbauer. NL4.3 Vpr Q65R, NL4.3 Vpr S79A, NL4.3 Vpr R80A were generated by site-directed mutagenesis (Promega) using the following primers:

NL4.3 VprQ65R fw:GTGGAAGCCATAATAAGAATTCTGCGACAACTGCTGTTTATCCATTTCAG

NL4.3 VprQ65R rv:CTGAAATGGATAAACAGCAGTTGTCGCAGAATTCTTATTATGGCTTCCAC

NL4.3 Vpr S79A fw: GAATTGGGTGTCGACATGCCAGAATAGGCGTTACTC

NL4.3 Vpr S79A rv:GAGTAACGCCTATTCTGGCATGTCGACACCCAATTC

NL4.3 Vpr R80A fw: GGTGTCGACATAGCGCAATAGGCGTTACTCG

NL4.3 Vpr R80A rv: CGAGTAACGCCTATTGCGCTATGTCGACACC.

All virus stocks were produced by plasmid transfection of HEK 293T cells with Fugene 6 (Promega). Supernatants were harvested at 48h and 72h, filtered, DNase treated, purified and concentrated by ultracentrifugation through a 25% sucrose cushion and resuspended in RPMI1640 with 10% FBS. Viral titres were determined by measuring reverse transcriptase activity by SG-PERT assay[51].

HIV-1 infection and cell-to-cell spread

For cell-to-cell spread experiments, activated primary CD4+ T cells (donor cells) were infected with 800mU reverse transcriptase per 10^6 cells of HIV-1 by spinoculation at 1200x*g* for 2h at room temperature and incubated in RPMI 20% FBS supplemented with 10IU/ml IL2 for 72h. HIV-1+ donor

CD4+ T cells were washed with medium, counted and cultured with primary CD4+ target T cells at

a 1:1 ratio in RPMI 20% FBS supplemented with 10IU/ml IL2 for up to 72h before analysis by flow

cytometry or FACS sorting. Uninfected target CD4+ T cells were pre-stained with 1-2nM CellTrace

FarRed dye (Invitrogen) prior to co-culture. For cell-to-cell spread into tonsil-derived lymphocytes,

total tonsil lymphocytes were cultured at a 4:1 ratio with HIV-1 infected or uninfected eFluor450-

labelled Jurkat T cells. For FACS sorting experiments, donor cells were pre-labeled with cell dye

eFluor450 (ThermoFisher). For transwell experiments, HIV-1 infected donor T cells were separated

from target T cells by a 0.4µm transwell insert (Corning). Experiments to quantify cell-to-cell versus

cell-free infection in the presence and absence of a transwell were performed in equivalent volumes

(600µl). For some experiments, FACS sorted infected resting CD4+ target T cells were returned into

cultured for up to 4 days. Infection levels were measured by intracellular Gag staining and flow

cytometry, and virus release into cell culture supernatant determined by SG-PERT[51]. At day 1 or day

4 post FACS sorting, resting CD4+ T cells were washed extensively and co-cultured at a 1:1 ratio

with uninfected eFluor450-labelled Jurkat T cells for 72h, when Jurkat T cell infection was measured

by Gag-staining. Where indicated, cultures were incubated in the presence of 20ng/ml IL7 (Miltenyi

Biotec), 20ng/ml IL15 (Peprotech), 20ng/ml IL12 (Peprotech) or 50ng/ml TGFβ (Peprotech). The

following inhibitors were added 30min before co-culture at the following concentrations: T20 (25-

50ng/ml, CFAR), Efavirenz (1µM, CFAR), Raltegravir (5µM, CFAR) and Ruxolitinib (50nM, Sigma).

For RNAi knockdown of DCAF1, primary CD4+T cells were activated for 4 days with 1µg/ml plate-

bound αCD3 antibody (cloneOKT3, Biolegend) in the presence of 2 µg/mL soluble αCD28

antibody(clone CD28.2, Biolegend). RNAi knockdown of DCAF1 was performed as described

before[52] using ON-TARGET plusHuman DCAF1 siRNA - SMARTpool (Dharmacon) and non-

targeting siRNA (Dharmacon) was used as a control. 2×10^6cells were electroporated with 200 pmol

siRNA using a NeonTransfection System (Thermo Fisher Scientific; three pulses, 10 ms, 1,600 V).

After 48h, DCAF1 knockdowns were confirmed by western blotting and cells used in cell-to-cell

spread experiments as described above.

Flow cytometry and FACS

For flow cytometry analysis, cells were washed in PBS and stained with fixable Zombie UV Live/Dead dye, Aqua Live/Dead dye or NIR Live/Dead dye (Biolegend) for 5 min at 37°C. Excess stain was quenched with FBS-complemented RPMI. When tonsil and lymph node lymphocytes were used, Live/Dead staining was quenched using human AB serum (Sigma) in RPMI. Cell surface staining was performed in PBS, complemented with 20% Super Bright Staining Buffer (ThermoFisher) when appropriate, at 4°C for 30min. Unbound antibody was washed off thoroughly and cells were fixed with 4% FA or PFA before intracellular staining. For intracellular detection of IFNγ in infected target CD4+ T cells after 72h of cell-to-cell spread, cells were treated throughout the co-culture with IL7 and Brefeldin A (Biolegend) treated for 6h before surface staining and fixation. Permeabilisation for intracellular staining was performed with IC perm buffer or FoxP3 Buffer set (Biolegend) according to the manufacturer's instructions. For detection of intracellular P-STAT5, cells were resuspended in ice cold True-Phos Perm buffer (Biolegend) and permeabilised for 48h at -20°C. Intracellular P-STAT5 staining was then performed in PBS with wash steps performed at 1800 rpm for 6 min at 4°C.The following antibody clones and fluorochromes were used: CD3 (UCHT1, Biolegend; Bv510, Bv711, FITC), CD8 (SK1, Biolegend; Bv605, PE), CD4 (SK3, Biolegend; APC/Fire750); CD45RA (HI100, Biolegend; Bv421, PE-Dazzle); CD45RO (UCHL1, Biolegend; PerCp-Cy5.5), CD69 (FN50, Biolegend; APC/Fire750, PE-Dazzle); CXCR6/CD186 (K041ES, Biolegend; PE-Dazzle); MCM2 (ab4461, ABCAM; was detected with a secondary anti-rabbit AlexaFluor488-tagged antibody); HLA-DR (L234, Biolegend; PerCp-Cy5.5); CD49a (TS2/7, Biolegend; PE-Cy7); PD-1 (EH12.2H7, Biolegend; PE-Cy7); Ki67 (Ki-67, Biolegend; Bv711, PE); Blimp-1 (6D3, BD Pharmingen; PE); CD101 (BB27, Biolegend; PE-Cy7); CX3CR1 (2A9-1, Biolegend; PE-Dazzle); CD103 (Ber-ACT8, Biolegend; Bv711); CD127 (AO19D5, Biolegend; PE-Cy7), IFNγ (B27, Biolegend; PE), Phospho-STAT5 (Clone 47/Stat5 (pY694), BD; PE) and HIV-1 Gag core antigen (FH190-1-1, Beckman Coulter; PE, FITC). All samples were acquired on either an BD Fortessa X20 or LSR II using BD FACSDiva software and analyzed using FlowJo v10 (Tree Star). FACS sorting was performed with a BD FACSAria III or BD FACSAria IIu Cell Sorter. Cells were either lysed immediately in RLT lysis buffer (Qiagen) with 1% β-mercaptoethanol (Sigma) and stored at -80C for later RNA extraction or resuspended in RPMI supplemented with 20% FBS and 10IU/ml IL2 and used immediately.

Western blotting

Virus-containing supernatants (normalised for equal loading by measuring RT activity) or fifteen micrograms of total CD4+ T cell protein lysate were separated by SDS-PAGE, transferred onto nitrocellulose and blocked in PBS with 0.05% Tween 20 (v/v) and 5% skimmed milk (w/v). Blots were probed with rabbit antisera raised against HIV-1 Gag p24 (cat# 0432 donated by Dr G. Reid and obtained from the CFAR), Vpr anti-serum (cat# 3951, NIH ARP), α−alpha-Tubulin (T6199, Sigma-Aldrich) and α -DCAF1 antibody (11612-1-AO, Proteintech), followed by goat anti-rabbit or goat anti-mouse IRdye 800CW or 680RD infrared secondary antibody (Abcam) and imaged using an Odyssey Infrared Imager (LI-COR Biosciences) and analysed with Image Studio Lite software.

Quantification of HIV-1 integration

To quantify integration of HIV-1 in resting T cells, nested Alu-gag quantitative PCR was performed as previously described[53]. Briefly, DNA was isolated from FACS sorted infected resting CD4+ memory T cells after 72h of cell-to-cell spread using the Qiagen Blood Mini Kit.

Integrated DNA was pre-amplified using 100nM Alu fw primer, 600nM HIV-1 Gag rv primer, 0.2mM dNTP, 1U Phusion Hot Start Flex (Promega), and 45ng DNA in 50µl reactions. Cycling conditions were: 94˚C for 30s, followed by 40 cycles of 94˚C for 10s, 55˚C for 30s, and 70˚C for 2.5min. For quantitation of HIV-1 integration, a second round real-time quantitative PCR was performed using the pre-amplified DNA. These samples were run alongside a standard curve of known dilutions of CEM cells containing integrated HIV-1 DNA. Reactions contained 0.25µM of RU5 fw and rv primers, and 0.2µM probes, 1x Qiagen Multiplex Mastermix, and 10µl pre-amplified DNA. Cyclin conditions were: 95˚C for 15min, followed by 50 cycles of 94˚C for 60s and 60˚C for 60s. 2LTR circles were measured by quantitative PCR[54]. Reactions contained 150ng DNA, 10µ 2LTR fw and rv primers, 10µM probe and 1x TaqMan Gene Expression Master Mix (ThermoFisher). Cycling conditions were: 95˚C for 15min, followed by 50 cycles of 95˚C for 15s and 60˚C for 90s. Reactions were performed using 7500 Real-Time PCR System (Applied Biosystems). The following primers and probes were used:

Alu fw: GCCTCCCAAAGTGCTGGGATTACAG

HIV-1 Gag rv: GTTCCTGCTATGTCACTTCC

RU5 fw: TTAAGCCTCAATAAAGCTTGCC

RU5 rv: GTTCGGGCGCCACTGCTAGA

RU5-WT probe: FAM-CCAGAGTCACACAACAGACGGGCACA-TAMRA

RU5-degenerate 1 probe: FAM-CCAGAGTCACATAACAGACGGGCACA-TAMRA

RU5-degenerate 2 probe: FAM-CCAGAGTCACACAACAGATGGGCACA-TAMRA

2LTR fw: AACTAGAGATCCCTCAGACCCTTTT

2LTR rv: CTTGTCTTCGTTGGGAGTGAAT

2LTR probe: FAM-CTAGAGATTTTCCACACTGAC-TAMRA

qRT-PCR

RNA was extracted from FACS sorted target memory CD4+ T cells with RNeasy Micro Kit (Qiagen)

according to the manufacturer's instructions. cDNA was synthesised using SuperScript IV with

random hexamer primers (Invitrogen) and qRT-PCR was performed using Fast SYBR Green Master

Mix and 7500 Real-Time PCR System (Applied Biosystems). Gene expression was determined

using the $2^{-\Delta\Delta Ct}$ method and normalised to GAPDH expression. The following primers were used:

GAPDH fw: ACATCGCTCAGACACCATG, rv: TGTAGTTGAGGTCAATGAAGGG;

CXCR6 fw: GACTATGGGTTCAGCAGTTTCA, rv:GGCTCTGCAACTTATGGTAGAAG;

PRDM1 fw: ATGCGGATATGACTCTGTGGA, rv: CTGAACCGAAGTACCGCCATC;

CD69 fw: ATTGTCCAGGCCAATACACATT, rv: CCTCTCTACCTGCGTATCGTTTT;

S1PR1 fw: TCTGCTGGCAAATTCAAGCGA, rv: GTTGTCCCCTTCGTCTTTCTG;

KLF2 fw: CTACACCAAGAGTTCGCATCTG; rv: CCGTGTGCTTTCGGTAGTG.

Whole transcriptome profiling by RNA-Sequencing

RNA was extracted from FACS sorted target memory CD4+ T cells with RNeasy Micro Kit (Qiagen)

according to the manufacturer's instructions. For preparation of RNA-Sequencing libraries, RNA

concentration was measured using the Qubit RNA High Sensitivity kit (Life Technologies) and quality

checked on the 4200 Tapestation using either the High Sensitivity or standard RNA ScreenTape

assay (Agilent Technologies), depending on the measured RNA concentrations. PolyA-tailed mRNA

was separated for sequencing during library preparation. Libraries were prepared using KAPA's mRNA HyperPrep kit (Roche Diagnostics) according to the manufacturer's instructions using an input of up to 200ng and a fragmentation incubation time of 8 minutes at 94°C. Samples were sequenced on Illumina's NextSeq500 (Illumina Cambridge) using a high output 75 cycle paired-end run. 24 libraries were multiplexed in the same run. Libraries were pooled in equimolar quantities, calculated from concentrations measured using the Qubit dsDNA High Sensitivity kit (Life Technologies) and fragment analysis using the D1000 High Sensitivity assay on the 4200 Tapestation (Agilent Technologies).

RNA sequencing data was quality assessed using FASTQC (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) before and after low-quality and adapter trimming using Trimmomatic[55]. Filtered reads were then pseudo-mapped using Kallisto[56] to the transcriptome available in Ensembl v.101 (http://aug2020.archive.ensembl.org/index.html). Per-transcript counts were imported and aggregated per gene using the TXimport R package[57]. The DESeq2 package[58] was used for data normalisation, outlier detection and differential gene expression analysis between biological groups. The DESeq2 results were ranked based on the log2 transformation of the adjusted p-values, to provide a pre-ranked list for Gene Set Enrichment Analysis (GSEA)[59] as described in the GSEA documentation. Pathway enrichment and upstream regulator analysis was performed using Gene Set Enrichment Analysis (GSEA)[59] and Ingenuity Pathway Analysis (IPA,) respectively. Heatmaps were generated using ClustVis (https://biit.cs.ut.ee/clustvis/) [60]

Transcriptomic comparison with published human T_{RM} cells

TPM data from previously published transcriptomes of human T_{RM} cells (GSE94964)[21] were summed on gene level with Ensembl gene ID, gene name, and gene biotype using tximport and BioMart[57,61]. TPM values <0.001 were adjusted to 0.001 as a lower limit of detection. These data were aligned to the transcriptomic data from the present study using gene symbol in an integrated Log_2 transformed data matrix and subjected to batch correction by study using Combat[62]. Expression of selected genes

previously identified to be up and downregulated in T_{RM}[21] were used to cluster the samples in both studies using 1-Spearman rank correlation with average linkage in ClustVis[60]. A transcriptional signature score for T_{RM} was derived from the difference between the sum of up and down-regulated genes in T_{RM} in the previously published signature. This score was used to evaluate the relative similarity of each transcriptome data set in the present project to T_{RM} and non-T_{RM} data.

Statistical analysis

Statistical analysis was performed using GraphPad Prism. Normally distributed data was analysed for statistical significance by two-tailed t- tests (when comparing two groups) or one-way ANOVA with Bonferroni or Dunnett's post-test (when comparing more than two groups). Data show the mean +/- the S.E.M with significance shown on the figures. Where appropriate, the median+IQR is shown and Kruskall-Wallis test was used to compare groups. Significance levels were defined as *, P < 0.05; **, P < 0.01 and ***, P < 0.001.

Methods References

46.   Thornhill, J. P. *et al.* CD32 expressing doublets in HIV-infected gut-associated lymphoid tissue are associated with a T follicular helper cell phenotype. *Mucosal Immunol.* (2019). doi:10.1038/s41385-019-0180-2

47.   Sloan, R. D. *et al.* Transcription of Preintegrated HIV-1 cDNA Modulates Cell Surface Expression of Major Histocompatibility Complex Class I via Nef. *J. Virol.* **85**, 2828–2836 (2011).

48.   Neil, S. J. D., Eastman, S. W., Jouvenet, N. & Bieniasz, P. D. HIV-1 Vpu promotes release and prevents endocytosis of nascent retrovirus particles from the plasma membrane. *PLoS Pathog.* **2**, 354–367 (2006).

49.   Trinité, B. *et al.* Suppression of Foxo1 activity and down-modulation of CD62L (L-selectin) in HIV-1 infected resting CD4 T cells. *PLoS One* **9**, (2014).

593   50.   Rasaiyaah, J. *et al.* HIV-1 evades innate immune recognition through specific cofactor

594   recruitment. *Nature* **503**, 402–405 (2013).

595   51.   Pizzato, M. *et al.* A one-step SYBR Green I-based product-enhanced reverse transcriptase

596   assay for the quantitation of retroviruses in cell culture supernatants. *J. Virol. Methods* **156**,

597   1–7 (2009).

598   52.   Mesner, D., Hotter, D., Kirchhoff, F. & Jolly, C. Loss of Nef-mediated CD3 down-regulation

599   in the HIV-1 lineage increases viral infectivity and spread. *Proc. Natl. Acad. Sci. U. S. A.*

600   **117**, 7382–7391 (2020).

601   53.   Liszewski, M. K., Yu, J. J. & O'Doherty, U. Detecting HIV-1 integration by repetitive-

602   sampling Alu-gag PCR. *Methods* **47**, 254–260 (2009).

603   54.   Apolonia, L. *et al.* Stable gene transfer to muscle using non-integrating lentiviral vectors.

604   *Mol. Ther.* **15**, 1947–1954 (2007).

605   55.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: A flexible trimmer for Illumina sequence

606   data. *Bioinformatics* **30**, 2114–2120 (2014).

607   56.   Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq

608   quantification. *Nat. Biotechnol.* **34**, 525–527 (2016).

609   57.   Soneson, C., Love, M. I. & Robinson, M. D. Differential analyses for RNA-seq: Transcript-

610   level estimates improve gene-level inferences [version 2; referees: 2 approved].

611   *F1000Research* **4**, 1–23 (2016).

612   58.   Anders, S. & Huber, W. Differential expression analysis for sequence count data. *Genome*

613   *Biol.* **11**, (2010).

614   59.   Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach for

615   interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 15545–

616   15550 (2005).

617   60.   Metsalu, T. & Vilo, J. ClustVis: A web tool for visualizing clustering of multivariate data using

618   Principal Component Analysis and heatmap. *Nucleic Acids Res.* **43**, W566–W570 (2015).

619   61.   Smedley, D. *et al.* The BioMart community portal: An innovative alternative to large,

620   centralized data repositories. *Nucleic Acids Res.* **43**, W589–W598 (2015).

621   62.   Leek, J. T., Johnson, W. E., Parker, H. S., Jaffe, A. E. & Storey, J. D. The SVA package for

622   removing batch effects and other unwanted variation in high-throughput experiments.

623   *Bioinformatics* **28**, 882–883 (2012).

Acknowledgements

This work was funded by a Wellcome Trust Investigator award (108079/Z/15/Z) to C.J. We are

grateful to members of the Jolly lab, as well as Greg Towers, Rebecca Sumner, Lorena Zulianai-

Alvarez, Lucy Thorne, Laura McCoy and Richard Milne for helpful discussions and critical reading of

the manuscript. We thank Jamie Evans (UCL) for assistance with sorting CD69-negative T cells and

Parisa Amjadi (Imperial College London Chelsea and Westminster Hospital) for sorting target T cells

from co-culture experiments and the Pathogen Genomics Unit (PGU) at UCL for RNAseq analysis.

We acknowledge the NIBSC Centre for AIDS Reagents, the NIH AIDS Reagent Program and

Imperial College NIHR Biomedical Research Centre, London, UK.

Author contributions

A.K.R., and C.J. conceived the project. A.K.R., and C.J. designed the experiments. A.K.R., M.S. and

D.M performed the experiments. A.K.R., C.J., L.P., M.K.M., M.S., D.M., A.G-A and M.N. analysed

the data. L.P. and M.K.M provided reagents. J.P.T., C.H., S.F., R.M, K.J.D., and A.S. provided

lymphoid tissue samples. A.G-A and M.N. performed the core T_{RM} transcriptional mapping analysis.

A.K.R., and C.J. prepared the manuscript. All authors provided critical review of the manuscript.

Competing interest

The authors declare they have no competing interests.

Figure legends:

Fig. 1 HIV-1 exploits cell-to-cell spread to preferentially infect resting memory CD4+ T cells.
HIV-1 NL4.3 infected (**a**) Jurkat or (**b**) mitogenically-activated primary CD4+ donor T cells co-cultured
with resting primary CD4+ target T cells separated by a 0.4μm transwell (cell-free) or direct co-culture
(cell-cell). Target cell infection was measured by intracellular staining for HIV-1 Gag protein.
Representative flow cytometry plots are shown. Bar graphs show mean of independent experiments
(n=4). (**c,d**) Cell-to-cell spread from activated primary donor CD4+ T cells to resting primary target
CD4+ T cells preferentially infects CD45RA- memory CD4+ T cells. A representative flow cytometry
plot and quantification is shown (n=4). (**e**) Quantification of infection performed as in (**c**) (n=11). (**f**)
HIV-1 infection of target CD4+ T cells as part of the total resting CD4+ T cell population (total)
compared to pre-isolated naive and memory CD4+ target T cells (isolated) (n=9). (**g**) Representative
flow cytometry plots of cell-to-cell infection of resting CD4+ T cells with CCR5-tropic HIV-1 NL4.3
BaL and transmitter founder viruses HIV-1 CH040 and CH077 as performed in (**c**). (**h**) Quantification
of infection of CXCR4 (X4) and CCR5 (R5)-tropic viruses (n=4) and (**i**) transmitter/founder viruses
HIV-1 CH040 and CH077 (n=7). (**j**) Cell-to-cell infection of resting CD4+ T cells is reduced by the
HIV-1 fusion-inhibitor T20 (n=6) and (**k**) the reverse transciptase inhibitor Efavirenz (n=6) measured
by intracellular Gag staining (MFI) or (**l**) HIV-1 LTR-driven GFP-reporter gene expression (n=4). (**m**)
HIV-1 infection downregulates CD4 expression. Shown are the percentage of CD4+ cells of the total
CD3+ target cell population (n=6). Resting CD4+ memory T cells were isolated after 72h of cell-to-
cell spread by FACS sorting and cultured for 4 days. HIV-1 infection was measured by intracellular
Gag staining (**n**) and virus release measured by culture supernatant RT activity (**o**) (n=5-7). T cells
from (**n**) were recovered at day 1 or 4 post isolation and cultured with uninfected eFluor450+ target
Jurkat T cells. Infection of Jurkat T cells was measured after 72h (**p**) (n=3). Data are the mean±SEM.
Paired two-tailed *t*-test or one-way ANOVA with Bonferroni post-test were used. For (**m**),
median+IQR is shown and Friedman test with Dunn's post-test was used. For (**o**), unpaired one-
tailed *t*-test was used. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; n.s., not significant.

Fig. 2 HIV-1 infection induces a T$_{RM}$-like phenotype in resting memory CD4+ T cells. (**a**) CD69 expression on resting memory CD4+ target T cells following co-culture with HIV-1 infected primary donor T cells or uninfected donor T cells (mock) (n=17). (**b**) Representative flow cytometry plots from (**a**). (**c**) CD69 expression on infected resting memory CD4+ T cells ± IL7 and T20 (n=7). (**d**) CD69 expression on infected resting memory CD4+ T cells ± IL7 and Ruxilitinib (n=8). (**e**) CD69 expression on infected resting memory CD4+ T cells in response to IL7 and IL15 (n=11). (**f**) CD69 expression on infected Gag+ resting memory CD4+ T cells and uninfected Gag- bystander cells in response to IL7 and IL15 (n=11). (**g**) CXCR6 surface expression from (**f**) (n=11). (**h**) Representative flow cytometry plots of CD69 and CXCR6 co-expression. (**i**) Co-expression of CD69 with CXCR6, CD49A or PD-1 on infected resting memory CD4+ T cells (n=5-7). (**j**) As for (**i**) in the presence of IL7 (n=4-7). (**k**) As for (**i**) comparing infected Gag+ memory CD4+ T cells and uninfected Gag- bystander cells. (**l**) and (**m**) Blimp-1 expression in CD69+ infected resting memory CD4+ T cells and infected CD69- cells in response to IL7 (n=8). (**n**) Total lymphocytes from cellularised tonsils co-cultured with HIV-1 infected Jurkat T cells. Infection of resting CD4+ T cells (live CD3+/CD8-/Ki67- lymphocytes) shown as CD45RO vs Gag. (**o**) CD69 and CXCR6 co-expression on infected Gag+ and uninfected Gag- tonsil resting memory CD4+ T cells ± IL7. (**p**) Total lymphocytes from mediastinal lymph nodes co-cultured with HIV-1 infected activated autologous LN-derived lymphocytes. Infection of resting CD4+ T cells (live CD3+/CD8-/Ki67- lymphocytes) shown as CD45RA vs Gag. (**q**) CD69 and CXCR6 co-expression on infected Gag+ and uninfected Gag- lymph node resting memory CD4+ T cells ±IL7. Data are the mean±SEM. Paired two-tailed *t*-test or one-way ANOVA with Bonferroni or Dunnett's post-test were used. *p<0.05 ; **, p<0.01; ***, p<0.001; n.s., not significant.

Fig. 3 Vpr drives HIV-1-induced T$_{RM}$-induction in resting memory CD4+ T cells. Resting memory CD4+ T cells were co-cultured with HIV-1 infected primary CD4+ T cells infected with HIV-1 WT or mutant viruses, or uninfected donor cells (mock). (**a**) CD69 upregulation in response to IL7 compared to mock (n=9). (**b**) CD69 expression on HIV-1 infected Gag+ resting memory CD4+ T cells compared to uninfected Gag- bystander cells (n=9). (**c**) Quantification of cell-to-cell spread of HIV-1 WT and ΔVpr to resting memory CD4+ T cells (n=9). (**d**) CD69/CXCR6/CD49a co-expression on resting

memory CD4+ T cells infected with HIV-1 WT or ΔVpr (n=9). (**e**) *CD69, CXCR6* and *PRDM1* (Blimp1) mRNA levels from FACS sorted infected resting memory CD4+ T cells. Fold change relative to unifected (mock) is shown (n=5). (**f**) IFNg expression by HIV-1 infected resting memory CD4+ T cells at 72h in response to IL7 (n=3). (**g**) CD69 and (**h**) CD69/CXCR6 co-expression in response to IL7 in the presence of integrase inhibitor Raltegravir (n=6). (**i**) Quantification of integrated provirus and 2LTR circles in FACS sorted target CD4+ memory T cells after 72h of cell-to-cell spread in the presence or absence of Raltegravir. (**j**) Western blot showing Vpr packaging into HIV-1 WT and Vpr-mutant virions. (**k**) CD69 upregulation in response to IL7 on resting memory CD4+ T cells infected with HIV-1 WT, ΔVpr or Vpr mutants (n=9). (**l**) as for (k) showing CD69 expression on HIV-1 infected Gag+ memory T cells compared to uninfected Gag- bystander cells. (**m**) Co-expression of CD69 and CXCR6 from (k) (n=9). Data are the mean±SEM. Paired two-tailed *t*-test or one-way ANOVA with Bonferroni or Dunnett's post-test were used. 2LTR circles (i) were compared by unpaired one-tailed *t*-test. *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$; n.s., not significant.

Fig. 4 Transcriptional profiling of HIV-1 infected resting memory CD4+ T cells. (**a**) Heatmap showing hierarchical clustering of 226 differentially expressed genes (DEG) of infected (HIV-1 WT) over uninfected (Mock) resting memory CD4+ T cells (adjusted p-value < 0.01, Fold change +/- 1.2). Mean log2 TPM of 4 biological repeats are shown. Cytokine indicates presence or absence of IL7. Virus indicates infection with HIV-1 WT, HIV-1 ΔVpr or uninfected (Mock) conditions. (**b**) Principal component analysis (PCA) of (a), with ellipses indicating 95% CI. (**c**) and (**d**) show scatter plots of mean log2 TPMs of DEGs from HIV-1 WT/Mock (grey circles) or HIV-1 ΔVpr/Mock (orange circles) in the absence (c) or presence (d) of IL7 (adjusted p-value < 0.01, Fold change +/- 1.2). Lines indicate line of identity (LOD). Genes above or below LOD are up or downregulated, respectively. (**e**) and (**f**) Venn diagrams showing overlap of DEGs comparing expression profiles of HIV-1 WT/Mock with (e) HIV-1 ΔVpr\Mock or (f) HIV-1 ΔVpr/HIV-1 WT. (**g**) GSEA was performed on expression profiles comparing HIV-1 WT / Mock (black) or HIV-1 ΔVpr/HIV-1 WT (grey). Normalised enrichment scores are shown for significantly enriched Hallmark gene sets are shown (FDR q-value<0.05 and NES>1.75). (**h**) and (**i**) top ten significantly enriched canonical pathways

predicted by ingenuity pathway (IPA) analysis of DEGs (h) HIV-1 WT/Mock or (i) HIV-1 ΔVpr/HIV-1

WT (adjusted p-value<0.05). (**j**) Cytokines and (**k**) transcription regulators predicted to be upstream

regulators by IPA of gene expression signatures HIV-1 WT/Mock (black) or HIV-1 ΔVpr/Mock

(grey), line indicates p=0.05.

Fig. 5 Vpr drives a T$_{RM}$-like transcriptomic program in HIV-1 infected resting memory CD4+

T cells. (**a**) Heatmap showing hierarchical clustering based on a T$_{RM}$ core gene expression

signature[21] that was performed to compare transcriptional profiles of HIV-1 infected resting

memory CD4 T cells (Reuschl et al, 2021) (Mock, HIV-1 WT, HIV-1 ΔVpr) with previously

described gene expression profiles (Kumar et al, 2017) of T$_{RM}$ (CD69 POS), non-T$_{RM}$ (CD69 NEG)

tissue-derived T cells (lung and spleen) and blood-derived CD69- (CD69 NEG) T cells. Cytokine

indicates presence or absence of IL7. Virus indicates infection with HIV-1 WT, HIV-1 ΔVpr or

uninfected (Mock) conditions. n/a, not applicable. (**b**) shows the T$_{RM}$ signature score for the

indicated conditions calculated based on (a). Subsets from[21] are indicated in red, shown are CD4+

or CD8+ T cells from lungs or spleens. T$_{RM}$+, CD69+ T cells; T$_{RM}$-, CD69- T cells. T$_{RM}$ signature

scores for resting CD4 memory T cells infected or uninfected are shown in the presence or

absence of IL7. Means are shown.

Extended data Fig. 1. (**a**) Experimental set-up schematic. (**b**) Flow cytometry gating strategy. (**c**)

Ki67, CD69 and HLA-DR expression of CD3+CD4+ T cells from unstimulated PBMCs (n=8). (**d**)

Ki67 expression on resting and activated primary CD4+ T cells. Representative flow cytometry plots.

(**e**) MCM2 expression on resting and activated primary CD4+ T cells. Representative flow cytometry

plots. (**f**) Resting or (**g**) mitogenically-activated primary target CD4+ T cells cultured with HIV-1

infected Jurkat T cells separated by a 0.4µm transwell or in direct co-culture. Target cell infection

levels was measured by intracellular staining for Gag. Representative flow cytometry plots are

shown. (**h**) Infection levels of target CD4+ T cells determined by intracellular Gag staining and flow cytometry (n=2). (**i**) Proportion of CD45RA+ naive and CD45RA- memory CD4+ T cells in unstimulated PBMCs (n=8). (**j**) Resting target CD4+ T cells were cultured with mock-treated or HIV-1-infected donor cells. Surface expression of CD45RA and CCR7 were measured after 72h of co-culture. Representative flow cytometry plots are shown. (**k**) Quantification of T cell subsets in infected (Gag+) and uninfected (Gag-) resting CD4+ T cells according to CD45RA/CD62L expression (n=3). T_{naive} = CD45RA+/CD62L+, T_{EMRA} = CD45RA+/CD62L-, T_{EM} = CD45RA-/CD62L-, T_{CM} = CD45RA-/CD62L+. Data are shown as mean±SEM.

Extended data Fig. 2. (**a**) Representative flow cytometry plots of CD45RA+ and CD45RA- CD4+ T cells pre- and post-isolation. (**b**) Representative histogram of intracellular Gag staining in resting naïve (CD45RA+) and memory (CD45RA-) CD4+ T cells after 72h of cell-to-cell spread ± T20. (**c**) Representative histogram of intracellular Gag-levels in resting naïve (CD45RA+) and memory (CD45RA-) CD4+ T cells after 72h of cell-to-cell spread ± Efavirenz. (**d**) Representative histogram of CD4 surface levels in resting naïve (CD45RA+) and memory (CD45RA-) CD3+ T cells after 72h of cell-to-cell spread. (**e**) Mean post-sort population purity of T cells from (Fig. 1 n-p) was 99.92% target cells of which 99.86% were memory T cells (n=5).

Extended data Fig. 3 (**a** and **b**) Representative flow cytometry plots showing (**a**) Ki67 and Gag, or (**b**) MCM2 and Gag staining of resting CD4+ T cells after co-culture with mock or HIV-1 infected primary donor CD4+ T cells. (**c**) FACS sorted CD69- resting naïve or memory CD4+ T cells co-cultured with HIV-1 infected primary CD4+ donor T cells and infection of targets measured by Gag staining (n=4). (**d**) FACS sorted CD69- CD4+ T cells co-cultured with HIV-1 infected primary CD4+ donor T cells. CD69 expression was measured on resting memory CD4+ T cells (n=4). (**e**) Total CD69 expression alongside CD69 with or without HLA-DR co-expression on infected resting memory T cells (n=6). (**f**) Expression of T_{RM}-markers CXCR6, CD49A and CD69 on resting CD4+ memory T cells from unstimulated PBMCs (n=8). (**g**) Quantification of cell-to-cell spread of HIV-1 WT to resting memory CD4+ T cells in the presence or absence of IL7 (n=10). (**h** and **i**) CD69 expression on

infected resting memory CD4+ target T cells in the presence or absence of (**h**) IL12 (n=4) or (**i**) TFGβ

(n=4). (**j**) CD69 expression on infected resting memory CD4+ T cells with IL7 added at the indicated

times post cell-mixing (n=5). (**k**) CD69/CXCR6 co-expression from (**j**) (n=5). (**l**) CD69/CD101 co-

expression on infected resting memory CD4+ T cells (n=3) (**m**) CD69 upregulation in response to

IL7 on resting memory CD4+ T cells infected HIV-1 NL4.3, or transmitter-founder viruses CH040

and CH077 comparing infected Gag+ and uninfected Gag- bystander cells (n=7). (**n**) CD69/CXCR6

co-expression on resting memory CD4+ T cells from (**i**) (n=7). (**o**) CD69/CX3CR1 co-expression on

infected resting memory CD4+ T cells (n=3). (**p**) CD103 expression and (**q**) CD69/CD103 co-

expression on infected resting memory CD4+ T cells (n=4). Data are the mean±SEM. Paired two-

tailed *t*-test or one-way ANOVA with Bonferroni post-test was used. For (**i**), median+IQR is shown

and Kruskall-Wallis test was used to compare groups *, p<0.05 ; **, p<0.01; ***, p<0.001; n.s., not

significant.

Extended data Fig. 4 Resting memory CD4+ T cells were co-cultured with HIV-1 infected primary

CD4+ T cells infected with HIV-1 WT or mutant viruses. Representative flow cytometry plots of HIV-

1 Gag and CD69 co-expression in the presence or absence of IL7 from three independent

experiments are shown (**a-c**).

Extended data Fig. 5 (**a**) CD69 surface expression on resting CD45RA- memory CD4+ T cells

following co-culture with primary CD4+ donor T cells infected with HIV-1 NL4.3 (WT), ΔVpr, ΔNef or

ΔVpu or uninfected (mock) donors (n=9). (**b**) CXCR6 expression from (**a**) (n=9). (**c**) CD69/CXCR6

co-expression from (**a**) (n=9). (**d**) As for (b) but cells were incubated in the presence of IL7 (n=9). (**e**)

CD69/CXCR6 surface co-expression from (d) (n=9). (**f**) Gag MFI of cell-to-cell spread of HIV-1 WT

and ΔVpr to resting memory CD4+ T cells (n=10). Correlation plot of CD69 MFI with Gag MFI in

absence (**g**) or presence (**h**) of IL7 (n=18). (**i**) *S1PR1* and *KLF2* mRNA levels in FACS sorted resting

memory CD4+ T cells from Fig. 3e. Fold change over mock is shown (n=5). (**j**) CD69 expression on

infected resting memory CD4+ T cells ± Ruxilitinib (n=4). (**k**) CD127 MFI on infected resting memory

CD4+ T cells ± IL7 (n=7). (**l**) Representative histogram of intracellular STAT5-phosphorylation in

infected resting memory T cells. (**m**) Quantification of (**l**) (n=10). Data are the mean±SEM. One-way

ANOVA with Dunnet's post-test was used. Statistical significance is shown relative to mock treated

cells (no HIV-1). R^2 in (g) and (h) was determined by simple linear regression. *, p<0.05 ; **, p<0.01;

***, p<0.001; n.s., not significant.

Extended data Fig. 6 (**a**) and (**b**) Venn diagrams showing overlap of DEGs comparing expression

profiles of HIV-1 WT+IL7/Mock+IL7 with (**e**) HIV-1 ΔVpr+IL7/\Mock+IL7 or (**f**) HIV-1 ΔVpr+IL7/HIV-1

WT+IL7. (**g**) GSEA was performed on expression profiles comparing HIV-1 WT+IL7/Mock+IL7

(black) or HIV-1 ΔVpr+IL7/HIV-1 WT+IL7 (grey). Normalised enrichment scores are shown for

significantly enriched Hallmark gene sets are shown (FDR q-value<0.05 and NES>1.75). (**h**) and (**i**)

top ten significantly enriched canonical pathways predicted by ingenuity pathway (IPA) analysis of

DEGs (h) HIV-1 WT+IL7/Mock+IL7 or (i) HIV-1 ΔVpr+IL7/HIV-1 WT+IL7 (adjusted p-value<0.05). (**j**)

Cytokines and (**k**) transcription regulators predicted to be upstream regulators by IPA of gene

expression signatures HIV-1 WT+IL7/Mock+IL7 (black) or HIV-1 ΔVpr+IL7/Mock+IL7 (grey), line

indicates p=0.05.

Extended data Fig. 7 (**a**) Western blot showing siRNA knockdown of DCAF1 in CD3/CD28-activated

CD4+ T cells 48h post transfection. Two representative samples are shown. (**b**) Number of live CD3+

Target T cells recovered after 72h of cell-to-cell spread into control or DCAF1 siRNA-treated T cells

(n=3).

Fig. 1 HIV-1 exploits cell-to-cell spread to preferentially infect resting memory CD4+ T cells. HIV-1 NL4.3 infected (**a**) Jurkat or (**b**) mitogenically-activated primary CD4+ donor T cells co-cultured with resting primary CD4+ target T cells separated by a 0.4μm transwell (cell-free) or direct co-culture (cell-cell). Target cell infection was measured by intracellular staining for HIV-1 Gag protein. Representative flow cytometry plots are shown. Bar graphs show mean of independent experiments (n=4). (**c,d**) Cell-to-cell spread from activated primary donor CD4+ T cells to resting primary target CD4+ T cells preferentially infects CD45RA-memory CD4+ T cells. A representative flow cytometry plot and quantification is shown (n=4). (**e**) Quantification of infection performed as in (**c**) (n=11). (**f**) HIV-1 infection of target CD4+ T cells as part of the total resting CD4+ T cell population (total) compared to pre-isolated naive and memory CD4+ target T cells (isolated) (n=9). (**g**) Representative flow cytometry plots of cell-to-cell infection of resting CD4+ T cells with CCR5-tropic HIV-1 NL4.3 BaL and transmitter founder viruses HIV-1 CH040 and CH077 as performed in (**c**). (**h**) Quantification of infection of CXCR4 (X4) and CCR5 (R5)-tropic viruses (n=4) and (**i**) transmitter/founder viruses HIV-1 CH040 and CH077 (n=7). (**j**) Cell-to-cell infection of resting CD4+ T cells is reduced by the HIV-1 fusion-inhibitor T20 (n=6) and (**k**) the reverse transciptase inhibitor Efavirenz (n=6) measured by intracellular Gag staining (MFI) or (**l**) HIV-1 LTR-driven GFP-reporter gene expression (n=4). (**m**) HIV-1 infection downregulates CD4 expression. Shown are the percentage of CD4+ cells of the total CD3+ target cell population (n=6). Resting CD4+ memory T cells were isolated after 72h of cell-to-cell spread by FACS sorting and cultured for 4 days. HIV-1 infection was measured by intracellular Gag staining (**n**) and virus release measured by culture supernatant RT activity (**o**) (n=5-7). T cells from (**n**) were recovered at day 1 or 4 post isolation and cultured with uninfected eFluor450+ target Jurkat T cells. Infection of Jurkat T cells was measured after 72h (**p**) (n=3). Data are the mean±SEM. Paired two-tailed *t*-test or one-way ANOVA with Bonferroni post-test were used. For (**m**), median+IQR is shown and Friedman test with Dunn's post-test was used. For (**o**), unpaired one-tailed *t*-test was used. *, p<0.05 ; **, p<0.01; ***, p<0.001; n.s., not significant.

Fig. 2 HIV-1 infection induces a T_RM-like phenotype in resting memory CD4+ T cells. (**a**) CD69 expression on resting memory CD4+ target T cells following co-culture with HIV-1 infected primary donor T cells or uninfected donor T cells (mock) (n=17). (**b**) Representative flow cytometry plots from (**a**). (**c**) CD69 expression on infected resting memory CD4+ T cells ± IL7 and T20 (n=7). (**d**) CD69 expression on infected resting memory CD4+ T cells ± IL7 and Ruxilitinib (n=8). (**e**) CD69 expression on infected resting memory CD4+ T cells in response to IL7 and IL15 (n=11). (**f**) CD69 expression on infected Gag+ resting memory CD4+ T cells and uninfected Gag- bystander cells in response to IL7 and IL15 (n=11). (**g**) CXCR6 surface expression from (**f**) (n=11). (**h**) Representative flow cytometry plots of CD69 and CXCR6 co-expression. (**i**) Co-expression of CD69 with CXCR6, CD49A or PD-1 on infected resting memory CD4+ T cells (n=5-7). (**j**) As for (**i**) in the presence of IL7 (n=4-7). (**k**) As for (**i**) comparing infected Gag+ memory CD4+ T cells and uninfected Gag- bystander cells. (**l**) and (**m**) Blimp-1 expression in CD69+ infected resting memory CD4+ T cells and infected CD69- cells in response to IL7 (n=8). (**n**) Total lymphocytes from cellularised tonsils co-cultured with HIV-1 infected Jurkat T cells. Infection of resting CD4+ T cells (live CD3+/CD8-/Ki67- lymphocytes) shown as CD45RO vs Gag. (**o**) CD69 and CXCR6 co-expression on infected Gag+ and uninfected Gag- tonsil resting memory CD4+ T cells ± IL7. (**p**) Total lymphocytes from mediastinal lymph nodes co-cultured with HIV-1 infected activated autologous LN-derived lymphocytes. Infection of resting CD4+ T cells (live CD3+/CD8-/Ki67- lymphocytes) shown as CD45RA vs Gag. (**q**) CD69 and CXCR6 co-expression on infected Gag+ and uninfected Gag- lymph node resting memory CD4+ T cells ±IL7. Data are the mean±SEM. Paired two-tailed *t*-test or one-way ANOVA with Bonferroni or Dunnett's post-test were used. *p<0.05 ; **, p<0.01; ***, p<0.001; n.s., not significant.

Fig. 3 Vpr drives HIV-1-induced T$_{RM}$-induction in resting memory CD4+ T cells. Resting memory CD4+ T cells were co-cultured with HIV-1 infected primary CD4+ T cells infected with HIV-1 WT or mutant viruses, or uninfected donor cells (mock). (**a**) CD69 upregulation in response to IL7 compared to mock (n=9). (**b**) CD69 expression on HIV-1 infected Gag+ resting memory CD4+ T cells compared to uninfected Gag- bystander cells (n=9). (**c**) Quantification of cell-to-cell spread of HIV-1 WT and ΔVpr to resting memory CD4+ T cells (n=9). (**d**) CD69/CXCR6/CD49a co-expression on resting memory CD4+ T cells infected with HIV-1 WT or ΔVpr (n=9). (**e**) *CD69*, *CXCR6* and *PRDM1* (Blimp1) mRNA levels from FACS sorted infected resting memory CD4+ T cells. Fold change relative to unifected (mock) is shown (n=5). (**f**) IFNg expression by HIV-1 infected resting memory CD4+ T cells at 72h in response to IL7 (n=3). (**g**) CD69 and (**h**) CD69/CXCR6 co-expression in response to IL7 in the presence of integrase inhibitor Raltegravir (n=6). (**i**) Quantification of integrated provirus and 2LTR circles in FACS sorted target CD4+ memory T cells after 72h of cell-to-cell spread in the presence or absence of Raltegravir. (**j**) Western blot showing Vpr packaging into HIV-1 WT and Vpr-mutant virions. (**k**) CD69 upregulation in response to IL7 on resting memory CD4+ T cells infected with HIV-1 WT, ΔVpr or Vpr mutants (n=9). (**l**) as for (**k**) showing CD69 expression on HIV-1 infected Gag+ memory T cells compared to uninfected Gag-bystander cells. (**m**) Co-expression of CD69 and CXCR6 from (**k**) (n=9). Data are the mean±SEM. Paired two-tailed *t*-test or one-way ANOVA with Bonferroni or Dunnett's post-test were used. 2LTR circles (**i**) were compared by unpaired one-tailed *t*-test. *, p<0.05 ; **, p<0.01; ***, p<0.001; n.s., not significant.

Fig. 4 Transcriptional profiling of HIV-1 infected resting memory CD4+ T cells. (**a**) Heatmap showing hierarchical clustering of 226 differentially expressed genes (DEG) of infected (HIV-1 WT) over uninfected (Mock) resting memory CD4+ T cells (adjusted p-value < 0.01, Fold change +/- 1.2). Mean log2 TPM of 4 biological repeats are shown. Cytokine indicates presence or absence of IL7. Virus indicates infection with HIV-1 WT, HIV-1 ΔVpr or uninfected (Mock) conditions. (**b**) Principal component analysis (PCA) of (a), with ellipses indicating 95% CI. (**c**) and (**d**) show scatter plots of mean log2 TPMs of DEGs from HIV-1 WT/Mock (grey circles) or HIV-1 ΔVpr/Mock (orange circles) in the absence (c) or presence (d) of IL7 (adjusted p-value < 0.01, Fold change +/-1.2). Lines indicate line of identity (LOD). Genes above or below LOD are up or downregulated, respectively. (**e**) and (**f**) Venn diagrams showing overlap of DEGs comparing expression profiles of HIV-1 WT/Mock with (e) HIV-1 ΔVpr/\Mock or (f) HIV-1 ΔVpr/HIV-1 WT. (**g**) GSEA was performed on expression profiles comparing HIV-1 WT / Mock (black) or HIV-1 ΔVpr/HIV-1 WT (grey). Normalised enrichment scores are shown for significantly enriched Hallmark gene sets are shown (FDR q-value<0.05 and NES>1.75). (**h**) and (**i**) top ten significantly enriched canonical pathways predicted by ingenuity pathway (IPA) analysis of DEGs (h) HIV-1 WT/Mock or (i) HIV-1 ΔVpr/HIV-1 WT (adjusted p-value<0.05). (**j**) Cytokines and (**k**) transcription regulators predicted to be upstream regulators by IPA of gene expression signatures HIV-1 WT/Mock (black) or HIV-1 ΔVpr/Mock (grey), line indicates p=0.05.

Fig. 5 Vpr drives a T_{RM}-like transcriptomic programme in HIV-1 infected resting memory CD4+ T cells. (**a**) Heatmap showing hierarchical clustering based on a T_{RM} core gene expression signature[21] that was performed to compare transcriptional profiles of HIV-1 infected resting memory CD4 T cells (Reuschl et al, 2021) (Mock, HIV-1 WT, HIV-1 ΔVpr) with previously described gene expression profiles (Kumar et al, 2017) of T_{RM} (CD69 POS), non-T_{RM} (CD69 NEG) tissue-derived T cells (lung and spleen) and blood-derived CD69- (CD69 NEG) T cells. Cytokine indicates presence or absence of IL7. Virus indicates infection with HIV-1 WT, HIV-1 ΔVpr or uninfected (Mock) conditions. n/a, not applicable. (**b**) shows the T_{RM} signature score for the indicated conditions calculated based on (**a**). Subsets from[21] are indicated in red, shown are CD4+ or CD8+ T cells from lungs or spleens. T_{RM}+, CD69+ T cells; T_{RM}-, CD69- T cells. T_{RM} signature scores for resting CD4 memory T cells infected or uninfected are shown in the presence or absence of IL7. Means are shown.

Extended data Fig. 1. (**a**) Experimental set-up schematic. (**b**) Flow cytometry gating strategy. (**c**) Ki67, CD69 and HLA-DR expression of CD3+CD4+ T cells from unstimulated PBMCs (n=8). (**d**) Ki67 expression on resting and activated primary CD4+ T cells. Representative flow cytometry plots. (**e**) MCM2 expression on resting and activated primary CD4+ T cells. Representative flow cytometry plots. (**f**) Resting or (**g**) mitogenically-activated primary target CD4+ T cells cultured with HIV-1 infected Jurkat T cells separated by a 0.4μm transwell or in direct co-culture. Target cell infection levels was measured by intracellular staining for Gag. Representative flow cytometry plots are shown. (**h**) Infection levels of target CD4+ T cells determined by intracellular Gag staining and flow cytometry (n=2). (**i**) Proportion of CD45RA+ naive and CD45RA- memory CD4+ T cells in unstimulated PBMCs (n=8). (**j**) Resting target CD4+ T cells were cultured with mock-treated or HIV-1-infected donor cells. Surface expression of CD45RA and CCR7 were measured after 72h of co-culture. Representative flow cytometry plots are shown. (**k**) Quantification of T cell subsets in infected (Gag+) and uninfected (Gag-) resting CD4+ T cells according to CD45RA/CD62L expression (n=3). T$_{naive}$ = CD45RA+/CD62L+, T$_{EMRA}$ = CD45RA+/CD62L-, T$_{EM}$ = CD45RA-/CD62L-, T$_{CM}$ = CD45RA-/CD62L+. Data are shown as mean±SEM.

Extended data Fig. 2. (**a**) Representative flow cytometry plots of CD45RA+ and CD45RA- CD4+ T cells pre- and post-isolation. (**b**) Representative histogram of intracellular Gag staining in resting naïve (CD45RA+) and memory (CD45RA-) CD4+ T cells after 72h of cell-to-cell spread ± T20. (**c**) Representative histogram of intracellular Gag-levels in resting naïve (CD45RA+) and memory (CD45RA-) CD4+ T cells after 72h of cell-to-cell spread ± Efavirenz. (**d**) Representative histogram of CD4 surface levels in resting naïve (CD45RA+) and memory (CD45RA-) CD3+ T cells after 72h of cell-to-cell spread. (**e**) Mean post-sort population purity of T cells from (Fig. 1 n-p) was 99.92% target cells of which 99.86% were memory T cells (n=5).

Extended data Fig. 3 (**a** and **b**) Representative flow cytometry plots showing (**a**) Ki67 and Gag, or (**b**) MCM2 and Gag staining of resting CD4+ T cells after co-culture with mock or HIV-1 infected primary donor CD4+ T cells. (**c**) FACS sorted CD69- resting naïve or memory CD4+ T cells co-cultured with HIV-1 infected primary CD4+ donor T cells and infection of targets measured by Gag staining (n=4). (**d**) FACS sorted CD69- CD4+ T cells co-cultured with HIV-1 infected primary CD4+ donor T cells. CD69 expression was measured on resting memory CD4+ T cells (n=4). (**e**) Total CD69 expression alongside CD69 with or without HLA-DR co-expression on infected resting memory T cells (n=6). (**f**) Expression of T_{RM}-markers CXCR6, CD49A and CD69 on resting CD4+ memory T cells from unstimulated PBMCs (n=8). (**g**) Quantification of cell-to-cell spread of HIV-1 WT to resting memory CD4+ T cells in the presence or absence of IL7 (n=10). (**h** and **i**) CD69 expression on infected resting memory CD4+ target T cells in the presence or absence of (**h**) IL12 (n=4) or (**i**) TFGβ (n=4). (**j**) CD69 expression on infected resting memory CD4+ T cells with IL7 added at the indicated times post cell-mixing (n=5). (**k**) CD69/CXCR6 co-expression from (**j**) (n=5). (**l**) CD69/CD101 co-expression on infected resting memory CD4+ T cells (n=3) (**m**) CD69 upregulation in response to IL7 on resting memory CD4+ T cells infected HIV-1 NL4.3, or transmitter-founder viruses CH040 and CH077 comparing infected Gag+ and uninfected Gag- bystander cells (n=7). (**n**) CD69/CXCR6 co-expression on resting memory CD4+ T cells from (**i**) (n=7). (**o**) CD69/CX3CR1 co-expression on infected resting memory CD4+ T cells (n=3). (**p**) CD103 expression and (**q**) CD69/CD103 co-expression on infected resting memory CD4+ T cells (n=4). Data are the mean±SEM. Paired two-tailed *t*-test or one-way ANOVA with Bonferroni post-test was used. For (**i**), median+IQR is shown and Kruskall-Wallis test was used to compare groups *, p<0.05 ; **, p<0.01; ***, p<0.001; n.s., not significant.

Extended data Fig. 4 Resting memory CD4+ T cells were co-cultured with HIV-1 infected primary CD4+ T cells infected with HIV-1 WT or mutant viruses. Representative flow cytometry plots of HIV-1 Gag and CD69 co-expression in the presence or absence of IL7 from three independent experiments are shown (**a-c**).

Extended data Fig. 5 (**a**) CD69 surface expression on resting CD45RA- memory CD4+ T cells following co-culture with primary CD4+ donor T cells infected with HIV-1 NL4.3 (WT), ΔVpr, ΔNef or ΔVpu or uninfected (mock) donors (n=9). (**b**) CXCR6 expression from (**a**) (n=9). (**c**) CD69/CXCR6 co-expression from (**a**) (n=9). (**d**) As for (b) but cells were incubated in the presence of IL7 (n=9). (**e**) CD69/CXCR6 surface co-expression from (d) (n=9). (**f**) Gag MFI of cell-to-cell spread of HIV-1 WT and ΔVpr to resting memory CD4+ T cells (n=10). Correlation plot of CD69 MFI with Gag MFI in absence (**g**) or presence (**h**) of IL7 (n=18). (**i**) *S1PR1* and *KLF2* mRNA levels in FACS sorted resting memory CD4+ T cells from Fig. 3e. Fold change over mock is shown (n=5). (**j**) CD69 expression on infected resting memory CD4+ T cells ± Ruxilitinib (n=4). (**k**) CD127 MFI on infected resting memory CD4+ T cells ± IL7 (n=7). (**l**) Representative histogram of intracellular STAT5-phosphorylation in infected resting memory T cells. (**m**) Quantification of (**l**) (n=10). Data are the mean±SEM. One-way ANOVA with Dunnet's post-test was used. Statistical significance is shown relative to mock treated cells (no HIV-1). R^2 in (g) and (h) was determined by simple linear regression. *, p<0.05 ; **, p<0.01; ***, p<0.001; n.s., not significant.

Extended data Fig. 6 (**a**) and (**b**) Venn diagrams showing overlap of DEGs comparing expression profiles of HIV-1 WT+IL7/Mock+IL7 with (**e**) HIV-1 ΔVpr+IL7/\Mock+IL7 or (**f**) HIV-1 ΔVpr+IL7/HIV-1 WT+IL7. (**g**) GSEA was performed on expression profiles comparing HIV-1 WT+IL7/Mock+IL7 (black) or HIV-1 ΔVpr+IL7/HIV-1 WT+IL7 (grey). Normalised enrichment scores are shown for significantly enriched Hallmark gene sets are shown (FDR q-value<0.05 and NES>1.75). (**h**) and (**i**) top ten significantly enriched canonical pathways predicted by ingenuity pathway (IPA) analysis of DEGs (h) HIV-1 WT+IL7/Mock+IL7 or (i) HIV-1 ΔVpr+IL7/HIV-1 WT+IL7 (adjusted p-value<0.05). (**j**) Cytokines and (**k**) transcription regulators predicted to be upstream regulators by IPA of gene expression signatures HIV-1 WT+IL7/Mock+IL7 (black) or HIV-1 ΔVpr+IL7/Mock+IL7 (grey), line indicates p=0.05.

Extended data Fig. 7 (**a**) Western blot showing siRNA knockdown of DCAF1 in CD3/CD28-activated CD4+ T cells 48h post transfection. Two representative samples are shown. (**b**) Number of live CD3+ Target T cells recovered after 72h of cell-to-cell spread into control or DCAF1 siRNA-treated T cells (n=3).