Genome-wide analysis of mitochondrial DNA copy number reveals multiple loci implicated in nucleotide metabolism, platelet activation, and megakaryocyte proliferation

Longchamps RJ[1*], Yang SY[1*], Castellani CA[1,2], Shi W[1], Lane J[3], Grove ML[4], Bartz TM[5], Sarnowski C[6], Burrows K[7,8], Guyatt AL[9], Gaunt TR[7,8], Kacprowski T[10], Yang J[11], De Jager PL[12,13], Yu L[11], CHARGE Aging and Longevity Group, Bergman A[14], Xia R[15], Fornage M[15,16], Feitosa MF[17], Wojczynski MK[17], Kraja AT[17], Province MA[17], Amin N[18], Rivadeneira F[19], Tiemeier H[18,20], Uitterlinden AG[18,19], Broer L[19], Van Meurs JBJ[18,19], Van Duijn CM[18], Raffield LM[21], Lange L[22], Rich SS[23], Lemaitre RN[24], Goodarzi MO[25], Sitlani CM[24], Mak ACY[26], Bennett DA[11], Rodriguez S[7,8], Murabito JM[27], Lunetta KL[6], Sotoodehnia N[28], Atzmon G[29], Kenny Y[30], Barzilai N[31], Brody JA[32], Psaty BM[33], Taylor KD[34], Rotter JI[34], Boerwinkle E[4,35], Pankratz N[3], Arking DE[1]

* Indicates authors contributed equally to this work.

McKusick-Nathans Institute, Department of Genetic Medicine, Johns Hopkins University School of Medicine, Baltimore, MD

Department of Pathology and Laboratory Medicine, Western University, London, ON, Canada

Department of Laboratory Medicine and Pathology, University of Minnesota Medical School, Minneapolis, MN

Human Genetics Center, Department of Epidemiology, Human Genetics, and Environmental Sciences, School of Public Health, The University of Texas Health Science Center at Houston, Houston, TX

Cardiovascular Health Research Unit, Departments of Medicine and Biostatistics, University of Washington, Seattle, WA

Department of Biostatistics, Boston University School of Public Health, Boston, MA

MRC Integrative Epidemiology Unit at the University of Bristol, University of Bristol, Oakfield House, Oakfield Grove, Bristol, UK

8 Population Health Sciences, Bristol Medical School, University of Bristol, Oakfield House,

Oakfield Grove, Bristol, UK

9 Department of Health Sciences, University of Leicester, University Road, Leicester, UK

10 Department of Functional Genomics, Interfaculty Institute for Genetics and Functional

Genomics, University of Greifswald, Greifswald, Germany; Division Data Science in

Biomedicine, Peter L. Reichertz Institute for Medical Informatics, TU Braunschweig and

Hannover Medical School, Brunswick, Germany

11 Rush Alzheimer's Disease Center & Department of Neurological Sciences, Rush University

Medical Center, Chicago, IL

12 Center for Translational and Systems Neuroimmunology, Department of Neurology,

Columbia University Medical Center, New York, NY

13 Program in Medical and Population Genetics, Broad Institute, Cambridge, MA

14 Department of Systems and Computational Biology, Albert Einstein College of Medicine,

Bronx, New York, USA

15 Institute of Molecular Medicine, The University of Texas health Science Center at Houston,

Houston TX

16 Human Genetics Center, The University of Texas Health Science Center at Houston,

Houston

17 Division of Statistical Genomics, Department of Genetics, Washington University School of

Medicine

18 Department of Epidemiology, Erasmus Medical Center, Rotterdam, The Netherlands

19 Department of Internal Medicine, Erasmus Medical Center, Rotterdam, The Netherlands

20 Department of Social and Behavioral Science, Harvard T.H. School of Public Health, Boston,

USA

21 Department of Genetics, University of North Carolina at Chapel Hill, Chapel Hill, NC

22 Department of Medicine, University of Colorado Denver, Anschutz Medical Campus, Aurora,

CO

23 Center for Public Health Genomics, University of Virginia, Charlottesville, Virginia, USA

24 Cardiovascular Health Research Unit, Department of Medicine, University of Washington,

Seattle, WA

25 Division of Endocrinology, Diabetes and Metabolism, Cedars-Sinai Medical Center, Los

Angeles, CA

26 Cardiovascular Research Institute and Institute for Human Genetics, University of California,

San Francisco, California

27 Boston University School of Medicine, Boston University, Boston, MA

28 Cardiovascular Health Research Unit, Division of Cardiology, University of Washington,

Seattle, WA

29 Department of Natural science, University of Haifa, Haifa, Israel; Departments of Medicine

and Genetics, Albert Einstein College of Medicine, Bronx, NY, 10461, USA.

30 Department of Epidemiology and Population Health, Albert Einstein College of Medicine,

Bronx, NY, 10461, USA.

31 Departments of Medicine and Genetics, Albert Einstein College of Medicine, Bronx, NY,

10461, USA.

32 Cardiovascular Health Research Unit, Department of Medicine, University of Washington,

Seattle, WA

33 Cardiovascular Health Research Unit, Departments of Epidemiology, Medicine and Health

Services, University of Washington, Seattle, WA

34 The Institute for Translational Genomics and Population Sciences, Department of Pediatrics,

The Lundquist Institute for Biomedical Innovation at Harbor-UCLA Medical Center, Torrance,

CA

35 Baylor College of Medicine, Human Genome Sequencing Center, Houston, TX

Abstract

Blood-derived mitochondrial DNA copy number (mtDNA-CN) is a minimally invasive proxy measure of mitochondrial function that exhibits both inter-individual and intercellular variation. While mtDNA-CN has been previously associated with various aging-related diseases, little is known about the genetic factors that may modulate this phenotype. We performed a genome-wide association study (GWAS) in 465,809 individuals of White (European) ancestry from the Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) consortium and the UK Biobank (UKB). We identified 129 SNPs with statistically significant, independent effects associated with mtDNA-CN across 96 loci. A combination of fine-mapping, variant annotation, co-localization, and gene set enrichment analyses were used to prioritize genes within each of the 129 independent sites. Putative causal genes were enriched for known mitochondrial DNA depletion syndromes ($p = 3.09 \times 10^{-15}$) and the gene ontology (GO) terms for mtDNA metabolism ($p = 1.43 \times 10^{-8}$) and mtDNA replication ($p = 1.2 \times 10^{-7}$). A clustering approach leveraged pleiotropy between mtDNA-CN associated SNPs and 42 mtDNA-CN associated phenotypes to identify functional domains, revealing five distinct groups, including platelet activation, megakaryocyte proliferation, and mtDNA metabolism. In conclusion, in a GWAS of mtDNA-CN conducted in >450,000 individuals, we identified SNPs within loci that implicate novel pathways that provide a framework for defining the underlying mechanisms involved in genetic control of mtDNA-CN.

Introduction

Mitochondria are the cellular organelles primarily responsible for producing the chemical energy required for metabolism, as well as signaling the apoptotic process, maintaining homeostasis, and synthesizing several macromolecules such as lipids, heme and iron-sulfur clusters[1,2].

Mitochondria possess their own genome (mtDNA); a circular, intron-free, double-stranded,

haploid, ~ 16.6 kb maternally inherited molecule encoding 37 genes vital for proper

mitochondrial function. Due to the integral role of mitochondria in cellular metabolism,

mitochondrial dysfunction is known to play a critical role in the underlying etiology of several

aging-related diseases[3–5].

Unlike the nuclear genome, a large amount of variation exists in the number of copies of mtDNA

present within cells, tissues, and individuals. The relative copy number of mtDNA (mtDNA-CN)

has been shown to be positively correlated with oxidative stress[6], energy reserves, and

mitochondrial membrane potential[7]. As a minimally invasive proxy measure of mitochondrial

dysfunction[8], decreased blood-derived mtDNA-CN has been previously associated with aging-

related disease states including frailty[9], cardiovascular disease[10–12], chronic kidney disease[13],

neurodegeneration[14,15], and cancer[16].

Although mtDNA-CN measured from whole blood presents itself as an easily accessible and

minimally invasive biomarker, cell type composition has been shown to be an important

confounder, complicating analyses[17,18]. For example, while platelets generally have fewer

mtDNA molecules than leukocytes, the lack of a platelet nuclear genome drastically skews

mtDNA-CN estimates. As a result, not only is controlling for cell composition extremely vital for

accurate mtDNA-CN estimation, but interpreting the results in relation to the impact of cell

composition becomes a necessity[18–20].

Although the comprehensive mechanism through which mtDNA-CN is modulated is largely

unknown[21,22], twin studies have estimated broad-sense heritability ~0.65, consistent with

moderate genetic control[23]. Several nuclear genes have been shown to directly modulate

mtDNA-CN, specifically those within the mtDNA replication machinery such as the mitochondrial

polymerase, *POLG* and *POLG2*[24,25], as well as the mitochondrial DNA helicase, *TWNK*, and the

mitochondrial single-stranded binding protein, *mtSSB*[26]. Furthermore, nuclear genes which

maintain proper mitochondrial nucleotide supply including *DGUOK* and *TK2* have also been

shown to regulate mtDNA-CN[27–29]. To further elucidate the genetic control over mtDNA-CN,

several genome-wide association studies (GWAS) of mtDNA-CN have been published[30–33],

including a study that was published while the current manuscript was in preparation, analyzing

~300,000 participants from the UK Biobank (UKB), and identifying 50 independent loci[33].

In the present study, we report mtDNA-CN GWAS results from 465,809 individuals across the

Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) consortium[34] and

the UK Biobank (UKB)[35]. Using multiple gene prioritization and functional annotation methods,

we assign genes to loci that reach genome-wide significance. Finally, we perform gene

expression analyses and gene-set enrichment through PHEWAS-based SNP-clustering to

identify functional domains related to mtDNA-CN.

Subjects and Methods

Study Populations

470,579 individuals participated in this GWAS, 465,809 of whom self-identified as White.

Participants were derived from 7 population-based cohorts representing the Cohorts for Heart

and Aging Research in Genetic Epidemiology (CHARGE) consortium (Avon Longitudinal Study

of Parents and Children [ALSPAC], Atherosclerosis Risk in Communities [ARIC], Cardiovascular

Health Study [CHS], Multi-Ethnic Study of Atherosclerosis [MESA], Religious Orders Study and

Memory and Aging Project [ROSMAP], Study of Health in Pomerania [SHIP]) and from the UK

Biobank (UKB) (Supplemental Table 1). Detailed descriptions of each participating cohort, their

quality control practices, study level analyses, and ethic statements are available in the

Supplemental Methods. All study participants provided written informed consent and all centers

obtained approval from their institutional review boards.

Methods for Mitochondrial DNA Copy Number Estimation (CHARGE cohorts)

qPCR

mtDNA-CN was determined using a quantitative PCR assay as previously described[32,36]. Briefly,

the cycle threshold (Ct) value of a nuclear-specific and mitochondrial-specific probe were

measured in triplicate for each sample. In CHS, a multiplex assay using the mitochondrial *ND1*

probe and nuclear *RPPH1* probe was used, whereas ALSPAC used a mitochondrial probe

targeting the D-Loop and a nuclear probe targeting *B2M*. In CHS, we observed plate effects, as

well as a linear increase in ΔCt due to the pipetting order of each replicate. These effects were

corrected in the analysis using linear mixed model regression, with pipetting order included as a

fixed effect and plate as a random effect to create a raw measure of mtDNA-CN. In ALSPAC,

run-to-run variability was controlled using 3 calibrator samples added to every plate, to allow for

adjustment by a per-plate calibration factor[32].

Microarray

Microarray probe intensities were used to estimate mtDNA-CN using the Genvisis software

package[37] as previously described[10,36]. Briefly, Genvisis uses the median mitochondrial probe

intensity across all homozygous mitochondrial SNPs as an initial estimate of mtDNA-CN.

Technical artifacts such as DNA input quality, DNA input quantity, and hybridization efficiency

were captured through either surrogate variable (SV) or principal component (PC) analyses.

SVs or PCs were adjusted for through stepwise linear regression by adding successive

components until each successive surrogate variable or principal component no longer

significantly improved the model.

Whole Genome Sequencing (ARIC)

Whole genome sequencing read counts were used to estimate mtDNA-CN as previously

described[36]. Briefly, the total number of reads in a sample were web scraped from the NCBI

sequence read archive. Mitochondrial reads were downloaded directly from dbGaP through

Samtools (1.3.1). There was no overlap between ARIC microarray and ARIC whole-genome

sequencing samples. A ratio of mitochondrial reads to total aligned reads was used as a raw

measure of mtDNA-CN.

Adjusting for Covariates

Each method described above represents a raw measure of mtDNA-CN, adjusted for technical

artifacts; however, several potential confounding variables (e.g., age, sex, blood cell

composition) have been identified previously[18]. Raw mtDNA-CN values were adjusted for white

blood cell count in ARIC, SHIP and CHS (which also adjusted for platelet count), depending on

available data. Standardized residuals (mean = 0, standard deviation = 1) of mtDNA-CN were

used after adjusting for covariates (Supplemental Table 1.

Estimation of Mitochondrial DNA Copy Number (UKB)

Due to the availability of more detailed cell count data, as well as a different underlying

biochemistry for the Affymetrix Axiom array compared to the genotyping arrays used in the

CHARGE cohorts, mtDNA-CN in the UKB was estimated differently (Supplemental Methods).

Briefly, mtDNA-CN estimates derived from whole exome sequencing data, available on ~50,000

individuals, were generated first using customized Perl scripts to aggregate the number of

mapped sequencing reads and correct for covariates through both linear and spline regression

models. Concurrently, mitochondrial probe intensities from the Affymetrix Axiom arrays,

available on the full ~500,000 UKB cohort, were adjusted for technical artifacts through principal

components generated from nuclear probe intensities. Probe intensities were then regressed

onto the whole exome sequencing mtDNA-CN metric, and beta estimates from that regression

were used to estimate mtDNA-CN in the full UKB cohort. Finally, we used a 10-fold cross

validation method to select the cell counts to include in the final model (Supplemental Table 2).

The final UKB mtDNA-CN phenotype is the standardized residuals (mean = 0, standard

deviation = 1) from a linear model adjusting for covariates (age, sex, cell counts) as described in

the Supplemental Methods.

Genome-Wide Association Study

For each individual cohort, regression analysis was performed with residualized mtDNA-CN as

the dependent variable adjusting for age, sex, and cohort-specific covariates (e.g., principal

components, DNA collection site, family structure, cell composition). Cohorts with multiple

mtDNA-CN estimation platforms were stratified into separate analyses. Ancestry-stratified meta-

analyses were performed using Metasoft software using the Han and Eskin random effects

model to control for unobserved heterogeneity due to differences in mtDNA-CN estimation

method[38]. Effect size estimates for SNPs were calculated using a random effect meta-analysis

from cohort summary statistics, as the Han and Eskin model relaxes the assumption under the

null hypothesis without modifying the effect size estimates that occur under the alternative

hypothesis[38]. In total, three complementary analyses were performed in self-identified White

individuals, (1) a meta-analysis using all available studies, (2) a meta-analysis of studies with

available data for cell count adjustments, and (3) an analysis of UKB-only data. As the vast

majority of samples are derived from the UKB study, and given the difficulty in interpreting effect

size estimates from a random effects model, further downstream analyses were all performed

using effect size estimates from UKB-only data.

Identification of Independent GWAS Loci

To identify the initial genome-wide significant (lead) SNPs in each locus, the most significant

SNP that passed genome-wide significance ($p < 5 \times 10^{-8}$) within a 1 Mb window was selected.

To avoid Type I error, SNPs were only retained for further analyses if there were either (a) at

least two genome-wide significant SNPs in the 1 Mb window or (b) if the lead SNP was directly

genotyped. Conditional analyses were performed in UKB, where the lead SNPs from the original

GWAS were used as additional covariates in order to identify additional independent

associations.

Fine-mapping

The susieR package was used to identify all potential causal variants for each independent

locus associated with mtDNA CN[39]. UKB imputed genotype data for unrelated White subjects

were used and variants were extracted using a 500 kb window around the lead SNP for each

locus with minor allele frequency (MAF) > 0.001. 95% credible sets (CS) of SNPs, containing a

potential causal variant within a locus, were generated. The minimum absolute correlation within

each CS is 0.5 and the scaled prior variance is 0.01. When the CS did not include the lead SNP

identified from the GWAS, some of the parameters were slightly relaxed [minimum absolute

correlation is 0.2, estimate prior variance is TRUE]. The SNP with the highest posterior inclusion

probability (PIP) within each CS was also identified (Supplemental Table 3). With a few

exceptions, final lead SNPs were selected by prioritizing initially identified SNPs unless the SNP

with the highest PIP had a PIP greater than 0.2 and was 1.75 times larger than the SNP with the

second highest PIP.

Functional Annotation and Gene Prioritization

Functional Annotation

ANNOVAR was used for functional annotation of variants identified in the fine-mapping step[40].

First, variants were converted to an ANNOVAR-ready format using the dbSNP version 150

database[41]. Then, variants were annotated with ANNOVAR using the RefSeq Gene database[42].

The annotation for each variant includes the associated gene and region (e.g., exonic, intronic,

intergenic). For intergenic variants, ANNOVAR provides flanking genes and the distance to

each gene. For exonic variants, annotations also include likely functional consequences (e.g.,

synonymous/nonsynonymous, insertion/deletion), the gene affected by the variant, and the

amino acid sequence change (Supplemental Table 4).

Co-localization Analyses

Co-localization analyses were performed using the approximate Bayes factor method in the R

package *coloc*[43]. Briefly, *coloc* utilizes eQTL data and GWAS summary statistics to evaluate

the probability that gene expression and GWAS data share a single causal SNP (colocalize).

Coloc returns multiple posterior probabilities; H0 (no causal variant), H1 (causal variant for gene

expression only), H2 (causal variant for mtDNA-CN only), H3 (two distinct causal variants), and

H4 (shared causal variant for gene expression and mtDNA-CN). In the event of high H4, we can

assume that the gene is potentially causal for the GWAS phenotype of interest. eQTL summary

statistics were obtained from the eQTLGen database[44]. Genes with significant associations with

lead SNPs were tested for co-localization using variants within a 500 kb window of the sentinel

SNP. Occasionally, some of the eQTLGen p-values for certain SNPs were identical due to R's

(ver 4.0.3) limitation in handling small numbers. To account for this, if the absolute value for a

SNP's z-score association with a gene was greater than 37.02, z-scores were rescaled so that

the largest z-score would result in a p-value of 5×10^{-300}. Additionally, a few clearly co-localized

genes did not result in high H4 PPs due to the strong effect for each phenotype of a single SNP

(Supplemental Figure 1), possibly due to differences in linkage disequilibrium (LD) between the

populations. To account for this, we summed mtDNA-CN GWAS p-values and eQTLGen p-

values for each SNP and removed the SNP with the lowest combined p-value. Co-localization

analyses were then repeated without the lowest SNP. Genes with H4 greater than 50% were

classified as genes with significant evidence of co-localization.

DEPICT

Gene prioritization was performed with Depict, an integrative tool that incorporates gene co-

regulation and GWAS data to identify the most likely causal gene at a given locus[45]. Across

GWAS SNPs which overlapped with the DEPICT database, we identified SNPs representing

119 independent loci with LD pruning defined as $p < 5$ x 10-8, r2 < 0.05 and > 500 kb from other

locus boundaries. Only genes with a nominal p-value of less than 0.05 were considered for

downstream prioritization.

Gene Assignment

To prioritize genes for each identified locus, we utilized functional annotations, eQTL co-

localization analyses, and DEPICT gene prioritization results (Supplemental Figure 2). First,

genes with missense variants within *SusieR* fine-mapped credible sets were assigned to loci. If

loci co-localized with a gene's expression with a posterior probability (PP) of greater than 0.50

and there were no other co-localized genes with a PP within 5%, the gene with the highest

posterior probability was assigned. If there was still no assigned gene, the most significant

DEPICT gene was assigned. If there was no co-localization or DEPICT evidence, the nearest

gene was assigned.

Gene Set Enrichment Analyses

Using the finalized gene list from the prioritization pipeline, GO and KEGG pathway enrichment

analyses were performed using the "goana" and "kegga" functions from the R package *limma*[46],

treating all known genes as the background universe[47]. Only one gene per locus was used for

"goana" and "kegga" gene set enrichment analysis, prioritizing genes assigned to primary

independent hits. If there were multiple assigned genes, one gene was randomly selected to

avoid biasing results through loci with multiple genes. To identify an appropriate p-value cutoff,

96 genes were randomly selected from the genome and run through the same enrichment

analysis. This permutation was repeated 1000 times to generate a null distribution of the

smallest p-values from each permutation. For cluster 5 gene set enrichment analyses,

permutation testing used 47 random genes. To ensure robustness of results, gene set

enrichment analysis was repeated 50 times with random selection of genes at loci with multiple

assigned genes. GO and KEGG terms that passed permutation cutoffs at least 40/50 times

were retained.

Gene-based Association Test

We used metaXcan, which employs gene expression prediction models to evaluate associations

between phenotypes and gene expression[48]. We obtained pre-calculated expression prediction

models and SNP covariance matrices, computed using whole blood from European ancestry

individuals in version 7 of the Genotype-Tissue expression (GTEx) database[49]. Using prediction

performance p-values of less than 0.05, a total of 6,285 genes were predicted. Of these genes,

74 passed Bonferroni correction of $p < 7.95 \times 10^{-6}$. Gene set enrichment analyses were

performed on Bonferroni-significant genes as previously described. REVIGO[50] was used on the

"medium" setting (allowed similarity = 0.7) to visualize significantly enriched GO terms.

We used a one-sided Fisher's exact test to test for enrichment of genes that have been

previously identified as causal for mtDNA depletion syndromes[51–53].

PHEWAS-based SNP Clustering

mtDNA-CN Phenome-wide Association Study (PHEWAS)

We used the PHEnome Scan ANalysis Tool (PHESANT)[54] to identify mtDNA-CN associated

quantitative traits in the UKB. Briefly, we tested for the association of mtDNA-CN with 869

quantitative traits (Supplemental Table 5), limiting analyses to 365,781 White, unrelated

individuals (used.in.pca.calculation=1), and excluding individuals with extreme cell count

measurements (see Supplementary Methods). Analyses were adjusted for age, sex, and

assessment center.

SNP Phenotype Associations

SNP genotypes were regressed on mtDNA-associated quantitative traits using linear

regression, adjusted for sex, age with a natural spline (df=2), assessment center, genotyping

array, and 40 principal components (provided as part of the UKB data download).

SNP Clustering

To identify distinct clusters of mtDNA-CN GWS SNPs based on phenotypic associations, beta

estimates from the SNP phenotype associations were first divided by the beta estimate of the

mtDNA-CN SNP-mtDNA-CN association, so that all SNP-phenotype associations are relative to

the mtDNA-CN increasing allele and scaled to the effect of the SNP on mtDNA-CN. The

adjusted beta estimates were subjected to a dimensionality reduction method, Uniform Manifold

and Approximation Projection (UMAP), as implemented in the R package _umap_[55]

(random_state=123, n_neighbors=10, n_components=2, n_epochs=5000). SNPs were

assigned to clusters using Density Based Clustering of Applications with Noise (DBSCAN) as

implemented in the R package _dbscan_[56] (minPts=3). Clusters represent groups of SNPs with

similar phenotypic associations.

To identify reproducible sub-clusters within cluster 5, we ran _umap_ 100 times, varying the initial

random state (n_neighbors=4, n_epochs=5000, min_dist=0.05, n_components=3). For each

umap run, sub-clusters were assigned using _dbscan_ as described above. Final SNP sub-cluster

assignments were determined by identifying SNPs that all mapped to the same cluster >50% of

the time (i.e., all pairwise correlations for each SNP within the cluster against all other SNPs in

the cluster was >50%). The final umap run for plotting in 2 dimensions used the following

parameters: random_state=123, n_neighbors=4, n_epochs=5000, min_dist=0.05.

Phenotype Enrichment and Permutation Testing

To test for enrichment of specific phenotypes within clusters, we compared the median mtDNA-

CN scaled phenotype beta estimates within the cluster to the median beta estimates for all

SNPs not in the cluster, with significance determined using 20,000 permutations in which cluster

assignment was permuted. For multi-test correction of the overall cluster and sub-cluster

analyses, we performed 300 permutations of the initial cluster assignment (separately for the

cluster and sub-cluster analyses), followed by the comparison of median beta estimates as

described above. We retained only the most significant result from across all phenotypes and

clusters from each of the 300 permutations, and then selected the 15[th] most significant value as

the study-wide threshold for multi-test corrected significance of $p < 0.05$.

All statistical analyses were performed using R version 4.0.3

Results

Sample Characteristics

The current study included 465,809 individuals of White (European) ancestry (53.9% female)

with an average age of 56.6 yrs (sd = 8.2 yrs) (Supplemental Table 1). Follow-up validation

analyses were performed in 4,770 Blacks (60.2% female) with an average age of 61.2 yrs (sd =

7.4 yrs). The majority of the data originated from the UKB (93%). The bulk of the DNA used for

mtDNA-CN estimation was derived from buffy coat (95.5%) while the rest was derived from

peripheral leukocytes (2.2%) or whole blood (2.3%). mtDNA-CN estimated from Affymetrix

genotyping arrays consisted of 97.9% of the data while the remainder was derived from qPCR-

(1.8%) and WGS (0.3%).

GWAS

Previous work has demonstrated that the method used to measure mtDNA-CN can impact the

strength of association[36]. To account for potential differences across studies due to the

different mtDNA-CN measurements used, as well the inclusion of blood cell counts in only a

subset of the cohorts, we took two approaches. First, we used a random effects model to

perform meta-analyses, allowing for different genetic effect size estimates across cohorts.

Second, we performed three complementary analyses in individuals who self-identified as

White: 1) meta-analysis of all available studies (n = 465,809); 2) meta-analysis of studies

with available data for cell count adjustment (n = 456,151); and 3) GWAS of UKB only (n =

440,266). 77 loci were significant in all three meta-analyses, and we identified 93 independent

loci, of which 92 were genome-wide significant in the UKB data alone (Supplemental Figure 3,

Figure 1). Given that > 90% of the samples come from the UKB study, and the challenge of

interpreting effect size estimates from a random effects model, downstream analyses all use

effect size estimates from the UKB only (Supplemental Table 6).

The most significant SNP associated with mtDNA-CN was a missense mutation in

LONP1 (p = 3.00×10^{-141}), a gene that encodes a mitochondrial protease that can directly bind

mtDNA, and has been shown to regulate *TFAM*, a transcription factor involved in mtDNA

replication and repair (for review see Gibellini *et al.*)[57].

Fine-mapping and Secondary Hits

To identify additional independent SNPs within novel loci whose effects were masked by the

original significant SNP, as well as identify additional loci, we took two approaches. First, a

conditional analysis adjusting for the top 93 SNPs from the initial (primary) GWAS run revealed

3 novel loci and 19 additional independent significant SNPs within existing loci. We also

performed fine-mapping with SuSiE[39] and discovered an additional 14 independent SNPs within

existing loci. The majority of loci had only one 95% credible set of SNPs; further, twenty of the

credible sets contained only one SNP. However, many of the credible sets contained greater

than 50 SNPs after fine-mapping, and 12 of the 122 credible sets had a missense SNP as the

SNP with the highest PIP in the set. Using these two methods, we identified in total 129

independent SNPs across 96 loci (Supplemental Figure 4).

Associations in African Ancestry (AA) Populations

Examining the 129 SNPs from the Whites-only analysis, 99 were available in the Blacks-

only meta-analysis (n = 4770). After multiple testing correction, one of these SNPs was

significant (rs73349121, p = 0.0001), 9 were nominally significant (p < 0.05, with 5 expected),

and 58/99 had a direction of effect that was consistent with the White-only analyses (one-sided

p = 0.04, Figure 2). Despite being under-powered, these results in the Black-only analyses

provide evidence for similar genetic effects in a different ancestry.

Gene Prioritization and Enrichment of mtDNA Depletion Syndrome Genes

We integrated results from three different gene prioritization and functional annotation methods

(ANNOVAR[40], COLOC[43] , and DEPICT[45]) so that loci with nonsynonymous variants in gene

exons were prioritized first, with eQTL co-localization results considered second (Supplemental

Table 7), and those from DEPICT (Supplemental Table 8) were considered last (Supplemental

Figure 2). For 20 loci, multiple genes were assigned as analyses could not identify a single

priority gene (Supplemental Table 9). We noted the identification of a number of mtDNA

depletion syndrome genes in the priority list and tested for enrichment of these known causal

genes using a one-sided Fisher's exact test. For this analysis, all genes for loci assigned to

multiple genes were used, and genes for all primary and secondary loci were considered. Our

gene prioritization approach identified 7 of 16 mtDNA depletion genes (Supplemental Table 10),

consistent with a highly significant enrichment (one-sided $p = 3.09 \times 10^{-15}$).

Gene Set Enrichment Analyses

To avoid bias from a single locus with multiple functionally related genes contributing to a false-

positive signal, only one gene per unique locus was used, prioritizing genes assigned to primary

loci. One gene was randomly selected for loci with multiple assigned genes. To test for

robustness of gene set enrichment results, random selection was repeated 50 times, and only

gene sets that were significantly enriched for at least 40 iterations were retained. In all, a total of

96 genes were utilized for GO term and KEGG pathway enrichment analyses. Using a

Bonferroni-corrected p-value cutoff, 15 gene sets were significantly enriched for all 50 iterations,

including mitochondrial DNA metabolic process, mitochondrial DNA replication, coagulation,

hemostasis, amyloid-beta clearance, and mitochondrial genome maintenance (Supplemental

Table 11). No KEGG terms were significant across multiple iterations.

MetaXcan Gene Expression Analysis

As a complementary approach to single-SNP analyses, we explored the associations between

mtDNA-CN and predicted gene expression using MetaXcan[48] MetaXcan incorporates multiple

SNPs within a locus along with a reference eQTL dataset to generate predicted gene

expression levels. As our study estimated mtDNA-CN derived from blood, we used whole blood

gene expression eQTLs from the Gene-Tissue Expression (GTEx) consortium[58] to predict gene

expression in the UKB dataset. We identified 6,285 genes that had a predicted performance p-

value of less than 0.05 (i.e., they had sufficient data to generate robust gene expression levels)

and were tested for association with mtDNA-CN. Of these genes, 74 were significantly

associated with mtDNA-CN ($p < 7.95 \times 10^{-6}$, Figure 3), including 8 that were not identified

through single-SNP analyses. Many of the significant genes have known mitochondrial

functions, notably the mtDNA transcription factor *TFAM* ($p = 1.09 \times 10^{-29}$) and mitochondrial

exonuclease *MGME1* ($p = 5.87 \times 10^{-23}$), genes known as causal for mtDNA depletion

syndromes[51,52]. Additionally, *LONP1*, *MRPL43*, and *BAK1*, are all genes with known

mitochondrial functions[59–61]. Bonferroni significant MetaXcan genes were used for gene

enrichment analysis, finding enrichment for "nucleobase metabolic process" (p = 1.47 x 10^{-4})

and "mitochondrial fusion" (p = 1.86 x 10^{-4}, Supplemental Figure 5).

PHEWAS-based SNP Clustering and Gene Set Enrichment

mtDNA-CN is associated with numerous quantitative and qualitative phenotypes, many of which

are relevant to aging-related disease[3–5,9,10,13–16]. We hypothesized that this pleiotropy may

reflect different underlying functional domains captured by mtDNA-CN that may be reflected in

GWAS-identified SNPs and their likely causal genes. To test this hypothesis, we used the UKB

data to identify quantitative traits associated with mtDNA-CN and selected 42 highly significant,

non-redundant traits to test for association with the mtDNA-CN GWAS SNPs (Supplemental

Table 5, in PHEWAS = 1). We clustered SNPs using the trait effect size (beta) divided by the

mtDNA-CN effect size estimate, so that all effects are standardized to the effect of the mtDNA-

CN increasing allele for each locus. We identified 5 clusters of SNPs (Figure 4A), with clusters

1, 2, and 3 containing SNPs in which the mtDNA-CN increasing allele is associated with

decreased platelet count (PLT) (Figure 4B), increased mean platelet volume (MPV) (Figure 4C),

and platelet distribution width (PDW) (Figure 4D), consistent with a role in platelet activation[62].

Cluster 4 is most strongly enriched for SNPs in which the mtDNA-CN increasing allele is

associated with increased PLT, plateletcrit (PLTCRIT, a measure of total platelet mass), serum

calcium (Figure 4E), serum phosphate, as well as decreased mean corpuscular volume (MCV)

and mean spherical cellular volume (Figure 4F) (Supplemental Table 12). The cluster 4

phenotypes, and supported by the genes identified for this cluster, implicate megakaryocyte

proliferation and proplatelet formation (*MYB*, *AK3*, *JAK2*)[63], and apoptosis and autophagy

(*BAK1*, *BCL2, TYMP*)[64]. Cluster 5 did not yield any specific trait enrichment (all significant

results reflected the strong enrichment observed in clusters 1-4); however, gene set enrichment

for this cluster identified multiple mtDNA-related gene ontology terms, including mitochondrial

genome maintenance, regulation of phospholipid efflux, and amyloid-beta clearance

(Supplemental Table 13). Cluster 5 contains a highly diverse set of SNP-trait associations. A

secondary clustering only with SNPs mapped to this cluster identified 9 sub-clusters

(Supplemental Figure 6A), 3 having clear evidence for enrichment of specific phenotypes.

Cluster 6 is enriched for decreased MCV (Supplemental Figure 6B). Cluster 7 is enriched for

decreased MPV (Supplemental Figure 6C), PDW, serum phosphate, serum calcium, and

PLTCRT. Cluster 10 is enriched for decreased apoA, apoB, total cholesterol (Supplemental

Figure 6D), LDL, and HDL, and increased vitamin D, pulse rate, and direct bilirubin. SNPs that

were not assigned to a cluster were enriched for decreased fasting blood glucose and maximum

workload from fitness testing (Supplemental Table 14).

Discussion

We conducted a GWAS for mtDNA-CN using 465,809 individuals from the CHARGE consortium

and the UKB. In addition to replicating two previously reported loci, we discovered 94 novel loci

and report multiple independent hits for 26 loci. Examining our GWS SNPs in a Black

population, we observed a concordant signal, suggesting that the genetic etiology of mtDNA-CN

may be broadly similar across populations. Using several functional follow-up methods, genes

were assigned for each identified independent hit and significant enrichment was observed for

genes involved in mitochondrial DNA metabolism, homeostasis, cell activation, and amyloid-

beta clearance. In total, we assigned 124 unique genes to independent GWAS signals

associated with mtDNA-CN. We also identified 8 additional genes whose predicted gene

expression is associated with mtDNA-CN that could not be mapped back to GWS loci. Finally,

using a clustering approach based on SNP associations with various mtDNA-CN associated

phenotypes, we were able to functionally categorize SNPs, providing insight into biological

pathways that impact mtDNA-CN. We note that during the preparation of this manuscript, a

GWAS including 295,150 unrelated individuals from the UK Biobank was published, which

reported 50 genome-wide significant regions[33]. While many of our loci overlap with their

findings, our study reports twice as many loci largely due to the increased power of our study.

We were able to identify a substantial proportion of the genes involved in mtDNA

depletion syndromes (7/16, p = 3.09 x 10^{-15} for enrichment), including *TWNK, TFAM, DGUOK,*

MGME1, RRM2B, TYMP, and *POLG*. mtDNA depletion syndromes can be broken down into 5

subtypes based on their constellation of phenotypes[65] , and with the exception of

cardiomyopathic subtypes (associated with mutations in *AGK* and *SLC25A4*), we were able to

identify at least 1 gene from the other 4 subtypes, suggesting that our mtDNA-CN measurement

in blood-derived DNA can identify genes widely relevant to non-blood phenotypes. This finding

is consistent with a large body of work showing that mtDNA-CN measured in blood is

associated with numerous aging-related phenotypes for which the primary tissue of interest is

not blood (e.g. chronic kidney disease[13], heart failure[11], and diabetes[66]). Also consistent with this

finding is recent work demonstrating that mtDNA-CN measured in blood is associated with

mtRNA expression across numerous non-blood tissues, suggesting a link between

mitochondrial function measured in blood and other tissues[67].

In addition to identifying the mtDNA depletion syndrome genes directly linked to

mitochondrial DNA metabolic processes, DNA replication, and genome maintenance, we also

identify genes which play a role in mitochondrial function. The top GWAS hit is a missense

mutation in *LONP1*, which encodes a mitochondrial protease that has been shown to cause

mitochondrial cytopathy and reduced respiratory chain activity[68,69]. Interestingly, this missense

mutation was recently found to be associated with mitochondrial tRNA methylation levels[70].

Additional genes known to impact mitochondrial function include *MFN1*, which encodes a

mediator of mitochondrial fusion[71,72], *STMP1*, which plays a role in mitochondrial respiration[73],

and *MRPS35*, which encodes a ribosomal protein involved in protein synthesis in the

mitochondrion[74,75].

Using a combination of gene-based tests and gene prioritization using functional

annotation, pathway analyses reveal enrichment for numerous mitochondrial related pathways,

as well as those involved in regulation of cell activation ($p < 3.65 \times 10^{-5}$), homeostatic processes

($p < 1.82 \times 10^{-5}$), and regulation of immune system processes ($p < 2.75 \times 10^{-5}$) (Supplemental

Table 11). These results provide additional evidence for the broad role played by mitochondria

in numerous aspects of cellular function. Of particular interest, the GO term for amyloid beta is

significantly enriched, reinforcing a link between mtDNA-CN and neurodegenerative disease[76–

78]. Previous work from our lab using the UKB has shown that increased mtDNA-CN is

associated with lower rates of prevalent neurodegenerative disease, as well as predictive for

decreased risk of incident neurodegenerative disease[67]. mtDNA-CN is also known to be

decreased in the frontal cortex of Alzheimer's disease (AD) patients[79]. Interestingly, the four

GWAS-identified genes driving the enrichment for amyloid-beta clearance are all related to

regulation of lipid levels, and lipid homeostasis within the brain is known to play an important

role in Alzheimer's disease[80]. *APOE*, one of the most well-known risk genes for Alzheimer's

disease, is a cholesterol carrier involved in lipid transport, and the ApoE-ε4 isoform involved in

AD pathogenesis is associated with mitochondrial dysfunction and oxidative distress in the

human brain[81]; *CD36* is a platelet glycoprotein which mediates the response to amyloid-beta

accumulation[82]; *LDLR* is a low-density lipoprotein receptor associated with AD[83]; and *ABCA7* is

a phospholipid transporter[84]. *ABCA7* loss of function variants are enriched in both AD and

Parkinson's disease (PD) patients[85], suggesting a broad role across neurodegenerative

diseases.

Given the integral role of mitochondria in cellular function, not just with ATP

formation/energy production, but signaling through ROS, and its key role in apoptosis, there is a

strong reason to *a priori* assume that genetic variants associated with mtDNA-CN are likely to

be highly pleiotropic. Indeed, mtDNA-CN itself is associated with numerous phenotypes

(Supplemental Table 5). Through our PHEWAS-based clustering approach using 42 mtDNA-CN

associated phenotypes, we uncovered phenotypic associations between five distinct clusters of

GWS mtDNA-CN associated SNPs. Cluster 1-3 were characterized by increased MPV, PDW,

and decreased PLT (note that measured MPV and PLT are generally inversely correlated to

maintain hemostasis), which are the hallmarks of platelet activation[62]. The link between platelets

and mtDNA-CN has typically revolved around platelet count, as platelets have functional

mitochondria, but do not have a nucleus. Given that the mtDNA-CN measurement is the ratio

between mtDNA and nuclear DNA, increased platelets, all else being equal, would directly

equate with increased mtDNA-CN. We note that the mtDNA-CN metric used in this GWAS was

adjusted for platelet count, likely increasing the ability to detect variants that impact mtDNA-CN

through increased platelet activation. Examining the genes within this cluster suggests role for

actin formation/regulation (*CARMIL1*, *TPM4*, *PACSIN2*)[86–88] and vesicular transport/endocytic

trafficking (*DNM3*, *EHD3*)[89,90] in platelet activation.

Cluster 4 is most strongly enriched for SNPs in which the mtDNA-CN increasing allele is

associated with increased PLT/PLTCRIT and serum calcium/phosphate. Examining the genes

assigned to the cluster, we implicate megakaryocyte proliferation and proplatelet formation

(*MYB*, *AK3*, *JAK2*)[63], and apoptosis and autophagy (*BAK1*, *BCL2, TYMP*)[64]. Megakaryocytes

are used to form proplatelets, and the process includes an important role for both intra- and

extracellular calcium levels[91]. A role for apoptosis, and specifically *BCL2*, in proplatelet

formation and platelet release has been suggested[92,93], however work in mice has suggested

that apoptosis does not play a direct role in these processes[94]. Nevertheless, apoptosis is

important for platelet lifespan[95].

Cluster 5 was particularly challenging to interpret, given that no particular phenotype was

enriched relative to the non-cluster 5 SNPs. We note that this cluster appeared to be enriched

for the mtDNA depletion syndrome genes, containing 6/7 genes identified in the GWAS, and

significantly enriched for GO Terms related mitochondrial DNA. Examining sub-clusters within

cluster 5 identified MCV as an important phenotype (while only sub-cluster 6 was formally

associated, looking at Figure 3B suggests more widespread clustering based on MCV). MCV is

a measure of the average volume of a red blood corpuscle, and the link between red blood

volume and mtDNA-CN is unlikely to be direct, given that red blood cells contain neither nuclei

nor mitochondria. Sub-cluster 7 is surprisingly enriched for variants associated with decreased

MPV/PDW and serum phosphate/calcium, inverse of what we observe for cluster 4. Finally, sub-

cluster 10 is strongly associated with lipids, and includes a number of genes known to be

associated with lipid levels (*LIPC*, *CETP*, *LDLR*, *APOE*). While lipids play a role in both energy

metabolism (largely through fatty acids) and cellular membrane formation, a link to mtDNA-CN

and/or mitochondrial function is not well-established. One potentially interesting result is

provided by Olkowicz and colleagues, who demonstrated that ApoE$^{-/-}$/LDLR$^{-/-}$ mice had

increased cardiac mitochondrial oxidative metabolism, with proteomic analysis suggesting

increased mitochondrial abundance in mouse hearts[96]. However, we note that our results show

an association between decreased lipids and increased mtDNA-CN, rather than the reverse,

shown in the Olkowicz study.

Although the question of causality is of great interest, this study was unable to determine

the directionality of effect between mtDNA-CN and phenotypes of interest, as we are

underpowered for Mendelian randomization (MR), with less than 1% of the variance in mtDNA-

CN explained by GWS loci when predicted into ARIC. As an additional limitation, we note that

despite the large sample size and numerous loci identified, we are likely missing a great deal of

the true signal, as previous studies have estimated mtDNA-CN heritability to be 65%[23]. Finally,

while we have adjusted our mtDNA-CN metric for a variety of confounders, it is important to

note that mtDNA-CN can be influenced by a variety of environmental factors including

smoking[97] and drugs, which have not been adjusted for in these analyses.

In summary, we performed the largest-to-date GWAS for mtDNA-CN, including almost

500,000 individuals. We identified distinct groups of SNPs associated with mtDNA-CN that are

related to platelet activation, megakaryocyte formation and apoptotic processes, and showed

clear enrichment for genes involved in mtDNA depletion and nucleotide regulation. Given the

role of mtDNA-CN in aging-related disease, this work begins to unravel the many varied

underlying mechanisms for genetic control of mtDNA-CN.

Supplemental Data

Supplemental Data include six supplemental figures, fourteen supplemental tables, and

supplemental methods.

Declaration of Interests

Psaty serves on the Steering Committee of the Yale Open Data Access Project funded by

Johnson & Johnson. All other authors declare no competing interests.

Acknowledgements

This work was supported by National Heart, Lung and Blood Institute, National Institutes of

Health (NIH) grants R01HL13573 and R01HL144569 (RJL, SYY, CAC, DEA), NIH grant P01-

AG027734 (GA, YK, NB, AB) and the National Center for Advancing Translational Sciences,

NIH, through Grant KL2TR002490 (LMR). The content is solely the responsibility of the authors

and does not necessarily represent the official views of the NIH. LMR was also funded by T32

HL129982.

This research was conducted using data from the Genotype-Tissue Expression (GTEx) project

(dbGaP accession: phs000424.**v8**.p2). The GTEx project was supported by the Common

Fund of the Office of the Director of the National Institutes of Health, and by NCI, NHGRI,

NHLBI, NIDA, NIMH, and NINDS.

This research was also conducted using the UK Biobank Resource under Application Number

17731.

The Atherosclerosis Risk in Communities study has been funded in whole or in part with Federal

funds from the National Heart, Lung, and Blood Institute, National Institutes of Health,

Department of Health and Human Services (contract numbers HSN268201700001I,

HHSN268201700002I, HHSN268201700003I, HHSN268201700004I and

HHSN268201700005I), R01HL087641, R01HL059367 and R01HL086694; National Human

Genome Research Institute contract U01HG004402; and National Institutes of Health contract

HHSN268200625226C. Funding support for "Building on GWAS for NHLBI diseases: the U.S.

CHARGE consortium" was provided by the NIH through the American Recovery and

Reinvestment Act of 2009 (ARRA) (5RC2HL102419). Sequencing was carried out at the Baylor

College of Medicine Human Genome Sequencing Center and supported by the National Human

Genome Research Institute grants U54 HG003273 and UM1 HG008898. The authors thank the

staff and participants of the ARIC study for their important contributions. Infrastructure was

partly supported by Grant Number UL1RR025005, a component of the National Institutes of

Health and NIH Roadmap for Medical Research.

This CHS research was supported by NHLBI contracts HHSN268201200036C,

HHSN268200800007C, HHSN268201800001C, N01HC55222, N01HC85079, N01HC85080,

N01HC85081, N01HC85082, N01HC85083, N01HC85086, 75N92021D00006; and NHLBI

grants U01HL080295, R01HL087652, R01HL105756, R01HL103612, R01HL120393, and

U01HL130114 with additional contribution from the National Institute of Neurological Disorders

and Stroke (NINDS). Additional support was provided through R01AG023629 from the National

Institute on Aging (NIA). A full list of principal CHS investigators and institutions can be found

at CHS-NHLBI.org. The provision of genotyping data was supported in part by the National

Center for Advancing Translational Sciences, CTSI grant UL1TR001881, and the National

Institute of Diabetes and Digestive and Kidney Disease Diabetes Research Center (DRC) grant

DK063491 to the Southern California Diabetes Endocrinology Research Center.

The FHS phenotype-genotype analyses were supported by the National Institute of Aging

(U34AG051418). This research was conducted in part using data and resources from the

Framingham Heart Study of the National Heart Lung and Blood Institute of the National

Institutes of Health and Boston University School of Medicine. This work was partially supported

by the National Heart, Lung and Blood Institute's Framingham Heart Study (Contract No. N01-

HC-25195, HHSN268201500001) and its contract with Affymetrix, Inc for genotyping services

(Contract No. N02-HL-6-4278). Genotyping, quality control and calling of the Illumina

HumanExome BeadChip in the Framingham Heart Study was supported by funding from the

National Heart, Lung and Blood Institute Division of Intramural Research (Daniel Levy and

Christopher J. O'Donnell, Principle Investigators). The authors thank the participants for their

dedication to the study. The authors are pleased to acknowledge that the computational work

reported on in this paper was performed on the Shared Computing Cluster which is

administered by Boston University's Research Computing Services. URL:

www.bu.edu/tech/support/research/.

MESA and the MESA SHARe projects are conducted and supported by the National Heart,

Lung, and Blood Institute (NHLBI) in collaboration with MESA investigators. Support for MESA

is provided by contracts 75N92020D00001, HHSN268201500003I, N01-HC-95159,

75N92020D00005, N01-HC-95160, 75N92020D00002, N01-HC-95161, 75N92020D00003,

N01-HC-95162, 75N92020D00006, N01-HC-95163, 75N92020D00004, N01-HC-95164,

75N92020D00007, N01-HC-95165, N01-HC-95166, N01-HC-95167, N01-HC-95168, N01-HC-

95169, UL1-TR-000040, UL1-TR-001079, and UL1-TR-001420. Funding for SHARe

genotyping was provided by NHLBI Contract N02-HL-64278. Genotyping was performed at

Affymetrix (Santa Clara, California, USA) and the Broad Institute of Harvard and MIT (Boston,

Massachusetts, USA) using the Affymetrix Genome-Wide Human SNP Array 6.0. The provision

of genotyping data was supported in part by the National Center for Advancing Translational

Sciences, CTSI grant UL1TR001881, and the National Institute of Diabetes and Digestive and

Kidney Disease Diabetes Research Center (DRC) grant DK063491 to the Southern California

Diabetes Endocrinology Research Center.

ROS/MAP is supported by the Translational Genomics Research Institute and National Institute

on Aging (NIA) through grants U01AG46152, U01AG61256, P30AG10161, R01AG17917,

RF1AG15819, R01AG30146.

SHIP and SHIP-TREND are part of the Community Medicine Research net of the University of

Greifswald, Germany, which is funded by the Federal Ministry of Education and Research (grants

no. 01ZZ9603, 01ZZ0103, and 01ZZ0403), the Ministry of Cultural Affairs as well as the Social

Ministry of the Federal State of Mecklenburg-West Pomerania, and the network 'Greifswald

Approach to Individualized Medicine (GANI_MED)' funded by the Federal Ministry of Education

and Research (grant 03IS2061A).

This research was funded in whole, or in part, by the Wellcome Trust [Grant ref: 217065/Z/19/Z].

For the purpose of Open Access, the author has applied a CC BY public copyright license to any

Author Accepted Manuscript version arising from this submission.

Web Resources

REVIGO was accessed at http://revigo.irb.hr/.

Data and Code Availability

All data used in this manuscript is available through either the UKBiobank and CHARGE

consortiums. Code and scripts are available in a zipped file at

https://www.arkinglab.org/resources/.

References

1. Wallace, D.C. (1992). Diseases of the Mitochondrial Dna. Annual Review of Biochemistry *61*,
1175–1212.

2. Vakifahmetoglu-Norberg, H., Ouchida, A.T., and Norberg, E. (2017). The role of mitochondria
in metabolism and cell death. Biochem. Biophys. Res. Commun. *482*, 426–431.

3. Herst, P.M., Rowe, M.R., Carson, G.M., and Berridge, M.V. (2017). Functional Mitochondria
in Health and Disease. Front Endocrinol (Lausanne) *8*,.

4. Dai, D.-F., Rabinovitch, P.S., and Ungvari, Z. (2012). Mitochondria and cardiovascular aging.
Circ. Res. *110*, 1109–1124.

5. Cui, H., Kong, Y., and Zhang, H. (2012). Oxidative stress, mitochondrial dysfunction, and
aging. J Signal Transduct *2012*, 646354.

6. Liu, C.-S., Tsai, C.-S., Kuo, C.-L., Chen, H.-W., Lii, C.-K., Ma, Y.-S., and Wei, Y.-H. (2003).
Oxidative stress-related alteration of the copy number of mitochondrial DNA in human
leukocytes. Free Radic Res *37*, 1307–1317.

7. Guha, M., and Avadhani, N.G. (2013). Mitochondrial retrograde signaling at the crossroads of
tumor bioenergetics, genetics and epigenetics. Mitochondrion *13*, 577–591.

8. Malik, A.N., and Czajka, A. (2013). Is mitochondrial DNA content a potential biomarker of
mitochondrial dysfunction? Mitochondrion *13*, 481–492.

9. Ashar, F.N., Moes, A., Moore, A.Z., Grove, M.L., Chaves, P.H.M., Coresh, J., Newman, A.B.,
Matteini, A.M., Bandeen-Roche, K., Boerwinkle, E., et al. (2015). Association of mitochondrial
DNA levels with frailty and all-cause mortality. J. Mol. Med. *93*, 177–186.

10. Ashar, F.N., Zhang, Y., Longchamps, R.J., Lane, J., Moes, A., Grove, M.L., Mychaleckyj,
J.C., Taylor, K.D., Coresh, J., Rotter, J.I., et al. (2017). Association of Mitochondrial DNA Copy
Number With Cardiovascular Disease. JAMA Cardiol *2*, 1247–1255.

11. Hong, Y.S., Longchamps, R.J., Zhao, D., Castellani, C.A., Loehr, L.R., Chang, P.P.,
Matsushita, K., Grove, M.L., Boerwinkle, E., Arking, D.E., et al. (2020). Mitochondrial DNA Copy
Number and Incident Heart Failure: The Atherosclerosis Risk in Communities (ARIC) Study.
Circulation *141*, 1823–1825.

12. Zhao, D., Bartz, T.M., Sotoodehnia, N., Post, W.S., Heckbert, S.R., Alonso, A.,
Longchamps, R.J., Castellani, C.A., Hong, Y.S., Rotter, J.I., et al. (2020). Mitochondrial DNA
copy number and incident atrial fibrillation. BMC Medicine *18*, 246.

13. Tin, A., Grams, M.E., Ashar, F.N., Lane, J.A., Rosenberg, A.Z., Grove, M.L., Boerwinkle, E.,
Selvin, E., Coresh, J., Pankratz, N., et al. (2016). Association between Mitochondrial DNA Copy
Number in Peripheral Blood and Incident CKD in the Atherosclerosis Risk in Communities
Study. J Am Soc Nephrol *27*, 2467–2473.

14. Pyle, A., Anugrha, H., Kurzawa-Akanbi, M., Yarnall, A., Burn, D., and Hudson, G. (2016).
Reduced mitochondrial DNA copy number is a biomarker of Parkinson's disease. Neurobiol.
Aging *38*, 216.e7-216.e10.

15. Wei, W., Keogh, M.J., Wilson, I., Coxhead, J., Ryan, S., Rollinson, S., Griffin, H., Kurzawa-
Akanbi, M., Santibanez-Koref, M., Talbot, K., et al. (2017). Mitochondrial DNA point mutations
and relative copy number in 1363 disease and control human brains. Acta Neuropathologica
Communications *5*, 13.

16. Reznik, E., Miller, M.L., Şenbabaoğlu, Y., Riaz, N., Sarungbam, J., Tickoo, S.K., Al-
Ahmadie, H.A., Lee, W., Seshan, V.E., Hakimi, A.A., et al. (2016). Mitochondrial DNA copy
number variation across human cancers. Elife *5*,.

17. Hurtado-Roca, Y., Ledesma, M., Gonzalez-Lazaro, M., Moreno-Loshuertos, R., Fernandez-
Silva, P., Enriquez, J.A., and Laclaustra, M. (2016). Adjusting MtDNA Quantification in Whole
Blood for Peripheral Blood Platelet and Leukocyte Counts. PLOS ONE *11*, e0163770.

18. Knez, J., Winckelmans, E., Plusquin, M., Thijs, L., Cauwenberghs, N., Gu, Y., Staessen,
J.A., Nawrot, T.S., and Kuznetsova, T. (2016). Correlates of Peripheral Blood Mitochondrial
DNA Content in a General Population. Am J Epidemiol *183*, 138–146.

19. Kumar, P., Efstathopoulos, P., Millischer, V., Olsson, E., Wei, Y.B., Brüstle, O., Schalling,
M., Villaescusa, J.C., Ösby, U., and Lavebratt, C. (2018). Mitochondrial DNA copy number is
associated with psychosis severity and anti-psychotic treatment. Sci Rep *8*, 12743.

20. Urata, M., Koga-Wada, Y., Kayamori, Y., and Kang, D. (2008). Platelet contamination
causes large variation as well as overestimation of mitochondrial DNA content of peripheral
blood mononuclear cells. Ann Clin Biochem *45*, 513–514.

21. Clay Montier, L.L., Deng, J.J., and Bai, Y. (2009). Number matters: control of mammalian
mitochondrial DNA copy number. J Genet Genomics *36*, 125–131.

22. Tang, Y., Schon, E.A., Wilichowski, E., Vazquez-Memije, M.E., Davidson, E., and King, M.P.
(2000). Rearrangements of Human Mitochondrial DNA (mtDNA): New Insights into the
Regulation of mtDNA Copy Number and Gene Expression. Mol Biol Cell *11*, 1471–1485.

23. Xing, J., Chen, M., Wood, C.G., Lin, J., Spitz, M.R., Ma, J., Amos, C.I., Shields, P.G.,
Benowitz, N.L., Gu, J., et al. (2008). Mitochondrial DNA content: its genetic heritability and
association with renal cell carcinoma. J. Natl. Cancer Inst. *100*, 1104–1112.

24. Carling, P.J., Cree, L.M., and Chinnery, P.F. (2011). The implications of mitochondrial DNA
copy number regulation during embryogenesis. Mitochondrion *11*, 686–692.

25. Harvey, A., Gibson, T., Lonergan, T., and Brenner, C. (2011). Dynamic regulation of
mitochondrial function in preimplantation embryos and embryonic stem cells. Mitochondrion *11*,
829–838.

26. Copeland, W.C. (2014). Defects of Mitochondrial DNA Replication. J Child Neurol *29*, 1216–
1224.

27. Mandel, H., Szargel, R., Labay, V., Elpeleg, O., Saada, A., Shalata, A., Anbinder, Y.,
Berkowitz, D., Hartman, C., Barak, M., et al. (2001). The deoxyguanosine kinase gene is
mutated in individuals with depleted hepatocerebral mitochondrial DNA. Nat. Genet. *29*, 337–
341.

28. Wang, L., Limongelli, A., Vila, M.R., Carrara, F., Zeviani, M., and Eriksson, S. (2005).
Molecular insight into mitochondrial DNA depletion syndrome in two patients with novel
mutations in the deoxyguanosine kinase and thymidine kinase 2 genes. Mol. Genet. Metab. *84*,
75–82.

29. Rusecka, J., Kaliszewska, M., Bartnik, E., and Tońska, K. (2018). Nuclear genes involved in
mitochondrial diseases caused by instability of mitochondrial DNA. J Appl Genet *59*, 43–57.

30. Cai, N., Li, Y., Chang, S., Liang, J., Lin, C., Zhang, X., Liang, L., Hu, J., Chan, W., Kendler,
K.S., et al. (2015). Genetic Control over mtDNA and Its Relationship to Major Depressive
Disorder. Curr Biol *25*, 3170–3177.

31. Workalemahu, T., Enquobahrie, D.A., Tadesse, M.G., Hevner, K., Gelaye, B., Sanchez, S.,
and Williams, M.A. (2017). Genetic Variations Related to Maternal Whole Blood Mitochondrial
DNA Copy Number: A Genome-Wide and Candidate Gene Study. J Matern Fetal Neonatal Med
30, 2433–2439.

32. Guyatt, A.L., Brennan, R.R., Burrows, K., Guthrie, P.A.I., Ascione, R., Ring, S.M., Gaunt,
T.R., Pyle, A., Cordell, H.J., Lawlor, D.A., et al. (2019). A genome-wide association study of
mitochondrial DNA copy number in two population-based cohorts. Human Genomics *13*, 6.

33. Hägg, S., Jylhävä, J., Wang, Y., Czene, K., and Grassmann, F. (2020). Deciphering the
genetic and epidemiological landscape of mitochondrial DNA abundance. Hum Genet.

34. Longchamps, R.J., Castellani, C.A., Yang, S.Y., Newcomb, C.E., Sumpter, J.A., Lane, J.,
Grove, M.L., Guallar, E., Pankratz, N., Taylor, K.D., et al. (2020). Evaluation of mitochondrial
DNA copy number estimation techniques. PLOS ONE *15*, e0228166.

35. MitoPipeline: Generating Mitochondrial copy number estimates from SNP array data in
Genvisis.

36. Han, B., and Eskin, E. (2011). Random-Effects Model Aimed at Discovering Associations in
Meta-Analysis of Genome-wide Association Studies. The American Journal of Human Genetics
88, 586–598.

37. Wang, G., Sarkar, A., Carbonetto, P., and Stephens, M. (2020). A simple new approach to
variable selection in regression, with application to genetic fine mapping. Journal of the Royal
Statistical Society: Series B (Statistical Methodology) *82*, 1273–1300.

38. Wang, K., Li, M., and Hakonarson, H. (2010). ANNOVAR: functional annotation of genetic
variants from high-throughput sequencing data. Nucleic Acids Research *38*, e164–e164.

39. Sherry, S.T., Ward, M., and Sirotkin, K. (1999). dbSNP-database for single nucleotide
polymorphisms and other classes of minor genetic variation. Genome Res *9*, 677–679.

40. O'Leary, N.A., Wright, M.W., Brister, J.R., Ciufo, S., Haddad, D., McVeigh, R., Rajput, B.,
Robbertse, B., Smith-White, B., Ako-Adjei, D., et al. (2016). Reference sequence (RefSeq)
database at NCBI: current status, taxonomic expansion, and functional annotation. Nucleic
Acids Res *44*, D733-745.

41. Giambartolomei, C., Vukcevic, D., Schadt, E.E., Franke, L., Hingorani, A.D., Wallace, C.,
and Plagnol, V. (2014). Bayesian Test for Colocalisation between Pairs of Genetic Association
Studies Using Summary Statistics. PLOS Genetics *10*, e1004383.

42. Võsa, U., Claringbould, A., Westra, H.-J., Bonder, M.J., Deelen, P., Zeng, B., Kirsten, H.,
Saha, A., Kreuzhuber, R., Kasela, S., et al. (2018). Unraveling the polygenic architecture of
complex traits using blood eQTL metaanalysis. BioRxiv 447367.

43. Pers, T.H., Karjalainen, J.M., Chan, Y., Westra, H.-J., Wood, A.R., Yang, J., Lui, J.C.,
Vedantam, S., Gustafsson, S., Esko, T., et al. (2015). Biological interpretation of genome-wide
association studies using predicted gene functions. Nature Communications *6*, 5890.

44. Smyth, G., Hu, Y., Ritchie, M., Silver, J., Wettenhall, J., McCarthy, D., Wu, D., Shi, W.,
Phipson, B., Lun, A., et al. (2021). limma: Linear Models for Microarray Data (Bioconductor
version: Release (3.12)).

45. Young, M.D., Wakefield, M.J., Smyth, G.K., and Oshlack, A. (2010). Gene ontology analysis
for RNA-seq: accounting for selection bias. Genome Biology *11*, R14.

46. Barbeira, A.N., Dickinson, S.P., Bonazzola, R., Zheng, J., Wheeler, H.E., Torres, J.M.,
Torstenson, E.S., Shah, K.P., Garcia, T., Edwards, T.L., et al. (2018). Exploring the phenotypic
consequences of tissue specific gene expression variation inferred from GWAS summary
statistics. Nat Commun *9*, 1825.

47. Barbeira, A.N., Pividori, M., Zheng, J., Wheeler, H.E., Nicolae, D.L., and Im, H.K. (2019).
Integrating predicted transcriptome from multiple tissues improves association detection. PLOS
Genetics *15*, e1007889.

48. Supek, F., Bošnjak, M., Škunca, N., and Šmuc, T. (2011). REVIGO summarizes and
visualizes long lists of gene ontology terms. PLoS One *6*, e21800.

49. Stiles, A.R., Simon, M.T., Stover, A., Eftekharian, S., Khanlou, N., Wang, H.L., Magaki, S.,
Lee, H., Partynski, K., Dorrani, N., et al. (2016). Mutations in TFAM, encoding mitochondrial
transcription factor A, cause neonatal liver failure associated with mtDNA depletion. Mol Genet
Metab *119*, 91–99.

50. Kornblum, C., Nicholls, T.J., Haack, T.B., Schöler, S., Peeva, V., Danhauser, K., Hallmann,
K., Zsurka, G., Rorbach, J., Iuso, A., et al. (2013). Loss-of-function mutations in MGME1 impair
mtDNA replication and cause multisystemic mitochondrial disease. Nat Genet *45*, 214–219.

51. El-Hattab, A.W., and Scaglia, F. (2013). Mitochondrial DNA depletion syndromes: review
and updates of genetic basis, manifestations, and therapeutic options. Neurotherapeutics *10*,
186–198.

52. Millard, L.A.C., Davies, N.M., Gaunt, T.R., Davey Smith, G., and Tilling, K. (2018). Software
Application Profile: PHESANT: a tool for performing automated phenome scans in UK Biobank.
Int J Epidemiol *47*, 29–35.

53. Konopka, T. (2020). umap: Uniform Manifold Approximation and Projection.

54. Hahsler, M., Piekenbrock, M., Arya, S., and Mount, D. (2019). dbscan: Density Based
Clustering of Applications with Noise (DBSCAN) and Related Algorithms.

55. Gibellini, L., De Gaetano, A., Mandrioli, M., Van Tongeren, E., Bortolotti, C.A., Cossarizza,
A., and Pinti, M. (2020). The biology of Lonp1: More than a mitochondrial protease. Int Rev Cell
Mol Biol *354*, 1–61.

56. GTEx Consortium (2013). The Genotype-Tissue Expression (GTEx) project. Nat Genet *45*,
580–585.

57. Liu, T., Lu, B., Lee, I., Ondrovicová, G., Kutejová, E., and Suzuki, C.K. (2004). DNA and
RNA binding by the mitochondrial lon protease is regulated by nucleotide and protein substrate.
J Biol Chem *279*, 13902–13910.

58. Sharma, M.R., Koc, E.C., Datta, P.P., Booth, T.M., Spremulli, L.L., and Agrawal, R.K.
(2003). Structure of the mammalian mitochondrial ribosome reveals an expanded functional role
for its component proteins. Cell *115*, 97–108.

59. Shimizu, S., Narita, M., and Tsujimoto, Y. (1999). Bcl-2 family proteins regulate the release
of apoptogenic cytochrome c by the mitochondrial channel VDAC. Nature *399*, 483–487.

60. Vagdatli, E., Gounari, E., Lazaridou, E., Katsibourlia, E., Tsikopoulou, F., and Labrianou, I.
(2010). Platelet distribution width: a simple, practical and specific marker of activation of
coagulation. Hippokratia *14*, 28–32.

61. PathCards :: Factors involved in megakaryocyte development and platelet production
Pathway and related pathways.

62. PathCards :: Apoptosis and Autophagy Pathway and related pathways.

63. Basel, D. (2020). Mitochondrial DNA Depletion Syndromes. Clin Perinatol *47*, 123–141.

64. DeBarmore, B., Longchamps, R.J., Zhang, Y., Kalyani, R.R., Guallar, E., Arking, D.E.,
Selvin, E., and Young, J.H. (2020). Mitochondrial DNA copy number and diabetes: the
Atherosclerosis Risk in Communities (ARIC) study. BMJ Open Diabetes Res Care *8*,.

65. Yang, S.Y., Castellani, C.A., Longchamps, R.J., Pillalamarri, V.K., O'Rourke, B., Guallar, E.,
and Arking, D.E. (2020). Blood-derived mitochondrial DNA copy number is associated with gene
expression across multiple tissues and is predictive for incident neurodegenerative disease.
BioRxiv 2020.07.17.209023.

66. Hannah-Shmouni, F., MacNeil, L., Brady, L., Nilsson, M.I., and Tarnopolsky, M. (2019).
Expanding the Clinical Spectrum of LONP1-Related Mitochondrial Cytopathy. Front Neurol *10*,
981.

67. Grainha, T.R.R., Jorge, P.A. da S., Pérez-Pérez, M., Pérez Rodríguez, G., Pereira,
M.O.B.O., and Lourenço, A.M.G. (2018). Exploring anti-quorum sensing and anti-virulence
based strategies to fight Candida albicans infections: an in silico approach. FEMS Yeast Res
18,.

68. Ali, A.T., Idaghdour, Y., and Hodgkinson, A. (2020). Analysis of mitochondrial m1A/G RNA
modification reveals links to nuclear genetic variants and associated disease processes.
Communications Biology *3*, 1–11.

69. Schrepfer, E., and Scorrano, L. (2016). Mitofusins, from Mitochondria to Metabolism. Mol
Cell *61*, 683–694.

70. Ishihara, N., Eura, Y., and Mihara, K. (2004). Mitofusin 1 and 2 play distinct roles in
mitochondrial fusion reactions via GTPase activity. J Cell Sci *117*, 6535–6546.

71. Zhang, D., Xi, Y., Coccimiglio, M.L., Mennigen, J.A., Jonz, M.G., Ekker, M., and Trudeau,
V.L. (2012). Functional prediction and physiological characterization of a novel short trans-
membrane protein 1 as a subunit of mitochondrial respiratory complexes. Physiol Genomics *44*,
1133–1140.

72. Cavdar Koc, E., Burkhart, W., Blackburn, K., Moseley, A., and Spremulli, L.L. (2001). The
small subunit of the mammalian mitochondrial ribosome. Identification of the full complement of
ribosomal proteins present. J Biol Chem *276*, 19363–19374.

73. Márquez-Jurado, S., Díaz-Colunga, J., das Neves, R.P., Martinez-Lorente, A., Almazán, F.,
Guantes, R., and Iborra, F.J. (2018). Mitochondrial levels determine variability in cell death by
modulating apoptotic gene expression. Nat Commun *9*, 389.

74. Dölle, C., Flønes, I., Nido, G.S., Miletic, H., Osuagwu, N., Kristoffersen, S., Lilleng, P.K.,
Larsen, J.P., Tysnes, O.-B., Haugarvoll, K., et al. (2016). Defective mitochondrial DNA
homeostasis in the substantia nigra in Parkinson disease. Nat Commun *7*, 13548.

75. Chen, C., Turnbull, D.M., and Reeve, A.K. (2019). Mitochondrial Dysfunction in Parkinson's
Disease-Cause or Consequence? Biology (Basel) *8*,.

76. Pinto, M., and Moraes, C.T. (2014). Mitochondrial genome changes and neurodegenerative
diseases. Biochim Biophys Acta *1842*, 1198–1207.

77. Rodríguez-Santiago, B., Casademont, J., and Nunes, V. (2001). Is mitochondrial DNA
depletion involved in Alzheimer's disease? Eur J Hum Genet *9*, 279–285.

78. Chew, H., Solomon, V.A., and Fonteh, A.N. (2020). Involvement of Lipids in Alzheimer's
Disease Pathology and Potential Therapies. Front Physiol *11*, 598.

79. Yin, J., Reiman, E.M., Beach, T.G., Serrano, G.E., Sabbagh, M.N., Nielsen, M., Caselli,
R.J., and Shi, J. (2020). Effect of ApoE isoforms on mitochondria in Alzheimer disease.
Neurology *94*, e2404–e2411.

80. El Khoury, J.B., Moore, K.J., Means, T.K., Leung, J., Terada, K., Toft, M., Freeman, M.W.,
and Luster, A.D. (2003). CD36 mediates the innate host response to beta-amyloid. J Exp Med
197, 1657–1666.

81. Lämsä, R., Helisalmi, S., Herukka, S.-K., Tapiola, T., Pirttilä, T., Vepsäläinen, S., Hiltunen,
M., and Soininen, H. (2008). Genetic study evaluating LDLR polymorphisms and Alzheimer's
disease. Neurobiol Aging *29*, 848–855.

82. Tomioka, M., Toda, Y., Mañucat, N.B., Akatsu, H., Fukumoto, M., Kono, N., Arai, H., Kioka,
N., and Ueda, K. (2017). Lysophosphatidylcholine export by human ABCA7. Biochim Biophys
Acta Mol Cell Biol Lipids *1862*, 658–665.

83. Nuytemans, K., Maldonado, L., Ali, A., John-Williams, K., Beecham, G.W., Martin, E., Scott,
W.K., and Vance, J.M. (2016). Overlap between Parkinson disease and Alzheimer disease in
ABCA7 functional variants. Neurol Genet *2*, e44.

84. Yang, C., Pring, M., Wear, M.A., Huang, M., Cooper, J.A., Svitkina, T.M., and Zigmond, S.H.
(2005). Mammalian CARMIL inhibits actin filament capping by capping protein. Dev Cell *9*, 209–
221.

85. Crabos, M., Yamakado, T., Heizmann, C.W., Cerletti, N., Bühler, F.R., and Erne, P. (1991).
The calcium binding protein tropomyosin in human platelets and cardiac tissue: elevation in
hypertensive cardiac hypertrophy. Eur J Clin Invest *21*, 472–478.

86. Kostan, J., Salzer, U., Orlova, A., Törö, I., Hodnik, V., Senju, Y., Zou, J., Schreiner, C.,
Steiner, J., Meriläinen, J., et al. (2014). Direct interaction of actin filaments with F-BAR protein
pacsin2. EMBO Rep *15*, 1154–1162.

87. Sever, S. (2002). Dynamin and endocytosis. Curr Opin Cell Biol *14*, 463–467.

88. Cai, B., Giridharan, S.S.P., Zhang, J., Saxena, S., Bahl, K., Schmidt, J.A., Sorgen, P.L.,
Guo, W., Naslavsky, N., and Caplan, S. (2013). Differential roles of C-terminal Eps15 homology
domain proteins as vesiculators and tubulators of recycling endosomes. J Biol Chem *288*,
30172–30180.

89. Di Buduo, C.A., Moccia, F., Battiston, M., De Marco, L., Mazzucato, M., Moratti, R., Tanzi,
F., and Balduini, A. (2014). The importance of calcium in the regulation of megakaryocyte
function. Haematologica *99*, 769–778.

90. De Botton, S., Sabri, S., Daugas, E., Zermati, Y., Guidotti, J.E., Hermine, O., Kroemer, G.,
Vainchenker, W., and Debili, N. (2002). Platelet formation is the consequence of caspase
activation within megakaryocytes. Blood *100*, 1310–1317.

91. Josefsson, E.C., James, C., Henley, K.J., Debrincat, M.A., Rogers, K.L., Dowling, M.R.,
White, M.J., Kruse, E.A., Lane, R.M., Ellis, S., et al. (2011). Megakaryocytes possess a
functional intrinsic apoptosis pathway that must be restrained to survive and produce platelets. J
Exp Med *208*, 2017–2031.

92. Josefsson, E.C., Burnett, D.L., Lebois, M., Debrincat, M.A., White, M.J., Henley, K.J., Lane,
R.M., Moujalled, D., Preston, S.P., O'Reilly, L.A., et al. (2014). Platelet production proceeds
independently of the intrinsic and extrinsic apoptosis pathways. Nat Commun *5*, 3455.

93. McArthur, K., Chappaz, S., and Kile, B.T. (2018). Apoptosis in megakaryocytes and
platelets: the life and death of a lineage. Blood *131*, 605–610.

94. Olkowicz, M., Tomczyk, M., Debski, J., Tyrankiewicz, U., Przyborowski, K., Borkowski, T.,
Zabielska-Kaczorowska, M., Szupryczynska, N., Kochan, Z., Smeda, M., et al. (2021).
Enhanced cardiac hypoxic injury in atherogenic dyslipidaemia results from alterations in the
energy metabolism pattern. Metabolism *114*, 154400.

95. Vyas, C.M., Ogata, S., Reynolds, C.F., Mischoulon, D., Chang, G., Cook, N.R., Manson,
J.E., Crous-Bou, M., De Vivo, I., and Okereke, O.I. (2020). Lifestyle and behavioral factors and
mitochondrial DNA copy number in a diverse cohort of mid-life and older adults. PLoS One *15*,
e0237235.

Figures

Figure 1. Manhattan plot of GWS loci from UKB-only analyses.

Manhattan plot showing genome-wide significant loci for the UK Biobank-only analyses.

Figure 2. Scatterplot displaying effect size estimates between Whites and Blacks GWAS
results for the 129 loci identified in the Whites analyses.

1001

1002   Scatterplot showing comparison between effect size estimates for White (European ancestry)
1003   and Black (African Ancestry) individuals. Color represents significance of effect for each locus
1004   in Blacks GWAS analyses.

1005

1006

1007 **Figure 3. Volcano plot of genes whose predicted gene expression is significantly**
1008 **associated with mtDNA-CN.**

1009

1010 Volcano plot showing genes whose predicted gene expression is significantly associated with
1011 mtDNA-CN. Red indicates positive associations, blue indicates negative associations. Three
1012 genes (ARRDC1, EHMT1, PNPLA7) had extreme effect size estimates greater than 0.3 but
1013 were non-significant and removed from the plot for readability.

1014

Figure 4. PHEWAS-based clustering of mtDNA-CN associated SNPs.

UMAP clusters created from PHEWAS associations for mtDNA-CN associated SNPs. (A) Five clusters were identified as labeled in the panel; orange indicates no cluster. (B-F) SNPs are colored based on their effect estimate size, standardized to the effect on mtDNA-CN (red = positive, blue = negative estimates), for (B) platelet count, (C) mean platelet volume, (D) platelet distribution width, (E) serum calcium levels, (F) mean spherical cellular volume.