Genetic analysis of mitochondrial DNA copy number and associated traits identifies loci implicated in

nucleotide metabolism, platelet activation, and megakaryocyte proliferation, and reveals a causal

association of mitochondrial function with mortality

Longchamps RJ[1*], Yang SY[1*], Castellani CA[1,2], Shi W[1], Lane J[3], Grove ML[4], Bartz

TM[5], Sarnowski C[6], Burrows K[7,8], Guyatt AL[9], Gaunt TR[7,8], Kacprowski T[10], Yang J[11], De Jager PL[12,13], Yu

L[11], CHARGE Aging and Longevity Group, Bergman A[14], Xia R[15], Fornage M[15,16], Feitosa

MF[17], Wojczynski MK[17], Kraja AT [17], Province MA [17], Amin N [18], Rivadeneira F[19], Tiemeier H[18,20],

Uitterlinden AG[18,19], Broer L[19], Van Meurs JBJ[18,19], Van Duijn CM[18], Raffield LM[21], Lange L[22], Rich

SS[23], Lemaitre RN[24], Goodarzi MO[25], Sitlani CM[24], Mak ACY[26], Bennett DA[11], Rodriguez S[7,8], Murabito

JM[27], Lunetta KL[6], Sotoodehnia N[28], Atzmon G[29], Kenny Y[30], Barzilai N[31], Brody JA[32], Psaty BM[33], Taylor

KD[34], Rotter JI[34], Boerwinkle E[4,35], Pankratz N[3], Arking DE[1]

* Indicates authors contributed equally to this work.

1 McKusick-Nathans Institute, Department of Genetic Medicine, Johns Hopkins University School of

Medicine, Baltimore, MD

2 Department of Pathology and Laboratory Medicine, Western University, London, ON, Canada

3 Department of Laboratory Medicine and Pathology, University of Minnesota Medical School,

Minneapolis, MN

4 Human Genetics Center, Department of Epidemiology, Human Genetics, and Environmental Sciences,

School of Public Health, The University of Texas Health Science Center at Houston, Houston, TX

5 Cardiovascular Health Research Unit, Departments of Medicine and Biostatistics, University of

Washington, Seattle, WA

6 Department of Biostatistics, Boston University School of Public Health, Boston, MA

7 MRC Integrative Epidemiology Unit at the University of Bristol, University of Bristol, Oakfield House,

Oakfield Grove, Bristol, UK

8 Population Health Sciences, Bristol Medical School, University of Bristol, Oakfield House, Oakfield

Grove, Bristol, UK

9 Department of Health Sciences, University of Leicester, University Road, Leicester, UK

10 Department of Functional Genomics, Interfaculty Institute for Genetics and Functional Genomics,

University of Greifswald, Greifswald, Germany; Division Data Science in Biomedicine, Peter L. Reichertz

Institute for Medical Informatics, TU Braunschweig and Hannover Medical School, Brunswick, Germany

11 Rush Alzheimer's Disease Center & Department of Neurological Sciences, Rush University Medical

Center, Chicago, IL

12 Center for Translational and Systems Neuroimmunology, Department of Neurology, Columbia

University Medical Center, New York, NY

13 Program in Medical and Population Genetics, Broad Institute, Cambridge, MA

14 Department of Systems and Computational Biology, Albert Einstein College of Medicine, Bronx, New

York, USA

15 Institute of Molecular Medicine, The University of Texas health Science Center at Houston, Houston

TX

16 Human Genetics Center, The University of Texas Health Science Center at Houston, Houston

17 Division of Statistical Genomics, Department of Genetics, Washington University School of Medicine

18 Department of Epidemiology, Erasmus Medical Center, Rotterdam, The Netherlands

19 Department of Internal Medicine, Erasmus Medical Center, Rotterdam, The Netherlands

20 Department of Social and Behavioral Science, Harvard T.H. School of Public Health, Boston, USA

21 Department of Genetics, University of North Carolina at Chapel Hill, Chapel Hill, NC

22 Department of Medicine, University of Colorado Denver, Anschutz Medical Campus, Aurora, CO

23 Center for Public Health Genomics, University of Virginia, Charlottesville, Virginia, USA

24 Cardiovascular Health Research Unit, Department of Medicine, University of Washington, Seattle,

WA

25 Division of Endocrinology, Diabetes and Metabolism, Cedars-Sinai Medical Center, Los Angeles, CA

26 Cardiovascular Research Institute and Institute for Human Genetics, University of California, San

Francisco, California

27 Boston University School of Medicine, Boston University, Boston, MA

28 Cardiovascular Health Research Unit, Division of Cardiology, University of Washington, Seattle, WA

29 Department of Natural science, University of Haifa, Haifa, Israel; Departments of Medicine and

Genetics, Albert Einstein College of Medicine, Bronx, NY, 10461, USA.

30 Department of Epidemiology and Population Health, Albert Einstein College of Medicine, Bronx, NY,

10461, USA.

31 Departments of Medicine and Genetics, Albert Einstein College of Medicine, Bronx, NY, 10461, USA.

32 Cardiovascular Health Research Unit, Department of Medicine, University of Washington, Seattle,

WA

33 Cardiovascular Health Research Unit, Departments of Epidemiology, Medicine and Health Services,

University of Washington, Seattle, WA

34 The Institute for Translational Genomics and Population Sciences, Department of Pediatrics, The

Lundquist Institute for Biomedical Innovation at Harbor-UCLA Medical Center, Torrance, CA

35 Baylor College of Medicine, Human Genome Sequencing Center, Houston, TX

Abstract

Mitochondrial DNA copy number (mtDNA-CN) measured from blood specimens is a minimally

invasive marker of mitochondrial function that exhibits both inter-individual and intercellular variation.

To identify genes involved in regulating mitochondrial function, we performed a genome-wide

association study (GWAS) in 465,809 White individuals from the Cohorts for Heart and Aging Research in

Genomic Epidemiology (CHARGE) consortium and the UK Biobank (UKB). We identified 133 SNPs with

statistically significant, independent effects associated with mtDNA-CN across 100 loci. A combination of

fine-mapping, variant annotation, and co-localization analyses were used to prioritize genes within each

of the 133 independent sites. Putative causal genes were enriched for known mitochondrial DNA

depletion syndromes (p = 3.09 x 10^{-15}) and the gene ontology (GO) terms for mtDNA metabolism (p =

1.43 x 10^{-8}) and mtDNA replication (p = 1.2 x 10^{-7}). A clustering approach leveraged pleiotropy between

mtDNA-CN associated SNPs and 41 mtDNA-CN associated phenotypes to identify functional domains,

revealing three distinct groups, including platelet activation, megakaryocyte proliferation, and mtDNA

metabolism. Finally, using mitochondrial SNPs, we establish causal relationships between mitochondrial

function and a variety of blood cell related traits, kidney function, liver function and overall (p = 0.044)

and non-cancer mortality (p = 6.56 x 10^{-4}).

Introduction

Mitochondria are the cellular organelles primarily responsible for producing the chemical energy

required for metabolism, as well as signaling the apoptotic process, maintaining homeostasis, and

synthesizing several macromolecules such as lipids, heme and iron-sulfur clusters[1,2]. Mitochondria

possess their own genome (mtDNA); a circular, intron-free, double-stranded, haploid, ~16.6 kb

maternally inherited molecule encoding 37 genes vital for proper mitochondrial function. Due to the

integral role of mitochondria in cellular metabolism, mitochondrial dysfunction is known to play a critical

role in the underlying etiology of several aging-related diseases[3–5].

Unlike the nuclear genome, a large amount of variation exists in the number of copies of mtDNA present

within cells, tissues, and individuals. The relative copy number of mtDNA (mtDNA-CN) has been shown

to be positively correlated with oxidative stress[6], energy reserves, and mitochondrial membrane

potential[7]. As a minimally invasive proxy measure of mitochondrial dysfunction[8], decreased mtDNA-CN

measured in blood has been previously associated with aging-related disease states including frailty[9],

cardiovascular disease[10–12], chronic kidney disease[13], neurodegeneration[14,15], and cancer[16].

Although mtDNA-CN measured from whole blood presents itself as an easily accessible and minimally

invasive biomarker, cell type composition has been shown to be an important confounder, complicating

analyses[17,18]. For example, while platelets generally have fewer mtDNA molecules than leukocytes, the

lack of a platelet nuclear genome drastically skews mtDNA-CN estimates. As a result, not only is

controlling for cell composition extremely vital for accurate mtDNA-CN estimation, interpreting the

results in relation to the impact of cell composition becomes a necessity[18–20].

Although the comprehensive mechanism through which mtDNA-CN is modulated is largely unknown[21,22],

twin studies have estimated a broad-sense heritability of ~0.65, consistent with moderate genetic

control[23]. Several nuclear genes have been shown to directly modulate mtDNA-CN, specifically those

within the mtDNA replication machinery such as the mitochondrial polymerase, *POLG* and *POLG2*[24,25], as

well as the mitochondrial DNA helicase, *TWNK*, and the mitochondrial single-stranded binding protein,

mtSSB[26]. Furthermore, nuclear genes which maintain proper mitochondrial nucleotide supply including

DGUOK and *TK2* have also been shown to regulate mtDNA-CN[27–29]. To further elucidate the genetic

control over mtDNA-CN, several genome-wide association studies (GWAS) of mtDNA-CN have been

published[30–33], including a study that was published while the current manuscript was in preparation,

analyzing ~300,000 participants from the UK Biobank (UKB), and identifying 50 independent loci[33].

In the present study, we report mtDNA-CN GWAS results from 465,809 individuals across the Cohorts

for Heart and Aging Research in Genomic Epidemiology (CHARGE) consortium[34] and the UK Biobank

(UKB)[35]. Using multiple gene prioritization and functional annotation methods, we assign genes to loci

that reach genome-wide significance. We perform a PHEWAS and group our genome-wide significant

SNPs into 3 clusters that represent distinct functional domains related to mtDNA-CN. Finally, we

leverage mitochondrial SNPs to establish causality between mitochondrial function and mtDNA-CN

associated traits.

Subjects and Methods

Study Populations

470,579 individuals participated in this GWAS, 465,809 of whom self-identified as White. Participants

were derived from 7 population-based cohorts representing the Cohorts for Heart and Aging Research in

Genetic Epidemiology (CHARGE) consortium (Avon Longitudinal Study of Parents and Children [ALSPAC],

Atherosclerosis Risk in Communities [ARIC], Cardiovascular Health Study [CHS], Multi-Ethnic Study of

Atherosclerosis [MESA], Religious Orders Study and Memory and Aging Project [ROSMAP], Study of

Health in Pomerania [SHIP]) and from the UK Biobank (UKB) (Supplemental Table 1). Detailed

descriptions of each participating cohort, their quality control practices, study level analyses, and ethic

statements are available in the Supplemental Methods. All study participants provided written informed

consent and all centers obtained approval from their institutional review boards.

Methods for Mitochondrial DNA Copy Number Estimation (CHARGE cohorts)

qPCR

mtDNA-CN was determined using a quantitative PCR assay as previously described[32,36]. Briefly, the cycle

threshold (Ct) value of a nuclear-specific and mitochondrial-specific probe were measured in triplicate

for each sample. In CHS, a multiplex assay using the mitochondrial *ND1* probe and nuclear *RPPH1* probe

was used, whereas ALSPAC used a mitochondrial probe targeting the D-Loop and a nuclear probe

targeting *B2M*. In CHS, we observed plate effects, as well as a linear increase in ΔCt due to the pipetting

order of each replicate. These effects were corrected in the analysis using linear mixed model

regression, with pipetting order included as a fixed effect and plate as a random effect to create a raw

measure of mtDNA-CN. In ALSPAC, run-to-run variability was controlled using 3 calibrator samples

added to every plate, to allow for adjustment by a per-plate calibration factor[32].

Microarray

Microarray probe intensities were used to estimate mtDNA-CN using the Genvisis software package[37] as

previously described[10,36]. Briefly, Genvisis uses the median mitochondrial probe intensity across all

homozygous mitochondrial SNPs as an initial estimate of mtDNA-CN. Technical artifacts such as DNA

input quality, DNA input quantity, and hybridization efficiency were captured through either surrogate

variable (SV) or principal component (PC) analyses. SVs or PCs were adjusted for through stepwise linear

regression by adding successive components until each successive surrogate variable or principal

component no longer significantly improved the model.

Whole Genome Sequencing (ARIC)

Whole genome sequencing read counts were used to estimate mtDNA-CN as previously described[36].

Briefly, the total number of reads in a sample were web scraped from the NCBI sequence read archive.

Mitochondrial reads were downloaded directly from dbGaP through Samtools (1.3.1). There was no

overlap between ARIC microarray and ARIC whole-genome sequencing samples. A ratio of mitochondrial

reads to total aligned reads was used as a raw measure of mtDNA-CN.

Adjusting for Covariates

Each method described above represents a raw measure of mtDNA-CN, adjusted for technical artifacts;

however, several potential confounding variables (e.g., age, sex, blood cell composition) have been

identified previously[18]. Raw mtDNA-CN values were adjusted for white blood cell count in ARIC, SHIP and

CHS (which also adjusted for platelet count), depending on available data. Standardized residuals (mean

= 0, standard deviation = 1) of mtDNA-CN were used after adjusting for covariates (Supplemental Table

1).

Estimation of Mitochondrial DNA Copy Number (UKB)

Due to the availability of more detailed cell count data, as well as a different underlying biochemistry for

the Affymetrix Axiom array compared to the genotyping arrays used in the CHARGE cohorts, mtDNA-CN

in the UKB was estimated differently (Supplemental Methods). Briefly, mtDNA-CN estimates derived

from whole exome sequencing data, available on ~50,000 individuals, were generated first using

customized Perl scripts to aggregate the number of mapped sequencing reads and correct for covariates

through both linear and spline regression models. Concurrently, mitochondrial probe intensities from

the Affymetrix Axiom arrays, available on the full ~500,000 UKB cohort, were adjusted for technical

artifacts through principal components generated from nuclear probe intensities. Probe intensities were

then regressed onto the whole exome sequencing mtDNA-CN metric, and beta estimates from that

regression were used to estimate mtDNA-CN in the full UKB cohort. Finally, we used a 10-fold cross

validation method to select the cell counts to include in the final model (Supplemental Table 2). The

final UKB mtDNA-CN metric is the standardized residuals (mean = 0, standard deviation = 1) from a

linear model adjusting for covariates (age, sex, cell counts) as described in the Supplemental Methods.

Genome-Wide Association Study

For each individual cohort, regression analysis was performed with residualized mtDNA-CN as the

dependent variable adjusting for age, sex, and cohort-specific covariates (e.g., principal components,

DNA collection site, family structure, cell composition). Cohorts with multiple mtDNA-CN estimation

platforms were stratified into separate analyses. Ancestry-stratified meta-analyses were performed

using Metasoft software using the Han and Eskin random effects model to control for unobserved

heterogeneity due to differences in mtDNA-CN estimation method[38]. Effect size estimates for SNPs

were calculated using a random effect meta-analysis from cohort summary statistics, as the Han and

Eskin model relaxes the assumption under the null hypothesis without modifying the effect size

estimates that occur under the alternative hypothesis[38]. In total, three complementary analyses were

performed in self-identified White individuals: (1) a meta-analysis using all available studies, (2) a meta-

analysis of studies with available data for cell count adjustments, and (3) an analysis of UKB-only data.

As the vast majority of samples are derived from the UKB study, and given the difficulty in interpreting

effect size estimates from a random effects model, further downstream analyses were all performed

using effect size estimates from UKB-only data. We additionally performed X chromosome analyses,

using only UKB data. X-chromosome analyses were stratified by sex (males = 194,151, females =

216,989), and summary statistics were meta-analyzed using METAL[39] to obtain the final effect estimates.

SNP Heritability Estimation

SNP heritability estimates were retrieved from BOLT-LMM[40]. To verify this metric, we used SumHer[41] to

calculate an independent heritability metric using summary statistics. The heritability model used in this

analysis was the BLD-LDAK model. The tagging file used is the pre-computed UK Biobank GBR version for

the corresponding heritability model. The summary statistics were filtered so that only single-character

reference and alternate alleles are allowed. Chr:BP combination duplicates were removed except the

first appearance. SNP heritability was then calculated and extracted from output files.

Identification of Independent GWAS Loci

To identify the initial genome-wide significant (lead) SNPs in each locus, the most significant SNP that

passed genome-wide significance ($p < 5 \times 10^{-8}$) within a 1 Mb window was selected. To avoid Type I

error, SNPs were only retained for further analyses if there were either (a) at least two genome-wide

significant SNPs in the 1 Mb window or (b) if the lead SNP was directly genotyped. Conditional analyses

were performed in UKB, where the lead SNPs from the original GWAS were used as additional covariates

in order to identify additional independent associations.

Comparisons with Hägg et al. 2020

To compare results with Hägg et al. 2020[33], summary statistics were obtained from their Supplementary

Table 4. Loci were identified as shared between the two GWAS if two lead SNPs were fewer than

500,000 base pairs apart from one another.

Fine-mapping

The susieR package was used to identify all potential causal variants for each independent locus

associated with mtDNA CN[42]. UKB imputed genotype data for unrelated White subjects were used and

variants were extracted using a 500 kb window around the lead SNP for each locus with minor allele

frequency (MAF) > 0.001. 95% credible sets (CS) of SNPs, containing a potential causal variant within a

locus, were generated. The minimum absolute correlation within each CS is 0.5 and the scaled prior

variance is 0.01. When the CS did not include the lead SNP identified from the GWAS, some of the

parameters were slightly relaxed [minimum absolute correlation is 0.2, estimate prior variance is TRUE].

The SNP with the highest posterior inclusion probability (PIP) within each CS was also identified

(Supplemental Table 3). With a few exceptions, final lead SNPs were selected by prioritizing initially

identified SNPs unless the SNP with the highest PIP had a PIP greater than 0.2 and was 1.75 times larger

than the SNP with the second highest PIP.

Functional Annotation and Gene Prioritization

Functional Annotation

ANNOVAR was used for functional annotation of variants identified in the fine-mapping step[43]. First,

variants were converted to an ANNOVAR-ready format using the dbSNP version 150 database[44]. Then,

variants were annotated with ANNOVAR using the RefSeq Gene database[45]. The annotation for each

variant includes the associated gene and region (e.g., exonic, intronic, intergenic). For intergenic

variants, ANNOVAR provides flanking genes and the distance to each gene. For exonic variants,

annotations also include likely functional consequences (e.g., synonymous/nonsynonymous,

insertion/deletion), the gene affected by the variant, and the amino acid sequence change

(Supplemental Table 4).

Co-localization Analyses

Co-localization analyses were performed using the approximate Bayes factor method in the R package

coloc[46]. Briefly, *coloc* utilizes eQTL data and GWAS summary statistics to evaluate the probability that

gene expression and GWAS data share a single causal SNP (colocalize). *Coloc* returns multiple posterior

probabilities; H0 (no causal variant), H1 (causal variant for gene expression only), H2 (causal variant for

mtDNA-CN only), H3 (two distinct causal variants), and H4 (shared causal variant for gene expression

and mtDNA-CN). In the event of high H4, we designate the gene as causal for the GWAS phenotype of

interest (mtDNA-CN). eQTL summary statistics were obtained from the eQTLGen database[47]. Genes with

significant associations with lead SNPs were tested for co-localization using variants within a 500 kb

window of the sentinel SNP. Occasionally, some of the eQTLGen p-values for certain SNPs were

identical due to R's (ver 4.0.3) limitation in handling small numbers. To account for this, if the absolute

value for a SNP's z-score association with a gene was greater than 37.02, z-scores were rescaled so that

the largest z-score would result in a p-value of 5×10^{-300}. Additionally, a few clearly co-localized genes

did not result in high H4 PPs due to the strong effect for each phenotype of a single SNP (Supplemental

Figure 1), possibly due to differences in linkage disequilibrium (LD) between the UKB and eQTLGen

populations. To account for this, we summed mtDNA-CN GWAS p-values and eQTLGen p-values for each

SNP and removed the SNP with the lowest combined p-value. Co-localization analyses were then

repeated without the lowest SNP. Genes with H4 greater than 50% were classified as genes with

significant evidence of co-localization.

DEPICT

Gene prioritization was performed with Depict, an integrative tool that incorporates gene co-regulation

and GWAS data to identify the most likely causal gene at a given locus[48]. Across GWAS SNPs which

overlapped with the DEPICT database, we identified SNPs representing 119 independent loci with LD

pruning defined as $p < 5 \times 10^{-8}$, $r^2 < 0.05$ and > 500 kb from other locus boundaries. Only genes with a

nominal p-value of less than 0.05 were considered for downstream prioritization.

Gene Assignment

To prioritize genes for each identified locus, we utilized functional annotations, eQTL co-localization

analyses, and DEPICT gene prioritization results (Supplemental Figure 2). First, genes with missense

variants within susieR fine-mapped credible sets were assigned to loci. If loci co-localized with a gene's

expression with a posterior probability (PP) of greater than 0.50 and there were no other co-localized

genes with a PP within 5%, the gene with the highest posterior probability was assigned. If there was still

no assigned gene, the most significant DEPICT gene was assigned. If there was no co-localization or

DEPICT evidence, the nearest gene was assigned.

Gene Set Enrichment Analyses

Using the finalized gene list from the prioritization pipeline, GO and KEGG pathway enrichment analyses

were performed using the "goana" and "kegga" functions from the R package *limma*[49], treating all

known genes as the background universe[50]. Only one gene per locus was used for "goana" and "kegga"

gene set enrichment analysis, prioritizing genes assigned to primary independent hits. If there were

multiple assigned genes, one gene was randomly selected to avoid biasing results through loci with

multiple functionally related genes. To identify an appropriate p-value cutoff, 100 genes were randomly

selected from the genome and run through the same enrichment analysis. This permutation was

repeated 1000 times to generate a null distribution of the smallest p-values from each permutation. For

cluster-specific gene set enrichment analyses, permutation testing used the same number of random

genes as the number of genes in each cluster. To ensure robustness of results, gene set enrichment

analysis was repeated 50 times with random selection of genes at loci with multiple assigned genes. GO

and KEGG terms that passed permutation cutoffs at least 40/50 times were retained.

Gene-based Association Test

We used metaXcan, which employs gene expression prediction models to evaluate associations

between phenotypes and gene expression[51]. We obtained pre-calculated expression prediction models

and SNP covariance matrices, computed using whole blood from European ancestry individuals in

version 7 of the Genotype-Tissue expression (GTEx) database[52]. Using prediction performance $p < 0.05$, a

total of 6,285 genes were predicted. Of these genes, 74 passed Bonferroni correction of $p < 7.95 \times 10^{-6}$.

Gene set enrichment analyses were performed on Bonferroni-significant genes as previously described.

REVIGO[53] was used on the "medium" setting (allowed similarity = 0.7) to visualize significantly enriched

GO terms.

We used a one-sided Fisher's exact test to test for enrichment of genes that have been previously

identified as causal for mtDNA depletion syndromes[54-56].

PHEWAS-based SNP Clustering

mtDNA-CN Phenome-wide Association Study (PHEWAS)

We used the PHEnome Scan ANalysis Tool (PHESANT)[57] to identify mtDNA-CN associated quantitative

traits in the UKB. Briefly, we tested for the association of mtDNA-CN with 869 quantitative traits

(Supplemental Table 5), limiting analyses to 365,781 White, unrelated individuals

(used.in.pca.calculation=1). As extreme cell count measurements could indicate individuals with active

infections or cancers, they were excluded from analysis (see Supplemental Methods). Analyses were

adjusted for age, sex, and assessment center.

SNP-Phenotype Associations

SNP genotypes were regressed on mtDNA-associated quantitative phenotypic traits using linear

regression, adjusted for sex, age with a natural spline (df = 2), assessment center, genotyping array, and

40 genotyping principal components (provided as part of the UKB data download).

SNP Clustering

To identify distinct clusters of mtDNA-CN GWS SNPs based on phenotypic associations, beta estimates

from the SNP-phenotype associations were first divided by the beta estimate of the mtDNA-CN SNP-

mtDNA-CN association, so that all SNP-phenotype associations are relative to the mtDNA-CN increasing

allele and scaled to the effect of the SNP on mtDNA-CN. The adjusted beta estimates were subjected to

a dimensionality reduction method, Uniform Manifold and Approximation Projection (UMAP), as

implemented in the R package *umap*[58] (random_state=123, n_neighbors=10, min_dist=0.001,

n_components=2, n_epochs=200). SNPs were assigned to clusters using Density Based Clustering of

Applications with Noise (DBSCAN) as implemented in the R package *dbscan*[59] (minPts=10). Robustness of

cluster assignment was established by varying n_neighors, min_dist, and random_state parameters.

Clusters represent groups of SNPs with similar phenotypic associations.

Phenotype Enrichment and Permutation Testing

To test for enrichment of *individual* phenotypes within clusters, we compared the median mtDNA-CN

scaled phenotype beta estimates within the cluster to the median beta estimates for all SNPs not in the

cluster, with significance determined using 20,000 permutations in which cluster assignment was

permuted. For multi-test correction *across all* phenotypes, we performed 300 permutations of the initial

cluster assignment, followed by the comparison of median beta estimates as described above. We

retained only the most significant result from across all phenotypes and clusters from each of the 300

permutations, and then selected the 15th most significant value as the study-wide threshold for multi-

test corrected significance of $p < 0.05$.

mtDNA Variant Association Analyses

Mitochondrial Variant Phasing and Imputation

Shapeit4 and Impute5 were used for UK Biobank mtDNA genotype phasing and imputation[60,61]. Phasing

and imputation were performed separately for each genotyping array (UKBB, UKBL), and restricted to

self-identified White individuals. The reference panel used for imputation analysis was the 1000

Genomes Project phase 3 mtDNA variants[62]. UK Biobank genotypes were coded to match the reference

panel allele. All genotype files, including the reference panel, were phased using Shapeit4 to fill in any

missing genotypes using the phasing iteration sequence "10b,1p,1b,1p,1b,1p,1b,1p,10m", where b is

burn-in iteration, p is pruning iteration, and m is main iteration. The –sequencing option was also used

due to the presence of multiple mtDNA variants in a very small region, analogous to sequencing data.

Phased UK Biobank genotypes were then imputed with the reference panel using Impute5 with

the following parameters: –pbwt-depth 8; –pbwt-cm 0.005; –no-threshold. All imputed variants were

functionally annotated using MSeqDR mvTools[63].

mtSNP Association Tests

Linear regressions stratified by genotyping array (UKBB, UKBL) were performed for each mtDNA SNP on

the 41 traits and mtDNA-CN, including the following covariates: age, age^2, sex, center, first 20

genotyping PCs. Only SNPs with MAF > 0.005 and imputation INFO score > 0.80 were included (UKBB,

n=223; UKBL, n=190; both, n=149). Results were then meta-analyzed using inverse variance weighting.

For association analyses between mitochondrial SNPs/haplotypes and mtDNA-CN, the mtDNA-CN metric

used was derived only from WES data, as mtDNA genotypes can subtly influence the estimates from

genotype array intensities.

Identification of Independent Genetic Effects

Single SNP study-wide significance was established by generating 300 normally distributed dummy

traits, and running single SNP tests using the UKBB data. The minimum SNP p-value for each dummy

trait was then selected, and the 15th most significant p-value from the 300 analyses was divided by 42

(41 real traits + mtDNA-CN), resulting in study-wide p-value threshold of $p < 9.5 \times 10^{-6}$. To identify a

subset of traits to perform credible set identification using SusieR (see above, **Fine Mapping**), SNPs were

first filtered based on the study-wide p-value threshold, and then then most significantly associated trait

was identified for each SNP. SusieR, (parameters: L=10, estimate_residual_variance=TRUE,

estimate_prior_variance=TRUE, check_z = FALSE) was then run for each of these traits using the UKBB

imputed data and summary association test statistics. A total of 7 credible sets were identified across

the 4 traits, two of which co-localized, resulting in 6 credible sets. Independence across the 6 credible

sets was tested using multivariate regression models, and requiring $p < 0.0005$ for at least one trait for a

SNP to remain in the model. SNPs MT73A_G and MT 7028C_T were in moderate high LD (r^2=0.67), but

based on conditional regression analyses as described in the main results, capture independent effects

and are associated with different traits.

Haplotype Generation and Analysis

Haplotype were constructed by concatenating SNPs across the 6 credible sets using SNPs directly

genotyped on both genotyping arrays. This required selecting a SNP with a lower PIP for 2 of the 6

credible sets (MT12612A_G replaced MT462C_T, r^2 = 0.81; MT10238T_C replaced MT4529A_T, r^2 =

0.89). Haplotypes with MAF < 0.005 were set to missing (n = 1607), resulting in 8 haplotypes, with the

most common haplotype set as reference. Significance for haplotype associations with each trait were

generated by an anova between regression models with and without the haplotypes. Covariates

included age, age^2, sex, center, first 40 genotyping PCs, and genotyping array.

Mortality analyses were run using Cox proportional hazards models, with covariates as above.

Individuals with external causes of death (ICD 10 Death Code categories V, W, X, Y) were censored at

time of death. Additionally, for non-cancer mortality analyses, cancer death (ICD 10 Death Code

categories C00-D48) were censored. For cancer mortality analyses, all death due to non-cancer cases

were censored at time of death.

Clustering for visualization was performed using the R package 'heatmaply', with default setting and

hclust_method="ward.D2".

All statistical analyses were performed using R version 4.0.3.

Results

Sample Characteristics

The current study included 465,809 White individuals (53.9% female) with an average age of 56.6 yrs (sd

= 8.2 yrs) (Supplemental Table 1). Follow-up validation analyses were performed in 4,770 Blacks (60.2%

female) with an average age of 61.2 yrs (sd = 7.4 yrs). The majority of the data originated from the UKB

(93%). The bulk of the DNA used for mtDNA-CN estimation was derived from buffy coat (95.5%) while

the rest was derived from peripheral leukocytes (2.2%), whole blood (2.3%), or brain (< 0.2%). mtDNA-

CN estimated from Affymetrix genotyping arrays consisted of 97.9% of the data while the remainder

was derived from qPCR (1.8%) and WGS (0.3%).

GWAS

Previous work has demonstrated that the method used to measure mtDNA-CN can impact the strength

of association[36]. To account for potential differences across studies due to the different mtDNA-CN

measurements used, as well as the inclusion of blood cell counts as covariates in only a subset of the

cohorts, we took two approaches. First, we used a random effects model to perform meta-analyses,

allowing for different genetic effect size estimates across cohorts. Second, we

performed three complementary analyses in individuals who self-identified as White: 1) meta-analysis of

all available studies (n = 465,809); 2) meta-analysis of studies with available data for cell count

adjustment (n = 456,151); and 3) GWAS of UKB only (n = 440,266) (Figure 1). 77 loci were significant in

all three meta-analyses, and we identified 93 independent loci that were significant in at least one of the

analyses. In the meta-analysis of UKB-only data, 92 of the total 93 loci were identified (Supplemental

Figure 3). Given that >90% of the samples come from the UKB study, and the challenge of interpreting

effect size estimates from a random effects model, downstream analyses all use effect size estimates

from the UKB only analyses (Supplemental Table 6), which showed no evidence for population

substructure inflating test statistics, with a genomic inflation factor of 1.09 (Supplemental Figure 4). SNP

heritability estimated from BOLT-LMM[40] for mtDNA-CN adjusted for age and sex was 10.5% while

heritability for mtDNA-CN adjusted for age, sex, and cell counts was 7.4%, implying that some of the

mtDNA-CN heritability observed in previous studies could be due to heritability of cell type composition.

We also used SumHer[41] as an alternative approach to calculating SNP heritability, which returned a

comparable estimate of 7.0% for the cell-count corrected mtDNA-CN metric.

The most significant SNP associated with mtDNA-CN was a missense mutation in *LONP1* (p =

3.00×10^{-141}), a gene that encodes a mitochondrial protease that can directly bind mtDNA, and has been

shown to regulate *TFAM*, a transcription factor involved in mtDNA replication and repair (for review see

Gibellini *et al.*)[64].

Meta-analysis of the sex-stratified X chromosome results identified four loci significantly

associated with mtDNA-CN, with directionality consistent across the male and female stratified analyses

(Supplemental Table 7).

Fine-mapping and Secondary Hits

To identify additional independent SNPs within novel loci whose effects were masked by the original

significant SNP, as well as identify additional loci, we took two approaches. First, a conditional analysis

adjusting for the top 93 SNPs from the initial (primary) GWAS run revealed 3 novel loci and 19 additional

independent significant SNPs within existing loci. We also performed fine-mapping with susieR[42] and

discovered an additional 14 independent SNPs within existing loci. The majority of loci had only one 95%

credible set of SNPs; further, twenty of the credible sets contained only one SNP. However, many of the

credible sets contained greater than 50 SNPs after fine-mapping, and 12 of the 122 credible sets had a

missense SNP as the SNP with the highest PIP in the set. Using these two methods, we identified in total

129 independent SNPs across 96 autosomal loci (Supplemental Figure 5), while susieR fine-mapping and

conditional analyses for the X-chromosome loci did not reveal any additional secondary signals.

Comparisons with Hägg et al. 2020

Out of the 50 loci reported in Hägg et al. 2020[33], we replicate 38 loci in our cell-count adjusted analyses

(Supplemental Table 8). As the two GWASs both use UK Biobank data, this replication is unsurprising.

Out of the 12 loci that were not genome-wide significant in our cell-count adjusted analyses, 11 were

significant when we did not adjust our mtDNA-CN metric for cell counts, suggesting that cell-type

composition may be driving these signals. The current manuscript also reports 62 additional loci that are

not in the Hägg et al. 2020 study. This is likely due to increased power, as the sample size used for the

current analyses is nearly twice as large.

Associations in Black Populations

Examining the 129 autosomal SNPs from the Whites-only analysis, 99 were available in the Blacks-

only meta-analysis (n = 4770). After multiple testing correction, one of these SNPs was significant

(rs73349121, p = 0.0001), 9 were nominally significant ($p < 0.05$, with 5 expected), and 58/99 had a

direction of effect that was consistent with the Whites-only analyses (one-sided p = 0.04, Figure

2). Despite being under-powered, these results in the Blacks-only analyses provide evidence for similar

genetic effects in a different ancestry group.

Gene Prioritization and Enrichment of mtDNA Depletion Syndrome Genes

We integrated results from three different gene prioritization and functional annotation methods

(ANNOVAR[43], COLOC[46] , and DEPICT[48]) so that loci with nonsynonymous variants in gene exons were

prioritized first, with eQTL co-localization results considered second (Supplemental Table 9), and those

from DEPICT (Supplemental Table 10) were considered last (Supplemental Figure 2). For 20 loci, multiple

genes were assigned as analyses could not identify a single priority gene (Supplemental Table 11). As

eQTLGen did not evaluate X chromosome variants, three of the four X-chromosome loci were assigned

to the nearest gene. SLC25A5 was assigned to rs392020, as the second highest PIP SNP was an exonic

nonsynonymous variant.

We noted the identification of a number of mtDNA depletion syndrome genes in the priority list

and tested for enrichment of these known causal genes using a one-sided Fisher's exact test. For this

analysis, all genes for loci assigned to multiple genes were used, and genes for all primary and secondary

loci were considered. Our gene prioritization approach identified 7 of 16 mtDNA depletion genes

(Supplemental Table 12), consistent with a highly significant enrichment (one-sided p = 3.09 x 10^{-15}).

Gene Set Enrichment Analyses

To avoid bias from a single locus with multiple functionally related genes contributing to a false-positive

signal, only one gene per unique locus was used, prioritizing genes assigned to primary loci. For loci with

multiple assigned genes, one gene was randomly selected for testing. To test for robustness of gene set

enrichment results, random selection was repeated 50 times, and only gene sets that were significantly

enriched for at least 40 iterations were retained. In all, a total of 100 genes were utilized for GO term

and KEGG pathway enrichment analyses. Using a Bonferroni-corrected p-value cutoff, 12 gene sets were

significantly enriched for all 50 iterations, including mitochondrial nucleoid, mitochondrial DNA

replication, and amyloid-beta clearance (Supplemental Table 13). No KEGG terms were significant across

multiple iterations.

MetaXcan Gene Expression Analysis

As a complementary approach to single-SNP analyses, we explored the associations between mtDNA-CN

and predicted gene expression using MetaXcan[51] MetaXcan incorporates multiple SNPs within a locus

along with a reference eQTL dataset to generate predicted gene expression levels. As our study

estimated mtDNA-CN derived from blood, we used whole blood gene expression eQTLs from the Gene-

Tissue Expression (GTEx) consortium[65] to predict gene expression in the UKB dataset. We identified

6,285 genes that had a predicted performance $p < 0.05$ (i.e., they had sufficient data to generate robust

gene expression levels) and were tested for association with mtDNA-CN. Of these genes, 74 were

significantly associated with mtDNA-CN ($p < 7.95 \times 10^{-6}$) (Figure 3, Supplemental Table 14), including 8

that were not identified through single-SNP analyses. Many of the significant genes have known

mitochondrial functions, notably the mtDNA transcription factor *TFAM* ($p = 1.09 \times 10^{-29}$) and

mitochondrial exonuclease *MGME1* ($p = 5.87 \times 10^{-23}$), genes known as causal for mtDNA depletion

syndromes[54,55]. Additionally, *LONP1*, *MRPL43*, and *BAK1*, are all genes with known mitochondrial

functions[66–68]. Bonferroni significant MetaXcan genes were used for gene enrichment analysis, finding

enrichment for "nucleobase metabolic process" ($p = 1.47 \times 10^{-4}$) and "mitochondrial fusion" ($p = 1.86 \times$

10^{-4}) (Supplemental Figure 6).

PHEWAS-based SNP Clustering and Gene Set Enrichment

mtDNA-CN is associated with numerous quantitative and qualitative phenotypes, many of which are

relevant to aging-related disease[3–5,9,10,13–16]. We hypothesized that this pleiotropy may reflect different

underlying functional domains captured by mtDNA-CN, and may be reflected in GWAS-identified SNPs

and their likely causal genes. To test this hypothesis, we used the UKB data to identify quantitative traits

associated with mtDNA-CN and selected 41 highly significant, non-redundant traits to test for

association with the mtDNA-CN GWAS SNPs (Supplemental Table 5, in PHEWAS = 1). We clustered SNPs

using the trait effect size (beta) divided by the mtDNA-CN effect size estimate, so that all effects are

standardized to the effect of the mtDNA-CN increasing allele for each locus. We identified 3 clusters of

SNPs (Supplemental Table 15, Figure 4A), with cluster 1 containing SNPs in which the mtDNA-CN

increasing allele is associated with decreased platelet count (PLT) (Figure 4B), increased mean platelet

volume (MPV) (Figure 4C), and platelet distribution width (PDW) (Figure 4D), consistent with a role in

platelet activation[69]. Cluster 2 is most strongly enriched for SNPs in which the mtDNA-CN increasing

allele is associated with increased PLT, plateletcrit (PCT, a measure of total platelet mass), serum

calcium (Figure 4E), serum phosphate, as well as decreased mean corpuscular volume (MCV) and mean

spherical cellular volume (Figure 4F) (Supplemental Table 16). The cluster 2 phenotypes implicate

megakaryocyte proliferation and proplatelet formation in addition to apoptosis and autophagy, and are

supported by the genes identified for this cluster (megakaryocyte proliferation and proplatelet

formation: *MYB*, *JAK2*[70], apoptosis and autophagy: *BAK1*, *BCL2, TYMP*)[71]. Gene set enrichment analysis

confirmed this, as cluster 2 genes are significantly enriched for extrinsic apoptosis signaling pathways in

the absence of ligand (Supplemental Table 17). Cluster 3 did not yield any specific trait enrichment (all

significant results reflected the strong enrichment observed in clusters 1-2); however, gene set

enrichment for this cluster identified multiple mtDNA-related gene ontology terms, including

mitochondrial DNA replication, gamma DNA polymerase complex, and mitochondrial nucleoid

(Supplemental Table 18).

Determination of causal associations between mitochondrial function and mtDNA-CN associated traits

The extensive pleiotropy and limited variance explained of nuclear DNA SNPs associated with mtDNA-CN

(<1% of the variance in mtDNA-CN explained by GWS loci when predicted into the ARIC cohort)

precludes the use of traditional Mendelian randomization (MR) approaches to establish causality

between mtDNA-CN and the 41 identified mtDNA-CN associated traits. As alternative approach, we

examined the association of mitochondrial SNPs with mtDNA-CN and the 41 traits, under the

assumption that these SNPs can only act through alteration of mitochondrial function, and thus a

significant association implies causality. Imputation and analyses of mitochondrial SNPs were run

stratified by genotyping array (see Methods), and then meta-analyzed using inverse-variance weighting.

After multi-test correction ($p < 9.5 \times 10^{-6}$), we identified 45 SNPs associated with 1 or more of the traits,

ranging from 1 to 6 traits per SNP (Supplementary Table 19). To identify independent effects, we first

identified the most significantly associated trait for each SNP, highlighting 4 traits (aspartate

aminotransferase, creatinine, MCV, PCT) in which to run susieR to identify independent credible sets.

We identified 6 independent effects across the four traits, with MCV credible set 4 and platelet credible

set 1 representing the same effect. We note that 2 of the SNPs are in moderately high LD (MT73A_G and

MT7028C_T, r^2=0.67), however, conditional analyses demonstrate that MT73A_G is associated with

creatinine, and not MCV, and the reverse is seen for MT7028C_T (Supplemental Table 20). Leveraging

the haploid nature of the mitochondrial genome, we selected the directly genotyped SNP with the

highest PIP from each credible set (Supplemental Table 21), and identified 8 haplotypes with MAF >

0.005 (Supplemental Table 22). Comparing linear regression models with and without the haplotypes in

the model, we identify 14 traits nominally associated, and 9 traits significantly associated after

Bonferroni correction, with mtDNA genetic variation (Supplemental Table 23, Figure 5). These results

causally implicate mitochondrial function in a variety of cell related traits (MCV, MSCV, MPV, PCT,

Platelet), kidney function (creatinine), liver function (aspartate and alanine aminotransferases) and

mtDNA-CN.

Association of mitochondrial function with mortality

We have previously shown that mtDNA-CN is associated with overall mortality[9]. As above, we

collectively tested the mitochondrial haplotypes for association with mortality not due to external

causes (e.g., no accidents, falls, see Methods ; n = 24,622, median follow-up time = 4318 days), and

found a nominally significant association with overall mortality (p = 0.044, Figure 6). Given the

conflicting reports between increased mitochondrial function and both increased and decreased cancer

risk[16,72,73], we looked separately at cancer (n = 13,231) and non-cancer mortality (n = 11,391). While

there was no association with cancer mortality (p = 0.74), we saw a highly significant association with

non-cancer mortality (p = 6.56×10^{-4}).

Discussion

We conducted a GWAS for mtDNA-CN using 465,809 individuals from the CHARGE consortium and the

UKB. We report 133 independent signals originating from 100 loci, the majority of which were not

identified in previous studies. Examining our GWS SNPs in a Black population, we observed a concordant

signal, suggesting that the genetic etiology of mtDNA-CN may be broadly similar across populations.

Using several functional follow-up methods, genes were assigned for each identified independent hit

and significant enrichment was observed for genes involved in mitochondrial DNA metabolism,

homeostasis, cell activation, and amyloid-beta clearance. In total, we assigned 128 unique genes to

independent GWAS signals associated with mtDNA-CN. We also identified 8 additional genes whose

predicted gene expression is associated with mtDNA-CN that could not be mapped back to GWS loci.

Finally, using a clustering approach based on SNP associations with various mtDNA-CN associated

phenotypes, we were able to functionally categorize SNPs, providing insight into biological pathways

that impact mtDNA-CN.

We note that during the preparation of this manuscript, a GWAS for mtDNA-CN performed in

295,150 unrelated individuals from the UK Biobank was published, which reported 50 genome-wide

significant regions[33]. Within our GWAS, we replicate 38 of these 50 genome-wide significant loci in our

cell count corrected analyses. An additional 11 out of the remaining 12 loci are genome-wide significant

when we do not adjust mtDNA-CN for cell count. While Hagg et. al adjust for cell type composition, this

difference suggests that their adjustment may not be fully capturing the effects of cell counts.

Additionally, our analyses report 59 additional loci that are not observed in the previous paper, largely

due to the increased power of our study.

We were able to identify a substantial proportion of the genes involved in mtDNA depletion

syndromes (7/16, p = 3.09 x 10^{-15} for enrichment), including *TWNK, TFAM, DGUOK*, *MGME1*, *RRM2B,*

TYMP, and *POLG*. mtDNA depletion syndromes can be broken down into 5 subtypes based on their

constellation of phenotypes[74] , and with the exception of cardiomyopathic subtypes (associated with

mutations in *AGK* and *SLC25A4*), we were able to identify at least 1 gene from the other 4 subtypes,

suggesting that our mtDNA-CN measurement in blood-derived DNA can identify genes widely relevant

to non-blood phenotypes. This finding is consistent with a large body of work showing that mtDNA-CN

measured in blood is associated with numerous aging-related phenotypes for which the primary tissue

of interest is not blood (e.g. chronic kidney disease[13], heart failure[11], and diabetes[75]). Also consistent

with this finding is recent work demonstrating that mtDNA-CN measured in blood is associated with

mtRNA expression across numerous non-blood tissues, suggesting a link between mitochondrial

function measured in blood and other tissues[76].

In addition to identifying the mtDNA depletion syndrome genes directly linked to mitochondrial

DNA metabolic processes, DNA replication, and genome maintenance, we also identify genes which play

a role in mitochondrial function. The top GWAS hit is a missense mutation in *LONP1*, which encodes a

mitochondrial protease that has been shown to cause mitochondrial cytopathy and reduced respiratory

chain activity[77,78]. Interestingly, this missense mutation was recently found to be associated with

mitochondrial tRNA methylation levels[79]. Additional genes known to impact mitochondrial function

include *MFN1*, which encodes a mediator of mitochondrial fusion[80,81], *STMP1*, which plays a role in

mitochondrial respiration[82], and *MRPS35*, which encodes a ribosomal protein involved in protein

synthesis in the mitochondrion[83,84].

Using a combination of gene-based tests and gene prioritization using functional annotation,

pathway analyses reveal enrichment for numerous mitochondrial related pathways, as well as those

involved in regulation of cell differentiation ($p < 1.08 \times 10^{-5}$), homeostatic processes ($p < 3.77 \times 10^{-6}$), and

cellular response to stress ($p < 3.49 \times 10^{-6}$) (Supplemental Table 13). These results provide additional

evidence for the broad role played by mitochondria in numerous aspects of cellular function. Of

particular interest, the GO term for amyloid beta is significantly enriched, reinforcing a link between

mtDNA-CN and neurodegenerative disease[85–87]. Previous work from our lab using the UKB has shown

that increased mtDNA-CN is associated with lower rates of prevalent neurodegenerative disease, and is

predictive for decreased risk of incident neurodegenerative disease[76]. mtDNA-CN is also known to be

decreased in the frontal cortex of Alzheimer's disease (AD) patients[88]. Interestingly, the four GWAS-

identified genes driving the enrichment for amyloid-beta clearance are all related to regulation of lipid

levels, and lipid homeostasis within the brain is known to play an important role in Alzheimer's

disease[89]. *APOE*, one of the most well-known risk genes for Alzheimer's disease, is a cholesterol carrier

involved in lipid transport, and the ApoE-ε4 isoform involved in AD pathogenesis is associated with

mitochondrial dysfunction and oxidative distress in the human brain[90]; *CD36* is a platelet glycoprotein

which mediates the response to amyloid-beta accumulation[91]; *LDLR* is a low-density lipoprotein receptor

associated with AD[92]; and *ABCA7* is a phospholipid transporter[93]. *ABCA7* loss of function variants are

enriched in both AD and Parkinson's disease (PD) patients[94], suggesting a broad role across

neurodegenerative diseases.

642   Given the integral role of mitochondria in cellular function, from ATP formation and energy

643  production, signaling through reactive oxygen species, and apoptosis mediation, there is a strong basis

644  to *a priori* assume that genetic variants associated with mtDNA-CN are likely to be highly pleiotropic.

645  Indeed, mtDNA-CN itself is associated with numerous phenotypes (Supplemental Table 5). Through our

646  PHEWAS-based clustering approach using 41 mtDNA-CN associated phenotypes, we uncovered

647  phenotypic associations between three distinct clusters of GWS mtDNA-CN associated SNPs. Cluster 1

648  was characterized by increased MPV, PDW, and decreased PLT (note that measured MPV and PLT are

649  generally inversely correlated to maintain hemostasis), which are the hallmarks of platelet activation[69].

650  The link between platelets and mtDNA-CN has typically revolved around platelet count, as platelets have

651  functional mitochondria, but do not have a nucleus. Given that the mtDNA-CN measurement is the ratio

652  between mtDNA and nuclear DNA, increased platelets, all else being equal, would directly equate with

653  increased mtDNA-CN. We note that the mtDNA-CN metric used in this GWAS was adjusted for platelet

654  count, likely increasing the ability to detect variants that impact mtDNA-CN through increased platelet

655  activation. Examining the genes within this cluster suggests roles for actin formation and regulation

656  (*TPM4*, *PACSIN2*)[95,96] and vesicular transport and endocytic trafficking (*DNM3*, *EHD3*)[97,98] in platelet

657  activation.

658    Cluster 2 is most strongly enriched for SNPs in which the mtDNA-CN increasing allele is

659  associated with increased PLT/PCT and serum calcium/phosphate. Examining the genes assigned to the

660  cluster, we implicate megakaryocyte proliferation and proplatelet formation (*MYB*, *JAK2*)[70], and

661  apoptosis and autophagy (*BAK1*, *BCL2, TYMP*)[71]. Megakaryocytes are used to form proplatelets, and the

662  process includes an important role for both intra- and extracellular calcium levels[99]. A role for apoptosis,

663  and specifically *BCL2*, in proplatelet formation and platelet release has been suggested[100,101], however

664  work in mice has suggested that apoptosis does not play a direct role in these processes[102].

665  Nevertheless, apoptosis is important for platelet lifespan[103].

Cluster 3 was particularly challenging to interpret, given that no particular phenotype was

enriched relative to the non-cluster 3 SNPs. We note that this cluster appeared to be enriched for the

mtDNA depletion syndrome genes, containing 6/7 genes identified in the GWAS, and significantly

enriched for GO Terms related mitochondrial DNA. Additionally, genes in cluster 3 were significantly

enriched for low-density lipoprotein particle binding, suggesting a role for lipid homeostasis. Closer

inspection of cluster 3 genes reveals a number of genes known to be associated with lipid levels (*LIPC*,

CETP, *LDLR*, *APOE*). While lipids play a role in both energy metabolism (largely through fatty acids) and

cellular membrane formation, a link to mtDNA-CN and/or mitochondrial function is not well-established.

A strong rationale for the study of mtDNA-CN is the underlying assumption that it reflects

mitochondrial function and is readily measured, often from existing data. A serious complication to the

interpretation of the role of mitochondrial function in various traits has been use of blood-derived

measurements, which can be confounded by differences in cell counts across individuals. Mendelian

randomization has been widely used to infer causality between traits (e.g. LDL and CAD)[104], but is only

robust under conditions of little to no pleiotropy[105] and its power is a function of variance explained. For

mtDNA-CN, the extensive pleiotropy and small amount of variance explained of GWS variants (<1%)

prevents the use of traditional MR approaches. As an alternative approach, we analyzed associations

between mitochondrial DNA variants and mtDNA-CN associated phenotypes. Presumably, variants

located on the mitochondrial genome are only able to modify phenotypes through modulating

mitochondrial function, allowing for causal inference. Our analyses revealed significant relationships

between mitochondrial variants and creatinine, aspartate aminotransferase, MCV, and PCT. Creatinine

and aspartate aminotransferase are markers of kidney and liver function respectively, and supporting

these findings, mtDNA-CN has been linked to both chronic kidney disease[13] and non-alcoholic fatty liver

disease[106]. We also find a highly significant association between mitochondrial variation and non-cancer

mortality, adding evidence for a causal relationship to previous findings showing mtDNA-CN is

associated with all-cause mortality[9].

Several limitations should be noted. First, despite the large sample size and numerous loci

identified, we are likely missing a great deal of the true signal, as our SNP heritability estimates through

SumHer and BOLT-LMM were 7.0% and 7.4% respectively, while previous studies have estimated

mtDNA-CN heritability to be 65%[23]. Finally, while we have adjusted our mtDNA-CN metric for a variety of

confounders, it is important to note that mtDNA-CN can be influenced by a variety of environmental

factors including smoking[107] and drugs, which have not been adjusted for in these analyses.

In summary, we performed the largest-to-date GWAS for mtDNA-CN, including almost 500,000

individuals. We identified three distinct groups of SNPs associated with mtDNA-CN that are related to

platelet activation, megakaryocyte formation and apoptotic processes, and showed clear enrichment for

genes involved in mtDNA depletion and nucleotide regulation. Additionally, we find that mitochondrial

variants are significantly associated with creatinine, aspartate aminotransferase, MCV, and PCT,

implying a causal relationship between mitochondrial function and these phenotypes. Finally, we

provide strong evidence that mitochondrial function is causal for non-cancer mortality. Given the role of

mtDNA-CN, and, by proxy, mitochondrial function in aging-related disease, this work begins to unravel

the many varied underlying mechanisms through which mitochondrial function impacts human health.

Supplemental Data

Supplemental Data include six supplemental figures, twenty-two supplemental tables, and supplemental

methods.

Declaration of Interests

Psaty serves on the Steering Committee of the Yale Open Data Access Project funded by Johnson &

Johnson. All other authors declare no competing interests.

Acknowledgements

This work was supported by National Heart, Lung and Blood Institute, National Institutes of Health (NIH)

grants R01HL13573 and R01HL144569 (RJL, SYY, CAC, DEA), NIH grant P01-AG027734 (GA, YK, NB, AB)

and the National Center for Advancing Translational Sciences, NIH, through Grant KL2TR002490 (LMR).

The content is solely the responsibility of the authors and does not necessarily represent the official

views of the NIH. LMR was also funded by T32 HL129982.

This research was conducted using data from the Genotype-Tissue Expression (GTEx) project (dbGaP

accession: phs000424.**v8**.p2). The GTEx project was supported by the Common Fund of the Office of the

Director of the National Institutes of Health, and by NCI, NHGRI, NHLBI, NIDA, NIMH, and NINDS.

This research was also conducted using the UK Biobank Resource under Application Number 17731.

The Atherosclerosis Risk in Communities study (dbGaP accession: phs000280.**v7**.p1) has been funded in

whole or in part with Federal funds from the National Heart, Lung, and Blood Institute, National

Institutes of Health, Department of Health and Human Services (contract numbers HSN268201700001I,

HHSN268201700002I, HHSN268201700003I, HHSN268201700004I and HHSN268201700005I),

R01HL087641, R01HL059367 and R01HL086694; National Human Genome Research Institute contract

U01HG004402; and National Institutes of Health contract HHSN268200625226C. Funding support for

"Building on GWAS for NHLBI diseases: the U.S. CHARGE consortium" was provided by the NIH through

the American Recovery and Reinvestment Act of 2009 (ARRA) (5RC2HL102419). Sequencing was carried

out at the Baylor College of Medicine Human Genome Sequencing Center and supported by the National

Human Genome Research Institute grants U54 HG003273 and UM1 HG008898. The authors thank the

staff and participants of the ARIC study for their important contributions. Infrastructure was partly

supported by Grant Number UL1RR025005, a component of the National Institutes of Health and NIH

Roadmap for Medical Research.

This CHS research was supported by NHLBI contracts HHSN268201200036C,

HHSN268200800007C, HHSN268201800001C, N01HC55222, N01HC85079, N01HC85080, N01HC85081,

N01HC85082, N01HC85083, N01HC85086, 75N92021D00006; and NHLBI grants U01HL080295,

R01HL087652, R01HL105756, R01HL103612, R01HL120393, and U01HL130114 with additional

contribution from the National Institute of Neurological Disorders and Stroke (NINDS). Additional

support was provided through R01AG023629 from the National Institute on Aging (NIA). A full list of

principal CHS investigators and institutions can be found at CHS-NHLBI.org. The provision of genotyping

data was supported in part by the National Center for Advancing Translational Sciences, CTSI

grant UL1TR001881, and the National Institute of Diabetes and Digestive and Kidney Disease Diabetes

Research Center (DRC) grant DK063491 to the Southern California Diabetes Endocrinology Research

Center.

The FHS phenotype-genotype analyses were supported by the National Institute of Aging

(U34AG051418). This research was conducted in part using data and resources from the Framingham

Heart Study of the National Heart Lung and Blood Institute of the National Institutes of Health and

Boston University School of Medicine. This work was partially supported by the National Heart, Lung and

Blood Institute's Framingham Heart Study (Contract No. N01-HC-25195, HHSN268201500001) and its

contract with Affymetrix, Inc for genotyping services (Contract No. N02-HL-6-4278). Genotyping, quality

control and calling of the Illumina HumanExome BeadChip in the Framingham Heart Study was

supported by funding from the National Heart, Lung and Blood Institute Division of Intramural Research

(Daniel Levy and Christopher J. O'Donnell, Principle Investigators). The authors thank the participants for

their dedication to the study. The authors are pleased to acknowledge that the computational work

reported on in this paper was performed on the Shared Computing Cluster which is administered by

Boston University's Research Computing Services. URL: www.bu.edu/tech/support/research/.

MESA and the MESA SHARe projects (dbGaP accession: phs000209.**v13**.p3) are conducted and

supported by the National Heart, Lung, and Blood Institute (NHLBI) in collaboration with MESA

investigators. Support for MESA is provided by contracts 75N92020D00001, HHSN268201500003I, N01-

HC-95159, 75N92020D00005, N01-HC-95160, 75N92020D00002, N01-HC-95161, 75N92020D00003,

N01-HC-95162, 75N92020D00006, N01-HC-95163, 75N92020D00004, N01-HC-95164, 75N92020D00007,

N01-HC-95165, N01-HC-95166, N01-HC-95167, N01-HC-95168, N01-HC-95169, UL1-TR-000040, UL1-TR-

001079, and UL1-TR-001420. Funding for SHARe genotyping was provided by NHLBI Contract N02-HL-

64278. Genotyping was performed at Affymetrix (Santa Clara, California, USA) and the Broad Institute of

Harvard and MIT (Boston, Massachusetts, USA) using the Affymetrix Genome-Wide Human SNP Array

6.0. The provision of genotyping data was supported in part by the National Center for Advancing

Translational Sciences, CTSI grant UL1TR001881, and the National Institute of Diabetes and Digestive

and Kidney Disease Diabetes Research Center (DRC) grant DK063491 to the Southern California Diabetes

Endocrinology Research Center.

ROS/MAP is supported by the Translational Genomics Research Institute and National Institute on Aging

(NIA) through grants U01AG46152, U01AG61256, P30AG10161, R01AG17917, RF1AG15819,

R01AG30146.

SHIP and SHIP-TREND are part of the Community Medicine Research net of the University of Greifswald,

Germany, which is funded by the Federal Ministry of Education and Research (grants no. 01ZZ9603,

01ZZ0103, and 01ZZ0403), the Ministry of Cultural Affairs as well as the Social Ministry of the Federal State

of Mecklenburg-West Pomerania, and the network 'Greifswald Approach to Individualized Medicine

(GANI_MED)' funded by the Federal Ministry of Education and Research (grant 03IS2061A).

This research was funded in whole, or in part, by the Wellcome Trust [Grant ref: 217065/Z/19/Z]. For the

purpose of Open Access, the author has applied a CC BY public copyright license to any Author Accepted

Manuscript version arising from this submission.

Web Resources

REVIGO was accessed at http://revigo.irb.hr/.

Data and Code Availability

All data used in this manuscript is available through either the UKBiobank and CHARGE consortiums.

Code and scripts are available in a zipped file at https://www.arkinglab.org/resources/.

References

1. Wallace, D.C. (1992). Diseases of the Mitochondrial Dna. Annual Review of Biochemistry *61*, 1175–
1212.

2. Vakifahmetoglu-Norberg, H., Ouchida, A.T., and Norberg, E. (2017). The role of mitochondria in
metabolism and cell death. Biochem. Biophys. Res. Commun. *482*, 426–431.

3. Dai, D.-F., Rabinovitch, P.S., and Ungvari, Z. (2012). Mitochondria and cardiovascular aging. Circ. Res.
110, 1109–1124.

4. Cui, H., Kong, Y., and Zhang, H. (2012). Oxidative stress, mitochondrial dysfunction, and aging. J Signal
Transduct *2012*, 646354.

5. Herst, P.M., Rowe, M.R., Carson, G.M., and Berridge, M.V. (2017). Functional Mitochondria in Health
and Disease. Front Endocrinol (Lausanne) *8*, 296.

6. Liu, C.-S., Tsai, C.-S., Kuo, C.-L., Chen, H.-W., Lii, C.-K., Ma, Y.-S., and Wei, Y.-H. (2003). Oxidative stress-
related alteration of the copy number of mitochondrial DNA in human leukocytes. Free Radic Res *37*,
1307–1317.

7. Guha, M., and Avadhani, N.G. (2013). Mitochondrial retrograde signaling at the crossroads of tumor
bioenergetics, genetics and epigenetics. Mitochondrion *13*, 577–591.

8. Malik, A.N., and Czajka, A. (2013). Is mitochondrial DNA content a potential biomarker of
mitochondrial dysfunction? Mitochondrion *13*, 481–492.

9. Ashar, F.N., Moes, A., Moore, A.Z., Grove, M.L., Chaves, P.H.M., Coresh, J., Newman, A.B., Matteini,
A.M., Bandeen-Roche, K., Boerwinkle, E., et al. (2015). Association of mitochondrial DNA levels with
frailty and all-cause mortality. J. Mol. Med. *93*, 177–186.

10. Ashar, F.N., Zhang, Y., Longchamps, R.J., Lane, J., Moes, A., Grove, M.L., Mychaleckyj, J.C., Taylor,
K.D., Coresh, J., Rotter, J.I., et al. (2017). Association of Mitochondrial DNA Copy Number With
Cardiovascular Disease. JAMA Cardiol *2*, 1247–1255.

11. Hong, Y.S., Longchamps, R.J., Zhao, D., Castellani, C.A., Loehr, L.R., Chang, P.P., Matsushita, K., Grove,
M.L., Boerwinkle, E., Arking, D.E., et al. (2020). Mitochondrial DNA Copy Number and Incident Heart
Failure: The Atherosclerosis Risk in Communities (ARIC) Study. Circulation *141*, 1823–1825.

12. Zhao, D., Bartz, T.M., Sotoodehnia, N., Post, W.S., Heckbert, S.R., Alonso, A., Longchamps, R.J.,
Castellani, C.A., Hong, Y.S., Rotter, J.I., et al. (2020). Mitochondrial DNA copy number and incident atrial
fibrillation. BMC Medicine *18*, 246.

13. Tin, A., Grams, M.E., Ashar, F.N., Lane, J.A., Rosenberg, A.Z., Grove, M.L., Boerwinkle, E., Selvin, E.,
Coresh, J., Pankratz, N., et al. (2016). Association between Mitochondrial DNA Copy Number in
Peripheral Blood and Incident CKD in the Atherosclerosis Risk in Communities Study. J Am Soc Nephrol
27, 2467–2473.

14. Pyle, A., Anugrha, H., Kurzawa-Akanbi, M., Yarnall, A., Burn, D., and Hudson, G. (2016). Reduced
mitochondrial DNA copy number is a biomarker of Parkinson's disease. Neurobiol. Aging *38*, 216.e7-
216.e10.

15. Wei, W., Keogh, M.J., Wilson, I., Coxhead, J., Ryan, S., Rollinson, S., Griffin, H., Kurzawa-Akanbi, M.,
Santibanez-Koref, M., Talbot, K., et al. (2017). Mitochondrial DNA point mutations and relative copy
number in 1363 disease and control human brains. Acta Neuropathologica Communications *5*, 13.

16. Reznik, E., Miller, M.L., Şenbabaoğlu, Y., Riaz, N., Sarungbam, J., Tickoo, S.K., Al-Ahmadie, H.A., Lee,
W., Seshan, V.E., Hakimi, A.A., et al. (2016). Mitochondrial DNA copy number variation across human
cancers. ELife *5*, e10769.

17. Hurtado-Roca, Y., Ledesma, M., Gonzalez-Lazaro, M., Moreno-Loshuertos, R., Fernandez-Silva, P.,
Enriquez, J.A., and Laclaustra, M. (2016). Adjusting MtDNA Quantification in Whole Blood for Peripheral
Blood Platelet and Leukocyte Counts. PLOS ONE *11*, e0163770.

18. Knez, J., Winckelmans, E., Plusquin, M., Thijs, L., Cauwenberghs, N., Gu, Y., Staessen, J.A., Nawrot,
T.S., and Kuznetsova, T. (2016). Correlates of Peripheral Blood Mitochondrial DNA Content in a General
Population. Am J Epidemiol *183*, 138–146.

19. Kumar, P., Efstathopoulos, P., Millischer, V., Olsson, E., Wei, Y.B., Brüstle, O., Schalling, M.,
Villaescusa, J.C., Ösby, U., and Lavebratt, C. (2018). Mitochondrial DNA copy number is associated with
psychosis severity and anti-psychotic treatment. Sci Rep *8*, 12743.

20. Urata, M., Koga-Wada, Y., Kayamori, Y., and Kang, D. (2008). Platelet contamination causes large
variation as well as overestimation of mitochondrial DNA content of peripheral blood mononuclear cells.
Ann Clin Biochem *45*, 513–514.

21. Clay Montier, L.L., Deng, J.J., and Bai, Y. (2009). Number matters: control of mammalian
mitochondrial DNA copy number. J Genet Genomics *36*, 125–131.

22. Tang, Y., Schon, E.A., Wilichowski, E., Vazquez-Memije, M.E., Davidson, E., and King, M.P. (2000).
Rearrangements of Human Mitochondrial DNA (mtDNA): New Insights into the Regulation of mtDNA
Copy Number and Gene Expression. Mol Biol Cell *11*, 1471–1485.

23. Xing, J., Chen, M., Wood, C.G., Lin, J., Spitz, M.R., Ma, J., Amos, C.I., Shields, P.G., Benowitz, N.L., Gu,
J., et al. (2008). Mitochondrial DNA content: its genetic heritability and association with renal cell
carcinoma. J. Natl. Cancer Inst. *100*, 1104–1112.

24. Carling, P.J., Cree, L.M., and Chinnery, P.F. (2011). The implications of mitochondrial DNA copy
number regulation during embryogenesis. Mitochondrion *11*, 686–692.

25. Harvey, A., Gibson, T., Lonergan, T., and Brenner, C. (2011). Dynamic regulation of mitochondrial
function in preimplantation embryos and embryonic stem cells. Mitochondrion *11*, 829–838.

26. Copeland, W.C. (2014). Defects of Mitochondrial DNA Replication. J Child Neurol *29*, 1216–1224.

27. Mandel, H., Szargel, R., Labay, V., Elpeleg, O., Saada, A., Shalata, A., Anbinder, Y., Berkowitz, D.,
Hartman, C., Barak, M., et al. (2001). The deoxyguanosine kinase gene is mutated in individuals with
depleted hepatocerebral mitochondrial DNA. Nat. Genet. *29*, 337–341.

28. Wang, L., Limongelli, A., Vila, M.R., Carrara, F., Zeviani, M., and Eriksson, S. (2005). Molecular insight
into mitochondrial DNA depletion syndrome in two patients with novel mutations in the
deoxyguanosine kinase and thymidine kinase 2 genes. Mol. Genet. Metab. *84*, 75–82.

29. Rusecka, J., Kaliszewska, M., Bartnik, E., and Tońska, K. (2018). Nuclear genes involved in
mitochondrial diseases caused by instability of mitochondrial DNA. J Appl Genet *59*, 43–57.

30. Cai, N., Li, Y., Chang, S., Liang, J., Lin, C., Zhang, X., Liang, L., Hu, J., Chan, W., Kendler, K.S., et al.
(2015). Genetic Control over mtDNA and Its Relationship to Major Depressive Disorder. Curr Biol *25*,
3170–3177.

31. Workalemahu, T., Enquobahrie, D.A., Tadesse, M.G., Hevner, K., Gelaye, B., Sanchez, S., and
Williams, M.A. (2017). Genetic Variations Related to Maternal Whole Blood Mitochondrial DNA Copy
Number: A Genome-Wide and Candidate Gene Study. J Matern Fetal Neonatal Med *30*, 2433–2439.

32. Guyatt, A.L., Brennan, R.R., Burrows, K., Guthrie, P.A.I., Ascione, R., Ring, S.M., Gaunt, T.R., Pyle, A.,
Cordell, H.J., Lawlor, D.A., et al. (2019). A genome-wide association study of mitochondrial DNA copy
number in two population-based cohorts. Human Genomics *13*, 6.

33. Hägg, S., Jylhävä, J., Wang, Y., Czene, K., and Grassmann, F. (2020). Deciphering the genetic and
epidemiological landscape of mitochondrial DNA abundance. Hum Genet.

34. Psaty Bruce M., O'Donnell Christopher J., Gudnason Vilmundur, Lunetta Kathryn L., Folsom Aaron R.,
Rotter Jerome I., Uitterlinden André G., Harris Tamara B., Witteman Jacqueline C.M., and Boerwinkle
Eric (2009). Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) Consortium.
Circulation: Cardiovascular Genetics *2*, 73–80.

35. Bycroft, C., Freeman, C., Petkova, D., Band, G., Elliott, L.T., Sharp, K., Motyer, A., Vukcevic, D.,
Delaneau, O., O'Connell, J., et al. (2018). The UK Biobank resource with deep phenotyping and genomic
data. Nature *562*, 203–209.

36. Longchamps, R.J., Castellani, C.A., Yang, S.Y., Newcomb, C.E., Sumpter, J.A., Lane, J., Grove, M.L.,
Guallar, E., Pankratz, N., Taylor, K.D., et al. (2020). Evaluation of mitochondrial DNA copy number
estimation techniques. PLOS ONE *15*, e0228166.

37. MitoPipeline: Generating Mitochondrial copy number estimates from SNP array data in Genvisis.

38. Han, B., and Eskin, E. (2011). Random-Effects Model Aimed at Discovering Associations in Meta-
Analysis of Genome-wide Association Studies. The American Journal of Human Genetics *88*, 586–598.

39. Willer, C.J., Li, Y., and Abecasis, G.R. (2010). METAL: fast and efficient meta-analysis of genomewide
association scans. Bioinformatics *26*, 2190–2191.

40. Loh, P.-R., Tucker, G., Bulik-Sullivan, B.K., Vilhjálmsson, B.J., Finucane, H.K., Salem, R.M., Chasman,
D.I., Ridker, P.M., Neale, B.M., Berger, B., et al. (2015). Efficient Bayesian mixed model analysis increases
association power in large cohorts. Nat Genet *47*, 284–290.

41. Speed, D. (2019). SumHer better estimates the SNP heritability of complex traits from summary
statistics. Nature Genetics *51*, 12.

42. Wang, G., Sarkar, A., Carbonetto, P., and Stephens, M. (2020). A simple new approach to variable
selection in regression, with application to genetic fine mapping. Journal of the Royal Statistical Society:
Series B (Statistical Methodology) *82*, 1273–1300.

43. Wang, K., Li, M., and Hakonarson, H. (2010). ANNOVAR: functional annotation of genetic variants
from high-throughput sequencing data. Nucleic Acids Research *38*, e164–e164.

44. Sherry, S.T., Ward, M., and Sirotkin, K. (1999). dbSNP-database for single nucleotide polymorphisms
and other classes of minor genetic variation. Genome Res *9*, 677–679.

45. O'Leary, N.A., Wright, M.W., Brister, J.R., Ciufo, S., Haddad, D., McVeigh, R., Rajput, B., Robbertse, B.,
Smith-White, B., Ako-Adjei, D., et al. (2016). Reference sequence (RefSeq) database at NCBI: current
status, taxonomic expansion, and functional annotation. Nucleic Acids Res *44*, D733-745.

46. Giambartolomei, C., Vukcevic, D., Schadt, E.E., Franke, L., Hingorani, A.D., Wallace, C., and Plagnol, V.
(2014). Bayesian Test for Colocalisation between Pairs of Genetic Association Studies Using Summary
Statistics. PLOS Genetics *10*, e1004383.

47. Võsa, U., Claringbould, A., Westra, H.-J., Bonder, M.J., Deelen, P., Zeng, B., Kirsten, H., Saha, A.,
Kreuzhuber, R., Kasela, S., et al. (2018). Unraveling the polygenic architecture of complex traits using
blood eQTL metaanalysis. BioRxiv 447367.

48. Pers, T.H., Karjalainen, J.M., Chan, Y., Westra, H.-J., Wood, A.R., Yang, J., Lui, J.C., Vedantam, S.,
Gustafsson, S., Esko, T., et al. (2015). Biological interpretation of genome-wide association studies using
predicted gene functions. Nature Communications *6*, 5890.

49. Smyth, G., Hu, Y., Ritchie, M., Silver, J., Wettenhall, J., McCarthy, D., Wu, D., Shi, W., Phipson, B., Lun,
A., et al. (2021). limma: Linear Models for Microarray Data (Bioconductor version: Release (3.12)).

50. Young, M.D., Wakefield, M.J., Smyth, G.K., and Oshlack, A. (2010). Gene ontology analysis for RNA-
seq: accounting for selection bias. Genome Biology *11*, R14.

51. Barbeira, A.N., Dickinson, S.P., Bonazzola, R., Zheng, J., Wheeler, H.E., Torres, J.M., Torstenson, E.S.,
Shah, K.P., Garcia, T., Edwards, T.L., et al. (2018). Exploring the phenotypic consequences of tissue
specific gene expression variation inferred from GWAS summary statistics. Nat Commun *9*, 1825.

52. Barbeira, A.N., Pividori, M., Zheng, J., Wheeler, H.E., Nicolae, D.L., and Im, H.K. (2019). Integrating
predicted transcriptome from multiple tissues improves association detection. PLOS Genetics *15*,
e1007889.

53. Supek, F., Bošnjak, M., Škunca, N., and Šmuc, T. (2011). REVIGO summarizes and visualizes long lists
of gene ontology terms. PLoS One *6*, e21800.

54. Stiles, A.R., Simon, M.T., Stover, A., Eftekharian, S., Khanlou, N., Wang, H.L., Magaki, S., Lee, H.,
Partynski, K., Dorrani, N., et al. (2016). Mutations in TFAM, encoding mitochondrial transcription factor
A, cause neonatal liver failure associated with mtDNA depletion. Mol Genet Metab *119*, 91–99.

55. Kornblum, C., Nicholls, T.J., Haack, T.B., Schöler, S., Peeva, V., Danhauser, K., Hallmann, K., Zsurka, G.,
Rorbach, J., Iuso, A., et al. (2013). Loss-of-function mutations in MGME1 impair mtDNA replication and
cause multisystemic mitochondrial disease. Nat Genet *45*, 214–219.

56. El-Hattab, A.W., and Scaglia, F. (2013). Mitochondrial DNA depletion syndromes: review and updates
of genetic basis, manifestations, and therapeutic options. Neurotherapeutics *10*, 186–198.

57. Millard, L.A.C., Davies, N.M., Gaunt, T.R., Davey Smith, G., and Tilling, K. (2018). Software Application
Profile: PHESANT: a tool for performing automated phenome scans in UK Biobank. Int J Epidemiol *47*,
29–35.

58. Konopka, T. (2020). umap: Uniform Manifold Approximation and Projection.

59. Hahsler, M., Piekenbrock, M., Arya, S., and Mount, D. (2019). dbscan: Density Based Clustering of
Applications with Noise (DBSCAN) and Related Algorithms.

60. Delaneau, O., Zagury, J.-F., Robinson, M.R., Marchini, J.L., and Dermitzakis, E.T. (2019). Accurate,
scalable and integrative haplotype estimation. Nat Commun *10*,.

61. Rubinacci, S., Delaneau, O., and Marchini, J. (2020). Genotype imputation using the Positional
Burrows Wheeler Transform. PLOS Genetics *16*, e1009049.

62. Auton, A., Abecasis, G.R., Altshuler, D.M., Durbin, R.M., Abecasis, G.R., Bentley, D.R., Chakravarti, A.,
Clark, A.G., Donnelly, P., Eichler, E.E., et al. (2015). A global reference for human genetic variation.
Nature *526*, 68–74.

63. Shen, L., Attimonelli, M., Bai, R., Lott, M.T., Wallace, D.C., Falk, M.J., and Gai, X. (2018). MSeqDR
mvTool: A mitochondrial DNA Web and API resource for comprehensive variant annotation, universal
nomenclature collation, and reference genome conversion. Hum Mutat *39*, 806–810.

64. Gibellini, L., De Gaetano, A., Mandrioli, M., Van Tongeren, E., Bortolotti, C.A., Cossarizza, A., and
Pinti, M. (2020). The biology of Lonp1: More than a mitochondrial protease. Int Rev Cell Mol Biol *354*, 1–
61.

65. GTEx Consortium (2013). The Genotype-Tissue Expression (GTEx) project. Nat Genet *45*, 580–585.

66. Liu, T., Lu, B., Lee, I., Ondrovicová, G., Kutejová, E., and Suzuki, C.K. (2004). DNA and RNA binding by
the mitochondrial lon protease is regulated by nucleotide and protein substrate. J Biol Chem *279*,
13902–13910.

67. Sharma, M.R., Koc, E.C., Datta, P.P., Booth, T.M., Spremulli, L.L., and Agrawal, R.K. (2003). Structure
of the mammalian mitochondrial ribosome reveals an expanded functional role for its component
proteins. Cell *115*, 97–108.

68. Shimizu, S., Narita, M., and Tsujimoto, Y. (1999). Bcl-2 family proteins regulate the release of
apoptogenic cytochrome c by the mitochondrial channel VDAC. Nature *399*, 483–487.

69. Vagdatli, E., Gounari, E., Lazaridou, E., Katsibourlia, E., Tsikopoulou, F., and Labrianou, I. (2010).
Platelet distribution width: a simple, practical and specific marker of activation of coagulation.
Hippokratia *14*, 28–32.

70. PathCards :: Factors involved in megakaryocyte development and platelet production Pathway and
related pathways.

71. PathCards :: Apoptosis and Autophagy Pathway and related pathways.

72. Yuan, Y., Ju, Y.S., Kim, Y., Li, J., Wang, Y., Yoon, C.J., Yang, Y., Martincorena, I., Creighton, C.J.,
Weinstein, J.N., et al. (2020). Comprehensive molecular characterization of mitochondrial genomes in
human cancers. Nature Genetics *52*, 342–352.

73. Mizumachi, T., Muskhelishvili, L., Naito, A., Furusawa, J., Fan, C.-Y., Siegel, E.R., Kadlubar, F.F., Kumar,
U., and Higuchi, M. (2008). Increased Distributional Variance of Mitochondrial DNA Content Associated
With Prostate Cancer Cells as Compared With Normal Prostate Cells. Prostate *68*, 408–417.

74. Basel, D. (2020). Mitochondrial DNA Depletion Syndromes. Clin Perinatol *47*, 123–141.

75. DeBarmore, B., Longchamps, R.J., Zhang, Y., Kalyani, R.R., Guallar, E., Arking, D.E., Selvin, E., and
Young, J.H. (2020). Mitochondrial DNA copy number and diabetes: the Atherosclerosis Risk in
Communities (ARIC) study. BMJ Open Diabetes Res Care *8*,.

76. Yang, S.Y., Castellani, C.A., Longchamps, R.J., Pillalamarri, V.K., O'Rourke, B., Guallar, E., and Arking,
D.E. (2021). Blood-derived mitochondrial DNA copy number is associated with gene expression across
multiple tissues and is predictive for incident neurodegenerative disease. Genome Res.

77. Hannah-Shmouni, F., MacNeil, L., Brady, L., Nilsson, M.I., and Tarnopolsky, M. (2019). Expanding the
Clinical Spectrum of LONP1-Related Mitochondrial Cytopathy. Front Neurol *10*, 981.

78. Grainha, T.R.R., Jorge, P.A. da S., Pérez-Pérez, M., Pérez Rodríguez, G., Pereira, M.O.B.O., and
Lourenço, A.M.G. (2018). Exploring anti-quorum sensing and anti-virulence based strategies to fight
Candida albicans infections: an in silico approach. FEMS Yeast Res *18*,.

79. Ali, A.T., Idaghdour, Y., and Hodgkinson, A. (2020). Analysis of mitochondrial m1A/G RNA
modification reveals links to nuclear genetic variants and associated disease processes. Communications
Biology *3*, 1–11.

80. Schrepfer, E., and Scorrano, L. (2016). Mitofusins, from Mitochondria to Metabolism. Mol Cell *61*,
683–694.

81. Ishihara, N., Eura, Y., and Mihara, K. (2004). Mitofusin 1 and 2 play distinct roles in mitochondrial
fusion reactions via GTPase activity. J Cell Sci *117*, 6535–6546.

82. Zhang, D., Xi, Y., Coccimiglio, M.L., Mennigen, J.A., Jonz, M.G., Ekker, M., and Trudeau, V.L. (2012).
Functional prediction and physiological characterization of a novel short trans-membrane protein 1 as a
subunit of mitochondrial respiratory complexes. Physiol Genomics *44*, 1133–1140.

83. Cavdar Koc, E., Burkhart, W., Blackburn, K., Moseley, A., and Spremulli, L.L. (2001). The small subunit
of the mammalian mitochondrial ribosome. Identification of the full complement of ribosomal proteins
present. J Biol Chem *276*, 19363–19374.

84. Márquez-Jurado, S., Díaz-Colunga, J., das Neves, R.P., Martinez-Lorente, A., Almazán, F., Guantes, R.,
and Iborra, F.J. (2018). Mitochondrial levels determine variability in cell death by modulating apoptotic
gene expression. Nat Commun *9*, 389.

85. Dölle, C., Flønes, I., Nido, G.S., Miletic, H., Osuagwu, N., Kristoffersen, S., Lilleng, P.K., Larsen, J.P.,
Tysnes, O.-B., Haugarvoll, K., et al. (2016). Defective mitochondrial DNA homeostasis in the substantia
nigra in Parkinson disease. Nat Commun *7*, 13548.

86. Chen, C., Turnbull, D.M., and Reeve, A.K. (2019). Mitochondrial Dysfunction in Parkinson's Disease-
Cause or Consequence? Biology (Basel) *8*,.

87. Pinto, M., and Moraes, C.T. (2014). Mitochondrial genome changes and neurodegenerative diseases.
Biochim Biophys Acta *1842*, 1198–1207.

88. Rodríguez-Santiago, B., Casademont, J., and Nunes, V. (2001). Is mitochondrial DNA depletion
involved in Alzheimer's disease? Eur J Hum Genet *9*, 279–285.

89. Chew, H., Solomon, V.A., and Fonteh, A.N. (2020). Involvement of Lipids in Alzheimer's Disease
Pathology and Potential Therapies. Front Physiol *11*, 598.

90. Yin, J., Reiman, E.M., Beach, T.G., Serrano, G.E., Sabbagh, M.N., Nielsen, M., Caselli, R.J., and Shi, J.
(2020). Effect of ApoE isoforms on mitochondria in Alzheimer disease. Neurology *94*, e2404–e2411.

91. El Khoury, J.B., Moore, K.J., Means, T.K., Leung, J., Terada, K., Toft, M., Freeman, M.W., and Luster,
A.D. (2003). CD36 mediates the innate host response to beta-amyloid. J Exp Med *197*, 1657–1666.

92. Lämsä, R., Helisalmi, S., Herukka, S.-K., Tapiola, T., Pirttilä, T., Vepsäläinen, S., Hiltunen, M., and
Soininen, H. (2008). Genetic study evaluating LDLR polymorphisms and Alzheimer's disease. Neurobiol
Aging *29*, 848–855.

93. Tomioka, M., Toda, Y., Mañucat, N.B., Akatsu, H., Fukumoto, M., Kono, N., Arai, H., Kioka, N., and
Ueda, K. (2017). Lysophosphatidylcholine export by human ABCA7. Biochim Biophys Acta Mol Cell Biol
Lipids *1862*, 658–665.

94. Nuytemans, K., Maldonado, L., Ali, A., John-Williams, K., Beecham, G.W., Martin, E., Scott, W.K., and
Vance, J.M. (2016). Overlap between Parkinson disease and Alzheimer disease in ABCA7 functional
variants. Neurol Genet *2*, e44.

95. Crabos, M., Yamakado, T., Heizmann, C.W., Cerletti, N., Bühler, F.R., and Erne, P. (1991). The calcium
binding protein tropomyosin in human platelets and cardiac tissue: elevation in hypertensive cardiac
hypertrophy. Eur J Clin Invest *21*, 472–478.

96. Kostan, J., Salzer, U., Orlova, A., Törö, I., Hodnik, V., Senju, Y., Zou, J., Schreiner, C., Steiner, J.,
Meriläinen, J., et al. (2014). Direct interaction of actin filaments with F-BAR protein pacsin2. EMBO Rep
15, 1154–1162.

97. Sever, S. (2002). Dynamin and endocytosis. Curr Opin Cell Biol *14*, 463–467.

98. Cai, B., Giridharan, S.S.P., Zhang, J., Saxena, S., Bahl, K., Schmidt, J.A., Sorgen, P.L., Guo, W.,
Naslavsky, N., and Caplan, S. (2013). Differential roles of C-terminal Eps15 homology domain proteins as
vesiculators and tubulators of recycling endosomes. J Biol Chem *288*, 30172–30180.

99. Di Buduo, C.A., Moccia, F., Battiston, M., De Marco, L., Mazzucato, M., Moratti, R., Tanzi, F., and
Balduini, A. (2014). The importance of calcium in the regulation of megakaryocyte function.
Haematologica *99*, 769–778.

100. De Botton, S., Sabri, S., Daugas, E., Zermati, Y., Guidotti, J.E., Hermine, O., Kroemer, G.,
Vainchenker, W., and Debili, N. (2002). Platelet formation is the consequence of caspase activation
within megakaryocytes. Blood *100*, 1310–1317.

101. Josefsson, E.C., James, C., Henley, K.J., Debrincat, M.A., Rogers, K.L., Dowling, M.R., White, M.J.,
Kruse, E.A., Lane, R.M., Ellis, S., et al. (2011). Megakaryocytes possess a functional intrinsic apoptosis
pathway that must be restrained to survive and produce platelets. J Exp Med *208*, 2017–2031.

102. Josefsson, E.C., Burnett, D.L., Lebois, M., Debrincat, M.A., White, M.J., Henley, K.J., Lane, R.M.,
Moujalled, D., Preston, S.P., O'Reilly, L.A., et al. (2014). Platelet production proceeds independently of
the intrinsic and extrinsic apoptosis pathways. Nat Commun *5*, 3455.

103. McArthur, K., Chappaz, S., and Kile, B.T. (2018). Apoptosis in megakaryocytes and platelets: the life
and death of a lineage. Blood *131*, 605–610.

104. Linsel-Nitschke, P., Götz, A., Erdmann, J., Braenne, I., Braund, P., Hengstenberg, C., Stark, K.,
Fischer, M., Schreiber, S., El Mokhtari, N.E., et al. (2008). Lifelong reduction of LDL-cholesterol related to
a common variant in the LDL-receptor gene decreases the risk of coronary artery disease--a Mendelian
Randomisation study. PLoS One *3*, e2986.

105. Lawlor, D.A., Harbord, R.M., Sterne, J.A.C., Timpson, N., and Davey Smith, G. (2008). Mendelian
randomization: using genes as instruments for making causal inferences in epidemiology. Stat Med *27*,
1133–1163.

106. Sookoian, S., Rosselli, M.S., Gemma, C., Burgueño, A.L., Gianotti, T.F., Castaño, G.O., and Pirola, C.J.
(2010). Epigenetic regulation of insulin resistance in nonalcoholic fatty liver disease: Impact of liver
methylation of the peroxisome proliferator–activated receptor γ coactivator 1α promoter. Hepatology
52, 1992–2000.

107. Vyas, C.M., Ogata, S., Reynolds, C.F., Mischoulon, D., Chang, G., Cook, N.R., Manson, J.E., Crous-
Bou, M., De Vivo, I., and Okereke, O.I. (2020). Lifestyle and behavioral factors and mitochondrial DNA
copy number in a diverse cohort of mid-life and older adults. PLoS One *15*, e0237235.

1074

1075   **Figures**

1076   **Figure 1. Manhattan plot of GWS loci from UKB-only analyses.**

1077

1078

1079   Manhattan plot showing genome-wide significant loci for the UK Biobank-only analyses.

1080

1081 **Figure 2. Scatterplot displaying effect size estimates between Whites and Blacks GWAS results for the**
1082 **129 autosomal SNPs identified in the Whites analyses.**

1083

1084 Scatterplot showing comparison between effect size estimates for White and Black individuals. Color
1085 represents significance of effect for each locus in Blacks GWAS analyses.

1086

1087

1088   **Figure 3. Volcano plot of genes whose predicted gene expression is significantly associated with**
1089   **mtDNA-CN.**

1090

1091   Volcano plot showing genes whose predicted gene expression is significantly associated with mtDNA-CN.
1092   Red indicates positive associations, blue indicates negative associations. Three genes (ARRDC1, EHMT1,
1093   PNPLA7) had extreme effect size estimates greater than 0.3 but were non-significant and removed from
1094   the plot for readability.

1095

1096   **Figure 4. PHEWAS-based clustering of mtDNA-CN associated SNPs.**

1097
1098
1099   **UMAP clusters created from PHEWAS associations for mtDNA-CN associated SNPs. (A) Three clusters**
1100   **were identified as labeled in the panel; orange indicates no cluster. (B-F) SNPs are colored based on**
1101   **their effect estimate size, standardized to the effect on mtDNA-CN (red = positive, blue = negative**
1102   **estimates), for (B) platelet count, (C) mean platelet volume, (D) platelet distribution width, (E) serum**
1103   **calcium levels, (F) mean spherical cellular volume.**
1104

 Figure 5. Associations between mtDNA-CN associated phenotypes and mitochondrial haplotypes.

1106
Mitochondrial haplotypes are significantly associated with mtDNA-CN associated traits, implying causal relationships between mitochondrial function and traits of interest. Haplotypes are notated in the following format: MT73_MT12612_MT7028_MT10238_MT13617_MT15257.

1110
1111

Figure 6. Associations between mitochondrial DNA haplotypes and morbidity (overall, cancer, and

noncancer)

 Mitochondrial haplotypes are significantly associated with overall morbidity, in particular, non-cancer
 mortality. Haplotypes are notated in the following format: MT73_ MT7028_MT10238_
 MT12612_MT13617_MT15257.