Estimating the fundamental niche: accounting for the uneven availability of existing climates

Jiménez, L.[1,*] and Soberón, J.[2]

[1]School of Life Sciences, University of Hawaii, Honolulu, HI 96822, USA

[2]Biodiversity Institute, University of Kansas, 1345 Jayhawk Blvd, Lawrence, KS 66045, USA

[*]Corresponding author: laurajj@hawaii.edu

March 16, 2021

Abstract

Studies that question important conceptual and methodological aspects of the field of ecological niche modeling (and species distribution modeling) have cast doubts on the validity of existing methodologies. One concern that has been broadly discussed is whether it is possible to estimate the fundamental niche of a species using presence data. Although it is well-recognized that the main limitation to doing so is that presence data come from the realized niche, which is only a subset of the fundamental niche, most of the existing methods for niche estimation lack the ability to overcome this limitation and therefore fit niches that resemble the realized niche. To obtain a more accurate estimate of the fundamental niche, we propose using the geographic region that is accessible to a species (based on its dispersal ability) to determine a sampling distribution in environmental space from which we can quantify the likelihood of observing a particular environmental combination in a sample of presence points. We incorporate this sampling distribution into a multivariate normal model (i.e., a Mahalanobis model) by creating a weight function that modifies the probabilities of observing different environmental combinations in a sample of presences. This modification accounts for the uneven availability of environmental conditions. We show that the parameters of this modified, weighted-normal model can be approximated with a maximum likelihood estimation approach; and then used to draw ellipsoids (confidence regions) that represent the fundamental niche of the species. We illustrate the application of

our model with two worked examples. First, we use presence data for an invasive species and an accessible area that includes only its native range to evaluate whether the fitted model predicts confirmed establishments of the species outside its native range. Second, we use presence data for closely related species with known accessible areas to demonstrate how the different dispersal abilities of the species constrain a classic Mahalanobis model. Overall, we show that accounting for the distribution of environmental conditions that are accessible to a species indeed affects the estimation of the ellipsoids used to model its fundamental niche.

Keywords: *fundamental niche; realized niche; environmental space; presence data; weighted distribution; accessible area*

1 Introduction

In recent years, there has been substantial progress in the fields of ecological niche modeling (ENM) and species distribution modeling (SDM) (Guisan et al., 2013). However, there is still debate about what aspects of the niche are estimated by these methods (Jiménez-valverde et al., 2008; Lobo, 2008; Warren, 2012). Specifically, conventional ENM/SDM approaches that are based on presence-only data estimate objects that, in the case of ENM, are between the realized and the fundamental niches (Peterson, Soberón, Pearson, et al., 2011), or, in the case of SDM, are between the actual and potential distributions of the species (Jiménez-valverde et al., 2008). Here, we propose a model to estimating fundamental niches that fits biologically realistic shapes and quantifies the availability of climates that are accessible for the species, as a way to account for biased presence samples.

The distinction between the fundamental and the realized niche, as proposed by Hutchinson (1957), is essential to understand what kind of objects are being estimated by the different correlative statistical models used in ENM/SDM. The fundamental niche of a species is the set of environmental conditions where, in the absence of biotic interactions, the population growth rate is positive (Peterson, Soberón, Pearson, et al., 2011). The realized niche is a subset of the fundamental niche that is determined by abiotic factors (environmental conditions), biotic factors, and dispersal limitations (Soberón, 2007). Estimating the fundamental niche of a species is of particular importance when using the estimated niche to model species distributions at other times or in different regions, such as when using ENM/SDM to predict the effects of climate change or the spread of invasive species (Tingley et al., 2014). However, estimating the fundamental niche of a species is also substantially more difficult than estimating the realized niche,

and it requires experimental data on the physiology of the species (Hoogenboom & Connolly, 2009; Jiménez et al., 2019).

The relationship between modeling niches and modeling geographic distributions is mediated by Hutchinson's duality (Colwell & Rangel, 2009), which is the relationship between environmental and geographic spaces. With the right resolution, a discrete set of geographic coordinates can be made to have a one-to-one relationship with a discrete set of environmental vectors (Aspinall & Lees, 1994; Soberón & Nakamura, 2009). This fundamental correspondence allows us to move back and forth between modeling niches and modeling geographic distributions.

As a consequence of Hutchinson's duality (Colwell & Rangel, 2009), and because species presence data come only from areas currently occupied by a species, a sample of presence records may not reflect all the environmental potentiality of a species (Jiménez-valverde et al., 2008; Lobo, 2008), and a niche estimated from geographic presence data therefore probably approximates the realized niche (Soberón & Nakamura, 2009). This means that correlative models designed to estimate the fundamental niche of a species using only presence data will be constrained by the imposed limitations of the set of environments where the species can be observed (Owens, Ribeiro, et al., 2020). Failing to acknowledge and (somehow) include this constraint in a model leads to severe uncertainties and drawbacks when trying to predict the effect of climate change on species distributions (Lobo, 2016) or on possible invasion scenarios for invasive species.

Species presence/absence data are often spatially biased and noisy. Several techniques have been developed for resolving some of the common types of problems in presence data (Chapman, 2005), such as a lack of accuracy in the reported coordinates, nomenclatural and taxonomic errors, and the presence of geographic or environmental outliers. We work under the assumption that a cleaning and preparation process to resolve these errors precedes the application of ENM/SDM methods. However, other types of bias remain in presence data. Here, we focus on the implicit bias created when defining the relevant spatial region for a study. Selecting different study regions produces different sampling universes, and the definition of the sampling region is an important part of the specification of a model. We explicitly define an "M" hypothesis to set the sampling universe as all the sites where we can observe the species as present (N. Barve et al., 2011). Under the"biotic-abiotic-movement" (BAM) framework (Peterson, Soberón, Pearson, et al., 2011), the region M contains all the sites that a species is hypothesized to have been able to reach from some past time. Not all sites in M have the environmental conditions needed to sustain viable populations, and it is known that some sites inside M could be sink

populations (N. Barve et al., 2011).

In geography (G-space), M is usually a connected and continuous set (i.e., a single polygon) describing a region of space that a species can reach via dispersal and migration. To perform practical computations, this set is first converted into a discrete grid of coordinates, and then the values for environmental variables at these coordinates are used to build an environmental space (E-space). Because the fundamental niche is defined in E-space, it makes sense to estimate niches in E-space. Hutchinson's duality permits the establishment of a one-to-one relationship between coordinates in G-space and multidimensional vectors in E-space, but this relationship does not preserve distance: nearby elements in G-space are not necessarily nearby in E-space, and vice versa. This creates a serious, and mostly ignored problem: uniform, random sampling in G-space does not imply uniform, random sampling in E-space.

It is important to stress that most ENM/SDM applications implicitly assume that any (relevant) combination of environmental conditions could be observed in a sample of species presences. However, recent studies have shown that acknowledging the effect of M on the sample can improve the performance of SDMs (Cooper & Soberón, 2018; Owens, Ribeiro, et al., 2020; Saupe, N. Barve, et al., 2017), although these studies primarily emphasized the modeling of geographical distributions. The key lesson from these works is that M limits the presence of a species to a discrete set of multivariate environmental combinations, and the probabilities of sampling different environments form this set are not uniform. If the goal of ENM/SDM is to get a better estimate closer of the fundamental niche using presence-only data, the empirical distribution of M in E-space (i.e., the distribution of environments that are accessible to the species) should therefore be incorporated in the statistical model to account for the uneven distribution of available sampling points. In other words, the E-space is not sampled uniformly when using presence data because M imposes biases in E.

In a previous contribution (Jiménez et al., 2019), we proposed a Bayesian argument for combining correlative techniques with information from physiological experiments to approximate to the fundamental niche of a species. However, in doing so, we noticed that our approach was limited by the fact that environmental combinations were not uniformly available in E-space. In this work, we consider the situation where the existing environmental combinations in E-space have different probabilities of being recorded as an environment where the species is present, and these probabilities are determined by M. We specifically consider the event of observing an environmental combination $\boldsymbol{x} \in \mathbb{R}^d$ (where the species was recorded as present) with a certain

probability of being included in the sample. In a typical random sampling scenario on the random variable $\boldsymbol{X}$, with a probability density function (pdf) $f(\boldsymbol{x}; \theta)$, the probability of selection is the same for each environmental combination, regardless of the value of $\boldsymbol{x}$. The pdf at $\boldsymbol{x}$ is therefore $f(\boldsymbol{x}; \theta)$. However, in a biased sampling scenario on $\boldsymbol{X}$, the probability of selecting an environmental combination is proportional to a predetermined weight function $w(\boldsymbol{x})$, implying that the pdf at the observation $\boldsymbol{x}$ is no longer $f(\boldsymbol{x}; \theta)$. Here, we propose a method of determining the weight function, $w(\boldsymbol{x})$, and the resulting pdf under biased sampling, which we then use to estimate a species' fundamental niche when $f(\boldsymbol{x}; \theta)$ is a multivariate normal density (i.e., a Mahalanobis model).

We illustrate the application of our proposed statistical model with two worked examples. In the first example, we estimate the fundamental niche of the Asian giant hornet *Vespa mandarinia*, an invasive species Europe that has recently been documented in the United States and Canada (Wilson et al., 2020). We first use presence records and an M hypothesis that only includes the native range of the hornet to fit a convex shape (an ellipsoid) and estimate the fundamental niche. We then evaluate the fitted model using the presence records from the invaded regions. In the second example, we again use ellipsoids, presence records, and species-specific M hypotheses to demonstrate how different M scenarios constrain a classic Mahalanobis model for different hummingbird species. We identify M scenarios under which we expect the Mahalanobis model to deviate from the fundamental niche and instead be closer to the realized niche of a species.

2   Materials and Methods

2.1   Modeling approach

Our aim was to account for the structure of the environmental space when attempting to estimate a fundamental niche. To address this problem, we follow Austin (2002) who suggested the inclusion of three major components in any modeling procedure in ecology: (1) an ecological model that describes the ecological assumptions to be incorporated into the analysis, and the ecological theory to be tested; (2) a statistical model that includes the statistical theory and methods used; and (3) a data model that accounts for how the data were collected or measured. The ecological model, described in the following section, includes a detailed definition of the fundamental niche as a function of both fitness and the relationship between fitness and the combinations of environmental conditions in which a species has been observed. The statistical

¹⁵³ model and the data model are addressed in subsequent sections.

The ecological model: relevant concepts in the study of the fundamental niche

¹⁵⁵ The fundamental niche of a species, N_F, is the set of all environmental conditions that permit

¹⁵⁶ the species to exist (Hutchinson, 1957; Peterson, Soberón, Pearson, et al., 2011). Let E $(\subseteq \mathbb{R}^d)$

¹⁵⁷ be a d-dimensional environmental space influencing fitness (measured, for example, as the finite

¹⁵⁸ rate of increase in a demographic response function). Furthermore, define a function, $\Lambda(\boldsymbol{x})$:

¹⁵⁹ $E \longrightarrow \mathbb{R}$, that relates each environmental combination, $\boldsymbol{x} \in E$, to fitness (Jiménez et al., 2019;

¹⁶⁰ Pulliam, 2000). If the fitness function has the right shape, there is a value of fitness, λ_{min},

¹⁶¹ that can be interpreted as minimum survivorship and defines the border of the fundamental

¹⁶² niche (Etherington & Omondiagbe, 2019; Jiménez et al., 2019). In other words, λ_{min} is the

¹⁶³ threshold above which the fitness is high enough to support a population; any combination of

¹⁶⁴ environmental values that results in a fitness lower than λ_{min} is outside the fundamental niche:

¹⁶⁵ $N_F = \{\boldsymbol{x} \in E | \Lambda(\boldsymbol{x}) \geq \lambda_{min}\}$. Notice that, using this notation, $\Lambda(\mu) = \lambda_{max}$.

¹⁶⁶ Assumptions about the shape of a species' response to an environmental variable (i.e.,

¹⁶⁷ the shape of the fitness function) are central to any predictive modeling effort (Austin, 2002;

¹⁶⁸ Soberón & Peterson, 2019). We assume that the contour surfaces of the fitness function, Λ,

¹⁶⁹ and, therefore, the boundary of the fundamental niche is convex (Drake, 2015). We specifically

¹⁷⁰ model these level surfaces using ellipsoids in multivariate space (Brown, 1984; Jiménez et al.,

¹⁷¹ 2019; Maguire, 1973) because these surfaces are simple and manageable convex sets defined by

¹⁷² (1) a vector μ that indicates the position of the optimal environmental conditions (the center of

¹⁷³ the ellipsoid) and (2) a covariance matrix Σ that defines the size and orientation of the ellipsoid

¹⁷⁴ in E. These two parameters can also be interpreted as the parameters of a multivariate normal

¹⁷⁵ density function, $f(\boldsymbol{x}; \mu, \Sigma)$, in which the corresponding random variable $\boldsymbol{X}$ represents a combi-

¹⁷⁶ nation environmental conditions in E where the species could be recorded as present (Jiménez

¹⁷⁷ et al., 2019). These ellipsoids are also known as Mahalanobis models (Farber & Kadmon, 2003)

¹⁷⁸ because the Mahalanobis distance defined by μ and Σ is equivalent the quadratic form that

¹⁷⁹ defines $f(\boldsymbol{x}; \mu, \Sigma)$. Ellipsoids representing the niche have been used to test the niche-center

¹⁸⁰ hypothesis (Osorio-Olvera et al., 2020), which predicts that the closer an environmental combi-

¹⁸¹ nation is to the vector μ, the higher the suitability value associated with that combination, and

¹⁸² therefore, the higher the abundance of the species there.

¹⁸³ We transform fitness values, $\Lambda(\boldsymbol{x})$, by calculating $f(\boldsymbol{x}; \mu, \Sigma)/f(\boldsymbol{x}; \hat{\mu}, \hat{\Sigma})$ (where $\hat{\mu}$ and $\hat{\Sigma}$ are

<div align="center">6</div>

estimates of μ and Σ, respectively) which produces a value between 0 and 1 that can be interpreted as a "suitability" index for the environmental combination $\boldsymbol{x}$. The central assumption here is that there is a monotonic transformation between $\Lambda(\boldsymbol{x})$ and $f(\boldsymbol{x}; \mu, \Sigma)$ in the sense that high values of fitness (near λ_{max}) correspond to high suitability values (near μ). We can therefore work with the normal model to delimit the environmental requirements of a species and test the niche-center hypothesis.

Other central concepts in the study of the fundamental niche are the existing niche (Jackson & Overpeck, 2000) and the realized niche (Hutchinson, 1957), which are each subsets of the fundamental niche. The existing niche represents the existing climatic conditions in the study region, and, the realized niche represents the subset of the fundamental niche that a species occupies as a result of biotic interactions with other species. We assume that the following relationships between three niche concepts – the fundamental niche N_F, the existing niche $N^*(t; M)$ and the realized niche N_R – are fulfilled (Peterson & Soberón, 2012; Peterson, Soberón, Pearson, et al., 2011):

$$N_F \supseteq N_F \cap E(t) = N^*(t; M) \supseteq N_R(t; M). \tag{1}$$

These relationships are illustrated in Figure 1 and constitute our ecological model. In theory, every environmental combination inside the fundamental niche ellipse is suitable for the species, and, if the species can reach a site with those environmental conditions, it could persist there indefinitely (in the absence of biotic factors such as predators). However, at a given time, t, only a discrete subset of the environmental combinations that can be mapped into E-space exist in geographic space. This subset constitutes the *existing niche* of the species. Note that this set of environmental combinations, which are contiguous points in E-space, are divided across different regions when represented in geographic space: some of the points are in North America (purple points) and others are in South America (green points). Species have dispersal limitations that may prevent them from colonizing all the environments in their existing niche. Thus, if a species is native to North America and is only able to reach the area shaded in orange (henceforth M), then (i) its realized niche will be a subset of purple points (because they are suitable environments that the species has access to) and (ii) the set of green points constitutes its potential niche (because they are suitable environments that are not accessible to the species). There is therefore only a discrete set of environmental combinations from the fundamental niche that is available to the species, and we expect to observe higher abundances of the species in the

environmental conditions that both exist and are close to the center of the fundamental niche ellipse.

Figure 1: Different subsets of environmental combinations that are of interest for niche estimation. Each grey point in geographic space corresponds to a grey point in E-space, and vice versa. The ellipse represents the border of a fundamental niche in which only the green and purple environmental combinations exist somewhere in geography (existing niche) and only the purple points are accessible to the species. The corresponding regions in geographic space are highlighted with the same colors.

The statistical and data models: two-stage sampling

Given a sample of environmental conditions where the species has been observed as present, $D = \{\boldsymbol{x}_1, ..., \boldsymbol{x}_n\}$, we propose a likelihood function for the parameters that describe the fundamental niche of the species ($\theta = (\mu, \Sigma)$). Suppose that the environmental space E is defined by d environmental variables (i.e., $D \subset E \subseteq \mathbb{R}^d$ and each point in E-space $\boldsymbol{x}_i$ has d coordinates). If E were a uniform grid of points embedded in $\mathbb{R}^d$, then the sample of conditions in which a species is present D could be considered to be a random sample of a multivariate normal variable with the density function

$$f(\boldsymbol{x}; \theta) = (2\pi)^{-d} |\Sigma|^{-1/2} \exp\left[-\frac{1}{2}(\boldsymbol{x} - \mu)^T \Sigma^{-1}(\boldsymbol{x} - \mu)\right], \tag{2}$$

This density function could theoretically be used to define a likelihood function and estimate μ and Σ.

This is unfortunately not the case, for two main reasons. First, the environmental combi-

nations that exist currently on the planet do not represent a uniform sample taken from the whole multivariate space E because Huthinson's duality does not preserve distances. If we take a uniform grid in geographic space and map it into environmental space, the resulting cloud of points will be concentrated in select regions, leaving other regions empty, as seen in Figure 1. Second, species occurrences can only come from E_M, or the set of environmental combinations associated with all the sites in the region that is accessible to the species M. The irregular shape of E_M induces a sampling bias such that the probability of recording the species as present in an environmental combination is no longer given directly by $f(\boldsymbol{x}; \mu, \Sigma)$, as these probabilities are affected by the availability of environmental conditions in E_M.

To account for the sampling bias induced by E_M, suppose that when the event $\{\boldsymbol{X} = \boldsymbol{x}\}$ occurs (i.e., when the species is recorded as present at a site with these environmental conditions), the probability of observing it changes depending on the observed $\boldsymbol{x}$. We represent this probability by $w(\boldsymbol{x})$. The set of observed environmental combinations, D, can therefore be considered as a random sample of the random variable $\boldsymbol{X}_w$, with probability density function

$$f_w(\boldsymbol{x}; \mu, \Sigma) = \frac{w(\boldsymbol{x}) f(\boldsymbol{x}; \mu, \Sigma)}{E[w(\boldsymbol{X})]}, \tag{3}$$

where

$$E[w(\boldsymbol{X})] = \int w(\boldsymbol{x}) f(\boldsymbol{x}; \mu, \Sigma) dx.$$

Note that $f_w(\cdot)$ is an example of a weighted density function in which $E[w(\boldsymbol{X})]$ is the normalizing factor, making the total probability function equal to unity (Lele & Keim, 2006; Patil & Ord, 1976; Patil & Rao, 1978). Patil & Rao (1978) call this normalizing factor the *visibility factor*, which captures the idea that samples from E are not uniform. The observed species presences can only come from E_M (the visible set of environmental combinations), which may include regions in $\mathbb{R}^d$ where the environmental combinations are abundant (associated to high values of $w(\boldsymbol{x})$). But, even though the points inside these abundant regions might not be close to μ (i.e., where $f(\boldsymbol{x}; \mu, \Sigma)$ is small), the probability of observing the species in these points could be higher than the probability associated to other points closer to μ (i.e., where $f(\boldsymbol{x}; \mu, \Sigma)$ is large) that do not exist in E_M (they are not visible), or whose weight $w(\boldsymbol{x})$ is too small.

On the other hand, equation 3 can also be described as the model resulting from a two-stage sampling design that accounts for the random process under which an environmental combination $\boldsymbol{x} \in D$ is observed (Patil & Rao, 1978). Suppose that nature produces a sample

of size N of environmental conditions inside the fundamental niche, with probabilities of being observed given by the density function $f(\boldsymbol{x}; \mu, \Sigma)$. This sample may contain any point in E. Because the species not only requires the right abiotic conditions to maintain a population, but also needs specific biotic conditions and can only disperse to a finite set of sites, the recorded sample will not include all the N observations. Instead, only a subsample of size $n < N$ is selected by drawing observations from the original sample (of size N) with a probabilities proportional to $w(\boldsymbol{x})$.

Let us illustrate this two-stage sampling design using the fundamental niche shown in Figure 1. Suppose that a species' fundamental niche is defined by a bivariate normal distribution with parameters:

$$\mu_1 = (-0.38, 1.85), \Sigma_1 = \begin{pmatrix} 0.0412 & 0.0366 \\ 0.0366 & 0.075 \end{pmatrix},$$

where the first dimension represents annual mean temperature (x-axis), and the second represents annual precipitation (y-axis). The level curves of the corresponding density function, $f(\boldsymbol{x}; \mu_1, \Sigma_1)$, are ellipses; they are plotted in the left panel of Figure 2, where the largest ellipse corresponds to the 99% confidence region. In the first stage of the sampling process, we generate a sample of size $N = 100$ from within the fundamental niche and plot this sample on top of the ellipses. As expected, most of the environmental combinations in this sample are close to the center of the ellipses (green points inside the ellipses plotted in the left panel of Figure 2). The middle panel of Figure 2 shows all the environmental combinations that exist inside the region that is accessible to the species, E_M, which are indicated by purple and orange points, in both Fig. 1 and Fig. 2). We identified the subset of points that both exist in E_M and are inside the 99% confidence region of $f(\boldsymbol{x}; \mu, \Sigma)$ and colored them purple. We then estimated the density function of the accessible environments, $\hat{h}(\cdot; E_M)$, using a kernel method. The resulting level curves of $\hat{h}(\cdot; E_M)$ correspond to the orange regions in the middle panel of Fig. 2. We used $\hat{h}(\cdot; E_M)$ to define the weights, $w(\boldsymbol{x}) = \hat{h}(\boldsymbol{x}; E_M)$, which were used in the second stage of the sampling process. Based on the assumption that not all the $N = 100$ points from the original sample exist in E_M, the second sampling stage involved selecting a subsample of size $n = 25$ from the first sample of $N = 100$ environments from the fundamental niche. The resulting sample, $D = \{\boldsymbol{x}_1, ..., \boldsymbol{x}_n\}$, is shown in the right panel of Figure 2 (purple triangles).

Note that if we use a simple likelihood approach based on the simulated sample to estimate the parameters μ_1 and Σ_1, which is equivalent to fitting a Mahalanobis model, we recover a 99%

confidence ellipse (violet ellipse in the right panel of Fig. 2) that is smaller than the theoretical fundamental niche of our species (largest, green ellipse in the right panel of Fig. 2). More importantly, the estimated center of this ellipse does not coincide with the optimal environmental conditions, μ_1. Therefore, if we estimate the parameters that describe the fundamental niche of the species without accounting for the distribution of available environments, we cannot claim that the model fully recovers the species' fundamental niche. Moreover, if we attempt to test the niche-center hypothesis using an ellipse recovered from a Mahalanobis model, we will expect high abundances around an environmental combination that is not the true optimum.

Figure 2: Left: A theoretical fundamental niche is represented as ellipsoids that correspond to the confidence regions of a normal distribution with parameters μ_1 and Σ_1. In the first stage of the sampling process, a sample of size $N = 100$ (green triangles) is simulated from this distribution. Middle: The environmental combinations accessible to the species, identified as E_M (orange and purple circles), and the contour levels of the kernel density function estimated using these points, $\hat{h}(\boldsymbol{x}; E_M)$ (orange regions). Right: In the second stage of the sampling process, a subsample of size $n = 25$ (purple triangles) is selected using the weights defined by $\hat{h}(\cdot; E_M)$, the points in E_M (orange circles), and the theoretical fundamental niche of the species (green ellipse).

With respect to the application of the two-stage sampling method, we assume that the sample of environmental combinations where the species of interest was observed as present, D, is a random sample of the random variable $\boldsymbol{X}_w$ with a probability density function given by Eq. 3. We can therefore define the likelihood function of the parameters of interest $\theta = (\mu, \Sigma)$ as follows:

$$\mathcal{L}(\theta|D) \propto \prod_{i=1}^{n} f_w(\boldsymbol{x}; \theta) = \prod_{i=1}^{n} \frac{w(\boldsymbol{x}) f(\boldsymbol{x}; \theta)}{E[w(\boldsymbol{X})]}, \tag{4}$$

where the function $w(\boldsymbol{x})$ is approximated using all the environmental combinations accessible to the species along with a kernel density procedure, and we obtain a Monte Carlo estimate for the expected value $E[w(\boldsymbol{X})]$ (which was defined as an integral; see Eq. 3). Note that the function $w(\boldsymbol{x})$ does not depend on the parameters of interest and can be ignored when maximizing the log-

likelihood function. Furthermore, because the analytical form of $w(\boldsymbol{x})$ is unknown, the analytical form of $E[w(\boldsymbol{X})]$ is also unknown. However, we can sample environmental combinations from the availability distribution by randomly choosing sites inside M and extracting the values of the environmental variables at those sampled sites. Using this sample, we can obtain Monte-Carlo estimates of $E[w(\boldsymbol{X})]$ for any fixed value of θ. This method has been used before in the statistical modeling literature and is called the method of simulated maximum likelihood (Lele & Keim, 2006; Robert & Casella, 1999). We can use this method to obtain a Monte Carlo estimate of the log-likelihood function as follows:

$$\ell(\theta|D) = \sum_{i=1}^{n} \log(f_w(\boldsymbol{x}_i; \theta)) = \sum_{i=1}^{n} \left\{ \log(f_w(\boldsymbol{x}_i; \theta)) - \log\left[\frac{1}{K} \sum_{j=1}^{K} f_w(\boldsymbol{x}_j^*; \theta)\right] \right\}, \qquad (5)$$

where $\boldsymbol{x}_j^*, j = 1, 2, ..., K$ is a random sample that was sampled with replacement from the distribution $w(\boldsymbol{x})$. The size of this sample, K, must be large enough to ignore Monte Carlo error. Once this sample is generated, we can apply standard optimization techniques to minimize the negative log-likelihood function in Eq. 5 and obtain the maximum likelihood estimators $\hat{\mu}$ and $\hat{\Sigma}$.

In summary, our proposed modeling procedure uses a sample of occurrences for the species of interest together with polygons that represent the geographic areas accessible to the species. Inside these polygons, we extract environmental values from sites chosen at random. This is done with two different purposes: (1) to estimate a kernel density used to define the weights in the likelihood function, and (2) to obtain a Monte Carlo estimate of the log-likelihood function. Once we have all the elements of the likelihood function, we calculate the maximum likelihood estimates of the parameters $\hat{\mu}$ and $\hat{\Sigma}$. These estimated parameters will allow us to plot ellipses in E-space that represent the border of the estimated fundamental niche. In all the examples we show, we plot the ellipses that correspond to the 99% confidence regions of the fitted multivariate normal distribution. We compare these ellipses to the ones that correspond to the 99% confidence region of a standard Mahalanobis model (Farber & Kadmon, 2003) with estimated parameters $\hat{\mu}_0 = \frac{1}{n} \sum_{i=1}^{n} \boldsymbol{x}_i$ and $\hat{\Sigma}_0 = \frac{1}{n} \sum_{i=1}^{n} (\boldsymbol{x}_i - \hat{\mu}_0)(\boldsymbol{x}_i - \hat{\mu}_0)^T$ (i.e., the maximum likelihood estimates under a multivariate normal model described in Eq. 2). We hypothesized that the two ellipses will be similar in cases where E_M covers most of the fundamental niche of the species (i.e., the overlap between these two sets is high) and the distribution of points in E_M is approximately uniform.

Finally, we project the resulting models back to geographic space. Once we have the maximum likelihood estimates of the parameters of interest, $\hat{\theta}$, we use the multivariate normal density function given in Eq. 2 to calculate a suitability index, that can be plotted in G-space as either a continuous value or a binary region defined by a threshold. For interpretation purposes, it is convenient to standardize this suitability index to the interval $(0, 1)$, which is easily done by dividing $f(\boldsymbol{x}; \theta)$ by its maximum value, $f(\boldsymbol{x}; \hat{\theta})$.

2.2 Data

Using our method to approximate fundamental niches requires three types of data: (*i*) species occurrence data, which should be thoroughly cleaned before being used; (*ii*) an M hypothesis or geographic polygon that encloses all the sites accessible to the species, taking into consideration the dispersal ability of the species and natural geographic barriers; and (*iii*) environmental layers, cropped to the study area, from which we can extract environmental data for the occurrences and for the sites inside M. In the following sections, we describe the datasets that we used to create our worked examples.

Occurrence data

We selected seven species to illustrate the use of the statistical model that we present here. The first species is the Asian giant hornet *Vespa mandarinia* (Matsuura & Sakagami, 1973; Matsuura, 1988). Because *V. mandarinia* is an invasive species, using it in our model can provide valuable insight into whether the niche estimated by our model is a good approximation of a true fundamental niche. In other words, we test whether the estimated fundamental niche contains the locations that this species has been able to invade. We also selected six species of hummingbirds for a separate analysis: *Amazilia chionogaster*, *Threnetes ruckeri*, *Sephanoides sephanoides*, *Basilinna leucotis*, *Colibri thalassinus*, and *Calypte costae*. These species were used because detailed M hypotheses are available for them.

The occurrence data for *V. mandarinia* was obtained from the Global Biodiversity Information Facility (GBIF) database (`https://www.gbif.org/`). We downloaded 1944 occurrence records for (GBIF, 2020), which, after undergoing a standard cleaning procedure (Cobos et al., 2018), were reduced to 170 presence records in the species' native range and one presence record in Europe. To avoid introducing an extra source of bias (Anderson, 2012), the cleaned dataset was spatially thinned by geographic distance (at least 50 km away), resulting in a final sample

of 46 presence records that were used to fit the niche models (see red points in Fig. 3).

For the hummingbird species, we used the cleaned occurrence records that were used by Cooper & Soberón (2018), which are available at `https://github.com/jacobccooper/trochilidae` and whose original source is also GBIF. Cooper & Soberón thoroughly revised and cleaned these presence records, and eliminated misidentified individuals and synonyms. They additionally defined the regions that were accessible to most of the extant hummingbird species and used those regions to fit species distribution models. Given the success of Cooper & Soberón at using these M hypothesis, we believed their data provided a great opportunity to test our proposed model.

Figure 4 shows the presence data of each hummingbird species. The sample sizes range from 148 observations for *A. chionogaster*, to 926 presences for *C. costae*. We selected these six species because they occupy different regions of the Americas, from the southern United States to the Patagonia. Given this wide geographic range, we believed it would be interesting to compare the estimated niches of all the species in E-space and evaluate how similar or different the optimal environmental conditions are among the different species. Although the different species are likely to occupy different regions of E-space, we suspected that their fundamental niches might share some environmental combinations (niche overlap).

M hypothesis

In the case of *V. mandarinia*, we defined the area accessible to the species as a combination of buffers that represent the species' dispersal ability and the elevation range where the species is known to occur (850 - 1900 m). First, we identified the region delineated by a buffer of 500 km around all occurrence records in the sample, which accounts for the dispersal ability of the hornets (Matsuura & Sakagami, 1973). Second, we clipped this region with an elevation layer to remove regions at elevations higher than 1900 m. The resulting polygon is outlined in blue in Figure 3. Note that this polygon includes some sites not on the continent; however, we only extracted the environmental values for inland sites within this blue M, and did not considered oceanic regions.

For the hummingbird species, we used the polygons generated by Cooper & Soberón (2018). Polygons were hypothesized based on the known species occurrences, topography, ecoregions, and where bounded by significant geographical barriers such as large rivers and mountains. By doing so, they took into account all the criteria that are know to yield more accurate models (N. Barve et al., 2011; Owens, Campbell, et al., 2013;

Owens, Ribeiro, et al., 2020; Saupe, V. Barve, et al., 2012). Figure 4 shows the six polygons for the different species. Note that some species, like *T. ruckeri*, occupy most of their accessible area (Fig. 4b), whereas others, like *B. leucotis* and *C. costae*, occupy only a fraction of it. In addition, there is wide diversity in range sizes, with *S. sephanoides* having a more restricted range compared to *C. thalassinus*. Given this geographic variation among species, we were interested in making similar comparisons in E-space.

Environmental variables

The climatic layers used to create the models for each species came from the WorldClim database (Hijmans et al., 2005). We used only two of the 19 variables available in this database: annual mean temperature (Bio1), and annual precipitation (Bio12), The two climatic variables variables that we selected are biologically meaningful for both the Asian hornet (Matsuura & Sakagami, 1973), and (Root, 1988) and the Hummingbirds. Furthermore, in the case of the hummingbirds, we wanted to compare their estimated niches. Because comparisons of estimated niches in E-spaces with different axes and/or dimensions are not possible, we only looked at these two dimensions of their niches. Both variables were recorded at 10 arcmin resolution. We clipped each climatic layer using the polygons that correspond to the M hypotheses for each species.

Figure 3: Occurrence records (red squares) of *V. mandarinia* in its native range, and the M hypothesis (regions delineated with blue lines) created from a combination of buffers and the known elevation range of the species. There were a total of 46 occurrence records after the data cleaning and thinning process.

All analyses were performed in R version 3.6.3 (R Core Team, 2020). We used several existing packages for data preparation, analysis, and visualization: ggplot2 (Wickham, 2016), ks (Duong,

2020), raster (Hijmans, 2020), rgdal (Bivand, Keitt, et al., 2020),rgeos (Bivand & Rundel, 2019),

scales (Wickham & Seidel, 2020), sf (Pebesma, 2018). We additionally created new functions

that can be used to reproduce our examples, and apply our methodology to other species. These

functions can be found at `https://github.com/LauraJim`.

Figure 4: Occurrence samples and M polygons for the six species of hummingbirds selected for this study. (a) *A. chionogaster* is represented in brown, (b) *T. ruckeri* in red, (c) *S. sephanoides* in pink, (d) *B. leucotis* in green, (e) *C. thalassinus* in pale blue, and (g) *C. costae* in yellow. The sample sizes (n) are given in each panel.

3 Results

3.1 Estimated fundamental niche of *Vespa mandarinia*

We randomly sampled 10,000 sites inside the accessible (native) region of the Asian giant hornet (shown in Fig. 3) and extracted their environmental values (E_M) so we could plot them in E-space. The resulting set E_M is shown in Figure 5 as a cloud of points in the background (grey open circles). We used this E_M and the 46 presence records of *V. mandarinia* (red points in Fig. 5) to determine the log-likelihood function (Eq. 5) of the parameters that describes its fundamental niche under the weighted-normal model, $\theta = (\mu, \Sigma)$. We maximized this log-likelihood function to obtain the maximum likelihood estimates (MLEs) $\hat{\mu}$ and $\hat{\Sigma}$. For comparison, we also used the 46 presence records only to obtain MLEs using the Mahalanobis model, $\hat{\mu}_0$ and $\hat{\Sigma}_0$. The resulting MLEs for both models are given in Table 1 and the corresponding 99% confidence regions are plotted in Figure 5.

Figure 5: Estimated ellipses (99% confidence regions) from the weighted-normal model(red) and the Mahalanobis model (purple), which are defined by $(\hat{\mu}, \hat{\Sigma})$ and $(\hat{\mu}_0, \hat{\Sigma}_0)$, respectively, as given in Table 1. Red points are the presences from the native range of *V. mandarinia* that were used to fit the models, and blue points are presences recorded outside the native range. The centers of the ellipses are indicated with a square of the same color as the corresponding model. The grey points represent E_M.

The MLEs of μ (the optimal environmental conditions) estimated by both the weighted-normal model and the Mahalanobis model, are highly similar (purple and red squares in Figure 5). However, the models predicted different limits for the fundamental niche of *V. mandarinia*. Specifically, the estimated N_F is larger under the Mahalanobis model (purple ellipse in Fig. 5), whereas the variance predicted by the weighted-normal model for annual precipitation is smaller, as shown in Table 1. Nevertheless, both estimated ellipses contain 45 out of the 46 initial presence points and they contain all the presences from the locations that *V. mandarinia* has invaded (blue points in Fig. 5). The presences from the invaded regions are not close to the center of the ellipses, but they are placed in a region of E_M where the environmental combinations are well represented in G-space.

We calculated a standardized suitability index for the Asian giant hornet using the MLEs from the weighted-normal model and Eq. 2. Using this index, we created a worldwide suitability map (Fig. 6) to visually assess if there are regions where *V. mandarinia* could theoretically establish based solely on the species' environmental requirements. As shown in Figure 6, the Pacific coast of the northern United States and southern Canada, where the species has already been confirmed, has a low to moderate suitability index. Similarly, most of Europe has a low to moderate suitability, particularly around the one site where the species was already recorded. Additional maps that are focused more closely on the West Coast of North America, Europe, and the native range of *V. mandarinia* are provided in the Supplementary Material. We also note that there are other regions that appear highly suitable for *V. mandarinia*, such as the eastern United States, the mountain ranges of Mexico and the Andes, and the Brazilian and Ethiopian highlands.

Table 1: MLEs of the weighted-normal model ($\hat{\mu}$ and $\hat{\Sigma}$) and the Mahalanobis model ($\hat{\mu}_0$ and $\hat{\Sigma}_0$) obtained from 46 presences of *V. mandarinia* inside its native range.

Model	Weighted-Normal		Mahalanobis	
Parameters	$\hat{\mu}$	$\hat{\Sigma}$	$\hat{\mu}_0$	$\hat{\Sigma}_0$
V. mandarinia	(170.21, 1012.19)	$\begin{pmatrix} 2223.45 & 8031.54 \\ 8031.54 & 39554.33 \end{pmatrix}$	(167.52, 992.97)	$\begin{pmatrix} 2348.47 & 9813.12 \\ 9813.12 & 52770.37 \end{pmatrix}$

Figure 6: Worldwide suitability map of *V. mandarinia.* Dark purple regions correspond to highly suitable sites where the environmental combinations in E-space are close to the center of the estimated fundamental niche, and light purple regions correspond to sites with low suitability whose environmental combinations in E-space are near the border of the fundamental niche.

3.2 Estimated fundamental niches of hummingbird species

In the case of the six hummingbird species that we chose for our analysis, the accessible areas (Ms) and the presence records were in different regions of G-space but also, of E-space. The corresponding E_M and presence points therefore occupied different regions (see grey clouds of points in Fig. 7). Using the M hypothesis and the presences, we estimated the parameters μ and Σ, which describe the fundamental niche of each species, using both the Mahalanobis model and the weighted-normal model. The resulting MLEs for each species are given in Table 2.

Unlike the *V. mandarinia* example, the ellipses estimated with the Mahalanobis model were smaller than the ellipses estimated by the weighted-normal model for all species except *C. thalassinus*. In other words, the weighted-normal model predicted broader fundamental niches for most of the studied hummingbirds. The estimated centers of the ellipses ($\hat{\mu}_0$ and $\hat{\mu}$) and their orientation with respect to the axes were similar for *A. chionogaster* and *B. leucotis*, however, there was a clear difference between the centers of the estimated ellipses for the *T. ruckeri, S. sephanoides, C. thalassinus*, and *C. costae*.

For all the hummingbird species, both the Mahalanobis and the weighed-normal models agreed on the sign of the covariance between the two environmental variables used to describe and compare the fundamental niches. The estimated ellipses for *S. sephanoides* were the most

different with respect to the magnitude of the estimated covariance between the two environ-

mental variables. We additionally noted that, with the exception of *C. thalassinus*, the ellipses

obtained with the weighted-normal model contained more presence points than the ellipses ob-

tained from a simple Mahalanobis model.

It is also worth noting that, under the Mahalanobis model, the estimated optimum tempera-

ture for four of the six hummingbird species was between 16 and 18 degrees Celsius (see Table 2).

The Mahalanobis model thus predicts that the species temperature optima are not substantially

different among these species, even though they differ in their precipitation optima. However,

under the weighted-normal model, the estimated optimum temperatures were clearly different

among all the species (see Figure 8).

Table 2: Maximum likelihood estimates of the parameters that determine the fundamental niche of the six hummingbird species. Parameters were obtained with the weighted-normal model (second column) and the Mahalanobis model (third column).

Model	Weighted-Normal		Mahalanobis	
Species	$\hat{\mu}$	$\hat{\Sigma}$	$\hat{\mu}_0$	$\hat{\Sigma}_0$
A. chionogaster	(155.20, 939.80)	$\begin{pmatrix} 1975.96 & 6702.73 \\ 6702.73 & 191251.29 \end{pmatrix}$	(161.72, 854.68)	$\begin{pmatrix} 1405.95 & 3874.59 \\ 3874.59 & 123768.21 \end{pmatrix}$
T. ruckeri	(232.92, 3187.12)	$\begin{pmatrix} 710.18 & -4647.29 \\ -4647.29 & 897405.47 \end{pmatrix}$	(244.71, 2894.13)	$\begin{pmatrix} 420.79 & -1367.22 \\ -1367.22 & 816368.46 \end{pmatrix}$
S. sephanoides	(125.88, 1341.95)	$\begin{pmatrix} 1877.22 & -11075.44 \\ -11075.44 & 296005.42 \end{pmatrix}$	(108.98, 1208.83)	$\begin{pmatrix} 1024.16 & -10577.69 \\ -10577.69 & 542467.87 \end{pmatrix}$
B. leucotis	(180.99, 1310.40)	$\begin{pmatrix} 1556.64 & 7922.37 \\ 7922.37 & 565818.85 \end{pmatrix}$	(177.08, 1199.03)	$\begin{pmatrix} 1163.15 & 6300.40 \\ 6300.40 & 281258.818 \end{pmatrix}$
C. thalassinus	(146.58, 1826.31)	$\begin{pmatrix} 2103.17 & 11601.23 \\ 11601.23 & 466300.43 \end{pmatrix}$	(177.43, 1640.71)	$\begin{pmatrix} 1821.22 & 13570.96 \\ 13570.96 & 646108.77 \end{pmatrix}$
C. costae	(197.17, 266.32)	$\begin{pmatrix} 2287.66 & -5866.60 \\ -5866.60 & 37195.04 \end{pmatrix}$	(177.87, 277.69)	$\begin{pmatrix} 1563.16 & -3495.43 \\ -3495.43 & 23727.06 \end{pmatrix}$

Figure 7: Estimated fundamental niches for the six species of hummingbirds. In all panels, the purple ellipse represents the estimated niche from a Mahalanobis model, and the other ellipse represent the estimated niche from our proposed weighted-normal model. The centers of both ellipses are marked with a purple square and a black square, respectively.

Figure 8: Comparison of the estimated fundamental niches (ellipses) for the six species of hummingbirds and the presence data used to fit the models. The centers of the ellipses are marked with a black circle

4 Discussion

In recent decades, we have seen a significant increase in the availability of species presence data and software for ecological niche modeling and species distribution modeling. At the same time, however, we have seen a concomitant increase in the number of studies that question important conceptual and methodological aspects of ENMs (Austin, 2002; Godsoe, 2010; Jiménez-valverde et al., 2008; Lobo, 2008). For that reason, we agree with Jiménez-valverde et al. (2008), who concluded that the lack of a solid conceptual background endangers the advancement of the field. The development and application of ENM/SDM should be rooted in a firm understanding of the technique's conceptual background. We intended to lead by example in this work by explicitly relating the ecological theory to the statistical method that we developed to estimate the fundamental niche of a species.

It is essential to acknowledge that the presence points of a species come from the realized niche. If our objective is to use presence data to approximate the fundamental niche, then we must account for the bias induced in the sample by the available E-space in the region of interest. In this study, we have therefore proposed defining the distribution of environments that are accessible to the species based on the dispersal ability of the species and geographic barriers to dispersal. This is a definition of M, the geographic area accessible to the species via dispersal (*sensu* Peterson, Soberón, Pearson, et al., 2011), but here we highlight the implications of sampling M in E-space, which are seldom discussed. By projecting M into E-space, we determined its empirical probability density function and used it as a weight function in the multivariate normal distribution, which we then used to estimate the fundamental niche of a species of interest. Our main result is that accounting for the shape of the E-space defined by an M indeed affects the estimation of the ellipsoids we use to model fundamental niches.

We illustrated the application of our proposed method with two examples. In the first example, we showed how presence data for invasive species, the Asian giant hornet, can be used to evaluate the estimated shape of the fundamental niche. We also showed how the fitted model can be projected back into G-space to get a suitability map beyond the region M. The estimated fundamental ellipse of *V. mandarinia* estimated using our model contained the sited the species is known to have invaded and the suitability map confirms that these known invaded regions are indeed suitable for the species.

In the second example, we showed how to use existing M hypotheses and presence data to approximate the fundamental niches of closely related species. We specifically examined six hummingbird species that according to phylogenetic taxonomies belong to different major clades: *T. ruckeri* in the Hermits (Phaethornithinae), *C. thalassinus* in the Mangoes (Polytmini), *S. sephanoides* in the Coquettes (Lophornithini), *C. costae* in the Bees (Mellisugini), *A. chianogaster* and *H. leucotis* in the Emeralds (Trochilini). The clades here were listed starting with the one having the oldest split and following the order in which the splitting continued (Hernández-Baños et al., 2014; McGuire et al., 2009). We compared the estimated fundamental niches (ellipses) of these six species with respect to the phylogenetic relationships among them and noted that *T. ruckeri*, which has the oldest split in the phylogeny, had the most different estimated fundamental niche (red ellipse in Fig. 8) among the six species. On the other hand, the other five species shared some regions in environmental space that belong to their fundamental niches. The same patterns were observed when comparing the centers of the

fundamental niche ellipses. These observations support the theory of conservatism of ecological niches, which predicts low niche differentiation between species over evolutionary time scales (Peterson, Soberón & Sánchez-Cordero, 1999).

The central idea in our modeling approach is that the set of accessible sites, represented in environmental space (E_M), restricts the sample space from which we can obtain occurrence records for a species. However, it is universal practice in ENM and SDM to use the environments in the occurrence data to fit niche models. Given that we use the distribution of E_M to inform the model about the uneven availability of environmental combinations, and, therefore, about the bias of the sample of presence points induced by M, the delimitation of M is crucial. However, there is no unique way to delimit the accessible area of a species (N. Barve et al., 2011). When outlining M, we need to simultaneously consider several factors: the natural history of the species, its dispersal characteristics, the geography of the landscape, the time span relevant to the species' presence, and any environmental changes that occurred during that time span, among other things. Although we do not favor any particular approach as the best method for estimating M, we acknowledge that all those factors have an important role in determining its shape. As a future work of this research project, a thorough analysis comparing different approaches to outlining M and testing their effects in recovering the fundamental niche of a species under the proposed model is needed. This could be done using virtual species, where we know the true parameters that shape the fundamental niche, or using invasive species, where the invaded locations are used as evaluation points to test the fitted model for the fundamental niche, as we did here in the example of *V. mandarinia*).

The presence of geographic sampling biases in primary biodiversity data and the implications of these biases (e.g., decreasing model performance) are broadly recognized (Kadmon et al., 2004; Meyer et al., 2016). Here, we focused on one specific type of bias that affects the estimation of a fundamental niche. However, other types of sampling biases, such as the accessibility of a site to observers, may also affect the estimation of the fundamental niche. In such cases, it would be relatively easy to incorporate the effect of other inherent sampling biases into our proposed model. We would still use a multivariate normal distribution to represent the hypothesis that the fundamental niche has a simple, convex shape, but we would modify the way we define the weights in the likelihood function (Eq. 4). For example, suppose we want to include the effects of two sources of bias that affect the presence of a species: the bias induced by M and the differences in sampling intensity across the landscape due to differences in

human accessibility (i.e., accessibility bias). Each source of bias will define a different sampling probability distribution across the environmental combinations from the region of interest. If the biases are independent, we can calculate a single, general sampling probability distribution as the product of the individual sampling probability distributions and substitute it in Eq. 4. This is, the function $w(\boldsymbol{x})$ can be defined as the product of two probabilities, the probability of observing $\boldsymbol{x}$ considering the distribution of points in E_M and the probability of observing $\boldsymbol{x}$ considering the accessibility bias. We proposed using a kernel to estimate the former, and Zizka et al. (2020) recently developed a method to quantify accessibility biases that could be used to determine the later.

Our proposed model to estimate fundamental niches can also be generalized in othert ways to represent more realistic scenarios. In the previous paragraph, we described an approach to include more than one type of bias but several other modifications are possible. For example, information from physiological experiments could be included if we consider the Bayesian approach proposed by Jiménez et al. (2019). To do so, we need to transform the likelihood function (Eq. 4) into a posterior distribution that uses the tolerance ranges obtained from physiological experiments to define the *a priori* distributions of the parameters μ and Σ ($g_1(\mu)$ and $g_2(\Sigma)$, respectively), as follows:

$$f(\mu, \Sigma; D) = \mathcal{L}(\mu, \Sigma | D) g_1(\mu) g_2(\Sigma) \tag{6}$$

where

$$\mathcal{L}(\mu, \Sigma | D) \propto \prod_{i=1}^{n} \frac{w(\boldsymbol{x}_i) f(\boldsymbol{x}_i; \mu, \Sigma)}{\sum_{y \in E} w(y) f(y; \mu, \Sigma)}. \tag{7}$$

Note that the function that determines the shape and size of the fundamental niche, $f(\cdot; \mu, \Sigma)$, can be modified to have asymmetrical confidence regions for species that are known to have asymmetrical response curves for the environmental variables under consideration (Jiménez et al., 2019). Furthermore, the likelihood and the Bayesian approaches can be applied in cases where it makes sense to include more than two environmental variables in the delimitation of a species' fundamental niche. The disadvantage of making these modifications to our proposed model is that they increase the algebraic complexity of the model and the number of parameters to be estimated, and more computational power is therefore needed to either maximize the likelihood function (if we do not include physiological tolerances and use the likelihood approach) or simulate from the posterior distribution (when considering physiological tolerances, using the Bayesian approach).

Acknowledgements

We thank the KU-ENM group at the Biodiversity Institute of the University of Kansas for providing valuable comments. We would also want to thank Maria E. Orive at the University of Kansas for her valuable comments. L.J. was supported by the Mexican Council of Science and Technology (CONACyT) grant 409052.

References

Anderson, R. P. (2012). Harnessing the world's biodiversity data: promise and peril in ecological niche modeling of species distributions. *Annals of the New York Academy of Sciences* 1260.1, pp. 66–80. DOI: `10.1111/j.1749-6632.2011.06440.x`.

Aspinall, R. & B. Lees (1994). Sampling and analysis of spatial environmental data. *Advances in GIS Research. Taylor and Francis, Southampton*, pp. 1086–1098.

Austin, M. (2002). Spatial prediction of species distribution: an interface between ecological theory and statistical modelling. *Ecological Modelling* 157.2-3, pp. 101–118. DOI: `10.1016/S0304-3800(02)00205-3`.

Barve, N., V. Barve, A. Jiménez-Valverde, A. Lira-Noriega, S. P. Maher, A. T. Peterson, J. Soberón & F. Villalobos (2011). The crucial role of the accessible area in ecological niche modeling and species distribution modeling. *Ecological Modelling* 222.11, pp. 1810–1819. DOI: `10.1016/j.ecolmodel.2011.02.011`.

Bivand, R., T. Keitt & B. Rowlingson (2020). *rgdal: Bindings for the 'Geospatial' Data Abstraction Library*. R package version 1.5-10. URL: `https://CRAN.R-project.org/package=rgdal`.

Bivand, R. & C. Rundel (2019). *rgeos: Interface to Geometry Engine - Open Source ('GEOS')*. R package version 0.5-2. URL: `https://CRAN.R-project.org/package=rgeos`.

Brown, J. H. (1984). On the Relationship between Abundance and Distribution of Species. *The American Naturalist* 124.2, pp. 255–279. DOI: `10.1093/ehr/cepl85`.

Chapman, A. D. (2005). Principles and methods of data cleaning–primary species and species-occurrence data, version 1.0. *Global Biodiversity Information Facility, Copenhagen* 75.

Cobos, M. E., L. Jiménez, C. Nuñez-Penichet, D. Romero-Alvarez & M. Simoes (2018). Sample data and training modules for cleaning biodiversity information. *Biodiversity Informatics* 13, pp. 49–50. DOI: `10.17161/bi.v13i0.7600`.

Colwell, R. K. & T. F. Rangel (2009). Hutchinson's duality: The once and future niche. *Proceedings of the National Academy of Sciences of the United States of America* 106.SUPPL. 2, pp. 19651–19658. DOI: 10.1073/pnas.0901650106.

Cooper, J. C. & J. Soberón (2018). Creating individual accessible area hypotheses improves stacked species distribution model performance. *Global Ecology and Biogeography* 27.1, pp. 156–165. DOI: 10.1111/geb.12678.

Drake, J. M. (2015). Range bagging: A new method for ecological niche modelling from presence-only data. *Journal of the Royal Society Interface* 12.107. DOI: 10.1098/rsif.2015.0086.

Duong, T. (2020). *ks: Kernel Smoothing.* R package version 1.11.7. URL: https://CRAN.R-project.org/package=ks.

Etherington, T. & O. Omondiagbe (2019). virtualNicheR: generating virtual fundamental and realised niches for use in virtual ecology experiments. *Journal of Open Source Software* 4.41, p. 1661. DOI: 10.21105/joss.01661.

Farber, O. & R. Kadmon (2003). Assessment of alternative approaches for bioclimatic modeling with special emphasis on the Mahalanobis distance. *Ecological Modelling* 160.1-2, pp. 115–130. DOI: 10.1016/S0304-3800(02)00327-7.

GBIF (2020). *Occurrence download.* Accessed: 07 May 2020. DOI: 10.15468/dl.kzcgc2. URL: https://www.gbif.org/.

Godsoe, W. (2010). I can't define the niche but i know it when i see it: A formal link between statistical theory and the ecological niche. *Oikos* 119.1, pp. 53–60. DOI: 10.1111/j.1600-0706.2009.17630.x.

Guisan, A. et al. (2013). Predicting species distributions for conservation decisions. *Ecology Letters* 16.12, pp. 1424–1435. DOI: 10.1111/ele.12189.

Hernández-Baños, B. E., L. E. Zamudio-Beltrán, L. E. Eguiarte-Fruns, J. Klicka & J. García-Moreno (2014). The Basilinna genus (Aves: Trochilidae): An evaluation based on molecular evidence and implications for the genus Hylocharis. *Revista Mexicana de Biodiversidad* 85.3, pp. 797–807. DOI: 10.7550/rmb.35769.

Hijmans, R. J. (2020). *raster: Geographic Data Analysis and Modeling.* R package version 3.1-5. URL: https://CRAN.R-project.org/package=raster.

Hijmans, R. J., S. E. Cameron, J. L. Parra, P. G. Jones & A. Jarvis (2005). Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology* 25.15, pp. 1965–1978. DOI: 10.1002/joc.1276.

Hoogenboom, M. O. & S. R. Connolly (2009). Defining fundamental niche dimensions of corals: synergistic effects of colony size, light, and flow. *Ecology* 90.3, pp. 767–780. DOI: 10.1890/07-2010.1.

Hutchinson, G. E. (1957). Concluding remarks. *Cold Sprig Harbor Symposia on Quantitative Biology.* Chap. 22, pp. 415–427.

Jackson, S. T. & J. T. Overpeck (2000). Responses of plant populations and communities to environmental changes of the late Quaternary. *Paleobiology* 26.4 SUPPL. Pp. 194–220. DOI: 10.1017/s0094837300026932.

Jiménez, L., J. Soberón, J. A. Christen & D. Soto (2019). On the problem of modeling a fundamental niche from occurrence data. *Ecological Modelling* 397.February, pp. 74–83. DOI: 10.1016/j.ecolmodel.2019.01.020.

Jiménez-valverde, A., J. M. Lobo & J. Hortal (2008). Not as good as they seem : the importance of concepts in species distribution modelling, pp. 885–890. DOI: 10.1111/j.1472-4642.2008.00496.x.

Kadmon, R., O. Farber & A. Danin (2004). Effect of roadside bias on the accuracy of predictive maps produced by bioclimatic models. *Ecological Applications* 14.2, pp. 401–413. DOI: 10.1890/02-5364.

Lele, S. R. & J. L. Keim (2006). Weighted distributions and estimation of resource selection probability functions. *Ecology* 87.12, pp. 3021–3028. DOI: 10.1890/0012-9658(2006)87[3021:WDAEOR]2.0.CO;2.

Lobo, J. M. (2008). More complex distribution models or more representative data. 82, pp. 14–19.

Lobo, J. M. (2016). The use of occurrence data to predict the effects of climate change on insects. *Current Opinion in Insect Science* 17, pp. 62–68. DOI: 10.1016/j.cois.2016.07.003.

Maguire, B. (1973). Niche Response Structure and the Analytical Potentials of Its Relationship to the Habitat Author (s): Bassett Maguire , Jr . Source : The American Naturalist , Vol . 107 , No . 954 (Mar . - Apr ., 1973), pp . 213-246 Published by : The University of C. *The American naturalist* 107.954, pp. 213–246.

Matsuura, M. & S. Sakagami (1973). A bionomic sketch of the giant hornet. *Vespa mandarinia.*

Matsuura, M. (1988). Ecological study on vespine wasps (Hymenoptera: Vespidae) attacking honeybee colonies: I. seasonal changes in the frequency of visits to apiaries by vespine wasps

and damage inflicted, especially in the absence of artificial protection. *Applied Entomology and Zoology* 23.4, pp. 428–440.

McGuire, J. A., C. C. Witt, J. V. Remsen, R. Dudley & D. L. Altshuler (2009). A higher-level taxonomy for hummingbirds. *Journal of Ornithology* 150.1, pp. 155–165. DOI: `10.1007/s10336-008-0330-x`.

Meyer, C., P. Weigelt & H. Kreft (2016). Multidimensional biases, gaps and uncertainties in global plant occurrence information. *Ecology letters* 19.8, pp. 992–1006. DOI: `10.1111/ele.12624`.

Osorio-Olvera, L., C. Yañez-Arenas, E. Martínez-Meyer & A. T. Peterson (2020). *Relationships between population densities and niche-centroid distances in North American birds*. DOI: `10.1111/ele.13453`.

Owens, H. L., L. P. Campbell, L. L. Dornak, E. E. Saupe, N. Barve, J. Soberón, K. Ingenloff, A. Lira-Noriega, C. M. Hensz, C. E. Myers & A. T. Peterson (2013). Constraints on interpretation of ecological niche models by limited environmental ranges on calibration areas. *Ecological Modelling* 263, pp. 10–18. DOI: `10.1016/j.ecolmodel.2013.04.011`.

Owens, H. L., V. Ribeiro, E. E. Saupe, M. E. Cobos, P. A. Hosner, J. C. Cooper, A. M. Samy, V. Barve, N. Barve, C. J. Muñoz-R. & A. T. Peterson (2020). Acknowledging uncertainty in evolutionary reconstructions of ecological niches. *Ecology and Evolution* 10.14, pp. 6967–6977. DOI: `10.1002/ece3.6359`.

Patil, G. & J. Ord (1976). On Size-Biased Sampling and Related Form-Invariant Weighted Distributions. *Sankhyā: The Indian Journal of StatisticsSeries B* 38.1, pp. 48–61.

Patil, G. & C. Rao (1978). Weighted Distributions and Size-Biased Sampling with Applications to Wildlife Populations and Human Families. *Biometrics* 34.2, pp. 179–189.

Pebesma, E. (2018). Simple Features for R: Standardized Support for Spatial Vector Data. *The R Journal* 10.1, pp. 439–446. DOI: `10.32614/RJ-2018-009`.

Peterson, A. T. & J. Soberón (2012). Integrating fundamental concepts of ecology, biogeography, and sampling into effective ecological niche modeling and species distribution modeling. *Plant Biosystems* 146.4, pp. 789–796. DOI: `10.1080/11263504.2012.740083`.

Peterson, A. T., J. Soberón, R. G. Pearson, R. P. Anderson, E. Martínez-Meyer, M. Nakamura & M. B. Araújo (2011). Ecological niches and geographic distributions. *Princeton University Press* 49.11, pp. 1–314. DOI: `10.5860/choice.49-6266`.

Peterson, A. T., J. Soberón & V. Sánchez-Cordero (1999). Conservatism of ecological niches in evolutionary time. *Science* 285.5431, pp. 1265–1267. DOI: `10.1126/science.285.5431.1265`.

Pulliam, H. R. (2000). On the relationship between niche and distribution. *Ecology Letters* 3.4, pp. 349–361. DOI: `10.1046/j.1461-0248.2000.00143.x`.

R Core Team (2020). *R: A Language and Environment for Statistical Computing.* R Foundation for Statistical Computing. Vienna, Austria. URL: `https://www.R-project.org/`.

Robert, C. & G. Casella (1999). *Monte Carlo statistical methods.* Springer Science & Business Media.

Root, T. (1988). Environmental Factors Associated with Avian Distributional Boundaries. *Journal of Biogeography* 15.3, p. 489. DOI: `10.2307/2845278`.

Saupe, E. E., V. Barve, C. E. Myers, J. Soberón, N. Barve, C. M. Hensz, A. T. Peterson, H. L. Owens & A. Lira-Noriega (2012). Variation in niche and distribution model performance: The need for a priori assessment of key causal factors. *Ecological Modelling* 237-238, pp. 11–22. DOI: `10.1016/j.ecolmodel.2012.04.001`.

Saupe, E. E., N. Barve, H. L. Owens, J. C. Cooper, P. A. Hosner & A. T. Peterson (2017). Reconstructing ecological niche evolution when niches are incompletely characterized. *Systematic Biology* 67.3, pp. 428–438. DOI: `10.1093/sysbio/syx084`.

Soberón, J. (2007). Grinnellian and Eltonian niches and geographic distributions of species. *Ecology Letters* 10.12, pp. 1115–1123. DOI: `10.1111/j.1461-0248.2007.01107.x`.

Soberón, J. & M. Nakamura (2009). Niches and distributional areas: Concepts, methods, and assumptions. *Proceedings of the National Academy of Sciences of the United States of America* 106.SUPPL. 2, pp. 19644–19650. DOI: `10.1073/pnas.0901637106`.

Soberón, J. & A. T. Peterson (2019). What is the shape of the fundamental Grinnellian niche? *Theoretical Ecology* May. DOI: `10.1007/s12080-019-0432-5`.

Tingley, R., M. Vallinoto, F. Sequeira & M. R. Kearney (2014). Realized niche shift during a global biological invasion. *Proceedings of the National Academy of Sciences* 111.28, pp. 10233–10238. DOI: `10.1073/pnas.1405766111`.

Warren, D. L. (2012). In defense of 'niche modeling'. *Trends in Ecology and Evolution* 27.9, pp. 497–500. DOI: `10.1016/j.tree.2012.03.010`.

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis.* Springer-Verlag New York. URL: `https://ggplot2.tidyverse.org`.

Wickham, H. & D. Seidel (2020). *scales: Scale Functions for Visualization*. R package version

1.1.1. URL: `https://CRAN.R-project.org/package=scales`.

Wilson, T. M., J. Takahashi, S.-E. Spichiger, I. Kim & P. van Westendorp (2020). First Reports of

Vespa mandarinia (Hymenoptera: Vespidae) in North America Represent Two Separate Ma-

ternal Lineages in Washington State, United States, and British Columbia, Canada. *Annals*

of the Entomological Society of America 113.6, pp. 468–472. DOI: `10.1093/aesa/saaa024`.

Zizka, A., A. Antonelli & D. Silvestro (2020). Sampbias, a Method for Quantifying Geographic

Sampling Biases in Species Distribution Data. *Ecography*, pp. 1–8. DOI: `10.1111/ecog.`

`05102`.

Supplementary materials

In the main text, we provided a worldwide suitability map for *V. mandarinia* obtained using the weighted-normal model that we have proposed for estimating the fundamental niche of a species. Here, we provide three additional maps focused on areas of concern for this invasive species: (1) the native range of the species in Figure S1; (2) Europe in Figure S2; and (3) the West Coast of North America in Figure S3. The same color scale used in Figure 6 is used in these figures: dark purple correspond to highly suitable sites where the environmental combinations in E-space are close to the center of the estimated fundamental niche, and light purple corresponds to sites with low suitability where the environmental combinations in E-space are near the border of the fundamental niche.

Figure S1: Suitability index obtained with the weighted-normal model plotted in the native range of *V. mandarinia*. The green squares are the occurrence points used to estimate the fundamental niche of the species.

Figure S2: Suitability index obtained with the weighted-normal model for *V. mandarinia* plotted in Europe. The red square is a confirmed occurrence point in Germany.

Figure S3: Suitability index obtained using the weighted-normal model for *V. mandarinia* plotted for the west coast of Canada and the United States. The red squares are confirmed occurrence points in each country.