TITLE: Artificial Sputum Medium Formulation Impacts *Pseudomonas aeruginosa*

Phenotype and Chemotype

AUTHORS: Rachel L. Neve[1] and Vanessa V. Phelan[2#]

AUTHOR AFFILIATIONS:

[1] Department of Immunology and Microbiology, School of Medicine, University of

Colorado - Anschutz Medical Campus, Department of Immunology and Microbiology,

Aurora, CO, 80045, USA

[2] Department of Pharmaceutical Sciences, Skaggs School of Pharmacy and

Pharmaceutical Sciences, University of Colorado - Anschutz Medical Campus, Aurora,

CO, 80045, USA

CORRESPONDING AUTHOR:

[#] Address correspondence to Vanessa V. Phelan, <u>vanessa.phelan@cuanschutz.edu</u>

ABSTRACT

Artificial sputum medium (ASM) is a class of *in vitro* bacterial culture medium intended

to mimic the nutritional environment of cystic fibrosis (CF) pulmonary mucus. One of the

most commonly studied microbes in ASM is *Pseudomonas aeruginosa,* a prevalent and

dangerous pathogen of the CF pulmonary microbiome. Many ASM formulations have

been reported in literature, with differing nutrient concentrations and availability. Here,

we show that common formulations of ASM yield different phenotypes and chemotypes

of *P. aeruginosa*. Further, we demonstrate that iron in commercial porcine gastric mucin

(PGM) is sufficient to alter production of *P. aeruginosa* siderophores in the chemically

defined ASM, synthetic CF medium 1 (SCFM1). These results highlight that the choice

of ASM formulation for *in vitro* investigations of microbial pathogenicity, physiology, and

interactions should be carefully considered.

IMPORTANCE

In vitro culture media are being developed to resemble the *in vivo* nutritional

environment more closely. These culture media are used to investigate microbial

pathogenicity and ecology in environments that are more reflective of disease states. In

cystic fibrosis (CF), a number of different artificial sputum media (ASM) formulations

have been created to recapitulate the CF lung environment. However, these ASM have

different sources and concentrations of nutrients. Here, we cultured *Pseudomonas*

aeruginosa in nine different formulations of ASM. *P. aeruginosa* is the primary pathogen

causing lung infection in CF.

35   We show that different ASM formulations lead to different phenotypes and chemotypes

36   by *P. aeruginosa* and one component of ASM, mucin, contains high levels of iron, which

37   may affect *P. aeruginosa* physiology.

Cystic fibrosis (CF) is a genetic disease characterized by the accumulation of thick bronchial mucus, which causes decreased mucociliary clearance and enables the growth of pathogenic bacteria and fungi. Pulmonary mucus is a nutritionally distinct environmental niche comprised of mucin, DNA, amino acids, lipids, and micronutrients, including metals (1, 2). To study the physiology of microbial pathogens and communities in the unique CF pulmonary environment, a variety of artificial sputum media (ASM) have been developed (3-14), and applied to investigate microbial physiology (3, 5, 15-17), biofilm morphology (10, 18), antibiotic susceptibility (19, 20), and interspecies interactions in the context of CF (21-23). However, disparate nutrient concentration and bioavailability between ASM formulations may affect microbial growth, metabolism, and virulence (3, 7, 20, 24).

Pseudomonas aeruginosa is one of the most common opportunistic pathogens affecting CF patients, with colonization starting in childhood and affecting more than half of adults (25). Chronic *P. aeruginosa* pulmonary infections cause progressive loss of lung function (26), leading to premature death (25). Due to its importance, *P. aeruginosa* is well-studied in terms of its genome plasticity (27), biofilm growth (28), adaptations (29), development of antibiotic resistance (27, 29-31), and production of virulence factors in the CF lung (32-34). While *P. aeruginosa* produces a vast array of virulence factors, key components of the interactions between *P. aeruginosa* and the environment are its specialized metabolites (3, 4, 6-8, 10, 11), including phenazines (PHZs), quinolones (AQs), rhamnolipids (RHLs), and siderophores. These specialized metabolites function as quorum sensing signals (35), electron shuttles (36), biosurfactants (37), iron scavenging molecules (35, 38-40), and antimicrobials (40-42).

Herein, we investigated the effect of nine ASM formulations on *P. aeruginosa*

PAO1 phenotype and specialized metabolite chemotype. We observed that the

phenotype of PAO1 was greatly altered in disparate ASM formulations. Using molecular

networking (MN) (43, 44) guided analysis of non-targeted metabolomics data, we

determined that all ASM formulations induced the production of a similar suite of

specialized metabolites. However, production of individual specialized metabolites and

summed total of molecular families were statistically different between ASM

formulations. Our results show that PAO1 cultured in ASM formulations with different

nutrient concentration and bioavailability yield different *P. aeruginosa* specialized

metabolism. Further, we show that commercial porcine gastric mucin (PGM), a

component of most ASM formulations, is as an unexpected source of iron, which alters

P. aeruginosa siderophore production and implicates the inclusion of complex

biomolecules in otherwise chemically defined media as sources of other nutrients.

RESULTS AND DISCUSSION

ASM formulation affects *Pseudomonas aeruginosa* phenotype. *In vitro* cultivation of

microbial pathogens has enabled thorough investigation of microbial physiology, gene

expression, and virulence factor production. However, most of these investigations have

been performed in standard laboratory media, which may not reflect the *in vivo*

nutritional environment in which pathogens exist. In cystic fibrosis (CF), an effort to

replicate the *in vivo* nutritional environment of the lung has resulted in the creation of a

wide variety of *in vitro* culture media, broadly referred to as artificial sputum media

(ASM). Most ASM formulations contain the major components of CF sputum, including

mucin, DNA, amino acids, and a lipid source (45). However, the exact composition of

ASM varies between formulations with alterations based upon nutritional measurements

from CF sputum, cited literature, personal observation, or cost of materials.

We hypothesized that culturing *P. aeruginosa* PAO1 in different ASM

formulations would yield phenotypes. To test this hypothesis, PAO1 was cultured in nine

commonly used ASM (3-8, 10, 11) (**Fig. 1 Timeline and composition alterations of**

common ASM formulations, Dataset S1 ASM Formulations). These ASM

formulations can be broken down into two lineages, complex and defined; both of which

have contain the major components of CF sputum (45). The complex ASM lineage is

derived from Soothill ASM (4). As Soothill ASM did not contain amino acids, Sriramulu

et al. complemented the medium with 250 mg/L of each canonical amino acid to

generate Romling ASM (9, 10). Winstanley ASM is compositionally identical to Romling,

except for its buffering capacity (6). ASMDM ASM (3) is modified from Romling ASM by

the addition of bovine serum albumin (BSA), and alterations in mucin and DNA

concentration. Derived from ASMDM ASM, Cordwell ASM (5) has reduced buffering

capacity, replacement of individual amino acids with casamino acids (Cas AA), and

replacement of DTPA with ferritin to replicate CF sputum iron availability more

accurately. SDSU ASM (8) is modified from Cordwell ASM by replacement of Cas AA

with MEM amino acids (MEM AA), removal of BSA, and increasing the mucin

concentration.

Recently, chemically defined Synthetic Cystic Fibrosis Medium (SCFM1) was

developed from the average concentrations of ions, free amino acids, glucose, and

lactate as measured from CF sputum samples (7). SCFM1 was supplemented with

DNA, bovine submaxillary mucin (BSM), *N*-acetyl glucosamine (GlcNAc), and

dioleoylphosphatidylcholine (DOPC) at concentrations similar to those observed in CF

sputum to create SCFM2 (11). SCFM2 was further modified by the addition of

p-aminobenzoic acid (PABA), nicotinamide adenine dinucleotide (NAD$^+$), adenine

(ADE), guanine (GUA), xanthine (XAN), and hypoxanthine (HX) to generate SCFM3, as

experimental results suggested that these metabolites are more available in CF sputum

than in SCFM2 (11). We used PGM was used as the mucin source as it is a cost

effective and common replacement of BSM (12-14).

PAO1 grew in all media (**Fig. S1 Growth of PAO1 in ASM**); however, both the

gross morphology and culture color varied between ASM formulations (**Fig. 2A PAO1**

phenotype and metabolome in ASM and LB). In Soothill ASM, the cultures were

turbid, but, unlike the other ASM formulations, no macrostructures were visible. The

Romling and Winstanley formulations both induced a brown macrostructure phenotype.

In ASMDM ASM, *P. aeruginosa* grew heterogeneously with a layer of growth at the air-

liquid interface, giving way to turbid growth further down the culture. In SDSU ASM,

PAO1 formed a green, floating structure near the center of the well that reached the

surface of the medium. PAO1 grew as a turbid culture with bright green coloration in

SCFM1; while in SCFM2 and SCFM3, PAO1 growth was more structured with a

distinctly blue color.

***P. aeruginosa* produces similar specialized metabolites in all ASM formulations.**

As metabolism often underlies phenotypic changes, we hypothesized that different ASM

formulations induced different chemotypes of PAO1. *P. aeruginosa* has a well

characterized specialized metabolome chemical signals produced by microorganisms to

interact with their environment (46, 47)), including six molecular families: homoserine

lactones (HSLs), rhamnolipids (RHLs), alkyl quinolones (AQs), phenazines (PHZs), and

the siderophores pyochelin (PCH) and pyoverdine (PVD) (48-50). These compounds

are chemically and functionally diverse and include quorum sensors, small molecule

virulence factors, and natural products. *P. aeruginosa* specialized metabolites have

been detected in CF sputum (32-34) and associated with pathogenicity and virulence

(24, 50, 51).

To determine if different ASM formulations induced the production of different

molecular families by *P. aeruginosa*, we analyzed non-targeted liquid chromatography

tandem mass spectrometry (LC-MS/MS) data of PAO1 grown in ASM and LB using the

classical molecular networking (CMN) workflow of the GNPS platform. Briefly, CMN

organizes metabolites into molecular families using MS/MS spectral similarity as a proxy

for structural similarity (43). CMN is best used for applications assessing the presence

or absence of a molecular signal in an experiment (43, 52, 53). Of the 1,824 nodes in

the network, 220 (12%) had spectral matches to the GNPS libraries, representing 120

unique compounds. These compounds included members of the PHZ, AQ, RHL

molecular families as well as PCH (**Fig. 2B PAO1 phenotype and metabolome in**

ASM and LB).

Closer examination of the CMN results revealed the presence of duplicate nodes

representing the same compound. Therefore, to generate a more accurate assessment

of unique MS1 mass-to-charge (*m/z)* values shared between cultures of PAO1, nodes

with the same precursor *m/z* value were combined and a binary presence/absence table

was created. Of 1,165 unique *m/z* values, 140 were present in all cultures (**Fig. 2C**

PAO1 phenotype and metabolome in ASM and LB), including values corresponding

to the PHZ, AQ, and RHL molecular families, suggesting that all ASM formulations have

the nutritional resources for PAO1 to produce these compounds. In addition to *m/z*

values shared between all cultures of PAO1, a number of *m/z* values were detected in

samples of PAO1 grown in specific media. PAO1 cultures in Soothill ASM had the most

m/z values unique to a single medium (**Fig. 2D PAO1 phenotype and metabolome in**

ASM and LB). GNPS library matches of these Soothill ASM associated *m/z* values

largely represent fragmentation of media components, likely due to relative higher

intensity of these peaks compared to other ASM formulations. The production levels of

PAO1 metabolites of Soothill ASM cultures did not require dilution to prevent saturation

of the MS detector.

ASM formulation alters the production levels of *P. aeruginosa* specialized

metabolites. As PAO1 specialized metabolite presence was generally conserved

across all ASM formulations, we applied the feature based molecular networking

(FBMN) to quantify and visualize metabolite production levels of PAO1 in the different

ASM formulations. FBMN integrates feature finding, a data reduction step in LC-MS

analysis, with MN (44). Feature finding generates a quantitative representation of all

metabolites within a metabolomics dataset by defining a feature as an *m/z*, retention

time (RT), and corresponding abundance. In FBMN, LC-MS/MS data is processed

feature finding to generate abundance from MS1 data, and one representative MS/MS

spectrum is generated for each feature (*m/z*-RT pair). Like in CMN, MS/MS spectra

from an experiment are organized into molecular families. However, in FBMN,

abundance data is imported into the molecular network as pie charts to provide rapid

visualization of differential abundance between metadata categories, such as ASM

formulation. Of the 492 nodes in the network, 80 (16%) had matches to GNPS library

spectra, representing 60 unique compounds. Statistical analysis of the feature table

revealed that the abundance of only two features were not statistically different between

PAO1 cultures in ASM formulations (**Dataset S2 Statistics of ASM Metabolism**

Results). *P. aeruginosa* production of individual and total PHZs, AQs, RHLs, PCH, and

PVDs, were all altered in different ASM formulations. Notably, HSLs were not detected

under our conditions.

PHZs are redox active virulence factors with various biological activities,

including promoting iron uptake (36, 40), anaerobic respiration (36), biofilm formation

(54, 55), and interspecies interactions (40, 48, 56, 57), among others (58-60). We

quantified four PHZs: 1-hydroxyphenazine (1-HP), pyocyanin (PYO), phenazine-1-

carboxamide (PCN), and phenazine-1-carboxylic acid (PCA) (**Fig. 3A Phenazines**).

While PHZs were detected in all PAO1 cultures, total PHZ production by PAO1 varied

between formulations with lowest production in Soothill ASM and highest production in

SCFM2 and SCFM3 (**Fig. 3B Phenazines**). The low levels of PHZs produced by PAO1

in Soothill ASM is likely due to the absence of amino acids in the medium. In addition to

variations in total PHZ production, the production of individual PHZs varied. In Soothill

ASM, PAO1 produced primarily PCA. PCA has been implicated in inducing *P.*

aeruginosa biofilm formation by enabling the reduction of Fe(III) to Fe(II) in a *P.*

aeruginosa mutant unable to produce the siderophores PVD and PCH (61). As Soothill

ASM is amino acid limited and restricts the precursor availability for siderophore

biosynthesis by *P. aeruginosa* (62-64), PCA may be produced to acquire iron from the

medium. *P. aeruginosa* produces higher levels of 1-HP in the Romling, Winstanley,

ASMDM, Cordwell, and SDSU ASM formulations. PYO is produced highest in SCFM2

and SCFM3, which contributes to the characteristic blue phenotype of these cultures.

Interestingly, despite similar production ratios of PHZs in the SCFM series of ASM (**Fig.**

3C Phenazines), the total PHZ production by PAO1 in SCFM1 is significantly lower

than in SCFM2 and SCFM3 (mean PHZ AUC SCFM1=3.28×10^4, SCFM2=1.38×10^5,

SCFM3=1.34×10^5, $p<0.0001$). SCFM2 and SCFM3 (11) are compositionally different

from SCFM1 by the addition of several components, including GlcNAc. An increase of

PYO production by *P. aeruginosa* was measured spectrographically upon addition of

GlcNAc (65). GlcNAc likely induces PYO production through a mechanism to sense and

respond to peptidoglycan*.*

Like PHZ production, *P. aeruginosa* production of individual and total AQs varied

depending upon ASM formulation. We quantified forty-three AQs from the PAO1 ASM

cultures and only one could not be annotated (**Fig. 4A Quinolones**). Total AQ

production by *P. aeruginosa* was highest in Romling, Winstanley, and ASMDM ASM

formulations, all at similar levels (mean = ~4×10^5), and lowest in Soothill and SDSU

ASM (**Fig. 4B Quinolones**). Different AQs have different roles in intra- and interspecies

interactions within the CF lung environment (66-70). 2-heptyl-4-hydroxyquinolone-N-

oxide (HQNO) is a potent inhibitor of the respiratory chain of microbial and host cells

(66, 67). 2-heptyl-4-hydroxyquinolone (HHQ) is an autoinducer of the *pqs* operon (68),

and 2-heptyl-3,4-dihydroxyquinolone (Pseudomonas Quinolone Signal, PQS)

potentiates HSL-dependent QS (69), biofilm formation (70), and siderophore production

(68). AQs were classified as HHQ-type, PQS-type, or HQNO-type based on their

MS/MS fragmentation patterns (56, 71). AQ production by PAO1 in all ASM

formulations was predominantly HHQ-type AQs (~44-76%), while HQNO-type AQs

accounted for ~15-41%, and PQS-type AQs were the consistently the least abundant

(~3-17%) (**Fig. 4C Quinolones**).

225   RHLs have functional roles in biofilm formation and maintenance (72, 73), motility

as biosurfactants (37), competition with fungal neighbors (41), and evasion of the host

immune response (74, 75). Six RHLs were quantified (**Fig. 5A Rhamnolipids**). Total

PAO1 RHL production was lower in Romling, SCFM2, and SCFM3 (means = $2.9x10^4$,

$1.8x10^4$, $2.2x10^4$, respectively) ASM formulations compared to the other versions of

ASM (range of means = $3.6x10^4$ - $4.0x10^4$) (**Fig. 5B Rhamnolipids**). The proportion of

mono- to di-RHL production was maintained across cultures in all ASM, with 30.0 -

43.8% mono-RHLs. The predominant fatty acid length was C10, accounting for 56.2 -

70.0% of RHLs produced (**Fig. 5C Rhamnolipids**). The production of RHLs is

dependent on the *rhl* QS system; specifically the binding of the autoinducer C4-HSL,

produced by RhlI, to the transcription factor RhlR (76). The RhlI/RhlR QS system is

activated by the LasI/R QS system and directs the regulation of many virulence

encoding genes (77, 78). Lower production levels of RHLs by PAO1 may indicate

altered QS signaling, which may affect the production of other virulence factors.

239   Strikingly, production of siderophores by *P. aeruginosa* was almost exclusive to

SCFM1 cultures (**Fig. 6 Siderophores**). Siderophores are metal scavenging virulence

factors (79, 80), secreted in low iron conditions to compete for environmental iron (81).

P. aeruginosa produces two siderophores, PVD and PCH (38), with PVD having a

higher affinity for iron than PCH (38). While PCH was identified in SCFM1 cultures

based upon a match to GNPS libraries, the nodes corresponding to the PVD molecular

family were prioritized for annotation based upon coloration of the pie chart, which

indicated high production by PAO1 in SCFM1 (**Fig. 6A Siderophores**). Although PVD

biosynthesis was reported to be upregulated in SCFM1 (7), we were surprised to detect

PVDs as the production of this molecular family by PAO1 is below our limit of detection

when cultured in most media. The rapid visualization of molecular families and their

production levels in FBMN enabled us to identify this molecular family in a non-biased

manner, when we were admittedly not expecting to observe its production.

Addition of commercial PGM to SCFM1 reduces PAO1 siderophore production

due to the presence of iron. As the production of PCH and PVD is triggered by low

environmental iron (82), the production of PCH and PVD by *P. aeruginosa* in SCFM1

was surprising as all ASM formulations contain similar concentrations of added iron.

Soothill ASM was measured to have 9.5 µM Fe^{2+} prior to addition of DTPA (15 µM) as

an iron chelator (4). Both Cordwell and SDSU ASM contain 0.000003% (w/v) ferritin,

which adds ~2.85 µM Fe^{2+} to the media. Furthermore, SCFM1, SCFM2, and SCFM3

formulations contain 3.60 µM Fe^{2+}. Taken together, the inability to detect siderophores

after PAO1 growth in all ASM except SCFM1 suggests these formulations contain an

undefined source of iron.

To replicate the CF lung environment more closely, SCFM1 was modified by the

addition of DNA, DOPC, GlcNAc, and mucin to create SCFM2 (11). As PGM was the

most complex additive to our SCFM2 formulation, we hypothesized that commercial

PGM was a source of contaminating iron in ASM. Mucins are a critical component of *in*

vitro culture media for modeling the microbial interactions of CF lung infections (13, 83,

84), as well as other mucosal body sites (85, 86). The mucin concentration of CF

sputum ranges from 16-30 mg/mL (1). In the lung, mucin is a source of nutrients for

inhabiting microbes (2, 87, 88) and also is involved in microbial adhesion (87),

aggregation (89, 90) and dispersion (83).

To test if the addition of PGM to SCFM1 altered siderophore production by *P.*

aeruginosa, PAO1 was cultured in SCFM1 and SCFM1 complemented with 0.5% (w/v)

PGM (**Fig. S2 Growth of PAO1 in SCFM1 and SCFM1 with PGM**). Quantification of

PCH and PVD molecular families revealed that the addition of PGM to SCFM1 was

sufficient to reduce siderophore production by *P. aeruginosa* (**Fig. 7A, 7B PGM iron**

content and impact on PAO1 siderophores). As measured by ICP-MS, 0.5% (w/v) of

PGM contains 30.6 μM iron, which is 8.5 times higher than the concentration of iron

added to the SCFM1 base medium (3.6 μM) (**Fig. 7C PGM iron content and impact**

on PAO1 siderophores). Due to increased PGM concentration, ASMDM and Cordwell

ASM contain ~60 μM iron and SDSU ASM contains ~117 μM.

Importance of ASM formulation on *in vitro* investigations. The differential

production of specialized metabolites by PAO1 grown in different ASM has important

implications for the interpretation of experimental results performed in this class of

media. It is well established that *P. aeruginosa* specialized metabolites play important

roles in virulence (36, 38), microbial interactions (40-42), and nutrient acquisition from

the environment (35, 38-40). While the conclusions from a single dataset are valid,

comparing the results between experiments performed in different ASM formulations

should be carefully considered. For example, PYO is produced in highest abundance by

PAO1 in SCFM2 and SCFM3. PYO is known to affect respiratory epithelial functions,

including cellular respiration, electron transport, and innate immune response

mechanisms (60, 91-94). If the culture supernatants of PAO1 in SCFM2 and SDSU

ASM were used to investigate the effect of PAO1 exometabolome on lung epithelial

cells, the results of those experiments would likely vary significantly due to the

differential production of PYO.

Similarly, many of the *P. aeruginosa* specialized metabolites are important in

interspecies microbial interactions. The PHZ 1-HP is important in interactions between

P. aeruginosa and the fungal pathogen *Aspergillus fumigatus* (40, 95). Treating *A.*

fumigatus cultures with supernatants from PAO1 grown in Romling ASM and SCFM2

would likely lead to different results. The *P. aeruginosa* AQ HQNO is known to alter *S.*

aureus susceptibility to antibiotics (96). Therefore, differential production of HQNO by *P.*

aeruginosa in different ASM could affect the results of antibiotic susceptibility assays

performed in different ASM formulations.

Last, siderophores are important for iron competition *in vivo* and loss of

siderophore production affects the production of other specialized metabolite molecular

families by *P. aeruginosa* (97). Inclusion of PGM in SCFM1 increases the concentration

of iron in the medium to a more clinically relevant level of iron (98-100), leading to

decreased production of PCH and PVD. However, the distinct iron sources in PGM are

unknown and may not represent those in the CF lung (99, 101). If investigators desire to

test iron concentration or source on *P. aeruginosa* and other microbes, PGM inclusion

and concentration in the medium may influence the outcome of their experiments.

CONCLUSION

In vitro model systems are attractive due to their broad accessibility and ability to probe

hypotheses thoroughly. The foundation of an *in vitro* investigation is the medium. In CF,

a broad suite of different ASM formulations has been created. In this study, we

demonstrated that differences in nutritional content between ASM formulations alter the

phenotype and chemotype of *P. aeruginosa* PAO1. Importantly, the concentration of

specialized metabolites, including members of the AQ, PHZ, RHL, and siderophore

molecular families, are impacted by ASM formulation. Given the role of these

metabolites in virulence, differences in ASM formulation may influence the overall

outcome of interactions among *P. aeruginosa*, the host, and other members of the CF

pulmonary microbiota. Commercial PGM was identified as a source of contaminating

bioavailable iron, influencing siderophore production by *P. aeruginosa*. In view of the

importance of mucin in replicating mucosal infection environments *in vitro,* further work

is needed to understand the contribution of commercial mucin to bioavailable nutrients

in culture media more fully. Therefore, careful consideration should be used when

comparing results across studies employing different ASM formulations.

MATERIALS AND METHODS

General: *Pseudomonas aeruginosa* PAO1 (MPAO1) was acquired from the University

of Washington, Seattle (102). All media components, concentrations, and commercial

sources for artificial sputum media (ASM) formulations are listed in **Dataset S1 ASM**

formulations (3-11). Miller Luria broth (LB) and porcine gastric mucin type III (PGM)

were purchased for Millipore Sigma. Organic solvents for extraction and liquid

chromatography tandem mass spectrometry (LC-MS/MS) analysis were purchased from

VWR (ethyl acetate (EtOAc), HiPerSolv Chromanorm purity) and Fisher Scientific

(methanol (MeOH), acetonitrile (ACN), formic acid (FA), and water; all Optima LC/MS

grade). PA Mix was created as a set of external standards for *P. aeruginosa* specialized

metabolite annotation. PA Mix contains 10 µM of each of the following:

1-hydroxyphenazine (1-HP, Tokyo Chemical Industry); pyocyanin (PYO, Sigma);

phenazine-1-carboxylic acid (PCA, Ark Pharm); phenazine-1-carboxamide (PCN, Ark

Pharm); 2-heptylquinolin-4(1H)-one (HHQ, Ark Pharm); 2-heptyl-4-hydroxyquinoline *N*-

oxide (HQNO, Cayman Chemical); 2-heptyl-3-hydroxyl-4-quinolone (Pseudomonas

Quinolone Signal, PQS, Chemodex); N-butyryl-l-homoserine lactone (C4-HSL, Cayman

Chemical); N-(3-oxododecanoyl)-l-homoserine lactone (3-oxo-C12-HSL, Millipore

Sigma); and a rhamnolipid mixture (RHL, AGAE Technologies, 90% pure).

Identification of commonly used ASM formulations: The Web of Science Core

Collection (Clarivate Analytics) was searched for papers with the 'TOPIC' terms:

"sputum medium", "sputum media", "cystic fibrosis medium", "cystic fibrosis media",

"sputa media", "sputa medium", "CF media", or "CF medium" in the title, abstract, or

keywords fields on May 10, 2020. All terms were searched as shown, including

quotation marks. The results of this search were analyzed using the Web of Science

tools 'Create Citation Report' and 'Analyze Results'. The Analyze Results tool was used

to download data corresponding to Web of Science Categories. The citation report was

used to select ASM formulations that were commonly cited.

Preparation of media: LB broth was prepared according to manufacturer instructions.

Commercial PGM was suspended in 1X 3-morpholinopropane-1-sulfonic acid (MOPS)

buffer (pH 7.0) to make a 10% (w/v) suspension, sterilized using a liquid autoclave cycle

(<20 minute sterilization time), and an aliquot was checked for sterility before use. ASM

formulations were prepared as reported (3-11) with the following modifications.

Antibiotics were not included in any medium. For the Cordwell formulation (5), the

reported 0.1% (w/v) agar was excluded. For SCFM2 and SCFM3 (11), bovine

submaxillary mucin (BSM) was replaced with PGM at the published concentration, a

common substitution (12-14). Each ASM formulation was brought to the published pH

(3-11). All media was stored in the dark at 4°C, checked for sterility prior to use, and

used within one month of preparation.

Cultivation of *P. aeruginosa* in LB and ASM formulations: PAO1 was inoculated

from a streak plate into 5 mL LB broth and incubated overnight at 37°C, shaking at 220

RPM. 1.98 mL of LB or ASM were inoculated with 20 µL of *P. aeruginosa* LB broth

culture (OD_{600} of 0.05; ~10^6 CFU/mL final concentration) in polystyrene 24 well plates

(6). For assessing *P. aeruginosa* growth and specialized metabolite production, each

medium was evaluated in its own 24 well plate (12 medium control wells and 12 growth

wells per medium). The media were inoculated and processed in three batches,

comprised of 2-4 ASM formulations and a parallel LB plate as a control. All plates were

covered and incubated statically at 37°C for 72 hours. For assessing the addition of

PGM to SCFM1 on *P. aeruginosa* growth and specialized metabolite production, *P.*

aeruginosa was cultured in SCFM1 and PGM-SCFM1 in a single polystyrene 24 well

plate (4 medium control wells and 4 growth wells per medium).

Processing of *P. aeruginosa* cultures in LB and ASM formulations: Gross

phenotypes of *P. aeruginosa* growth were photographed using a top-down view with a

Stemi 508 stereo microscope with an Axiocam 105 color camera (Zeiss). Following

visualization, 100 µL cellulase (*Aspergillus niger*, 100 mg/mL in sterile water; Sigma)

was added to each well and aggregates were mechanically disrupted by pipetting using

wide-bore pipette tips (USA Scientific), followed by water bath sonication (Branson) for

10 minutes, and a secondary mechanical disruption by pipetting using wide-bore pipette

tips. Disrupted samples were aliquoted for measurements of growth (CFUs), pH, and

metabolomics analysis. Three representative biological replicates from each culture

condition were serially diluted, spotted onto LB agar, incubated at 37°C overnight, and

counted to determine CFU/mL (**Fig. S1 Growth of PAO1 in ASM**).

Processing of *P. aeruginosa* cultures in SCFM1 and PGM-SCFM1: Gross

phenotypes of *P. aeruginosa* growth were photographed using a top-down view with a

Stemi 508 stereo microscope with an Axiocam 105 color camera (Zeiss). Following

visualization, aggregates were mechanically disrupted by pipetting using wide-bore

pipette tips (USA Scientific). Disrupted samples were aliquoted for measurements of

growth (CFUs), pH, and metabolomics analysis. All four biological replicates from each

culture condition were serially diluted, spotted onto LB agar, incubated at 37°C

overnight, and counted to determine CFU/mL (**Fig. S2 Growth of PAO1 in SCFM1**

with and without PGM).

Sample preparation for LC-MS/MS: Each sample aliquot was chemically disrupted

with an equal volume of 1:1 EtOAc:MeOH. The samples were dried, resuspended in

100% MeOH containing 1 µM glycocholic acid (Calbiochem, 100.1% pure), diluted as

needed in 100% MeOH containing 1 µM glycocholic acid, and centrifuged for 10 min at

4,000 RPM (Thermo Sorvall ST 40R) to remove non-soluble particulates. Samples were

diluted as follows for measurement of *P. aeruginosa* specialized metabolite production

in ASM formulations and LB: not diluted: Soothill ASM; 5-fold dilution: Romling ASM,

Cordwell ASM, SCFM1, SCFM2, and SCFM3; 10-fold dilution: Winstanley ASM,

ASMDM, SDSU ASM, and LB. All samples for measurement of *P. aeruginosa*

specialized metabolite production in SCFM1 and SCFM + PGM were diluted 10-fold.

LC-MS/MS analysis: Mass spectrometry data acquisition for all samples was

performed using a Bruker Daltonics Maxis II HD qTOF mass spectrometer equipped

with a standard electrospray ionization (ESI) source. The mass spectrometer was tuned

by infusion of Tuning Mix ESI-TOF (Agilent Technologies) at a 3 µL/min flow rate. For

accurate mass measurements, a wick saturated with Hexakis(1H,1H,2H-difluoroethoxy)

phosphazene ions (Apollo Scientific, *m/z* 622.1978) located within the source was used

as a lock mass internal calibrant. Samples were introduced by an Agilent 1290 UPLC

using a 10 µl injection volume. Extracts were separated using a Phenomenex Kinetex

2.6 µm C18 column (2.1 mm x 50 mm) using a 9 minute, linear water-ACN gradient

(from 98:2 to 2:98% water:ACN) containing 0.1% FA at a flow rate of 0.5 mL/min. The

mass spectrometer was operated in data dependent positive ion mode, automatically

switching between full scan MS and MS/MS acquisitions. Full scan MS spectra (*m/z* 50

- 1500) were acquired in the TOF and the top five most intense ions in a particular scan

were fragmented via collision induced dissociation (CID) using the stepping function in

the collision cell. LC-MS/MS data for PA mix were acquired under identical conditions.

Bruker Daltonics CompassXport was used to apply lock mass calibration and convert

the LC-MS/MS data from .d format to .mzXML format.

Classical molecular networking (CMN) of *P. aeruginosa* specialized metabolism in

LB and ASM formulations: A classical molecular network was created from the

.mzXML files using the Molecular Networking workflow (version release 22) of the

GNPS web platform (gnps.ucsd.edu,

https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=92a1c1f712904db4961fe0ff0dee3c2

b) (43). Briefly, the data was filtered by removing all MS/MS fragment ions ± 17 Da of

the precursor *m/z*. MS/MS spectra were window filtered by choosing only the top 6

fragment ions in each ± 50 Da window throughout the spectrum. The precursor ion

mass tolerance was set to 0.1 Da and the MS/MS fragment ion tolerance to 0.2 Da. A

network was then created where edges were filtered to have a cosine score above 0.7

and at least 5 matched fragment peaks between corresponding nodes. Further, edges

between two nodes were kept in the network only if each of the nodes appeared in each

other's respective top 10 most similar nodes. Finally, the maximum size of a molecular

family was set to 100, and the lowest scoring edges were removed from molecular

families until the molecular family was below this threshold. The spectra in the network

were then searched against the GNPS spectral libraries. The library spectra were

filtered in the same manner as the input data. All matches kept between the network

spectra and library spectra were required to have a cosine score above 0.6 and at least

5 matched peaks. The molecular network was visualized using Cytoscape (version

3.7.1) (103).

Creation of UpSet plots from CMN results. The network table generated *via* CMN

was downloaded from GNPS. As duplicate nodes representing the same metabolite

were present in the network, precursor *m/z* values were rounded to the nearest

hundredths and nodes with the same *m/z* values were combined to generate a list of

unique *m/z* values. A binary presence/absence table was created to reflect whether

MS/MS spectra corresponding to each *m/z* value was detected in each medium. The

binary table was visualized using the UpSetR Shiny App or in R using the UpSet

package (104, 105).

Feature based molecular networking (FBMN) of *P. aeruginosa* specialized

metabolism in LB and ASM formulations: MZmine (version 2.53) (106) was used to

perform feature finding on the .mzXML files. The output was a feature table containing

area under the curve (AUC) integration values for each feature (*m/z*-RT pair) and an

.mgf file containing MS/MS spectra for each feature. The feature table was corrected for

sample dilutions prior to submission to GNPS. A metadata table was created containing

sample file names and attributes according to GNPS specifications. The Feature

Networking workflow (version release 22) (44) was applied to the dilution-corrected

feature table, .mgf file, and metadata file *via* the GNPS web platform (gnps.ucsd.edu,

https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=baba5df5ab1b437980f81b4e64bcffc

0) (43). Briefly, the data was filtered by removing all MS/MS fragment ions ± 17 Da of

the precursor *m/z*. MS/MS spectra were window filtered by choosing only the top 6

fragment ions in each ± 50 Da window throughout the spectrum. The precursor ion

mass tolerance was set to 0.05 Da and the MS/MS fragment ion tolerance was set to

0.1 Da. A network of nodes connected by edges was then created. Edges were filtered

to have a cosine score above 0.75. Further, edges between two nodes were kept in the

network only if each of the nodes appeared in each other's respective top 5 most similar

nodes. Finally, the maximum size of a molecular family was set to 50, and the lowest

scoring edges were removed from molecular families until the molecular family was

below this threshold. The spectra in the network were then searched against the GNPS

spectral libraries. The library spectra were filtered in the same manner as the input data.

All matches kept between the network spectra and library spectra were required to have

a cosine score above 0.6 and at least 5 matched peaks. Sum normalization and mean

aggregation were applied. The molecular network was visualized using Cytoscape

(version 3.7.1) (103).

Feature annotation of *P. aeruginosa* specialized metabolites from FBMN results:

All annotated features corresponding to *P. aeruginosa* specialized metabolites identified

from FBMN analysis are listed in **Table S1. ASM metabolite annotation and mass**

defect. Annotations of features corresponding to commercial standards (level 1

annotation (107)) were confirmed by comparing the experimental data (exact mass,

MS/MS, and retention time) with data acquired for the compound in PA mix using

Bruker Daltonics DataAnalysis v4.1 (Build 362.7). Putative annotation of specialized

metabolites based on matches to the GNPS libraries (level 2 annotation (107)) and

molecular families (level 3 annotation (107)) were validated by comparing the

experimental data (exact mass, MS/MS) to reported data (56) and putative structures,

respectively (56, 71) (**Fig. S3 Chromatograms of PVD family molecules from PAO1**

in SCFM1).

Metabolite, Molecular Family, and Sub-structure Class Annotation: Multiple

features (different adducts, in-source fragments, and/or in-source dimers) can represent

the same metabolite. Therefore, sum normalized AUC values for features

corresponding to the same metabolite of interest (identified by co-elution and mass

defect from predicted exact mass of the compound adduct, fragment, and/or dimer)

were summed to provide the total AUC for a metabolite. Sum normalized AUC values

for metabolites representing the same sub-structure class based upon MS/MS

fragmentation pattern (e.g. HHQ-type AQs or C10-C10 containing RHLs) were summed

to provide the total AUC for a sub-structure class. Sum normalized AUC values for all

identified metabolites representing the same molecular family based upon MS/MS

fragmentation patterns (e.g. AQs or RHLs) were summed to provide the total AUC for a

molecular family**.**

Targeted quantitation of *P. aeruginosa* siderophores in SCFM1 and PGM-SCFM1

cultures. MZmine (version 2.53) (106) was used to perform feature finding on the

.mzXML files for *P. aeruginosa* growth in SCFM1 and PGM-SCFM1. Sum normalized

AUC values for features corresponding to PCH, PVD, or their biosynthetic relatives

(identified by co-elution and mass defect from predicted exact mass of the compound

adduct, fragment, and/or dimer) were summed to provide the total AUC for each

metabolite (**Table S2 SCFM1 and mucin siderophore annotation and mass defect**).

Sum normalized AUC values for all identified metabolites representing the same

molecular family based upon MS/MS fragmentation patterns were summed to provide

the total AUC for the PCH or PVD molecular families.

ICP-MS analysis of PGM: Metals were quantified from 2.5% (w/v) PGM suspended in

1X MOPS using inductively coupled plasma mass spectrometry (ICP-MS) analysis at

the University of Nebraska - Lincoln Spectroscopy and Biophysics Core facility.

Samples were diluted 1:1 with nitric acid, digested overnight, and diluted an additional

10-fold before analysis conducting ICP-MS (Agilent 7500 cx). Gallium (Ga), 50 ppb, was

added to each sample as an internal standard. Samples were analyzed in technical

triplicate with a blank wash run between each sample. Only concentrations above the

limit of quantification were retained. To determine the amount of iron added to media by

the addition of PGM, or 1X MOPS when used in place of PGM (**Fig. 7 PGM iron**

content and impact on PAO1 siderophores), the concentration of PGM was brought

to that present in most ASM formulations (0.5% w/v) and converted to micromolar

concentration.

Statistical analysis: Statistical comparison of FBMN features of *P. aeruginosa* cultured

in ASM formulations was conducted on the sum normalized feature table (**Dataset S2**

ASM statistical analysis) using ANOVA with Tukey's correction for multiple

comparisons in MetaboAnalyst 4.0 (108-111) (**Dataset S2 ASM statistical analyses**).

Statistical comparison of individual metabolites, sub-structure classes, and molecular

families performed using ANOVA with Tukey's multiple comparisons correction in

GraphPad Prism (version 8.4.0) (**Dataset S2 ASM statistical analyses**). Statistical

comparison of PCH and PVD production by *P. aeruginosa* in SCFM1 and SCFM1+PGM

was conducted using a two tailed, unpaired t-test with Welch's correction for unequal

standard deviations in GraphPad Prism (version 8.4.3). For all analyses, *P* values less

than 0.05 were considered statistically significant.

Data availability: All mass spectrometry data including the raw files, .mzXML files,

metadata table, and Cytoscape files are available via MassIVE (accession numbers:

MSV000086721 (LB and ASM formulations – media control, *P. aeruginosa* culture, QC,

and PA mix data) and MSV000086723 (SCFM1 and PGM-SCFM1 – media control, *P.*

aeruginosa culture, QC, and PA mix data)).

ACKNOWLEDGEMENTS:

We are very grateful to Mingxun Wang (University of California, San Diego) for

providing input and beta versions of the networking software for metabolomics analysis;

REFERENCES

1. Sanders NN, Van Rompaey E, De Smedt SC, Demeester J. 2001. Structural Alterations of Gene Complexes by Cystic Fibrosis Sputum. American Journal of Respiratory and Critical Care Medicine 164:486-493.

2. Schwab U, Abdullah LH, Perlmutt OS, Albert D, Davis CW, Arnold RR, Yankaskas JR, Gilligan P, Neubauer H, Randell SH, Boucher RC. 2014. Localization of Burkholderia cepacia complex bacteria in cystic fibrosis lungs and interactions with Pseudomonas aeruginosa in hypoxic mucus. Infection and Immunity 82:4729-4745.

3. Fung C, Naughton S, Turnbull L, Tingpej P, Rose B, Arthur J, Hu HH, Harmer C, Harbour C, Hassett DJ, Whitchurch CB, Manos J. 2010. Gene expression of Pseudomonas aeruginosa in a mucin-containing synthetic growth medium mimicking cystic fibrosis lung sputum. Journal of Medical Microbiology 59:1089-1100.

4. Ghani M, Soothill JS. 1997. Ceftazidime, gentamicin, and rifampicin, in combination, kill biofilms of mucoid Pseudomonas aeruginosa. Canadian Journal of Microbiology 43:999-1004.

5. Hare NJ, Soe CZ, Rose B, Harbour C, Codd R, Manos J, Cordwell SJ. 2012. Proteomics of Pseudomonas aeruginosa Australian Epidemic Strain 1 (AES-1)

Cultured under Conditions Mimicking the Cystic Fibrosis Lung Reveals Increased

Iron Acquisition via the Siderophore Pyochelin. Journal of Proteome Research

11:776-795.

6.   Kirchner S, Fothergill JL, Wright EA, James CE, Mowat E, Winstanley C. 2012.

Use of Artificial Sputum Medium to Test Antibiotic Efficacy Against Pseudomonas

aeruginosa in Conditions More Relevant to the Cystic Fibrosis Lung. Jove-

Journal of Visualized Experiments doi:ARTN e3857 10.3791/3857.

7.   Palmer KL, Aye LA, Whiteley M. 2007. Nutritional cues control Pseudomonas

aeruginosa multicellular Behavior in cystic fibrosis sputum. Journal of

Bacteriology 189:8079-8087.

8.   Quinn RA, Whiteson K, Lim YW, Salamon P, Bailey B, Mienardi S, Sanchez SE,

Blake D, Conrad D, Rohwer F. 2015. A Winogradsky-based culture system

shows an association between microbial fermentation and cystic fibrosis

exacerbation (vol 9, pg 1024, 2015). Isme Journal 9:1052-1052.

9.   Sriramulu DD. 2010. Artifical Sputum Medium. Protocol Exchange

doi:doi:10.1038/protex.2010.212.

10.   Sriramulu DD, Lunsdorf H, Lam JS, Romling U. 2005. Microcolony formation: a

novel biofilm model of Pseudomonas aeruginosa for the cystic fibrosis lung.

Journal of Medical Microbiology 54:667-676.

11.   Turner KH, Wessel AK, Palmer GC, Murray JL, Whiteley M. 2015. Essential

genome of Pseudomonas aeruginosa in cystic fibrosis sputum. Proceedings of

the National Academy of Sciences of the United States of America 112:4110-

4115.

12.   Homa M, Sandor A, Toth E, Szebenyi C, Nagy G, Vagvolgyi C, Papp T. 2019. In

vitro Interactions of Pseudomonas aeruginosa With Scedosporium Species

Frequently Associated With Cystic Fibrosis. Front Microbiol 10:441.

13.   Schick A, Kassen R. 2018. Rapid diversification of Pseudomonas aeruginosa in

cystic fibrosis lung-like conditions. Proc Natl Acad Sci U S A 115:10714-10719.

14.   Wong A, Rodrigue N, Kassen R. 2012. Genomics of adaptation during

experimental evolution of the opportunistic pathogen Pseudomonas aeruginosa.

PLoS Genet 8:e1002928.

15.   Behrends V, Geier B, Williams HD, Bundy JG. 2013. Direct Assessment of

Metabolite Utilization by Pseudomonas aeruginosa during Growth on Artificial

Sputum Medium. Applied and Environmental Microbiology 79:2467-2470.

16.   Darch SE, Simoska O, Fitzpatrick M, Barraza JP, Stevenson KJ, Bonnecaze RT,

Shear JB, Whiteley M. 2018. Spatial determinants of quorum signaling in a

<i>Pseudomonas aeruginosa</i> infection model. Proceedings of the National

Academy of Sciences 115:4779-4784.

17.   Behrends V, Ebbels TMD, Williams HD, Bundy JG. 2009. Time-Resolved

Metabolic Footprinting for Nonlinear Modeling of Bacterial Substrate Utilization.

Applied and Environmental Microbiology 75:2453-2463.

18.   Haley CL, Colmer-Hamood JA, Hamood AN. 2012. Characterization of biofilm-

like structures formed by Pseudomonas aeruginosa in a synthetic mucus

medium. BMC Microbiology 12:181.

19.   Diaz Iglesias Y, Van Bambeke F. 2020. Activity of Antibiotics against

Pseudomonas aeruginosa in an *In Vitro* Model of Biofilms in the

Context of Cystic Fibrosis: Influence of the Culture Medium. Antimicrobial Agents

and Chemotherapy 64.

20.   Stevens DA, Moss RB, Hernandez C, Clemons KV, Martinez M. 2016. Effect of

Media Modified To Mimic Cystic Fibrosis Sputum on the Susceptibility of

Aspergillus fumigatus, and the Frequency of Resistance at One Center.

Antimicrob Agents Chemother 60:2180-4.

21.   Lightly TJ, Phung RR, Sorensen JL, Cardona ST. 2017. Synthetic cystic fibrosis

sputum medium diminishes *Burkholderia cenocepacia* antifungal activity

against *Aspergillus fumigatus* independently of phenylacetic acid

production. Canadian Journal of Microbiology 63:427-438.

22. Kirchhoff L, Weisner A-K, Schrepffer M, Hain A, Scharmann U, Buer J, Rath P-M,

Steinmann J. 2020. Phenotypical Characteristics of the Black Yeast Exophiala

dermatitidis Are Affected by Pseudomonas aeruginosa in an Artificial Sputum

Medium Mimicking Cystic Fibrosis–Like Conditions. Frontiers in Microbiology 11.

23. Limoli DH, Whitfield GB, Kitao T, Ivey ML, Davis MR, Grahl N, Hogan DA,

Rahme LG, Howell PL, O'Toole GA, Goldberg JB. 2017. *Pseudomonas

aeruginosa* Alginate Overproduction Promotes Coexistence with

Staphylococcus aureus in a Model of Cystic Fibrosis Respiratory Infection.

mBio 8.

24. Colmer-Hamood JA, Dzvova N, Kruczek C, Hamood AN. 2016. In Vitro Analysis

of Pseudomonas aeruginosa Virulence Using Conditions That Mimic the

Environment at Specific Infection Sites, p 151-191, Progress in Molecular Biology

and Translational Science, vol 142. Elsevier.

25. Foundation CF. 2019. 2018 Annual Data Report. Bethesda, Maryland.

26. Konstan MW, Wagener JS, VanDevanter DR, Pasta DJ, Yegin A, Rasouliyan L,

Morgan WJ. 2012. Risk factors for rate of decline in FEV1 in adults with cystic

fibrosis. Journal of Cystic Fibrosis 11:405-411.

27. Bianconi I, Jeukens J, Freschi L, Alcalá-Franco B, Facchini M, Boyle B, Molinaro

A, Kukavica-Ibrulj I, Tümmler B, Levesque RC, Bragonzi A. 2015. Comparative

genomics and biological characterization of sequential Pseudomonas aeruginosa

isolates from persistent airways infection. BMC Genomics 16.

28.   Ciofu O, Tolker-Nielsen T, Jensen PØ, Wang H, Høiby N. 2015. Antimicrobial

resistance, respiratory tract infections and role of biofilms in lung infections in

cystic fibrosis patients. Advanced Drug Delivery Reviews 85:7-23.

29.   Winstanley C, O'Brien S, Brockhurst MA. 2016. Pseudomonas aeruginosa

Evolutionary Adaptation and Diversification in Cystic Fibrosis Chronic Lung

Infections. Trends in Microbiology 24:327-337.

30.   Feliziani S, Luján AM, Moyano AJ, Sola C, Bocco JL, Montanaro P, Canigia LF,

Argaraña CE, Smania AM. 2010. Mucoidy, Quorum Sensing, Mismatch Repair

and Antibiotic Resistance in Pseudomonas aeruginosa from Cystic Fibrosis

Chronic Airways Infections. PLoS ONE 5:e12669.

31.   Mouton JW, den Hollander JG, Horrevorts AM. 1993. Emergence of antibiotic

resistance amongst *Pseudomonas aeruginosa* isolates from patients with

cystic fibrosis. Journal of Antimicrobial Chemotherapy 31:919-926.

32.   Martin LW, Reid DW, Sharples KJ, Lamont IL. 2011. *Pseudomonas* siderophores in the sputum of patients

Type="Italic">Pseudomonas</Emphasis> siderophores in the sputum of patients

with cystic fibrosis. BioMetals 24:1059-1067.

33.   Quinn RA, Phelan VV, Whiteson KL, Garg N, Bailey BA, Lim YW, Conrad DJ,

Dorrestein PC, Rohwer FL. 2016. Microbial, host and xenobiotic diversity in the

cystic fibrosis sputum metabolome. The ISME Journal 10:1483-1498.

34.   Workentine ML, Sibley CD, Glezerson B, Purighalla S, Norgaard-Gron JC,

Parkins MD, Rabin HR, Surette MG. 2013. Phenotypic Heterogeneity of

Pseudomonas aeruginosa Populations in a Cystic Fibrosis Patient. PLoS ONE

8:e60225.

35.   Diggle SP, Matthijs S, Wright VJ, Fletcher MP, Chhabra SR, Lamont IL, Kong X,

Hider RC, Cornelis P, Cámara M, Williams P. 2007. The Pseudomonas

aeruginosa 4-Quinolone Signal Molecules HHQ and PQS Play Multifunctional

Roles in Quorum Sensing and Iron Entrapment. Chemistry & Biology 14:87-96.

36.   Wang Y, Kern SE, Newman DK. 2010. Endogenous Phenazine Antibiotics

Promote Anaerobic Survival of Pseudomonas aeruginosa via Extracellular

Electron Transfer. Journal of Bacteriology 192:365-369.

37.   Caiazza NC, Shanks RMQ, O'Toole GA. 2005. Rhamnolipids Modulate

Swarming Motility Patterns of Pseudomonas aeruginosa. Journal of Bacteriology

187:7351-7361.

38.   Cornelis P. 2010. Iron uptake and metabolism in pseudomonads. Applied

Microbiology and Biotechnology 86:1637-1645.

39. Cox CD, Adams P. 1985. Siderophore activity of pyoverdin for Pseudomonas

aeruginosa. Infection and Immunity 48:130-138.

40. Briard B, Bomme P, Lechner BE, Mislin GLA, Lair V, Prévost M-C, Latgé J-P,

Haas H, Beauvais A. 2015. Pseudomonas aeruginosa manipulates redox and

iron homeostasis of its microbiota partner Aspergillus fumigatus via phenazines.

Scientific Reports 5.

41. Briard B, Rasoldier V, Bomme P, ElAouad N, Guerreiro C, Chassagne P,

Muszkieta L, Latgé J-P, Mulard L, Beauvais A. 2017. Dirhamnolipids secreted

from Pseudomonas aeruginosa modify anjpegungal susceptibility of Aspergillus

fumigatus by inhibiting β1,3 glucan synthase activity. The ISME Journal 11:1578-

1591.

42. Costello A, Reen FJ, O'Gara F, Callaghan M, McClean S. 2014. Inhibition of co-

colonizing cystic fibrosis-associated pathogens by Pseudomonas aeruginosa and

Burkholderia multivorans. Microbiology (Reading, England) 160:1474-1487.

43. Wang M, Carver JJ, Phelan VV, Sanchez LM, Garg N, Peng Y, Nguyen DD,

Watrous J, Kapono CA, Luzzatto-Knaan T, Porto C, Bouslimani A, Melnik AV,

Meehan MJ, Liu W-T, Crüsemann M, Boudreau PD, Esquenazi E, Sandoval-

Calderón M, Kersten RD, Pace LA, Quinn RA, Duncan KR, Hsu C-C, Floros DJ,

Gavilan RG, Kleigrewe K, Northen T, Dutton RJ, Parrot D, Carlson EE, Aigle B,

Michelsen CF, Jelsbak L, Sohlenkamp C, Pevzner P, Edlund A, McLean J, Piel J,

Murphy BT, Gerwick L, Liaw C-C, Yang Y-L, Humpf H-U, Maansson M, Keyzers

RA, Sims AC, Johnson AR, Sidebottom AM, Sedio BE, et al. 2016. Sharing and

community curation of mass spectrometry data with Global Natural Products

Social Molecular Networking. Nature Biotechnology 34:828-837.

44.   Nothias LF, Petras D, Schmid R, Duhrkop K, Rainer J, Sarvepalli A, Protsyuk I,

Ernst M, Tsugawa H, Fleischauer M, Aicheler F, Aksenov AA, Alka O, Allard PM,

Barsch A, Cachet X, Caraballo-Rodriguez AM, Da Silva RR, Dang T, Garg N,

Gauglitz JM, Gurevich A, Isaac G, Jarmusch AK, Kamenik Z, Kang KB, Kessler

N, Koester I, Korf A, Le Gouellec A, Ludwig M, Martin HC, McCall LI, McSayles

J, Meyer SW, Mohimani H, Morsy M, Moyne O, Neumann S, Neuweger H,

Nguyen NH, Nothias-Esposito M, Paolini J, Phelan VV, Pluskal T, Quinn RA,

Rogers S, Shrestha B, Tripathi A, van der Hooft JJJ, et al. 2020. Feature-based

molecular networking in the GNPS analysis environment. Nat Methods

doi:10.1038/s41592-020-0933-6.

45.   La Rosa R, Johansen HK, Molin S. 2019. Adapting to the Airways: Metabolic

Requirements of Pseudomonas aeruginosa during the Infection of Cystic Fibrosis

Patients. Metabolites 9.

46.   Phelan VV, Liu WT, Pogliano K, Dorrestein PC. 2011. Microbial metabolic

exchange--the chemotype-to-phenotype link. Nat Chem Biol 8:26-35.

47.   Schmidt R, Ulanova D, Wick LY, Bode HB, Garbeva P. 2019. Microbe-driven

chemical ecology: past, present and future. ISME J 13:2656-2663.

48.   Phelan VV, Moree WJ, Aguilar J, Cornett DS, Koumoutsi A, Noble SM, Pogliano

K, Guerrero CA, Dorrestein PC. 2014. Impact of a Transposon Insertion in phzF2

on the Specialized Metabolite Production and Interkingdom Interactions of

Pseudomonas aeruginosa. Journal of Bacteriology 196:1683-1693.

49.   Phelan VV, Fang J, Dorrestein PC. 2015. Mass Spectrometry Analysis of

Pseudomonas aeruginosa Treated with Azithromycin. J Am Soc Mass Spectrom

26:873-7.

50.   Condren AR, Kahl LJ, Boelter G, Kritikos G, Banzhaf M, Dietrich LEP, Sanchez

LM. 2020. Biofilm Inhibitor Taurolithocholic Acid Alters Colony Morphology,

Specialized Metabolism, and Virulence of Pseudomonas aeruginosa. ACS Infect

Dis 6:603-612.

51.   Depke T, Thöming JG, Kordes A, Häussler S, Brönstrup M. 2020. Untargeted

LC-MS Metabolomics Differentiates Between Virulent and Avirulent Clinical

Strains of Pseudomonas aeruginosa. Biomolecules 10:1041.

52.   Purves K, Macintyre L, Brennan D, Hreggviðsson G, Kuttner E, Ásgeirsdóttir M,

Young L, Green D, Edrada-Ebel R, Duncan K. 2016. Using Molecular Networking

for Microbial Secondary Metabolite Bioprospecting. Metabolites 6:2.

53.   Caraballo-Rodríguez AM, Dorrestein PC, Pupo MT. 2017. Molecular inter-

kingdom interactions of endophytes isolated from Lychnophora ericoides.

Scientific Reports 7.

54.   Das T, Manefield M. 2012. Pyocyanin Promotes Extracellular DNA Release in

Pseudomonas aeruginosa. PLoS ONE 7:e46718.

55.   Das T, Manefield M. 2013. Phenazine production enhances extracellular DNA

release via hydrogen peroxide generation in <i>Pseudomonas aeruginosa</i>.

Communicative & Integrative Biology 6:e23570.

56.   Moree WJ, Phelan VV, Wu CH, Bandeira N, Cornett DS, Duggan BM, Dorrestein

PC. 2012. Interkingdom metabolic transformations captured by microbial imaging

mass spectrometry. Proceedings of the National Academy of Sciences of the

United States of America 109:13811-13816.

57.   Noto MJ, Burns WJ, Beavers WN, Skaar EP. 2017. Mechanisms of Pyocyanin

Toxicity and Genetic Determinants of Resistance in Staphylococcus aureus.

Journal of Bacteriology 199.

58. Das T, Kutty SK, Tavallaie R, Ibugo AI, Panchompoo J, Sehar S, Aldous L,

Yeung AWS, Thomas SR, Kumar N, Gooding JJ, Manefield M. 2015. Phenazine

virulence factor binding to extracellular DNA is important for Pseudomonas

aeruginosa biofilm formation. Scientific Reports 5.

59. Wilson R, Pitt T, Taylor G, Watson D, MacDermot J, Sykes D, Roberts D, Cole P.

1987. Pyocyanin and 1-hydroxyphenazine produced by Pseudomonas

aeruginosa inhibit the beating of human respiratory cilia in vitro. Journal of

Clinical Investigation 79:221-229.

60. Allen L, Dockrell DH, Pattery T, Lee DG, Cornelis P, Hellewell PG, Whyte MKB.

2005. Pyocyanin Production by <i>Pseudomonas aeruginosa</i> Induces

Neutrophil Apoptosis and Impairs Neutrophil-Mediated Host Defenses In Vivo.

The Journal of Immunology 174:3643-3649.

61. Wang Y, Wilks JC, Danhorn T, Ramos I, Croal L, Newman DK. 2011. Phenazine-

1-Carboxylic Acid Promotes Bacterial Biofilm Development via Ferrous Iron

Acquisition. Journal of Bacteriology 193:3606-3617.

62. Schalk IJ, Guillon L. 2013. Pyoverdine biosynthesis and secretion in

<i>Pseudomonas aeruginosa</i> : implications for metal homeostasis:

Pyoverdine biosynthesis. Environmental Microbiology 15:1661-1673.

63. Gasser V, Guillon L, Cunrath O, Schalk IJ. 2015. Cellular organization of

siderophore biosynthesis in Pseudomonas aeruginosa: Evidence for

siderosomes. Journal of Inorganic Biochemistry 148:27-34.

64. Cox CD, Rinehart KL, Moore ML, Cook JC. 1981. Pyochelin: novel structure of

an iron-chelating growth promoter for Pseudomonas aeruginosa. Proceedings of

the National Academy of Sciences 78:4256-4260.

65. Korgaonkar AK, Whiteley M. 2011. Pseudomonas aeruginosa enhances

production of an antimicrobial in response to N-acetylglucosamine and

peptidoglycan. J Bacteriol 193:909-17.

66. Van Ark G, Berden JA. 1977. Binding of HQNO to beef-heart sub-mitochondrial

particles. Biochimica et Biophysica Acta (BBA) - Bioenergetics 459:119-137.

67. Hazan R, Que Yok A, Maura D, Strobel B, Majcherczyk Paul A, Hopper Laura R,

Wilbur David J, Hreha Teri N, Barquera B, Rahme Laurence G. 2016. Auto

Poisoning of the Respiratory Chain by a Quorum-Sensing-Regulated Molecule

Favors Biofilm Formation and Antibiotic Tolerance. Current Biology 26:195-206.

68. Rampioni G, Falcone M, Heeb S, Frangipani E, Fletcher MP, Dubern J-F, Visca

P, Leoni L, Cámara M, Williams P. 2016. Unravelling the Genome-Wide

Contributions of Specific 2-Alkyl-4-Quinolones and PqsE to Quorum Sensing in

Pseudomonas aeruginosa. PLOS Pathogens 12:e1006029.

69. McKnight SL, Iglewski BH, Pesci EC. 2000. The Pseudomonas Quinolone Signal

Regulates rhl Quorum Sensing in Pseudomonas aeruginosa. Journal of

Bacteriology 182:2702-2708.

70. Guo Q, Kong W, Jin S, Chen L, Xu Y, Duan K. 2014. PqsR-dependent and

PqsR-independent regulation of motility and biofilm formation by PQS in

<i>Pseudomonas aeruginosa</i> PAO1: Regulation of motility and biofilm by

PQS. Journal of Basic Microbiology 54:633-643.

71. Lépine F, Milot S, Déziel E, He J, Rahme LG. 2004. Electrospray/mass

spectrometric identification and analysis of 4-hydroxy-2-alkylquinolines (HAQs)

produced by Pseudomonas aeruginosa. Journal of the American Society for

Mass Spectrometry 15:862-869.

72. Davey ME, Caiazza NC, O'Toole GA. 2003. Rhamnolipid Surfactant Production

Affects Biofilm Architecture in Pseudomonas aeruginosa PAO1. Journal of

Bacteriology 185:1027-1036.

73. Glick R, Gilmour C, Tremblay J, Satanower S, Avidan O, Déziel E, Greenberg

EP, Poole K, Banin E. 2010. Increase in Rhamnolipid Synthesis under Iron-

Limiting Conditions Influences Surface Motility and Biofilm Formation in

Pseudomonas aeruginosa. Journal of Bacteriology 192:2973-2980.

74.   McClure CD, Schiller NL. 1996. Inhibition of Macrophage Phagocytosis by

806   Pseudomonas aeruginosa Rhamnolipids In Vitro and In Vivo. Current

807   Microbiology 33:109-117.

808   75.   Zulianello L, Canard C, Köhler T, Caille D, Lacroix J-S, Meda P. 2006.

809   Rhamnolipids Are Virulence Factors That Promote Early Infiltration of Primary

810   Human Airway Epithelia by Pseudomonas aeruginosa. Infection and Immunity

811   74:3134-3147.

812   76.   Ochsner UA, Reiser J. 1995. Autoinducer-mediated regulation of rhamnolipid

813   biosurfactant synthesis in Pseudomonas aeruginosa. Proceedings of the National

814   Academy of Sciences 92:6424-6428.

815   77.   Schuster M, Greenberg EP. 2007. Early activation of quorum sensing in

816   Pseudomonas aeruginosa reveals the architecture of a complex regulon. BMC

817   Genomics 8:287.

818   78.   Whiteley M, Lee KM, Greenberg EP. 1999. Identification of genes controlled by

819   quorum sensing in Pseudomonas aeruginosa. Proceedings of the National

820   Academy of Sciences 96:13904-13909.

821   79.   Meyer JM, Neely A, Stintzi A, Georges C, Holder IA. 1996. Pyoverdin is essential

822   for virulence of Pseudomonas aeruginosa. Infection and Immunity 64:518-523.

823   80.   Takase H, Nitanai H, Hoshino K, Otani T. 2000. Impact of Siderophore

824   Production onPseudomonas aeruginosa Infections in Immunosuppressed Mice.

825   Infection and Immunity 68:1834-1839.

826   81.   Sriyosachati S, Cox CD. 1986. Siderophore-mediated iron acquisition from

827   transferrin by Pseudomonas aeruginosa. Infection and Immunity 52:885-891.

828   82.   Ochsner UA, Vasil AI, Vasil ML. 1995. Role of the ferric uptake regulator of

829   Pseudomonas aeruginosa in the regulation of siderophores and exotoxin A

830   expression: purification and activity on iron-regulated promoters. Journal of

831   bacteriology 177:7194-7201.

832   83.   Co JY, Cárcamo-Oyarce G, Billings N, Wheeler KM, Grindy SC, Holten-Andersen

833   N, Ribbeck K. 2018. Mucins trigger dispersal of Pseudomonas aeruginosa

834   biofilms. npj Biofilms and Microbiomes 4.

835   84.   Wheeler KM, Cárcamo-Oyarce G, Turner BS, Dellos-Nolan S, Co JY, Lehoux S,

836   Cummings RD, Wozniak DJ, Ribbeck K. 2019. Mucin glycans attenuate the

837   virulence of Pseudomonas aeruginosa in infection. Nature Microbiology 4:2146-

838   2154.

839   85.   Paone P, Cani PD. 2020. Mucus barrier, mucins and gut microbiota: the

840   expected slimy partners? Gut 69:2232-2243.

86. Wagner CE, Wheeler KM, Ribbeck K. 2018. Mucins and Their Role in Shaping

the Functions of Mucus Barriers. Annual Review of Cell and Developmental

Biology 34:189-215.

87. Hoffman CL, Lalsiamthara J, Aballay A. 2020. Host Mucin Is Exploited by

Pseudomonas aeruginosa To Provide Monosaccharides Required for a

Successful Infection. mBio 11.

88. Flynn JM, Niccum D, Dunitz JM, Hunter RC. 2016. Evidence and Role for

Bacterial Mucin Degradation in Cystic Fibrosis Airway Disease. PLOS Pathogens

12:e1005846.

89. Darch SE, Kragh KN, Abbott EA, Bjarnsholt T, Bull JJ, Whiteley M. 2017. Phage

Inhibit Pathogen Dissemination by Targeting Bacterial Migrants in a Chronic

Infection Model. mBio 8.

90. Secor PR, Michaels LA, Ratjen A, Jennings LK, Singh PK. 2018. Entropically

driven aggregation of bacteria by host polymers promotes antibiotic tolerance in

Pseudomonas aeruginosa. Proceedings of the National Academy of

Sciences 115:10780-10785.

91. Hao Y, Kuang Z, Walling BE, Bhatia S, Sivaguru M, Chen Y, Gaskins HR, Lau

GW. 2012. Pseudomonas aeruginosa pyocyanin causes airway goblet cell

hyperplasia and metaplasia and mucus hypersecretion by inactivating the

transcriptional factor FoxA2: Pseudomonas pyocyanin inactivates airway FoxA2. Cellular Microbiology 14:401-415.

92. Kanthakumar K, Taylor G, Tsang KW, Cundell DR, Rutman A, Smith S, Jeffery PK, Cole PJ, Wilson R. 1993. Mechanisms of action of Pseudomonas aeruginosa pyocyanin on human ciliary beat in vitro. Infection and Immunity 61:2848-2853.

93. O'Malley YQ, Reszka KJ, Spitz DR, Denning GM, Britigan BE. 2004. *Pseudomonas aeruginosa* pyocyanin directly oxidizes glutathione and decreases its levels in airway epithelial cells. American Journal of Physiology-Lung Cellular and Molecular Physiology 287:L94-L103.

94. Schwarzer C, Fischer H, Kim E-J, Barber KJ, Mills AD, Kurth MJ, Gruenert DC, Suh JH, Machen TE, Illek B. 2008. Oxidative stress caused by pyocyanin impairs CFTR Cl− transport in human bronchial epithelial cells. Free Radical Biology and Medicine 45:1653-1662.

95. Moree WJ, Phelan VV, Wu C-H, Bandeira N, Cornett DS, Duggan BM, Dorrestein PC. 2012. Interkingdom metabolic transformations captured by microbial imaging mass spectrometry. Proceedings of the National Academy of Sciences 109:13811-13816.

96.   Orazi G, Jean-Pierre F, O'Toole GA. 2020. *Pseudomonas aeruginosa*

PA14 Enhances the Efficacy of Norfloxacin against *Staphylococcus*

aureus Newman Biofilms. Journal of Bacteriology 202.

97.   Minandri F, Imperi F, Frangipani E, Bonchi C, Visaggio D, Facchini M, Pasquali

P, Bragonzi A, Visca P. 2016. Role of Iron Uptake Systems in Pseudomonas

aeruginosa Virulence and Airway Infection. Infection and Immunity 84:2324-

2335.

98.   Kilbourn JP. 1984. Composition of sputum from patients with cystic fibrosis.

Current Microbiology 11:19-22.

99.   Hunter RC, Asfour F, Dingemans J, Osuna BL, Samad T, Malfroot A, Cornelis P,

Newman DK. 2013. Ferrous Iron Is a Significant Component of Bioavailable Iron

in Cystic Fibrosis Airways. mBio 4.

100.   Reid DW, Carroll V, O'May C, Champion A, Kirov SM. 2007. Increased airway

iron as a potential factor in the persistence of Pseudomonas aeruginosa infection

in cystic fibrosis. European Respiratory Journal 30:286-292.

101.   Ghio AJ, Roggli VL, Soukup JM, Richards JH, Randell SH, Muhlebach MS. 2013.

Iron accumulates in the lavage and explanted lungs of cystic fibrosis patients.

Journal of Cystic Fibrosis 12:390-398.

102.   Jacobs MA, Alwood A, Thaipisuttikul I, Spencer D, Haugen E, Ernst S, Will O,

896   Kaul R, Raymond C, Levy R, Chun-Rong L, Guenthner D, Bovee D, Olson MV,

897   Manoil C. 2003. Comprehensive transposon mutant library of Pseudomonas

898   aeruginosa. Proc Natl Acad Sci U S A 100:14339-44.

899   103.   Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N,

900   Schwikowski B, Ideker T. 2003. Cytoscape: A software environment for

901   integrated models of biomolecular interaction networks. Genome Research

902   13:2498-2504.

903   104.   Conway JR, Lex A, Gehlenborg N. 2017. UpSetR: an R package for the

904   visualization of intersecting sets and their properties. Bioinformatics 33:2938-

905   2940.

906   105.   Lex A, Gehlenborg N, Strobelt H, Vuillemot R, Pfister H. 2014. UpSet:

907   Visualization of Intersecting Sets. IEEE Trans Vis Comput Graph 20:1983-92.

908   106.   Pluskal T, Castillo S, Villar-Briones A, Oresic M. 2010. MZmine 2: Modular

909   framework for processing, visualizing, and analyzing mass spectrometry-based

910   molecular profile data. Bmc Bioinformatics 11.

911   107.   Sumner LW, Amberg A, Barrett D, Beale MH, Beger R, Daykin CA, Fan TWM,

912   Fiehn O, Goodacre R, Griffin JL, Hankemeier T, Hardy N, Harnly J, Higashi R,

913   Kopka J, Lane AN, Lindon JC, Marriott P, Nicholls AW, Reily MD, Thaden JJ,

Viant MR. 2007. Proposed minimum reporting standards for chemical analysis.

Metabolomics 3:211-221.

108. Chong J, Soufan O, Li C, Caraus I, Li S, Bourque G, Wishart DS, Xia J. 2018.

MetaboAnalyst 4.0: towards more transparent and integrative metabolomics

analysis. Nucleic Acids Res 46:W486-W494.

109. Chong J, Xia J. 2018. MetaboAnalystR: an R package for flexible and

reproducible analysis of metabolomics data. Bioinformatics 34:4313-4314.

110. Chong J, Yamamoto M, Xia J. 2019. MetaboAnalystR 2.0: From Raw Spectra to

Biological Insights. Metabolites 9.

111. Pang Z, Chong J, Li S, Xia J. 2020. MetaboAnalystR 3.0: Toward an Optimized

Workflow for Global Metabolomics. Metabolites 10:186.

FIGURES

Figure 1 Timeline of ASM formulations. ASM formulations can be categorized into

complex (open circles) or defined (closed circles) media. Each formulation is plotted at

the year it was first published. Nutrient alterations between formulations are indicated by

arrows. Compared to preceding formulation: + = added, - = removed, ↑ = increased, ↓ =

decreased, → = different source. AA = amino acids, BSA = bovine serum albumin, Cas

AA = casamino acids, MEM AA = MEM amino acids, DOPC = 1,2-dioleoyl-sn-glycero-3-

phosphocholine, GlcNAc = *N*-acetylglucosamine, PABA = 4-aminobenzoic acid, NAD^+ =

nicotinamide adenine dinucleotide, ADE = adenine, GUA = guanine, XAN = xanthine,

HX = hypoxanthine. Recipes for ASM formulations are in detailed in *Dataset S1 ASM*

formulations.

 Figure 2 PAO1 phenotype and metabolome in ASM and LB. (A) Representative

phenotypic images of *P. aeruginosa* PAO1 grown in ASM. LB was used as a positive

control. **(B)** Classical molecular network of the exometabolome of *P. aeruginosa* PAO1

grown in ASM. LB was used as a positive control. Nodes representing MS/MS detected

in all media are blue. Nodes representing MS/MS detected in a single medium are red.

The primary molecular families of *P. aeruginosa* are highlighted. Nodes not connected

to any other node (singletons) were removed for clarity. The full network can be

accessed via the link in 'Materials and Methods'. **(C & D)** Upset plots of number of

unique *m/z* values detected in the 10 media arranged by decreasing frequency (C) and

increasing degree (D). Intersections shown were limited to improve visualization. Upset

plot data analysis is detailed in *Supplemental Dataset – Upset Plot Generation and*

Results.

Figure 3 Phenazines Production in ASM. (A) Phenazine molecular family visualized

with feature based molecular networking. Each node represents a feature, and relative

abundance of each feature among ASM depicted by a pie chart. Pie charts are colored

counterclockwise starting at three o'clock: Soothill red; Romling orange; Winstanley

yellow; ASMDM green; Coldwell aqua; SDSU light blue; SCFM1 blue; SCFM2 purple;

SCFM3 pink. GNPS library matches are indicated by black outlines. (B) Total

normalized AUC for the phenazine molecular family in each ASM formulation depicted

by box and whisker plot with individual samples shown (n=12 per ASM formulation). (C)

Normalized AUC of each phenazine, stacked to show the phenazine profile of PAO1 in

each ASM formulation. (D) Phenazine production in each ASM formulation as a percent

of total phenazine production in the medium. (C, D) Phenazines from bottom to top of

bars: PCA, phenazine-1-carboxylic acid, purple; PCN, phenazine-1-carboxamide,

green; PYO, pyocyanin, red; 1-HP, 1-hydroxyphenazine, blue.

Figure 4 Quinolone Production in ASM. (A) Quinolone molecular family visualized

with feature based molecular networking. Each node represents a feature, and relative

abundance of each feature among ASM is depicted with a pie chart. Pie charts are

colored counterclockwise starting at three o'clock: Soothill red; Romling orange;

Winstanley yellow; ASMDM green; Cordwell aqua; SDSU light blue; SCFM1 blue;

SCFM2 purple; SCFM3 pink. GNPS library matches are indicated by black outlines. (B)

Total normalized AUC for the quinolone molecular family in each ASM formulation

depicted by box and whisker plot with individual samples shown (n=12 per ASM

formulation). (C) Quinolone subfamily production in each ASM formulation as a percent

of total quinolone production in the medium. Quinolone subfamilies from top to bottom

of bars: HHQ-type, blue; HQNO-type, red; PQS-type, green; Mixed, purple; Other,

orange.

Figure 5 Rhamnolipid Production in ASM. (A) Rhamnolipid molecular family

visualized with feature based molecular networking. Each node represents a feature,

and relative abundance of each feature among ASM is depicted with a pie chart. Pie

charts are colored counterclockwise starting at three o'clock: Soothill red; Romling

orange; Winstanley yellow; ASMDM green; Cordwell aqua; SDSU light blue; SCFM1

blue; SCFM2 purple; SCFM3 pink. GNPS library matches are indicated by black

outlines. (B) Total normalized AUC for the rhamnolipid molecular family in each ASM

formulation depicted by box and whisker plot with individual samples shown (n=12 per

ASM formulation). (C) Rhamnolipid production in each ASM formulation as a percent of

total rhamnolipid production in the medium. Rhamnolipids from top to bottom of bars:

Rha-C10-C10, blue; Rha-C10-C12:1, green; Rha-C10-C12, red; Rha-Rha-C10-C10,

purple; Rha-Rha-C10-C12:1, black; Rha-Rha-C10-C12, orange.

Figure 6 Siderophore Production in ASM. (A) PCH and PVD siderophore molecular

families visualized with feature based molecular networking. Each node represents a

feature, and relative abundance of each feature among ASM is depicted with a pie

chart. Pie charts are colored counterclockwise starting at three o'clock: Soothill red;

Romling orange; Winstanley yellow; ASMDM green; Cordwell aqua; SDSU light blue;

SCFM1 blue; SCFM2 purple; SCFM3 pink. (B) Total normalized AUC for the PCH

molecular family in each ASM formulation depicted by box and whisker plot with

individual samples shown (n=12 per ASM formulation). (C) Total normalized AUC for

the PVD molecular family in each ASM formulation depicted by box and whisker plot

with individual samples shown (n=12 per ASM formulation).

1007

1008

1009

Figure 7 PGM iron content and impact on PAO1 siderophores. (A, B) Production of

P. aeruginosa siderophores (A) pyochelin (PCH) and (B) pyoverdine (PVD) after growth

in SCFM1 (black circles) and SCFM1 with mucin (+PGM, porcine gastric mucin; gray

squares). The area under the curve (AUC) of four biological replicates are designated

by box and whisker plots. Sample groups were compared using a two tailed, unpaired t

test with Welch's correction, (A) *, p=0.0114; (B) *, p=0.0269. **(C)** Iron (^{56}Fe) was

quantified from 1X MOPS and PGM by ICP-MS and the amount of iron added to

SCFM1 by each component was determined. The iron concentration (µM) of six

technical replicates are designated by box and whisker plots. The iron concentration

added to SCFM1 (3.6 µM) is designated by the dotted line.

Supplemental Material

Figure S1 Growth of PAO1 in ASM. PAO1 was grown in nine ASM formulations and

LB. Samples were prepared in three batches with LB growth included in each batch as a

positive control. (A) Enumeration PAO1 of CFUs after growth in ASM formulations and

LB. The means, standard error of the mean, and values of three biological replicates are

shown. (B) Measurement of pH of PAO1 samples after growth in ASM formulations and

LB. The mean and standard error of the mean and values of three biological replicates

(open circles) are shown. The pH of each medium at inoculation is represented by a filled

triangle.

Figure S2 Growth of PAO1 in SCFM1 with and without PGM. (A) Enumeration PAO1

of CFUs after growth in each SCFM1 and SCFM1 with PGM. The means, standard error

of the mean, and values of four biological replicates are shown, $p<0.0001$ as determined

by a two-tailed Student's *t* test. **(B)** Measurement of pH of PAO1 samples after growth in

each SCFM1 and SCFM1 with PGM. The mean, standard error of the mean, and values

of four biological replicates are shown, the dotted line represents the pH of the medium

(pH = 7).

Figure S3 Pyovderdine chromatograms. Chromatogram traces of extracted ion

chromatograms for the *m/z* values 667.3094 (PVD E [M+2H], blue); 693.7660 (PVD E

[M+Fe+H], orange); 676.3333 (Ferribactin [M+2H], green); 505.2437 (red), and 661.3262

(purple) are shown with a tolerance of 20 ppm. Traces are all from a single representative

SCFM1 sample from the dataset of PAO1 grown in ASM formulations visualized using

the GNPS LCMS Dashboard (v0.25).

Table S1 ASM metabolite annotation and mass defect. Provides a table containing

metabolite annotation, mass defect, and level of annotation from PAO1 cultures in ASM.

Table S2 SCFM1 and mucin siderophore annotation and mass defect. Provides a

table containing metabolite annotation, mass defect, and level of annotation from PAO1

cultures in SCFM1 with and without mucin.

1047

1048