TITLE: Commercial porcine gastric mucin contributes to variation in production of small

molecule virulence factors by *Pseudomonas aeruginosa* when cultured in different

formulations of artificial sputum medium

AUTHORS: Rachel L. Neve[1], Brent D. Carrillo[2], and Vanessa V. Phelan[2#]

AUTHOR AFFILIATIONS:

[1] Department of Immunology and Microbiology, School of Medicine, University of

Colorado - Anschutz Medical Campus, Aurora, CO, 80045, USA

[2] Department of Pharmaceutical Sciences, Skaggs School of Pharmacy and

Pharmaceutical Sciences, University of Colorado - Anschutz Medical Campus, Aurora,

CO, 80045, USA

CORRESPONDING AUTHOR:

[#] Address correspondence to Vanessa V. Phelan, vanessa.phelan@cuanschutz.edu

ABSTRACT

Site-specific *in vitro* culture media are being developed to investigate microbial

pathogenicity and ecology in nutrient environments that are more reflective of disease.

For microbial ecology research in cystic fibrosis (CF), different artificial sputum media

(ASM) formulations have been created to recapitulate the nutrient availability of the CF

lung environment. However, these ASM formulations vary in concentration of amino

acids, mucin, and other niche-specific compounds. Here, we measured the differential

production of small molecule virulence factors by *Pseudomonas aeruginosa*, the

predominant pathogen in CF pulmonary infections, cultured in nine different ASM

formulations via liquid chromatography tandem mass spectrometry (LC-MS/MS) and

molecular networking. We show that different ASM formulations lead to different

phenotypes and metabolic profiles of *P. aeruginosa* and commercial porcine gastric

mucin (PGM) contains a myriad of contaminants, including iron, which affect *P.*

aeruginosa physiology.

IMPORTANCE

Different media formulations aiming to replicate *in vivo* infection environments contain

different nutrients, which affects interpretation of experimental results. Inclusion of

undefined components, such as commercial porcine gastric mucin (PGM), in an

otherwise chemically defined medium can alter the nutrient content of the medium in

unexpected ways and influence experimental outcomes.

In vitro cultivation systems are being developed and applied to investigate interactions

between members of host-associated microbiota using native-like nutritional

environments. These systems are beneficial to researchers because they enable higher

resolution analysis of microbiome community structure and function by removing

confounders associated with host cells, immune response, and variability in nutritional

intake.[1,2] Foundational to the utility of *in vitro* culture systems to investigate the

microbiota is the replication of host sites using physiologically relevant media. However,

across laboratories, different variations of site-specific culture media are being used in

experimental investigations. A representative example of multiple media formulations

created and used to interrogate site-specific microbial interactions is artificial sputum

media (ASM), a class of media aiming to replicate the nutritional environment

encountered by the pulmonary microbiome of persons with cystic fibrosis (CF).[3-14] ASM

have been applied to investigate microbial physiology[3,5,15-17], biofilm morphology[10,18],

antibiotic susceptibility[19,20], and interspecies interactions in the context of CF[21-23].

CF is a hereditary disease characterized by the accumulation of thick and sticky

bronchial mucus in the lungs, which enables growth of opportunistic pathogens.[24]

Pulmonary mucus is a nutritionally distinct environmental niche comprised of mucin,

DNA, amino acids, lipids, and micronutrients, including metals[25,26]. Reflective of the

variability of measurements of nutrient concentration in CF sputum and bronchoalveolar

lavage fluid (BALF)[27,28], each ASM formulation differs in the concentration mucin, DNA,

amino acids, lipids, and metals, affecting overall nutrient composition of the media.[25,26]

Variations in ASM formulation are driven by literature precedent, direct measurement of

nutrients from clinical samples, personal observations, and economic cost of materials.[3-

[14] Importantly, disparate nutrient concentration and bioavailability between ASM

formulations may affect microbial growth, metabolism, and virulence[3,7,20,29]. Even

though the formulations of the individual ASM are distinct, experimental results from

different formulations claim to provide physiologically relevant depictions of pathogen

growth and physiology similar to that in CF sputum.[3,7,30]

Herein, we perform comparative metabolomics of the *P. aeruginosa* specialized

metabolome grown in nine different ASM formulations. *P. aeruginosa* is the primary

bacterial opportunistic pathogen infecting CF lungs and contributes to lung function

decline.[31,32] In the CF lung, *P. aeruginosa* grows as biofilms, exopolysaccharide

encased communities, which are recalcitrant to innate immune response and

antibiotics.[33] Within these biofilms, *P. aeruginosa* produces a suite of small molecule

virulence factors that enable it to establish chronic infections by damaging host cells,

outcompeting other members of the pulmonary microbiome, and developing as

biofilms.[30,34-36] As *P. aeruginosa* specialized metabolites are associated with virulence

phenotypes, their measurement can be used to understand how different nutrient

availability in ASM formulations affect *P. aeruginosa* virulence.[37,38] We show that

different ASM formulations induce differential production of *P. aeruginosa* PAO1 small

molecule virulence factors, including members of the phenazine, quinolone,

rhamnolipid, pyochelin, and pyoverdine molecular families, which is partially driven by

the inclusion of commercial porcine gastric mucin (PGM) in the media.

Results and Discussion

The nutrient availability of different ASM formulations influences *Pseudomonas aeruginosa* PAO1 phenotype and secondary metabolic profile. To test the hypothesis that different ASM formulations would result in differential production of *P. aeruginosa* specialized metabolites, strain PAO1 was statically cultured in nine different ASM formulations.[3-8,10,11] We chose the nine ASM formulations based upon citation count, prior use in our research, and if the formulation was a predecessor to other formulations. Broadly, these ASM can be separated into two lineages, complex and chemically defined. The complex lineage originates with Soothill ASM[4], which was modified to yield Romling[9,10], Winstanley[6], ASMDM[3], Cordwell[5], and SDSU[8] ASM formulations (Figure 1). Soothill ASM was created using literature values for key nutrients identified in CF sputum, taking into account economic cost of materials.[4] This base medium was modified by other researchers by addition of amino acids[6,9,10], bovine serum albumin (BSA)[6], alterations of mucin[3,8] and DNA[3] concentration, and replacement of amino acid[5,8] or iron source.[5] Concurrent to the development of the complex ASM lineage, the chemically defined synthetic CF medium (SCFM) series of ASM was created from the average concentrations of ions, free amino acids, glucose, and lactate measured from CF sputum samples (Figure 1).[7] To more closely resemble the *in vivo* CF lung environment, the SCFM1 base medium was complemented with DNA, mucin, 1,2-Dioleoyl-sn-glycero-3-phosphocholine (DOPC) as a lipid source, and *N*-acetyl glucosamine (GlcNAc) as well as various nucleosides to create SCFM2 and SCFM3, respectively.[11]

The gross morphology of the PAO1 biofilms and color of the cultures varied

between media (Figure 2A), despite similar levels of growth (Figure 2B). The Soothill

ASM PAO1 cultures were turbid, while the other eight ASM formulations induced the

development of biofilm macrostructures. Addition of amino acids to the Soothill ASM to

create Romling and Winstanley ASM led to the formation of a brown macrostructure

phenotype by PAO1. Nutritional alterations of Romling ASM to create ASMDM,

Cordwell, and SDSU ASM formulations led to assorted phenotypes of PAO1 growth,

with structurally varied biofilms with diverse coloration. In ASMDM, *P. aeruginosa* grew

heterogeneously with a layer of growth at the air-liquid interface, giving way to turbid

growth further down the culture. In SDSU ASM, PAO1 formed a green, floating structure

near the center of the well that reached the surface of the medium. PAO1 cultures in

SCFM2 and SCFM3 were highly similar, with larger biofilm structures than in SCFM1

and a distinctly blue coloration, likely from high levels of pyocyanin.[39]

As the color of *P. aeruginosa* cultures is largely dependent upon the abundance

of its small molecular virulence factors, we applied comparative metabolomics to PAO1

cultures in the nine ASM formulations. Principal component analysis was used to

visualize the similarity within and between the metabolome of PAO1 cultures in different

ASM formulations (Figure 2C). Broadly, the metabolomes of biological replicates

clustered together, with SDSU ASM cultures showing the highest between-replicate

variability, likely due to the high concentration of mucin in the medium, which may affect

metabolome extraction efficiency due to the high viscosity of the samples.[8] In the PCA,

samples clustered into three groups, largely based upon their formulations. The

nutritionally distinct SCFM ASM formulations clustered separately from the Soothill ASM

derived media. Within the Soothill ASM lineage samples, two distinct clusters were

formed; one cluster encompassing Romling, Winstanley, and ASMDM ASM and the

other containing Cordwell, SDSU, and Soothill ASM. The latter three ASM formulations

have different amino acid content than the former (Figure 1), which likely affects the

production levels of *P. aeruginosa* small molecule virulence factors, as amino acids are

biosynthetic precursors of these metabolites[40-42]. Out of 490 molecular features

detected, all but two were statistically significantly different between the samples (Figure

2D). The number of statistically different features produced in each pairwise interaction

(Figure 2E) corresponds well to the PCA results, with cultures from nutritionally related

media yielding the most similar metabolomes.

ASM formulation variation stimulates differential production of small molecule

virulence factors by PAO1. Since statistical analysis of the metabolomics data was

unable to identify specific metabolic features that differentiated *P. aeruginosa*

metabolism in the different ASM formulations, we used feature based molecular

networking (FBMN) to annotate, quantify, and visualize the production of small molecule

virulence factors produced by PAO1 in all nine ASM formulations.[43,44] *P. aeruginosa* has

a well-characterized specialized metabolome, including phenazines, quinolones,

rhamnolipids, and the siderophores pyoverdine and pyochelin.[37,45,46] *P. aeruginosa*

specialized metabolites have been detected in CF samples[30,34,35,47-53] and are

associated with pathogenicity and virulence.[29,37,38] As nutrient concentration varies

between ASM formulations, alteration in production levels of these metabolites by *P.*

aeruginosa may influence interpretation of virulence-related phenotypes.

Phenazines are redox-active metabolites with various biological activities,

including roles in promoting iron uptake[54,55], anaerobic respiration[54], biofilm

formation[56,57], and interspecies interactions.[45,55,58,59] We quantified the production of four

phenazines: 1-hydroxphenazine (1-HP), pyocyanin (PYO), phenazine-1-carboxamide

(PCN), and phenazine-1-carboxylic acid (PCA) (Figure 3). Two metabolic features have

matches to the GNPS spectral libraries for PYO (Figure 3A). These features have

different chromatographic retention times, but identical MS/MS fragmentation patterns,

suggesting they are structurally related. While 1-HP, PCA, and PYO have been

detected in CF sputum[30,52], their detection across samples has been spurious and UV

detection may not be sufficient to differentiate phenazine production from other

components of sputum.[60] Therefore, phenazine production is most commonly measured

from pure cultures of CF clinical isolates.[61] Phenazines were quantified from all PAO1

cultures. However, the abundance of specific phenazines (Figure 3A), total phenazine

production (Figure 3B), and the composition of the phenazine molecular family varied

between ASM formulations (Figure 3C). Phenazine production was lowest in Soothill

ASM and highest in SCFM2 and SCFM3 (Figure 3B). Soothill ASM does not contain

amino acids,[4] restricting the amino acid precursor availability for the biosynthesis of

siderophores by *P. aeruginosa*.[62-64] In this medium, PAO1 may produce a higher

proportion of PCA relative to other phenazines (Figure 3D) in order to acquire iron from

the media as PCA has been shown to reduce Fe^{3+} to Fe^{2+} in a *P. aeruginosa* mutant

unable to produce pyochelin and pyoverdine.[65] PYO is produced at the highest levels by

PAO1 grown in SCFM2 and SCFM3 (Figure 3A/C), which contributes to the blue

phenotype of these cultures (Figure 2A).[39] SCFM2 and SCFM3 are compositionally

different from SCFM1 by the addition of several components, including *N*-acetyl

glucosamine (GlcNAc).[11] Addition of GlcNAc to SCFM1 was shown to increase

production of PYO by *P. aeruginosa*, likely through a mechanism to sense and respond

to the presence of peptidoglycan.[66]

P. aeruginosa is known to produce over 50 quinolones[67] in three structural sub-

classes based on the characterized molecules: 2-heptyl-4-quinolone (HHQ)[68], 2-heptyl-

3,4-dihydroxyquinoline (Pseudomonas Quorum Signal, PQS)[69] and 2- heptyl-4-

hydroxyquinoline N-oxide (HQNO).[70] Different quinolones have different roles in intra-

and interspecies interactions within the CF lung environment.[71-75] HHQ and PQS are

important quorum sensing signaling metabolites participating in regulating the

production of a large suite of virulence factors, including PYO, and many genes related

to iron limitation.[68,69,76-78] HQNO lacks regulatory activity, but is an important metabolite

in inter-species interactions with *Staphylococcus aureus* through inhibition of the

respiratory chain[71,72] and enhancement of antibiotic tolerance.[79-81] While we quantified

43 quinolones from the PAO1 cultures in ASM (Figure 4A), only 13 account for 85-95%

of the total quinolones produced in ASM. Like phenazine production, levels of individual

and total quinolones produced by PAO1 varied between ASM formulations, with PAO1

cultured in Romling, Winstanley, and ASMDM ASM formulations producing the highest

levels of quinolones (Figure 4B). These three formulations have the highest

concentration of the aromatic amino acids tryptophan, tyrosine, and phenylalanine.

Importantly, tryptophan levels are low in CF sputum;[7] therefore, these media may not

replicate the amino acid availability *in vivo.* Higher concentrations of tryptophan may

alter quinolone production as it is catabolized via the kynurenine pathway to generate

anthranilate, a precursor of quinolone biosynthesis.[42,82] Both tyrosine and phenylalanine

are important carbon sources in CF sputum and have been shown to induce PQS

biosynthesis in SCFM1.[7,83]

In all ASM, PAO1 produced predominantly HHQ-type quinolones (44-76%), while

HQNO-type were produced at 15-41% total quinolones, and PQS-type were the

consistently the lowest abundant quinolones (3-17%) (Fig. 4C).The most abundant alkyl

chain lengths were heptyl (C7), nonyl (C9), and nonenyl (C9:1), accounting for 33-60%,

24-41%, and 10-21% of the quinolones produced by PAO1 in each ASM, respectively.

HHQ- and HQNO-type quinolones have been suggested for use as diagnostic

biomarkers of *P. aeruginosa* infection in adults and children with CF[50] as they are the

most abundant quinolone congeners in sputum[30,50,84,85], explant lung tissue[86,87], BALF[50],

and serum[50]. Importantly, ASM formulation affects the ratio of C7, C9:1, and C9

quinolones produced by PAO1, most prominently for HHQ- and HQNO-type quinolones.

For example, PAO1 cultured in Soothill ASM produces 2.5-fold more C7-HQ (HHQ)

than C9-HQ (NHQ), while in SCFM2 PAO1 produces similar levels of both HHQ-type

quinolones. It is unknown whether the C9 and C9:1 quinolone congeners have different

biological function than their C7 analogues. The difference in quinolone production

specificity between Soothill and SCFM ASM lineages may be the result of different

sources of lipid precursors. Soothill-derived ASM includes egg yolk emulsion as a lipid

source, while SCFM-derived ASM include a single lipid species (1,2-Dioleoyl-sn-

glycero-3-phosphocholine, DOPC), although commercial PGM contains lipids as well

(Figure 8).

Rhamnolipids are important virulence-associated glycolipids with functional roles

in biofilm formation and maintenance[88,89], evasion of host immune response[90,91],

competition with fungal neighbors[92], transport of PQS[93], and as biosurfactants[94].

Rhamnolipids have been detected in CF sputum[30,47,84,85] as well as explant lung

tissue[86,87], and rhamnolipid production levels are higher in some virulent *P. aeruginosa*

clinical isolates.[38] Structurally, *P. aeruginosa* rhamnolipids consist of 1-2 rhamnose units

bound to two fatty acids, yielding mono- and di-rhamnolipids, respectively. We

quantified six rhamnolipids, three mono-rhamnolipids and three di-rhamnolipids (Figure

5A). Unlike phenazine and quinolone production by PAO1 in the different ASM

formulations, rhamnolipid production was somewhat consistent across most cultures,

with lowest levels of production by PAO1 in SCFM2 and SCFM3 (Figure 5B). The

proportion of mono- to di-rhamnolipids produced by PAO1 was maintained across all

ASM cultures (Figure 5C), with di-rhamnolipids accounting for 56-70% of total

rhamnolipids. The predominant fatty acid chain length was C10, accounting for 56-70%

of the rhamnolipids produced by PAO1 in ASM, which reflects the predominant

rhamnolipids detected from CF sputum[30,84,85] and explant lung samples.[86,87]

The most striking difference in specialized metabolite production by PAO1 in

different ASM formulations was the production of the siderophores pyochelin and

pyoverdine, which were produced by PAO1 in SCFM1 cultures (Figure 6). While

pyochelin was annotated by a MS/MS spectral match to the GNPS libraries[43], the nodes

corresponding to the pyoverdine molecular family were prioritized for annotation based

upon coloration of the pie chart, indicating high production by PAO1 in SCFM2 (Figure

6A). Siderophores are metal scavenging virulence factors[95,96] secreted in low iron

conditions by microbes to acquire environmental iron.[97-100] Pyoverdine is considered the

primary siderophore of *P. aeruginosa* as it has a higher affinity for iron than pyochelin,

displaces iron from host proteins, and is required for virulence in acute infection

models.[95,96,98,101-103] However, pyochelin production by *P. aeruginosa* typically occurs

first as pyoverdine production is only initiated by *P. aeruginosa* when iron

concentrations are extremely low.[104] Both pyochelin and pyoverdine have been

detected in CF sputum[84,85], with production of pyoverdine highly dependent on

strain[34,105]. Although pyoverdine biosynthesis was reported to be upregulated in SCFM1

by *P. aeruginosa*[7], we were surprised to detect pyoverdines as the production of this

molecular family by PAO1 is below our limit of detection when cultured in most media.

The high abundance of pyochelin and pyoverdine from SCFM1 PAO1 cultures suggests

that only this ASM formulation is iron limited, which does not align with the published

media compositions. Prior to addition of the iron chelator diethylenetriaminepentaacetic

acid (DPTA) to the media, Soothill ASM was measured to have an Fe^{2+} concentration of

9.5 µM[4], and the same iron concentration was presumed for Romling, Winstanley, and

ASMDM ASM formulations. Both Cordwell and SDSU ASM contain 0.000003% (w/v)

ferritin[5,8], which adds ~2.85 µM Fe^{3+} to the media. Last, the SCFM series of ASM all

include 3.60 µM Fe^{2+} from addition of iron sulfate.[7,11] As SCFM2 nutrient availability

differs from SCFM1 by only the addition of GlcNAc, DNA, DOPC, and mucin[11], these

results suggest that one of these additives is an undefined source of iron.

Commercial porcine gastric mucin (PGM) contains contaminants which affect

PAO1 specialized metabolite production. As mucin was the most complex additive to

our SCFM2 media, we hypothesized that it was a potential source of contaminating iron.

Mucins are high molecular weight glycoproteins responsible for the viscoelastic

properties of mucus.[106,107] In CF, loss of function of the cystic fibrosis transmembrane

conductance regulator (CFTR) leads to altered ion transport and accumulation of thick

bronchial mucus in the lungs.[25,108] The altered biophysical properties of pulmonary

mucus in CF contributes to the establishment of chronic infections by pathogens,

including *P. aeruginosa*.[109-111] Due to its importance in CF pathogenesis, inclusion of

mucin in ASM is critical to fully replicating the *in vivo* nutritional[26,112,113] and

structural[112,114-116] CF pulmonary environment.[13,116,117] However, despite the limitations

associated with altered physical and chemical properties of the mucins associated with

the purification process and inclusion of a number of mucin-bound impurities[118-120], the

most economical choice is commercial porcine gastric mucin (PGM), which is often

used as a component of ASM without further purification.

Indeed, SCFM1 complemented with 0.5% (w/v) PGM (PGM-SCFM1) was

sufficient to alter production of the phenazine, quinolone, and rhamnolipid molecular

families as well as the siderophores, pyochelin and pyoverdine by PAO1 compared to

SCFM1 (Figure 7). Overall phenazine production was increased in PAO1 cultures in

PGM-SCFM1 (Figure 7A), with increased abundance of 1-HP, PYO, and PCA. Unlike

phenazine abundance, quinolone production by PAO1 was similar in the two media

(Figure 7B). However, the composition of the quinolone molecular family was altered

with PGM addition, with an increase in C7-HQ (HHQ) and a decrease in C9:1-HQ-OH

and C9-QNO (NQNO). Importantly, PQS-type quinolones were not detected in this

experiment. Surprisingly, although HHQ production increased with PGM addition,

C9-HQ (NHQ) production did not. Conversely, while C9-QNO (NQNO) production

decreased, C7-QNO (HQNO) production remained the same. Rhamnolipid production

was decreased in PGM-SCFM1 PAO1 cultures (Figure 7C). Lastly, addition of PGM to

SCFM1 was sufficient to reduce siderophore production by PAO1 (Figure 7D). The

changes in specialized metabolite production by PAO1 in PGM-SCFM1 compared to

the SCFM1 cultures parallels the differential production of these metabolites between

SCFM1 and SCFM2 cultures, indicating that the other additives are minimally

contributing to these alterations.

As the composition of commercial mucins is largely undefined, the alterations in

production of small molecule virulence factors by PAO1 cultured in PGM-SCFM1 could

be due to the presence of mucin, assorted iron sources, GlcNAc, or another component

in the PGM. Confirming our suspicion that commercial PGM contains iron, ICP-MS

analysis revealed 0.5% PGM contains approximately 30 μM iron (Figure 7E). The

abundance of all metals quantified from PGM are summarized in Figure 8A. The

concentration of iron in commercial PGM is 8.5 times higher than the concentration of

iron added to SCFM1 medium (3.6 μM). As ASMDM and Cordwell ASM formulations

contain 1.0% (w/v) PGM and SDSU ASM contains 2.0% (w/v) PGM, the levels of iron in

these media correspond to two and four times higher than the other ASM formulations,

respectively. Inclusion of PGM in SCFM1 increases the concentration of iron in the

medium to a more clinically relevant level of iron.[121-123] However, the distinct iron

sources in PGM are unknown and may not represent those in the CF lung.[122,124]

Due to the role of iron concentration in regulating specialized metabolite

production by *P.* aeruginosa, we are confident that the high concentration of iron

contributed by PGM is affecting the production levels of these metabolites in PGM-

SCFM1.[89,125,126] However, during analysis of the FBMN of SCFM1 and PGM-SCFM1

cultures, several molecular features matched to lipid spectra in the GNPS libraries,

which could only be present in the PGM, as they were not separately added to the

media. Analysis of lipophilic extracts of commercial PGM using classical molecular

networking (CMN) revealed that in addition to contamination with a variety of metals,

PGM contains amino acids, small peptides, and lipids (Figure 8B). Although SCFM1 is

tryptophan limited, tryptophan was detected in PGM. Therefore, addition of PGM to the

medium increases the tryptophan concentration, which can serve as a source of

anthranilate in quinolone biosynthesis.[127] Like amino acids, bioactive host lipids may

play important roles in shaping CF lung ecology, including eicosanoids, sphingolipids,

and ceramides.[128,129] PGM contains a wide variety of lipids (Figure 8B), including those

most highly detected in CF sputum samples.[30,128] Detection of these contaminants

within commercial PGM suggest that inclusion of crude PGM in culture medium will

likely yield null results when investigating the mechanistic roles of these metabolites (as

well as those not detected under our extraction conditions) on microbial physiology,

virulence, and ecology.

Further research will be required to establish what specific components of

commercial PGM are responsible for alteration in PAO1 specialized metabolite

production and to develop cost-effective methods for purification of mucin from crude

preparations. The mucin concentration of CF sputum ranges from 16-30 mg/mL.[25] While

methods have been established for isolating pure PGM from pig stomach, only ~65 mg

of PGM can be purified from a single stomach.[118] To produce 1 L of an ASM formulation

containing 0.5% (w/v) mucin (5 mg/mL), this purification would require processing of at

least 77 pig stomachs. Other laboratories have established procedures for purifying

crude commercial mucins[130], but without performing analytical analysis to measure all

potential contaminants, it is impossible to assess the purity of the mucins.

CONCLUSION

In vitro model systems are attractive due to their broad accessibility and ability to probe

hypotheses thoroughly. The foundation of any *in vitro* investigation is the medium. In

CF, a broad suite of different ASM formulations has been created. In this study, we

demonstrated that differences in nutrient content between ASM formulations alter the

phenotype and metabolomics profile of *P. aeruginosa* PAO1. Importantly, the

concentration of specialized metabolites, including members of the phenazine,

quinolone, rhamnolipids, and siderophore molecular families, are impacted by ASM

formulation. Given the role of these metabolites in virulence, differences in ASM

formulation may influence the overall outcome of interactions of *P. aeruginosa* with host

cells and other members of the CF pulmonary microbiota. Commercial PGM was

identified as a source of contaminating bioavailable metals, lipids, amino acids, and

small peptides. Addition of commercial PGM to the chemically defined SCFM1 ASM

formulation was sufficient to alter production of small molecule virulence factors by

PAO1. In view of the importance of mucin in replicating mucosal infection environments

in vitro, further work is needed to understand the contribution of individual components

of commercial mucin to microbial physiology and virulence, as well as development of

cost-effective means to purify mucins from crude commercial preparations.

MATERIALS AND METHODS

General: *Pseudomonas aeruginosa* PAO1 (MPAO1) was acquired from the University of Washington, Seattle[113,131]. All media components, concentrations, and commercial sources for artificial sputum media (ASM) formulations are listed in **Dataset S1 ASM formulations**[3-11]. Miller Luria broth (LB) and porcine gastric mucin type III (PGM) were purchased from Millipore Sigma. Organic solvents for extraction and liquid chromatography tandem mass spectrometry (LC-MS/MS) analysis were purchased from VWR (ethyl acetate (EtOAc), HiPerSolv Chromanorm purity) and Fisher Scientific (methanol (MeOH), acetonitrile (ACN), formic acid (FA), and water; all Optima LC/MS grade). PA Mix was created as a set of external standards for *P. aeruginosa* specialized metabolite annotation. PA Mix contains 10 µM of each of the following: 1-hydroxyphenazine (1-HP, Tokyo Chemical Industry); pyocyanin (PYO, Sigma); phenazine-1-carboxylic acid (PCA, Ark Pharm); phenazine-1-carboxamide (PCN, Ark Pharm); 2-heptylquinolin-4(1H)-one (HHQ, Ark Pharm); 2-heptyl-4-hydroxyquinoline *N*-oxide (HQNO, Cayman Chemical); 2-heptyl-3-hydroxyl-4-quinolone (Pseudomonas Quinolone Signal, PQS, Chemodex); N-butyryl-l-homoserine lactone (C4-HSL, Cayman Chemical); N-(3-oxododecanoyl)-l-homoserine lactone (3-oxo-C12-HSL, Millipore Sigma); and a rhamnolipid mixture (RHL, AGAE Technologies, 90% pure).

Identification of commonly used ASM formulations: The Web of Science Core Collection (Clarivate Analytics) was searched for papers with the 'TOPIC' terms: "sputum medium", "sputum media", "cystic fibrosis medium", "cystic fibrosis media", "sputa media", "sputa medium", "CF media", or "CF medium" in the title, abstract, or keywords fields on May 10, 2020. All terms were searched as shown, including

quotation marks. The results of this search were analyzed using the Web of Science

tools 'Create Citation Report' and 'Analyze Results'. The Analyze Results tool was used

to download data corresponding to Web of Science Categories. The citation report was

used to select ASM formulations that were commonly cited.

Preparation of media: LB broth was prepared according to manufacturer instructions.

Commercial PGM was suspended in 1X 3-morpholinopropane-1-sulfonic acid (MOPS)

buffer (pH 7.0) to make a 10% (w/v) suspension, sterilized using a liquid autoclave cycle

(<20 minute sterilization time), and an aliquot was checked for sterility before use. ASM

formulations were prepared as reported[3-11] with the following modifications. Antibiotics

were not included in any medium. For the Cordwell formulation[5], the reported 0.1%

(w/v) agar was excluded. For SCFM2 and SCFM3[11], bovine submaxillary mucin (BSM)

was replaced with PGM at the published concentration, a common substitution[12-14].

Each ASM formulation was brought to the published pH[3-11]. All media was stored in the

dark at 4°C, checked for sterility prior to use, and used within one month of preparation.

Cultivation of *P. aeruginosa* in LB and ASM formulations: PAO1 was inoculated

from a streak plate into 5 mL LB broth and incubated overnight at 37°C, shaking at 220

RPM. 1.98 mL of LB or ASM were inoculated with 20 µL of *P. aeruginosa* LB broth

culture (OD_{600} of 0.05; ~10^6 CFU/mL final concentration) in polystyrene 24 well plates[6].

For assessing *P. aeruginosa* growth and specialized metabolite production, each

medium was evaluated in its own 24 well plate (12 medium control wells and 12 growth

wells per medium). The media were inoculated and processed in three batches,

comprised of 2-4 ASM formulations and a parallel LB plate as a control. All plates were

covered and incubated statically at 37°C for 72 hours. For assessing the addition of

PGM to SCFM1 on *P. aeruginosa* growth and specialized metabolite production, *P.*

aeruginosa was cultured in SCFM1 and PGM-SCFM1 in a single polystyrene 24 well

plate (4 medium control wells and 4 growth wells per medium).

Processing of *P. aeruginosa* cultures in LB and ASM formulations: Gross

phenotypes of *P. aeruginosa* growth were photographed using a top-down view with a

Stemi 508 stereo microscope with an Axiocam 105 color camera (Zeiss). Following

visualization, 100 µL cellulase (*Aspergillus niger*, 100 mg/mL in sterile water; Sigma)

was added to each well and aggregates were mechanically disrupted by pipetting using

wide-bore pipette tips (USA Scientific), followed by water bath sonication (Branson) for

10 minutes, and a secondary mechanical disruption by pipetting using wide-bore pipette

tips. Disrupted samples were aliquoted for measurements of growth (CFUs) and

metabolomics analysis. Three representative biological replicates from each culture

condition were serially diluted, spotted onto LB agar, incubated at 37°C overnight, and

counted to determine CFU/mL.

Processing of *P. aeruginosa* cultures in SCFM1 and PGM-SCFM1: Aggregates were

mechanically disrupted by pipetting using wide-bore pipette tips (USA Scientific).

Disrupted samples were aliquoted for measurements of growth (CFUs) and

metabolomics analysis. All four biological replicates from each culture condition were

serially diluted, spotted onto LB agar, incubated at 37°C overnight, and counted to

determine CFU/mL.

ASM sample preparation for LC-MS/MS: Each sample aliquot was chemically

disrupted with an equal volume of 1:1 EtOAc:MeOH. The samples were dried,

resuspended in 100% MeOH containing 1 µM glycocholic acid (Calbiochem, 100.1%

pure), diluted as needed in 100% MeOH containing 1 µM glycocholic acid, and

centrifuged for 10 min at 4,000 RPM (Thermo Sorvall ST 40R) to remove non-soluble

particulates. Samples were diluted as follows for measurement of *P. aeruginosa*

specialized metabolite production in ASM formulations and LB: not diluted: Soothill

ASM; 5-fold dilution: Romling ASM, Cordwell ASM, SCFM1, SCFM2, and SCFM3; 10-

fold dilution: Winstanley ASM, ASMDM, SDSU ASM, and LB. All samples for

measurement of *P. aeruginosa* specialized metabolite production in SCFM1 and PGM-

SCFM1 were diluted 10-fold.

PGM sample preparation for LC-MS/MS: 50 µL of 10% (w/v) PGM in 1X MOPS was

chemically extracted in technical triplicate with 200 µL of 1:1 EtOAc:MeOH and following

the Bligh-Dyer extraction protocol.[132] The samples were dried and resuspended in

100% MeOH containing 1 µM glycocholic acid, centrifuged for 10 min at 4,000 RPM,

and diluted 10-fold.

LC-MS/MS data collection: Mass spectrometry data acquisition for all samples was

performed using a Bruker Daltonics Maxis II HD qTOF mass spectrometer equipped

with a standard electrospray ionization (ESI) source. The mass spectrometer was tuned

by infusion of Tuning Mix ESI-TOF (Agilent Technologies) at a 3 µL/min flow rate. For

accurate mass measurements, a wick saturated with Hexakis(1H,1H,2H-difluoroethoxy)

phosphazene ions (Apollo Scientific, *m/z* 622.1978) located within the source was used

as a lock mass internal calibrant. Samples were introduced by an Agilent 1290 UPLC

using a 10 µl injection volume for ASM samples and a 5 µl injection volume for PGM

samples. Extracts were separated using a Phenomenex Kinetex 2.6 µm C18 column

(2.1 mm x 50 mm) using a 9 minute, linear water-ACN gradient (from 98:2 to 2:98%

water:ACN) containing 0.1% FA at a flow rate of 0.5 mL/min. The mass spectrometer

was operated in data dependent positive ion mode, automatically switching between full

scan MS and MS/MS acquisitions. Full scan MS spectra (*m/z* 50 - 1500) were acquired

in the TOF and the top five most intense ions in a particular scan were fragmented via

collision induced dissociation (CID) using the stepping function in the collision cell.

LC-MS/MS data for PA mix were acquired under identical conditions. Bruker Daltonics

CompassXport was used to apply lock mass calibration and convert the LC-MS/MS

data from .d format to .mzXML format.

Feature based molecular networking (FBMN) of *P. aeruginosa* specialized

metabolism in LB and ASM formulations: MZmine (version 2.53)[133] was used to

perform feature finding on the .mzXML files associated with this experiment. The output

was a feature table containing area under the curve (AUC) integration values for each

feature (*m/z*-RT pair) and an .mgf file containing MS/MS spectra for each feature. The

feature table was corrected for sample dilutions prior to submission to GNPS. The

Feature Networking workflow (version release 22)[44] was applied to the dilution-

corrected feature table, .mgf file, and a metadata file aligning to GNPS specifications

using the GNPS web platform[43]. Briefly, the data was filtered by removing all MS/MS

fragment ions ± 17 Da of the precursor *m/z*. MS/MS spectra were window filtered by

choosing only the top 6 fragment ions in each ± 50 Da window throughout the spectrum.

The precursor ion mass tolerance was set to 0.05 Da and the MS/MS fragment ion

tolerance was set to 0.1 Da. A network of nodes connected by edges was then created.

Edges were filtered to have a cosine score above 0.75. Further, edges between two

nodes were kept in the network only if each of the nodes appeared in each other's

respective top 5 most similar nodes. Finally, the maximum size of a molecular family

was set to 50, and the lowest scoring edges were removed from molecular families until

the molecular family was below this threshold. The spectra in the network were then

searched against the GNPS spectral libraries. The library spectra were filtered in the

same manner as the input data. All matches kept between the network spectra and

library spectra were required to have a cosine score above 0.6 and at least 5 matched

peaks. Sum normalization and mean aggregation were applied. The molecular network

(https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=baba5df5ab1b437980f81b4e64bcff

c0) was visualized using Cytoscape (version 3.7.1)[134].

FBMN and targeted quantification of *P. aeruginosa* specialized metabolites in

SCFM1 and PGM-SCFM1 cultures. MZmine (version 2.53)[133] was used to perform

feature finding on the .mzXML files associated with this experiment. The output was a

feature table containing area under the curve (AUC) integration values for each feature

(*m/z*-RT pair) and an .mgf file containing MS/MS spectra for each feature. The feature

table was corrected for sample dilutions prior to submission to GNPS. The Feature

Networking workflow (version release 28.1)[44] was applied to the dilution-corrected

feature table, .mgf file, and a metadata file aligning to GNPS specifications using the

GNPS web platform[43]. Briefly, the data was filtered by removing all MS/MS fragment

ions ± 17 Da of the precursor *m/z*. MS/MS spectra were window filtered by choosing

only the top 4 fragment ions in each ± 50 Da window throughout the spectrum. The

precursor ion mass tolerance was set to 0.02 Da and the MS/MS fragment ion tolerance

was set to 0.02 Da. A network of nodes connected by edges was then created. Edges

were filtered to have a cosine score above 0.7. Further, edges between two nodes were

kept in the network only if each of the nodes appeared in each other's respective top 10

most similar nodes. Finally, the maximum size of a molecular family was set to 100, and

the lowest scoring edges were removed from molecular families until the molecular

family was below this threshold. The spectra in the network were then searched against

the GNPS spectral libraries. The library spectra were filtered in the same manner as the

input data. All matches kept between the network spectra and library spectra were

required to have a cosine score above 0.7 and at least 4 matched peaks. Sum

normalization and mean aggregation were applied. The molecular network

(https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=5624bf8b3174483a89cf733d08ce0f

ff) was visualized using Cytoscape (version 3.7.1)[134].

Feature annotation of *P. aeruginosa* specialized metabolites: Annotations of

features corresponding to commercial standards (level 1 annotation[135]) were confirmed

by comparing the experimental data (exact mass, MS/MS, and retention time) with data

acquired for the compound in PA mix using Bruker Daltonics DataAnalysis v4.1 (Build

362.7). Putative annotation of specialized metabolites based on matches to the GNPS

libraries (level 2 annotation[135]) and molecular families (level 3 annotation[135]) were

validated by comparing the experimental data (exact mass, MS/MS) to reported data[58]

and putative structures, respectively[58]. Annotated features corresponding to *P.*

aeruginosa specialized metabolites identified from FBMN analysis of LB and ASM

samples and SCFM1 vs PGM-SCFM1 are listed in **Table S1. PA in ASM metabolite**

annotation and mass defect and **Table S2. PA in SCFM1/PGM-SCFM1 metabolite**

annotation and mass defect, respectively. Multiple features (different adducts, in-

source fragments, and/or in-source dimers) can represent the same metabolite.

Therefore, sum normalized AUC values for features corresponding to the same

metabolite of interest (identified by co-elution and mass defect from predicted exact

mass of the compound adduct, fragment, and/or dimer) were summed to provide the

total AUC for a metabolite. Sum normalized AUC values for metabolites representing

the same sub-structure class based upon MS/MS fragmentation pattern (e.g. HHQ-type

AQs or C10-C10 containing RHLs) were summed to provide the total AUC for a sub-

structure class. Sum normalized AUC values for all identified metabolites representing

the same molecular family based upon MS/MS fragmentation patterns (e.g. AQs or

RHLs) were summed to provide the total AUC for a molecular family**.**

ICP-MS analysis of PGM: Metals were quantified from 2.5% (w/v) PGM suspended in

1X MOPS using inductively coupled plasma mass spectrometry (ICP-MS) analysis at

the University of Nebraska - Lincoln Spectroscopy and Biophysics Core facility.

Samples were diluted 1:1 with nitric acid, digested overnight, and diluted an additional

10-fold before conducting ICP-MS (Agilent 7500 cx). Gallium (Ga), 50 ppb, was added

to each sample as an internal standard. Samples were analyzed in technical triplicate

with a blank wash run between each sample. Only concentrations above the limit of

quantification were retained. To determine the amount of iron added to media by the

addition of PGM, or 1X MOPS when used in place of PGM, the concentration of PGM

was brought to that present in most ASM formulations (0.5% w/v) and converted to

micromolar concentration.

Classical molecular networking of PGM lipophilic extracts: A classical molecular

network was created from the .mzXML files using the Molecular Networking workflow

(version release 27) of the GNPS web platform (gnps.ucsd.edu,

https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=fc3ee81541aa4a60af38b15f087896

20)[43,67]. Briefly, the data was filtered by removing all MS/MS fragment ions ± 17 Da of

the precursor *m/z*. MS/MS spectra were window filtered by choosing only the top 6

fragment ions in each ± 50 Da window throughout the spectrum. The precursor ion

mass tolerance was set to 0.05 Da and the MS/MS fragment ion tolerance to 0.5 Da. A

network was then created where edges were filtered to have a cosine score above 0.7

and at least 4 matched fragment peaks between corresponding nodes. Further, edges

between two nodes were kept in the network only if each of the nodes appeared in each

other's respective top 10 most similar nodes. Finally, the maximum size of a molecular

family was set to 100, and the lowest scoring edges were removed from molecular

families until the molecular family was below this threshold. The spectra in the network

were then searched against the GNPS spectral libraries. The library spectra were

filtered in the same manner as the input data. All matches kept between the network

spectra and library spectra were required to have a cosine score above 0.6 and at least

4 matched peaks. The molecular network was visualized using Cytoscape (version

3.7.1)[134].

Statistical analysis: Statistical comparison of FBMN features of *P. aeruginosa* cultured

in ASM formulations was conducted on the sum normalized feature table using ANOVA

with Tukey's correction for multiple comparisons in MetaboAnalyst 4.0[136-139]. Statistical

comparison of individual metabolites, sub-structure classes, and molecular families

performed using ANOVA with Tukey's multiple comparisons correction in GraphPad

Prism (version 8.4.0). For all analyses, $P < 0.05$ were considered statistically significant.

All statistical results are summarized in Dataset S2 ASM statistical analyses.

Data availability: All mass spectrometry data including the raw files, .mzXML files,

metadata tables, MZmine settings, and Cytoscape files are available via MassIVE:

MSV000086721 (LB and ASM formulations – media control, *P. aeruginosa* culture, QC,

and PA mix data; corresponds to Figures 2-6); MSV000086723 (SCFM1 and PGM-

SCFM1 – media control, *P. aeruginosa* culture, QC, and PA mix data; corresponds to

Figure 7); and MSV000087153 (PGM lipophilic extracts; corresponds to Figure 8).

ACKNOWLEDGEMENTS:

We are very grateful to Mingxun Wang (University of California, San Diego) for

providing input and beta versions of the networking software for metabolomics analysis;

Laura Sanchez, Jacob Porter, and Katherine Zink (University of Illinois, Chicago) for

providing initial collection of metabolomics data; and members of the Phelan lab for

thoughtful discussion about and suggestions on improving this manuscript.

REFERENCES

1   Fritz, J. V., Desai, M. S., Shah, P., Schneider, J. G. & Wilmes, P. From meta-
      omics to causality: experimental models for human microbiome research.
      Microbiome **1**, 14, doi:10.1186/2049-2618-1-14 (2013).

2   Coenye, T. & Nelis, H. J. In vitro and in vivo model systems to study microbial
      biofilm formation. *J Microbiol Methods* **83**, 89-105,
      doi:10.1016/j.mimet.2010.08.018 (2010).

3   Fung, C. *et al.* Gene expression of Pseudomonas aeruginosa in a mucin-
      containing synthetic growth medium mimicking cystic fibrosis lung sputum. *J Med
      Microbiol* **59**, 1089-1100, doi:10.1099/jmm.0.019984-0 (2010).

4   Ghani, M. & Soothill, J. S. Ceftazidime, gentamicin, and rifampicin, in
      combination, kill biofilms of mucoid Pseudomonas aeruginosa. *Canadian Journal
      of Microbiology* **43**, 999-1004, doi:DOI 10.1139/m97-144 (1997).

5   Hare, N. J. *et al.* Proteomics of Pseudomonas aeruginosa Australian Epidemic
      Strain 1 (AES-1) Cultured under Conditions Mimicking the Cystic Fibrosis Lung
      Reveals Increased Iron Acquisition via the Siderophore Pyochelin. *J Proteome
      Res* **11**, 776-795, doi:10.1021/pr200659h (2012).

6   Kirchner, S. *et al.* Use of Artificial Sputum Medium to Test Antibiotic Efficacy
      Against Pseudomonas aeruginosa in Conditions More Relevant to the Cystic
      Fibrosis Lung. *Jove-J Vis Exp,* doi:ARTN e3857 10.3791/3857 (2012).

7   Palmer, K. L., Aye, L. A. & Whiteley, M. Nutritional cues control Pseudomonas
      aeruginosa multicellular Behavior in cystic fibrosis sputum. *J Bacteriol* **189**, 8079-
      8087, doi:10.1128/Jb.01138-07 (2007).

594   8   Quinn, R. A. *et al.* A Winogradsky-based culture system shows an association

595   between microbial fermentation and cystic fibrosis exacerbation (vol 9, pg 1024,

596   2015). *Isme J* **9**, 1052-1052, doi:10.1038/ismej.2014.266 (2015).

597   9   Sriramulu, D. D. Artifical Sputum Medium. *Protocol Exchange*,

598   doi:doi:10.1038/protex.2010.212 (2010).

599   10   Sriramulu, D. D., Lunsdorf, H., Lam, J. S. & Romling, U. Microcolony formation: a

600   novel biofilm model of Pseudomonas aeruginosa for the cystic fibrosis lung. *J*

601   *Med Microbiol* **54**, 667-676, doi:10.1099/jmm.0.45969-0 (2005).

602   11   Turner, K. H., Wessel, A. K., Palmer, G. C., Murray, J. L. & Whiteley, M.

603   Essential genome of Pseudomonas aeruginosa in cystic fibrosis sputum. *P Natl*

604   *Acad Sci USA* **112**, 4110-4115, doi:10.1073/pnas.1419677112 (2015).

605   12   Homa, M. *et al.* In vitro Interactions of Pseudomonas aeruginosa With

606   Scedosporium Species Frequently Associated With Cystic Fibrosis. *Front*

607   *Microbiol* **10**, 441, doi:10.3389/fmicb.2019.00441 (2019).

608   13   Schick, A. & Kassen, R. Rapid diversification of Pseudomonas aeruginosa in

609   cystic fibrosis lung-like conditions. *Proc Natl Acad Sci U S A* **115**, 10714-10719,

610   doi:10.1073/pnas.1721270115 (2018).

611   14   Wong, A., Rodrigue, N. & Kassen, R. Genomics of adaptation during

612   experimental evolution of the opportunistic pathogen Pseudomonas aeruginosa.

613   *PLoS Genet* **8**, e1002928, doi:10.1371/journal.pgen.1002928 (2012).

614   15   Behrends, V., Geier, B., Williams, H. D. & Bundy, J. G. Direct Assessment of

615   Metabolite Utilization by Pseudomonas aeruginosa during Growth on Artificial

Sputum Medium. *Applied and Environmental Microbiology* **79**, 2467-2470, doi:10.1128/AEM.03609-12 (2013).

Darch, S. E. *et al.* Spatial determinants of quorum signaling in a Pseudomonas aeruginosa infection model. *PNAS* **115**, 4779-4784, doi:10.1073/pnas.1719317115 (2018).

Behrends, V., Ebbels, T. M. D., Williams, H. D. & Bundy, J. G. Time-Resolved Metabolic Footprinting for Nonlinear Modeling of Bacterial Substrate Utilization. *Applied and Environmental Microbiology* **75**, 2453-2463, doi:10.1128/AEM.01742-08 (2009).

Haley, C. L., Colmer-Hamood, J. A. & Hamood, A. N. Characterization of biofilm-like structures formed by Pseudomonas aeruginosa in a synthetic mucus medium. *BMC Microbiology* **12**, 181, doi:10.1186/1471-2180-12-181 (2012).

Diaz Iglesias, Y. & Van Bambeke, F. Activity of Antibiotics against Pseudomonas aeruginosa in an In Vitro Model of Biofilms in the Context of Cystic Fibrosis: Influence of the Culture Medium. *Antimicrobial Agents and Chemotherapy* **64**, doi:10.1128/AAC.02204-19 (2020).

Stevens, D. A., Moss, R. B., Hernandez, C., Clemons, K. V. & Martinez, M. Effect of Media Modified To Mimic Cystic Fibrosis Sputum on the Susceptibility of Aspergillus fumigatus, and the Frequency of Resistance at One Center. *Antimicrob Agents Chemother* **60**, 2180-2184, doi:10.1128/AAC.02649-15 (2016).

Lightly, T. J., Phung, R. R., Sorensen, J. L. & Cardona, S. T. Synthetic cystic fibrosis sputum medium diminishes Burkholderia cenocepacia antifungal activity

against Aspergillus fumigatus independently of phenylacetic acid production.

Canadian Journal of Microbiology **63**, 427-438, doi:10.1139/cjm-2016-0705

(2017).

22 Kirchhoff, L. *et al.* Phenotypical Characteristics of the Black Yeast Exophiala

dermatitidis Are Affected by Pseudomonas aeruginosa in an Artificial Sputum

Medium Mimicking Cystic Fibrosis–Like Conditions. *Frontiers in Microbiology* **11**,

doi:10.3389/fmicb.2020.00471 (2020).

23 Limoli, D. H. *et al.* Pseudomonas aeruginosa Alginate Overproduction Promotes

Coexistence with Staphylococcus aureus in a Model of Cystic Fibrosis

Respiratory Infection. *mBio* **8**, doi:10.1128/mBio.00186-17 (2017).

24 Kreda, S. M., Davis, C. W. & Rose, M. C. CFTR, mucins, and mucus obstruction

in cystic fibrosis. *Cold Spring Harb Perspect Med* **2**, a009589,

doi:10.1101/cshperspect.a009589 (2012).

25 Sanders, N. N., Van Rompaey, E., De Smedt, S. C. & Demeester, J. Structural

Alterations of Gene Complexes by Cystic Fibrosis Sputum. *American Journal of*

Respiratory and Critical Care Medicine **164**, 486-493,

doi:10.1164/ajrccm.164.3.2011041 (2001).

26 Schwab, U. *et al.* Localization of Burkholderia cepacia complex bacteria in cystic

fibrosis lungs and interactions with Pseudomonas aeruginosa in hypoxic mucus.

Infect. Immun. **82**, 4729-4745, doi:10.1128/IAI.01876-14 (2014).

27 La Rosa, R., Johansen, H. K. & Molin, S. Adapting to the Airways: Metabolic

Requirements of Pseudomonas aeruginosa during the Infection of Cystic Fibrosis

Patients. *Metabolites* **9**, doi:10.3390/metabo9100234 (2019).

28   Wolak, J. E., Esther, C. R., Jr. & O'Connell, T. M. Metabolomic analysis of

bronchoalveolar lavage fluid from cystic fibrosis patients. *Biomarkers* **14**, 55-60,

doi:10.1080/13547500802688194 (2009).

29   Colmer-Hamood, J. A., Dzvova, N., Kruczek, C. & Hamood, A. N. in *Progress in*

Molecular Biology and Translational Science Vol. 142   151-191 (Elsevier, 2016).

30   Quinn, R. A. *et al.* Microbial, host and xenobiotic diversity in the cystic fibrosis

sputum metabolome. *The ISME Journal* **10**, 1483-1498,

doi:10.1038/ismej.2015.207 (2016).

31   Foundation, C. F. 2018 Annual Data Report. (Bethesda, Maryland, 2019).

32   Konstan, M. W. *et al.* Risk factors for rate of decline in FEV1 in adults with cystic

fibrosis. *J Cyst Fibros* **11**, 405-411, doi:10.1016/j.jcf.2012.03.009 (2012).

33   Ciofu, O., Tolker-Nielsen, T., Jensen, P. Ø., Wang, H. & Høiby, N. Antimicrobial

resistance, respiratory tract infections and role of biofilms in lung infections in

cystic fibrosis patients. *Advanced Drug Delivery Reviews* **85**, 7-23,

doi:10.1016/j.addr.2014.11.017 (2015).

34   Martin, L. W., Reid, D. W., Sharples, K. J. & Lamont, I. L. Pseudomonas

siderophores in the sputum of patients with cystic fibrosis. *Biometals* **24**, 1059-

1067, doi:10.1007/s10534-011-9464-z (2011).

35   Workentine, M. L. *et al.* Phenotypic Heterogeneity of Pseudomonas aeruginosa

Populations in a Cystic Fibrosis Patient. *PLoS ONE* **8**, e60225,

doi:10.1371/journal.pone.0060225 (2013).

36   Sharon, G. *et al.* Specialized Metabolites from the Microbiome in Health and

Disease. *Cell Metabolism* **20**, 719-730, doi:10.1016/j.cmet.2014.10.016 (2014).

37   Condren, A. R. *et al.* Biofilm Inhibitor Taurolithocholic Acid Alters Colony

Morphology, Specialized Metabolism, and Virulence of Pseudomonas

aeruginosa. *ACS Infect Dis* **6**, 603-612, doi:10.1021/acsinfecdis.9b00424 (2020).

38   Depke, T., Thoming, J. G., Kordes, A., Haussler, S. & Bronstrup, M. Untargeted

LC-MS Metabolomics Differentiates Between Virulent and Avirulent Clinical

Strains of Pseudomonas aeruginosa. *Biomolecules* **10**,

doi:10.3390/biom10071041 (2020).

39   Blankenfeldt, W. & Parsons, J. F. The structural biology of phenazine

biosynthesis. *Curr Opin Struct Biol* **29**, 26-33, doi:10.1016/j.sbi.2014.08.013

(2014).

40   Ringel, M. T. & Brüser, T. The biosynthesis of pyoverdines. *Microbial Cell* **5**, 424-

437, doi:10.15698/mic2018.10.649 (2018).

41   Mavrodi, D. V. *et al.* Functional Analysis of Genes for Biosynthesis of Pyocyanin

and Phenazine-1-Carboxamide from Pseudomonas aeruginosa PAO1. *J*

Bacteriol **183**, 6454-6465, doi:10.1128/JB.183.21.6454-6465.2001 (2001).

42   Farrow, J. M., 3rd & Pesci, E. C. Two distinct pathways supply anthranilate as a

precursor of the Pseudomonas quinolone signal. *J Bacteriol* **189**, 3425-3433,

doi:10.1128/JB.00209-07 (2007).

43   Wang, M. *et al.* Sharing and community curation of mass spectrometry data with

Global Natural Products Social Molecular Networking. *Nature Biotechnology* **34**,

828-837, doi:10.1038/nbt.3597 (2016).

44   Nothias, L. F. *et al.* Feature-based molecular networking in the GNPS analysis

environment. *Nat Methods*, doi:10.1038/s41592-020-0933-6 (2020).

45   Phelan, V. V. *et al.* Impact of a Transposon Insertion in phzF2 on the Specialized

Metabolite Production and Interkingdom Interactions of Pseudomonas

aeruginosa. *J Bacteriol* **196**, 1683-1693, doi:10.1128/JB.01258-13 (2014).

46   Phelan, V. V., Fang, J. & Dorrestein, P. C. Mass Spectrometry Analysis of

Pseudomonas aeruginosa Treated with Azithromycin. *J Am Soc Mass Spectrom*

26, 873-877, doi:10.1007/s13361-015-1101-6 (2015).

47   Kownatzki, R., Tummler, B. & Doring, G. Rhamnolipid of Pseudomonas

aeruginosa in sputum of cystic fibrosis patients. *Lancet* **1**, 1026-1027,

doi:10.1016/s0140-6736(87)92286-0 (1987).

48   Collier, D. N. *et al.* A bacterial cell to cell signal in the lungs of cystic fibrosis

patients. *FEMS Microbiol Lett* **215**, 41-46, doi:10.1111/j.1574-

6968.2002.tb11367.x (2002).

49   Singh, P. K. *et al.* Quorum-sensing signals indicate that cystic fibrosis lungs are

infected with bacterial biofilms. *Nature* **407**, 762-764, doi:10.1038/35037627

(2000).

50   Barr, H. L. *et al.* Diagnostic and prognostic significance of systemic alkyl

quinolones for P. aeruginosa in cystic fibrosis: A longitudinal study. *J Cyst Fibros*

16, 230-238, doi:10.1016/j.jcf.2016.10.005 (2017).

51   Barr, H. L. *et al.* Pseudomonas aeruginosa quorum sensing molecules correlate

with clinical status in cystic fibrosis. *Eur Respir J* **46**, 1046-1054,

doi:10.1183/09031936.00225214 (2015).

52   Wilson, R. *et al.* Measurement of Pseudomonas aeruginosa phenazine pigments

in sputum and assessment of their contribution to sputum sol toxicity for

731   respiratory epithelium. *Infect Immun* **56**, 2515-2517, doi:10.1128/IAI.56.9.2515-

732   2517.1988 (1988).

733 53 Haas, B., Kraut, J., Marks, J., Zanker, S. C. & Castignetti, D. Siderophore

734   presence in sputa of cystic fibrosis patients. *Infect Immun* **59**, 3997-4000,

735   doi:10.1128/IAI.59.11.3997-4000.1991 (1991).

736 54 Wang, Y., Kern, S. E. & Newman, D. K. Endogenous Phenazine Antibiotics

737   Promote Anaerobic Survival of Pseudomonas aeruginosa via Extracellular

738   Electron Transfer. *J Bacteriol* **192**, 365-369, doi:10.1128/JB.01188-09 (2010).

739 55 Briard, B. *et al.* Pseudomonas aeruginosa manipulates redox and iron

740   homeostasis of its microbiota partner Aspergillus fumigatus via phenazines.

741   *Scientific Reports* **5**, doi:10.1038/srep08220 (2015).

742 56 Das, T. & Manefield, M. Pyocyanin Promotes Extracellular DNA Release in

743   Pseudomonas aeruginosa. *PLoS ONE* **7**, e46718,

744   doi:10.1371/journal.pone.0046718 (2012).

745 57 Das, T. & Manefield, M. Phenazine production enhances extracellular DNA

746   release via hydrogen peroxide generation in <i>Pseudomonas aeruginosa</i>.

747   *Communicative & Integrative Biology* **6**, e23570, doi:10.4161/cib.23570 (2013).

748 58 Moree, W. J. *et al.* Interkingdom metabolic transformations captured by microbial

749   imaging mass spectrometry. *P Natl Acad Sci USA* **109**, 13811-13816,

750   doi:10.1073/pnas.1206855109 (2012).

751 59 Noto, M. J., Burns, W. J., Beavers, W. N. & Skaar, E. P. Mechanisms of

752   Pyocyanin Toxicity and Genetic Determinants of Resistance in Staphylococcus

753   aureus. *J Bacteriol* **199**, doi:10.1128/JB.00221-17 (2017).

60 Glasser, N. R., Hunter, R. C., Liou, T. G., Newman, D. K. & Mountain West, C. F.

C. I. Refinement of metabolite detection in cystic fibrosis sputum reveals heme

correlates with lung function decline. *PLoS One* **14**, e0226578,

doi:10.1371/journal.pone.0226578 (2019).

61 Vilaplana, L. & Marco, M. P. Phenazines as potential biomarkers of

Pseudomonas aeruginosa infections: synthesis regulation, pathogenesis and

analytical methods for their detection. *Anal Bioanal Chem* **412**, 5897-5912,

doi:10.1007/s00216-020-02696-4 (2020).

62 Schalk, I. J. & Guillon, L. Pyoverdine biosynthesis and secretion in Pseudomonas

aeruginosa: implications for metal homeostasis: Pyoverdine biosynthesis.

Environmental Microbiology **15**, 1661-1673, doi:10.1111/1462-2920.12013

(2013).

63 Gasser, V., Guillon, L., Cunrath, O. & Schalk, I. J. Cellular organization of

siderophore biosynthesis in Pseudomonas aeruginosa: Evidence for

siderosomes. *Journal of Inorganic Biochemistry* **148**, 27-34,

doi:10.1016/j.jinorgbio.2015.01.017 (2015).

64 Cox, C. D., Rinehart, K. L., Moore, M. L. & Cook, J. C. Pyochelin: novel structure

of an iron-chelating growth promoter for Pseudomonas aeruginosa. *PNAS* **78**,

4256-4260, doi:10.1073/pnas.78.7.4256 (1981).

65 Wang, Y. *et al.* Phenazine-1-Carboxylic Acid Promotes Bacterial Biofilm

Development via Ferrous Iron Acquisition. *J Bacteriol* **193**, 3606-3617,

doi:10.1128/JB.00396-11 (2011).

66   Korgaonkar, A. K. & Whiteley, M. Pseudomonas aeruginosa enhances

production of an antimicrobial in response to N-acetylglucosamine and

peptidoglycan. *J Bacteriol* **193**, 909-917, doi:10.1128/JB.01175-10 (2011).

67   Lepine, F., Milot, S., Deziel, E., He, J. & Rahme, L. G. Electrospray/mass

spectrometric identification and analysis of 4-hydroxy-2-alkylquinolines (HAQs)

produced by Pseudomonas aeruginosa. *J Am Soc Mass Spectrom* **15**, 862-869,

doi:10.1016/j.jasms.2004.02.012 (2004).

68   Deziel, E. *et al.* Analysis of Pseudomonas aeruginosa 4-hydroxy-2-

alkylquinolines (HAQs) reveals a role for 4-hydroxy-2-heptylquinoline in cell-to-

cell communication. *Proc Natl Acad Sci U S A* **101**, 1339-1344,

doi:10.1073/pnas.0307694100 (2004).

69   Pesci, E. C. *et al.* Quinolone signaling in the cell-to-cell communication system of

Pseudomonas aeruginosa. *Proc Natl Acad Sci U S A* **96**, 11229-11234,

doi:10.1073/pnas.96.20.11229 (1999).

70   Machan, Z. A., Taylor, G. W., Pitt, T. L., Cole, P. J. & Wilson, R. 2-Heptyl-4-

hydroxyquinoline N-oxide, an antistaphylococcal agent produced by

Pseudomonas aeruginosa. *J Antimicrob Chemother* **30**, 615-623,

doi:10.1093/jac/30.5.615 (1992).

71   Van Ark, G. & Berden, J. A. Binding of HQNO to beef-heart sub-mitochondrial

particles. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **459**, 119-137,

doi:10.1016/0005-2728(77)90014-7 (1977).

72   Hazan, R. *et al.* Auto Poisoning of the Respiratory Chain by a Quorum-Sensing-

Regulated Molecule Favors Biofilm Formation and Antibiotic Tolerance. *Current*

Biology **26**, 195-206, doi:10.1016/j.cub.2015.11.056 (2016).

73   Rampioni, G. *et al.* Unravelling the Genome-Wide Contributions of Specific 2-

Alkyl-4-Quinolones and PqsE to Quorum Sensing in Pseudomonas aeruginosa.

PLOS Pathogens **12**, e1006029, doi:10.1371/journal.ppat.1006029 (2016).

74   McKnight, S. L., Iglewski, B. H. & Pesci, E. C. The Pseudomonas Quinolone

Signal Regulates rhl Quorum Sensing in Pseudomonas aeruginosa. *J Bacteriol*

182, 2702-2708, doi:10.1128/JB.182.10.2702-2708.2000 (2000).

75   Guo, Q. *et al.* PqsR-dependent and PqsR-independent regulation of motility and

biofilm formation by PQS in Pseudomonas aeruginosa PAO1: Regulation of

motility and biofilm by PQS. *Journal of Basic Microbiology* **54**, 633-643,

doi:10.1002/jobm.201300091 (2014).

76   Lin, J., Cheng, J., Wang, Y. & Shen, X. The Pseudomonas Quinolone Signal

(PQS): Not Just for Quorum Sensing Anymore. *Front Cell Infect Microbiol* **8**, 230,

doi:10.3389/fcimb.2018.00230 (2018).

77   Gallagher, L. A., McKnight, S. L., Kuznetsova, M. S., Pesci, E. C. & Manoil, C.

Functions required for extracellular quinolone signaling by Pseudomonas

aeruginosa. *J Bacteriol* **184**, 6472-6480, doi:10.1128/jb.184.23.6472-6480.2002

(2002).

78   Diggle, S. P., Cornelis, P., Williams, P. & Camara, M. 4-quinolone signalling in

Pseudomonas aeruginosa: old molecules, new perspectives. *Int J Med Microbiol*

296, 83-91, doi:10.1016/j.ijmm.2006.01.038 (2006).

79   Hoffman, L. R. *et al.* Selection for Staphylococcus aureus small-colony variants

due to growth in the presence of Pseudomonas aeruginosa. *Proc Natl Acad Sci*

U S A **103**, 19890-19895, doi:10.1073/pnas.0606756104 (2006).

80   Lightbown, J. W. An antagonist of streptomycin and dihydrostreptomycin

produced by Pseudomonas aeruginosa. *J Gen Microbiol* **11**, 477-492,

doi:10.1099/00221287-11-3-477 (1954).

81   Orazi, G. & O'Toole, G. A. Pseudomonas aeruginosa Alters Staphylococcus

aureus Sensitivity to Vancomycin in a Biofilm Model of Cystic Fibrosis Infection.

mBio **8**, doi:10.1128/mBio.00873-17 (2017).

82   Bortolotti, P. *et al.* Tryptophan catabolism in Pseudomonas aeruginosa and

potential for inter-kingdom relationship. *BMC Microbiol* **16**, 137,

doi:10.1186/s12866-016-0756-x (2016).

83   Palmer, K. L., Mashburn, L. M., Singh, P. K. & Whiteley, M. Cystic fibrosis

sputum supports growth and cues key aspects of Pseudomonas aeruginosa

physiology. *J Bacteriol* **187**, 5267-5277, doi:10.1128/JB.187.15.5267-5277.2005

(2005).

84   Quinn, R. A. *et al.* Neutrophilic proteolysis in the cystic fibrosis lung correlates

with a pathogenic microbiome. *Microbiome* **7**, 23, doi:10.1186/s40168-019-0636-

3 (2019).

85   Raghuvanshi, R. *et al.* High-Resolution Longitudinal Dynamics of the Cystic

Fibrosis Sputum Microbiome and Metabolome through Antibiotic Therapy.

mSystems **5**, doi:10.1128/mSystems.00292-20 (2020).

86   Melnik, A. V. *et al.* Molecular and Microbial Microenvironments in Chronically

Diseased Lungs Associated with Cystic Fibrosis. *mSystems* **4**,

doi:10.1128/mSystems.00375-19 (2019).

87   Garg, N. *et al.* Three-Dimensional Microbiome and Metabolome Cartography of a

Diseased Human Lung. *Cell Host Microbe* **22**, 705-716 e704,

doi:10.1016/j.chom.2017.10.001 (2017).

88   Davey, M. E., Caiazza, N. C. & O'Toole, G. A. Rhamnolipid Surfactant Production

Affects Biofilm Architecture in Pseudomonas aeruginosa PAO1. *J Bacteriol* **185**,

1027-1036, doi:10.1128/JB.185.3.1027-1036.2003 (2003).

89   Glick, R. *et al.* Increase in Rhamnolipid Synthesis under Iron-Limiting Conditions

Influences Surface Motility and Biofilm Formation in Pseudomonas aeruginosa. *J*

Bacteriol **192**, 2973-2980, doi:10.1128/JB.01601-09 (2010).

90   McClure, C. D. & Schiller, N. L. Inhibition of Macrophage Phagocytosis by

Pseudomonas aeruginosa Rhamnolipids In Vitro and In Vivo. *Current*

Microbiology **33**, 109-117, doi:10.1007/s002849900084 (1996).

91   Zulianello, L. *et al.* Rhamnolipids Are Virulence Factors That Promote Early

Infiltration of Primary Human Airway Epithelia by Pseudomonas aeruginosa.

Infect. Immun. **74**, 3134-3147, doi:10.1128/IAI.01772-05 (2006).

92   Briard, B. *et al.* Dirhamnolipids secreted from Pseudomonas aeruginosa modify

anjpegungal susceptibility of Aspergillus fumigatus by inhibiting β1,3 glucan

synthase activity. *The ISME Journal* **11**, 1578-1591, doi:10.1038/ismej.2017.32

(2017).

93   Calfee, M. W., Shelton, J. G., McCubrey, J. A. & Pesci, E. C. Solubility and

bioactivity of the Pseudomonas quinolone signal are increased by a

Pseudomonas aeruginosa-produced surfactant. *Infect Immun* **73**, 878-882,

doi:10.1128/IAI.73.2.878-882.2005 (2005).

94   Caiazza, N. C., Shanks, R. M. Q. & O'Toole, G. A. Rhamnolipids Modulate

Swarming Motility Patterns of Pseudomonas aeruginosa. *J Bacteriol* **187**, 7351-

7361, doi:10.1128/JB.187.21.7351-7361.2005 (2005).

95   Meyer, J. M., Neely, A., Stintzi, A., Georges, C. & Holder, I. A. Pyoverdin is

essential for virulence of Pseudomonas aeruginosa. *Infect. Immun.* **64**, 518-523,

doi:10.1128/IAI.64.2.518-523.1996 (1996).

96   Takase, H., Nitanai, H., Hoshino, K. & Otani, T. Impact of Siderophore

Production onPseudomonas aeruginosa Infections in Immunosuppressed Mice.

Infect. Immun. **68**, 1834-1839, doi:10.1128/IAI.68.4.1834-1839.2000 (2000).

97   Sriyosachati, S. & Cox, C. D. Siderophore-mediated iron acquisition from

transferrin by Pseudomonas aeruginosa. *Infect. Immun.* **52**, 885-891,

doi:10.1128/IAI.52.3.885-891.1986 (1986).

98   Meyer, J. M. Pyoverdines: pigments, siderophores and potential taxonomic

markers of fluorescent Pseudomonas species. *Arch Microbiol* **174**, 135-142,

doi:10.1007/s002030000188 (2000).

99   Ravel, J. & Cornelis, P. Genomics of pyoverdine-mediated iron uptake in

pseudomonads. *Trends Microbiol* **11**, 195-200, doi:10.1016/s0966-

842x(03)00076-3 (2003).

100   Visca, P., Imperi, F. & Lamont, I. L. Pyoverdine siderophores: from biogenesis to

887   biosignificance. *Trends Microbiol* **15**, 22-30, doi:10.1016/j.tim.2006.11.004

888   (2007).

889   101   Cornelis, P. Iron uptake and metabolism in pseudomonads. *Applied Microbiology*

890   *and Biotechnology* **86**, 1637-1645, doi:10.1007/s00253-010-2550-2 (2010).

891   102   Albrechtgary, A. M., Blanc, S., Rochel, N., Ocaktan, A. Z. & Abdallah, M. A.

892   Bacterial Iron Transport - Coordination Properties of Pyoverdin Paa, a Peptidic

893   Siderophore of Pseudomonas-Aeruginosa. *Inorg Chem* **33**, 6391-6402, doi:DOI

894   10.1021/ic00104a059 (1994).

895   103   Cornelis, P. & Dingemans, J. Pseudomonas aeruginosa adapts its iron uptake

896   strategies in function of the type of infections. *Front Cell Infect Microbiol* **3**, 75,

897   doi:10.3389/fcimb.2013.00075 (2013).

898   104   Dumas, Z., Ross-Gillespie, A. & Kummerli, R. Switching between apparently

899   redundant iron-uptake mechanisms benefits bacteria in changeable

900   environments. *Proc Biol Sci* **280**, 20131055, doi:10.1098/rspb.2013.1055 (2013).

901   105   Kang, D. *et al.* Pyoverdine-Dependent Virulence of Pseudomonas aeruginosa

902   Isolates From Cystic Fibrosis Patients. *Front Microbiol* **10**, 2048,

903   doi:10.3389/fmicb.2019.02048 (2019).

904   106   Wagner, C. E., Wheeler, K. M. & Ribbeck, K. Mucins and Their Role in Shaping

905   the Functions of Mucus Barriers. *Annu Rev Cell Dev Biol* **34**, 189-215,

906   doi:10.1146/annurev-cellbio-100617-062818 (2018).

907   107   Pinzon Martin, S., Seeberger, P. H. & Varon Silva, D. Mucins and Pathogenic

908   Mucin-Like Molecules Are Immunomodulators During Infection and Targets for

Diagnostics and Vaccines. *Front Chem* **7**, 710, doi:10.3389/fchem.2019.00710 (2019).

Morrison, C. B., Markovetz, M. R. & Ehre, C. Mucus, mucins, and cystic fibrosis. *Pediatr Pulmonol* **54 Suppl 3**, S84-S96, doi:10.1002/ppul.24530 (2019).

Rose, M. C. & Voynow, J. A. Respiratory tract mucin genes and mucin glycoproteins in health and disease. *Physiol Rev* **86**, 245-278, doi:10.1152/physrev.00010.2005 (2006).

Henke, M. O., John, G., Germann, M., Lindemann, H. & Rubin, B. K. MUC5AC and MUC5B mucins increase in cystic fibrosis airway secretions during pulmonary exacerbation. *Am J Respir Crit Care Med* **175**, 816-821, doi:10.1164/rccm.200607-1011OC (2007).

Henderson, A. G. *et al.* Cystic fibrosis airway secretions exhibit mucin hyperconcentration and increased osmotic pressure. *J Clin Invest* **124**, 3047-3060, doi:10.1172/JCI73469 (2014).

Hoffman, C. L., Lalsiamthara, J. & Aballay, A. Host Mucin Is Exploited by <i>Pseudomonas aeruginosa</i> To Provide Monosaccharides Required for a Successful Infection. *mBio* **11**, doi:10.1128/mBio.00060-20 (2020).

Flynn, J. M., Niccum, D., Dunitz, J. M. & Hunter, R. C. Evidence and Role for Bacterial Mucin Degradation in Cystic Fibrosis Airway Disease. *PLOS Pathogens* **12**, e1005846, doi:10.1371/journal.ppat.1005846 (2016).

Darch, S. E. *et al.* Phage Inhibit Pathogen Dissemination by Targeting Bacterial Migrants in a Chronic Infection Model. *mBio* **8**, doi:10.1128/mBio.00240-17 (2017).

115 Secor, P. R., Michaels, L. A., Ratjen, A., Jennings, L. K. & Singh, P. K.

Entropically driven aggregation of bacteria by host polymers promotes antibiotic

tolerance in <i>Pseudomonas aeruginosa</i>. *Proceedings of the National*

Academy of Sciences **115**, 10780-10785, doi:10.1073/pnas.1806005115 (2018).

116 Co, J. Y. *et al.* Mucins trigger dispersal of Pseudomonas aeruginosa biofilms. *npj*

Biofilms and Microbiomes **4**, doi:10.1038/s41522-018-0067-0 (2018).

117 Wheeler, K. M. *et al.* Mucin glycans attenuate the virulence of Pseudomonas

aeruginosa in infection. *Nature Microbiology* **4**, 2146-2154, doi:10.1038/s41564-

019-0581-8 (2019).

118 Schomig, V. J. *et al.* An optimized purification process for porcine gastric mucin

with preservation of its native functional properties. *Rsc Adv* **6**, 44932-44943,

doi:10.1039/c6ra07424c (2016).

119 Lieleg, O., Lieleg, C., Bloom, J., Buck, C. B. & Ribbeck, K. Mucin biopolymers as

broad-spectrum antiviral agents. *Biomacromolecules* **13**, 1724-1732,

doi:10.1021/bm3001292 (2012).

120 Kocevar-Nared, J., Kristl, J. & Smid-Korbar, J. Comparative rheological

investigation of crude gastric mucin and natural gastric mucus. *Biomaterials* **18**,

677-681, doi:10.1016/s0142-9612(96)00180-9 (1997).

121 Kilbourn, J. P. Composition of sputum from patients with cystic fibrosis. *Current*

Microbiology **11**, 19-22, doi:10.1007/BF01567570 (1984).

122 Hunter, R. C. *et al.* Ferrous Iron Is a Significant Component of Bioavailable Iron

in Cystic Fibrosis Airways. *mBio* **4**, doi:10.1128/mBio.00557-13 (2013).

123   Reid, D. W., Carroll, V., O'May, C., Champion, A. & Kirov, S. M. Increased airway

iron as a potential factor in the persistence of Pseudomonas aeruginosa infection

in cystic fibrosis. *Eur Respir J* **30**, 286-292, doi:10.1183/09031936.00154006

(2007).

124   Ghio, A. J. *et al.* Iron accumulates in the lavage and explanted lungs of cystic

fibrosis patients. *J Cyst Fibros* **12**, 390-398, doi:10.1016/j.jcf.2012.10.010 (2013).

125   Nguyen, A. T., Jones, J. W., Ruge, M. A., Kane, M. A. & Oglesby-Sherrouse, A.

G. Iron Depletion Enhances Production of Antimicrobials by Pseudomonas

aeruginosa. *J Bacteriol* **197**, 2265-2275, doi:10.1128/JB.00072-15 (2015).

126   Reinhart, A. A. & Oglesby-Sherrouse, A. G. Regulation of Pseudomonas

aeruginosa Virulence by Distinct Iron Sources. *Genes (Basel)* **7**,

doi:10.3390/genes7120126 (2016).

127   Palmer, G. C., Jorth, P. A. & Whiteley, M. The role of two Pseudomonas

aeruginosa anthranilate synthases in tryptophan and quorum signal production.

Microbiology (Reading) **159**, 959-969, doi:10.1099/mic.0.063065-0 (2013).

128   Yang, J., Eiserich, J. P., Cross, C. E., Morrissey, B. M. & Hammock, B. D.

Metabolomic profiling of regulatory lipid mediators in sputum from adult cystic

fibrosis patients. *Free Radic Biol Med* **53**, 160-171,

doi:10.1016/j.freeradbiomed.2012.05.001 (2012).

129   Scholte, B. J. *et al.* Oxidative stress and abnormal bioactive lipids in early cystic

fibrosis lung disease. *J Cyst Fibros* **18**, 781-789, doi:10.1016/j.jcf.2019.04.011

(2019).

130   Sturmer, R., Harder, S., Schluter, H. & Hoffmann, W. Commercial Porcine

Gastric Mucin Preparations, also Used as Artificial Saliva, are a Rich Source for

the Lectin TFF2: In Vitro Binding Studies. *Chembiochem* **19**, 2598-2608,

doi:10.1002/cbic.201800622 (2018).

131   Jacobs, M. A. *et al.* Comprehensive transposon mutant library of Pseudomonas

aeruginosa. *Proc Natl Acad Sci U S A* **100**, 14339-14344,

doi:10.1073/pnas.2036282100 (2003).

132   Bligh, E. G. & Dyer, W. J. A rapid method of total lipid extraction and purification.

Can J Biochem Physiol **37**, 911-917, doi:10.1139/o59-099 (1959).

133   Pluskal, T., Castillo, S., Villar-Briones, A. & Oresic, M. MZmine 2: Modular

framework for processing, visualizing, and analyzing mass spectrometry-based

molecular profile data. *Bmc Bioinformatics* **11**, doi:Artn 395 10.1186/1471-2105-

11-395 (2010).

134   Shannon, P. *et al.* Cytoscape: A software environment for integrated models of

biomolecular interaction networks. *Genome Res* **13**, 2498-2504,

doi:10.1101/gr.1239303 (2003).

135   Sumner, L. W. *et al.* Proposed minimum reporting standards for chemical

analysis. *Metabolomics* **3**, 211-221, doi:10.1007/s11306-007-0082-2 (2007).

136   Chong, J. *et al.* MetaboAnalyst 4.0: towards more transparent and integrative

metabolomics analysis. *Nucleic Acids Res* **46**, W486-W494,

doi:10.1093/nar/gky310 (2018).

137   Chong, J. & Xia, J. MetaboAnalystR: an R package for flexible and reproducible

998   analysis of metabolomics data. *Bioinformatics* **34**, 4313-4314,

999   doi:10.1093/bioinformatics/bty528 (2018).

1000   138   Chong, J., Yamamoto, M. & Xia, J. MetaboAnalystR 2.0: From Raw Spectra to

1001   Biological Insights. *Metabolites* **9**, doi:10.3390/metabo9030057 (2019).

1002   139   Pang, Z., Chong, J., Li, S. & Xia, J. MetaboAnalystR 3.0: Toward an Optimized

1003   Workflow for Global Metabolomics. *Metabolites* **10**, 186,

1004   doi:10.3390/metabo10050186 (2020).

1005

1006   **FIGURES AND LEGENDS**

1007

1008   **Figure 1. Publication timeline of ASM formulations.** Each formulation is plotted at

1009   the year it was first published. Open circles indicate media derived from the Soothill

1010   ASM recipe. Closed circles indicate SCFM ASM lineage. Nutrient alterations between

formulations are indicated by arrows. Compared to preceding formulation: + = added, -

= removed, ↑ = increased, ↓ = decreased, → = different source. AA = amino acids, BSA

= bovine serum albumin, Cas AA = casamino acids, MEM AA = MEM amino acids,

DOPC = 1,2-dioleoyl-sn-glycero-3-phosphocholine, GlcNAc = *N*-acetyl glucosamine,

PABA = 4-aminobenzoic acid, NAD+ = nicotinamide adenine dinucleotide, ADE =

adenine, GUA = guanine, XAN = xanthine, HX = hypoxanthine. Recipes for ASM

formulations are in detailed in Dataset S1 ASM formulations.

Figure 1. *P. aeruginosa* growth and global metabolism in ASM. (A) Phenotypes, (B)

growth of representative replicates, (C) principal component analysis (PCA) score plots

of metabolic profiles, (D) One-way ANOVA plot of metabolic features, and (E) pairwise

number of statistically different metabolic features of PAO1 cultures in ASM

formulations. Samples as labeled and colored as indicated. n=2-3 for CFU/mL growth

assessment. n=12 for metabolite profiling. Statistically significant metabolic features (p

< 0.05) are colored red in the one-way ANOVA plot.

Figure 3. Phenazine production by PAO1 in ASM. (A) Phenazine molecular family

visualized with feature based molecular networking, colored as in (B). (B) Total

normalized area under the curve (AUC) for the phenazine molecular family in each ASM

formulation. Box plots represent the 25-75th percentile, with a line at the median. Error

bars indicate the minimum to maximum. Individual samples values shown (n=12 per

1034 ASM formulation)*.* (C) Profile of total phenazine production*,* colored as in (D). (D)

1035 Proportion of total phenazines*.* 1-HP: 1-hydroxyphenazine; PYO: pyocyanin; PCN:

1036 phenazine-1-carboxamide; PCA: phenazine-1-carboxylic acid.

1037

1038

Figure 4. Quinolone production by PAO1 in ASM. (A) Quinolone molecular family

visualized with feature based molecular networking, colored as indicated in (B). Nodes

in the network are annotated by alkyl chain length and unsaturation, followed by

structural sub-class assignment. HQ, PQS, and QNO denotes structural relatedness to

HHQ, PQS, and HQNO, respectively. Features representing metabolites that could not

be distinguished based upon MS/MS fragmentation patterns are indicated and referred

to as 'mixed' in (C). Annotations with –OH designation indicate hydroxylation at

unknown position within the molecule and are referred to as 'other' in (C). (B) Total

normalized area under the curve (AUC) for the quinolone molecular family in each ASM

formulation. Box plots represent the 25-75th percentile, with a line at the median. Error

bars indicate the minimum to maximum. Individual samples values shown (n=12 per

ASM formulation). (C) Proportion of total quinolones based upon structural sub-class.

Quinolone structural sub-classes are colored as indicated. HHQ: 2-heptyl-4-quinolone;

PQS: 2-heptyl-3,4-dihydroxyquinoline; HQNO: 2- heptyl-4-hydroxyquinoline *N*-oxide.

1053

1054

Figure 5. Rhamnolipid production by PAO1 in ASM. (A) Rhamnolipid molecular

family (sodiated) visualized with feature based molecular networking, colored as

indicated in (B). Nodes in the network are annotated by rhamnose units (Rha) followed

by fatty acid chain lengths. Only sodium adducts are shown. (B) Total normalized area

under the curve (AUC) for the quinolone molecular family in each ASM formulation. Box

plots represent the 25-75th percentile, with a line at the median. Error bars indicate the

minimum to maximum. Individual samples values shown (n=12 per ASM formulation).

(C) Proportion of total rhamnolipid production, colored as indicated.

1063

1064

Figure 6. Siderophore production by PAO1 in ASM. (A) Pyochelin and pyoverdine

molecular families visualized with feature based molecular networking. SCFM1 cultures

are blue. Nodes in the network are annotated by adducts. (B) Total normalized area

under the curve (AUC) for pyochelin in each ASM formulation (n=12 per ASM

formulation). (C) Total normalized AUC for pyoverdine molecular family in each ASM

formulation (n=12 per ASM formulation). For (C) and (D), box plots represent the 25-

75th percentile, with a line at the median. Error bars indicate the minimum to maximum.

1072

1073

Figure 7. Effect of 0.5% PGM on production of small molecule virulence factors by

PAO1. (A) Left: Total normalized AUC for phenazine molecular family, colored as

indicated. Right: Normalized AUC for phenazine molecular family members. (B) Left:

Total normalized AUC for phenazine molecular family, colored as indicated and only

most abundant quinolones included. Right: Normalized AUC for quinolone molecular

family members with largest alterations in production. (C) Left: Total normalized AUC for

rhamnolipid molecular family, colored as indicated. Right: Normalized AUC for

rhamnolipid molecular family members with largest alterations in production. (D)

Normalized AUC for pyochelin and pyoverdine molecular family. (E) Concentration of

iron in 0.5% PGM and vehicle control (MOPS). Dashed line indicates the concentration

of iron included in the SCFM1 ASM formulation. For (A)-(D), n=4 biological replicates.

For all stacked bar charts, bars represent the mean and error bars indicate standard

deviation of the mean. All box plots represent the 25-75th percentile, with a line at the

median, and error bars indicating the minimum and maximum.

1088

1089

1090 **Figure 8. Contaminants in Commercial PGM.** (A) Metals detected in PGM. Line

1091 indicates the median of six technical replicates. (B) Classical molecular network of

1092 lipophilic extracts of commercial PGM. Colored as indicated.