Versatile mixotrophs employ CO$_2$ fixation as fill-up function to dominate in modern groundwater

Martin Taubert[1,+], Will A. Overholt[1], Beatrix M. Heinze[1], Georgette Azemtsop Matanfack[2,5], Rola Houhou[2,5], Nico Jehmlich[3], Martin von Bergen[3,4], Petra Rösch[2], Jürgen Popp[2,5], Kirsten Küsel[1,6]

[1]Aquatic Geomicrobiology, Institute of Biodiversity, Friedrich Schiller University Jena, Dornburger Str. 159, 07743 Jena, Germany

[2]Institute of Physical Chemistry and Abbe Center of Photonics, Friedrich Schiller University Jena, Helmholtzweg 4, 07743 Jena, Germany

[3]Department of Molecular Systems Biology, Helmholtz Centre for Environmental Research – UFZ, Permoserstrasse 15, 04318 Leipzig, Germany

[4]Institute of Biochemistry, Faculty of Biosciences, Pharmacy and Psychology, University of Leipzig, Brüderstraße 32, 04103 Leipzig, Germany

[5]Leibniz-Institute of Photonic Technology, Albert-Einstein-Straße 9, 07745 Jena, Germany

[6]German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5E, 04103 Leipzig, Germany

[+]Corresponding author: Martin Taubert; Tel.: +49 3641 949459; Fax: +49 3641 949402; E-mail: martin.taubert@uni-jena.de

Keywords: Chemolithoautotrophy, groundwater, [13]CO$_2$ stable isotope probing, genome-resolved metaproteomics, metagenomics, Raman microspectroscopy

Short title: Chemolithoautotrophy in shallow groundwater

Abstract

The processes for organic carbon inputs in ecosystems without photosynthetic primary production are poorly constrained, by data almost entirely derived from metagenomics studies. In modern groundwater, such studies reported a high abundance of putative chemolithoautotrophs. Here we quantitatively and temporally tracked the flux of CO_2-derived carbon in a groundwater microbial community stimulated with reduced sulfur compounds, by combining $^{13}CO_2$-stable isotope probing, multi-omics and Raman microspectroscopy. We demonstrate that metabolically flexible mixotrophs drove the recycling of organic carbon and supplemented their carbon requirements by chemolithoautotrophy alongside the preferential uptake of available organic compounds. These mixotrophs, including *Hydrogenophaga*, *Polaromonas* and *Dechloromonas*, dominated the microbial community, being five times more abundant than strict autotrophs based on metagenomics coverage. Due to their activity, 43% of total microbial carbon was replaced by ^{13}C within 21 days, increasing to 80% after 70 days. Sixteen of 31 taxa investigated, including both autotrophs and heterotrophs, were supported by sulfur oxidation pathways, highlighting its importance as an energy source in an environment with little available organic carbon. The versatile strategy employed by these organisms might also be successful in other oligotrophic habitats, like the upper ocean or boreal lakes, explaining the high abundance of mixotrophs found there.

From soils to deep-sea sediments, the majority of cells on Earth have to survive in environments

without photosynthetic primary production [1]. To understand the global carbon cycle, it is critical to

know to what extent these cells depend on allochthonous carbon for a heterotrophic lifestyle, or on

chemolithoautotrophy. The determination of CO_2 fixation rates and turnover is challenging in these

habitats, but metagenomics studies recovered thousands of metagenome-assembled genomes

(MAGs) from the organisms present to shed light on their metabolic capabilities [2-5].

Modern groundwater, having entered the subsurface less than 50 years ago [6], represents a

transitionary ecosystem, connecting surface habitats dominated by photosynthetic carbon with the

deep subsurface that lacks this energy source [7-9]. There, metagenomics studies have unveiled the

presence of a variety of organisms with the metabolic potential for CO_2 fixation and for the utilization

of inorganic electron donors [4, 10-13]. Such organisms with the ability to grow

chemolithoautotrophically constitute between 12 and 47% of the microbial communities in

groundwater [14-17]. These discoveries shake the paradigm that modern groundwater is primarily

fueled by surface input of organic material and dominated by heterotrophs. We thus hypothesize

that modern groundwater ecosystems are dominated by chemolithoautotrophic primary production.

For validating this hypothesis, we established Stable Isotope Cluster Analysis (SIsCA) to provide a

time-resolved and quantitative assessment of the flow of CO_2-derived carbon through the

groundwater microbial food web. By combining stable isotope probing (SIP) [18, 19] with genome-

resolved metaproteomics, we leveraged the high sensitivity of Orbitrap mass spectrometry in SIP-

metaproteomics to acquire highly accurate quantitative data on [13]C incorporation [20, 21]. The novel

SIsCA method then employs a dimensionality reduction approach to integrate the molecule-specific

temporal dynamics in the acquired isotopologue patterns from this [13]C-SIP time-series experiment, to

discern trophic fluxes between individual members of a microbial community.

We employed this approach to unravel the role of chemolithoautotrophy for the groundwater

microbiome by amending groundwater microcosms with [13]CO_2. In modern groundwater, different

inorganic electron donors like reduced nitrogen, iron and sulfur can fuel the chemolithoautotrophic

primary production, while hydrogen is rather of relevance in the deeper subsurface [9, 22-24].

Reduced sulfur compounds released from aquifer rocks by weathering create an energy source

independent of surface inputs [25-27]. Organisms with the genetic potential to oxidize reduced sulfur

compounds are widespread in groundwater, but little is known about their lifestyle [12, 13, 28].

Hence, we stimulated sulfur oxidizers in groundwater microcosms by the addition of thiosulfate,

which is commonly found in the environment [29]. Under these conditions favoring lithotrophic

growth, we expected chemolithoautotrophy to be the main source of organic carbon, and a

unidirectional carbon flux from autotrophs to heterotrophs in the groundwater microbiome. By

mapping the quantitative information derived from SIsCA to MAGs, we were able to characterize

carbon utilization and trophic interactions of the active autotrophic and heterotrophic key players in

the groundwater microbiome over a period of 70 days. This accurate resolution of the carbon fluxes

through a microbial community and the determination of taxon-specific microbial activities have the

power to provide novel insights into the metabolic strategies and trophic interactions of microbes.

Materials and Methods

Groundwater sampling & setup of groundwater microcosms

Groundwater was obtained in June 2018 from the Hainich Critical Zone Exploratory (CZE) well H41 (51.1150842N, 10.4479713E), accessing an aquifer assemblage in 48 m depth in a trochite limestone stratum. Groundwater from this well is oxic with average dissolved oxygen concentrations of 5.0 ± 1.5 mg L^{-1}, pH 7.2, < 0.1 mg L^{-1} ammonium, 1.9 ± 1.5 mg L^{-1} dissolved organic carbon, and 70.8 ± 12.7 mg L^{-1} total inorganic carbon [27, 30]. The recharge area of this aquifer assemblage is located in a beech forest (*Fagus sylvatica*). In total, 120 L of groundwater was sampled using a submersible pump (Grundfos MP1, Grundfos, Bjerringbro, Denmark). To collect biomass from the groundwater, 5 L each were filtered through twenty 0.2-µm Supor filters (Pall Corporation, Port Washington, NY, USA). To replace the natural background of inorganic carbon in the groundwater by defined concentrations of ^{12}C or ^{13}C, two times 3 L of filtered groundwater were acidified to pH 4 in 5-L-bottles to remove the bicarbonate present. Following that, ^{12}C- or ^{13}C-bicarbonate was dissolved in the groundwater to a final concentration of 400 mg L^{-1}, corresponding to 79 mg C L^{-1}, close to the *in situ* concentration. The pH of the groundwater samples was adjusted to 7.2 by addition of ^{12}C- or ^{13}C-CO_2 to the headspace of the bottles.

For the ^{13}C-SIP experiment, eighteen microcosms were set up by transferring one filter each into a 500-ml-bottle with 300 ml of treated groundwater as described, 9 microcosms with water containing ^{12}C-bicarbonate, 9 microcosms with water containing ^{13}C-bicarbonate. Additionally, two microcosms were set up by transferring one filter each into a 1-L-bottle with 350 ml untreated groundwater. One of these bottles was supplemented with 150 ml D_2O (final concentration 30% v:v), the second bottle was supplemented with 150 ml H_2O. To all microcosms, sodium thiosulfate was added to a final concentration of 2.5 mM, and ammonium chloride was added to a final concentration of 15 µM. All microcosms were incubated in the dark at 15 °C with shaking at 100 rpm.

Hydrochemical analyses

During the incubation, in the 18 microcosms supplemented with [12]C- or [13]C-bicarbonate, concentrations of oxygen, thiosulfate and sulfate were monitored at regular intervals. Oxygen concentrations were determined using the contactless fiber-optic oxygen sensor system Fibox 4 trace with SP-PSt3-SA23-D5-YOP-US dots (PreSens Precision Sensing GmbH, Regensburg, Germany). Measurements were taken in three [12]C microcosms and three [13]C microcosms every two days for the first three weeks, and every week thereafter. Thiosulfate concentration was determined in a colorimetric titration assay using iodine [31]. Samples from all microcosms were measured every 4 to 7 days. For each measurement, 1 ml of sample was mixed with 2 mg potassium iodide, then 10 µl of zinc iodide-starch-solution (4 g L^{-1} starch, 20 g L^{-1} zinc chloride and 2 g L^{-1} zinc iodide) and 10 µl of 17% (v:v) phosphoric acid were added. Titration was performed using 0.005 N iodine solution in steps of 5 µl until formation of a faint blue color. The concentration of thiosulfate ($c_{thiosulfate}$ in mg L^{-1}) was calculated according to equation (I), where V_{iodine} is the volume of iodine solution added and V_{sample} is the sample volume:

$$c_{thiosulfate} = \frac{V_{iodine} \times 561}{V_{sample}}$$
(I)

The concentration of sulfate was determined using a turbidimetric assay [32] in samples from all microcosms every 4 to 7 days. For each measurement, 1 ml of sample, standard (50 µM to 1000 µM potassium sulfate) or blank (dH$_2$O) was mixed with 0.4 ml 0.5 M HCl and 0.2 ml BaCl$_2$-gelatin reagent (0.5 g gelatin and 8 g BaCl$_2$ in 200 ml dH$_2$O). After incubation for 1 h in the dark, the absorbance at 420 nm was measured in a DR3900 spectrophotometer (HACH, Düsseldorf, Germany). No differences in thiosulfate or oxygen consumption or sulfate formation were observed between the [12]C- and [13]C-bicarbonate supplemented microcosms.

Detection of cellular activity by Raman microspectroscopy

The two microcosms supplemented with D$_2$O or H$_2$O were sampled regularly during the first 7 weeks of incubation for single cell Raman analysis, to quantify the incorporation of deuterium in the

biomolecules of active cells as carbon-deuterium-(C-D)-bonds. For Raman spectroscopic analysis, 1

ml sample was filtered through a 0.5-μm-filter and subsequently washed three times by centrifuging

at 10,000g for 2 min, discarding the supernatant and re-suspending the pellet in 1 ml of sterile

ddH$_2$O. Afterwards, the pellet was re-suspended in 50 μl ddH$_2$O, and 10 μl of the final bacterial

suspension were placed on a nickel foil (Raman substrate) and air-dried at room temperature.

Microbial cells were located by dark field microscopy. The measurements were performed with a

Raman microscope (BioParticleExplorer 0.5, rap.ID Particle Systems GmbH) with an excitation

wavelength of 532 nm from a solid-state frequency-doubled Nd:YAG module (Cobolt Samba 25 mW)

and a laser power of 13 mW at the sample. The laser light was focused with an x100 objective

(Olympus MPLFLN 100xBD) and with a spot size <1 μm laterally. The 180°-backscattered light was

diffracted by a single-stage monochromator (Horiba Jobin Yvon HE 532) with a 920 line mm^{-1} grating

and then registered with a thermoelectrically cooled CCD camera (Andor DV401-BV) resulting in a

spectral resolution of about 8 cm^{-1}. An integration time of 5 s was used per Raman spectrum (-57 to

3203 cm^{-1}).

Pre-processing and analysis of Raman data

The data pre-processing and statistical analysis were carried out in the software GNU R [33]. First,

cosmic spikes were detected and removed from the spectra [34]. Subsequently, a wavenumber

calibration was applied using 4-acetamidophenol standard spectra [35], while an intensity calibration

was performed using the SRM2242 standard material [36, 37]. The fluorescence contribution was

removed from the spectra using the asymmetric least-squares (ALS) baseline correction method [38].

Finally, the spectra were vector normalized and subjected to dimensionality reduction using principal

component analysis (PCA). Five principal components were used to build a linear discriminant

analysis (LDA) classification model to differentiate between deuterium labeled and non-labeled

bacterial cells. The deuterium uptake was expressed as the C-D ratio A(C-D) / [A(C-D) + A(C-H)], which

was calculated by integrating the areas of the C-H (2800 - 3100 cm^{-1}) and C-D (2040 - 2300 cm^{-1})

stretching vibration bands. The detection and quantification of deuterium incorporation from D$_2$O in

microbial cells was used to assess metabolic activity and decide on time points for sampling of

microcosms.

Sampling, DNA and protein extraction

After 21, 43 and 70 days of incubation, microbial biomass from the microcosms was recovered by

filtration of the water phase through 0.2-μm Supor filters (Pall Corporation). The filters used for

biomass enrichment before incubation were retrieved from the microcosms as well, and combined

with the filters used for the water phase. Combined DNA and protein extraction was performed using

a phenol/chloroform-based protocol as previously described [39]. Details on 16S rRNA gene amplicon

sequencing as well as of quantitative SIP of DNA samples can be found in the Supplement.

Metagenomic analysis

To obtain genomes of the key organisms in the microcosms, metagenomic sequencing was

performed on DNA samples from 4 selected [12]C microcosms, one replicate after 21 days of

incubation and 43 days of incubation, and two replicates after 70 days of incubation. These samples

were selected to cover the majority of taxonomic diversity, based on principal component analysis of

the 16S rRNA gene amplicon sequencing data. DNA sizing, quantitation, integrity, and purity were

determined using the Agilent 2100 Bioanalyzer system (Santa Clara, CA, USA). Library preparation

was performed with a NEBNext Ultra II DNA Lib Prep Kit (New England Biolabs, Ipswich, MA, USA) as

recommended by the manufacturer, followed by multiplexed sequencing on one flow cell of an

Illumina NextSeq 500 system (300 cycles) to generate 150-bp paired-end reads.

Raw sequencing data was quality filtered using BBDuk [40] following assembly using metaSPAdes

v3.13.0 [41]. Using contigs with more than 1000 bp, three different binning algorithm were used to

obtain genomic bins: MaxBin 2.0 v2.2.7 [42], MetaBAT 2 v2.12.1 [43], and BinSanity v0.2.7 [44]. Bin

refinement was performed using the MetaWRAP pipeline v1.1.3 [45]. Only bins with more than 50%

completeness and less than 10% contamination were considered. Bins were classified using GTDB-Tk

v0.3.2 [46] and completeness parameters were assessed using CheckM v1.0.12 [47]. Bins from

different samples were dereplicated using FastANI v1.0 [48]. The Prokka pipeline v1.13.3 [49] was

used for calling and functional annotation of gene sequences, and for translation into amino acid

sequences for metaproteomics analysis. Selected bins of key organisms covered by the

metaproteomics analysis were manually refined using Anvi'o v6.1 [50] to obtain the final

metagenome-assembled genomes (MAGs). Normalized coverage values for all MAGs were calculated

by dividing raw coverage values by the relative abundance of *rpoB* genes of each metagenome. The

rpoB gene abundances were determined using ROCker [51]. Table S1 provides an overview of the

curated MAGs referred to in the study.

Metaproteomics analysis

Proteins extracted from the microcosms were first subjected to SDS polyacrylamide gel

electrophoresis, followed by in-gel tryptic cleavage as previously described [39]. After reconstitution

in 0.1% formic acid (v:v), LC-MS/MS analysis was performed on a Q Exactive HF instrument (Thermo

Fisher Scientific, Waltham, MA, USA) equipped with a TriVersa NanoMate source (Advion Ltd., Ithaca,

NY, USA) in LC chip coupling mode. Raw data files were analyzed with Proteome Discoverer

(v1.4.1.14, Thermo Fisher Scientific, Waltham, MA, USA) using the Sequest HT search algorithm.

Amino acid sequences derived from translation of the genes from all contigs of the metagenomes

were used as reference database for protein identification, to allow the assignment of identified

proteins to MAGs. The following parameters were used: The enzyme specificity was set to trypsin,

with two missed cleavage allowed. A 5 ppm peptide ion tolerance and a 0.05 Da MS/MS tolerance

were used. Oxidation (methionine) and carbamidomethylation (cysteine) were selected as

modifications. Peptides were considered identified when they scored a q-value < 1% based on a

decoy database and when their peptide rank was 1. Functional classification of peptides was done

according to the gene annotation by Prokka [49], and taxonomic classification based on the

dereplicated and refined MAGs described above.

Stable Isotope Cluster Analysis to determine carbon utilization

To quantify the incorporation of ^{13}C, peptide identifications from samples of ^{12}C microcosms were mapped to mass spectra of the corresponding ^{13}C-labeled samples, by comparison of expected peptide mass, chromatographic retention time and MS/MS fragmentation pattern. The molecular masses of peptides were calculated based on their amino acid sequence. Isotopologue patterns of labeled peptides were extracted manually from mass spectral data using the Xcalibur Qual Browser (v3.0.63, Thermo Fisher Scientific, Waltham, MA, USA), and the ^{13}C incorporation was calculated as previously described [21]. We expected that microbes fixing CO_2 would exhibit a high (> 90%) ^{13}C RIA, while microbes utilizing organic carbon should feature a lower ^{13}C RIA with an increasing trend over time.

The classical approach of calculating the most probable ^{13}C relative isotope abundance (RIA) of a peptide does not take into account the information contained in isotopologue patterns, which provide detailed information about the carbon utilization of an organism. To include this information in the analysis, we developed Stable Isotope Cluster Analysis (SIsCA). SIsCA was performed using R [33]. Scripts for SIsCA are available on github (https://github.com/m-taubert/SIsCA). In this approach, measured isotopologue patterns were compared with 21 predicted isotopologue patterns with varying ^{13}C RIA in 5% intervals, from 0% to 100% ^{13}C RIA. For each comparison, the coefficient of determination (R^2) was calculated. With this approach, all information from isotopologue patterns is retained, while still data from different peptides is comparable, time series can be integrated, and the dataset can easily be used for downstream statistical analysis. To differentiate lifestyles of different microbes, R^2 values were averaged between samples obtained from replicate microcosms and peptides assigned to the same MAG. The resulting datasets of 21 R^2 values per time point for each MAG were compared using principal component analysis in the software package vegan [52], and clusters of MAGs were defined manually and validated by testing for overlapping confidence intervals.

Generation times of individual taxa were calculated by comparing the relative intensity of unlabeled

and labeled peptide signals in mass spectrometric data as previously described [20]. First, the

number of doubling was calculated according to equation (II), where n is the number of doublings,

I_{12_C} is the signal intensity of the unlabeled peptide and I_{13_C} is the signal intensity of the labeled

peptide.

$$n = \log_2 \frac{I_{12_C} + I_{13_C}}{I_{13_C}} \qquad\qquad\qquad (II)$$

If the mass spectrometric signals of labeled and unlabeled peptides were overlapping, the

monoisotopic peak was used to determine the total abundance of unlabeled peptide based on the

natural distribution of heavy isotopes, as previously described [21]. The generation time t_d was

subsequently calculated following equation (III), with Δt being the incubation time.

$$t_d = \frac{\Delta t}{n} \qquad\qquad\qquad (III)$$

Results

Sulfur oxidation by active groundwater microbes

The response of the groundwater microbiota to thiosulfate resulted in increasing rates of sulfur

oxidation. Within the first three weeks of incubation, thiosulfate consumption rates in the

microcosms were slowest, with 1.7 ± 1.9 µmol d^{-1} (mean ± SD), and where matched by an equally

slow oxygen consumption of 5.5 ± 2.0 µmol d^{-1} (Figure S1). Nevertheless, Raman microspectroscopic

analysis showed that more than 95% of cells were active already within 12 days of incubation, based

on the formation of a distinct C-D-band at a wavelength position between 2,100 and 2,200 cm^{-1} in

single-cell Raman spectra from the microcosm amended with D_2O (Figure 1, Figure S2). The C-D band

formation indicated metabolic activity, as new biomolecules were synthesized by incorporating

deuterium from D_2O into carbon-deuterium bonds. The relative intensity of the C-D-band increased

from 18.3% after 12 days to 25.7% after 47 days of incubation (median values), indicating continued

microbial growth and cross-feeding of deuterium-labeled organic carbon. Within 70 days,

consumption rates reached 7.2 ± 2.0 µmol d^{-1} for thiosulfate and 12.8 ± 3.2 µmol d^{-1} for oxygen,

representing a significant increase (p = 0.0006 (thiosulfate) and p = 0.0013 (oxygen), Student's t-test,

two-tailed) (Figure S1). Sulfate was produced with rates between 8.1 and 9.6 µmol d^{-1} with no

significant changes through the entire duration of the experiment. The observed stoichiometry for

oxygen:thiosulfate:sulfate was approximately 2.8:1:2.6 over the entire time course of incubation,

close to the theoretical values of 2:1:2 for oxygen-dependent thiosulfate oxidation.

Organism-specific ^{13}C incorporation reveals distinct lifestyles

To quantify the carbon utilization dynamics of the microbial key players, we conducted genome-

resolved SIP-metaproteomics based on the metagenome-assembled genomes (MAGs) acquired from

incubations after 21, 43, and 70 days. Our novel SIsCA (Stable Isotope Cluster Analysis) approach

revealed five clusters among the 31 most abundant MAGs with distinct carbon utilization patterns

(Figure 2, Figure S3). Organisms represented by the MAGs in cluster I that were related to

Thiobacillus (*Burkholderiales*) exhibited a stable ^{13}C relative isotope abundance (RIA) of 95% over 21,

43, and 70 days (Figure 2), resulting from exclusive assimilation of CO_2. By comparing the signal

intensity of ^{12}C and ^{13}C peptides, the generation times of these strict autotrophs were found to be

below two days at the first timepoint (Figure 3), highlighting the rapid production of new, ^{13}C-labeled

biomass from $^{13}CO_2$. These strict autotrophs made up only 11% of the five clusters, based on the

normalized coverage of the metagenomics dataset, and 3.2 ± 3.1% (mean ± sd) of the total

community, based on 16S rRNA gene profiles (Figure S4, Figure S5, Supplementary results).

Organisms in cluster II, related to *Methyloversatilis*, *Polaromonas* and *Dechloromonas* (all

Burkholderiales), displayed an intermediate ^{13}C RIA of 65% after 21 days of incubation (Figure 2),

suggesting that these organisms used labeled organic carbon from primary production as well as

unlabeled organic carbon from the groundwater. After 43 and 70 days, their ^{13}C RIA had increased

substantially to 91%, indicating a switch to chemolithoautotrophic growth, potentially due to

limitation of organic carbon. These mixotrophs were more than twice as abundant as cluster I,

representing 26% of the normalized coverage, and they showed generation times from 2 to 4 days

(Figure 3).

In cluster III, average ^{13}C RIAs increased from 65 to 76% and in cluster IV from 18 to 53% from 21 to

70 days of incubation (Figure 2), indicating heterotrophic growth in part on organic carbon from

chemolithoautotrophic primary production of clusters I and II. The observed increase in ^{13}C RIA

reflected the increasing labeling of the organic carbon in the microcosms due to $^{13}CO_2$ fixation.

Variations in the ^{13}C RIAs between species suggested different extents of cross-feeding on the

chemolithoautotrophically produced organic carbon, potentially due to preferences of different

organic carbon compounds. Cluster III comprised the largest part of the community, representing

28% of the normalized coverage, and cluster IV made up 20%. Most of the organisms in these

clusters, related to various *Burkholderiales* and other bacterial groups, exhibited generation times of

3 to 4 days (Figure 3). Members of cluster III affiliated with *Hydrogenophaga*, *Vitreoscilla* and

Rubrivivax, however, showed faster growth comparable with cluster I.

In cluster V, average ^{13}C RIAs reached 6% after 21 days of incubation and did not change at the later

time points, suggesting that the organisms were active only in the first period, where they displayed

a heterotrophic lifestyle. Nevertheless, these organisms still represented 15% of the normalized

coverage of all clusters. Generation times in this period were slightly longer and displayed more

variability than for clusters III and IV, ranging from 3.5 days for one *Acidovorax* species to eight days

for *Aquabacterium* (Figure 3).

Approximately 43% of carbon in the microbial community was replaced by ^{13}C after 21 days of

incubation, based on the corresponding peptide RIAs of all analyzed MAGs. This portion increased

further to 68% after 43 days and 80% after 70 days of incubation. Quantitative DNA-SIP confirmed

the increasing labeling the groundwater community by $^{13}CO_2$-derived carbon by an increase in the

number and buoyant density shift of ^{13}C-labeled OTUs (Figure S6) (see Supplement). Using SIsCA, we

revealed the carbon flux from autotrophic cluster I to mixotrophic cluster II, as well as the cross-

feeding of the heterotrophic clusters III to V on $^{13}CO_2$-derived organic carbon. Recycling of organic

carbon likely played a role across the entire groundwater microbiome.

Functional characterization of MAGs reveals putative mixotrophs

All putative autotrophs detected employed the Calvin-Benson-Bassham (CBB) cycle for CO_2 fixation

(Figure 4). Subunits of the key enzyme ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO)

were detected for 15 of the 31 MAGs. In 14 of these, further enzymes of the CBB cycle were present.

No other complete CO_2 fixation pathway was found. For the Wood-Ljungdahl pathway, either no

carbon monoxide dehydrogenase/acetyl-CoA synthase or no accessory enzymes were detected, and

only few accessory enzymes of the reductive tricarboxylic acid cycle or the 3-hydroxypropanoate

cycle were identified. Interestingly, proteins of the CBB cycle were not only present in strict or

facultative autotrophs related to *Thiobacillus* from cluster I or *Methyloversatilis*, *Polaromonas* and

Dechloromonas from cluster II, but also in heterotrophic organisms affiliated with genera

Hydrogenophaga, *Rhodoferax, Paucibacter* and *Rubrivivax*, from cluster III and IV. Hence, the ability

for mixotrophic growth was present in a larger number of organisms in the groundwater microcosms.

These mixotrophs constituted more than 50% of the microbial abundance over all clusters,

suggesting that they were of higher importance for the carbon cycling in the microcosms than strict

autotrophs.

MAGs express pathways for the utilization of reduced sulfur compounds

Sixteen of the MAGs obtained expressed proteins for sulfur oxidation via the Sox or the Dsr enzyme

system (Figure 4). Organisms from clusters II to IV, affiliated with the genera *Methyloversatilis*,

Dechloromonas, *Hydrogenophaga*, *Rhodoferax* and other *Betaproteobacteriales*, were exclusively

using the Sox system, which allows the complete oxidation of thiosulfate to sulfate, without any free

intermediates [28]. The corresponding MAGs exhibited gene clusters sharing a conserved

soxCDYZAXB gene order (Figure S7), featuring the core components of the Kelly-Friedrich pathway

[53, 54]. Further genes, including *soxVW*, *soxEF*, *soxTRS*, and *soxH*, were typically found in different

regions of the MAGs, showing variable arrangements. Organisms from cluster I, affiliated with

Thiobacillus, produced enzymes from both the Sox and the Dsr system. The corresponding MAGs

contained a truncated *soxXYZAB* gene cluster, lacking the genes *soxCD* required for oxidation of the

sulfane group of thiosulfate, and hence likely utilized the branched thiosulfate oxidation pathway. In

this pathway, the Dsr enzyme system, acting in reverse in *Thiobacillus* [55], oxidizes the sulfane-

derived sulfur atom to sulfite, with elemental sulfur as intermediate [28]. The MAGs in cluster I

contained the conserved operon structure *dsrABEFHCMKLJOPNR*, including the genes *dsrEFH* and

dsrL typical for sulfur oxidizers, but lacking the gene *dsrD* indicative of sulfate reducers [29]. In

addition, these organisms also expressed the *aprAB* and *sat* genes encoding Adenosine-5'-

phosphosulfate reductase and ATP sulfurylase, which can likewise act in reverse to oxidize sulfite to

sulfate [56]. Hence, facultative chemolithoautotrophs in the groundwater microcosms used the Sox

system to oxidize thiosulfate to sulfate, whereas obligate chemolithoautotrophs utilized an

incomplete version of the Sox system to oxidize the sulfone group and the Dsr/Apr/Sat system to

oxidize the sulfane group of thiosulfate.

Use of alternative electron acceptors and donors in sulfur oxidizers

For 15 of the MAGs of sulfur oxidizers, cytochrome c oxidase and other enzymes of the respiratory

chain were detected, and for 12 of them, enzymes for nitrate reduction (nitrate reductase, nitrite

reductase, and nitric oxide reductase) were found (Figure 4). Multiple sulfur oxidizers affiliated with

Dechloromonas and *Rhodoferax* had both pathways expressed simultaneously. Proteins for ammonia

oxidation, ammonia monooxygenase and hydroxylamine oxidoreductase, were produced by various

organisms in clusters I to IV, such as *Thiobacillus* and *Methyloversatilis*. Furthermore, MAG_77

(*Thiobacillus*), MAG_55 (*Dechloromonas*) and MAG_7 (*Hydrogenophaga*) expressed [NiFe]-

hydrogenase genes.

MAGs salvage a variety of organic carbon sources

MAGs of clusters I, II, and III showed an increasing versatility to utilize organic carbon compounds.

While the strict autotrophs in cluster I only expressed pathways for sugar degradation, clusters II to V

expressed utilization pathways and transporters targeting simple sugars (Glycolysis, pentose

phosphate pathway), amino acids (TCA cycle), fatty acids (beta-oxidation), C_1 compounds, and

aromatic compounds (Figure 4). The TCA cycle was one of the most abundant metabolic modules in

clusters II to V. Pathways for toluene and ethylbenzene utilization were expressed by organisms

affiliated with *Dechloromonas* and *Rhizobacter* (*Betaproteobacteriales*), respectively. Enzymes for

naphthalene and catechol utilization were detected, e.g., for organisms related to *Hydrogenophaga*

and *Pseudomonas*. The utilization of C_1 compounds was primarily detected for organisms related to

Methyloversatilis, which possessed methanol dehydrogenase, formate dehydrogenase and various

enzymes for tetrahydromethanopterin-dependent C_1-cycling. Enzymes for degradation of complex

carbohydrates such as starch and chitin were produced by organisms related to *Microbacterium* and

Sediminibacterium.

In mixotrophs and heterotrophs from clusters II to V, peptides of import systems for amino acids and

carboxylic acids (e.g. alpha-keto acids, C_4-dicarboxylates, lactate) were highly abundant (Figure 4).

The microorganisms from clusters III to V, which were exclusively growing heterotrophically in our

microcosms, possessed further transport systems for carbohydrates and nucleotides, and showed

the broadest variety of import functions. For the obligate autotrophs in cluster I, only transporters

targeting cations (mostly iron transporters) and phosphate were found.

Discussion

Mixotrophs, and not obligate chemolithoautotrophs, were the most abundant active organisms in

our groundwater microcosms. This was contrary to our hypothesis that strict chemolithoautotrophic

sulfur oxidizers would be stimulated to dominate the carbon cycling. Instead, we observed a highly

diverse community, where strict chemolithoautotrophs only comprised 3%. This is astonishing as

thiosulfate and oxygen were available in sufficient amounts throughout the whole experiment,

favoring chemolithoautotrophic growth. The high sensitivity of Raman spectroscopy revealed an

immediate activity of the groundwater microbes without a discernable lag phase, despite low sulfur

oxidation rates at the start of incubation.

Our novel SIsCA approach allowed us to identify active microbes and their expressed metabolic

functions, and to quantify the carbon uptake and further transfer within a diverse community over

time (Figure 5). This approach goes far beyond other studies that pointed to the importance of

chemolithoautotrophy in groundwater based on functional gene and metagenomics data [10, 11, 14,

57], as we shed light on how CO_2-derived carbon is assimilated and cycled by the groundwater

microbiota. Within 21 days, 43% of the total groundwater microbial biomass consisted of CO_2-

derived carbon, increasing to 80% after 70 days. This rapid enrichment of CO_2-derived carbon did not

occur via a fixed, linear path from chemolithoautotrophs to heterotrophs, but through complex

trophic interactions dominated by mixotrophs that heavily rely on the recycling of organic carbon.

More broadly, microbes appear to be well adapted to the oligotrophic conditions, characteristic for

the modern groundwater of the Hainich CZE, by being flexible, versatile, and able to utilize all

nutrient sources available.

This opportunistic strategy to use autotrophy to fill up their carbon demand allowed the mixotrophs

in our groundwater microcosms to dominate the microbial community despite the excess of reduced

sulfur to fuel chemolithoautotrophy. The abundant mixotrophs strongly preferred heterotrophic

growth to the fixation of CO_2. This is presumably a consequence of the higher metabolic cost of

carbon assimilation via the CBB cycle [58, 59]. The ability to fix CO_2, however, allows them to succeed

when suitable organic carbon becomes limited in oligotrophic systems. The mixotrophs of cluster II,

for example, switched from heterotrophic growth to CO_2 fixation at later time points, likely due to

such limitations. The mixotrophs of cluster III expressed pathways for autotrophic growth as well, but

were never required to fix CO_2, as they were able to access a larger repertoire of carbon sources.

The ability to utilize CO_2 fixation as a fill-up function for organic carbon proved highly advantageous

for mixotrophs. With generation times of 4 days to less than 2 days, mixotrophs were able to grow

considerably faster than organisms of cluster IV and V that were restricted to a purely heterotrophic

lifestyle, exhibiting generation times up to 8 days. Surprisingly, heterotrophs were also able to use

reduced sulfur compounds in the groundwater microcosms, suggesting a chemolithoheterotrophic

lifestyle. In modern groundwater habitats, the constant release of reduced sulfur during weathering

makes sulfur oxidation an attractive alternative to the oxidation of organic compounds for energy

conservation. The energetic requirement for CO_2 fixation is higher than the potential gain from

organic carbon oxidation. The most efficient strategy for both mixotrophs and heterotrophs is to gain

most energy from sulfur oxidation, preserving the precious organic carbon for anabolic

requirements.

The diversity of organic carbon utilization showed an inverse, gradual relationship to the CO_2 fixation

activity of the organisms in our groundwater microcosms. On one end of the gradient were the strict

autotrophs of cluster I, relying only on CO_2 as carbon source. No transporters for organic carbon from

these organisms were detected, and their metabolic capabilities were restricted to simple sugars,

likely to allow the utilization of carbon assimilated via the CBB cycle [58]. On the other end of the

gradient were organisms from clusters IV and V, exclusively assimilating organic carbon. To succeed

without autotrophic CO_2 fixation capabilities, these organisms were required to express a high range

of organic carbon transport and assimilation pathways. The most successful organisms, however,

were the mixotrophs of cluster II and III in the middle of the gradient, allowing them to become 5-

fold more abundant than the strict autotrophs. These organisms combined a range of abilities for

organic carbon assimilation with CO_2 fixation as a fill-up function, providing independence from

dynamics in the organic carbon supply in the groundwater.

Organisms affiliated with *Burkholderiales*, related to the key mixotrophic taxa in our groundwater

microcosms, were found to have the highest abundance of RuBisCO-encoding transcripts at our

groundwater site [14]. For taxa like *Polaromonas*, *Dechloromonas*, *Hydrogenophaga*, and

Rhodoferax, the ability for sulfur oxidation has been suggested based on genomic evidence [60-63].

However, chemolithoautotrophic growth on reduced sulfur compounds has not been observed for

these genera in pure culture so far. Our study demonstrates that these organisms can use reduced

sulfur as energy source, and *Polaromonas*, *Dechloromonas* and potentially *Hydrogenophaga* used it

to fuel autotrophic growth. These sulfur oxidizers expressed pathways for both aerobic respiration

and denitrification, although no nitrate was added to the microcosms and nitrate concentrations in

the groundwater are only around 10 mg/L at this well [27]. Constitutive production of denitrification

enzymes might be energetically cheaper than the regulation of gene expression [64], while allowing

to flexibly switch their terminal electron acceptor when oxygen is limited. This ability also explains

the frequent presence of these organisms in groundwater and other aquatic systems where the

oxidation of thiosulfate linked to denitrification plays an important role [65-67].

Some sulfur oxidizers in the groundwater microcosms were expressing proteins to use reduced

nitrogen compounds and hydrogen as electron donor, agreeing with previous experimental evidence

for their growth on H_2 and CO_2 [68-70]. Also other H_2 oxidizing organisms are known to grow

chemolithoautotrophically on reduced sulfur compounds [71], suggesting that further bacterial taxa

might harbor sulfur-oxidizing mixotrophs with a similar, versatile lifestyle.

The *Thiobacillus*-related organisms from cluster I are known to be obligate autotrophs, as they

feature an incomplete TCA cycle that prevents heterotrophic growth [72]. Performing thiosulfate and

hydrogen-driven denitrification, *Thiobacillus* constituted up to 50% of an enrichment culture

obtained from the groundwater of the Hainich CZE [66]. *In situ*, however, *Thiobacillus* typically are

only present in low abundance [14], suggesting a lack of competitiveness with the metabolically

flexible mixotrophs in modern groundwater, whereas they are known to be more dominant in

deeper, more CO_2-rich systems [10]. Thiobacillus can store the elemental sulfur produced as

intermediate by the Dsr enzyme system in periplasmic granules [72, 73]. This storage might allow the

organism to withstand times where no reduced sulfur compounds in the groundwater are available.

The diverse metabolic potential of sulfur-oxidizing microorganisms in groundwater seems to be a

common trait for this group [4, 13, 16]. Being a mixotrophic sulfur oxidizer has two advantages: first,

they are independent from surface input dynamics, since reduced sulfur compounds are released

from weathering of interspersed pyrite minerals [25-27]. Second, the employed repertoire of

metabolic pathways enables them to tailor the organic carbon availability in modern groundwater to

their anabolic requirements. This includes carbohydrate degradation pathways for surface-derived

plant polymers [8, 74], amino acid and nucleotide uptake systems for microbially-derived carbon [75,

76], C_1 metabolic functions for C_1 carbon compounds from biomass degradation [77], and

hydrocarbon degradation pathways for rock-derived carbon [78, 79]. We propose that such a

strategy, combining the utilization of a readily available energy source and versatile carbon

assimilation pathways, should also be successful in other oligotrophic systems, potentially explaining

the success of mixotrophs in habitats such as oligotrophic lakes or the upper ocean [80, 81].

Conclusions

Our novel SIsCA analysis enabled us to quantitatively and temporally resolve the carbon flux from

$^{13}CO_2$ through individual key players in a groundwater microbial community. We observed that

mixotrophs were most successful in this oligotrophic environment because they employed CO_2

fixation as a fill-up function to supplement their organic carbon requirements. CO_2-derived carbon

rapidly entered the microbial biomass through a complex trophic network relying on the efficient

recycling of organic carbon. As a further strategy to cope with low organic carbon levels, autotrophic,

mixotrophic and heterotrophic organisms were able to utilize reduced sulfur compounds as energy

sources, allowing them to preserve organic carbon for assimilation. A variety of carbon assimilation

pathways enabled mixotrophs and heterotrophs to make optimal use of the scarce organic carbon

compounds present in the oligotrophic groundwater. The combination of versatile carbon

assimilation pathways and inorganic energy sources is likely a key strategy for the success of

microbes in modern groundwater as well as other oligotrophic environments. Furthermore, our

study demonstrates how next-generation physiology approaches like SIsCA are essential to test

hypotheses established in metagenomics studies and enhance our understanding of microbial

survival strategies and their role in ecosystem functioning.

Declarations
479

Ethics approval and consent to participate
480

481   Not applicable

Consent for publication
482

483   Not applicable

Availability of data and materials
484

485   Metagenomic and amplicon sequencing data that support the findings of this study have been

486   deposited in NCBI with the BioProject accession PRJNA633367.

487   The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium

488   via the PRIDE [82] partner repository with the dataset identifier PXD024889.

Competing interests
489

490   The authors declare that they have no competing interests.

Funding
491

This work was financially supported by the Deutsche Forschungsgemeinschaft via the Collaborative

Research Centre AquaDiva (CRC 1076 AquaDiva - Project-ID 218627073) of the Friedrich-Schiller-

University Jena and the German Center for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig.

Martin Taubert gratefully acknowledges funding by the DFG under Germany's Excellence Strategy -

EXC 2051 - Project-ID 390713860. Climate chambers to conduct experiments under controlled

temperature conditions were financially supported by the Thüringer Ministerium für Wirtschaft,

Wissenschaft und Digitale Gesellschaft (TMWWDG; project B 715-09075). Martin von Bergen and

Nico Jehmlich are grateful for support by the UFZ-funded platform for metabolomics and proteomics

(MetaPro). The funding bodies played no role in the design of the study and collection, analysis, and

interpretation of data and in writing the manuscript.

Authors' contributions

503   MT and KK conceived and designed the study. MT conducted the microcosm experiments and

504   molecular biology work. GAM, RH, PR and JP conducted the Raman microspectroscopic analysis. NJ

505   and MvB conducted mass spectrometric analysis for metaproteomics. MT analyzed the

506   metagenomics data with help of WAO and BMH, and analyzed the SIP-metaproteomics data. MT

507   wrote the manuscript with contributions from all other authors.

Acknowledgements

We are grateful to Robert Lehmann, Falko Gutmann, Heiko Minkmar, Jens Wurlitzer and Lena

Carstens for help with field and lab work, and sampling of groundwater. We thank Julian Hniopek for

his help with the Raman measurements. We thank Ivonne Görlich and Marco Groth from the Core

Facility DNA sequencing of the Leibniz Institute on Aging - Fritz Lipmann Institute in Jena for their

help with Illumina sequencing. We further thank Daniel Desirò and Martin Hölzer for assistance with

sequencing organization.

References

1.   Flemming HC, Wuertz S. Bacteria and archaea on Earth and their abundance in biofilms. Nat

Rev Microbiol. 2019;17:247-260.

2.   Wrighton KC, Thomas BC, Sharon I, Miller CS, Castelle CJ, VerBerkmoes NC, et al.

Fermentation, hydrogen, and sulfur metabolism in multiple uncultivated bacterial phyla.

Science. 2012;337:1661-1665.

3.   Albertsen M, Hugenholtz P, Skarshewski A, Nielsen KL, Tyson GW, Nielsen PH. Genome

sequences of rare, uncultured bacteria obtained by differential coverage binning of multiple

metagenomes. Nat Biotechnol. 2013;31:533-538.

4.   Anantharaman K, Brown CT, Hug LA, Sharon I, Castelle CJ, Probst AJ, et al. Thousands of

microbial genomes shed light on interconnected biogeochemical processes in an aquifer

system. Nat Commun. 2016;7:1-11.

5.   Parks DH, Rinke C, Chuvochina M, Chaumeil PA, Woodcroft BJ, Evans PN, et al. Recovery of

nearly 8,000 metagenome-assembled genomes substantially expands the tree of life. Nat

Microbiol. 2017;2:1533-1542.

6.   Gleeson T, Befus KM, Jasechko S, Luijendijk E, Cardenas MB. The global volume and

distribution of modern groundwater. Nat Geosci. 2016;9:161-167.

7.   Akob DM, Küsel K. Where microorganisms meet rocks in the Earth's Critical Zone.

Biogeosciences. 2011;8:3531-3543.

8.   Griebler C, Lueders T. Microbial biodiversity in groundwater ecosystems. Freshwater Biol.

2009;54:649-677.

9.   Bell E, Lamminmäki T, Alneberg J, Andersson AF, Qian C, Xiong WL, et al. Active sulfur cycling

in the terrestrial deep subsurface. ISME J. 2020;14:1260-1272.

10. Emerson JB, Thomas BC, Alvarez W, Banfield JF. Metagenomic analysis of a high carbon

dioxide subsurface microbial community populated by chemolithoautotrophs and bacteria

and archaea from candidate phyla. Environ Microbiol. 2016;18:1686-1703.

11. Probst AJ, Ladd B, Jarett JK, Geller-McGrath DE, Sieber CMK, Emerson JB, et al. Differential

depth distribution of microbial function and putative symbionts through sediment- hosted

aquifers in the deep terrestrial subsurface. Nat Microbiol. 2018;3:328-336.

12.　Anantharaman K, Hausmann B, Jungbluth SP, Kantor RS, Lavy A, Warren LA, et al. Expanded

diversity of microbial groups that shape the dissimilatory sulfur cycle. ISME J. 2018;12:1715-

1728.

13.　Wegner CE, Gaspar M, Geesink P, Herrmann M, Marz M, Küsel K. Biogeochemical regimes in

shallow aquifers reflect the metabolic coupling of the elements nitrogen, sulfur, and carbon.

Appl Environ Microb. 2019;85:1-18.

14.　Herrmann M, Rusznyak A, Akob DM, Schulze I, Opitz S, Totsche KU, et al. Large fractions of

CO_2-fixing microorganisms in pristine limestone aquifers appear to be involved in the

oxidation of reduced sulfur and nitrogen compounds. Appl Environ Microb. 2015;81:2384-

2394.

15.　Probst AJ, Castelle CJ, Singh A, Brown CT, Anantharaman K, Sharon I, et al. Genomic

resolution of a cold subsurface aquifer community provides metabolic insights for novel

microbes adapted to high CO_2 concentrations. Environ Microbiol. 2017;19:459-474.

16.　Jewell TNM, Karaoz U, Brodie EL, Williams KH, Beller HR. Metatranscriptomic evidence of

pervasive and diverse chemolithoautotrophy relevant to C, S, N and Fe cycling in a shallow

alluvial aquifer. ISME J. 2016;10:2106-2117.

17.　Handley KM, Bartels D, O'Loughlin EJ, Williams KH, Trimble WL, Skinner K, et al. The complete

genome sequence for putative H-2- and S-oxidizer *Candidatus* Sulfuricurvum sp., assembled

de novo from an aquifer-derived metagenome. Environ Microbiol. 2014;16:3443-3462.

18.　Neufeld JD, Vohra J, Dumont MG, Lueders T, Manefield M, Friedrich MW, et al. DNA stable-

isotope probing. Nat Protoc. 2007;2:860-866.

19.   von Bergen M, Jehmlich N, Taubert M, Vogt C, Bastida F, Herbst FA, et al. Insights from

quantitative metaproteomics and protein-stable isotope probing into microbial ecology.

ISME J. 2013;7:1877-1885.

20.   Taubert M, Vogt C, Wubet T, Kleinsteuber S, Tarkka MT, Harms H, et al. Protein-SIP enables

time-resolved analysis of the carbon flux in a sulfate-reducing, benzene-degrading microbial

consortium. ISME J. 2012;6:2291-2301.

21.   Taubert M, Baumann S, von Bergen M, Seifert J. Exploring the limits of robust detection of

incorporation of ^{13}C by mass spectrometry in protein-based stable isotope probing (protein-

SIP). Anal Bioanal Chem. 2011;401:1975-1982.

22.   Einsiedl F, Mayer B. Hydrodynamic and microbial processes controlling nitrate in a fissured-

porous karst aquifer of the Franconian Alb, Southern Germany. Environ Sci Technol.

2006;40:6697-6702.

23.   Schlesinger WH. On the fate of anthropogenic nitrogen. P Natl Acad Sci USA. 2009;106:203-

208.

24.   McCollom TM, Seewald JS. Serpentinites, hydrogen, and life. Elements. 2013;9:129-134.

25.   Rimstidt JD, Vaughan DJ. Pyrite oxidation: A state-of-the-art assessment of the reaction

mechanism. Geochim Cosmochim Ac. 2003;67:873-880.

26.   Schippers A, Jozsa PG, Sand W. Sulfur chemistry in bacterial leaching of pyrite. Appl Environ

Microb. 1996;62:3424-3431.

27.   Kohlhepp B, Lehmann R, Seeber P, Küsel K, Trumbore SE, Totsche KU. Aquifer configuration

and geostructural links control the groundwater quality in thin-bedded carbonate-siliciclastic

alternations of the Hainich CZE, central Germany. Hydrol Earth Syst Sc. 2017;21:6091-6116.

28.   Ghosh W, Dam B. Biochemistry and molecular biology of lithotrophic sulfur oxidation by

taxonomically and ecologically diverse Bacteria and Archaea. FEMS Microbiol Rev.

2009;33:999-1043.

29.   Grimm F, Franz B, Dahl C. Thiosulfate and sulfur oxidation in purple sulfur bacteria.

In:editors. Microbial Sulfur Metabolism. Springer; 2008: 101-116.

30.   Schwab VF, Herrmann M, Roth VN, Gleixner G, Lehmann R, Pohnert G, et al. Functional

diversity of microbial communities in pristine aquifers inferred by PLFA- and sequencing-

based approaches. Biogeosciences. 2017;14:2697-2714.

31.   DEV-D15. Deutsche Einheitsverfahren zur Wasser-, Abwasser- und Schlammuntersuchung.

Physikalische, chemische, biologische und bakteriologische Verfahren., vol. DEV-D15.

Weinheim, Germany: VCH Verlagsgesellschaft; 1975.

32.   Tabatabai M. A rapid method for determination of sulfate in water samples. Environmental

Letters. 1974;7:237-243.

33.   R Core Team. R: A language and environment for statistical computing. In: R Foundation for

Statistical Computing. Vienna, Austria: R Core Team; 2019.

34.   Ryabchykov O, Bocklitz T, Ramoji A, Neugebauer U, Foerster M, Kroegel C, et al.

Automatization of spike correction in Raman spectra of biological samples. Chemometr Intell

Lab. 2016;155:1-6.

35.   Dörfer T, Bocklitz T, Tarcea N, Schmitt M, Popp J. Checking and improving calibration of

Raman spectra using chemometric approaches. Z Phys Chem. 2011;225:753-764.

36.   Bocklitz TW, Dörfer T, Heinke R, Schmitt M, Popp J. Spectrometer calibration protocol for

Raman spectra recorded with different excitation wavelengths. Spectrochim Acta A.

2015;149:544-549.

37.   Guo SX, Heinke R, Stöckel S, Rösch P, Bocklitz T, Popp J. Towards an improvement of model

transferability for Raman spectroscopy in biological applications. Vib Spectrosc. 2017;91:111-

118.

38.   Liland KH, Mevik BH, Canteri R. Baseline correction of spectra. (R package). ver. 1.3-0. 2020.

39.   Taubert M, Stöckel S, Geesink P, Girnus S, Jehmlich N, von Bergen M, et al. Tracking active

groundwater microbes with D_2O labelling to understand their ecosystem function. Environ

Microbiol. 2018;20:369-384.

40.   Bushnell B. BBTools software package. URL http://sourceforgenet/projects/bbmap. 2014.

41.   Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. metaSPAdes: a new versatile metagenomic

assembler. Genome Res. 2017;27:824-834.

42.   Wu YW, Simmons BA, Singer SW. MaxBin 2.0: an automated binning algorithm to recover

genomes from multiple metagenomic datasets. Bioinformatics. 2016;32:605-607.

43.   Kang DWD, Li F, Kirton E, Thomas A, Egan R, An H, et al. MetaBAT 2: an adaptive binning

algorithm for robust and efficient genome reconstruction from metagenome assemblies.

Peerj. 2019;7:1-13.

44.   Graham ED, Heidelberg JF, Tully BJ. BinSanity: unsupervised clustering of environmental

microbial assemblies using coverage and affinity propagation. Peerj. 2017;5:1-19.

45.   Uritskiy GV, DiRuggiero J, Taylor J. MetaWRAP - a flexible pipeline for genome-resolved

metagenomic data analysis. Microbiome. 2018;6:1-13.

46.   Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify genomes with

the Genome Taxonomy Database. Bioinformatics. 2019;36:1925-1927.

47.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the

632   quality of microbial genomes recovered from isolates, single cells, and metagenomes.

633   Genome Res. 2015;25:1043-1055.

634   48.   Jain C, Rodriguez RL, Phillippy AM, Konstantinidis KT, Aluru S. High throughput ANI analysis of

635   90K prokaryotic genomes reveals clear species boundaries. Nat Commun. 2018;9:1-8.

636   49.   Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics. 2014;30:2068-

637   2069.

638   50.   Eren AM, Esen OC, Quince C, Vineis JH, Morrison HG, Sogin ML, et al. Anvi'o: an advanced

639   analysis and visualization platform for 'omics data. Peerj. 2015;3:1-29.

640   51.   Orellana LH, Rodriguez-R LM, Konstantinidis KT. ROCker: accurate detection and

641   quantification of target genes in short-read metagenomic data sets by modeling sliding-

642   window bitscores. Nucleic Acids Res. 2017;45:e14.

643   52.   Dixon P. VEGAN, a package of R functions for community ecology. J Veg Sci. 2003;14:927-

644   930.

645   53.   Friedrich CG, Rother D, Bardischewsky F, Quentmeier A, Fischer J. Oxidation of reduced

646   inorganic sulfur compounds by bacteria: Emergence of a common mechanism? Appl Environ

647   Microb. 2001;67:2873-2882.

648   54.   Kelly DP, Shergill JK, Lu WP, Wood AP. Oxidative metabolism of inorganic sulfur compounds

649   by bacteria. Anton Leeuw Int J G. 1997;71:95-107.

650   55.   Beller HR, Letain TE, Chakicherla A, Kane SR, Legler TC, Coleman MA. Whole-genome

651   transcriptional analysis of chemolithoautotrophic thiosulfate oxidation by *Thiobacillus*

652   *denitrificans* under aerobic versus denitrifying conditions. J Bacteriol. 2006;188:7005-7015.

56. Beller HR, Chain PSG, Letain TE, Chakicherla A, Larimer FW, Richardson PM, et al. The

genome sequence of the obligately chemolithoautotrophic, facultatively anaerobic

bacterium *Thiobacillus denitfificans*. J Bacteriol. 2006;188:1473-1488.

57. Alfreider A, Vogt C, Geiger-Kaiser M, Psenner R. Distribution and diversity of autotrophic

bacteria in groundwater systems based on the analysis of RubisCO genotypes. Syst Appl

Microbiol. 2009;32:140-150.

58. Berg IA. Ecological aspects of the distribution of different autotrophic CO_2 fixation pathways.

Appl Environ Microb. 2011;77:1925-1936.

59. Quayle JR, Ferenci T. Evolutionary aspects of autotrophy. Microbiol Rev. 1978;42:251-273.

60. Mattes TE, Alexander AK, Richardson PM, Munk AC, Han CS, Stothard P, et al. The genome of

Polaromonas sp. strain JS666: Insights into the evolution of a hydrocarbon- and xenobiotic-

degrading bacterium, and features of relevance to biotechnology. Appl Environ Microb.

2008;74:6405-6416.

61. Salinero KK, Keller K, Feil WS, Feil H, Trong S, Di Bartolo G, et al. Metabolic analysis of the soil

microbe *Dechloromonas aromatica* str. RCB: indications of a surprisingly complex life-style

and cryptic anaerobic pathways for aromatic degradation. BMC Genomics. 2009;10:1-23.

62. Kämpfer P, Schulze R, Jäckel U, Malik KA, Amann R, Spring S. *Hydrogenophaga defluvii* sp.

nov. and *Hydrogenophaga atypica* sp. nov., isolated from activated sludge. Int J Syst Evol

Micr. 2005;55:341-344.

63. Jin CZ, Zhuo Y, Wu XW, Ko SR, Li TH, Jin FJ, et al. Genomic and metabolic insights into

denitrification, sulfur oxidation, and multidrug efflux pump mechanisms in the bacterium

Rhodoferax sediminis sp. nov. Microorganisms. 2020;8:262.

64.   Geisel N. Constitutive versus responsive gene expression strategies for growth in changing

676   environments. Plos One. 2011;6:e27033.

677   65.   Kumar S, Herrmann M, Thamdrup B, Schwab VF, Geesink P, Trumbore SE, et al. Nitrogen loss

678   from pristine carbonate-rock aquifers of the Hainich Critical Zone Exploratory (Germany) is

679   primarily driven by chemolithoautotrophic anammox processes. Front Microbiol.

680   2017;8:1951.

681   66.   Kumar S, Herrmann M, Blohm A, Hilke I, Frosch T, Trumbore SE, et al. Thiosulfate- and

682   hydrogen-driven autotrophic denitrification by a microbial consortium enriched from

683   groundwater of an oligotrophic limestone aquifer. FEMS Microbiol Ecol. 2018;94:fiy141.

684   67.   Sun SS, Liu J, Zhang MP, He SB. Thiosulfate-driven autotrophic and mixotrophic

685   denitrification processes for secondary effluent treatment: Reducing sulfate production and

686   nitrous oxide emission. Bioresource Technol. 2020;300:122651.

687   68.   Sizova M, Panikov N. *Polaromonas hydrogenivorans* sp. nov., a psychrotolerant hydrogen-

688   oxidizing bacterium from Alaskan soil. Int J Syst Evol Micr. 2007;57:616-619.

689   69.   Shrout JD, Scheetz TE, Casavant TL, Parkin GF. Isolation and characterization of autotrophic,

690   hydrogen-utilizing, perchlorate-reducing bacteria. Appl Microbiol Biot. 2005;67:261-268.

691   70.   Willems A, Busse J, Goor M, Pot B, Falsen E, Jantzen E, et al. *Hydrogenophaga*, a new genus

692   of hydrogen-oxidizing bacteria that includes *Hydrogenophaga flava* comb. nov. (formerly

693   *Pseudomonas flava*), *Hydrogenophaga palleronii* (formerly *Pseudomonas palleronii*),

694   *Hydrogenophaga pseudoflava* (formerly *Pseudomonas pseudoflava* and *Pseudomonas*

695   *carboxydoflava*), and *Hydrogenophaga taeniospiralis* (formerly *Pseudomonas taeniospiralis*).

696   Int J Syst Bacteriol. 1989;39:319-333.

71.　Sano R, Kameya M, Wakai S, Arai H, Igarashi Y, Ishii M, et al. Thiosulfate oxidation by a

thermo-neutrophilic hydrogen-oxidizing bacterium, *Hydrogenobacter thermophilus*. Biosci

Biotech Bioch. 2010;74:892-894.

72.　Boden R, Hutt LP, Rae AW. Reclassification of *Thiobacillus aquaesulis* (Wood & Kelly, 1995) as

Annwoodia aquaesulis gen. nov., comb. nov., transfer of *Thiobacillus* (Beijerinck, 1904) from

the *Hydrogenophilales* to the *Nitrosomonadales*, proposal of *Hydrogenophilalia* class. nov

within the '*Proteobacteria*', and four new families within the orders *Nitrosomonadales* and

Rhodocyclales. Int J Syst Evol Micr. 2017;67:1191-1205.

73.　Katayama-Fujimura Y, Tsuzaki N, Hirata A, Kuraishi H. Polyhedral inclusion-bodies

(Carboxysomes) in *Thiobacillus* species with reference to the taxonomy of the genus

Thiobacillus. J Gen Appl Microbiol. 1984;30:211-222.

74.　Küsel K, Totsche KU, Trumbore SE, Lehmann R, Steinhäuser C, Herrmann M. How deep can

surface signals be traced in the Critical Zone? Merging biodiversity with biogeochemistry

research in a Central German Muschelkalk landscape. Front Earth Sci. 2016;4:32.

75.　Roth VN, Lange M, Simon C, Hertkorn N, Bucher S, Goodall T, et al. Persistence of dissolved

organic matter explained by molecular changes during its passage through soil. Nat Geosci.

2019;12:755-761.

76.　Herrmann M, Wegner CE, Taubert M, Geesink P, Lehmann K, Yan LJ, et al. Predominance of

Cand. Patescibacteria in groundwater is caused by their preferential mobilization from soils

and flourishing under oligotrophic conditions. Front Microbiol. 2019;10:1407.

77.　Gray CM, Monson RK, Fierer N. Emissions of volatile organic compounds during the

decomposition of plant litter. J Geophys Res-Biogeo. 2010;115:G03015.

78.   Benk SA, Yan LJ, Lehmann R, Roth VN, Schwab VF, Totsche KU, et al. Fueling diversity in the

720   subsurface: Composition and age of dissolved organic matter in the Critical Zone. Front Earth

721   Sci. 2019;7:296.

722   79.   Schwab VF, Nowak ME, Elder CD, Trumbore SE, Xu XM, Gleixner G, et al. ^{14}C-free carbon is a

723   major contributor to cellular biomass in geochemically distinct groundwater of shallow

724   sedimentary bedrock aquifers. Water Resour Res. 2019;55:2104-2121.

725   80.   Eiler A. Evidence for the ubiquity of mixotrophic bacteria in the upper ocean: Implications

726   and consequences. Appl Environ Microb. 2006;72:7431-7437.

727   81.   Hansson TH, Grossart HP, del Giorgio PA, St-Gelais NF, Beisner BE. Environmental drivers of

728   mixotrophs in boreal lakes. Limnol Oceanogr. 2019;64:1688-1705.

729   82.   Perez-Riverol Y, Csordas A, Bai JW, Bernal-Llinares M, Hewapathirana S, Kundu DJ, et al. The

730   PRIDE database and related tools and resources in 2019: improving support for quantification

731   data. Nucleic Acids Res. 2019;47:D442-D450.

Figure and Table legends

Figure 1: Quantification of deuterium incorporation by single cell Raman microspectroscopy.

Boxplots depict the relative intensity of the Raman C-D-band determined as A(C-D) / [A(C-D) + A(C-H)] from single cell Raman spectra. Spectra were obtained from samples of groundwater microcosms with 30% D_2O (shaded) or with H_2O only (empty) at different time points of incubation. Boxes show median, first and third quartile. Whiskers indicate 5[th] and 95[th] percentile. Outliers are depicted as dots. At least n=147 spectra were obtained at each time point.

Figure 2: Clustering of selected MAGs based on carbon utilization. (A) Stable isotope cluster analysis based on PCA of ^{13}C incorporation profiles over incubation time obtained by SIP-metaproteomics of samples from the ^{13}C-microcosms. Each point represents an organism associated with one MAG. Clusters of MAGs are indicated with Latin numbers. Ellipses depict 95% confidence intervals. Only MAGs are shown where at least two replicates of ^{13}C incorporation patterns per time point could be acquired. (B) Representative ^{13}C incorporation profiles of MAGs marked with an asterisk are given for each cluster. Heatmaps depict the extent of ^{13}C incorporation in peptides of the corresponding MAG after 21 (T1), 43 (T2) and 70 days (T3) of incubation in 5%-intervals, ranging from 0 to 100% ^{13}C relative isotope abundance.

Figure 3: Generation times of groundwater organisms in the microcosms. Given values were determined for the first 3 weeks of incubation, based on the relative abundance of ^{12}C and ^{13}C peptides. Shown are mean and standard deviation based on n ≥ 4 replicate determinations. Colored horizontal lines indicate average generation time for each cluster. bdl: generation time was below the detection limit of 2 days. na: no quantification of the generation time was possible.

Figure 4: Metabolic functions of selected MAGs. The sizes of the bubbles correspond to the total number of peptides for each MAG and each functional category identified at any time point. Metabolic functions are grouped into CO_2 fixation (red), sulfur cycling (yellow), nitrogen cycling (green), aerobic respiration and ATP synthesis (blue), organic carbon utilization (black), and import

functions (brown). The taxonomic categories "other" include peptides that were assigned to multiple

MAGs affiliated with the same genus. Only MAGs included in the stable isotope cluster analysis are

shown. RuBisCO: ribulose-1,5-bisphosphate carboxylase/oxygenase, CODH/ACS: carbon monoxide

dehydrogenase/acetyl-CoA synthase, TCA cycle: tricarboxylic acid cycle.

Figure 5: Carbon flux between microbial clusters. Red arrow inlays illustrate the portion of $^{13}CO_2$-

derived carbon assimilated by each microbial cluster after 21, 43 and 70 days. Overall arrow width

scales with total amount of carbon assimilated based on the relative abundance of the respective

microbial cluster in the metagenomics analysis. Fading grey arrows indicate uptake of unlabeled

organic carbon from the groundwater. Checkboxes highlight the presence and activity of metabolic

functions for CO_2 fixation, utilization of organic carbon and sulfur oxidation.

Figure 1

Figure 2

777 **Figure 3**

Figure 4

Figure 5