An image-computable model on how endogenous and exogenous attention differentially alter visual perception

Michael Jigo[1][*], David J. Heeger[1,2] & Marisa Carrasco[1,2]

[1] Center for Neural Science and [2] Department of Psychology
New York University, New York, NY 10003

[*] **Corresponding author information**:
Michael Jigo (michael.jigo@nyu.edu)

Author Contributions
M.J., D.J.H. and M.C. conceived the project; M.J. implemented the model with input from D.J.H. and M.C.; M.C. provided all the data to be modelled; M.J. wrote the manuscript with guidance and supervision from M.C. All three authors edited the manuscript.

Acknowledgements
This work was supported by National Institutes of Health RO1-EY019693 to M.C. We thank Michael Landy, Jonathan Winawer, Antoine Barbot, Hsin-Hung Li, as well as Antonio Fernández, Nina Hanning, Marc Himmelburg, Luke Huszar & Yong-Jun Lin and other members of the Carrasco lab for their helpful comments and discussion.

ABSTRACT

Attention alters perception across the visual field. Typically, endogenous (voluntary) and exogenous (involuntary) attention similarly improve performance in many visual tasks, but they have differential effects in some tasks. Extant models of visual attention assume that the effects of these two types of attention are identical and consequently do not explain differences between them. Here, we develop a model of spatial resolution and attention that distinguishes between endogenous and exogenous attention. We focus on texture-based segmentation as a model system because it has revealed a clear dissociation between both attention types. For a texture for which performance peaks at parafoveal locations, endogenous attention improves performance across eccentricity, whereas exogenous attention improves performance where the resolution is low (peripheral locations) but impairs it where the resolution is high (foveal locations) for the scale of the texture. Our model emulates sensory encoding to segment figures from their background and predict behavioral performance. To explain attentional effects, endogenous and exogenous attention require separate operating regimes across visual detail (spatial frequency). Our model reproduces behavioral performance across several experiments and simultaneously resolves three unexplained phenomena: (1) the parafoveal advantage in segmentation, (2) the uniform improvements across eccentricity by endogenous attention and (3) the peripheral improvements and foveal impairments by exogenous attention. Overall, we unveil a computational dissociation between each attention type and provide a generalizable framework for predicting their effects on perception across the visual field.

INTRODUCTION

Endogenous and exogenous spatial attention prioritize subsets of visual information and facilitate their processing without concurrent eye movements (1-3). Selection by endogenous attention is goal-driven and adapts to task demands whereas exogenous attention transiently and automatically orients to salient stimuli (1-3). In most visual tasks both types of attention typically improve visual perception similarly (e.g., acuity (4-6), visual search (7, 8), perceived contrast (9-11)). Consequently, models of visual attention do not distinguish between endogenous and exogenous attention (e.g., (12-19)). However, stark differences also exist. Each attention type differentially modulates neural responses (20, 21) and fundamental properties of visual processing, including temporal resolution (22, 23), texture sensitivity (24), sensory tuning (25), contrast sensitivity (26) and spatial resolution (27-34).

The effects of endogenous and exogenous attention are dissociable during texture segmentation, a visual task constrained by spatial resolution (reviews(1-3)). Whereas endogenous attention optimizes spatial resolution to improve the detection of an attended texture (32-34), exogenous attention reflexively enhances resolution even when detrimental to perception (27-31, 34). Extant models of attention do not explain these well-established effects.

Two main hypotheses have been proposed to explain how attention alters spatial resolution. Psychophysical studies ascribe attentional effects to modulations of spatial frequency (SF) sensitivity (30, 33). Neurophysiological (13, 35, 36) and neuroimaging (37, 38) studies bolster the idea that attention modifies spatial profiles of neural receptive fields (2). Both hypotheses provide qualitative predictions of attentional effects but do not specify their underlying neural computations.

Differences between endogenous and exogenous attention are well established in segmentation tasks and thus provide an ideal model system to uncover their separate roles in altering perception. Texture-based segmentation is a fundamental process of mid-level vision that isolates regions of local structure to extract figures from their background (39-41). Successful segmentation hinges on the overlap between the visual system's spatial resolution and the levels of detail (i.e., SF) encompassed by the texture (39, 41, 42). Consequently, the ability to distinguish between adjacent textures varies as resolution declines toward the periphery (43-46). Each attention type differentially alters texture segmentation, demonstrating that their effects shape spatial resolution (reviews(1-3)).

Current models of texture segmentation do not explain performance across eccentricity and the

distinct modulations by attention. Conventional models treat segmentation as a feedforward

process that encodes the elementary features of an image (e.g., SF and orientation), transforms

them to reflect the local structure (e.g., regions of similarly oriented bars), then pools across space

to emphasize texture-defined contours (39, 41, 47). Few of these models account for variations in

resolution across eccentricity (46, 48, 49) or endogenous (but not exogenous) attentional

modulations (18, 50). All others postulate that segmentation is a 'preattentive' (42) operation

whose underlying neural processing is impervious to attention (39, 41, 46-49).

Here, we develop a computational model in which feedforward processing and attentional gain

contribute to segmentation performance. We augment a conventional model of texture processing

(39, 41, 47). Our model varies with eccentricity and includes contextual modulation within local

regions in the stimulus via normalization (51), a canonical neural computation (52). The defining

characteristic of normalization is that an individual neuron is (divisively) suppressed by the

summed activity of neighboring neurons responsive to different aspects of a stimulus. We model

attention as multiplicative gains (attentional gain factors (15)) that vary with eccentricity and SF.

Attention shifts sensitivity toward fine or coarse spatial scales depending on the range of SFs

enhanced.

Our model is image-computable, which allowed us to reproduce behavior directly from grayscale

images used in psychophysical experiments (6, 26, 27, 29-33). The model explains three

signatures of texture segmentation hitherto unexplained within a single computational framework

(**Figure 1**). (i) The central performance drop (CPD) (27-34, 43-46) (**Figure 1A**), i.e., the parafoveal

advantage of segmentation over the fovea. (ii) The improvements in the periphery and impairments

at foveal locations induced by exogenous attention (27-32, 34) (**Figure 1B**). (iii) The equivalent

improvements across eccentricity by endogenous attention (32-34) (**Figure 1C**).

Whereas our analyses focused on texture segmentation, our model is general and can be applied

to other visual phenomena. We show that the model predicts contrast sensitivity across SF and

eccentricity as well as the effects of attention on contrast sensitivity and acuity; i.e. in tasks in

which both endogenous and exogenous attention have similar or differential effects on

performance. To preview our results, model comparisons revealed that normalization is necessary

to elicit the CPD and that separate profiles of gain enhancement across SF (26) generate the

effects of exogenous and endogenous attention on texture segmentation. A preferential high-SF

enhancement reproduces the impairments by exogenous attention due to a shift in visual

sensitivity toward details too fine to distinguish the target at foveal locations. The transition from

impairments to improvements in the periphery results from exogenous attentional gain gradually

shifting to lower SFs that are more amenable for target detection. Improvements by endogenous

attention result from a uniform enhancement of SFs that encompass the target, optimizing visual

sensitivity for the attended stimulus across eccentricity.

124
Figure 1. Signatures of texture segmentation.
(**A**) Central performance drop. Shaded region depicts the magnitude of the central performance
drop. Identical axis labels are omitted in panels B and C.
(**B**) Exogenous attention modulation. Exogenous attention improves segmentation performance in
the periphery and impairs it near the fovea.
(**C**) Endogenous attention modulation. Endogenous attention improves segmentation performance
across eccentricity.

RESULTS

Image-computable model of attention and spatial resolution

We developed an observer model based on established principles of neural computation (51, 52), pattern (53, 54) and texture vision (39, 41, 47) and attentional modulation (15). The model incorporates elements of the Reynolds-Heeger normalization model of attention (NMA) (15) and illuminates how attention alters contrast and texture sensitivity across SF and eccentricity. We implement: (i) SF-tuned gain modulation to emulate the decline in contrast sensitivity and peak SF preference with eccentricity. (ii) Spatial summation of normalized inputs to generate texture selectivity. (iii) Separate attentional gain profiles across SF to reproduce effects of exogenous and endogenous attention. The model is composed of four components: stimulus drive, attentional gain, suppressive drive and spatial summation (**Figure 2A**). Following NMA, attention adjusts the gain on the stimulus drive before normalization. For a full description of the model, see **Methods**.

Stimulus drive. We simulate bottom-up responses of a collection of linear receptive fields (RFs), each jointly tuned to spatial position, SF and orientation. Images are processed through a filter bank (55) covering the visual field at several SFs and orientations using bandwidths compatible with neurophysiological (54) and psychophysical (53) measurements. Filter outputs are combined across quadrature phase (56), yielding contrast energy images corresponding to different SFs and orientations. These outputs simulate the responses of complex cells in primary visual cortex (54, 56). The gain on individual RFs varies as a function of SF and eccentricity preference (**Figure 2A, green**). Following the behavior of individual neurons (54) and pattern vision (53), gain modulation is narrowly tuned to high SFs near the fovea and progressively shifts to low SFs with eccentricity. Consequently, the stimulus drive reflects local spectral energy within each patch in an image, filtered through feature-selective RFs that vary with eccentricity.

Attentional gain. Attention is implemented as a gain control mechanism that scales the gain on the stimulus drive (15). The magnitude of attentional gain is largest at the cued location (**Figure 2A, orange**) and varies with the eccentricity and SF preference of each RF. Motivated by findings of psychophysical experiments that manipulated endogenous and exogenous attention (26), two SF-tuned profiles are assessed—narrow and broad. The narrow profile selectively enhances a small range of SFs at each eccentricity (**Figure 2A, blue**); the broad profile uniformly enhances SFs (**Figure 2A, red**).

Suppressive drive. Suppression operates via divisive normalization (51, 52). Normalized responses

are proportional to the attention-scaled stimulus drive divided by a normalization pool plus a

constant σ^2 that increases with eccentricity. This constant adjusts the model's overall sensitivity to

contrast (i.e., contrast gain; **Figure 2A, black**). The normalization pool consists of the attention-

scaled stimulus drive across nearby spatial locations (surround suppression (57)), uniformly across

orientation (cross-orientation suppression (58)) and across preferred and neighboring SFs (cross-

frequency suppression (59)) of individual RFs. Such broad suppressive pools are supported by

physiological (57, 58, 60) and psychophysical (59, 61, 62) findings and models of visual processing

(51).

Spatial summation. Normalized responses are weighted and summed across space within each SF

and orientation filter. Spatial summation followed normalization (63), which accentuated texture-

defined contours within the image. The size of pooling regions scale with the SF preference of

each RF (39, 41) (**Figure 2A, purple**); larger for low than for high SFs. This implements an inverse

relation between the integration area of individual RFs and their SF tuning.

Target discriminability. The model generated measures of discriminability (d′) in a texture

segmentation task (**Figure 2B**). The model generated population responses to two texture images.

One contained a target patch whose orientation differed from its surround (target-present) and the

other consisted of uniform orientation throughout (target-absent). The vector length (i.e., Euclidean

norm) of the difference between population responses indexed d′. This measure is proportional to

behavioral performance, assuming the addition of normally distributed noise after normalization.

189
190
Figure 2. Image computable model of attention and spatial resolution
(**A**) Model structure. A filter bank of linear receptive fields decomposes an image. Filter responses
are squared and summed across quadrature-phase pairs (odd, even), yielding contrast energy
outputs. *SF gain* scales contrast energy across SF and eccentricity (green box). The solid black
line depicts the center frequency of the tuning function (f_{stim}); insets display the full SF tuning
function at a single eccentricity. The *stimulus drive* characterizes contrast energy at each pixel in
the image, filtered through feature-selective and eccentricity-dependent receptive fields. *Attentional*
gain multiplicatively scales the stimulus drive at a circumscribed region within the image (orange
circle in left panel) and varies across SF and eccentricity. The center SF of attentional gain varies
with eccentricity (solid black lines in blue and red boxes). Across SF, attentional gain follows either
a narrow profile (blue box) or a broad profile (red box), each centered on a given frequency (f_{narrow}
or f_{broad}). The *suppressive drive* comprises the attention-scaled stimulus drive pooled across a local
neighborhood of positions, SFs and uniformly across orientation. Contrast gain, σ^2, adjusts
suppression magnitude across eccentricity. *Spatial summation* follows normalization (purple box)
and generates the *population response*. Pooling area varies inversely with SF tuning. Variables
displayed within the square brackets depict model parameters fit to behavior.
(**B**) Target discriminability. Population responses for texture images with (present) or without
(absent) a target patch are computed. The vector magnitude of their difference produces a metric
proportional to d′, assuming independent and identically distributed Gaussian output noise.

Texture stimuli, behavioral protocol and optimization strategy

Stimuli. Model parameters were constrained by data from ten published psychophysical experiments. Exogenous attention was manipulated in six (27, 29-32) (**Figure 3A-F**) and endogenous attention in four experiments (32, 33) (**Figure 3G-J**). In each experiment, observers distinguished a patch of one orientation embedded within a background of differing orientation at several possible eccentricities.

Behavioral protocol. Performance was typically measured with a two-interval forced choice protocol (**Figure 3K**). Observers maintained fixation at the display's center while viewing two intervals of texture stimuli, one of which randomly contained a target texture. Different pre-cues at their optimal timing manipulated exogenous or endogenous attention. Brief peripheral pre-cues manipulated exogenous attention and appeared before both intervals, but near the upcoming target location in the interval containing the target (27-32, 34). Symbolic pre-cues manipulated endogenous attention. Pre-cues appeared near fixation and indicated the target location in the target-present interval (32, 33). Attention effects were determined relative to a neutral condition, in which observers distributed attention across all possible target locations. Behavioral performance displayed the three signatures of texture segmentation: (i) The CPD emerged in the neutral condition (**Figure 1A**). (ii) Peripheral pre-cues improved performance in the periphery and impaired it at foveal locations (**Figure 1B**). (iii) Central, symbolic pre-cues improved performance at all eccentricities (**Figure 1C**).

Optimization. To identify the computations that underlie each signature, we separately fit the model to three subsets of behavioral data. First, the CPD was isolated from attentional effects by fitting to the neutral condition from all ten experiments. Second, exogenous attentional effects were assessed by fitting to neutral and peripheral cueing conditions from the six exogenous attention experiments. Third, endogenous attentional effects were assessed by fitting to neutral and central cueing conditions from the four endogenous attention experiments. The model was jointly fit to each subset of data, with model parameters shared among experiments within a subset (**Table S2-S4**).

Figure 3. Texture stimuli and a typical texture segmentation behavioral protocol.
Target-present texture stimuli used in (**A-F**) exogenous attention and (**G-J**) endogenous attention
experiments, displayed at their respective spatial scales. Textures displayed include:
(**A**) Fine and (**B**) coarse-scale textures used in Yeshurun & Carrasco, 1998 (27); (**C**) Talgar &
Carrasco, 2002 (29) with targets placed on the vertical meridian; (**D**) Carrasco, Loula & Ho, 2006
(30) wherein observers discriminated the target's orientation; (**E**) Yeshurun & Carrasco, 2008 (31)
where the cue's size was manipulated; (**F**) Experiment 2 of Yeshurun, Montagna & Carrasco, 2008
(32) with targets placed on the horizontal meridian; (**G**) Experiment 1 of Yeshurun, Montagna &
Carrasco, 2008 (32) with targets placed on the horizontal meridian; (**H**) Experiment 3 and (**I**)
Experiment 4 of Yeshurun, Montagna & Carrasco, 2008 (32) wherein fine and coarse-scale
textures were displayed, respectively; and (**J**) Barbot & Carrasco, 2017 (33) with targets placed on
the intercardinal meridians.
(**K**) Two-interval forced choice protocol typically used to assess texture segmentation performance.
EXO corresponds to exogenous attention and ENDO to endogenous attention. Numbers denote
the representative timing information for each pre-cue—peripheral (blue) and central (red)—and
their corresponding inter-stimulus intervals (ISI). Neutral pre-cues equally distributed attention to all
possible target locations. Valid peripheral pre-cues appeared near the upcoming target location
whereas valid central pre-cues symbolically indicated the upcoming target location. In the
displayed example, the number "3" and the adjacent line indicate that the target would appear at a
peripheral eccentricity in the right visual hemifield.
259

Contextual modulation and spatial summation mediate the CPD

260
To identify the computations mediating the CPD, we fit the model to group-average performance
across all experiments' neutral condition (103 data points). 15 model parameters constrained
performance (**Table S2**). To account for differences in contrast sensitivity due to variable display
properties among experiments (e.g., mean luminance), foveal contrast gain (g_σ; **Figure 2A**) was
independently determined for each of ten experiments (10 parameters). Two separate parameters
determined foveal SF preference (t_T)–one shared among exogenous attention studies and another
among endogenous attention studies. The remaining three parameters–SF bandwidth (b_T), the
gradual increase in contrast gain (m_σ) and the progressive shift to lower SFs with eccentricity (m_T)–
were shared among all experiments. Attentional gain was not included for these fits.
270
The model reproduced the CPD and its dependence on texture scale (**Figure 4**). For a fine-scale
texture—characterized by narrow, densely spaced lines—performance peaked within the
parafovea (4 deg) and declined toward the fovea and periphery (**Figure 4A**). Differences between
target-present and target-absent stimuli were largest within the 2 cpd filter (**Figure 4A, middle**).
This filter best differentiated the target patch from a homogenous texture; we denote its center SF
as f_{fine}. A coarser texture was best distinguished by lower SFs (1 cpd, f_{coarse}), which exaggerated
the CPD, moving peak performance to a farther eccentricity (~6 deg; **Figure 4B**). The CPD was
well-fit in all experiments (**Figure 4C**); 77% of the variance was explained (95% bootstrapped CI =
[70 80]), with the best-fitting regression line falling close to the unity line.
280
Previous models qualitatively matched the CPD through spatial summation (46, 48, 49), but
ignored the contributions of contextual modulation via normalization. To assess the contribution of
each operation to behavior, we compared the full model to variants that either lacked components
of the suppressive drive (cross-orientation, cross-frequency, and/or surround suppression) or
spatial summation (**Figure 4D**). We restricted contextual modulation (-context) by separately
limiting the pool of orientations (-θ), SFs (-f), spatial positions (-x,y) or all simultaneously (-all) such
that suppressive modulations due to featural attributes and/or spatial positions outside each
receptive field's tuning were removed. The final variant lacked spatial summation (-sum), which
resulted in a population response that consisted of only normalized inputs. Removing spatial
summation attenuates the response to regions of similar orientation (e.g., target patch). Each
model was fit to behavioral performance in the neutral condition across all experiments and
compared using Akaike Information Criterion (AIC) (64) and Bayesian Information Criterion (BIC)
(65).
294

Removing contextual modulation or spatial summation attenuated the CPD (**Figure S1**). We

measured model performance relative to the full model, which yielded ΔAIC and ΔBIC scores;

positive values represent a decrease in model performance. We use "M" and "CI" to denote the

median and 95% confidence interval of the bootstrapped distribution. Model performance fell

without cross-orientation suppression (ΔAIC: M=4.8, CI=[-0.1 9.7]; ΔBIC: M=4.6, CI=[-0.2 9.6]),

cross-frequency suppression (ΔAIC: M=7.9, CI=[2.7 13.2]; ΔBIC: M=7.7, CI=[2.4 13.8]), surround

suppression (ΔAIC: M=5.4, CI=[0.03 11.0]; ΔBIC: M=5.4, CI=[-0.1 11.5]), and without all forms of

contextual modulation (ΔAIC: M=17.0, CI=[11.5 22.1]; ΔBIC: M=16.9, CI=[11.6 22.4]). Without

spatial summation, model performance decreased as well (ΔAIC: M=37.8, CI=[33.3 42.6]; ΔBIC:

M=37.8, CI=[33.1 42.8]). Thus, reliable reproduction of the CPD requires both contextual

modulation and spatial summation.

Figure 4. Contextual modulation and spatial summation mediate the CPD

(**A**) *Left*. Fit to Experiment 1 in Yeshurun & Carrasco, 1998 (27). Dots (n=18) and error bars depict

group-average performance and ±1 SEM. The black line and shaded regions depict the median

and 68% bootstrapped confidence interval of model fits. The gray vertical bar on the x-axis

indicates the eccentricity of peak performance. The inset shows the textures stimulus.

Middle. The matrix depicts the absolute value of differences between target-present and target-
absent population responses, normalized by the maximum across eccentricity and averaged
across orientation and space. f_{fine} denotes the SF filter with the largest difference between
population responses. We use absolute differences only to visualize the SFs that drove
discriminability.
Right. Spatial distribution of the absolute value of differences between target-present and target-
absent population responses. Each panel depicts a subset of receptive fields centered on the
fovea and tuned to one of three SFs (4, 2, 1 cpd) and an orientation of 30°.
(**B**) Fit to Experiment 2 (n=18) in Yeshurun & Carrasco, 1998 (27). The model jointly fits neutral
performance with parameters shared among all ten experiments, including the data shown in *A*.
Visualization follows the conventions in *A*. Note that eccentricity (x-axis) is twice that of *A*. f_{coarse}
denotes the SF filter that best distinguished the coarse-scale target.
(**C**) Goodness-of-fit for the neutral condition across ten experiments (n=103). Each dot depicts the
measured (x-axis) and predicted (y-axis) performance at a given eccentricity. The solid line and
shaded area depict the best-fitting regression line and its 95% confidence interval. The dashed line
indicates the unity line y=x.
(**D**) Model comparisons using AIC and BIC. Positive values indicate models underperforming,
relative to the full model. '-context' describes restrictions of contextual modulation: '-θ' denotes the
variant without cross-orientation suppression, '-f' without cross-frequency suppression, '-x,y'
without surround suppression and '-all' devoid of all contextual modulation. '-sum', denotes the
model variant without spatial summation. The dots and error bars denote the median and 95%
confidence interval of the bootstrap distribution.
336
337
338
Narrow high-SF enhancement generates exogenous attention effects

The model predicted behavior in neutral and peripheral cueing conditions across six experiments

(146 data points). Exogenous attention was modeled as a narrow SF gain profile (**Figure 2, blue**),

motivated by psychophysical measurements (26). 14 free parameters constrained model behavior

(**Table S3**). Model parameters that determined neutral cueing performance—foveal contrast gain

(g_σ), SF tuning (t_T), SF bandwidth (b_T), the increase in contrast gain (m_σ) and the decline in SF

preference with eccentricity (m_T)—were configured identically as described above. Four

parameters, shared among experiments, determined attentional gain–foveal SF preference (a_N),

the gradual shift to lower SFs with eccentricity (m_N), SF bandwidth (b_N) and amplitude (γ_N).

Consequently, attention operated identically on each texture stimulus, with the spatial spread of

attention fixed across experiments (see **Methods**).

The model reproduced the central impairments, peripheral improvements and their variation with

texture scale. For a fine-scale texture, the narrow SF profile yielded improvements within the

parafovea (4-12°), impairments across a small range of central eccentricities (0-2°) and shifted

peak performance toward the periphery (~6°; **Figure 5A**). For the coarser texture, the same

attention profile generated improvements in the periphery (8-22°), impairments within the

parafovea (0-8°) and shifted peak performance farther toward the periphery (~15°; **Figure 5B**).

A gradual shift of attentional gain toward lower SFs (26) reproduced the transition from

impairments to improvements across eccentricity (**Figure 5C**). At the fovea, attentional gain was

centered on a SF (4 cpd) higher than those distinguishing the fine- (2 cpd, f_{fine}) or coarse-scale (1

cpd, f_{coarse}) textures. As a result, the population response shifted away from the target and impaired

performance. With increasing eccentricity, attentional gain progressively overlapped the SF of each

target, improving performance. Attention enhanced the fine-scale target SF within the parafovea

(4-12°) then enhanced the coarse-scale target at farther eccentricities (8-22°). Overall, across the

six experiments, the model explained 77% of the variance (95% bootstrapped CI = [49 82]; **Figure**

5D).

Attentional gain on SFs higher than the target yielded impairments at foveal locations. This pattern

was consistent across all six experiments (**Figure 5E**). Consequently, the overlap between fine-

(f_{fine}) or coarse-scale (f_{coarse}) targets and the SF tuning of attentional gain was minimal at the fovea

and peaked in the periphery (**Figure 5F**). This mismatch between the SF tuning of attention (f_{narrow})

and the target is suggested to be driven by exogenous attention operating above intrinsic SF

preferences at each eccentricity (26). We corroborated this relation. We compared f_{narrow} to the

model's baseline SF tuning, indexed by the peak SF of the stimulus drive (f_{stim}, **Figure 2A**).

Consistent with empirical measurements, we found that the narrow SF profile preferred SFs higher

than baseline tuning (**Figure S2**).

Figure 5. Narrow high-SF enhancement generates exogenous attention effects
(**A**) Fit to Experiment 1 in Yeshurun & Carrasco, 1998 (27). The dots (n=36) depict group-average
performance and error bars denote ±1 SEM. The solid lines and shaded regions indicate the
median and 68% confidence intervals of the bootstrapped distribution of model fits. The vertical
blue bar on the x-axis indicates the eccentricity of peak performance with peripheral cues.
(**B**) Fit to Experiment 2 (n=36) in Yeshurun & Carrasco, 1998 (27). The model jointly fits
performance on neutral and peripheral cue conditions with parameters shared among all six
experiments, including the data shown in *A*. Visualization follows the conventions in *A*.
(**C**) Best-fitting narrow gain profile. The matrix depicts attentional gain across eccentricity,
normalized by the maximum and averaged across space and orientation. Matrix visualization and
the panels on the right follow the conventions of **Figure 4A**.
(**D**) Goodness-of-fit for neutral and peripheral-cued performance (n=146). Plotted as in **Figure 4C.**
(**E**) SF preference of the narrow attentional gain profile (f_{narrow}) and the SF that best distinguished
fine- (f_{fine}) and coarse-scale targets (f_{coarse}). The solid lines and shaded areas indicate the median
and 68% bootstrapped confidence interval. The shaded area for f_{coase} overlaps the solid line.
(**F**) Normalized magnitude of attentional gain on the fine- and coarse-scale target SF across
eccentricity (median and 68% confidence interval of bootstrapped distribution).

Broad SF enhancements yield endogenous attention effects

The model predicted group-average data from neutral and central cueing conditions across four experiments (60 data points). Endogenous attention was modeled as a broad gain profile (**Figure 2A, red**) (26). 12 free parameters constrained model behavior (**Table S4**). Four parameters, shared among experiments, determined attentional gain–foveal SF preference (a_B), the decline in SF preference with eccentricity (m_B), SF bandwidth (b_B) and amplitude (γ_B).

The model reproduced improvements across eccentricity for both fine- (**Figure 6A**) and coarse-scale textures (**Figure 6B**). To generate these improvements, attentional gain encompassed the target SF for each texture scale (**Figure 6C**). Across all four experiments, the model explained 89% of the variance (95% bootstrapped CI [67 92]; **Figure 6D**).

Endogenous attention effects were reproduced by a broad SF attentional gain that was centered near the target SF across eccentricity (f_{broad} in **Figure 6E**). This contrasts with the narrow SF gain profile that modulated higher SFs at central locations to reproduce exogenous attention effects (**Figure 5E**). Although the center SF of attention declined with eccentricity, the modulation profile's plateau ensured that it overlapped both fine- and coarse-scale target SFs across eccentricity (**Figure 6F**). Psychophysical measurements of attentional effects on contrast sensitivity (26) suggest that the SF range enhanced by endogenous attention is centered near those intrinsically preferred by an observer at each eccentricity. However, our model fits to texture segmentation experiments revealed that attentional gain enhanced lower SFs than baseline tuning (f_{stim}) at central locations (**Figure S3**).

Figure 6. Broad SF enhancements yield endogenous attention effects
(**A**) Fit to Experiment 3 in Yeshurun, Montagna & Carrasco, 2008 (32). The dots (n=18) depict group-average performance and error bars denote ±1 SEM. The solid lines and shaded regions indicate the median and 68% confidence intervals of the bootstrapped distribution of model fits. The vertical red bar on the x-axis indicates the eccentricity of peak performance with peripheral cues.
(**B**) Fit to Experiment 4 (n=18) in Yeshurun, Montagna & Carrasco, 2008 (32). The model jointly fits performance on neutral and central cue conditions with parameters shared among all four experiments, including the data shown in *A*. Visualization follows the conventions in *A*.
(**C**) Best-fitting broad gain profile. Plotted as in **Figure 5C**.
(**D**) Goodness-of-fit for neutral and central-cued performance (n=60). Plotted as in **Figure 4C.**
(**E**) SF preference of the broad attentional gain profile (f_{broad}) and the SF that best distinguished fine- (f_{fine}) and coarse-scale targets (f_{coarse}). The solid lines and shaded areas indicate the median and 68% bootstrapped confidence interval.
(**F**) Normalized magnitude of attentional gain on the fine- and coarse-scale target SF across eccentricity (median and 68% confidence interval of bootstrapped distribution).

Different SF gain profiles govern exogenous and endogenous attention effects

We directly assessed whether different SF gain profiles—narrow or broad—generate the effects of

exogenous and endogenous attention. In addition, we compared the efficacy of SF-tuned gain

against a model wherein the spatial extent of attention varied across experiments while the gain

across SF was uniform. The spatial spread of attention is a key factor of the NMA (15), which

posits that its extent relative to the stimulus size helps reconcile apparent discrepancies between

each attention type's effects on contrast sensitivity. These predictions have been empirically tested

and confirmed (66). By comparing the narrow and broad SF models to the spatial extent model, we

directly assessed the separate contributions of SF gain and the spatial spread of attention to

segmentation performance (**Figure 7**).

Tuned SF gain modulation reproduced the effects of attention. The spatial extent alone was

insufficient to capture the effects of either exogenous (ΔAIC: M=21.2, CI=[18.8 26.0]; ΔBIC:

M=31.7, CI=[27.9 34.9]; **Figure 7A**) or endogenous attention (ΔAIC: M=11.4, CI=[3.9 18.9]; ΔBIC:

M=13.5, CI=[5.7 20.8]; **Figure 7B**). For exogenous attention, the narrow profile outperformed the

broad profile (ΔAIC: M=39.1, CI=[35.5 42.5]; ΔBIC: M=39.1, CI=[35.9 42.5]; **Figure 7A**). For

endogenous attention, the broad profile outperformed the narrow profile (ΔAIC: M=25.4, CI=[17.8

32.7]; ΔBIC: M=25.5, CI=[18.0 32.7]; **Figure 7B**). Decrements in model performance manifested as

an inability to capture impairments or improvements at eccentricities demarcating the CPD (**Figure

S4**). Thus, these model comparisons substantiate psychophysical measurements (25, 26):

exogenous and endogenous attention effects are best explained by different attentional gain

profiles across SF.

Figure 7. Different SF gain profiles govern exogenous and endogenous attention effects.
AIC and BIC model comparisons for different regimes of attentional modulation for (**A**) exogenous
attention and (**B**) endogenous attention. The dots and error bars represent the median and 95%
confidence intervals of the bootstrap distributions.
468

A parsimonious explanation for several experimental manipulations in texture segmentation

Figure 8 depicts behavioral data for a variety of texture segmentation experiments. Whereas we

focus on the impact of texture scale in **Figure 5 and 6**, the model is general. It jointly accounted for

multiple target locations (vertical, **Figure 8A**; horizontal, **Figure 8C-E**; and intercardinal meridians,

Figure 8F), behavioral tasks (orientation discrimination, **Figure 8B**) and attentional manipulations

(cue size, **Figure 8C**). Although the model was fit using texture images with fixed positions and

orientations (**Figure 3**), it behaved similarly for textures with randomly jittered elements (**Figure**

S5). Overall, the proposed model provides a parsimonious explanation for and a quantitative match

to segmentation performance (**Figure 8**).

479
480
Figure 8. A parsimonious explanation for several experimental manipulations in texture
segmentation
(**A-D**) Narrow SF gain profile fit to exogenous attention experiments. The model jointly fits these
data and those displayed in **Figure 5**, with parameters shared among all six experiments. Insets in
each panel depict the same textures displayed in **Figures 3C-F**, respectively**. (E-F)** Broad SF gain
profile fit to endogenous attention experiments. The model jointly fits these data and those
displayed in **Figure 6**, with parameters shared among all four experiments. Insets in each panel
depict the same textures displayed in **Figure 3G and 3J**, respectively. The dots and error bars
depict group-average and ±1 SEM. The solid lines and shaded regions depict the median and 68%
confidence intervals of the bootstrapped distribution of model fits.

Model predictions generalize to basic visual tasks

To test whether this model generalizes to other basic visual tasks, we applied it to tasks mediated by acuity (6) and contrast sensitivity (26), with no additional model parameters (**Figure 9**). These studies separately manipulated exogenous and endogenous attention and highlight how attention effects depend on the stimulus and task. In the acuity task, observers discriminated the location of a small gap (<1°) in a Landolt square (**Figure S6A**) whereas contrast sensitivity was measured with gratings in an orientation discrimination task (**Figure S7A**).

The model reproduced the improvements to acuity and contrast sensitivity for each attention type. On the one hand, both exogenous and endogenous attention improve acuity similarly (6). Model simulations yielded consistent visual acuity improvements for both exogenous (**Figure 9A**) and endogenous (**Figure 9B**) attention, despite different SF gain profiles underlying each attention type. On the other hand, each type of attention alters contrast sensitivity across SF differently (26). Model simulations captured the differences between exogenous (**Figure 9C**) and endogenous attention (**Figure 9D**). The model reproduced the narrow SF bandwidth of exogenous attention that is centered on SFs higher than baseline tuning preferences (**Figure S7D**). It also captured the broad SF modulation by endogenous attention that spanned SFs above and below baseline tuning (**Figure S7E**). Attention effects derived from our observer model closely matched descriptive fits to the data from (26) (**Figure 9C-D**).

The attention parameters were consistent across tasks (**Table S6**). The SF bandwidth of endogenous attentional gain consistently spanned a larger range than exogenous attention (**Table S6**, SF bw). Moreover, the rate at which SF selectivity declined with eccentricity also differed. The peak SF decreased with eccentricity (**Table S6,** SF slope), but less so for exogenous than endogenous attention, indicating that exogenous attention consistently enhanced SFs higher than the peak SF of the stimulus drive (**Figure S2**). Lastly, we observed tradeoffs between the amplitude and spatial spread of attention (**Table S6**). In the acuity task, the amplitude was large (>8) and the spatial spread was narrower (0.6°) than the stimulus (1°), whereas in contrast sensitivity, the amplitude was lower (<1.5) and the spatial spread was broader (>5°) than the stimulus (4°). Texture segmentation yielded intermediate values wherein the amplitude was ~4 for a fixed spread of 4°. Independent of attentional effects, differences in the experimental protocol and stimuli used across experiments resulted in subtle differences in the best-fitting model parameters for contrast gain and the stimulus drive. Importantly, similar attention parameters reproduce endogenous and exogenous attention effects in a variety of visual tasks.

525
526
Figure 9. Model predictions generalize to other basic visual tasks. The effects of (**A**) exogenous
and (**B**) endogenous attention on gap discrimination thresholds in an acuity task. Data from (6).
Lower thresholds indicate higher acuity. Bars depict group-average thresholds in neutral and valid
cueing conditions. Error bars are ±1 SEM. Dots depict model-derived gap thresholds for the acuity
task. (**C**) Exogenous and (**D**) endogenous attention effects on contrast sensitivity across SF and
eccentricity, quantified as the ratio between valid and neutral contrast sensitivity. Data from (26).
Values above 1 indicate an attentional enhancement of contrast sensitivity. The dots and error bars
depict the group-average and ±1 SEM. The vertical black lines show baseline SF preferences
measured in the neural condition (**Figure S7**). The solid colored lines show model fits to the data
whereas lightly shaded lines are descriptive fits to the data from (26). In all panels, the narrow SF
profile was fit to exogenous attention effects whereas the broad SF profile was fit to endogenous
attention effects.
539

DISCUSSION

We used texture segmentation as a model system to dissociate endogenous and exogenous attention. To this end, we developed an image-computable model that reproduces human segmentation performance and the modulations by each attention type. This model links neural computations to three visual phenomena. (i) Divisive normalization and spatial summation mediate the CPD (27-34, 43-46). (ii) Narrow high-SF enhancement drives exogenous attentional effects (27-32, 34)*.* (iii) Broad SF gain drives endogenous attentional modulations (32-34).

Normalization models of attention have described how spatial attention affects neural responses and behavior (e.g., (14, 15, 17)). Our model adopts the same algorithm specified by the Reynolds-Heeger normalization model of attention (15) (NMA)—attentional gain modulates the stimulus drive before divisive normalization. Predictions by NMA have been empirically confirmed with psychophysical experiments (66). These experiments equated seemingly distinct effects of endogenous and exogenous attention on contrast sensitivity by manipulating and accounting for the spatial extent of attention.

Here, we demonstrate a critical limitation of extant models of attention. Their predictions do not extend to the differential effects on spatial resolution and do not explain the dissociation between endogenous and exogenous attention. Although the spatial extent of attention is critical for explaining effects on contrast sensitivity, our model comparisons demonstrate that it is not vital for reproducing attention effects on texture segmentation ('spatial extent' model in **Figure 7** and **Figure S4**). These results corroborate empirical evidence that manipulating the spread of attention during texture segmentation does not yield shifts between the typical effects of endogenous and exogenous attention (31).

To capture the effects of attention on texture segmentation we implemented: (i) Eccentricity-dependent and SF-tuned multiplicative gains that emulate neural (54) and psychophysical (53) SF selectivity. (ii) Spatial summation, which emphasizes textural contours (39, 41, 47). (iii) Distinct SF gain profiles for endogenous and exogenous attention (25, 26) that scale responses prior to normalization (15), thereby adjusting the balance between fine and coarse-scale visual sensitivity. The model's distinct SF profiles instantiate a computational dissociation between each attention type that substantiates their differential impact on sensory processing.

The necessity for different SF profiles is supported by empirical evidence (25, 26) and provides

insights toward the distinct roles of endogenous and exogenous attention in guiding visual

behavior. Previous models (e.g. (14, 15, 17)) demonstrate that both forms of attention improve low-

level visual processes that encode elementary features (e.g., contrast, orientation, motion). Here,

we show that attention differentially interacts with normalization to shape the competition inherent

in mid-level processes such as texture segmentation. Exogenous attention preferentially enhances

a narrow range of high SFs. Consequently, its effects prioritize fine-grained visual details at the

expense of competing coarse-scale features within a stimulus. In contrast, endogenous attention

consistently improves mid-level processing by broadly enhancing sensory encoding across fine

and coarse spatial scales. The computations underlying mid-level processing bridge the gap

between sensory encoding and object recognition (39-42). Therefore, the distinct impact by each

type of attention and their computational differences at this processing stage have broad

implications for natural visual behavior.

The model provides a computational framework for understanding the mechanisms underlying

established effects of exogenous attention on spatial resolution (27-34) (reviews (1-3)). Previous

studies offered qualitative descriptions that exogenous attention automatically increases spatial

resolution (27-32, 34) (reviews (1-3)) with concomitant costs in temporal resolution (22) attributed

to an engagement of parvocellular neurons (22, 67). Here, we develop an observer model that

anchors these qualitative descriptions onto established neural computations. In doing so, we

corroborate previous psychophysical experiments that found a similar high-SF preference of

exogenous attention (25, 26, 30, 68), specify how attentional gain changes across the visual field

and demonstrate its computational validity for explaining effects on perception.

We also provide converging evidence that exogenous attention alters perception inflexibly. By

comparing the model's exogenous attentional gain on textures to empirical measurements made

with gratings (26), we found that it consistently operates above intrinsic (i.e., baseline) SF

preferences despite large differences in stimuli (**Figure S2**). These findings suggest that in addition

to exogenous attentional effects being invariant to cue validity (8) and sometimes detrimental to

perception (27-32, 34), its operating range across SF is also invariant to the type of stimulus being

attended.

The model provides insights on the mechanisms underlying endogenous attention effects on

spatial resolution. Previous research has established that endogenous attention modulates texture

segmentation (18, 32-34, 69) and its impact has been described as an optimization of spatial

resolution (reviews (1-3)). We propose that a broad SF gain control mechanism yields these
perceptual improvements. Our proposal complements previous reports that endogenous attention
uniformly excludes noise across SF (70), but seemingly conflicts with an earlier explanation that
endogenous attention suppresses sensitivity to high SFs to improve texture segmentation (33).
However, suppressed high-SF sensitivity at foveal locations would decrease cross-frequency
suppression (59, 61) and result in an effective dominance of lower SFs, which is compatible with
our findings (**Figure S3**).
616
617   Moreover, we provide converging evidence of the flexibility of endogenous attention. We found that
618   the model's SF preference during texture segmentation differed from those measured with gratings
619   (26). This discrepancy suggests that the impact of endogenous attention depends on the
620   properties of the attended stimulus and the nature of the task, consistent with the notion of a
621   flexible endogenous attentional mechanism (8, 32-34).
622
The effects of attention depend on divisive normalization. Without normalization, we could not
reliably capture the CPD, which served as the foundation of our analyses. Previous studies
demonstrate that when the pool of SFs contributing to normalization is restricted, the CPD is
attenuated (30, 33, 44). However, existing models of the CPD (46, 48, 49) relate the phenomenon
solely to an increase in receptive field size with eccentricity. Our model directly links the summation
area of receptive fields to their SF tuning. Consequently, the dominant summation area increases
with eccentricity as SF preferences decrease. Despite implementing an increase in receptive field
size, we could not capture the CPD without accounting for the surrounding context via
normalization.
632
Additionally, we demonstrate that spatial constraints mediate the CPD independently from
limitations in temporal processing across eccentricity. The proposal that the CPD may result from
slow information accrual at the fovea, which yields poor performance particularly when a backward
mask limits processing time (43), has been criticized (45, 46, 71). We note that our model accounts
equally well for the findings of texture segmentation studies regardless of whether they contained
or omitted a mask, which minimized temporal contributions to task performance (**Table S5**).
Importantly, both endogenous and exogenous attention speed information accrual (72) across the
visual field (73, 74) and across different levels of cue validity (8). Thus, effects of attention on
temporal processing would predict similar improvements by each attention type on the CPD, a
prediction clearly contradicted by the modeled studies here (27, 29-33).
643

The computations implemented in the model are based on the known properties of the human and non-human primate visual system. The stimulus drive simulates bottom-up responses of phase-invariant complex cells in V1 (56) that vary with SF and eccentricity (53, 54). The model's response to texture is generated through pooling bottom-up inputs, consistent with the gradual emergence of texture selectivity along the visual hierarchy (75-77).

Exogenous attentional gain in the model result in changes to texture sensitivity; however, little is known about the neural underpinnings of these effects. There are sparse demonstrations of exogenous attentional modulations in visuo-occipital areas and beyond (20, 21, 78-80). Transcranial magnetic stimulation of early visual cortex reveals that its activity plays a key role in the generation of exogenous attention effects (81). However, future studies are required to determine how the SF gain modulation we report manifests in neural populations.

In contrast, it is established that endogenous attention modulates cortical responses (1, 2, 13, 18, 20, 21, 36-38, 82, 83). During texture segmentation tasks, endogenous attention selectively enhances V1 and V4 responses to the embedded figure, suggesting that attention spreads across the target object to facilitate its segmentation (18). Our model provides complementary evidence that endogenous attention optimizes SF sensitivity to improve segmentation across texture scale. Yet, it is unclear how neural activity generates these SF modulations. Neuroimaging (37, 38) and electrophysiological (13, 36) recordings demonstrate that spatial tuning profiles are altered by endogenous attention. Such changes are consistent with, but not necessary for, the modulations of spatial resolution we report.

Few computational models have implemented possible ways in which attention alters spatial resolution. Some have proposed that attention modifies how finely a spatial region is analyzed. Such changes are either driven by an attention field that adjusts the spatial profile of receptive fields (13) or by attracting receptive fields toward and contracting them around the attended location (19). Other models suggest an attentional prioritization that selectively tunes responses for a given spatial location and attenuates responses to surrounding regions (12, 16). However, these models neither account for differences across eccentricity nor explain attentional shifts toward fine or coarse spatial scales. Critically, these models do not distinguish between endogenous and exogenous attention. In contrast to these previous models, we do not propose any modifications to the structure of receptive fields. Instead, we attribute changes in spatial resolution to modulations of SF, a fundamental dimension of early visual processing.

The fact that our model operates on arbitrary images facilitates its generalization to other visual

stimuli and tasks. We show that the model reproduces the differential endogenous and exogenous

attention effects on contrast sensitivity (**Figure 9C-D**). Notably, the model recreates behavior in

visual acuity tasks where the improvements by each attention type are similar (**Figure 9A-B**).

Unlike texture segmentation, acuity tasks always benefit from heightened spatial resolution, which

obscures differences between these two attention types. Recent studies that compared both

attention types head-to-head with the same observers, stimuli and task found that they produced

similar behavioral effects but modulated neural activity differently in the temporo-parietal junction

(20) and occipital cortex (21). Our model is consistent with these findings and highlights that

differences in the underlying computations can yield similar perceptual effects between

endogenous and exogenous attention depending on the stimulus and task.

Future work may extend the model to other visual phenomena. For instance, it could capture the

differential effects by each attention type on second-order texture perception (28, 34), second-

order texture contrast sensitivity (24) and temporal resolution (22, 23, 67). Lastly, it is unknown

how interactions between both forms of attention may affect mid-level processes like texture

segmentation. Endogenous attention attenuates the transient effects of exogenous attention on

stimulus discriminability when both are deployed concurrently (84). Therefore, it is possible that

endogenous attentional benefits will outweigh the costs induced by exogenous attention when both

are deployed simultaneously during texture segmentation. Although the experimental designs of

the studies we have modeled cannot address this open question, our model framework may

facilitate predictions of the perceptual consequences when both forms of attention are deployed.

In conclusion, we reproduce signatures of texture segmentation (27-34, 43-46) and characterize

the contributions of attention to a process commonly considered 'preattentive' (39, 41, 42, 44-49).

Moreover, we reveal the neural computations that underlie how attention modifies spatial resolution

(1-3). Attention scales sensitivity to high and/or low SFs, adjusting the balance between fine and

coarse-scale spatial resolution. Exogenous attention preferentially enhances fine details whereas

endogenous attention uniformly enhances fine and coarse features to optimize task performance.

Because the model distinguishes between endogenous and exogenous attention, varies with

stimulus eccentricity, flexibly implements psychophysical tasks and operates on arbitrary grayscale

images, it provides a general-purpose tool for assessing theories of vision and attention across the

visual field.

METHODS
713
714 **Model**
715
716 We developed an observer model that simulates the response of a collection of receptive fields
717 (RFs) each narrowly tuned to spatial position (x,y), orientation (θ) and SF (f). Responses varied
718 with eccentricity (α). The population response (R) is generated by four components: the stimulus
719 drive (E), attentional gain (A), suppressive drive (S and σ), and spatial summation (F), where *
720 represents convolution:
721

$$R(f, \theta, x, y) = \frac{E(f, \theta, x, y)A(f, \alpha)}{\sigma^2(\alpha) + S(f, \theta, x, y)} * F(f) \tag{1}$$

722
723 All model parameters are given in **Table S1**.
724
725 <u>Stimulus drive</u>
726 The stimulus drive characterizes responses of linear RFs in the absence of suppression, attention
727 and spatial summation. A steerable pyramid (55) decomposed stimulus images into several SF and
728 orientation subbands, defined by weighted sums of the image (i.e., linear filters). Weights were
729 parameterized by raised-cosine functions that evenly tiled SFs, orientations and positions.
730
The number of SF and orientation subbands are parameters that can be flexibly chosen. We used
a set of 30 subbands comprising five SF bands and six orientation bands. The size of the stimulus
image and the subband bandwidth determine the total number of SF subbands. In our simulations,
images were 160 x 160 pixels (see **Stimulus generation**) and SF bandwidth (i.e., full-width at half-
maximum, FWHM) was 1 octave, which allowed for five different SF subbands. The chosen
bandwidth is comparable to empirical tuning curves measured in primate electrophysiological
recordings (85) and human psychophysical (53) measurements. The FWHM orientation bandwidth
(60°) is comparable to physiological tuning curves measured in primates (86). Using narrower (30°)
or wider (90°) bandwidths yielded similar results supporting the same conclusions.
740
741 The pyramid includes RFs in quadrature phase. We computed a 'contrast energy' response (56),
742 (i.e., the sum of squared responses across phase) which depends on the local spectral energy at
743 each SF, orientation and position in the image. Contrast energy is fundamental to texture
744 perception models (39, 41, 47) and we denote it as $C(f, \theta, x, y)$.
745

746 *SF gain*. Human (26, 53, 87) and non-human primate (54) contrast sensitivity is narrowly tuned to

747 SF. SF tuning shifts from high to low SFs with eccentricity. To model this behavior, contrast energy

748 was multiplied point-by-point by a SF gain function, *T*, defined by a log-parabola (88, 89):

$$T(f, \alpha) = \exp(-[\frac{\log_2(\frac{f}{\lambda_T(\alpha)})}{b_T}]^2) \qquad (2)$$

751 where α denotes the eccentricity of a RF and b_T determines the function's SF bandwidth. The

752 preferred SF (λ_T) at a given eccentricity is given by:

$$\lambda_T(\alpha) = 2^{t_T - m_T \alpha} + t_{min} \qquad (3)$$

SF preferences converge onto a single value in the far periphery, t_{min} (87). The preferred SF at

the fovea is given by $2^{t_T} + t_{min}$ and progressively shifts towards t_{min} at the rate m_T. Whereas t_T

varied during simulations (see **Table S1-S4**), t_{min} was fixed at 0.5 cpd because texture stimuli

produced minimal contrast energy below that SF subband. Allowing t_{min} to vary yielded similar

results supporting the same conclusions.

In sum, the stimulus drive (*E*) characterizes the contrast energy responses that vary with SF and

eccentricity, computed as:

$$E(f, \theta, x, y) = T(f, \alpha)C(f, \theta, x, y) \qquad (4)$$

764 <u>Attentional gain</u>

765 Attention is implemented as an attentional gain field, *A*, that multiplies the stimulus drive point-by-

766 point as in the Reynolds-Heeger normalization model of attention (15). Attentional gain was

767 uniform across orientation. Across SF and position, gain was distributed according to cosine

768 window functions, *w*:

$$w(z; \mu, b) = \begin{cases} 0.5 + 0.5\cos(\frac{\pi[z - \mu]}{b}) & \square \\ 0 & \mu - b < z > \mu + b \end{cases} \qquad (5)$$

where μ defined its center and *b* defined its FWHM. The units of μ and *z* depended on the

dimension: for SF each variable was in units of log$_2$-transformed cycles per degree and for position

they were in units of degrees of visual angle. The window was defined on a logarithmic axis for SF

but on a linear axis for position. SF and spatial position functions were multiplied, point-by-point, to
characterize the full distribution of attentional gain.

Spatial spread. Attentional gain was centered on the target location. In our simulations, the target
fell along the horizontal meridian at eccentricity α_{targ} (see **Stimulus generation**). The product of
two cosine functions (*w*, equation 5) defined the spread of attention: one varied as a function of *x*
and another as a function of *y*, each with an identical width b_{pos}. Widths did not vary across
eccentricity. A_{pos} defined the spatial spread of attention:

$$A_{pos}(x, y) = w(x; \alpha_{targ}, b_{pos})w(y; 0, b_{pos}) \tag{6}$$

The precise spatial spread of attention is controversial (90) and can change based on task
demands (66, 91). Critically, it has not been explicitly manipulated during texture segmentation
tasks by varying the target's spatial uncertainty. Such a protocol has been used to test predictions
of the NMA and has been demonstrated to adjust the size of the attention field (66). Instead, a
previous study (31) measured exogenous attention effects while manipulating the size of a
peripheral pre-cue. The authors found that exogenous attention altered performance as long as the
cue was the same or smaller than the target size. In our simulations, the spread of attention was
fixed at a FWHM of 4° (**Table S1**) because it encompassed the largest target size used to
constrain model parameters (**Table S5**). As a result, the spatial extent of attention was identical
across eccentricity and experiments. Similar results were observed when the spread was fixed at
2° and 3°. However, in the model variant wherein the spatial extent could change (see **Model**
alternatives), the FWHM of attentional spread (b_{pos}) was free to vary between experiments.

SF gain profile. We implemented two gain profiles: narrow and broad (26).

Narrow profile. In the narrow model (A_N), attentional gain was bandpass across SF. Attentional
gain peaked at a given SF, λ_N, and fell gradually toward neighboring frequencies within its
bandwidth, b_N, characterized by a cosine function:

$$A_N(f, \alpha) = w(f; \lambda_N(\alpha), b_N) \tag{7}$$

806   The center SF of attentional gain profiles (λ_N for narrow, λ_B for broad) varied with eccentricity:

$$\lambda_N(\alpha) = 2^{a_N - m_N \alpha} \tag{8}$$

809   where a_N (or a_B for the broad profile) defined the center frequency at the fovea, which gradually
810   changed with eccentricity at the rate m_N (m_B for broad).

812   *Broad profile*. The broad profile (A_B) implemented broadband attentional gain, characterized by the
813   sum of three overlapping cosine functions:

$$A_B(f, \alpha) = w_1 + w_2 + w_3 \tag{9}$$

where $w_1 = w(f; \lambda_B(\alpha), b_B)$, $w_2 = w(f; \lambda_B(\alpha) - b_B, b_B)$, and $w_3 = w(f; \lambda_B(\alpha) + b_B, b_B)$. The
bandwidth of each function was given by b_B. Relative to the center SF, λ_B, the adjacent functions
were centered $\pm b_B$ apart, ensuring that their sum yielded a plateau spanning b_B octaves and a
FWHM of $1.5 b_B$.

In sum, attentional gain multiplicatively scaled the stimulus drive uniformly across orientation, but
differently across SF and eccentricity given by:

$$A(f, \alpha) = \gamma_B A_{pos} A_B \tag{10}$$

where A_{pos} and A_B (or A_N) were four-dimensional matrices characterizing attentional gain across
position, SF and orientation. γ_B (or γ_N) defined attentional amplitude. To simulate the neutral cueing
condition, amplitude was set to 1. In addition, to assess the explanatory power of the spatial
spread of attention (see **Model alternatives**), A_B (or A_N) were set to 1 and only γ and A_{pos} varied.

<u>Suppressive drive</u>
The suppressive drive comprised contextual modulation, computed through pooling the attention-
scaled stimulus drive (15) across nearby positions, all orientations and neighboring SFs. This
pooling procedure implemented lateral interactions between RFs and was computed via
convolution (15). Convolution kernels were cosine window functions (*w*, equation 5).

The bandwidth of the SF kernel, δ_f, equaled 1 octave:

$$K_f = \begin{cases} 1 & f_i - \delta_f \leq f_i \leq f_i + \delta_f \\ 0 & otherwise \end{cases} \tag{11}$$

840   where f_i denotes the center SF of a subband. This kernel summed contrast energy within and ±1

841   octave around each SF subband.

843   The bandwidth of the orientation kernel, δ_θ, equaled 180°, which encompassed all orientation

844   subbands:

$$K_\theta = \begin{cases} 1 & \theta_i - \delta_\theta \leq \theta_i \leq \theta_i + \delta_\theta \\ 0 & otherwise \end{cases} \tag{12}$$

847   where θ_i denotes the center orientation of a steerable pyramid subband. This kernel summed

848   contrast energy across all orientations.

850   Spatial position kernels were determined by multiplying two cosine windows:

$$K_{pos}(x, y; f) = w(x; 0, \delta_{pos}) w(y; 0, \delta_{pos}) \tag{13}$$

853   One window varied across x, another across y and their centers traversed across the image during

854   convolution. The two-dimensional kernel summed to unity, which computed the average energy

855   within the pooled area. Kernel width, δ_{pos}, equaled $\frac{2}{f}$ and was inversely proportional to subband SF

856   f and yielded two-dimensional spatial kernels, K_{pos}. Kernel widths were identical across eccentricity.

858   Contextual modulation was characterized via separable convolution:

$$S(f, \theta, x, y) = K_f * (K_\theta * (K_{pos} * [E(f, \theta, x, y)A(f, \alpha)])) \tag{14}$$

861   where * denotes convolution of the suppression kernels, K. Suppression magnitude was adjusted

862   across eccentricity by σ^2, which controlled the level of contrast at which neural responses reached

863   half-maximum and is referred to as contrast gain. Contrast gain was implemented as an

864   exponential function across eccentricity (26, 87):

$$\sigma^2(\alpha) = 10^{2(g_\sigma - m_\sigma \alpha)} \tag{15}$$

where g_σ and m_σ are free parameters that determine contrast gain at the fovea and the rate at

which it varies with eccentricity, respectively.

<u>Spatial summation</u>

Following divisive normalization, responses were weighted and summed across space, within each

SF and orientation subband. Summation was accomplished via convolution by cosine windows, *F*,

computed using equation (13). The width of each filter scaled with SF: narrow (wide) regions of

space were pooled for high (low) SFs (39) and did not vary with eccentricity.

Decision mechanism

We used signal detection theory to relate population responses to behavioral performance (*d'*). The

available signal *s* was computed as the Euclidean norm of the difference between target-present

(**r**$_t$) and target-absent (**r**$_n$) neural population responses: $s = ||r_t - r_n||$. Performance on a

discrimination task is proportional to the neural responses given the assumption of additive,

independent and identically distributed (IID) noise. An alternative model with Poisson noise and a

maximum-likelihood decision rule yields the same linkage between neural response and behavioral

performance (92, 93). The signal and noise magnitude (σ_n) defined behavioral performance $d' =$

$\frac{s}{\sigma_n}$. $\sigma_n = \overline{r_{neutral}} \Big/ \overline{s_{neutral}}$ where $\overline{r_{neutral}}$ denotes the observed neutral performance averaged across

eccentricity and $\overline{s_{neutral}}$ denotes the eccentricity-average of the signal. This ratio scaled the

model's predicted behavioral performance to match the observed data.

Model fitting

Models were optimized by minimizing the residual sum of squared error between model and

behavioral *d'* using Bayesian adaptive direct search (BADS (94)). When applicable, performance

data for a psychophysical experiment were converted from proportion correct, *p*, to *d'* with the

assumption of no interval bias (95): $d' = \sqrt{2}z(p)$ where *z* denotes the inverse normal distribution.

Although performance on 2IFC tasks can exhibit biases between intervals (96), our conversion

algorithm operated uniformly across eccentricity, which preserved the performance variation (i.e.,

the CPD) critical for the goals of this study.

Optimization strategy

Central performance drop

We fit the model jointly to performance on the neutral condition of all 10 texture segmentation experiments (103 data points). Peripheral and central cueing conditions (for exogenous and endogenous attentional conditions, respectively) were excluded to isolate the CPD. 15 free parameters fit all 103 data points (**Table S2**). Ten separate free parameters independently controlled the minimum contrast gain at the fovea (g_σ; equation 15) for each of the 10 experiments. Sensitivity to contrast and SF varies for stimuli placed at isoeccentric locations around the visual field; it is higher at the horizontal meridian and decreases gradually towards the vertical meridian (97-100). Whereas 5 out of 6 exogenous attention experiments used targets placed on the horizontal meridian, 3 out of 4 endogenous attention experiments used targets presented along the intercardinal meridians (**Table S5**). Because SF selectivity depends on stimulus polar angle, two parameters separately determined the highest preferred SF (t_T; equation 3)—one shared among exogenous attention experiments and the other shared among endogenous attention experiments. Alternatively, we could have fit separate parameters for horizontal, vertical and intercardinal meridians. However, this approach would have added a third free parameter, reducing the parsimony of the model. The configuration we used yielded reasonably good fits.

The remaining three parameters were shared among all experiments. Each controlled the bandwidth (b_T; equation 2) of the tuning function T, the gradual shift toward lower SFs with eccentricity (m_T; equation 3) and the increase in contrast gain across eccentricity (m_σ; equation 15).

Attentional modulation

To generate the effects of attention, the model was fit separately to exogenous and endogenous attention experiments. We jointly fit the model to neutral and valid conditions of each experiment.

Exogenous attention. All six exogenous attention experiments were fit jointly (146 data points) with 14 free parameters (**Table S3**). Minimum contrast gain at the fovea (g_σ; equation 15) was determined independently for each of six experiments, yielding six free parameters. The remaining eight parameters were shared among all experiments. Four determined the stimulus drive: its bandwidth (b_T; equation 2), the highest preferred SF at the fovea (t_T; equation 3), the shift to lower SFs with eccentricity (m_T; equation 3) and the slope of contrast gain across eccentricity (m_σ; equation 15). The remaining four controlled the narrow SF attentional gain profile, specifically its bandwidth (b_N; equation 7), center SF (λ_N; equation 8), the shift to lower SFs with eccentricity (m_N; equation 8), and its amplitude (γ_N; equation 10).

Endogenous attention. All four endogenous attention experiments were fit jointly (60 data points) with 12 free parameters (**Table S4**). Minimum contrast gain at the fovea (g_σ; equation 15) was determined independently for each experiment, which yielded four free parameters. The remaining eight parameters were shared among experiments, as described above for exogenous attention.

Model alternatives

To assess whether contextual modulation and spatial summation are critical for the CPD, we implemented five model variants. Individual components of the suppressive drive were iteratively removed: cross-orientation suppression ('-θ'), cross-frequency suppression ('-f'), surround suppression ('-x,y') and all components simultaneously ('-all'). In a separate variant, spatial summation was removed ('-sum'). We fit each variant separately to neutral performance data from all ten psychophysical experiments using the configuration described in **Optimization strategy, Central performance drop**.

In the '-all' model, each RF was suppressed by its own response, simulating an extremely narrow suppressive pool. Specifically, the extent of suppressive pools (δ_f, δ_θ, δ_{pos}; equations 11-13) were set to 0. As a result, the contributions of surround, cross-orientation and cross-frequency suppression were absent. The other contextual modulation variants only had a single parameter set to 0 (e.g., δ_f for cross-frequency suppression). The '-sum' variant removed spatial summation (i.e., *F* in equation 1) from the model.

We additionally compared the efficacy of each attentional gain profile across SF—narrow or broad—in generating the effects of exogenous and endogenous attention by fitting each profile to exogenous and endogenous attention experiments. To assess the explanatory power of the spatial extent of attention, a third model was compared in which the spatial spread of attention (b_{pos}, equation 6) varied between experiments and the gain across SF was uniform. Each model fit followed the configurations described in **Parameter configuration, Attentional modulation**.

Model comparisons

We compared models using AIC (64) and BIC (65). The difference in AIC/BIC values between model variants indexed model performance. '-θ', '-f', '-x,y', '-all' and '-sum' models were compared to the full model. Additionally, narrow and broad SF gain profiles as well as the spatial extent model were compared.

Stimulus generation

Target-present and target-absent textures were re-created to match the stimulus parameters used in each psychophysical study (**Table S5**). For all stimuli, each pixel subtended 0.03125° (i.e., 32 pixels/°), roughly matching the spatial resolution of a 1280 × 960 monitor display placed 57 cm away from the observer.

The full texture stimulus used in each experiment typically spanned the entire display. We generated 5°-wide square cutouts of the texture stimulus, centered on the target location. Because the model implemented visual sensitivity that varied with eccentricity, but was uniform at isoeccentric locations, all targets were assumed to be presented along the horizontal eccentricity for simplicity (as in equation 6).

Each texture array was composed of lines oriented 135°. The target comprised a patch of lines that were oriented 45°. One study was an exception (30) because the texture array comprised vertical lines (0°) and the target patch contained lines tilted ±8° (**Figure 3D**). In this study, observers' performed an orientation discrimination task by reporting the orientation of the target presented on each trial. To simulate orientation discrimination performance, the target-present and target-absent stimuli always contained a patch but their orientation differed.

To avoid edge artifacts, texture stimuli were windowed by the sum of three cosine window functions (as in equation 8) centered on the target that produced a uniform plateau covering the central 3.75 deg and fell off with cosine edges. Pixel intensities in each stimulus were constrained between 0 and 1.

Textures used to fit the model were generated without spatial or orientation jittering. In additional simulations, the stimuli of two representative experiments were jittered. The stimuli for Experiment 1 in (27) were spatially jittered (0.3 deg jitter), and the stimuli in Experiment 4 in (32), were jittered spatially (0.34 deg jitter) and in orientation (55° bandwidth). Jitter parameters were compatible with those specified in each study.

Resampling procedures

We obtained confidence intervals on the parameter estimates, model predictions and AIC/BIC values by bootstrapping the data and refitting the model 100 times per configuration (**Optimization strategy**) and for each model variant (**Model alternatives**). Bootstrap samples were generated by

1002   drawing and fitting random samples from Gaussian distributions centered on group-average

1003   performance at a given eccentricity, with the SEM for each study defining the distribution's width.

1004

1005   To generate bootstrap samples for simulations with jittered texture stimuli, the model was first fit to

1006   the data for each experiment using a non-jittered texture. Then, the model parameters were fixed

1007   and jittered stimuli were input to the model. This procedure allowed us to assess how a fixed model

1008   behaved with variable texture inputs. One hundred unique jittered stimuli were presented to the

1009   model.

1010

Cross-validation procedure

To characterize how the operating range of exogenous and endogenous attention varied with

eccentricity, relative to baseline tuning preferences (**Figure S2-S3**), we fit polynomials to empirical

measurements made by (26). Leave-one-subject-out cross-validation determined the best-fitting

polynomial order. Specifically, the ratio, in octaves, between the peak SF of the neutral contrast

sensitivity function and the preferred SF of attentional modulation were computed for individual

observers. Eccentricities were aggregated between each of the two experiments conducted. The

ratio for one observer was set aside, and the remaining were averaged. Zero to second-order

polynomials were fit to the group-average ratio across eccentricities. The sum of squared error to

the left-out data point indexed cross-validation error. This process was iterated until each

observation was left-out once, resulting in 19 total iterations. The best-fitting polynomial order was

defined as one that produced the lowest median cross-validation error across all iterations.

1023

Model generalizability to basic visual tasks

We applied the same observer model to behavioral data from tasks mediated by acuity (6) and

contrast sensitivity (26). The model was configured identically to what is described in the **Model**

section above and the same model parameters were fit to behavioral data using BADS (94). To

simulate the Neutral condition, attentional gain was not included in the model. Narrow SF and

broad SF gain profiles were used to simulate all exogenous and endogenous attentional effects,

respectively.

1031

<u>Acuity</u>

The modeling strategy for the acuity task is outlined in **Figure S6**. Landolt squares were inputted to

the model with stimulus parameters that matched those described in (6). The squares were 1°-

wide Landolt squares with a line thickness of 0.05°. Images were padded with 0.5° of empty space

on each side to avoid edge artifacts. Model responses were computed for Landolt squares with a

small gap (<1°) on the top or bottom. The Eucledian norm of the difference between responses

indexed localization performance in the task. The model was evaluated at the eccentricity tested in

the experiment (9.375°) and at 10 linearly spaced gap sizes (0-30 arcmin). We characterized the

full psychometric function by interpolating between gap sizes. Interpolation was used to reduce

computational load; similar psychometric functions were generated when the model was evaluated

at finer intervals. The available signal for discrimination was scaled so that the maximum d'

equaled 2 and gap thresholds were quantified as the gap size needed to attain $d'=1$. For each

attention type, 10 free parameters were fit to 14 gap thresholds (7 observers x 2 cueing conditions

(Neutral, Valid)).

1046

<u>Contrast sensitivity</u>

The modeling strategy for the contrast sensitivity task is outlined in **Figure S7**. Tilted gratings

(±45°) were inputted to the model with stimulus parameters that matched those described in (26).

Gratings were windowed by a cosine function with a FWHM of 2°, had one of 6 SFs (0.5, 1, 2, 4

and 8 cpd) and were simulated at each of the four eccentricities tested (0°, 3°, 6° and 12°). We

omitted the highest SF tested in (26) because it fell outside the range of SF subbands (0.5-8 cpd)

used to simulate texture segmentation performance. Grating images were padded with 0.5° of

empty space on each side to avoid edge artifacts.

1055

To simulate the signal available to an observer in the orientation discrimination task, we computed

the Eucledian norm of the difference between orthogonal gratings. This procedure was repeated

for each grating SF and eccentricity. Model population responses were evaluated at 7 log-spaced

levels of contrast that were interpolated to characterize the full contrast response function (**Figure**

S7C). Similar contrast response functions were produced when the model was evaluated at finer

contrast steps. We scaled the available signal by the magnitude of internal noise to yield stimulus

discriminability (**Decision mechanism**). Because internal noise varies with SF (101), the available

signal was scaled such that the maximum d' at the fovea equaled 2 for each SF. Contrast

thresholds were then determined as the level of contrast required to reach $d'=1$ and their inverse

indexed contrast sensitivity. For each attention type, 10 free parameters were fit to 360 contrast

thresholds (9 observers x 2 cueing conditions (neutral, valid) x 4 eccentricities x 5 SFs).

1067

1068

1069

1070

1071

REFERENCES

1. M. Carrasco, Visual attention: The past 25 years. *Vision Research* **51**, 1484-1525 (2011).
2. K. Anton-Erxleben, M. Carrasco, Attentional enhancement of spatial resolution: linking behavioural and neurophysiological evidence. *Nature Reviews Neuroscience* **14**, 188-200 (2013).
3. M. Carrasco, A. Barbot, How attention affects spatial resolution. *Cold Spring Harbor Symposia on Quantitative Biology* **79**, 149-160 (2014).
4. Y. Yeshurun, M. Carrasco, Spatial attention improves performance in spatial resolution tasks. *Vision research* **39**, 293-306 (1999).
5. M. Carrasco, P. E. Williams, Y. Yeshurun, Covert attention increases spatial resolution with or without masks: Support for signal enhancement. *Journal of Vision* **2**, 4 (2002).
6. B. Montagna, F. Pestilli, M. Carrasco, Attention trades off spatial acuity. *Vision Research* **49**, 735-745 (2009).
7. M. Carrasco, Y. Yeshurun, The contribution of covert attention to the set-size and eccentricity effects in visual search. *Journal of Experimental Psychology: Human Perception and Performance* **24**, 673-692 (1998).
8. A. M. Giordano, B. McElree, M. Carrasco, On the automaticity and flexibility of covert attention: A speed-accuracy trade-off analysis. *Journal of Vision* **9**, 30-30 (2009).
9. M. Carrasco, S. Ling, S. Read, Attention alters appearance. *Nature neuroscience* **7**, 308-313 (2004).
10. S. Ling, M. Carrasco, Transient covert attention does alter appearance: A reply to Schneider (2006). *Perception & Psychophysics* **69**, 1051-1058 (2007).
11. T. Liu, J. Abrams, M. Carrasco, Voluntary attention enhances contrast appearance. *Psychological science* **20**, 354-362 (2009).
12. J. K. Tsotsos *et al.*, Modeling visual attention via selective tuning. *Artificial intelligence* **78**, 507-545 (1995).
13. T. Womelsdorf, K. Anton-Erxleben, S. Treue, Receptive Field Shift and Shrinkage in Macaque Middle Temporal Area through Attentional Gain Modulation. *Journal of Neuroscience* **28**, 8934-8944 (2008).
14. G. M. Boynton, A framework for describing the effects of attention on visual responses. *Vision research* **49**, 1129-1143 (2009).
15. J. H. Reynolds, D. J. Heeger, The Normalization Model of Attention. *Neuron* **61**, 168-185 (2009).
16. J. K. Tsotsos, *A computational perspective on visual attention* (MIT Press, 2011).
17. A. M. Ni, S. Ray, J. H. Maunsell, Tuned normalization explains the size of attention modulations. *Neuron* **73**, 803-813 (2012).
18. J. Poort *et al.*, The Role of Attention in Figure-Ground Segregation in Areas V1 and V4 of the Visual Cortex. *Neuron* **75**, 143-156 (2012).
19. O. Baruch, Y. Yeshurun, Attentional attraction of receptive fields can explain spatial and temporal effects of attention. *Visual Cognition* **22**, 704-736 (2014).
20. L. Dugué, E. P. Merriam, D. J. Heeger, M. Carrasco, Specific Visual Subregions of TPJ Mediate Reorienting of Spatial Attention. *Cerebral Cortex* **28**, 2375-2390 (2018).
21. L. Dugué, E. P. Merriam, D. J. Heeger, M. Carrasco, Differential impact of endogenous and exogenous attention on activity in human visual cortex. *Scientific Reports* **10**, 21274 (2020).
22. Y. Yeshurun, L. Levy, Transient spatial attention degrades temporal resolution. *Psychological Science* **14**, 225-231 (2003).
23. E. Hein, B. Rolke, R. Ulrich, Visual attention and temporal discrimination: Differential effects of automatic and voluntary cueing. *Visual Cognition* **13**, 29-50 (2006).
24. A. Barbot, M. S. Landy, M. Carrasco, Differential effects of exogenous and endogenous attention on second-order texture contrast sensitivity. *Journal of Vision* **12**, 6-6 (2012).

25. A. Fernandez, S. Okun, M. Carrasco, Differential effects of endogenous and exogenous
attention on sensory tuning. *bioRxiv* 10.1101/2021.04.03.438325 (2021).
26. M. Jigo, M. Carrasco, Differential impact of exogenous and endogenous attention on the
contrast sensitivity function across eccentricity. *Journal of Vision* **20**, 11 (2020).
27. Y. Yeshurun, M. Carrasco, Attention improves or impairs visual performance by enhancing
spatial resolution. *Nature* **396**, 72 (1998).
28. Y. Yeshurun, M. Carrasco, The locus of attentional effects in texture segmentation. *Nature*
Neuroscience **3**, 622-627 (2000).
29. C. P. Talgar, M. Carrasco, Vertical meridian asymmetry in spatial resolution: Visual and
attentional factors. *Psychonomic Bulletin & Review* **9**, 714-722 (2002).
30. M. Carrasco, F. Loula, Y.-X. Ho, How attention enhances spatial resolution: Evidence from
selective adaptation to spatial frequency. *Attention, Perception, & Psychophysics* **68**, 1004-
1012 (2006).
31. Y. Yeshurun, M. Carrasco, The effects of transient attention on spatial resolution and the
size of the attentional cue. *Perception & Psychophysics* **70**, 104-113 (2008).
32. Y. Yeshurun, B. Montagna, M. Carrasco, On the flexibility of sustained attention and its
effects on a texture segmentation task. *Vision Research* **48**, 80-95 (2008).
33. A. Barbot, M. Carrasco, Attention modifies spatial resolution according to task demands.
Psychological science **28**, 285-296 (2017).
34. M. Jigo, M. Carrasco, Attention alters spatial resolution by modulating second-order
processing. *Journal of Vision* **18**, 2 (2018).
35. J. Moran, R. Desimone, Selective attention gates visual processing in the extrastriate
cortex. *Science* **229**, 782-784 (1985).
36. T. Womelsdorf, K. Anton-Erxleben, F. Pieper, S. Treue, Dynamic shifts of visual receptive
fields in cortical area MT by spatial attention. *Nature Neuroscience* **9**, 1156-1160 (2006).
37. J. Fischer, D. Whitney, Attention narrows position tuning of population responses in V1.
Current Biology **19**, 1356-1361 (2009).
38. Barrie P. Klein, Ben M. Harvey, Serge O. Dumoulin, Attraction of position preference by
spatial attention throughout human visual cortex. *Neuron* **84**, 227-237 (2014).
39. M. S. Landy, N. Graham, "Visual Perception of Texture" in The Visual Neurosciences. (MIT
Press, Cambrige, MA, 2004), pp. 1106-1118.
40. P. R. Roelfsema, Cortical algorithms for perceptual grouping. *Annu. Rev. Neurosci.* **29**,
203-227 (2006).
41. M. S. Landy, "Texture analysis and perception" in The New Visual Neurosciences*,* J. S.
Werner, L. M. Chalupa, Eds. (MIT Press, Cambrige, MA, 2013), pp. 639-652.
42. J. D. Victor, M. M. Conte, C. F. Chubb, Textures as probes of visual processing. *Annual*
review of vision science **3**, 275-296 (2017).
43. L. Kehrer, Central performance drop on perceptual segregation tasks. *Spatial vision* **4**, 45-
62 (1989).
44. K. Morikawa, Central performance drop in texture segmentation: the role of spatial and
temporal factors. *Vision Research* **40**, 3517-3526 (2000).
45. C. Potechin, R. Gurnsey, Backward masking is not required to elicit the central performance
drop. *Spatial vision* **16**, 393-406 (2003).
46. R. Gurnsey, D. Di Lenardo, C. Potechin, Backward masking and the central performance
drop. *Vision Research* **44**, 2587-2596 (2004).
47. J. R. Bergen, E. H. Adelson, Early vision and texture perception. *Nature* **333**, 363-364
(1988).
48. L. Kehrer, The central performance drop in texture segmentation: A simulation based on a
spatial filter model. *Biological Cybernetics* **77**, 297-305 (1997).
49. L. Kehrer, C. Meinecke, A space-variant filter model of texture segregation: Parameter
adjustment guided by psychophysical data. *Biological Cybernetics* **88**, 183-200 (2003).

50. A. Thielscher, H. Neumann, A computational model to link psychophysics and cortical cell
activation patterns in human texture processing. *Journal of Computational Neuroscience*
22, 255-282 (2007).
51. D. J. Heeger, Normalization of cell responses in cat striate cortex. *Visual Neuroscience* **9**,
181-197 (1992).
52. M. Carandini, D. J. Heeger, Normalization as a canonical neural computation. *Nature*
Reviews Neuroscience 10.1038/nrn3136 (2011).
53. N. V. S. Graham, *Visual pattern analyzers*, Oxford psychology series (Oxford University
Press, New York, 1989), pp. 646.
54. R. L. DeValois, K. K. DeValois, *Spatial vision*, Oxford psychology series (Oxford Univ.
Press, New York, ed. 2nd ed, 1990), pp. 381.
55. E. P. Simoncelli, W. T. Freeman, E. H. Adelson, D. J. Heeger, Shiftable multiscale
transforms. *IEEE Transactions on Information Theory* **38**, 587-607 (1992).
56. E. H. Adelson, J. R. Bergen, Spatiotemporal energy models for the perception of motion.
Journal of the Optical Society of America A **2**, 284 (1985).
57. J. R. Cavanaugh, W. Bair, J. A. Movshon, Nature and interaction of signals from the
receptive field center and surround in macaque V1 neurons. *Journal of Neurophysiology* **88**,
2530-2546 (2002).
58. G. J. Brouwer, D. J. Heeger, Cross-orientation suppression in human visual cortex. *Journal*
of Neurophysiology **106**, 2108-2119 (2011).
59. Y. Petrov, M. Carandini, S. McKee, Two distinct mechanisms of suppression in human
vision. *Journal of Neuroscience* **25**, 8704-8707 (2005).
60. N. J. Priebe, D. Ferster, Mechanisms underlying cross-orientation suppression in cat visual
cortex. *Nature Neuroscience* **9**, 552-561 (2006).
61. D. Sagi, S. Hochstein, Lateral inhibition between spatially adjacent spatial-frequency
channels? *Perception & Psychophysics* **37**, 315-322 (1985).
62. T. S. Meese, D. J. Holmes, Spatial and temporal dependencies of cross-orientation
suppression in human vision. *Proc. R. Soc. B.* **274**, 127-136 (2007).
63. Z. M. Westrick, M. S. Landy, Pooling of first-order inputs in second-order vision. *Vision*
Research **91**, 108-117 (2013).
64. H. Akaike, A new look at the statistical model identification. *IEEE Trans. Automat. Contr.*
19, 716-723 (1974).
65. G. Schwarz, Estimating the dimension of a model. *Annals of statistics* **6**, 461-464 (1978).
66. K. Herrmann, L. Montaser-Kouhsari, M. Carrasco, D. J. Heeger, When size matters:
attention affects performance by contrast or response gain. *Nature Neuroscience* **13**, 1554-
1559 (2010).
67. Y. Yeshurun, Isoluminant stimuli and red background attenuate the effects of transient
spatial attention on temporal resolution. *Vision Research* **44**, 1375-1387 (2004).
68. N. Megna, F. Rocchi, S. Baldassi, Spatio-temporal templates of transient attention revealed
by classification images. *Vision Research* **54**, 39-48 (2012).
69. C. Casco, A. Grieco, G. Campana, M. P. Corvino, G. Caputo, Attention modulates
psychophysical and electrophysiological response to visual texture segmentation in
humans. *Vision research* **45**, 2384-2396 (2005).
70. Z.-L. Lu, B. A. Dosher, Spatial attention excludes external noise without changing the
spatial frequency tuning of the perceptual template. *Journal of Vision* **4**, 10 (2004).
71. R. Gurnsey, P. Pearson, D. Day, Texture segmentation along the horizontal meridian:
nonmonotonic changes in performance with eccentricity. *Journal of Experimental*
Psychology: Human Perception and Performance **22**, 738 (1996).
72. M. Carrasco, B. McElree, Covert attention accelerates the rate of visual information
processing. *Proceedings of the National Academy of Sciences* **98**, 5363-5367 (2001).
73. M. Carrasco, A. M. Giordano, B. McElree, Temporal performance fields: Visual and
attentional factors. *Vision research* **44**, 1351-1365 (2004).

74. M. Carrasco, A. M. Giordano, B. McElree, Attention speeds processing across eccentricity:
Feature and conjunction searches. *Vision research* **46**, 2028-2040 (2006).
75. J. Larsson, M. S. Landy, D. J. Heeger, Orientation-Selective Adaptation to First- and
Second-Order Patterns in Human Visual Cortex. *Journal of Neurophysiology* **95**, 862-881
(2006).
76. Y. El-Shamayleh, J. A. Movshon, Neuronal Responses to Texture-Defined Form in
Macaque Visual Area V2. *Journal of Neuroscience* **31**, 8543-8555 (2011).
77. L. E. Hallum, M. S. Landy, D. J. Heeger, Human primary visual cortex (V1) is selective for
second-order spatial frequency. *Journal of Neurophysiology* **105**, 2121-2131 (2011).
78. T. Liu, F. Pestilli, M. Carrasco, Transient attention enhances perceptual performance and
fMRI response in human visual cortex. *Neuron* **45**, 469-477 (2005).
79. L. Busse, S. Katzner, S. Treue, Temporal dynamics of neuronal modulation during
exogenous and endogenous shifts of visual attention in macaque area MT. *Proceedings of*
the National Academy of Sciences **105**, 16380-16385 (2008).
80. F. Wang, M. Chen, Y. Yan, L. Zhaoping, W. Li, Modulation of Neuronal Responses by
Exogenous Attention in Macaque Primary Visual Cortex. *Journal of Neuroscience* **35**,
13419-13429 (2015).
81. A. Fernández, M. Carrasco, Extinguishing Exogenous Attention via Transcranial Magnetic
Stimulation. *Current Biology* 10.1016/j.cub.2020.07.068 (2020).
82. J. H. Reynolds, L. Chelazzi, Attentional modulation of visual processing. *Annu. Rev.*
Neurosci. **27**, 611-647 (2004).
83. F. Pestilli, M. Carrasco, D. J. Heeger, J. L. Gardner, Attentional enhancement via selection
and pooling of early sensory responses in human visual cortex. *Neuron* **72**, 832-846 (2011).
84. M. A. Grubb, A. L. White, D. J. Heeger, M. Carrasco, Interactions between voluntary and
involuntary attention modulate the quality and temporal dynamics of visual processing.
Psychonomic Bulletin & Review **22**, 437-444 (2015).
85. R. L. De Valois, D. G. Albrecht, L. G. Thorell, Spatial frequency selectivity of cells in
macaque visual cortex. *Vision Research* **22**, 545-559 (1982).
86. D. L. Ringach, R. M. Shapley, M. J. Hawken, Orientation Selectivity in Macaque V1:
Diversity and Laminar Dependence. *The Journal of Neuroscience* **22**, 5639-5651 (2002).
87. J. S. Pointer, R. F. Hess, The contrast sensitivity gradient across the human visual field:
With emphasis on the low spatial frequency range. *Vision research* **29**, 1133-1151 (1989).
88. A. B. Watson, A. J. Ahumada, A standard model for foveal detection of spatial contrast.
Journal of Vision **5**, 6 (2005).
89. L. A. Lesmes, Bayesian adaptive estimation of the contrast sensitivity function: The quick
CSF method. *Journal of Vision* **10**, 1-21 (2010).
90. Y. Yeshurun, The spatial distribution of attention. *Current Opinion in Psychology* **29**, 76-81
(2019).
91. S. V. d. Stigchel *et al.*, The limits of top-down control of visual attention. *Acta Psychologica*
132, 201-212 (2009).
92. M. Jazayeri, J. A. Movshon, Optimal representation of sensory information by neural
populations. *Nature Neuroscience* **9**, 690-696 (2006).
93. F. Pestilli, S. Ling, M. Carrasco, A population-coding model of attention's influence on
contrast response: Estimating neural effects from psychophysical data. *Vision research* **49**,
1144-1153 (2009).
94. L. Acerbi, W. J. Ma, Practical Bayesian optimization for model fitting with Bayesian adaptive
direct search. *Advances in neural information processing systems* **30**, 1836-1846 (2017).
95. N. A. Macmillan, C. D. Creelman, *Detection theory: A user's guide* (Lawrence Erlbaum
Associates, Mahwah, NJ, 2005).
96. Y. Yeshurun, M. Carrasco, L. T. Maloney, Bias and sensitivity in two-interval forced choice
procedures: Tests of the difference model. *Vision Research* **48**, 1837-1851 (2008).

97.   M. Carrasco, C. P. Talgar, E. L. Cameron, Characterizing visual performance fields: Effects
of transient covert attention, spatial frequency, eccentricity, task and set size. *Spatial vision*
15, 61-75 (2001).
98.   E. L. Cameron, J. C. Tai, M. Carrasco, Covert attention affects the psychometric function of
contrast sensitivity. *Vision Research* **42**, 949-967 (2002).
99.   M. M. Himmelberg, J. Winawer, M. Carrasco, Stimulus-dependent contrast sensitivity
asymmetries around the visual field. *Journal of vision* **20**, 18-18 (2020).
100.   A. Barbot, S. Xue, M. Carrasco, Asymmetries in visual acuity around the visual field.
Journal of Vision **21**, 2-2 (2021).
101.   D. Silvestre, A. Arleo, R. Allard, Internal noise sources limiting contrast sensitivity. *Scientific*
Reports **8**, 2596 (2018).
1287

1292
1293 Michael Jigo[1][*], David J. Heeger[1,2] & Marisa Carrasco[1,2]
1294
1295 [1] Center for Neural Science and [2] Department of Psychology
1296 New York University, New York, NY 10003
1297
1298
1299
1300 [*] **Corresponding author information**:
1301 Michael Jigo (michael.jigo@nyu.edu)
1302

1303
1304
1305

Figure S1. Model variants fit to the neutral condition of all texture segmentation experiments.
Each row depicts the behavioral data from (**A**) Yeshurun & Carrasco, 1998 (27), Experiment 1; (**B**)
Yeshurun & Carrasco, 1998 (27), Experiment 2; (**C**) Talgar & Carrasco, 2002 (29); (**D**) Carrasco,
Loula & Ho, 2006 (30); (**E**) Yeshurun & Carrasco, 2008 (31); (**F**) Yeshurun, Montagna & Carrasco,
2008 (32), Experiment 2; (**G**) Yeshurun, Montagna & Carrasco, 2008 (32), Experiment 1; (**H**)
Yeshurun, Montagna & Carrasco, 2008 (32), Experiment 3; (**I**) Yeshurun, Montagna & Carrasco,
2008 (32), Experiment 4; (**J**) Barbot & Carrasco, 2017 (33). Each column shows the fit of different
model variants arranged in order of best-to-worst according to the model comparisons displayed in
Figure 4D: 'full' denotes the full model, '-θ' lacks cross-orientation suppression, '-x,y' lacks
surround suppression, '-f' lacks cross-frequency suppression, '-all' lacks all contextual modulation
and '-sum' lacks spatial summation. Dots and error bars denote group-average performance and
±1 SEM. The solid lines depict the median and shaded regions depict 68% confidence intervals of
the bootstrapped distribution of model fits. Values in top-right of each panel denote the median R^2
of the bootstrapped distribution of model fits. Negative R^2 values indicate a model fit that captures
less variance in the data than a horizontal line passing through the mean d′ across eccentricity.

1321
1322
Figure S2. Spatial frequency operating range of exogenous attention.
(**A**) Peak spatial frequency of baseline contrast sensitivity (CSF) and exogenous attentional
modulation from Jigo & Carrasco, 2020 (26). Estimates were based on human contrast sensitivity,
measured psychophysically with narrowband gratings. (**B**) Ratio (in octaves) of attentional and
baseline peak spatial frequency tuning across eccentricity. Positive values denote an attentional
preference for spatial frequencies higher than baseline. The solid line depicts the best-fitting
second-order polynomial (i.e., parabola). Polynomial order was determined using leave-one-
subject-out cross-validation (**Methods, Cross-validation procedure**). Dots in *A* and *B* depict
group-average and error bars depict ±1 SEM. (**C**) Peak spatial frequency of the stimulus drive
(stim) and the narrow SF attention gain profile (narrow). Estimates were derived from model fits to
texture segmentation performance across all six exogenous attention experiments. (**D**) Ratio of the
preferred spatial frequency for the stimulus drive and attentional gain. Solid lines indicate the
median and shaded areas denotes 68% confidence interval of bootstrapped distribution in *C* and
D.

1337
1338
Figure S3. Spatial frequency operating range of endogenous attention.
(**A**) The peak spatial frequency of baseline contrast sensitivity (CSF) and the center frequency of
broad endogenous attentional modulation from Jigo & Carrasco, 2020 (26). Estimates were based
on human contrast sensitivity, measured psychophysically with narrowband gratings. (**B**) Ratio (in
octaves) of attentional and baseline and spatial frequency preferences across eccentricity.
Negative values denote an attentional preference for spatial frequencies lower than baseline. The
solid line depicts the best-fitting zero-order polynomial (i.e., constant). Polynomial order was
determined using leave-one-subject-out cross-validation (**Methods, Cross-validation procedure**).
Dots in *A* and *B* depict group-average and error bars depict ±1 SEM (**C**) The center spatial
frequency of the stimulus drive (stim) and the broad attentional gain profile (broad). Estimates were
derived from model fits to texture segmentation performance across all six endogenous attention
experiments. (**D**) Ratio of the preferred spatial frequency for the stimulus drive and attentional gain.
Solid lines indicate the median and shaded areas denote 68% confidence intervals of the
bootstrapped distribution in *C* and *D*.

Figure S4. Attention model variants fit to behavioral data of all ten experiments. Each row depicts a different experiment: (**A**) Yeshurun & Carrasco, 1998 (27), Experiment 1; (**B**) Yeshurun & Carrasco, 1998 (27), Experiment 2; (**C**) Talgar & Carrasco, 2002 (29); (**D**) Carrasco, Loula & Ho, 2006 (30); (**E**) Yeshurun & Carrasco, 2008 (31); (**F**) Yeshurun, Montagna & Carrasco, 2008 (32), Experiment 2; (**G**) Yeshurun, Montagna & Carrasco, 2008 (32), Experiment 1; (**H**) Yeshurun, Montagna & Carrasco, 2008 (32), Experiment 3; (**I**) Yeshurun, Montagna & Carrasco, 2008 (32), Experiment 4; (**J**) Barbot & Carrasco, 2017 (33). Each column depicts a different attentional gain model. The numbers in the top-right of each panel denote the median R^2 of the bootstrapped distribution of model fits.

1364
1365
Figure S5. Model fits to jittered texture stimuli.
(**A**) Predicted performance for Experiment 1 in Yeshurun & Carrasco, 1998 (27) using texture
stimuli generated with line elements spatially jittered within the stimulus parameters of the
experiment (**Methods, Stimulus generation**). An example jittered stimulus is shown in the top-
right. Solid lines indicate the median and shaded regions depict 68% confidence intervals of the
bootstrap distributions of model predictions. Gray and blue ticks on x-axis indicate peak of
performance in the neutral and peripheral cueing condition, respectively. To generate the bootstrap
distributions, model parameters were fixed to those that jointly captured all exogenous
experiments. Then new jittered texture stimuli were input to the model on each iteration.
(**B**) Predicted performance for Experiment 4 in Yeshurun, Montagna & Carrasco, 2008 (32) using
texture stimuli generated with line elements whose orientation and spatial location were randomly
jittered within the parameters of the experiment (**Methods, Stimulus generation**). Solid lines
indicate the median and shaded regions depict 68% confidence intervals of bootstrap distributions
of model predictions. Gray and red ticks on x-axis indicate peak of performance in the neutral and
central cueing condition, respectively. To generate the bootstrap distribution, model parameters
were fixed to those that jointly captured all endogenous experiments, then new jittered texture
stimuli were input to the model on each iteration.
1383

1384
1385
Figure S6. Behavioral protocol and modeling strategy for an acuity task.
(**A**) Behavioral protocol adapted from (6). Observers performed a standard Landolt acuity task. The
gap size in each 1°-wide Landolt square varied on a trial-by-trial basis and gap thresholds were
measured in conditions where attention was distributed across both target locations (neutral) or
focused at a single location (valid). Peripheral cues manipulated exogenous attention (EXO)
whereas central, symbolic cues manipulated endogenous attention (ENDO). On each trial, two
Landolt squares appeared on one of the two main diagonals of the visual field at 9.375°. Observers
judged whether a gap appeared at the top or bottom of the Landolt square indicated by a response
cue displayed at the end of the trial. The response cue equated uncertainty of the target's location
between neutral and valid cueing conditions. Gap thresholds indexed visual acuity in each
condition. The timing information for peripheral (blue) and central (red) cueing conditions is given
below each trial segment. (**B**) We modeled localization performance in this task by computing the
discriminability (d′) between two Landolt squares, each with a gap at the top or bottom of the
stimulus. (**C**) Model-derived discriminability across a range of gap sizes allowed for the creation of
psychometric functions. To simulate neutral discriminability, attentional gain was not included in the
model. We modeled discriminability for valid conditions with the narrow SF profile for exogenous
attention and the broad SF profile for endogenous attention. Gap thresholds (vertical lines) were
quantified as the gap size that resulted in d′=1 (horizontal line). We fit the model's thresholds in
each cueing condition to the behavioral data shown in **Figure 9A-B**.
1405
1406

1407
1408
Figure S7. Behavioral protocol and modeling strategy for a contrast sensitivity task. (**A**) Behavioral
protocol adapted from (26). Observers performed an orientation discrimination task on 4°-wide
grating stimuli that varied in their contrast, SF and eccentricity. Grating contrast varied on a trial-by-
trial basis and contrast thresholds were measured in conditions where attention was distributed
across all possible target locations (neutral) or focused at a single location (valid). Peripheral cues
manipulated exogenous attention (EXO) whereas central, symbolic cues manipulated endogenous
attention (ENDO). On each trial, a single grating appeared along the horizontal meridian at 0°, 3°,
6° and 12° of eccentricity. The grating was tilted ±45° from vertical. After the onset of a response

cue, observers judged whether the grating was oriented clockwise (CW) or counter-clockwise (CCW) from vertical. The response cue equated uncertainty of the target's location between neutral and valid cueing conditions. The timing information for peripheral (blue) and central (red) cueing conditions is given below each trial segment. (**B**) We modeled orientation discrimination performance by computing the discriminability (d′) between two gratings, each tilted ±45° from vertical. Stimulus discriminability was simulated across a range of contrast levels for eccentricities and SFs tested in (26). (**C**) We simulated contrast response functions for each cueing condition. To simulate the neutral condition, attentional gain was not included in the model. Discriminability in the valid condition was modeled with the narrow SF profile for exogenous attention (i.e., peripheral cueing condition) and the broad SF profile for endogenous attention (i.e., central cueing condition). Contrast thresholds (vertical lines) were quantified as the level of contrast that resulted in d′=1 (horizontal line). The inverse of contrast thresholds indexed contrast sensitivity and were fit to the behavioral data. (**D**) Contrast sensitivity functions for neutral (left) and peripheral cueing conditions (right). The dots and error bars depict group-average contrast sensitivity and 68% confidence intervals for each eccentricity tested in (26). The solid lines are model fits to the behavioral data. (**E**) Contrast sensitivity functions for neutral (left) and central cueing conditions (right). Visualization conventions follow those in *D*. The vertical black lines in **Figure 9C-D** depict the peak SF of neutral contrast sensitivity functions, which indexed observers' baseline tuning preferences. The ratio between valid and neutral contrast sensitivity indexed attentional effects across SF, shown in **Figure 9C-D**.

Table S1. Model parameters. Bolded entries indicate model components.

Parameter	Description
Stimulus drive	
t_T	peak SF (log$_2$-cpd)
t_{min}	minimum preferred SF (cpd); *fixed at 0.5*
m_T	SF tuning change across eccentricity (octaves/°)
b_T	SF FWHM bandwidth (octaves)
Contrast gain	
g_σ	gain at fovea
m_σ	slope along eccentricity
Normalization pool	
δ_f	SF pool bandwidth (octaves); *fixed at 1*
δ_θ	Orientation pool bandwidth (°); *fixed at 180*
δ_{pos}	Spatial pool width (°); *fixed at $\frac{2}{f}$*
Spatial summation	
δ_{pos}	Summation pool width (°); *fixed at $\frac{2}{f}$*
Attentional gain profile	
a_N or a_B	attentional SF tuning at the fovea (log$_2$-cpd)
m_N or m_B	slope of SF tuning across eccentricity (octaves/°)
γ_N or γ_B	amplitude
b_N or b_B	SF FWHM bandwidth (octaves)
b_{pos}	spatial spread (°); *fixed at FWHM of 4*

Table S2. Free parameters for the fits to the neutral condition of all ten texture segmentation

experiments. The mapping between the experiment labels (a-j) and the respective references is

given below the table and are consistent across all tables; italicized text describe the manipulation

conducted in a given experiment. Bold values indicate the median and values within square

brackets denote the 95% CI of the bootstrapped distribution of fitted parameters. min. = minimum;

SF = spatial frequency; bw = bandwidth.

1445

		Contrast gain		Stimulus drive		
parameter		g_σ	m_σ	t_T	m_T	b_T
description		min.	slope	SF peak	SF slope	SF bw
Experiment	a	**2.3** [2.2 2.5]	**-0.09** [-0.09 -0.08]	**1.9** [1.5 2.4]	**-0.8** [-0.9 -0.6]	**1.7** [1.6 2.0]
	b	**2.3** [2.1 2.5]				
	c	**2.7** [2.5 2.8]				
	d	**2.7** [2.6 2.8]				
	e	**2.2** [2.0 2.5]				
	f	**2.7** [2.4 2.8]				
	g	**1.9** [1.5 2.3]		**2.8** [2.4 3.1]		
	h	**1.8** [1.7 2.0]				
	i	**2.6** [2.5 2.8]				
	j	**2.7** [2.6 2.8]				

1446
[a] Yeshurun & Carrasco, 1998 (27). *Fine-scale texture; experiment 1.*
[b] Yeshurun & Carrasco, 1998 (27). *Coarse-scale texture; experiment 2.*
[c] Talgar & Carrasco, 2002 (29). *Target meridian: lower vertical.*
[d] Carrasco, Loula & Ho, 2006 (30). *Orientation discrimination: baseline adaptation.*
[e] Yeshurun & Carrasco, 2008 (31). *Attentional cue size: cue size 1.*
[f] Yeshurun, Montagna & Carrasco, 2008 (32). *Target meridian: horizontal; experiment 2.*
[g] Yeshurun, Montagna & Carrasco, 2008 (32). *Target meridian: horizontal; experiment 1*
[h] Yeshurun, Montagna & Carrasco, 2008 (32). *Fine-scale texture; experiment 3.*
[i] Yeshurun, Montagna & Carrasco, 2008 (32). *Coarse-scale texture; experiment 4.*
[j] Barbot & Carrasco, 2017 (33). *Target meridian: intercardinal; baseline adaptation.*
1457

1458 **Table S3**. Free parameters for the fits to neutral and peripheral cueing conditions of the six

1459 exogenous attention experiments. Bold values indicate the median and values in square brackets

1460 depict the 95% CI of the bootstrapped distribution of fitted parameters. min. = minimum; bw =

1461 bandwidth; amp. = amplitude.

1462

		Contrast gain		Stimulus drive			Narrow gain profile			
parameter		g_σ	m_σ	t_T	m_T	b_T	a_N	m_N	b_N	γ_N
description		*min.*	*slope*	*SF peak*	*SF slope*	*SF bw*	*SF peak*	*SF slope*	*SF bw*	*amp.*
Experiment	a	**1.9** [1.7 2.2]								
	b	**1.7** [1.5 2.1]								
	c	**2.6** [2.2 2.8]	**-0.05** [-0.07 -0.04]	**2.1** [1.8 2.8]	**-0.7** [-0.9 -0.5]	**1.6** [1.4 2.1]	**2.3** [1.8 3.0]	**-0.2** [-0.3 -0.01]	**2.9** [1.5 4.8]	**4.3** [3.0 7.8]
	d	**2.1** [2.8 2.5]								
	e	**1.5** [1.5 2.0]								
	f	**2.0** [1.7 2.7]								

1463

1464 **Table S4**. Free parameters for the fits to the neutral and central cueing conditions of the four

1465 endogenous attention experiments. Bold values indicate the median and values within square

1466 brackets depict the 95% CI of the bootstrapped distribution of fitted parameters. min. = minimum;

1467 bw = bandwidth; amp. = amplitude.

1468

		Contrast gain		Stimulus drive			Broad gain profile			
parameter		g_σ	m_σ	t_T	m_T	b_T	a_B	m_B	b_B	γ_B
description		*min.*	*slope*	*SF peak*	*SF slope*	*SF bw*	*SF peak*	*SF slope*	*SF bw*	*amp.*
Experiment	g	**1.6** [1.5 2.1]	**-0.09** [-0.1 -0.06]	**3.4** [2.8 4.0]	**-0.8** [-1.0 -0.5]	**1.3** [1.1 1.6]	**1.7** [0.5 2.3]	**-0.2** [-0.3 -0.1]	**4.5** [2.1 6.6]	**3.1** [2.2 6.2]
	h	**1.7** [1.5 1.9]								
	i	**2.5** [2.2 2.8]								
	j	**1.7** [1.5 1.9]								

1469

1470 **Table S5**. Stimulus parameters for each texture segmentation experiment.
1471

Experiment	Line spacing horz × vert	Line size width × height	Target size width × height	Target meridian	Backward mask
a	0.68° × 0.71°	0.1° × 0.4°	1.97° × 2.03°	horizontal	yes
b	1.36° × 1.42°	0.2° × 0.8°	3.78° × 3.84°	horizontal	yes
c	0.71° × 0.68°	0.1° × 0.4°	2.03° × 1.97°	vertical	yes
d	1.43° × 1.37°	0.2° × 1°	4.09° × 3.97°	horizontal	yes
e	0.68° × 0.71°	0.1° ×0.7°	2.28° × 2.34°	horizontal	yes
f	0.68° × 0.71°	0.1° × 0.7°	2.28° × 2.34°	horizontal	yes
g	0.68° × 0.71°	0.1° × 0.7°	2.28° × 2.34°	horizontal	yes
h	0.46° × 0.46°	0.1° × 0.2°	1.34° × 1.34°	intercardinal	no
i	0.91° × 0.91°	0.2° × 0.4°	2.47° × 2.47°	intercardinal	no
j	0.4° × 0.4°	0.1° × 0.24°	1.03° × 1.84°	intercardinal	no

1472

1473 **Table S6**. Best-fitting parameters for acuity and contrast sensitivity tasks. min. = minimum; bw =
1474 bandwidth; amp. = amplitude; acuity = acuity experiment (6); CS = contrast sensitivity experiment
1475 (26). The 95% confidence intervals show parameter values for fits to texture segmentation
1476 experiments, split by attention type.
1477

		Contrast gain		Stimulus drive			Attentional gain				
parameter		g_σ	m_σ	t_T	m_T	b_T	a	m	b	γ	b_{pos}
description		*min.*	*slope*	*SF peak*	*SF slope*	*SF bw*	*SF peak*	*SF slope*	*SF bw*	*amp.*	*spread*
EXO	95% CI	[1.5 2.8]	[-0.07 -0.05]	[1.8 2.8]	[-0.9 -0.5]	[1.4 2.1]	[1.8 3.0]	[-0.3 -0.01]	[1.5 4.8]	[3.0 7.8]	-
	Acuity	2	-0.07	1.8	-0.8	1.4	2.8	-0.03	1.5	8	0.6
	CS	2.5	-0.03	1.1	-0.2	1.4	1.5	-0.01	2.3	1.5	5.2
ENDO	95% CI	[1.5 2.8]	[-0.1 -0.06]	[2.8 4.0]	[-1.0 -0.5]	[1.1 1.6]	[0.5 2.3]	[-0.3 -0.1]	[2.1 6.6]	[2.2 6.2]	-
	Acuity	1.5	-0.09	3	-0.9	1.1	2	-0.3	5.3	15.0	0.6
	CS	2.5	-0.04	1.1	-0.2	1.5	2.6	-0.2	4.2	1.2	5.2

1478