Connectional Asymmetry of the Inferior Parietal Lobule Shapes Hemispheric Specialization in

Humans, Chimpanzees, and Rhesus Macaques

Luqi Cheng[1,2,3], Yuanchao Zhang[1], Gang Li[2,3,4], Jiaojian Wang[1,5], Chet C. Sherwood[6], Gaolang

Gong[7,8], Linzhong Fan[2,3,4,9,*], Tianzi Jiang[1,2,3,4,9,*]

[1]Key Laboratory for NeuroInformation of Ministry of Education, School of Life Science and Technology, University

of Electronic Science and Technology of China, Chengdu 610054, China

[2]Brainnetome Center, Institute of Automation, Chinese Academy of Sciences, Beijing 100190, China

[3]National Laboratory of Pattern Recognition, Institute of Automation, Chinese Academy of Sciences, Beijing 100190,

China

[4]School of Artificial Intelligence, University of Chinese Academy of Sciences, Beijing 100190, China

[5]Center for Language and Brain, Shenzhen Institute of Neuroscience, Shenzhen 518057, China

[6]Department of Anthropology and Center for the Advanced Study of Human Paleobiology, The George Washington

University, Washington, DC 20052, USA

[7]State Key Laboratory of Cognitive Neuroscience and Learning & IDG/McGovern Institute for Brain Research,

Beijing Normal University, Beijing 100875, China

[8]Beijing Key Laboratory of Brain Imaging and Connectomics, Beijing Normal University, Beijing 100875, China

[9]CAS Center for Excellence in Brain Science and Intelligence Technology, Institute of Automation, Chinese

Academy of Sciences, Beijing 100190, China

Corresponding Author: **Tianzi Jiang**, Institute of Automation, Chinese Academy of Sciences, Beijing 100190,

China. Email: jiangtz@nlpr.ia.ac.cn, Phone: 010 - 82544778, Fax: 010 - 82544777.

Co-Corresponding Author: Lingzhong Fan, Institute of Automation, Chinese Academy of Sciences, Beijing

100190, China. Email: lingzhong.fan@ia.ac.cn, Phone: 010 - 82544523.

Abstract

The inferior parietal lobule (IPL) is one of the most expanded cortical regions in humans relative to other primates. It is also among the most structurally and functionally asymmetric regions in the human cerebral cortex. Whether the structural and connectional asymmetries of IPL subdivisions differ across primate species and how this relates to functional asymmetries remain unclear. We identified IPL subregions that exhibited positive allometric in both hemispheres, scaling across rhesus macaque monkeys, chimpanzees, and humans. The patterns of IPL subregions asymmetry were similar in chimpanzees and humans, but no IPL asymmetries were evident in macaques. Among the comparative sample of primates, humans showed the most widespread asymmetric connections in the frontal, parietal, and temporal cortices, constituting leftward asymmetric networks that may provide an anatomical basis for language and tool use. Unique human asymmetric connectivity between the IPL and primary motor cortex might be related to handedness. These findings suggest that structural and connectional asymmetries may underlie hemispheric specialization of the human brain.

Keywords: inferior parietal lobule; brain asymmetry; brain evolution; anatomical connectivity; parcellation

Introduction

The association cortex has expanded greatly in size and exhibits modified connectivity patterns in human brain evolution (Orban *et al.*, 2006; Mars *et al.*, 2017; Ardesch *et al.*, 2019; Van Essen *et al.*, 2019). Compared with the primary sensory and motor cortical regions, the association cortex displays disproportionate expansion in conjunction with overall neocortical volume enlargement across primates (Chaplin *et al.*, 2013). Accordingly, association areas comprise a large percentage of the neocortex in human brains (Orban *et al.*, 2006; Van Essen and Dierker, 2007; Donahue *et al.*, 2018). Functional and neuroanatomical asymmetries are also pronounced in the human brain, appearing to be more extreme compared with other primate species, especially in the association cortex (Chance and Crow, 2007). Nevertheless, cerebral asymmetry exists not only in humans but also in nonhuman primates (Gómez-Robles *et al.*, 2013; Hopkins, 2013). For example, olive baboons and chimpanzees showed population-level leftward volumetric asymmetry in the planum temporale, which is thought to be homologous to part of Wernicke's area in humans and may have played a facilitating role in the evolution of spoken language (Spocter *et al.*, 2010; Marie *et al.*, 2018). Comparative studies on brain asymmetry are crucial for understanding the evolution and function of the modern human brain.

Language and complex tool use, which show considerable lateralization in the human brain, are considered to be universal features of humans (Johnson-Frey *et al.*, 2005; Lewis, 2006; Binder *et al.*, 2009). These specialized functions all involve the inferior parietal lobule (IPL), an area of the association cortex that represents a zone of topographical convergence in the brain (Johnson-Frey, 2004; Binder *et al.*, 2009). Moreover, the IPL is one of the most expanded regions in humans compared with nonhuman primates (Orban *et al.*, 2006; Van Essen and Dierker, 2007; Kaas, 2012; Ardesch *et al.*, 2019). The functional diversity and expansion of the IPL imply that it contains subdivisions that may

have been elaborated or developed in the ancestors of modern humans, allowing new abilities such as

extensive tool use and communication using gestures (Kaas, 2012). However, due to the scarcity of

data, different criteria, and methodological limitations for defining regions or subregions (Mars *et al.*,

2017), whether the internal organization of the IPL differs across species and how this relates to

different asymmetric functions remain unclear.

A major challenge for neuroscience is to translate results obtained using one method and in one species

to other methods and other species. Although the IPL has been subdivided into distinct subregions

using cytoarchitecture and this technique has provided invaluable information, cellular microstructure

alone is insufficient to completely represent brain organization, especially long-range connections,

which are the major determinant of regional specialization (Passingham *et al.*, 2002; Caspers *et al.*,

2006). Furthermore, histological methods with postmortem brains cannot be readily scaled to large

populations. Recent advances in diffusion magnetic resonance imaging (MRI), which allow the

quantitative mapping of whole-brain neural connectivity *in vivo*, provide an alternative technique

called connectivity-based parcellation to subdivide specific regions of the brain or even the entire

cortex (Fan *et al.*, 2016; Eickhoff *et al.*, 2018). In previous studies, this technique was successfully

used to characterize IPL subdivisions in different species as well as to perform cross-species

comparisons (Mars *et al.*, 2011; Wang *et al.*, 2020).

Previous studies have assessed asymmetries of the IPL using local characteristics, such as cortical

volume, thickness, and surface area (Croxson *et al.*, 2018; Kong *et al.*, 2018). However, although such

regional asymmetries have been identified, additional analyses are needed to address the architecture

of neural connectivity (Ocklenburg *et al.*, 2016). A recent "connectomic hypothesis for the

hominization of the brain" suggests neural network organization as an intermediate anatomical and

functional phenotype between the genome and cognitive capacities, which are extensively modified in

the human brain (Changeux *et al.*, 2020). The functions and interactions of brain regions are

determined by their anatomical connections (Passingham *et al.*, 2002). Therefore, identifying

connectional asymmetries may provide new insights into the structural and functional specializations

of the human brain.

This study investigated asymmetries of IPL subregions in terms of both structure and anatomical

connectivity in rhesus macaques, chimpanzees, and humans. We first used connectivity-based

parcellations to subdivide the IPL to reveal consistent cross-species topographical organization. We

then investigated the volumetric allometric scaling and asymmetries of the IPL subregions across

species. Using vertex-, region of interest (ROI)-, and tract-wise analyses, we examined asymmetries

of the IPL subregions in terms of their connectivity profiles and subcortical white matter pathways to

identify evolutionary changes.

Results

Connectivity-based parcellation

For each species, a data-driven connectivity-based parcellation was applied to group the vertices in the

IPL into functionally distinct clusters based on anatomical connectivity (**Figure 1**). Because spectral

clustering does not require a specific number of clusters, we iterated the number of subregions from

two to twelve to search for the optimal number of subregions. To accomplish this, we identified the

optimal number of subregions of the IPL by choosing the maximum number of subregions that showed

a coherent topological organization across all species while balancing that by the minimum number of

subregions that could be identified based on their cytoarchitectural definitions in macaques,

110 chimpanzees, and humans (Pandya and Seltzer, 1982; Reyes, 2017). The two- to five-cluster solutions

111 are shown in **Supplementary Figure 1**. The two- to four-cluster solutions showed a consistent rostral-

112 caudal pattern in all three species, but in the five-cluster solution a ventral cluster emerged in

113 chimpanzees and a dorsal cluster emerged in humans. The four-cluster solution revealed a rostral-

114 caudal topological pattern that was consistent with previous parcellations based on cytoarchitecture

115 and anatomical connectivity (Pandya and Seltzer, 1982; Caspers *et al.*, 2006; Mars *et al.*, 2011; Fan *et*

116 *al.*, 2016). Also, the cytoarchitectural definition of macaques revealed four subregions in the IPL

117 (Pandya and Seltzer, 1982), which was fewer than the seven cytoarchitectural subregions of the human

118 IPL (Caspers *et al.*, 2006). Although the four-cluster solution was not the finest, especially in humans,

119 it contained potentially valuable information about the differences between species. Furthermore, the

120 aim of our research was not to find the "best" cluster solution for the IPL but to identify an appropriate

121 parcellation that could shed light on the lateralization of the structure and connectivity of the IPL and

122 its subregions in this particular sample of three primate species. As such, we chose four clusters as the

123 optimal solution for the cross-species comparison.

124 It is widely accepted that the IPL contains two major cytoarchitectural divisions across species, the

125 anterior (PF) and posterior (PG) areas (von Economo and Koskinas, 1925; Von Bonin, 1947; Bailey *et*

126 *al.*, 1950). Our results were consistent with this two-way parcellation and refined it into four

127 subdivisions, specifically, two anterior clusters (the C1 and C2) in the PF and two posterior clusters

128 (the C3 and C4) in the PG. In macaques and chimpanzees, the IPL was previously parcellated into four

129 distinct areas based on histology (Pandya and Seltzer, 1982; Reyes, 2017) in keeping with our four-

130 cluster solution. In humans, the IPL was cytoarchitecturally parcellated into seven distinct areas.

131 Although we proposed a four-cluster solution that has fewer areas than the cytoarchitectural map, it is

also consistent with it (Caspers *et al.*, 2006). Specifically, the rostral anterior cluster (C1) is similar to

the PFt and part of the PFop area defined using cytoarchitecture by Caspers et al. (2006), the caudal

anterior cluster (C2) corresponds to the PF and PFm areas, the rostral posterior cluster (C3) is similar

to the PGa area, and the caudal posterior cluster (C4) is similar to the PGp area. Our results did not

include the PFcm area because it is located deep in the parietal operculum. Given the limited

descriptions of subdivisions and connectivity of the IPL in chimpanzees, our parcellation of the IPL

can depict the subregions and connectivity of the IPL in chimpanzees from an evolutionary perspective.

To assess which hemisphere was dominant with respect to a given function of the human IPL

subregions, we decoded the functions of the human IPL subregions from the Neurosynth database

(Yarkoni *et al.*, 2011) and calculated differences in the correlation values between the left and right

corresponding subregions (**Supplementary Figure 2**). The term *tool* showed a much higher

correlation with the left C1 than with the right C1, suggesting that the left C1 is more involved in tool

use. Terms such as *tool* and *semantics* showed relatively high correlations with the left C2, whereas

terms such as *nogo* and *inhibition* showed relatively high correlations with the right C2, suggesting

that the left C2 is more involved in tool use and language whereas the right C2 is more involved in

executive function. Terms such as *retrieval*, *episodic*, *recollection*, *memories*, and *coherent* showed

relatively high correlations with the left C3, whereas terms such as *nogo*, *inhibition*, and *beliefs* were

correlated with the right C3, suggesting that the left C3 is more involved in memory and language

whereas the right C3 is more involved in executive and social cognitive functions. Terms such as

episodic and *coherent* showed relatively high correlations with the left C4, whereas terms such as

spatial, *attention*, *mentalizing*, and *relevance* showed relatively high correlations with the right C4,

suggesting that the left C4 could be more involved in memory and language whereas the right C4 could

be more involved in spatial attention and social functions.

Figure 1. Framework of the connectivity-based brain parcellation for macaques, chimpanzees, and

humans. (**A**) Defining the seed masks of the inferior parietal lobule (IPL) in surface space according

to the gyri and sulci. (**B**) Connectivity-based parcellation using anatomical connectivity. Probabilistic

tractography was applied by sampling 5000 streamlines at each vertex within the seed mask. Whole-

brain connectivity profiles were used to generate a connectivity matrix with each row representing the

connectivity profile of each seed vertex. Next, a correlation matrix was calculated as a measure of

similarity between the seed vertices. Then, a group similarity matrix was calculated by averaging the

correlation matrix across subjects and spectral clustering was applied to it. (**C**) Parcellation results of

the IPL across species. The entire framework was applied independently for each hemisphere and each

species.

Allometric scaling and structural asymmetry of IPL subregions

When examining the relationship of the volume of each of the IPL subregions scaled against the total

grey matter volume, the scaling of all the IPL subregions showed positive allometry (all slopes > 1)

(**Figure 2A**). A statistical analysis revealed no significant differences between the slopes of each pair

of the bilateral IPL subregions. The asymmetry indices (AIs) for the IPL subregions were calculated

and are shown in **Figure 2B**. The macaques showed no significant asymmetry after Bonferroni

correction for any of the subregions. The chimpanzees and humans both displayed leftward asymmetry

in the rostral IPL (the C1 and C2, all $p < .001$) and rightward asymmetry in the caudal IPL (the C3 and

C4, all $p < .001$).

Figure 2. Structural allometric scaling and asymmetries of the inferior parietal lobule (IPL) subregions

across species. (**A**) Volumes of the IPL subregions plotted against total cortical gray matter volume

(GMV). Solid lines represent the best fit using mean macaque, chimpanzee, and human data points;

dotted lines represent 95% confidence intervals. (**B**) Volumetric asymmetries of the IPL subregions.

Negative asymmetry index indicates leftward asymmetry and positive index indicates rightward

asymmetry. * denotes significance at the Bonferroni corrected level of $p < .05$. The error bars indicate

the standard error of the mean.

Connectional asymmetries of IPL subregions

To investigate the connectional asymmetries of the IPL subregions, we first calculated the connectivity

profiles of the left and right subregions in macaques, chimpanzees, and humans using probabilistic

tracking (**Supplementary Figure 3**). Visualization of the connectivity patterns of the IPL did not show

significant interhemispheric asymmetry in macaque monkeys or chimpanzees but did in humans,

especially in connections with the inferior frontal gyrus (IFG) and lateral temporal cortex. A vertex-

wise analysis was then performed to examine the connectional asymmetry of each subregion for each

species by calculating the AIs between its connectivity profiles for the two hemispheres (**Figure 3**).

Additionally, ROI- and tract-wise analyses were used to examine the asymmetry of the cortical regions

and subcortical white matter pathways connected to the subregions, respectively (**Figure 4;**

connectivity values shown in **Supplementary Figure 4, 5**). No significant asymmetries were found in

macaques in any of the statistical analyses after correction for multiple comparisons.

In chimpanzees, the C1 showed significant leftward asymmetry mainly in connections with the anterior

middle frontal gyrus (MFG), anterior IFG, planum temporale, and insula. The C2 showed significant

leftward asymmetric connections with the insula and rightward asymmetric connections with the

superior parietal lobule (SPL) and superior longitudinal fasciculus 2 (SLF2). The C3 showed

significant leftward asymmetric connections with the anterior superior temporal gyrus (STG), anterior

superior temporal sulcus (aSTS), and occipitotemporal area and rightward asymmetric connections

with the SPL and posterior cingulate gyrus (PCC). The C4 showed significant leftward asymmetric

connections with the anterior STG (aSTG) and rightward asymmetry with the SPL and PCC.

In humans, the C1 showed significant leftward asymmetric connections with the ventral premotor and

motor cortices and insula, which was consistent with regional leftward asymmetric connections with

the precentral gyrus (PreG) and insula. The C1 also showed significant leftward asymmetric

connections with the posterior MFG, aSTG, and posterior middle temporal gyrus (MTG) and rightward

asymmetric connections with the orbital part of the IFG, posterior STS, and dorsal precuneus. The C2

showed significant leftward asymmetric connections with the posterior MFG, ventral premotor and

motor cortices, SPL, anterior temporal lobe, and posterior MTG, which was consistent with regional

leftward asymmetric connections with the IFG, PreG, postcentral gyrus (PostG), SPL, and STG and

was supported by leftward asymmetric subcortical connections with the SLF2, SLF3, and arcuate

fasciculus (AF). The C2 also showed rightward asymmetric connections with the orbital part of the

IFG and posterior cingulate sulcus. The C3 showed significant leftward asymmetry mainly in the

connections with the anterior IFG, SPL, and almost all the lateral temporal cortex, which was

consistent with regional leftward asymmetric connections with the MTG and inferior temporal gyrus

(MTG/ITG). The C3 also showed rightward asymmetric connections with the IFG, which was

supported by leftward asymmetric subcortical connections with the SLF3. The C4 showed significant

leftward asymmetry mainly in the connections with the IFG and anterior and posterior temporal cortex.

The C4 also showed significant regional leftward asymmetric connections with the PreG, PostG, and

SPL in the ROI-wise analysis.

Figure 3. Connectional asymmetries of the IPL subdivisions in the vertex-wise analyses across species.

Effect size (Cohen's *d*) related to asymmetric connections of IPL subdivisions displayed on the left

hemisphere of a species-specific standard brain (leftward asymmetry: yellow, rightward asymmetry:

blue) for each species for areas showing significance at the *p* < .05 level corrected for multiple

comparisons using false discovery rate correction. PreG, precentral gyrus; SPL, superior parietal lobule;

aSTG, anterior superior temporal gyrus; aSTS, anterior superior temporal sulci; PT, planum temporale;

VPMC, ventral premotor cortex; pMTG, posterior middle temporal gyrus; IFG, inferior frontal gyrus;

Ins, insula.

Figure 4. (**A**) Connectional asymmetries of IPL subdivisions in the region of interest (ROI)-wise

analyses across species. Connectional asymmetry was calculated for the connections between each IPL

subregion and eleven ROIs. (**B**) Connectional asymmetries of the IPL subdivisions in the tract-wise

analysis across species. Connectional asymmetry was calculated for the connections between each IPL

subregion and the seven tracts. For all plots, the four quadrants of each circle correspond to the four

IPL subregions. The outermost circles represent ROIs or tracts. The three inner circles from inside to

outside represent macaques, chimpanzees, and humans, respectively. For all plots, only the

connectivity showing a significance at a Bonferroni corrected level of $p < .05$ are displayed. SFG,

superior frontal gyrus; IFG, inferior frontal gyrus; CGa, anterior cingulate gyrus; Orb, orbitofrontal

cortex; PreG, precentral gyrus; PostG, postcentral gyrus; SPL, superior parietal lobule; STG, superior

temporal gyrus; MTG/ITG, middle temporal gyrus and inferior temporal gyrus; Ins, insula; SLF1,

SLF2, SLF3, the three branches of the superior longitudinal fasciculus; AF, arcuate fasciculus; MdLF,

middle longitudinal fasciculus; ILF, inferior longitudinal fasciculus; IFOF, inferior fronto-occipital

fasciculus.

Discussion

In the present study, we investigated asymmetries of the IPL in the structure and connectivity of rhesus

macaques, chimpanzees, and humans. In the structural analysis, the IPL and its subregions exhibited a

similar pattern of positive allometric scaling between hemispheres. In addition, the chimpanzees and

humans shared similar asymmetric patterns in the IPL subregions, i.e., left asymmetry in the anterior

part and right asymmetry in the posterior part, whereas macaques did not display asymmetry. In the

connectivity analysis, the chimpanzees showed some connectional asymmetric regions including the

SPL, insula, planum temporale, aSTG, and aSTS. The humans showed widespread connectional

asymmetric regions including the primary motor and premotor cortices, SPL, insula, and the entire

lateral temporal lobe. These regions are associated with language, tool use, and handedness, suggesting

a potential relationship between the connectional asymmetry and the functional hemispheric

specialization of the human brain.

Positive allometric scaling and structural asymmetry of IPL subregions

Brain allometry describes the quantitative scaling relationship between changes in the size of one

structure relative another structure, often the whole brain or cerebral cortex (Mars *et al.*, 2017; Smaers

et al., 2017). Previous allometric studies suggested that the association cortex (prefrontal, temporal,

and parietal regions) scales with positive allometry (i.e., increases in size disproportionally, or more

rapidly) across primates (Passingham and Smaers, 2014; Mars *et al.*, 2017). Utilizing parcellation-

based delineations, a recent study provided evidence that human brains have a greater proportion of

prefrontal cortex gray matter volume than other primates (Donahue *et al.*, 2018) and other studies

demonstrate that human prefrontal expansion is greater than would be expected from allometric scaling

in nonhuman primates (Passingham and Smaers, 2014; Smaers *et al.*, 2017), although some conflicting

analyses remain (Gabi *et al.*, 2016). In the present study, we used macro-anatomical boundaries to

identify the boundaries of the IPL and a connectivity-based parcellation approach to subdivide the IPL,

which helped to reveal its internal organization. We found that the bilateral IPL subregions exhibited

consistent, positive allometric scaling, which suggests that allometric scaling of the internal

organization of the IPL was similar and was also consistent between homotopic regions during the

evolution of the IPL in anthropoid primates. With only three species in the sample, our dataset does

not allow us to use phylogenetic comparative statistical methods or determine whether human IPL

subregions fall significantly above allometric expectations from nonhuman primates; future research

that incorporates a broad phylogenetic sample of diverse primate brains would be necessary.

We found that chimpanzees and humans showed a similar dichotomous asymmetric pattern in their

IPL subregions, i.e., leftward asymmetry in the anterior portion (the C1 and C2) and rightward

asymmetry in the posterior portion (the C3 and C4), but macaques did not show any asymmetry. The

result in humans is consistent with a recent study using data from a large consortium showing leftward

asymmetry in the supramarginal gyrus and rightward asymmetry in the angular gyrus in terms of

surface area (Kong *et al.*, 2018). The divergent volumetric asymmetries suggest functional

heterogeneities of the IPL and emphasize the importance of analyzing subregions within the IPL. The

shared asymmetric pattern also suggests that divergences in the internal organization of the IPL

evolved prior to the common ancestor of chimpanzees and humans and after the common ancestor of

the three species.

Connectional asymmetries underlying human language and complex tool use

Recent neuroimaging studies have highlighted specific brain regions and pathways that may be

necessary for tool use (Lewis, 2006; Stout and Chaminade, 2012). We found that humans showed

leftward asymmetric connectivity between the IPL (the C2) and the primary motor cortex, ventral

premotor cortex, SPL, and posterior MTG, all of which were activated in tasks related to tool use and

might constitute a cortical network underlying complex tool use (Lewis, 2006). In addition, portions

of this network appeared to represent part of a system that is tightly linked with language systems. The

interaction between the tool use system and the language system, though with a clear left hemisphere

bias, is responsible for representing semantic knowledge about familiar tools and their uses and for

acquiring the skills necessary to perform these actions (Johnson-Frey, 2004; Lewis, 2006; Stout and

Chaminade, 2012; Mars *et al.*, 2017). Several theories suggest that the evolutionary path leading to

language and tool use in humans may be built upon modifications of circuits that subserve gestures

and imitation (Lewis, 2006). Macaques are thought to emulate the goals and intentions of others,

whereas chimpanzees can also imitate certain specific actions, but humans have an even stronger bias

for high-fidelity copying of precise sequences of actions, which has been called "overimitation" (Hecht

et al., 2013). Our findings provide a potential explanation for these phenomena in that the macaques

showed no asymmetric network connections, the chimpanzees showed a few asymmetric connections,

but the humans showed a large number of asymmetric connections. These species differences in

leftward asymmetric connections involving language and tool use may reflect human specializations

for language and complex tool use.

Unlike the humans, who showed considerable leftward asymmetry connectivity between the IPL and

the lateral temporal cortex, the chimpanzees showed few leftward asymmetric connections between

the IPL and the temporal cortex, including the planum temporale, aSTG, and aSTS, while macaques

showed symmetric connections between the IPL and temporal cortex. The planum temporale is

considered to include part of Wernicke's area homolog (Spocter *et al.*, 2010), and displays leftward

anatomical asymmetry in humans and great apes (Gannon *et al.*, 1998; Hopkins *et al.*, 1998). Recent

work suggested a left-hemispheric size predominance of the planum temporale in olive baboons, a

nonhominid primate species (Marie *et al.*, 2018). We speculate that this planum temporale asymmetry

may not be the only prominent characteristic related to language lateralization. The patterns from

symmetry in macaques to asymmetry in humans and chimpanzees in the present study provide a

possible new evidence that neural connectivity asymmetry may underlie the roots of language

specialization, with the initial emergence of hemispheric specializations in apes which are elaborated

even further in human brain evolution. In addition, identifying increased asymmetric connections

between the IPL and planum temporale in human brains compared to chimpanzees and macaques

reinforces the evidence that the evolutionary origin of human language capacities is related to further

left hemispheric specialization of neural substrates for auditory processing that are shared with other

primates (Balezeau *et al.*, 2020). Since the aSTG and aSTS have been implicated in semantic and

phonologic processing in humans (Vigneau *et al.*, 2006), the leftward asymmetric connections of the

IPL with the aSTG and aSTS may be relevant to the evolution of human language processing. Our

results suggest an evolutionary trajectory for the connectivity of the IPL with the temporal cortex; that

is, the connectivity started as symmetric in macaque monkeys, began to develop asymmetrically in

chimpanzees, and finally achieved the greatest degree of asymmetry and is refinement in humans. This

sequence may support the emergence of language and language-related functions.

Species-specific differences in asymmetric connectivity in chimpanzees and humans

Species-specific differences in asymmetric connectivity between the IPL and SPL were found in

chimpanzees and humans, with leftward asymmetry in the former and rightward asymmetry in the

latter, whereas no asymmetry of this connectivity was found in macaques. These species differences

in hemispheric asymmetry may reflect evolutionary changes responsible for adaptations or the

production of new abilities in the human brain. Structurally, in chimpanzees, right anatomical

asymmetry in the white matter below the SPL (Hopkins *et al.*, 2010) may increase the right

connectivity between the IPL and SPL compared with the left side. In humans, the leftward volumetric

asymmetry in the SPL (Goldberg *et al.*, 2013), together with leftward volumetric asymmetry in the IPL

(the C2), may support the leftward asymmetric connectivity. Functionally, interaction between the IPL

and SPL is crucial for tool use, which is dominant in the left hemisphere, and visuospatial function,

which is dominant in in the right (Lewis, 2006; De Schotten *et al.*, 2011b; Catani *et al.*, 2017). As for

tool use, in contrast to the relatively simple tools used by chimpanzees and other species, humans can

create complex artifacts through a sequence of actions that may incorporate multiple parts, reflecting

a deep understanding of the kinematics of our bodies, the mechanical properties of surrounding objects,

and the unique demands of the external environments in which we live (Povinelli *et al.*, 2000; Johnson-

Frey, 2004). In addition, complex tool use requires the SPL to code the location of the limbs relative

to other body parts during planning and executing tool-use movements or hand gestures (Wolpert *et*

al., 1998; Johnson-Frey *et al.*, 2005; Lewis, 2006). Leftward asymmetric connectivity between the IPL

and SPL may have provided a connectional substrate for complex tool use during human evolution.

As for visuospatial functions, the rightward asymmetric connectivity between the IPL and SPL in

chimpanzees may indicate that visuospatial functions are dominant in the right hemisphere and had

already been lateralized to the right hemisphere from the common ancestor with macaques. During

evolution, these lateralized functions may be retained in the human brain. Meanwhile, the lateralized

directional reversal of this connectivity from the right to the left hemisphere may reflect evolutionary

adaptations for the emergence of new abilities, such as sophisticated and complex tool making and use.

Human unique asymmetric connectivity of IPL subregions

Unlike the chimpanzees and macaques, humans showed leftward asymmetry in the connection

between the rostral IPL (the C1 and C2) and the primary motor cortex, which is consistent with a larger

neuropil volume in the left primary motor cortex than in the right side (Amunts *et al.*, 1996).

Meanwhile, the leftward asymmetric volume of the anterior IPL and the primary motor cortex may

also increase the neural connectivity between these two regions in the left hemisphere compared with

the right side. Such a leftward connection is thought to be related to handedness and hand manual skills

(Amunts *et al.*, 1996; Amunts *et al.*, 1997). In contrast to humans, chimpanzees and macaques did not

show any asymmetric connectivity between the IPL and the primary motor cortex. A more recent study

reported that, in olive baboons, contralateral hemispheric sulcus depth asymmetry of the central sulcus

related to the motor hand area is correlated with the direction and degree of hand preference, as

measured by a bimanual coordinated tube task, but only about 41% of them were classified as right-

handed and 33% were classified as left-handed (Margiotoudi *et al.*, 2019). Although previous studies

have shown that chimpanzees exhibit population-level handedness in the use of tools and a

corresponding asymmetry in the primary motor cortex, inferior frontal cortex, and parietal operculum

(Gilissen and Hopkins, 2013; Hopkins *et al.*, 2017), they do not show handedness as a more universal

trait or exhibit manual dexterity to the same extent as humans. One possible explanation is that humans

developed the asymmetric connectivity that became the structural basis for specific behaviors of

handedness and hand skills during evolution.

An unexpected finding was that in humans the IPL, particularly the C3, showed rightward asymmetric

connectivity with the IFG. Since the IPL and the IFG are interconnected through the SLF3, which is

strongly rightward asymmetric (De Schotten *et al.*, 2011b), it may also increase the connection between

the IPL and IFG in the right hemisphere. Functionally, the left IFG is involved in various aspects of

language functions, including speech production and semantic, syntactic, and phonological processing

(Wang *et al.*, 2020), whereas the right IFG is associated with various cognitive functions, including

attention, motor inhibition, and social cognitive processes (Hartwigsen *et al.*, 2019). Our result of

rightward asymmetry in this connectivity seems to be associated with attention and social function,

but not language, although language dominance in the left hemisphere is considered to be a common

characteristic in humans.

The widespread asymmetric connections of the IPL in humans compared with the other two primates

is in keeping with the inter-hemispheric independence hypothesis, in which, during evolution, brain

size expansion led to hemispheric specialization due to time delays in neuron signaling over increasing

distances, resulting in decreased inter-hemispheric connectivity and increased intra-hemispheric

connectivity (Ringo *et al.*, 1994; Phillips *et al.*, 2015). While having more cortical neurons (local

characteristics) in one hemisphere than the other seems to be a necessary condition for asymmetries of

complex and flexible behaviors, it is not a full condition for such behaviors. Given that a function or

behavior in an area is determined by its connectivity or networks in which it is involved (Passingham

et al., 2002), the widespread lateralized connections may provide the human brain with the increased

computational capacity necessary for processing language and complex tool use and may play a

facilitating role in human cognitive and behavioral specialization.

Methodological considerations

The three levels of analyses, i.e., the vertex-wise, ROI-wise, and tract-wise analyses, were performed

to provide a full description of the connectivity asymmetry. However, it should be noted that some

analyses produced results that were not completely consistent with each other. In humans, the

connectivity of the IPL with SLF3 and AF was left-lateralized in the C2 while right-lateralized in the

C3. The previous studies assessed the asymmetry of the SLF3 and AF with local characteristics such

as cortical volume, voxel count, and FA and their average across all the voxels in the tracts (de Schotten

et al., 2011a; De Schotten *et al.*, 2011b; Kamali *et al.*, 2014), the SLF3 and AF were usually found to

have a single pattern, e.g., leftward asymmetry, rightward asymmetry, or symmetry. However, our

results seem to indicate two different asymmetric patterns for the SLF3 and AF, both of which connect

the IPL subregion C2 and C3 to the IFG (De Schotten *et al.*, 2011b; Hecht *et al.*, 2015; Barbeau *et al.*,

2020). Furthermore, these connectivity asymmetries matched well with the ROI-wise and tract-wise

analyses. The leftward connectivity asymmetry of the human C2 with the SLF3 and AF using the tract-

wise approach corresponds to that of the human C2 with the IFG and precentral gyrus using the ROI-

wise approach. The rightward connectivity asymmetry of the human C3 with the SLF3 and AF using

the tract-wise approach corresponds to that of the human C3 with the IFG using the ROI-wise approach.

The SLF3, located at the ventrolateral SLF, connects to the IPL, especially the anterior part, and from

there predominantly to the ventral premotor and prefrontal areas (De Schotten *et al.*, 2011b; Kamali *et*

al., 2014; Barbeau *et al.*, 2020). The C2 and C3 appear to separate the SLF3 into two finer components,

one connecting the posterior IFG and anterior IPL with leftward asymmetry, and one connecting the

anterior IFG to the posterior IPL with rightward asymmetry. The two types of connectivity patterns are

consistent with previous studies using invasive tract-tracing findings in macaque monkeys and resting-

state functional connectivity results in humans to study frontal and parietal connectivity (Petrides and

Pandya, 2009; Margulies and Petrides, 2013). Our results indicated that both cortical areas, such as the

IFG, and subcortical tracts, such as the SLF3 and AF, have at least two distinct subcomponents.

The inconsistency was observed when significant ROI-wise connectivity asymmetry was found but

few or no significant tract-wise connectivity asymmetries were found. For example, the human C1

showed ROI-wise connectivity asymmetry with the precentral gyrus and insula but no significant tract-

wise connectivity asymmetry. The IPL is connected to the precentral gyrus mainly through the SLF3,

which in turn is connected to not only the precentral gyrus but also the IFG and MFG in the prefrontal

cortex (De Schotten *et al.*, 2011b; Kamali *et al.*, 2014; Hecht *et al.*, 2015; Barbeau *et al.*, 2020). The

connectivity between the C1 and the precentral gyrus may include only a portion of the SLF3; this

may have diluted the laterality effect from the SLF3 because it may include other pathways that were

not in our selected tracts and could, thus, have affected the observed lateralization. In other words, the

traditionally defined major fiber tracts are not a single bundle but, instead, contain many

subcomponents. Therefore, the patterns of lateralization might not yet have been fully explored in our

study. A recent work also suggested that the SLF2, SLF3, and AF could be separated into several

branches based on their projections into the prefrontal and/or temporal areas (Barbeau *et al.*, 2020).

This may be true for the other major fiber tracts, such as the ILF (Latini *et al.*, 2017), uncinate

fasciculus (Hau *et al.*, 2017), and cingulum bundle (Jones *et al.*, 2013). On the other hand, brain regions,

such as the IFG were connected to many fiber tracts, including the SLFII, SLFIII, and AF. Hence, we

did not find tract-wise connectivity asymmetry that corresponded to the ROI-wise connectivity

asymmetry in the human C1. This was also the case for the chimpanzee and human C4. The creation

of a finer tract atlas should be a priority for future work because this would help to map the tract-wise

connectivity asymmetry at a higher resolution.

In conclusion, we identified similar topographical maps of the IPL to study structural and connectional

asymmetry in macaques, chimpanzees, and humans. We found that the structural asymmetry of the

IPL was independent of the allometric scaling of this region. The connectional analysis revealed that

humans had the largest connectional asymmetries of IPL subregions compared to macaques and

chimpanzees. The regions showing larger asymmetric connections with the human IPL were associated

with language, complex tool use, and handedness, which provided potential anatomical substrates for

functional and behavioral lateralization in humans. The opposite asymmetric connection between the

IPL and SPL in chimpanzees and humans may reflect distinct species-specific modifications to cortical

circuits during the course of ape and human evolution.

Materials and methods

Human data

Data from 40 right-handed healthy adults (age: 22–35, 18 males) were randomly selected from the

S500 subjects release of the Human Connectome Project (HCP) database (Van Essen *et al.*, 2013)

(http://www.humanconnectome.org/study/hcp-young-adult/). T1-weighted (T1w) MPRAGE images

(resolution: 0.7mm isotropic, slices: 256; field of view: 224 × 320; flip angle: 8°), and diffusion-

weighted images (DWI) (resolution: 1.25mm isotropic; slices: 111; field of view: 210 × 180; flip angle:

78°; b-values: 1000, 2000, and 3000 s/mm^2) were collected on a 3 T Skyra scanner (Siemens, Erlangen,

Germany) using a 32-channel head coil.

Chimpanzee data

Data from 27 adult chimpanzees (*Pan troglodytes*, 14 males) were made available by the National

Chimpanzee Brain Resource (http://www.chimpanzeebrain.org, supported by the NIH National

Institute of Neurological Disorders and Stroke). Data, including T1w and DWI, were acquired at the

Yerkes National Primate Research Center (YNPC) on a 3T MRI scanner under propofol anesthesia (10

mg/kg/h) using previously described procedures (Chen *et al.*, 2013). All procedures were carried out

in accordance with protocols approved by YNPRC and the Emory University Institutional Animal Care

and Use Committee (Approval no. YER-2001206).

DWI were acquired using a single-shot spin-echo echo-planar sequence for each of 60 diffusion

directions ($b = 1000$ s/mm^2, repetition time 5900 ms; echo time 86 ms; 41 slices; 1.8 mm isotropic

resolution). DWI with phase-encoding directions (left–right) of opposite polarity were acquired to

correct for susceptibility distortion. For each repeat of a set of DWI, five $b = 0$ s/mm^2 images were also

acquired with matching imaging parameters. T1w images were also acquired for each subject (218

slices, resolution: 0.7x0.7x1mm).

Macaque data

Data from 8 male adult rhesus macaque monkeys (*Macaca mulatta*) were obtained from

TheVirtualBrain (Shen *et al.*, 2019). All surgical and experimental procedures were approved by the

Animal Use Subcommittee of the University of Western Ontario Council on Animal Care (AUP no.

2008-125) and followed the Canadian Council of Animal Care guidelines. Surgical preparation and

anesthesia as well as imaging acquisition protocols have been previously described (Shen *et al.*, 2019).

Images were acquired using a 7-T Siemens MAGNETOM head scanner. Two diffusion-weighted scans

were acquired for each animal, with each scan having opposite phase encoding in the superior-inferior

direction at 1 mm isotropic resolution, allowing for correction of susceptibility-related distortion. For

five animals, the data were acquired with 2D EPI diffusion, while for the remaining three animals, a

multiband EPI diffusion sequence was used. In all cases, data were acquired with $b = 1000$ s/mm^2, 64

directions, 24 slices. Finally, a 3D T1w image was also collected for each animal (128 slices, resolution:

0.5 mm isotropic).

Image preprocessing

The human T1w structural data had been preprocessed following the HCP's minimal preprocessing pipeline (Glasser *et al.*, 2013), while the chimpanzee and monkey T1w structural data had been preprocessed following the HCP's nonhuman preprocessing pipelines described in previous studies (Glasser *et al.*, 2013; Donahue *et al.*, 2018). Briefly, the processing pipeline included imaging alignment to standard volume space using FSL, automatic anatomical surface reconstruction using FreeSurfer, and registration to a group average surface template space using the Multimodal Surface Matching (MSM) algorithm (Robinson *et al.*, 2014). Human volume data were registered to Montreal Neurological Institute (MNI) standard space and surface data were transformed into surface template space (fs_LR). Chimpanzee volume and surface data were registered to the Yerkes29 chimpanzee template (Donahue *et al.*, 2018). Macaque volume and surface data were registered to the Yerkes19 macaque template (Donahue *et al.*, 2018).

Preprocessing of the diffusion-weighted images was performed in a similar way in the human, chimpanzee, and macaque datasets using FSL. FSL's DTIFIT was used to fit a diffusion tensor model for each of the three datasets. Following preprocessing, voxel-wise estimates of the fiber orientation distribution were calculated using Bedpostx, allowing for three fiber orientations for the human dataset and two fiber orientations for the chimpanzee and macaque datasets due to the b-value in the diffusion data.

Definition of the IPL

The IPL, located at the lateral surface of the ventral posterior parietal lobe, is surrounded by several sulci including the Sylvian fissure, superior temporal sulcus (STS), and intraparietal sulcus (IPS) (von

Economo and Koskinas, 1925; Von Bonin, 1947; Bailey *et al.*, 1950; Pandya and Seltzer, 1982). In the

absence of detailed homologous definitions, it is necessary to use cytoarchitectonic delineations and

macroscopic boundaries, such as gyri and sulci, that can be reliably identified in all species as the

boundaries of the IPL. The region of interest (ROI) of the IPL was manually drawn on the standard

surface template using Connectome Workbench (Glasser *et al.*, 2013). In the present study, we

restricted the ROI to the lateral surface of the IPL and excluded the cortex buried in the sulci, especially

the lateral bank of the IPS and the upper bank of the Sylvian fissure. Rostrally, the IPL borders the

vertical line between the Sylvian fissure and the rostral lip of the IPS. Dorsally, the IPL borders the

lateral bank of the IPS. Ventrally, the anterior ventral IPL borders the upper bank of the Sylvian fissure.

The border of the posterior and ventral IPL is formed by the extension of the Sylvian fissure to the top

end of the STS in chimpanzees and macaques but by the extension of the Sylvian fissure to the posterior

end of the IPS in humans.

Connectivity-based parcellation

We used a data-driven connectivity-based parcellation framework modified from Fan *et al* (2016)

(**Figure 1**). All steps in the framework were processed on surface data because the surface-based

method has advantages, such as cortical areal localization (Coalson *et al.*, 2018), over the traditional

approach and because the use of surface meshes is a straightforward way to improve existing

tractography processing pipelines, such as the precise locations of streamline seeding and termination

(St-Onge *et al.*, 2018). The surface ROI was first registered to native surface using MSM (Robinson

et al., 2014). The probabilistic tractography was performed on the native mesh representing the

gray/white matter interface using Probtrackx. The pial surfaces were used as stop masks to prevent

streamlines from crossing sulci. 5000 streamlines were seeded from each of the white matter surface

vertices in the seed region to estimate its whole-brain connectivity profile and were downsampled to

5 mm isotropic voxels to construct the native connectivity M-by-N, a matrix between all the IPL

vertices (M) and the brain voxels (N). Based on the native connectivity matrix, a symmetric cross-

correlation M-by-M matrix was calculated to quantify the similarity between the connectivity profiles

of each IPL vertex. A group cross-correlation matrix was calculated by averaging the cross-correlation

matrix across subjects.

Data-driven spectral clustering was applied to the group cross-correlation matrix to define the

anatomical boundaries of the IPL. Spectral clustering can capture clusters that have complicated shapes,

making them suitable for parcellating the structure of complicated brain regions such as the IPL. In

addition, the spectral clustering algorithm was successfully used to establish the Brainnetome Atlas

(Fan *et al.*, 2016). However, the number of clusters must be defined by the experimenter when using

this method. In the current study, we explored from two to twelve parcellations.

Volumetric analysis of the IPL

The cortical gray matter volumetric measurements were calculated using Freesurfer. Total cortical

volumes were determined by the space between the white and pial surfaces in native space. Each

subregion drawn on standard surface space was registered to native surface space using an existing

mapping between the two meshes. The volume of the IPL and its subregions was determined by

averaging all the vertices for each subject.

Functional decoding of each subregion of the human IPL

Each subregion was first mapped to MNI volume space using a ribbon-constrained method in Connectome Workbench. To decode the functions of each subregion, we used the automated meta-analysis database, Neurosynth (Yarkoni *et al.*, 2011) to identify the terms that were the most associated with each subregion. The top five non-anatomical terms with the highest correlation values were kept for all subregions and redundant terms, such as '*semantic*' and '*semantics*', were only considered once. For simplicity, we only showed the positive correlations found by decoding because negative correlations do not directly inform us about the functions of the subregions. The lateralization for each term was obtained by calculating the difference in the correlation values of the subregions between the left and right hemispheres.

Mapping anatomical connectivity profiles

To map the whole-brain anatomical connectivity pattern for each cluster, we performed probabilistic tractography by drawing 5000 samples from each vertex in each cluster. The resulting tractograms were log-transformed, normalized by the maximum, and then projected onto surface space using the 'surf_proj' command in FSL to obtain tractograms in surface space. The surface tractograms were smoothed using a 4 mm kernel for humans, 3 mm kernel for chimpanzees, and 2 mm kernel for macaques. We subsequently averaged the surface tractograms across subjects for the left and right hemispheres separately to obtain population tractograms, which were thresholded by a value of 0.5 for humans, 0.2 for chimpanzees, and 0.3 for macaques due to data quality. The resultant population tractograms represented approximately twenty percent of the non-zero vertexes in the non-thresholded population tractograms and were used for the vertex-wise and ROI-wise comparisons. The volumetric

tractograms were used for the tract-wise comparison.

Vertex-wise analysis

For each subregion, we restricted the analysis to the group mask defined by the combination of the left

and mirrored right population tractograms described above. We here used the connectivity probabilistic

value to quantify the connectivity between the IPL and each vertex of the rest of the brain. A higher

value in the vertex means a higher likelihood of being connected to the IPL than other vertices.

ROI-wise analysis

Although previous studies have devoted much effort to establishing homologous regions in primates,

these are still limited to a few regions, particularly in chimpanzees. To make comparisons across

species possible, here we used the common principle of macroscopic anatomical boundaries based on

the gyri and sulci to define ROIs in the cerebral cortex. Specifically, the Desikan–Killiany–Tourville

(DKT) atlas was used for humans (Desikan *et al.*, 2006), a modified DKT atlas for the chimpanzees,

and the Neuromaps atlas for the macaques (Rohlfing *et al.*, 2012). Because the Neuromaps atlas is

volumetric, we first mapped it to surface space for the subsequent calculations. A total of eleven

cortical ROIs were chosen for each hemisphere: the superior frontal gyrus (SFG), inferior frontal gyrus

(IFG, a combination of the pars triangularis and pars opercularis in humans and chimpanzees), anterior

cingulate gyrus (CGa, a combination of the rostral and caudal anterior-cingulate in humans and

chimpanzees), orbitofrontal cortex (Orb), precentral gyrus (PreG), postcentral gyrus (PostG), superior

parietal lobule (SPL), precuneus, superior temporal gyrus (STG), middle temporal gyrus and inferior

temporal gyrus (MTG/ITG), and insula. The MTG/ITG was a combination of the MTG and ITG in

598 humans and chimpanzees due to the absence of the MTG in macaques. The connectional value for

599 each ROI was calculated by averaging all vertices in the ROI on the individual surface tractogram for

600 each subregion.

601 **Tract-wise analysis**

602 To investigate which subcortical fiber tracts are associated with lateralization of cortical areas

603 connected to the IPL, we analyzed the lateralization of the subcortical white matter tracts connected to

604 the IPL across species. A total of seven tracts were chosen: the three branches of the superior

605 longitudinal fasciculus, arcuate fasciculus, middle longitudinal fasciculus, inferior longitudinal

606 fasciculus, inferior fronto-occipital fasciculus. The automated tractographic protocols for tracts for

607 each species were from previous studies (Bryant *et al.*, 2020) and these tracts were reconstructed using

608 the Xtract tool (Warrington *et al.*, 2020). The mean value for each tract was calculated by averaging

609 all voxels in the tract in the individual volumetric tractogram for each subregion.

610 **Statistical analysis**

611 To investigate the allometric relationship between the volume of each of the IPL subregions and the

612 total gray matter volume using log-transformed data (Donahue *et al.*, 2018), linear regression was

613 performed by pooling the human, chimpanzee, and macaque data for each of the IPL subregions,

614 separately. To test whether the scaling regression slopes differed significantly between the two

615 hemispheres, we performed an ANCOVA for comparisons across the two regression slopes for each

616 plot.

617 In all the analyses of the structural and connectional asymmetries (i.e., volumetric, vertex-wise, ROI-

wise, and tract-wise), the asymmetry index (AI) was defined as the difference between values for the

left and right hemispheres according to the formula AI = 2 x (R − L) / (R + L). For the vertex-wise

analysis, a one-sample t test was performed at each vertex on the group mask for each species using

PALM, with 5000 permutations with a sign-flip strategy (Winkler *et al.*, 2014). The statistically

significant level was set at false discovery rate corrected $p < .05$. The effect sizes (Cohen's d) were

displayed on the average surface. For the volumetric, ROI-wise, and tract-wise analysis, a two-sided

Wilcoxon signed-rank test was performed for each subregion. Bonferroni correction was then used for

multiple comparisons for seeds, ROIs or tracts, and species, with statistical significance set at $p < .05$.

Acknowledgments

This work was partially supported by the National Natural Science Foundation of China (Grant Nos.

91432302, 82072099, and 31620103905), the Science Frontier Program of the Chinese Academy of

Sciences (Grant No. QYZDJ-SSW-SMC019), Beijing Municipal Science & Technology Commission

(Grant Nos. Z161100000216139 and Z171100000117002), the Guangdong Pearl River Talents Plan

(2016ZT06S220), Key-Area Research and Development Program of Guangdong Province

(2018B030333001), the Youth Innovation Promotion Association, the Beijing Advanced Discipline

Fund, and the National Science Foundation (SMA-1542848). The National Chimpanzee Brain

Resource was supported by NIH - National Institute of Neurological Disorders and Stroke (NIH Grant

No. NS092988). We thank Rhoda E. and Edmund F. Perozzi, PhDs, for English language and editing

assistance.

Author contributions

T.J. and L.F. designed the study. L.C. and G.L. analyzed the data. C.C.S collected the chimpanzee data.

L.C. wrote the first draft of the manuscript. T.J. supervised the study. All authors revised and approved

the manuscript.

Data availability

The datasets analyzed during the current study are available at https://www.humanconnectome.org,

http://www.chimpanzeebrain.org, http://openneuro.org/datasets/ds001875/versions/1.0.3, and

http://www.neurosynth.org.

Competing interests

The authors declare no competing interests.

References

Amunts K, Schlaug G, Jäncke L, Steinmetz H, Schleicher A, Dabringhaus A, Zilles K. Motor cortex

and hand motor skills: structural compliance in the human brain. Human Brain Mapping 1997; 5(3):

206-15.

Amunts K, Schlaug G, Schleicher A, Steinmetz H, Dabringhaus A, Roland PE, Zilles K. Asymmetry

in the human motor cortex and handedness. NeuroImage 1996; 4(3): 216-22.

Ardesch DJ, Scholtens LH, Li L, Preuss TM, Rilling JK, van den Heuvel MP. Evolutionary expansion

of connectivity between multimodal association areas in the human brain compared with chimpanzees.

Proceedings of the National Academy of Sciences 2019; 116(14): 7101-6.

Bailey P, Bonin GV, McCulloch WS. The isocortex of the chimpanzee. Urbana: University of Illinois

Press; 1950.

Balezeau F, Wilson B, Gallardo G, Dick F, Hopkins W, Anwander A, Friederici AD, Griffiths TD,

Petkov CI. Primate auditory prototype in the evolution of the arcuate fasciculus. Nature Neuroscience

2020; 23(5): 611-4.

Barbeau EB, Descoteaux M, Petrides M. Dissociating the white matter tracts connecting the temporo-

parietal cortical region with frontal cortex using diffusion tractography. Scientific reports 2020; 10(1):

1-13.

Binder JR, Desai RH, Graves WW, Conant LL. Where is the semantic system? A critical review and

meta-analysis of 120 functional neuroimaging studies. Cerebral Cortex 2009; 19(12): 2767-96.

Bryant KL, Li L, Eichert N, Mars RB. A comprehensive atlas of white matter tracts in the chimpanzee.

PLoS Biology 2020; 18(12): e3000971.

Caspers S, Geyer S, Schleicher A, Mohlberg H, Amunts K, Zilles K. The human inferior parietal cortex:

673 cytoarchitectonic parcellation and interindividual variability. NeuroImage 2006; 33(2): 430-48.

674 Catani M, Robertsson N, Beyh A, Huynh V, de Santiago Requejo F, Howells H, Barrett RLC, Aiello

675 M, Cavaliere C, Dyrby TB, Krug K, Ptito M, D'Arceuil H, Forkel SJ, Dell'Acqua F. Short parietal lobe

676 connections of the human and monkey brain. Cortex 2017; 97: 339-57.

677 Chance SA, Crow TJ. Distinctively human: cerebral lateralisation and language in Homo sapiens. The

678 Journal of Anthropological Sciences 2007; 85: 83-100.

679 Changeux J-P, Goulas A, Hilgetag CC. A connectomic hypothesis for the hominization of the brain.

680 Cerebral Cortex 2020; 31(5): 24252449.

681 Chaplin TA, Yu H-H, Soares JG, Gattass R, Rosa MG. A conserved pattern of differential expansion

682 of cortical areas in simian primates. The Journal of Neuroscience 2013; 33(38): 15120-5.

683 Chen X, Errangi B, Li L, Glasser MF, Westlye LT, Fjell AM, Walhovd KB, Hu X, Herndon JG, Preuss

684 TM, Rilling JK. Brain aging in humans, chimpanzees (Pan troglodytes), and rhesus macaques (Macaca

685 mulatta): magnetic resonance imaging studies of macro- and microstructural changes. Neurobiology

686 of Aging 2013; 34(10): 2248-60.

687 Coalson TS, Van Essen DC, Glasser MF. The impact of traditional neuroimaging methods on the spatial

688 localization of cortical areas. Proceedings of the National Academy of Sciences 2018; 115(27): E6356-

689 E65.

690 Croxson PL, Forkel SJ, Cerliani L, Thiebaut de Schotten M. Structural variability across the primate

691 brain: a cross-species comparison. Cerebral Cortex 2018; 28(11): 3829-41.

692 de Schotten MT, Bizzi A, Dell'Acqua F, Allin M, Walshe M, Murray R, Williams SC, Murphy DG,

693 Catani M. Atlasing location, asymmetry and inter-subject variability of white matter tracts in the

694 human brain with MR diffusion tractography. NeuroImage 2011a; 54(1): 49-59.

De Schotten MT, Dell'Acqua F, Forkel S, Simmons A, Vergani F, Murphy DG, Catani M. A lateralized

brain network for visuo-spatial attention. Nature Precedings 2011b: 1-.

Desikan RS, Ségonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, Buckner RL, Dale AM,

Maguire RP, Hyman BT, Albert MS, Killiany RJ. An automated labeling system for subdividing the

human cerebral cortex on MRI scans into gyral based regions of interest. NeuroImage 2006; 31(3):

968-80.

Donahue CJ, Glasser MF, Preuss TM, Rilling JK, Van Essen DC. Quantitative assessment of prefrontal

cortex in humans relative to nonhuman primates. Proceedings of the National Academy of Sciences

2018; 115(22): E5183-E92.

Eickhoff SB, Yeo BT, Genon S. Imaging-based parcellations of the human brain. Nature Reviews

Neuroscience 2018; 19(11): 672-86.

Fan L, Li H, Zhuo J, Zhang Y, Wang J, Chen L, Yang Z, Chu C, Xie S, Laird AR, Fox PT, Eickhoff

SB, Yu C, Jiang T. The human Brainnetome Atlas: a new brain atlas based on connectional architecture.

Cerebral Cortex 2016; 26(8): 3508-26.

Gabi M, Neves K, Masseron C, Ribeiro PF, Ventura-Antunes L, Torres L, Mota B, Kaas JH, Herculano-

Houzel S. No relative expansion of the number of prefrontal neurons in primate and human evolution.

Proceedings of the National Academy of Sciences 2016; 113(34): 9617-22.

Gannon PJ, Holloway RL, Broadfield DC, Braun AR. Asymmetry of chimpanzee planum temporale:

humanlike pattern of Wernicke's brain language area homolog. Science 1998; 279(5348): 220-2.

Gilissen EP, Hopkins WD. Asymmetries of the parietal operculum in chimpanzees (Pan troglodytes)

in relation to handedness for tool use. Cerebral Cortex 2013; 23(2): 411-22.

Glasser MF, Sotiropoulos SN, Wilson JA, Coalson TS, Fischl B, Andersson JL, Xu J, Jbabdi S, Webster

M, Polimeni JR, Van Essen DC, Jenkinson M. The minimal preprocessing pipelines for the Human

Connectome Project. NeuroImage 2013; 80: 105-24.

Goldberg E, Roediger D, Kucukboyaci NE, Carlson C, Devinsky O, Kuzniecky R, Halgren E, Thesen

T. Hemispheric asymmetries of cortical volume in the human brain. Cortex 2013; 49(1): 200-10.

Gómez-Robles A, Hopkins WD, Sherwood CC. Increased morphological asymmetry, evolvability and

plasticity in human brain evolution. Proceedings of the Royal Society B: Biological Sciences 2013;

280(1761): 20130575.

Hartwigsen G, Neef NE, Camilleri JA, Margulies DS, Eickhoff SB. Functional segregation of the right

inferior frontal gyrus: evidence from coactivation-based parcellation. Cerebral Cortex 2019; 29(4):

1532-46.

Hau J, Sarubbo S, Houde JC, Corsini F, Girard G, Deledalle C, Crivello F, Zago L, Mellet E, Jobard

G. Revisiting the human uncinate fasciculus, its subcomponents and asymmetries with stem-based

tractography and microdissection validation. Brain Structure and Function 2017; 222(4): 1645-62.

Hecht EE, Gutman DA, Bradley BA, Preuss TM, Stout D. Virtual dissection and comparative

connectivity of the superior longitudinal fasciculus in chimpanzees and humans. Neuroimage 2015;

108: 124-37.

Hecht EE, Gutman DA, Preuss TM, Sanchez MM, Parr LA, Rilling JK. Process versus product in

social learning: comparative diffusion tensor imaging of neural systems for action execution–

observation matching in macaques, chimpanzees, and humans. Cerebral Cortex 2013; 23(5): 1014-24.

Hopkins WD. Neuroanatomical asymmetries and handedness in chimpanzees (Pan troglodytes): a case

for continuity in the evolution of hemispheric specialization. Annals of the New York Academy of

Sciences 2013; 1288(1): 17.

Hopkins WD, Marino L, Rilling JK, MacGregor LA. Planum temporale asymmetries in great apes as

revealed by magnetic resonance imaging (MRI). NeuroReport 1998; 9(12): 2913-8.

Hopkins WD, Meguerditchian A, Coulon O, Misiura M, Pope S, Mareno MC, Schapiro SJ. Motor skill

for tool-use is associated with asymmetries in Broca's area and the motor hand area of the precentral

gyrus in chimpanzees (Pan troglodytes). Behavioural Brain Research 2017; 318: 71-81.

Hopkins WD, Taglialatela JP, Nir T, Schenker NM, Sherwood CC. A voxel-based morphometry

analysis of white matter asymmetries in chimpanzees (Pan troglodytes). Brain, Behavior and Evolution

2010; 76(2): 93-100.

Johnson-Frey SH. The neural bases of complex tool use in humans. Trends in Cognitive Sciences 2004;

8(2): 71-8.

Johnson-Frey SH, Newman-Norlund R, Grafton ST. A distributed left hemisphere network active

during planning of everyday tool use skills. Cerebral Cortex 2005; 15(6): 681-95.

Jones DK, Christiansen KF, Chapman R, Aggleton JP. Distinct subdivisions of the cingulum bundle

revealed by diffusion MRI fibre tracking: implications for neuropsychological investigations.

Neuropsychologia 2013; 51(1): 67-78.

Kaas JH. The evolution of neocortex in primates.   Progress in Brain Research: Elsevier; 2012. p. 91-

102.

Kamali A, Flanders AE, Brody J, Hunter JV, Hasan KM. Tracing superior longitudinal fasciculus

connectivity in the human brain using high resolution diffusion tensor tractography. Brain Structure

and Function 2014; 219(1): 269-81.

Kong X-Z, Mathias SR, Guadalupe T, Glahn DC, Franke B, Crivello F, Tzourio-Mazoyer N, Fisher

SE, Thompson PM, Francks C. Mapping cortical brain asymmetry in 17,141 healthy individuals

worldwide via the ENIGMA Consortium. Proceedings of the National Academy of Sciences 2018; 115(22): E5154-E63.

Latini F, Mårtensson J, Larsson E-M, Fredrikson M, Åhs F, Hjortberg M, Aldskogius H, Ryttlefors M. Segmentation of the inferior longitudinal fasciculus in the human brain: A white matter dissection and diffusion tensor tractography study. Brain Research 2017; 1675: 102-15.

Lewis JW. Cortical networks related to human use of tools. The Neuroscientist 2006; 12(3): 211-31.

Margiotoudi K, Marie D, Claidière N, Coulon O, Roth M, Nazarian B, Lacoste R, Hopkins WD, Molesti S, Fresnais P. Handedness in monkeys reflects hemispheric specialization within the central sulcus. An in vivo MRI study in right-and left-handed olive baboons. Cortex 2019; 118: 203-11.

Margulies DS, Petrides M. Distinct parietal and temporal connectivity profiles of ventrolateral frontal areas involved in language production. Journal of Neuroscience 2013; 33(42): 16846-52.

Marie D, Roth M, Lacoste R, Nazarian B, Bertello A, Anton J-L, Hopkins WD, Margiotoudi K, Love SA, Meguerditchian A. Left brain asymmetry of the planum temporale in a nonhominid primate: Redefining the origin of brain specialization for language. Cerebral Cortex 2018; 28(5): 1808-15.

Mars R, Passingham R, Neubert F, Verhagen L, Sallet J. Evolutionary specializations of human association cortex. Kaas, JH (ed), Evolution of nervous systems (2nd ed, vol 4) 2017: 185-200.

Mars RB, Jbabdi S, Sallet J, O'Reilly JX, Croxson PL, Olivier E, Noonan MP, Bergmann C, Mitchell AS, Baxter MG, Behrens TEJ, Johansen-Berg H, Tomassini V, Miller KL, Rushworth MFS. Diffusion-Weighted Imaging Tractography-Based Parcellation of the Human Parietal Cortex and Comparison with Human and Macaque Resting-State Functional Connectivity. The Journal of Neuroscience 2011; 31(11): 4087-100.

Ocklenburg S, Friedrich P, Güntürkün O, Genç E. Intrahemispheric white matter asymmetries: the

missing link between brain structure and functional lateralization? Reviews in the Neurosciences 2016; 27(5): 465-80.

Orban GA, Claeys K, Nelissen K, Smans R, Sunaert S, Todd JT, Wardak C, Durand J-B, Vanduffel W. Mapping the parietal cortex of human and non-human primates. Neuropsychologia 2006; 44(13): 2647-67.

Pandya DN, Seltzer B. Intrinsic connections and architectonics of posterior parietal cortex in the rhesus monkey. Journal of Comparative Neurology 1982; 204(2): 196-210.

Passingham RE, Smaers JB. Is the prefrontal cortex especially enlarged in the human brain? Allometric relations and remapping factors. Brain, Behavior and Evolution 2014; 84(2): 156-66.

Passingham RE, Stephan KE, Kötter R. The anatomical basis of functional localization in the cortex. Nature Reviews Neuroscience 2002; 3(8): 606-16.

Petrides M, Pandya DN. Distinct parietal and temporal pathways to the homologues of Broca's area in the monkey. PLoS Biol 2009; 7(8): e1000170.

Phillips KA, Stimpson CD, Smaers JB, Raghanti MA, Jacobs B, Popratiloff A, Hof PR, Sherwood CC. The corpus callosum in primates: processing speed of axons and the evolution of hemispheric asymmetry. Proceedings of the Royal Society B: Biological Sciences 2015; 282(1818): 20151535.

Povinelli DJ, Reaux JE, Theall LA, Giambrone S, Humphrey N. Folk physics for apes: The chimpanzee's theory of how the world works: Oxford University Press Oxford; 2000.

Reyes LD. Tool-use and the chimpanzee brain: an investigation of gray and white matter, and a focused study of inferior parietal microstructure: The George Washington University; 2017.

Ringo JL, Doty RW, Demeter S, Simard PY. Time is of the essence: a conjecture that hemispheric specialization arises from interhemispheric conduction delay. Cerebral Cortex 1994; 4(4): 331-43.

Robinson EC, Jbabdi S, Glasser MF, Andersson J, Burgess GC, Harms MP, Smith SM, Van Essen DC,

Jenkinson M. MSM: a new flexible framework for Multimodal Surface Matching. NeuroImage 2014;

100: 414-26.

Rohlfing T, Kroenke CD, Sullivan EV, Dubach MF, Bowden DM, Grant K, Pfefferbaum A. The

INIA19 template and NeuroMaps atlas for primate brain image parcellation and spatial normalization.

Frontiers in Neuroinformatics 2012; 6: 27.

Shen K, Bezgin G, Schirner M, Ritter P, Everling S, McIntosh AR. A macaque connectome for large-

scale network simulations in TheVirtualBrain. Scientific data 2019; 6(1): 1-12.

Smaers JB, Gómez-Robles A, Parks AN, Sherwood CC. Exceptional evolutionary expansion of

prefrontal cortex in great apes and humans. Current Biology 2017; 27(5): 714-20.

Spocter MA, Hopkins WD, Garrison AR, Bauernfeind AL, Stimpson CD, Hof PR, Sherwood CC.

Wernicke's area homologue in chimpanzees (Pan troglodytes) and its relation to the appearance of

modern human language. Proceedings of the Royal Society B: Biological Sciences 2010; 277(1691):

2165-74.

St-Onge E, Daducci A, Girard G, Descoteaux M. Surface-enhanced tractography (SET). NeuroImage

2018; 169: 524-39.

Stout D, Chaminade T. Stone tools, language and the brain in human evolution. Philosophical

Transactions of the Royal Society B: Biological Sciences 2012; 367(1585): 75-87.

Van Essen DC, Dierker DL. Surface-based and probabilistic atlases of primate cerebral cortex. Neuron

2007; 56(2): 209-25.

Van Essen DC, Donahue CJ, Coalson TS, Kennedy H, Hayashi T, Glasser MF. Cerebral cortical folding,

parcellation, and connectivity in humans, nonhuman primates, and mice. Proceedings of the National

Academy of Sciences 2019; 116(52): 26173-80.

Van Essen DC, Smith SM, Barch DM, Behrens TE, Yacoub E, Ugurbil K, Consortium W-MH. The

WU-Minn human connectome project: an overview. NeuroImage 2013; 80: 62-79.

Vigneau M, Beaucousin V, Herve P-Y, Duffau H, Crivello F, Houde O, Mazoyer B, Tzourio-Mazoyer

N. Meta-analyzing left hemisphere language areas: phonology, semantics, and sentence processing.

NeuroImage 2006; 30(4): 1414-32.

Von Bonin G, and Percival Bailey. The neocortex of Macaca mulatta. Urbana: University of Illinois

Press; 1947.

von Economo CF, Koskinas GN. Die cytoarchitektonik der hirnrinde des erwachsenen menschen.

Wien: Springer; 1925.

Wang J, Yang Y, Zhao X, Zuo Z, Tan L-H. Evolutional and developmental anatomical architecture of

the left inferior frontal gyrus. NeuroImage 2020; 222: 117268.

Warrington S, Bryant KL, Khrapitchev AA, Sallet J, Charquero-Ballester M, Douaud G, Jbabdi S,

Mars RB, Sotiropoulos SN. XTRACT-Standardised protocols for automated tractography in the human

and macaque brain. NeuroImage 2020: 116923.

Winkler AM, Ridgway GR, Webster MA, Smith SM, Nichols TE. Permutation inference for the

general linear model. NeuroImage 2014; 92: 381-97.

Wolpert DM, Goodbody SJ, Husain M. Maintaining internal representations: the role of the human

superior parietal lobe. Nature Neuroscience 1998; 1(6): 529-33.

Yarkoni T, Poldrack RA, Nichols TE, Van Essen DC, Wager TD. Large-scale automated synthesis of

human functional neuroimaging data. Nature Methods 2011; 8(8): 665-70.