AGES: An auxin-inducible, GAL4-compatible, gene expression system for *Drosophila*

Colin D. McClure[1,2,3*], Amira Hassan[1,3], Aneisha Duggal[1], Chee Ying Sia[1] and Tony D. Southall[1*]

[1]Department of Life Sciences, Imperial College London, Sir Ernst Chain Building, London, SW7 2AZ, UK

[2] Present address: Queen's University Belfast, Northern Ireland Technology Centre, Belfast, BT9 5AH, UK

[3]These authors contributed equally to this work

[*]Correspondence to t.southall@imperial.ac.uk and c.mcclure@qub.ac.uk

Abstract

The ability to control transgene expression, both spatially and temporally, is essential for studying model organisms. In *Drosophila*, spatial control is primarily provided by the GAL4/UAS system, whilst temporal control relies on a temperature-sensitive GAL80 (which inhibits GAL4) and drug-inducible systems. However, these are not ideal. Shifting temperature can impact on many physiological and behavioural traits, and the current drug-inducible systems are either leaky, toxic, incompatible with existing GAL4-driver lines, or do not generate effective levels of expression. Here we describe the Auxin-inducible Gene Expression System (AGES). AGES relies on the auxin-dependent degradation of a ubiquitously expressed GAL80, and therefore, is compatible with existing GAL4-driver lines. Water-soluble auxin is added to fly food at a low, non-lethal, concentration, which induces expression comparable to uninhibited GAL4 expression. The system works in both larvae and adults, providing a stringent, non-lethal, cost-effective, and convenient method for temporally controlling GAL4 activity in *Drosophila*.

Introduction

The ability to manipulate the expression of specific genes in model organisms has been the cornerstone of genetics research over the last 50 years. It is from such studies that geneticists have been able to elucidate fundamental biological process, such as those underlying neurodegenerative diseases (Lu and Vogel, 2009), oncogenic mechanisms (Villegas, 2019), and neural circuitry involved in memory formation (Cognigni et al., 2018).

The fruit fly, *Drosophila melanogaster*, has been at the forefront of genetics since its cultivation in the lab in the early 1900s, by geneticist Thomas Morgan (Morgan, 1910). In the 100 years since, a plethora of genetic tools have been developed for this model organism, enabling geneticists to tightly control and manipulate gene expression, allowing them to study the roles of genes in development, physiology, and behaviour. One of the most prominent developments is the GAL4/UAS system, which has been an incredibly powerful tool for the spatial control of transgenes in *Drosophila* (Brand and Perrimon, 1993). This system relies on a cloned

or endogenous promoter to drive the expression of the yeast transcription factor GAL4, which can then bind to the UAS (upstream activating sequence) sites upstream of a transgene and activate its expression. Since the development of this system, *Drosophila* researchers have created thousands of genetically distinct lines to enable expression of transgenes in specific populations of cells (for example (Jenett et al., 2012; Robie et al., 2017)). The majority of these lines are publicly available at the Bloomington Drosophila Stock Center (BDSC) and Vienna Drosophila Resource Center (VDRC).

GAL4 activity can be regulated by GAL80, a protein which antagonises GAL4 by inhibiting its activation domain (Lue et al., 1987), and thus the ability of the GAL4 to induce transcript expression (Lee and Luo, 1999; Suster et al., 2004). The generation of a fly line that ubiquitously expressed a temperature-sensitive version of GAL80 (GAL80^{ts}) (Matsumoto et al., 1978) was a significant advance for the field, as it allowed temporal control of GAL4-induced transgene expression (McGuire et al., 2003). Flies can be kept at temperatures permissive of transcript-expression (29^{o}C) or temperatures that inhibit expression (18^{o}C), offering researchers to control not only where, but when transgenes are expressed. The versatility of this system enabled researchers to determine time-specific effects of gene function, and is a system that is widely employed within the *Drosophila* community.

Although the impact of this technology on the field cannot be understated, it is becoming increasingly evident that using a temperature shift to induce gene expression is far from optimal for some experiments, especially in fields such as behaviour, neuropathy, and ageing. Flies, as ectotherms, are highly dependent on their ambient temperature, which affects a wide range of physiological and behavioural traits (Abram et al., 2017). These include sleep (Beckwith and French, 2019), territorial success (Zamudio et al., 1995), lifespan (McCabe and Partridge, 1997), development rate (Danjuma et al., 2014), immunity (Hunt et al., 2016), metabolism (Schou et al., 2017) and even their microbiome (Moghadam et al., 2018). Therefore, the use of a significant temperature shift to induce the expression is far from ideal in these contexts.

As an alternative approach, a range of drug-inducible systems have been developed, where the drug can be administered to flies in their diet to activate transcript expression (Barwell et al., 2017; Kogenaru and Isalan, 2018; Osterwalder et al., 2001; Potter et al., 2010; Sethi and Wang, 2017). These systems each have advantages and disadvantages (see Table 1) and are not often compatible with the GAL4 lines, thus requiring the creation (and optimisation) of their own drivers. In addition, the systems that are compatible with existing GAL4 lines can affect the fly's physiology or survival, thereby limiting their use, and the certainty of any findings when employed.

EXPRESSION SYSTEM	REFERENCE	ADVANTAGES	DISADVANTAGES
GENESWITCH	(Osterwalder et al., 2001)	Non-toxic for adult flies	Not compatible with existing GAL4 lines Leaky expression (Poirier et al., 2008; Scialo et al., 2016) Larval exposure impacts on adult sleep patterns (Li and Stavropoulos, 2016) Drug is unsafe to handle for female researchers Drug is expensive
QF SYSTEM	(Potter et al., 2010)	Non-toxic for flies	Not compatible with existing GAL4 lines
TET-OFF GAL80	(Barwell et al., 2017)	GAL4/UAS compatible	Long induction time (> 5 days) Requires x2 copies of the GAL80 transgene Affects microbiota
TMP INDUCIBLE	(Kogenaru and Isalan, 2018; Sethi and Wang, 2017)	GAL4/UAS compatible	Currently not compatible for general use with existing GAL4 lines. TMP drug is only dissolvable in DMSO (affects survival of larvae) or has to be added as a dry powder to food.

Table 1 – Main advantages and disadvantages of the most common drug-inducible gene expression systems currently available to the *Drosophila* community.

The auxin-degron system, first identified in *Arabidopsis*, involves the auxin-dependent ubiquitination, and subsequent degradation, of proteins which are tagged with a specific auxin-inducible degron (AID) sequence (Dharmasiri et al., 2005; Dharmasiri and Estelle, 2002; Li et al., 2019). Its natural function allows for the prompt elimination of Aux/IAA transcription factors in plants and has been adapted for use in other species, most notably *C. elegans*, to artificially target proteins for rapid degradation (Zhang et al., 2015). In this system, the F-box protein TIR1, an auxin-receptor, binds with conserved proteins Skp1 and Cullin to form a Skp1–Cul1–F-box (SCF) E3 ligase complex which ubiquitinates proteins tagged with an AID sequence in an auxin-dependent manner. The specificity and versatility of the system has made it an effective tool in multiple systems mice (Yesbolatova et al., 2020), and human cell culture (Li et al., 2019). It has also been employed in flies to induce the degradation of specific *Drosophila* proteins (Trost et al., 2016).

Here, we apply the auxin-degron system to enable both temporal and spatial control of transgene expression when combined with the GAL4/UAS system in *Drosophila*. This was achieved by creating a ubiquitously expressed GAL80 fused with AID tags, such that GAL80 is degraded in the presence of the auxin phytohormone (Figure 1). We demonstrate that this system works in both larvae (5 mM auxin) and adult flies (10 mM), which are concentrations that do not affect fly survival or development. This Auxin-inducible Gene Expression System (AGES) is safe to handle, cheap, easy to prepare, provides tuneable and stringent (i.e. not leaky) expression, and most importantly is compatible with the existing GAL4-driver lines developed by the *Drosophila* community.

Results

Generation of an auxin-degradable GAL80 *Drosophila* line

To generate an auxin inducible system for control of GAL4 activity in *Drosophila,* we designed a transgene

that would ubiquitously express two proteins, TIR1, and AID-tagged GAL80. The T2A sequence between TIR1

and AID-tagged GAL80 is a "cleavage" element that is known to work efficiently in *Drosophila* cell culture

(Daniels et al., 2014). The *AtTIR1-T2A-AID-GAL80-AID* sequence was codon optimized, synthesized, and

cloned downstream of the *αTub84B* promoter in *pattB* (Figure 1 and Figure 1 – figure supplement 1)*.* The AID

tag is from the minimal degron of *IAA17* (Morawska and Ulrich, 2013), whose fusion to luciferase resulted in

a fusion protein with short half-life in plants (~ 10 min) (Dreher et al., 2006). The *αTub84B* (tubulin) promoter

drives expression of the bicistronic sequence in all cells, all of the time, and (in the absence of auxin) will

inhibit any GAL4 activity. When present, auxin will tether the TIR1 to the AID sequences, triggering the

degradation of GAL80 and the release of GAL4 inhibition (Figure 1).

Figure 1. Design and action of the AGES system. A tubulin promoter ubiquitously expresses the bicistronic TIR1 and AGES cassette. AID-tagged GAL80 is cleaved from the TIR1 due to the presence of the T2A sequence. In the absence of auxin, GAL80 can inhibit GAL4 activity, whilst the presence of auxin induces degradation of GAL80, allowing GAL4 to drive the expression of UAS-transgene(s).

AGES allows induction of GAL4 activity in adult flies

We combined the AGES system with *c564-GAL4* to see if we could get inducible expression in fat body. A

nuclear localised GFP (nls-GFP) was used as a reporter of GAL4 activity. We first examined adult female flies

that were placed on food containing different concentrations of water-soluble auxin (K-NAA). A negative

control consisting of fly food without auxin (0 mM auxin) was included to investigate whether there was any

leaky GAL4 activity in absence of auxin. After 24 hours, GFP fluorescence in the abdomen were assayed.

Note that there is some naturally occurring autofluorescence in the fly abdomen. At 5 mM and 10 mM, GFP

fluorescence is clearly detectable in the abdomen of live female flies (Figure 2A). Based on this assay, 10 mM

 for 24 hours results in ~60% of the GFP fluorescence levels seen in the positive control (Figure 2B). At 1 mM

 there is no detectable GFP above background fluorescence, and the levels are indistinguishable from both

 the 0 mM and the negative control (lacking both the GAL4 driver and the *UAS-nls-GFP*). To provide an

 alternative and more direct assay of transgene induction, quantitative PCR (qPCR) was used to measure the

 levels of *GFP* mRNA (Figure 2C). Here 5 mM and 10 mM show levels of mRNA equivalent to the positive

 control, while 0 mM showed no significant difference when compared with the negative control. Shorter

 exposure times of adults to auxin containing food were tested (data not shown), however, 24 hours is the

 minimal amount of time required for proper induction of GAL4 activity.

Figure 2. AGES effectively induces GAL4 activity in *Drosophila* adults. A) Ventral images of live females that express GAL4 in fat body tissue. Ingestion of food containing auxin (>= 5 mM for 24 hours) induces GAL4 activity and the expression of GFP. B) Quantification of GFP levels. Pixel intensity thresholding was performed to isolate abdomens as regions of interest. The average pixel intensities from 6 replicates were quantified and analysed using one-way ANOVA (****, $p < 0.0001$). C) qPCR data for *GFP* mRNA levels using different concentrations of auxin. Values were normalised to housekeeping gene *RpL4* (*Ribosomal Protein L4*) and relative expression levels (compared to the negative control) were calculated using the ΔΔCt method. Y-axis displaying ΔΔCt values and statistics done using one-way ANOVA (****, $p < 0.0001$).

Adult males eat less than females (Wong et al., 2009), therefore, this will impact on auxin ingestion and possibly on transgene induction. As with females, we examined both GFP fluorescence levels in the abdomen, and GFP mRNA levels in male flies (Figure 2 – figure supplement 1). Although reduced compared to females, there is still a robust induction of GFP expression.

For AGES to work with GAL4 lines expressed in the central nervous system, auxin must be able to pass the selectively-permeable glial membrane, akin to the mammalian blood-brain barrier (Limmer et al., 2014). To test this, we used *elav-GAL4* (expressed in all neurons) and examined the expression of GFP in the adult brain. Similar to the fat body *c564-GAL4* driver, we observe robust expression of GFP, when flies are fed auxin food (10 mM) for 24 hours (Figure 3). An anti-Elav antibody was included as a counterstain. Furthermore, this demonstrates that AGES works in the central nervous system and that auxin can cross the blood brain barrier in *Drosophila*.

Figure 3. AGES allows induction of GAL4 activity in the *Drosophila* brain. Confocal images of adult brains stained with anti-GFP and anti-Elav. Adult were fed food containing 5 mM auxin for 24 hours. Scale bars represent 20 µm.

AGES allows induction of GAL4 activity in *Drosophila* larvae

In addition to the need for temporal control of GAL4 activity in the adult, researchers might also want to induce expression at specific points during larval development. To test if AGES works in larvae, we used the *c564-GAL4* fat body driver and tested multiple auxin concentrations (Figure 4A). Here we observed that expression could be induced at lower concentrations of auxin. When using food containing no auxin, there is no detectable GFP, whereas with 0.1 mM auxin, GFP is expressed, although at lower levels than with higher concentrations of auxin (Figure 4A). Also, at 0.1 mM, expression is not ubiquitous within the fat body tissue

(Figure 4B). Strong expression is observed for concentrations of 1 mM and above (Figure 4A). Larvae require at least 10 hours of exposure to auxin food before reaching high levels of expression (Figure 4C).

Figure 4. AGES allows induction of GAL4 activity in the *Drosophila* larvae. Bright field images of 3rd instar larvae. A) GFP fluorescence in fat body tissue after induction (for 24 hours) on food with different concentrations of auxin. B) Sporadic GFP expression when using 0.1 mM auxin). C) Representative images of larvae fed on 1 mM auxin and imaged at various time intervals.

Effects of auxin on development and adult survival

For a universally applicable drug-inducible system, the drug should not impact on development or mortality of the flies. We tested whether continuous exposure to different concentrations of auxin would impact on developmental timing. Time to pupation (from egg-laying) is unaffected for concentrations up to 5 mM, however, 10 mM caused a delay of approximately one day (Figure 5A). 10 mM auxin causes some developmental delay, although it has no impact on the survival of the flies through these developmental stages (compared to a no auxin control) (Figure 5B). These data indicate that 10 mM auxin should be avoided for larval induction, however, this is not an issue as > 1 mM is sufficient for robust induction of expression (Figure 4A) and 5 mM does not cause any developmental delay. Concentrations of 5 mM and 10 mM auxin can induce expression in adults (Figure 2 and Figure 2 – figure supplement 1). Survival assays on both male and female adults show that concentrations of 5 mM and 10 mM have no effect on survival and lifespan (Figure 5C and 5D). Therefore, in summary, 5 mM auxin is optimal for use in larvae, whilst 10 mM auxin is best for use in adults.

Figure 5. Impact of auxin on developmental timing and survival. A) Time taken from egg-laying to pupation. B) Survival across developmental stages (larval to pupal and pupal to adult) with 10 mM auxin. C) Survival of adult females during continuous exposure to auxin. D) Survival of adult males during continuous exposure to auxin. Logrank test and weighted Gehan-Breslow-Wilcoxon model (ns) were used for the adult survival assays.

Discussion

The GAL4/UAS/GAL80[ts] systems enable *Drosophila* researchers to control the expression of transgenes both spatially and temporally. However, the impact of the required temperature shift on physiology and behaviour to activate/deactivate the system, can affect the experimental results and subsequent conclusions (Abram et al., 2017). Drug inducible systems have been developed as an alternative to the GAL80[ts] method, however, these are not ideal, limited by issues such as leaky expression, long induction times or incompatibility with existing GAL4 lines. Here we present AGES, an auxin-inducible gene expression system, which eliminates the need for temperature shifts to manipulate temporal expression, and does not possess any of the drawbacks of the current drug-inducible systems.

A significant benefit of AGES over existing systems is its compatibility with existing GAL4-driver lines. Currently, the most prevalent drug-inducible expression systems, i.e. the GeneSwitch (Osterwalder et al.,

2001) and the Q-system (Potter et al., 2010), require the generation of their own driver lines. AGES uses a GAL80 fusion protein, which allows the use of the system alongside the majority of existing GAL4-driver lines, including split-GAL4 lines (Luan et al., 2006; Pfeiffer et al., 2010), whereby suppression of the GAL4-UAS activation is achieved in all cells where auxin is absent. As with all GAL80-based methods of GAL4 control, the GAL4 line must contain the standard GAL4 activation domain (rather than a VP16 or p65 domain) for GAL80 to be affective. We have demonstrated the system using two driver lines, *c564-GAL4* and *elav-GAL4*, which are utilised for fat body and neuron-specific expression respectively. Two different lines were chosen to assess whether the auxin hormone used (K-NAA) could freely access various cell-types within the fruit fly, particularly the central nervous system where auxin must pass the selectively-permeable glial membrane, akin to the mammalian blood-brain barrier (Limmer et al., 2014). Our results demonstrate that feeding flies auxin-containing food (5 mM) for 24 hours is sufficient to drive expression when using both *c564-GAL4* and *elav-GAL4*. Therefore, given that auxin can transverse the blood-brain barrier, this system should, in theory, be compatible with all other GAL4 lines.

Successful gene driver systems require negligible expression when un-induced. While some current systems are known to exhibit leaky expression (Scialo et al., 2016), AGES demonstrates an almost complete repression of GAL4 activity when un-induced. Using GFP fluorescence as a readout, 0 mM auxin does not show any greater signal than the negative control (Figure 4A). Also, the level of *GFP* mRNA without auxin showed no significant difference when compared to the negative control. The AID tags on GAL80 allow the rapid, and tuneable, degradation of the GAL80 protein, providing users with the option to determine the strength of expression by adjusting the concentration of auxin in the food, or by limiting the exposure time of the flies to auxin.

AGES is also a system that is safe for both humans and flies alike. Auxin hormone carries no hazardous concerns for human handling, and no lethal effects were identified for the effective concentrations of 10 mM or below for both adult and larval stages (Figure 5). Continuous exposure of larvae to 10 mM auxin does result in some developmental delay; however, it does not affect survival. Higher concentrations of auxin result in increased mortality rates (data not shown) and should be avoided. Users are encouraged to consider this if attempting to optimise exposure conditions for specific experiments, or GAL4 driver lines that have not been assessed in this paper. Overall, based on our data, we recommend the use of 5 mM auxin and 10 mM for optimal induction in larvae and adults, respectively.

It is also paramount that inducible systems are easy and cheap to apply, to facilitate their practical use in diverse experimental settings. While expenses of the GeneSwitch system average £0.31 (GBP) per vial, AGES costs ~120x less at £0.0025 per vial (authors' estimate). Furthermore, this water-soluble auxin (K-NAA) can be easily added to the food immediately before the food is poured into vials or bottles, requiring no additional safety measures, enabling ease of production. In our experience, auxin-containing food can be stored 4 °C

for up to 4 weeks where the hormone's potency still persists. Further studies are required to assess the stability and effectiveness of auxin when stored for longer periods of time, or in differing conditions.

Despite the various advantages that AGES offers the fly community, there are still limitations to consider. These include the level of expression obtained and the response time of the system (24 hours) and that induced levels are not always at the maximum levels obtained without the system. For example, as male adults eat less than their female counterparts, lower expression is observed in males (Figure S2). However, as GAL4 induced expression is inherently strong, reduced transgene expression for most experiments is unlikely to be an issue. Furthermore, a 24 hour induction time, although not instant, is both practical and convenient for researchers performing experiments in the lab.

The sensitivity of the system is directly related to the amount of auxin present within the tissue, and thus concerns are raised regarding the fly's age and sex, as these factors impact feeding rates (Carey et al., 2006; Wong et al., 2009). As all experimental adult flies were aged to 5 days post-eclosion, the effects of age on the system are yet to be determined, although lower expression is to be expected in older flies. To address these concerns, alterations to the auxin application could be made, such as a longer exposure (> 24 hours), which as we have shown does not impact on survival (Figure 5D and 5E).

In summary, AGES offers the fly community a cheap, safe, and easy system for temporal transgene expression using existing GAL4 lines. Furthermore, it does not require a shift in temperature changes for induction of gene expression. It is a particularly promising tool for research fields where researchers want to avoid using temperature shifts (e.g. aging, behavioural genetics and neuropathology), and will undoubtedly have wide-ranging benefits for multiple fields of study.

Materials and Methods

Generation of *AGES* line

The *AtTIR1-T2A-AID-GAL80-AID* sequence was synthesized by Twist Biosciences (twistbioscience.com) (Figure 1 – Supplement 1). The *GAL80* sequence (from *S. cerevisiae*) and the *AtTIR1* sequence (Zhang et al., 2015) were codon optimized for *Drosophila* for more effective translation. The *αTub84B* promoter was amplified from a *tubulin-eGFP* plasmid (gift from M. Dionne) using the following primers: tub_FWD: GATATCAAGCTTGCACAGGTCC and tub-RV: GTACCTTCACGCTGTGGATGAGG. The *αTub84B* and *AtTIR1-T2A-AID-GAL80-AID* sequences were cloned into *pattB* (Bischof et al., 2007) using Gibson assembly (Gibson et al., 2009). Successful clones were sequenced verified. Annotated sequence is in Figure 1 – figure supplement 2. Microinjection was performed by *Cambridge Fly Facility* using the VK00040 line, which has an *attB* site at location att3B (87B10) on chromosomal arm 3R. Injected adult males were collected and mated to w^{1118} virgin females to identify transgenics (orange eyes). *tub-TIR1-T2A-AID-GAL80-AID* will be available at Bloomington Drosophila Stock Centre or, alternatively, from the corresponding authors.

Fly stocks and food

In this study *c564-GAL4, UAS-nls-GFP / Cyo-Actin-GAL4-GFP* flies were used to perform fluorescent reporter experiments in the adult and larval fat body, and for quantification of *GFP* mRNA expression. Moreover, these flies were used in adult survival assays. *CantonS* flies were used in the developmental survival and developmental timeline experiments. For fluorescent reporter experiments in the adult brain we used *elavGAL4, UAS-nls-GFP*. Flies were kept at 25 °C on and standard *Drosophila* food (recipe in supplementary material) was supplemented (just before being aliquoted into vials/bottles/plates) with auxin (K-NAA (1-Naphthaleneacetic acid potassium salt) available from Phytotech (#N610) or Glentham Life Sciences (GK2088)) at varying concentrations.

Immunohistochemistry

Adult brains were dissected in 1x PBS and fixed for 25 minutes at room temperature in 4% formaldehyde (methanol free) in 0.3% Triton X-100 PBS (PBST). They were washed four times for 1 hour with 0.3% PBST. Normal Goat Serum (2 % in PBST) was used for tissue blocking (RT, 15 min- 1 hour) and subsequent overnight primary antibody incubation. All tissue washes were done in PBST.

Primary antibodies used were chicken anti-GFP (Abcam #13970, 1:2000), and rat anti-Elav 1:500 (Developmental Studies Hybridoma Bank, DSHB). Secondary antibodies used include Alexa Fluor® 488 and 545 at concentration 1:200 (Life technologies) and tissue was incubated for 1.5 hours at room temperature. Tissue was mounted on standard glass slides in Vectashield Mounting medium (Vector Laboratories). Brains were imaged using a Zeiss LSM 510 microscope. Analysis of acquired images was done using Fiji (Schindelin et al., 2012).

Imaging and Image Analysis

Live imaging of GFP in adult flies and larvae was performed using a Nikon SMZ 1500 microscope. In larvae, to identify earliest auxin-induction effects, animals were then placed on 1 mM auxin- supplemented food and imaged at 0, 1, 5, 10 and 25 hour intervals. Adults were placed on food containing varying auxin concentrations for 24 hours prior to imaging.

Live GFP levels were quantified in 6 animals per condition (including negative and positive controls). Binary images were created by using a threshold (Outsu's thresholding) and the Wand Tool in Fiji was used to trace the GFP-positive area as region of interest (ROI). The ROI corresponded to abdomens in adults or whole larvae. Mean pixel intensity per ROI was calculated using the Measure plugin in Fiji for a total of 6 biological replicates. Statistical significance analysed using ONE-way ANOVA. Parametric t-test used to compare individual conditions with respective controls. Statistical tests and plots were performed using the software GraphPad Prism version 9 for Windows.

Developmental and Survival Assays

All animals were kept at 25 °C. To determine the effects of auxin on larval development, 10 mated Canton-S flies were allowed to lay eggs on fly food containing 0, 1, 5, or 10 mM of auxin for a maximum of 24 hours, or until ~50 eggs were counted. 5 replicates were obtained for each concentration, except for 0 mM where 4 replicates were used. Daily emergence of L3 larvae, and presence of pupae were recorded to determine the time required for egg-to-L3, and egg-to-pupae development for each concentration.

Adult survival assays were performed on *c564-GAL4, UAS-nls-GFP / AGES* flies on three replicates, with 20 flies per replicate. Male and female animals were separated 5 days post-eclosion. Each replicate was continuously exposed to 0 mM, 5mM and 10 mM Auxin food. Death was scored daily until all flies were deceased. Statistical tests and plots were performed using the software GraphPad Prism version 9 for Windows.

RNA extraction and RT qPCR

In order to quantify whether levels of *GFP* mRNA expression changed in the presence of Auxin, the *c564-GAL4, UAS-nls-GFP/ AGES* flies were placed on 0 mM, 5 mM and 10 mM auxin-supplemented food for 24 hrs. The *c564-GAL4, UAS-nls-GFP/ +* flies were used as a positive control whilst the *AGES* flies were used as a negative control (lacking both the GAL4 driver and the *UAS-nls-GFP*). The 0 mM was included to investigate whether there was any leaky GAL4 activity in absence of auxin.

Total RNA was extracted from whole adult flies, 3 per replicate per condition, using a standard TRIzol ® extraction protocol (Scientific, 2016). DNA was degraded using RNAse-free DNAse (Thermo Scientific) and cDNA synthesis was performed using the iScript™ cDNA synthesis kit (Bio-Rad), following manufacturer's instructions. Real time quantitative PCR was performed with iTaq™ Universal SYBR® Green Supermix (Bio-Rad) using the StepOnePlus™ Real-Time PCR System (Applied Biosystems).

The gene *RpL4* (*Ribosomal Protein L4*) was used as reference gene. The *GFP* mRNA expression levels were calculated using the ΔΔCt method (ΔCt = Ct [mean Ct of reference gene] – Ct [target]; ΔΔCt = ΔCt [target] – mean ΔCt [control]) (Livak and Schmittgen, 2001). Target refers to the auxin concentrations (0mM, 5mM, 10mM) and control refers to the *AGES* flies. Primers used: RpL4_FW - 5'-TCCACCTTGAAGAAGGGCTA-3', RpL4_RV – 5'-TTGCGGATCTCCTCAGACTT-3', GFP_FW – 5'-GAGCTGTACAAGAGCAGGCA-3', GFP_RV – 5'-GTTGACGGCGTTTTCGTTCA-3'.

Acknowledgements

Thanks to Marc Dionne (Imperial College London) for providing the *tubulin-eGFP* plasmid, and to Christian Lehner (University of Zurich) for providing the *pMT-OsTIR1-P2A-H2B-aid-eYFP* plasmid as well invaluable insights into the molecular biology of the auxin-degradation system in *Drosophila*. Furthermore, thank you to the entire Southall lab for helpful discussions and pre-reading the manuscript. This work was funded by Wellcome Trust Investigator grant 104567 to T.D.S. and an Innovation Grant from the Society of Developmental Biology to C.D.M.

Figure Legends

Figure 1. Design and action of the AGES system. A tubulin promoter ubiquitously expresses the bicistronic TIR1 and AGES cassette. AID-tagged GAL80 is cleaved from the TIR1 due to the presence of the T2A sequence. In the absence of auxin, GAL80 can inhibit GAL4 activity, whilst the presence of auxin induces degradation of GAL80, allowing GAL4 to drive the expression of UAS-transgene(s).

Figure 2. AGES effectively induces GAL4 activity in *Drosophila* adults. A) Ventral images of live females that express GAL4 in fat body tissue. Ingestion of food containing auxin (>= 5 mM for 24 hours) induces GAL4 activity and the expression of GFP. B) Quantification of GFP levels. Pixel intensity thresholding was performed to isolate abdomens as regions of interest. The average pixel intensities were quantified and analysed using one-way ANOVA (****, $p < 0.0001$). C) qPCR data for *GFP* mRNA levels using different concentrations of auxin. Values were normalised to housekeeping gene RpL4 (Ribosomal Protein L4) and relative expression levels (compared to the negative control) were calculated using the ΔΔCt method. Y-axis displaying ΔΔCt values and statistics done using one-way ANOVA (****, $p < 0.0001$).

Figure 3. AGES allows induction of GAL4 activity in the *Drosophila* brain. Confocal images of adult brains stained with anti-GFP and anti-Elav. Scale bars represent 20 μm.

Figure 4. AGES allows induction of GAL4 activity in the *Drosophila* larvae. Bright field images of 3rd instar larvae. A) GFP fluorescence in fat body tissue after induction (for 24 hours) on food with different concentrations of auxin. B) Sporadic GFP expression when using 0.1 mM auxin). C) Representative images of larvae fed on 1 mM auxin and imaged at various time intervals.

Figure 5. Impact of auxin on developmental timing and survival. A) Time taken from egg-laying to pupation. B) Survival across developmental stages (larval to pupal and pupal to adult) with 10 mM auxin. C) Survival of adult females during continuous exposure to auxin. D) Survival of adult males during continuous exposure to auxin. Logrank test and weighted Gehan-Breslow-Wilcoxon model (ns) were used for the adult survival assays.

Figure 1 – figure supplement 1. Plasmid map of the AGES plasmid. Map of the *pattB-tubP-AtTIR1-T2A-miniAID-Gal80-miniAID-SV40* plasmid (generated using Snapgene). The full sequence is in Figure 1 – figure supplement 2.

Figure 1 – figure supplement 2.

Genbank (.gb) sequence file for *pattB-tubP-AtTIR1-T2A-miniAID-Gal80-miniAID-SV40*

Figure 2 – figure supplement 1.

AGES effectively induces GAL4 activity in *Drosophila* adult males. A) Ventral images of live males that express GAL4 in fat body tissue. Ingestion of food containing auxin (24 hours) induces GAL4 activity and the expression of GFP. B) Quantification of GFP levels. Pixel intensity thresholding was performed to isolate abdomens as regions of interest. The average pixel intensity intensities were quantified and analysed using Ordinary one-way ANOVA (**, $p < 0.01$, ****, $p < 0.0001$). C) qPCR data for *GFP* mRNA levels using different concentrations of auxin. Values were normalised to housekeeping gene *RpL4* (*Ribosomal Protein L4*) and relative expression levels were calculated using the ΔΔCt method. Y-axis displaying ΔΔCt values and statistics done using Ordinary one-way ANOVA (**, $p < 0.01$).

References

Abram, P.K., Boivin, G., Moiroux, J., Brodeur, J., 2017. Behavioural effects of temperature on ectothermic animals: unifying thermal physiology and behavioural plasticity. *Biol Rev Camb Philos Soc* **92**, 1859-1876.

Barwell, T., DeVeale, B., Poirier, L., Zheng, J., Seroude, F., Seroude, L., 2017. Regulating the UAS/GAL4 system in adult Drosophila with Tet-off GAL80 transgenes. *PeerJ* **5**, e4167.

Beckwith, E.J., French, A.S., 2019. Sleep in Drosophila and Its Context. *Front Physiol* **10**, 1167.

Bischof, J., Maeda, R.K., Hediger, M., Karch, F., Basler, K., 2007. An optimized transgenesis system for Drosophila using germ-line-specific phiC31 integrases. *Proc Natl Acad Sci U S A* **104**, 3312-3317.

Brand, A.H., Perrimon, N., 1993. Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development* **118**, 401-415.

Carey, J.R., Papadopoulos, N., Kouloussis, N., Katsoyannos, B., Muller, H.G., Wang, J.L., Tseng, Y.K., 2006. Age-specific and lifetime behavior patterns in Drosophila melanogaster and the Mediterranean fruit fly, Ceratitis capitata. *Exp Gerontol* **41**, 93-97.

Cognigni, P., Felsenberg, J., Waddell, S., 2018. Do the right thing: neural network mechanisms of memory formation, expression and update in Drosophila. *Curr Opin Neurobiol* **49**, 51-58.

Daniels, R.W., Rossano, A.J., Macleod, G.T., Ganetzky, B., 2014. Expression of multiple transgenes from a single construct using viral 2A peptides in Drosophila. *PLoS One* **9**, e100637.

Danjuma, S., Thaochan, N., Permkam, S., Satasook, C., 2014. Effect of temperature on the development and survival of immature stages of the carambola fruit fly, Bactrocera carambolae, and the Asian papaya fruit fly, Bactrocera papayae, reared on guava diet. *J Insect Sci* **14**, 126.

Dharmasiri, N., Dharmasiri, S., Estelle, M., 2005. The F-box protein TIR1 is an auxin receptor. *Nature* **435**, 441-445.

Dharmasiri, S., Estelle, M., 2002. The role of regulated protein degradation in auxin response. *Plant Mol Biol* **49**, 401-409.

Dreher, K.A., Brown, J., Saw, R.E., Callis, J., 2006. The Arabidopsis Aux/IAA protein family has diversified in degradation and auxin responsiveness. *Plant Cell* **18**, 699-714.

Gibson, D.G., Young, L., Chuang, R.Y., Venter, J.C., Hutchison, C.A., 3rd, Smith, H.O., 2009. Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat Methods* **6**, 343-345.

Hunt, V.L., Zhong, W., McClure, C.D., Mlynski, D.T., Duxbury, E.M., Keith Charnley, A., Priest, N.K., 2016. Cold-seeking behaviour mitigates reproductive losses from fungal infection in Drosophila. *J Anim Ecol* **85**, 178-186.

Jenett, A., Rubin, G.M., Ngo, T.T., Shepherd, D., Murphy, C., Dionne, H., Pfeiffer, B.D., Cavallaro, A., Hall, D., Jeter, J., Iyer, N., Fetter, D., Hausenfluck, J.H., Peng, H., Trautman, E.T., Svirskas, R.R., Myers, E.W., Iwinski, Z.R., Aso, Y., DePasquale, G.M., Enos, A., Hulamm, P., Lam, S.C., Li, H.H., Laverty, T.R., Long, F., Qu, L., Murphy, S.D., Rokicki, K., Safford, T., Shaw, K., Simpson, J.H., Sowell, A., Tae, S., Yu, Y., Zugates, C.T., 2012. A GAL4-driver line resource for Drosophila neurobiology. *Cell Rep* **2**, 991-1001.

Kogenaru, M., Isalan, M., 2018. Drug-Inducible Control of Lethality Genes: A Low Background Destabilizing Domain Architecture Applied to the Gal4-UAS System in Drosophila. *ACS Synth Biol* **7**, 1496-1506.

Lee, T., Luo, L., 1999. Mosaic analysis with a repressible cell marker for studies of gene function in neuronal morphogenesis. *Neuron* **22**, 451-461.

Li, Q., Stavropoulos, N., 2016. Evaluation of Ligand-Inducible Expression Systems for Conditional Neuronal Manipulations of Sleep in Drosophila. *G3 (Bethesda)* **6**, 3351-3359.

Li, S., Prasanna, X., Salo, V.T., Vattulainen, I., Ikonen, E., 2019. An efficient auxin-inducible degron system with low basal degradation in human cells. *Nat Methods* **16**, 866-869.

Limmer, S., Weiler, A., Volkenhoff, A., Babatz, F., Klambt, C., 2014. The Drosophila blood-brain barrier: development and function of a glial endothelium. *Front Neurosci* **8**, 365.

Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25**, 402-408.

Lu, B., Vogel, H., 2009. Drosophila models of neurodegenerative diseases. *Annu Rev Pathol* **4**, 315-342.

Luan, H., Lemon, W.C., Peabody, N.C., Pohl, J.B., Zelensky, P.K., Wang, D., Nitabach, M.N., Holmes, T.C., White, B.H., 2006. Functional dissection of a neuronal network required for cuticle tanning and wing expansion in Drosophila. *J Neurosci* **26**, 573-584.

Lue, N.F., Chasman, D.I., Buchman, A.R., Kornberg, R.D., 1987. Interaction of GAL4 and GAL80 gene regulatory proteins in vitro. *Mol Cell Biol* **7**, 3446-3451.

Matsumoto, K., Toh-e, A., Oshima, Y., 1978. Genetic control of galactokinase synthesis in Saccharomyces cerevisiae: evidence for constitutive expression of the positive regulatory gene gal4. *J Bacteriol* **134**, 446-457.

McCabe, J., Partridge, L., 1997. An Interaction between Environmental Temperature and Genetic Variation for Body Size for the Fitness of Adult Female Drosophila Melanogaster. *Evolution* **51**, 1164-1174.

McGuire, S.E., Le, P.T., Osborn, A.J., Matsumoto, K., Davis, R.L., 2003. Spatiotemporal rescue of memory dysfunction in Drosophila. *Science* **302**, 1765-1768.

Moghadam, N.N., Thorshauge, P.M., Kristensen, T.N., de Jonge, N., Bahrndorff, S., Kjeldal, H., Nielsen, J.L., 2018. Strong responses of Drosophila melanogaster microbiota to developmental temperature. *Fly (Austin)* **12**, 1-12.

Morawska, M., Ulrich, H.D., 2013. An expanded tool kit for the auxin-inducible degron system in budding yeast. *Yeast* **30**, 341-351.

Morgan, T.H., 1910. Sex Limited Inheritance in Drosophila. *Science* **32**, 120-122.

Osterwalder, T., Yoon, K.S., White, B.H., Keshishian, H., 2001. A conditional tissue-specific transgene expression system using inducible GAL4. *Proc Natl Acad Sci U S A* **98**, 12596-12601.

Pfeiffer, B.D., Ngo, T.T., Hibbard, K.L., Murphy, C., Jenett, A., Truman, J.W., Rubin, G.M., 2010. Refinement of tools for targeted gene expression in Drosophila. *Genetics* **186**, 735-755.

Poirier, L., Shane, A., Zheng, J., Seroude, L., 2008. Characterization of the Drosophila gene-switch system in aging studies: a cautionary tale. *Aging Cell* **7**, 758-770.

Potter, C.J., Tasic, B., Russler, E.V., Liang, L., Luo, L., 2010. The Q system: a repressible binary system for transgene expression, lineage tracing, and mosaic analysis. *Cell* **141**, 536-548.

Robie, A.A., Hirokawa, J., Edwards, A.W., Umayam, L.A., Lee, A., Phillips, M.L., Card, G.M., Korff, W., Rubin, G.M., Simpson, J.H., Reiser, M.B., Branson, K., 2017. Mapping the Neural Substrates of Behavior. *Cell* **170**, 393-406 e328.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.Y., White, D.J., Hartenstein, V., Eliceiri, K., Tomancak, P., Cardona, A., 2012. Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**, 676-682.

Schou, M.F., Kristensen, T.N., Pedersen, A., Karlsson, B.G., Loeschcke, V., Malmendal, A., 2017. Metabolic and functional characterization of effects of developmental temperature in Drosophila melanogaster. *Am J Physiol Regul Integr Comp Physiol* **312**, R211-R222.

Scialo, F., Sriram, A., Stefanatos, R., Sanz, A., 2016. Practical Recommendations for the Use of the GeneSwitch Gal4 System to Knock-Down Genes in Drosophila melanogaster. *PLoS One* **11**, e0161817.

Scientific, T.F., 2016. Available: https://tools.thermofisher.com/content/sfs/manuals/trizol_reagent.pdf.

Sethi, S., Wang, J.W., 2017. A versatile genetic tool for post-translational control of gene expression in Drosophila melanogaster. *Elife* **6**.

Suster, M.L., Seugnet, L., Bate, M., Sokolowski, M.B., 2004. Refining GAL4-driven transgene expression in Drosophila with a GAL80 enhancer-trap. *Genesis* **39**, 240-245.

Trost, M., Blattner, A.C., Lehner, C.F., 2016. Regulated protein depletion by the auxin-inducible degradation system in Drosophila melanogaster. *Fly (Austin)* **10**, 35-46.

Villegas, S.N., 2019. One hundred years of Drosophila cancer research: no longer in solitude. *Dis Model Mech* **12**.

Wong, R., Piper, M.D., Wertheim, B., Partridge, L., 2009. Quantification of food intake in Drosophila. *PLoS One* **4**, e6063.

Yesbolatova, A., Saito, Y., Kitamoto, N., Makino-Itou, H., Ajima, R., Nakano, R., Nakaoka, H., Fukui, K., Gamo, K., Tominari, Y., Takeuchi, H., Saga, Y., Hayashi, K.I., Kanemaki, M.T., 2020. The auxin-inducible degron 2 technology provides sharp degradation control in yeast, mammalian cells, and mice. *Nat Commun* **11**, 5701.

452 **Zamudio, K.R., Huey, R.B., Crill, W.D.**, 1995. Bigger isn't always better: body size, developmental and
453 parental temperature and male territorial success in Drosophila melanogaster. *Animal Behaviour* **49**, 671-
454 677.

455 **Zhang, L., Ward, J.D., Cheng, Z., Dernburg, A.F.**, 2015. The auxin-inducible degradation (AID) system enables
456 versatile conditional protein depletion in C. elegans. *Development* **142**, 4374-4384.