Title: Comprehensive analysis of male-free reproduction in *Monomorium triviale*

(Formicidae: Myrmicinae)

Authors: Naoto Idogawa[1]*, Tomonori Sasaki[2, 3], Kazuki Tsuji[3] and Shigeto Dobata[1, 4]*

Affiliations:

[1] Laboratory of Insect Ecology, Graduate School of Agriculture, Kyoto University,

Kitashirakawa-Oiwakecho, Sakyo-ku, Kyoto 606-8502, Japan.

[2] Research and Development Department, Fumakilla Limited, 1-11-13, Umehara,

Hatsukaichi, Hiroshima 739-0494, Japan.

[3] Department of Subtropical Agro-Environmental Sciences, Faculty of Agriculture, University

of the Ryukyus, 1, Senbaru, Nishihara, Okinawa 903-0213, Japan.

[4] Department of General Systems Studies, Graduate School of Arts and Sciences, The

University of Tokyo, 3-8-1, Komaba, Meguro-ku, Tokyo 153-8902, Japan.

***Corresponding authors**

E-mail: idogawa.naoto.na@alumni.tsukuba.ac.jp

E-mail: dobata@g.ecc.u-tokyo.ac.jp

Abstract

We report comprehensive evidence for obligatory thelytokous parthenogenesis in an ant *Monomorium triviale*. This species is characterized by distinct queen–worker dimorphism with strict reproductive division of labor: queens produce both workers and new queens without mating, whereas workers are completely sterile. We collected 333 nests of this species from 14 localities and three laboratory-reared populations in Japan. All wild queens dissected had no sperm in their spermathecae. Laboratory observation confirmed that virgin queens produced workers without mating. Furthermore, microsatellite genotyping showed identical heterozygous genotypes between mothers and their respective daughters, suggesting an extremely low probability of sexual reproduction. Microbial analysis detected no bacterial genera that are known to induce thelytokous parthenogenesis in Hymenoptera. Finally, the lack of variation in partial sequences of mitochondrial DNA among individuals sampled from across Japan suggests recent rapid spread or selective sweep. *M. triviale* would be a promising model system of superorganism-like adaptation through comparative analysis with well-studied sexual congeners, including the pharaoh ant *M. pharaonis*.

Introduction

Hymenopteran insects are characterized by haplo-diploid sex determination systems, in which females are derived from fertilized diploid eggs and males are derived from unfertilized haploid eggs via arrhenotokous parthenogenesis [1]. However, the production of diploid females from unfertilized eggs—known as thelytokous parthenogenesis—has been reported sporadically across the Hymenoptera [2]. Thelytoky has provided an opportunity for empirical testing of evolutionary hypotheses to account for the near-ubiquity of sex among eukaryotes [3]. The last decade has seen an increasing number of known thelytokous Hymenoptera, especially in eusocial species [4]. In studies taking advantage of this simplified mode of reproduction, thelytokous social insects have been acknowledged as a model system in behavioral ecology (e.g., [5–8]) and sociogenomics (e.g., [9]).

Previous studies of thelytokous parthenogenesis of ants have identified three distinct categories [10, 11]: (type I) queens produce workers sexually but daughter queens via thelytoky; (type II) the queen caste is usually lost, males are absent, and workers produce workers via thelytoky; and (type III) queens produce both workers and queens via thelytoky, males are usually absent, and workers are sterile.

In this paper, we report comprehensive evidence of the third type of thelytokous parthenogenesis, namely obligatory thelytoky by queens of *M. triviale* [12]. The genus *Monomorium* is one of the most species-rich genera among ants [13] and has a worldwide distribution [14]. With the marked exception of tramp species such as the pharaoh ant *M. pharaonis* [15], most *Monomorium* species remain to be investigated [16]. *M. triviale* is reported from East Asia, encompassing Japan, South Korea and mainland China [14]. Neither males nor inseminated females have been found to date; this has been referred to in several

studies [1, 4, 17–23]. Surprisingly, however, direct evidence of thelytoky is lacking. We

examined the reproductive system of *M. triviale* through multiple approaches, namely

dissection of spermathecae of field-collected queens, direct observation of virgin queen

reproduction, and microsatellite genotyping of mothers and daughters. Furthermore, we

analyzed for the presence of parthenogenesis-inducing microsymbionts and within-species

phylogenetic relationships among populations in Japan.

Materials & Methods

Colony sampling

A total of 333 nests of *M. triviale* were collected from 14 local populations in Japan

(Fig 1, Table 1). In the field, nests were found in small gaps in dead plant bodies such as

rotten roots, dead twigs, hollow bamboo sticks, and acorns. Because of the small nest size

(about 200 workers on average) it was easy to collect the whole nest. During the field

investigation, males of *M. triviale* were never found. Colonies were transferred into artificial

nests in the laboratory as soon as possible and were kept at 25 °C until the following

experiments. In addition, laboratory-reared nests and ethanol-preserved samples originating

from three other sites and provided by other researchers were examined (Fig 1).

Fig 1. Localization of the 14 sampling sites (1 to 14) and the three sites of origin of

laboratory-reared populations (15 to 17) in Japan. Open circles indicate site locations. Site

names, numbers of nests collected, and types of analyses are described in **Table 1**.

83 **Table 1. Sampling sites of *Monomorium triviale* populations and experiments performed**

	Locality	Latitude	Longitude	Collection dates	No. of nests	Experiments
1	Kyoto City, Kyoto Prefecture	35.060087	135.788488	19/04/2017 to 27/11/2017 (24 times)	176	SP, VR, SEQ
2	Tsuchiura City, Ibaraki Prefecture	36.077927	140.165473	20/06/2017, 09/10/2018	28	SP, VR, SEQ
3	Chofu City, Tokyo Metropolis	35.668182	139.549061	16/06/2017, 27/09/2018	28	SP, VR, SEQ
4	Tsukuba City, Ibaraki Prefecture	36.100415	140.101297	19/06/2017, 08/10/2018	26	SP, VR, SEQ
5	Otsu City, Shiga Prefecture	34.970262	135.956026	10/06/2017	26	SP, VR, SEQ
6	Matsudo City, Chiba Prefecture	35.774787	139.899362	18/06/2017, 28/09/2018	22	SP, VR, SEQ
7	Higashiosaka City, Osaka Prefecture	34.665748	135.671267	09/11/2017	5	SP, SEQ
8	Okazaki City, Aichi Prefecture	34.941529	137.175594	03/09/2017	1	SP, SEQ
9	Sanda City, Hyogo Prefecture	34.914575	135.165764	09/05/2017	1	VR, SEQ
10	Kaizu City, Gifu Prefecture	35.22	136.63	06/06/2018	1	SP
11	Tobishima Island, Sakata City, Yamagata Prefecture	39.19	139.55	20/09/2018	2	SP
12	Takamatsu City, Kagawa Prefecture	34.364349	134.098205	17/06/2017	16	SEQ
13	Matsuyama City, Ehime Prefecture	33.845539	132.765722	19/09/2018	2	SEQ
14	Takashima City, Shiga Prefecture	35.3677	135.9168	30/05/2017	1	SEQ
15	Kanonji City, Kagawa Prefecture	34.12	133.66	Reared in Laboratory	–	SEQ
16	Kagamihara City, Gifu Prefecture	35.40	136.85	Reared in Laboratory	–	SEQ
17	Matsue City, Shimane Prefecture	35.47	133.05	Reared in Laboratory	–	SEQ
				Total	**333**	

The nests from Kaizu City (locality no. 10) and Tobishima Island (no. 11) were provided by Dr. K.

Ohkawara. Samples from Higashiosaka City (no. 7) and Takashima City (no. 14) were provided by Mr.

K. Sadahiro and Dr. T. Nozaki, respectively. Ethanol-preserved samples from Kanonji City,

Kagamihara City, and Matsue City (nos. 15 to 17) were maintained in the laboratory for several years

and were provided by Dr. F. Ito. Experiments on the samples from the 17 sites are abbreviated as

follows: SP: dissection of wild queens' spermathecae; VR: observation of virgin queen reproduction;

SEQ: phylogenetic analysis based on mtDNA sequencing.

Dissection of wild queens

To confirm the reproductive status of the queens from wild nests, we dissected 63

individuals from 38 nests of 10 *M. triviale* populations within 6 months after collection (Table

2). First, the queen was immobilized by being soaked in 70% ethanol for 3 min. The body was

then transferred to a 30-mm petri dish filled with distilled water, and the internal reproductive

organs were pulled out from the end of the abdomen by using precision forceps under a

binocular microscope (SZ40; OLYMPUS Optical, Tokyo, Japan). The mating status of the

queen was determined by the presence or absence of sperm in the spermatheca. As an

indicator of oviposition, the ovary's yellow body was checked. As a positive control [18], 11

queens of a congeneric species, *M. intrudens*, were dissected in the same manner.

Table 2. Dissected queens from each sampling site

	Locality	No. of nests	No. of queens	Insemination		Yellow body	
				yes	no	yes	unclear
1	Kyoto City, Kyoto Prefecture	3	10	0	10	9	1
2	Tsuchiura City, Ibaraki Prefecture	8	8	0	8	7	1
3	Chofu City, Tokyo Metropolis	8	8	0	8	5	3
4	Tsukuba City, Ibaraki Prefecture	8	8	0	8	3	5

	Locality						
5	Otsu City, Shiga Prefecture	2	9	0	9	9	0
6	Matsudo City, Chiba Prefecture	4	8	0	8	8	0
7	Higashiosaka City, Osaka Prefecture	1	1	0	1	1	0
8	Okazaki City, Aichi Prefecture	1	8	0	8	8	0
10	Kaizu City, Gifu Prefecture	1	1	0	1	1	0
11	Tobishima Island, Sakata City, Yamagata Prefecture	2	2	0	2	1	1
	Total	38	63	0	63	52	11
–	*Monomorium intrudens*, Kyoto City, Kyoto Prefecture	4	11	10	1	11	0

103   The numbers of individuals possessing a yellow body and presumed to have experienced oviposition are

104   also shown. As a positive control, data for the sexual congener *M. intrudens* are shown in the bottom row.

Rearing experiment

Each wild nest was put into a plastic container (68 × 39 × 15 mm) with gypsum on

the bottom. The nests were kept in the laboratory at 25 °C, and water and food were

replenished every 3 days. The nests were fed mealworms, *Tenebrio molitor*, cut into

approximately 5-mm lengths. During July and August 2017, a total of 44 queen broods

(larvae or pupae) from 21 nests of seven populations were produced (Table 3). Each queen

was isolated with 10 nestmate workers in a plastic container (36 × 36 × 14 mm). These nests

were kept under the same conditions as the source nests. Reproduction by these virgin queens

was observed for 6 months. Finally, all remaining queens were dissected and confirmed to

have no sperm in their spermathecae (Table 4).

Table 3. Thelytokous worker production in the rearing experiment

Locality	No. of nests	No. of queens isolated	No. of queens surviving for 6 months	No. of workers produced

1	Kyoto City, Kyoto Prefecture	6	10	6	17
2	Tsuchiura City, Ibaraki Prefecture	3	7	2	11
3	Chofu City, Tokyo Metropolis	3	5	5	26
4	Tsukuba City, Ibaraki Prefecture	3	5	5	28
5	Otsu City, Shiga Prefecture	3	8	8	52
6	Matsudo City, Chiba Prefecture	2	7	8	28
9	Sanda City, Hyogo Prefecture	1	2	2	15
	Total	**21**	**44**	**36**	**177**

Table 4. Dissection of virgin queens from the rearing experiment

	Locality	No. of queens dissected	Insemination		Yellow body	
			yes	no	yes	unclear
2	Tsuchiura City, Ibaraki Prefecture	2	0	2	2	0
3	Chofu City, Tokyo Metropolis	5	0	5	3	2
4	Tsukuba City, Ibaraki Prefecture	5	0	5	2	3
5	Otsu City, Shiga Prefecture	8	0	8	3	5
6	Matsudo City, Chiba Prefecture	7	0	7	2	5
9	Sanda City, Hyogo Prefecture	2	0	2	1	1
	Total	**29**	**0**	**29**	**13**	**16**

Microsatellite analysis

To provide genetic evidence of thelytoky, 33 queens (from 17 nests of seven populations; all used in the rearing experiment) were examined (Table 5). They were genotyped at microsatellite locus *Mp-1* (f: GCCAATGGTTTAATCCCTCA; r: TCATACTGCGTGTGCCTTTC), originally developed from *M. pharaonis* [24]. Daughter workers produced from these virgin queens (174 individuals in total) were also genotyped and were compared with their mothers. The thorax of each individual was crushed and placed in a 0.2-mL microtube filled with 100 µL of a DNA extraction reagent (PrepMan Ultra Reagent, Applied Biosystems, Foster City, CA, USA). The polymerase chain reaction (PCR) cocktail contained 1 µL of template DNA, 0.3 µL of 25 mM $MgCl_2$, 0.3 µL of 10 mM dNTPs, 1.5 µL

of 10× PCR Buffer, 0.1 µL of 5 U/µL Taq DNA Polymerase (QIAGEN, Valencia, CA, USA),

0.2 µL of U19 fluorescent dye, and 1.0 µL of each primer pair, to which distilled water was

added to make a total volume of 15.2 µL. The PCR program consisted of an initial step of

94 °C for 180 s, followed by 35 cycles of 94 °C for 30 s, 57 °C for 60 s and 72 °C for 60 s,

with a final step of 72 °C for 10 min. PCR products were mixed with Hi-Di formamide and

GS-600 LIZ size standard and were analyzed by using a 3500 Series Genetic Analyzer and

GeneMapper 5.0 software (Applied Biosystems).

Table 5. Microsatellite analysis data

	Locality	No. of nests	No. of queens	No. of workers	Genotype	Prob. that all offspring were heterozygous
1	Kyoto City, Kyoto Prefecture	4	4	15	222/234	$(1/2)^{15} = 3.05 \times 10^{-5}$
2	Tsuchiura City, Ibaraki Prefecture	1	2	11	220/232	$(1/2)^{11} = 4.88 \times 10^{-4}$
3	Chofu City, Tokyo Metropolis	3	5	26	222/232	$(1/2)^{26} = 1.49 \times 10^{-8}$
4	Tsukuba City, Ibaraki Prefecture	3	5	28	220/232	$(1/2)^{28} = 3.73 \times 10^{-9}$
5	Otsu City, Shiga Prefecture	3	8	52	220/228	$(1/2)^{52} = 2.22 \times 10^{-16}$
6	Matsudo City, Chiba Prefecture	2	7	27	220/234	$(1/2)^{27} = 7.45 \times 10^{-9}$
9	Sanda City, Hyogo Prefecture	1	2	15	222/238	$(1/2)^{15} = 3.05 \times 10^{-5}$
	Total	**17**	**33**	**174**	−	$\mathbf{(1/2)^{174} = 4.18 \times 10^{-53}}$

138   Genotypes at the *Mp-1* locus are shown. Under the assumption of sexual reproduction, the probability that

139   all daughter workers would be heterozygous was calculated.

Microbial analysis

To assess potential infection of *M. triviale* by thelytoky-inducing bacteria, we

performed high-throughput amplicon sequencing by using whole bodies of adults. In August

2020, three *M. triviale* nests were collected in Takaragaike Park, Kyoto, Japan (lat 35.060087,

long 135.788488). All colonies were transported to our laboratory and moved to plastic

containers (68 × 39 × 30 mm) with gypsum on the bottom. They were maintained at 25 °C,

and water replenishment and mealworm-feeding were performed every 3 days. In October

2020, ten queens and 100 workers were picked up from each nest. The queens or workers

from each nest were stored as a group in acetone at −30 °C. The pooled individuals

represented one biological replicate (i.e., three replicates per caste). Whole bodies in each

replicate were air-dried and pooled in a 1.5-mL plastic tube, and DNA was extracted with a

QIAamp DNA Micro Kit (QIAGEN). We followed the manufacturer's instructions and added

extra steps of thermal cycling and lysozyme to the protocol to ensure the lysis of

Gram-positive bacterial cell walls: After being crushed in 180 μL ATL buffer, samples in

plastic tubes first underwent two cycles of −80 °C for 30 min and 50 °C for 5 min; then, under

room temperature, we added 2 μL of lysozyme from egg white (Nakalai Tesque, Kyoto,

Japan; 20 μg/μL TE buffer) to each sample and incubated the samples at 37 °C for 30 min.

16S rRNA amplicon sequencing and the subsequent data analysis were performed according

to in-house workflow by Bioengineering Lab. Co., Ltd. (Sagamihara, Kanagawa, Japan). The

V4 region was amplified from 1 ng of template DNA by using ExTaq HS (Takara Bio, Otsu,

Shiga, Japan) polymerase and the 515f–806r primer pair. Sequences were determined by

using a MiSeq system with a MiSeq reagent kit v3 (Illumina, San Diego, CA, USA), which

generated 2 × 300-bp paired-end reads.

Demultiplexing (on the basis of a perfect match with the primer sequences used), adapter

trimming, and quality filtering (phred score ≥ 20; primer sequences, 50 bp on both 3′-ends,

noise and chimeric reads were removed) of the paired-end reads were performed by using a

Fastx toolkit (ver. 0.0.14, http://hannonlab.cshl.edu/fastx_toolkit/) and dada2 plugin for

Qiime2 (ver. 2020.8) [25], resulting in representative sequences and operational taxonomic

unit (OTU) tables. Assignment of appropriate taxa (confidence level > 0.7) was performed by

using the default setting of the feature-classifier plugin for Qiime2 against the EzBioCloud

16S reference database (https://www.ezbiocloud.net/).

Phylogenetic analysis

To investigate intraspecific diversity and determine the phylogenetic position of *M.*

triviale, phylogenetic analysis was performed. A 639-bp region of the *cytochrome oxidase I*

(*COI*) sequence was determined by using PCR with a combination of primers, namely

LCO1490 (GGTCAACAAATCATAAGATATTGG) and HCO2198

(TAAACTTCAGGGTGACCAAAAAATCA) [26]. One worker randomly chosen from each

of 15 local populations (Table 1) of *M. triviale* and a single worker of the sexual species, *M.*

intrudens, collected in Kihoku-cho, Mie, Japan (lat 34.21, long 136.33), were sequenced.

Protocols for DNA extraction and the PCR mix were the same as for the microsatellite

analysis. The thermal cycle consisted of an initial denaturation at 94 °C for 1 min, 35 cycles

of denaturation at 94 °C for 30 s, annealing at 50 °C for 30 s, extension at 72 °C for 60 s, and

a final extension at 72 °C for 10 min. PCR products were ethanol-precipitated and sequenced

in both directions by using a BigDye Terminator v3.1 cycle sequencing kit on a 3500 Series

Genetic Analyzer (Applied Biosystems). A total of 16 sequences were submitted to the DNA

Data Bank of Japan under accession numbers LC592050 to LC592065; see Fig 3). To identify

the phylogenetic position of *M. triviale* after recent advances in systematics of the subfamily

Myrmicinae [27], a congeneric species, *M. pharaonis* (GenBank accession number

GU710434) and two former *Monomorium* species, *Syllophopsis sechellensis* (EF609858) and

Erromyrma latinodis (GU709833) were added to the analysis. The red imported fire ant,

Solenopsis invicta (HQ928672), was used as an outgroup. The 16 sequences, together with

those of the above four species, were aligned by using ClustalW [28] in MEGA 10.0 [29].

Maximum likelihood analyses were conducted as implemented in MEGA 10.0 by using a

GTR+I+Γ model with node support assessed with 1000 bootstrap replicates.

Results

Dissection of wild queens

All 63 dissected queens of *M. triviale* had no sperm in their spermathecae (Fig 2,

Table 2). Yellow bodies, suggestive of queen oviposition experience, were identified in 52

individuals across all 10 sampling sites. In contrast, 10 out of 11 *M. intrudens* queens

dissected as positive controls had sperm in their spermathecae.

Fig 2. Reproductive organs of (a) *Monomorium triviale* and (b) *M. intrudens* queens. (a)

M. triviale: Translucent (empty) spermatheca (spt), ovarioles with oocytes and obvious

yellow bodies (yb) indicating that virgin queens were reproductively mature and laid eggs. **(b)**

M. intrudens: Opaque (filled with sperm) spermatheca, well-developed ovarioles, and yellow

bodies indicating that the queen is sexually reproducing. Yellow arrowheads indicate yellow

bodies.

Rearing experiment

212   Eight queens died before producing daughter workers (Table 3). Thirty-six virgin

213   queens from seven local populations survived, and each produced at least one worker. In total,

214   177 workers emerged during the experimental period; no males were produced. All dissected

215   queens had empty spermathecae (Table 4). Despite all the queens producing workers, only 13

216   of 29 queens dissected had obvious yellow bodies.

Microsatellite analysis

Analysis of 33 queens and 174 workers showed that all individuals were

heterozygous at *Mp-1* and that the genotypes of all the daughter workers were identical to

those of their mothers. In addition, all individuals within the same local population had the

same genotype. Comparison of genotypes between mothers and their daughters enables us to

infer the occurrence of sexual reproduction. A potential male mate of a heterozygous mother

(genotype AB) either shares or does not share the same allele (A or B) of *Mp-1* as the mother.

We can rule out the latter possibility on the basis of the daughter genotypes. In the former

case, the mother's sexually produced daughters should have two genotypes in the expected

ratio of 1:1. One is the same as their mother's (AB) and the other is homozygous at one of the

two mother alleles (AA when the father's genotype is A, or BB when the father's genotype is

B). Under sexual reproduction, the observed bias toward the daughter's same heterozygous

genotypes as their mother would be extremely rare. (The exact binomial probabilities are

given in Table 5.) Therefore, we can reject the possibility of sexual reproduction among the

individuals that we genotyped.

Microbial analysis
13

MiSeq sequencing and Qiime2-based analysis yielded 8364 to 46,817 reads per

biological replicate and a total of 267 OTUs (summarized in S1 Table). Among them, 263

OTUs (covering 99.9% to 100% of the reads) were classified as bacterial, and 117 OTUs

(92.4% to 99.5% of the reads, excepting 54.7% of the reads from one queen replicate from

nest B, see below) were classified at least to the phylum level. No reads were assigned to

bacterial genera that are known to induce thelytokous parthenogenesis in Hymenoptera, i.e.,

Cardinium, *Rickettsia*, and *Wolbachia* [1].

In two nests (A and C) out of the three replicates for both queens and workers, the

most abundant reads were classified as from *Spiroplasma platyhelix* (88.4% and 67.4% of

total reads from queen and worker replicates, respectively, of nest A; 91.5% and 91.1%,

respectively, for nest C). The replicates obtained from nest B showed a different pattern of *S.*

platyhelix abundance: 0% from the queen replicate and 13.2% from the worker replicate. The

absence of the *S. platyhelix* sequence from the nest B queen replicate was likely associated

with the relatively small number of total reads (8364 vs. >30,000, S1 Table).

Phylogenetic analysis

The 639-bp partial sequences of *COI* were completely identical among all

individuals representing 15 populations of *M. triviale* (Fig 3). These individuals were placed

in the monophyletic group with *M. intrudens* and *M. pharaonis*. Although the genus

Monomorium has recently been revealed to be a polyphyletic group [26, 30], our result

suggests that *M. triviale* belongs to the genus *Monomorium sensu stricto*.

Fig 3 Maximum likelihood tree of 15 populations of *M. triviale* and related species, based

on *COI* sequences. The number at each branch of the phylogenetic tree represents the

bootstrap percentage (1000 replicates). GenBank accession codes follow the taxon names.

Scale bar: 0.1 substitutions per site.

Discussion

Our comprehensive investigation allows us to add *M. triviale* to the list of

parthenogenetic ants. Production of daughters by unmated queens (indicative of type III

thelytoky) was corroborated on the basis of multiple lines of evidence: (i) absence of males

and inseminated queens in the field-collected nests; (ii) worker production by laboratory-

reared virgin queens; (iii) unfilled spermathecae of queens that produced workers; and (iv)

identical genotypes of mother queens and daughter workers. These features were common to

all the locations we tested, suggesting that thelytoky is a dominant mode of reproduction

across Japanese populations of *M. triviale*.

It should be noted here that rare occurrences of males have been reported in other

parthenogenetic ants (type II), such as *Ooceraea biroi* [19] and *Pristomyrmex punctatus* [31].

In addition, queens of *M. triviale* retain undegenerated spermathecae [18], suggesting that

they have low level of specialization to male-free reproduction. Moreover, geographic

variation in sexual and asexual reproduction has been reported in some thelytokous ant

species (types II and III), such as *Mycocepurus smithii*, *Myrmecina nipponica*, and

Platythyrea punctata [17, 21, 32]. Whether and how often sexual reproduction occurs in *M.*

triviale, especially in populations outside Japan, would be an interesting topic for future

study.

As major cytological mechanisms of thelytoky, terminal fusion automixis and central

fusion automixis have usually been considered [33]. Under terminal fusion automixis, cells

derived from the same cell at the first meiosis are fused to a diploid after the second meiosis.

If there is no recombination between chromosomes, the offspring have a homozygous

genotype. Under central fusion automixis, different cells at the first meiosis are fused. If

recombination does not occur, the offspring have the same genotype as the mother. Many

thelytokous species of social Hymenopteran insects adopt the central fusion type of automixis

[1]. The identical heterozygous genotypes that we observed here between the mother queen

and her daughter workers in *M. triviale* are consistent with central fusion automixis.

Our exploratory analysis of bacterial communities in *M. triviale* provides basic

information for future studies of the host–symbiont relationship. No evidence was found for

infection with thelytoky-inducing bacteria in this species, confirming previous reports of no

cases of microorganism-induced thelytoky in ants [10, 24, 34, 35]. *Spiroplasma* has been

detected in various ant species [36], including another thelytokous species, *Mycocepurus*

smithii (type III thelytoky) [37]. In *Solenopsis*, a genus closely related to *Monomorium*,

Spiroplasma has been detected as a dominant bacterial taxon in whole bodies of adult workers

of *S. geminata* [38], thus drawing parallels with our finding. It is noteworthy that type I

thelytoky has recently been found in polygynous colonies of this species [39]. *Spiroplasma* is

well known as a sex ratio distorter in *Drosophila* [40], and its role in the host's reproductive

biology deserves further study in ants.

In our phylogenetic analysis, the extremely low levels of diversity observed in the *COI* sequences suggest the possibility of rapid spread or selective sweep in the recent past [41]. Our results support the concept that this species is a member of the genus *Monomorium sensu stricto*. This phylogenetic status is advantageous in that the study designs established in *M. pharaonis*, a well-studied model system (e.g., [42–47]), will be applicable to future comparative analyses.

Thelytokous parthenogenesis is often considered as a trait overrepresented among introduced or invasive ant species [48]. Although some studies have categorized *M. triviale* as invasive [1, 19], no exotic distribution has been reported so far [14] and there is insufficient information to support the possibility of an introduced origin of the Japanese population of *M. triviale*. The known distribution range of this species is far more limited than those of successful invasive congeners such as *M. pharaonis* and *Monomorium floricola* [15, 49]. A preference for disturbed and urban habitat is shared widely among invasive ants [50]. Ito et al. [50] listed *M. triviale* along with *Strumigenys membranifera*, *P. punctatus*, and *O. biroi* as thelytokous ants found in "open, disturbed areas.'' Among these species, the life history of *M. triviale* is most poorly known. In our study, *M. triviale* nests were found even in a thicket dominated by deciduous broad-leaved trees, which is not typical of "open, disturbed areas." Nevertheless, the above information does not rule out the possibility that *M. triviale* will become invasive in the future. Future studies of this species' social structure, such as queen numbers and the mode of colony foundation, will help to evaluate the potential invasion risk of this species as an ecological consequence of its life history.

Acknowledgments

We are grateful to Kenji Matsuura who allowed us to use his laboratory. Fuminori Ito

provided valuable advice. We thank Riou Mizuno, Yu Hisasue, Kazuya Takeda, and Kunio

Sadahiro for field assistance. We also thank Kyosuke Ohkawara, Hiroyuki Shimoji, and

Tomonari Nozaki for providing us with *M. triviale* nests.

References

1.   Rabeling C, Kronauer DJC. Thelytokous Parthenogenesis in eusocial hymenoptera.

     Annu Rev Entomol. 2013;58: 273–292. doi:10.1146/annurev-ento-120811-153710

2.   Engelstädter J. Constraints on the evolution of asexual reproduction. BioEssays.

     2008;30: 1138–1150. doi:10.1002/bies.20833

3.   Neiman M, Schwander T. Using parthenogenetic lineages to identify advantages of

     sex. Evol Biol. 2011;38: 115–123. doi:10.1007/s11692-011-9113-z

4.   Goudie F, Oldroyd BP. The distribution of thelytoky, arrhenotoky and androgenesis

     among castes in the eusocial Hymenoptera. Insectes Soc. 2018;65: 5–16.

     doi:10.1007/s00040-017-0597-0

5.   Hartmann A, Wantia J, Torres JA, Heinze J. Worker policing without genetic

     conflicts in a clonal ant. Proc Natl Acad Sci. 2003;100: 12836–12840.

     doi:10.1073/pnas.2132993100

6.   Ravary F, Lecoutey E, Kaminski G, Châline N, Jaisson P. Individual Experience

     alone can generate lasting division of labor in ants. Curr Biol. 2007;17: 1308–1312.

     doi:10.1016/j.cub.2007.06.047

7.   Dobata S, Tsuji K. Public goods dilemma in asexual ant societies. Proc Natl Acad

     Sci. 2013;110: 16056–16060. doi:10.1073/pnas.1309010110

346   8.   Bernadou A, Busch J, Heinze J. Diversity in identity: behavioral flexibility,

347   dominance, and age polyethism in a clonal ant. Behav Ecol Sociobiol. 2015;69:

348   1365–1375. doi:10.1007/s00265-015-1950-9

349   9.   Oxley PR, Ji L, Fetter-Pruneda I, McKenzie SK, Li C, Hu H, et al. The genome of

350   the Clonal raider ant *Cerapachys Biroi*. Curr Biol. 2014;24: 451–458.

351   doi:10.1016/j.cub.2014.01.018

352   10.   Himler AG, Caldera EJ, Baer BC, Fernandez-Marin H, Mueller UG. No sex in

353   fungus-farming ants or their crops. Proc R Soc B Biol Sci. 2009;276: 2611–2616.

354   doi:10.1098/rspb.2009.0313

355   11.   Rabeling C, Lino-Neto J, Cappellari SC, Dos-Santos IA, Mueller UG, Bacci M.

356   Thelytokous parthenogenesis in the fungus-gardening ant *Mycocepurus smithii*

357   (Hymenoptera: Formicidae). PLoS One. 2009;4. doi: 10.1371/journal.pone.0006781

358   12.   Wheeler, William Morton HS. The ants of Japan. Bulletin of the AMNH. Bull

359   AMNH. 1906; v. 22, art.

360   13.   Bolton B. An online catalog of the ants of the world. 2020 (accessed 17 December,

361   2020). Available from: http://antcat.org.

362   14.   Guénard B, Weiser MD, Gómez K, Narula N, Economo EP. The Global Ant

363   Biodiversity Informatics (GABI) database: Synthesizing data on the geographic

364   distribution of ant species (Hymenoptera: Formicidae). Myrmecological News.

365   2017;24: 83–89.

366   15.   Wetterer JK. Worldwide spread of the pharaoh ant, *Monomorium pharaonis*

367   (Hymenoptera: Formicidae). Myrmecological News. 2010;13: 115–129.

16.   Pontieri L, Linksvayer TA. *Monomorium*. Encyclopedia of Social Insects. Cham:

Springer International Publishing; 2019. pp. 1–6. doi:10.1007/978-3-319-90306-

4_171-1

17.   Rabeling C, Gonzales O, Schultz TR, Bacci M, Garciad MVB, Verhaaghe M, et al.

Cryptic sexual populations account for genetic diversity and ecological success in a

widely distributed, asexual fungus-growing ant. Proc Natl Acad Sci U S A.

2011;108: 12366–12371. doi:10.1073/pnas.1105467108

18.   Gotoh A, Billen J, Tsuji K, Sasaki T, Ito F. Histological study of the spermatheca in

three thelytokous parthenogenetic ant species, *Pristomyrmex punctatus*, *Pyramica*

membranifera and *Monomorium triviale* (Hymenoptera: Formicidae). Acta Zool.

2012;93: 200–207. doi:10.1111/j.1463-6395.2010.00498.x

19.   Kronauer DJC, Pierce NE, Keller L. Asexual reproduction in introduced and native

populations of the ant *Cerapachys biroi*. Mol Ecol. 2012;21: 5221–5235.

doi:10.1111/mec.12041

20.   Masuko K. Thelytokous Parthenogenesis in the Ant *Strumigenys hexamera*

(Hymenoptera: Formicidae). Ann Entomol Soc Am. 2013;106: 479–484.

doi:10.1603/an12144

21.   Masuko K. Thelytokous Parthenogenesis in the Ant *Myrmecina nipponica*

(Hymenoptera: Formicidae). Zoolog Sci. 2014;31: 582–586. doi:10.2108/zs140050

22.   van der Kooi CJ, Schwander T. On the fate of sexual traits under asexuality. Biol

Rev. 2014;89: 805–819. doi:10.1111/brv.12078

23.   Lee CC, Hsu SF, Yang CCS, Lin CC. Thelytokous parthenogenesis in the exotic

dacetine ant *Strumigenys rogeri* (Hymenoptera: Formicidae) in Taiwan. Entomol Sci.

2018;21: 28–33. doi:10.1111/ens.12277

24.   The invasion biology and sociogenetics of pharaoh ants. PhD Thesis, University of

Copenhagen, København, Denmark.

25.   Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al.

Reproducible, interactive, scalable and extensible microbiome data science using

QIIME 2. Nat Biotechnol. 2019;37: 852–857. doi:10.1038/s41587-019-0209-9

26.   Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R. DNA primers for amplification

of mitochondrial cytochrome c oxidase subunit I from diverse metazoan

invertebrates. Mol Mar Biol Biotechnol. 1994;3: 294–9. Available:

http://www.ncbi.nlm.nih.gov/pubmed/7881515

27.   Ward PS, Brady SG, Fisher BL, Schultz TR. The evolution of myrmicine ants:

Phylogeny and biogeography of a hyperdiverse ant clade (Hymenoptera:

Formicidae). Syst Entomol. 2015;40: 61–81. doi:10.1111/syen.12090

28.   Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: Improving the sensitivity of

progressive multiple sequence alignment through sequence weighting, position-

specific gap penalties and weight matrix choice. Nucleic Acids Res. 1994;22: 4673–

4680. doi:10.1093/nar/22.22.4673

29.   Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular evolutionary

genetics analysis across computing platforms. Mol Biol Evol. 2018;35: 1547–1549.

doi:10.1093/molbev/msy096

30.   Sparks KS, Andersen AN, Austin AD. A multi-gene phylogeny of Australian

Monomorium Mayr (Hymenoptera: Formicidae) results in reinterpretation of the

genus and resurrection of *Chelaner* Emery. Invertebr Syst. 2019;33: 225–236.

doi:10.1071/IS16080

31.   Yamada A, Eguchi K. Description of the male genitalia of *Pristomyrmex punctatus*

(Smith, 1860) (Hymenoptera, Formicidae, Myrmicinae). Asian Myrmecology.

2016;8: 87–94. doi:10.20362/am.008010

32.   Kellner K, Seal JN, Heinze J. Sex at the margins: Parthenogenesis vs. facultative and

obligate sex in a Neotropical ant. J Evol Biol. 2013;26: 108–117.

doi:10.1111/jeb.12025

33.   Pearcy M, Hardy O, Aron S. Thelytokous parthenogenesis and its consequences on

inbreeding in an ant. Heredity (Edinb). 2006;96: 377–382.

doi:10.1038/sj.hdy.6800813

34.   Grasso DA, Wenseleers T, Mori A, Le Moli F, Billen J. Thelytokous worker

reproduction and lack of *Wolbachia* infection in the harvesting ant *Messor capitatus*.

Ethol Ecol Evol. 2000;12: 309–314. doi:10.1080/08927014.2000.9522803

35.   Wenseleers T, Billen J. No evidence for *Wolbachia*-induced parthenogenesis in the

social Hymenoptera. J Evol Biol. 2000;13: 277–280. doi:10.1046/j.1420-

9101.2000.00168.x

36.   Kautz S, Rubin BER, Moreau CS. Bacterial infections across the ants: Frequency and

prevalence of *Wolbachia*, *Spiroplasma*, and *Asaia*. Psyche (London). 2013.

doi:10.1155/2013/936341

37.   Sen R, Ishak HD, Estrada D, Dowd SE, Hong E, Mueller UG. Generalized antifungal

activity and 454-screening of *Pseudonocardia* and *Amycolatopsis* bacteria in nests of

fungus-growing ants. Proc Natl Acad Sci U S A. 2009;106: 17805–17810.

doi:10.1073/pnas.0904827106

38.   Ishak HD, Plowes R, Sen R, Kellner K, Meyer E, Estrada DA, et al. Bacterial

Diversity in *Solenopsis invicta* and *Solenopsis geminata* Ant colonies characterized

by 16S amplicon 454 pyrosequencing. Microb Ecol. 2011;61: 821–831.

doi:10.1007/s00248-010-9793-4

39.   Lacy KD, Shoemaker DW, Ross KG. Joint Evolution of Asexuality and Queen

Number in an Ant. Curr Biol. 2019;29: 1394-1400.e4. doi:10.1016/j.cub.2019.03.018

40.   Paredes JC, Herren JK, Schüpfer F, Marin R, Claverol S, Kuo CH, et al. Genome

sequence of the *Drosophila melanogaster* male-killing *Spiroplasma* strain MSRO

endosymbiont. MBio. 2015;6. doi:10.1128/mBio.02437-14

41.   Avise JC, others. Phylogeography: the history and formation of species. Harvard

university press; 2000.

42.   Beekman M, Sumpter DJT, Ratnieks FLW. Phase transition between disordered and

ordered foraging in pharaoh's ants. Proc Natl Acad Sci U S A. 2001;98: 9703–9706.

doi:10.1073/pnas.161285298

43.   Jefford CW, Tang Q, Zaslona A, Tang Q, Zaslona A. Short, Enantiogenic Syntheses

of (-)-Indolizidine 167B and (+)-Monomorine. J Am Chem Soc. 1991;113: 3513–

3518. doi:10.1021/ja00009a043

44.   Jackson DE, Holcombe M, Ratnieks FLW. Trail geometry gives polarity to ant

foraging networks. Nature. 2004;432: 907–909. doi:10.1038/nature03105

45.   Robinson EJH, Jackson DE, Holcombe M, Ratnieks FLW. Insect communication:

"No entry" signal in ant foraging. Nature. 2005;438: 442. doi:10.1038/438442a

46.   Warner MR, Mikheyev AS, Linksvayer TA. Genomic signature of kin selection in an

ant with obligately sterile workers. Mol Biol Evol. 2017;34: 1780–1787.

doi:10.1093/molbev/msx123

47.   Oliveira RC, Warson J, Sillam-Dussès D, Herrera-Malaver B, Verstrepen K, Millar

JG, et al. Identification of a queen pheromone mediating the rearing of adult sexuals

in the pharaoh ant *Monomorium pharaonis*. Biol Lett. 2020;16: 20200348.

doi:10.1098/rsbl.2020.0348

48.   Trible W, McKenzie SK, Kronauer DJC. Globally invasive populations of the clonal

raider ant are derived from Bangladesh. Biol Lett. 2020;16: 20200105.

doi:10.1098/rsbl.2020.0105

49.   Wetterer JK. Worldwide spread of the flower ant, *Monomorium floricola*

(hymenoptera: Formicidae). Myrmecological News. 2009;13: 19–27.

50.   Rabitsch W. The hitchhiker's guide to alien ant invasions. BioControl. 2011;56: 551–

572. doi:10.1007/s10526-011-9370-x

51.   Ito F, Touyama Y, Gotoh A, Kitahiro S, Billen J. Thelytokous parthenogenesis by

queens in the dacetine ant *Pyramica membranifera* (Hymenoptera: Formicidae).

Naturwissenschaften. 2010;97: 725–728. doi:10.1007/s00114-010-0688-5

Supporting information

477 **S1 Table. Microbial taxa and their relative abundances (% of total reads) per each *M.***

478 ***triviale* sample**

Funding

481 This work was supported by a Japan Society for the Promotion of Science (JSPS) Research

482 Fellowship for Young Scientists to NI (19J22242) and a grant from the Secom Science and

483 Technology Foundation to SD.

Figure_1

Figure_2

1. *Monomorium triviale* Kyoto LC592056
2. *M. triviale* Tsuchiura LC592064
3. *M. triviale* Chofu LC592062
4. *M. triviale* Tsukuba LC592050
5. *M. triviale* Ohtsu LC592057
6. *M. triviale* Matsudo LC592063
7. *M. triviale* Higashiosaka LC592055
8. *M. triviale* Okazaki LC592060
9. *M. triviale* Sanda LC592054
12. *M. triviale* Takamatsu LC592053
13. *M. triviale* Matsuyama LC592051
14. *M. triviale* Takashima LC592058
15. *M. triviale* Kannonji LC592052
16. *M. triviale* Kagamihara LC592061
17. *M. triviale* Matsue LC592059
M. intrudens LC592065
M. pharaonis GU710434
Erromyrma latinodis GU709833
Syllophopsis sechellensis EF609858
Solenopsis invicta HQ928672

100
62
94
59

0.1

Figure_3