Comparison of mouse models reveals the molecular bases for psychotic illness in Prader-Willi syndrome

Simona Zahova[1], Trevor Humby[2], Jennifer R. Davies[1], Joanne E. Morgan[1], Anthony R. Isles[1]*

[1]MRC Centre for Neuropsychiatric Genetics and Genomics, School of Medicine, Cardiff University, Hadyn Ellis Building, Maindy Road, Cardiff, CF24 4HQ, U.K.; ZahovaS@cardiff.ac.uk

[2]School of Psychology, Cardiff University, Tower Building, 70 Park Place, Cardiff, CF10 3AT, U.K. HumbyT@cardiff.ac.uk

* *Correspondence:* Anthony R. Isles; IslesAR1@cardiff.ac.uk; MRC Centre for Neuropsychiatric Genetics and Genomics, School of Medicine, Cardiff University, Hadyn Ellis Building, Maindy Road, Cardiff, CF24 4HQ, U.K.

Abstract

Prader-Willi Syndrome (PWS) is a neurodevelopmental disorder caused by mutations affecting paternal chromosome 15q11-q13, characterized by hypotonia, hyperphagia, impaired cognition and behavioural problems. Psychotic illness is a challenging problem for individuals with PWS, and has different rates of prevalence in distinct PWS genotypes. Previously, we demonstrated behavioural and cognitive endophenotypes of relevance to psychiatric illness in a mouse model for one of the associated PWS genotypes, namely PWS-IC, in which deletion of the imprinting centre leads to loss of paternal imprinted gene expression and over-expression of *Ube3a*. Here we examine the broader gene expression changes that are specific to the psychiatric endophenotypes seen in this model. To do this we compared the brain transcriptomic profile of the PWS-IC mouse to the PWS-cr model that carries a deletion of the PWS minimal critical interval spanning the snoRNA *Snord116* and *Ipw*. Firstly, we examined the same behavioural and cognitive endophenotypes of relevance to psychiatric illness in the PWS-cr mice. PWS-cr exhibit no differences in locomotor activity, sensory-motor gating and attention of relevance to psychotic illness, and previously seen in the PWS-IC mice. RNA-seq analysis of neonatal brain revealed a greater number of transcriptional changes between PWS-IC and wild-type littermates, than between PWS-cr and wild-type littermates. Moreover, the differentially expressed genes in PWS-IC brain were enriched for GWAS variants associated with episodes of psychotic illness but, interestingly, not schizophrenia. These data illustrate the molecular pathways that may underpin psychotic illness in PWS and have implication for potential therapeutic interventions.

Introduction

Prader-Willi syndrome (PWS) is a neurodevelopmental disorder caused by disrupted expression of the paternally imprinted genes on chromosome 15q11-q13. The disruption of this locus is associated with hypotonia, hyperphagia, mild to moderate learning disability, as well as a range of psychiatric conditions (1, 2). Upwards of 20% of PWS patients are formally diagnosed with major depressive disorder (MDD), while 50-70% score highly on questionaires for anxious and depressive episodes (3). Phenotypes of reduced attention and poor impulse control are also highly prevalent and are reported to be intensified in children with PWS compared to their peers with learning disability only (4, 5). Furthermore, between 16% and 32% of patients experience episodes of psychosis, including delusional ideation and hallucinations (3, 5-7).

The prevalence and variation of psychiatric endophenotypes in PWS are influenced by genotype, which can vary among individuals. The 15q11-q13 imprinted locus includes several genes expressed only from the paternally inherited copy of chromosome 15 (*MKRN3, MAGEL2, NECDIN, NPAP1, SNURF-SNRPN*, cluster of non-coding RNAs), as well as two genes expressed only from the maternally inherited copy of chromosome 15 (*UBE3A, ATP10*) (8). Roughly, PWS genotypes can be divided into two main subtypes. The first are those that lead to loss of paternal gene expression from 15q11-q13 and include most common PWS genotype (75%) in which individuals carrying a paternally-derived deletion spanning the whole interval (delPWS). The delPWS genotype has been further refined in recent years as several case studies have reported paternal microdeletions of a few of the non-coding genes, which induce hyperphagia and hypotonia (9-11). Since these phenotypes are considered the core features of PWS, the deleted non-coding genes define a PWS critical interval.

The second main subtype is maternal uniparental disomy of chromosome 15 (mUPD15), found in about 25% of PWS patients. Population studies of individuals with PWS report increased incidence and severity of psychosis in those with mUPD15 compared to delPWS, including a higher rate of diagnoses of schizophrenia and bipolar disorder (12, 13). In addition to complete loss of expression of all paternally expressed genes (PEGs), mUPD15 leads to an overexpression of the two maternally expressed genes (MEGs) in the 15q11-q13 imprinted interval (8). A similar molecular profile is also imprinting centre (IC) deletion genotype, which, although less well-studied due to its rarity (<5% of all PWS), also shows increased incidence of psychotic illness (14). These findings have led to the suggestion that it is the overexpression of MEGs in these genotypes that might be the main contributors to psychiatric illness in PWS (8), an idea supported by the increased incidence of psychotic illness in maternal 15q duplication syndrome (15). Nevertheless, delPWS patients can also exhibit depression, anxiety and, albeit with reduced incidence and severity, psychotic episodes (16). Consequently, it is possible that the PEGs of the PWS locus also contribute. However, as yet the mechanism leading to psychiatric illness remain unclear.

To address this, we have used mouse models to investigate behavioural and cognitive phenotypes of relevance to psychiatric illness seen in PWS. Previously, we have demonstrated that a PWS imprinting centre deletion (PWS-IC) mouse, which models the rare imprinting centre genotype and demonstrates loss of expression of PEGs and overexpression of MEGs matching that observed in individuals with mUPD15, recapitulates many of the behavioural and cognitive deficits of relevance to PWS, including hypoactivity, abnormal sensory-motor gating, reduced attention and greater impulsivity (17, 18). Here, we attempt to examine the molecular changes that may underpin these psychiatric endophenotypes specifically by comparing the brain gene expression changes in PWS-IC

mice with those in another model, carrying a deletion of the critical interval (PWS-cr) that eliminates expression of the non-coding PEGs *Snord116* and *Ipw*. Firstly, we established that the behavioural and cognitive endophenotypes of relevance to psychiatric illness are limited to the PWS-IC mice by conducting the same behavioural characterisation of PWS-cr mice. We then used RNA-seq to investigate the molecular bases of the differences in behaviour between the two mouse models for PWS. We show that the differentially expressed genes (DEGs) and isoform changes seen in PWS-IC brain, but not PWS-cr, are enriched for common genetic variants associated with psychotic episode occurrences. Interestingly, the DEGs and isoform changes in PWS-IC were not enriched for genetic variants associated with schizophrenia, providing molecular support for the observed aetiology of psychotic illness in PWS. Our findings shed light on a molecular basis of behavioural and cognitive problems seen in PWS, and may have implications for the treatment of psychotic illness in individuals with genetic lesions at 15q11-q13.

Results

Elevated plus maze and open field test

The elevated plus maze (EPM) and open field (OF) were used to investigate anxiety-related phenotypes in 5 min and 10 min sessions respectively. As expected, on the EPM all mice spent significantly more time of the 5 min session in the closed arms of the maze compared to the more anxiogenic open arms of the maze ($\chi^2_{(1)}$=138.98, p<0.001), but no effect of genotype on percentage of time spent in the open arms of the EPM ($F_{(3,50)}$=0.132; p=0.361) (Figure 1A), or on distance travelled through the area of the maze ($F_{(3,\ 50)}$=2.352, p=0.174). Number of head dips and stretch attends were also recorded as a measure of explorative behaviour through the EPM, and showed no significant differences between PWS-cr mice and their wild type littermates ($F_{(3,\ 50)}$=0.7832, p=0.989; $F_{(3,\ 50)}$=1.933, p=0.151).

During the OF test, mice spent significantly more time of the 10 min session in the outer areas of the arena than in the more anxiogenic centre square $\chi^2_{(1)}$=452.2, p<0.001), but there was no effect of genotype on percentage of time spent in the centre square ($F_{(3,\ 50)}$=0.084; p=0.361) (Figure 1B) or on distance travelled through the arena ($F_{(3,\ 50)}$=4.55; p=0.472). Overall, the data collected from the EPM and OF show no differences in behaviour between PWS-cr mice and their wild type littermates. Our results suggest no obvious indications of anxiety phenotype in the PWS-cr mouse model.

Locomotor activity

Locomotor activity (LMA) was measured in custom made chambers fitted with infrared beams tracking the movement of the mice. The test was run in a two-hour session for two consecutive days. There was no effect of genotype on total number of beam breaks made over the span of the two sessions $\chi^2_{(2)}$=4.625, p=0.099), or on total number of runs made from one end of the chamber to the other $\chi^2_{(2)}$=1.237, p=0.539). Further analysis of the data in 30 min time bins showed that all animals exhibited habituation to the environment within and between sessions, as indicated by the steady decrease of number of runs (Figure 1C; $\chi^2_{(3)}$=347.75, p<0.001; $\chi^2_{(1)}$=49.953, p<0.001) and beam breaks (Figure 1D; $\chi^2_{(3)}$=197.83, p<0.001; $\chi^2_{(1)}$=33.562, p<0.001) over time. However, there was no statistically significant difference in habituation behaviour between the two genotypes $\chi^2_{(2)}$=3.342, p=0.342; $\chi^2_{(2)}$=1.761, p=0.222). Overall, the results from this test showed no locomotor activity phenotype in the PWS-cr mice.

Figure 1. Behaviours recorded at the elevated plus maze, open field arena and locomotor activity test. The elevated plus maze task (a) showed no effect of genotype on percentage of time spent in the open arms of the maze (PWS-cr=23, WT=31). Similarly, data from the open field test (b) showed no difference in percentage of time spent in the centre square of the arena between PWS-cr mice (n=23) and their wild type littermates (n=31). Locomotor activity was tested in custom made chambers with fitted infrared beams, which tracked the movement of the animals in two-hour sessions on two consecutive days. Number of beam breaks (d) and number of runs from one end of the chamber to the other (c) were considered the main measures of activity. The test showed decreased locomotor activity between and within sessions (p<0.001), which is indicative of habituation to the environment. There was no effect of genotype on the behaviour examined by this test (PWS-cr=23, WT=31).

Acoustic startle response and pre-pulse inhibition

Acoustic startle response (ASR) and pre-pulse inhibition (PPI) to acoustic startle response were measured at pulse intensities of 120 dB and 105 dB. A total of thirteen ASR-alone pulses were presented to the mice at the two intensities, and both showed a significantly reduced startle response in the PWS-cr mice (Figure 2A; $\chi2_{(2)}$=6.5292, p=0.03821; Figure 2B; $\chi2_{(2)}$=14.562, p<0.001). Pre-pulses of 8 dB and 16 dB played 70 ms prior to the 120 dB and 105 dB pulses inhibited the startle response up to mean average of 74.09% (±1.84% SEM) and 56.1% (±4.86% SEM) respectively, indicating operational sensory-motor gating in both groups of mice (Figure 2C, 2D). Genotype had no effect on pre-pulse inhibition at either pulse intensity ($\chi2_{(2)}$=5.376, p=0.251; $\chi2_{(2)}$=2.556, p=0.635). Overall, the results from the ASR and PPI tests show no sensory-motor gating phenotypes in the PWS-cr mouse model, but a reduced startle response to acoustic stimuli.

Following the 120dB and 105dB ASR and PPI schedules, mice were exposed to a gradual increasing ASR-alone pulse intensity, ranging from 70dB to 120dB. As expected, the increasing intensity

resulted in increasing response ($\chi^2_{(12)}$=253.28, p<0.001). These data suggest that the difference in ASR between PWS-cr and wild type mice emerges at >100dB intensity (Supplementary Figure 1). However, there was no effect of genotype ($\chi^2_{(12)}$=20.49, p=0.058) and no interaction between genotype and pulse intensity ($\chi^2_{(10)}$=9.27, p=0.507).

Figure 2. Acoustic startle and pre-pulse inhibition. The startle response to a series of 13 acoustic stimuli at 120 dB (a) and 105 dB (b) was significantly reduced in PWS-cr mice (n=23) compared to their wild type littermates (n=31) (* indicates p<0.05, *** indicates p<0.001). Pre-pulse inhibition of the acoustic startle response was measured by presenting noises of 8 dB and 16 dB 70 ms before the pulses of 120 dB (c) and 105 (d). The pre-pulse inhibition data showed operational sensory motor gating, which was not affected by genotype (PWS-cr=23, WT=31).

5-choice serial reaction time task

Behaviours relating to visuo-spatial attention and control response were investigated in the 5-CSRTT, where mice were trained to respond correctly to stimuli in order to receive condensed milk dilution as a reward. Prior examination showed an increased preference towards condensed milk dilution over water in all animals ($\chi^2_{(3)}$=68.461, p<0.001), thus validating its use as a motivational reward. There was no significant distinction between the preference to the reward between PWS-cr and wild type (Figure 3A; $\chi^2_{(2)}$=0.5827, p=0.7473) and thus no indications of increased interest in the PWS-cr mice towards the reward, which was an important consideration in the interpretation of the results from the operant tasks.

The average number of sessions required to reach baseline performance in the 5-CSRTT did not differ between PWS-cr (mean 98.6, ±7.3 SEM) and wild type (mean 90.8, ±5.7 SEM) mice ($F_{3,40}$=0.703, p=0.802). Performance at baseline conditions of the 5-CSRTT showed no effect of genotype on task accuracy (Figure 3B; $F_{(3, 42)}$=0.127, p=0.724), omissions (Figure 3C, $F_{(3, 42)}$=0.104;

p=0.749), premature responses (Figure 3D; $F_{(3, 40)}$=0.515, p=0.361) and all other parameters recorded at these conditions of the task.

Figure 3. Condensed milk test and 5-CSRTT performance at baseline conditions. Preference to the condense milk reward over water was tested over the span of days 3 to 6 from the condense milk test (a). Over the course of these four days the preference to the reward increased significantly (p<0.001) as expected for the purposes of the 5-CSRTT, and this tendency was not affected by genotype (PWS-cr=21, WT=22). Performance at baseline conditions of the 5-CSRTT also showed no difference between PWS-cr mice (n=20) and their wild type littermates (n=23) in accuracy of performance (b), percentage omissions (b) and number of premature responses (b).

The conditions of the 5-CSRTT were manipulated in order to tease out subtler aspects of attention and impulsivity. Behaviour was inspected at stimulus duration of shortened presentation, all introduced in pseudorandom order within one 5-CSRTT session. The shortened stimulus duration task lowered the overall task accuracy ($\chi^2_{(3)}$=41.114, p<0.001), and this performance decline was not influenced by genotype (Figure 4A; $\chi^2_{(1)}$=1.642, p=0.2). There was no effect of genotype on percentage omissions (Figure 4B; $\chi^2_{(1)}$= 0.357, p=0.551) or any of the other parameters that were investigated in the task.

An increase of the inter-trial intervals was introduced in a different session in order to induce premature responses in the task and tease out a potential impulsivity phenotype. The inter-trial interval increase lead to a rise in number of premature responses as predicted ($\chi^2_{(3)}$=32.131, p<0.001), but this tendency was not affected by genotype (Figure 4C; $\chi^2_{(2)}$= 4.966, p=0.084). The percentage of omissions at this task was also not significantly different between PWS-cr mice and their wild type littermates (Figure 4D; $\chi^2_{(2)}$= 0.015, p=0.992), and neither were any of the other examined parameters. Overall, the data collected from the 5-CSRTT showed no indications of an attention or impulsivity phenotype in the PWS-cr mouse model.

Figure 4. Shortened stimulus duration and increased inter-trial intervals at the 5-CSRTT.
Shortening the stimulus duration at the 5-CSRTT reduced performance accuracy (p<0.001) (a) and increased percentage of trial omissions (p<0.001) (b) but revealed no difference between the behaviour of PWS-cr mice (n=15) and their wild type littermates (n=21). Increasing the inter-trial intervals lead to an increase in premature responses (p<0.001), which was not affected by genotype (PWS-cr=18, WT=24) (c). Omissions were also not affected by genotype in this task manipulation (d).

RNA-sequencing of whole brain neonatal tissue

The results from a behavioural study demonstrated a vastly different behavioural and cognitive profile in the PWS-cr mice compared to the previously examined PWS-IC mouse. In order to investigate the gene expression changes underlying these behavioural difference, we conducted an RNA-sequencing assay on whole brain tissue from both mouse models. While all behavioural tasks were conducted on adult individuals, for the RNA-sequencing study we took tissue from neonatal mice, as genetic risk for psychiatric illness has been shown to manifest from early developmental stages (19, 20). Furthermore, qPCR assay of genes from within the PWS critical interval in wild-type whole brain tissue at three developmental stages (E13.5, E18.5, P0), showed that the expression peaks at birth (Supplementary Figure 2).

PWS-cr differentially expressed genes and isoforms

Visualization of the aligned RNA-sequencing reads in the UCSC browser confirmed that the deletion in the PWS-cr mouse model mode spans all copies of *Snord116* snoRNA and five of the six *Ipw* exons (Supplementary Figure 3). However, differential gene expression analysis showed only very subtle differences between the PWS-cr mice and their wild type littermates after Benjamini-Hochberg adjustment for multiple testing. In total, there were seven differentially expressed genes (DEGs) in brain tissue from the PWS-cr mouse model (Figure 5a and Appendix 3). Among them was the *Snhg14* non-coding RNA, which acts as a host for the critical interval (z=21.846, padj<0.001). The *Mafa* gene

encoding a transcription factor that regulates pancreatic beta cell-specific expression of the insulin gene was upregulated in the absence of the critical interval (z=-7.014, padj<0.001). Notably, the *Necdin* growth suppressor which is one of the imprinted genes of the PWS locus, was upregulated in the PWS-cr mouse brain tissue (z=-8.621, padj<0.001). The remaining four DEGs were predicted genes of unknown function, two of which (*Gm44559* and *Gm44562*) fall within intron 4/12 of the *Ube3A*, and *Gm44831* which falls into *Ipw* in the critical interval.

In addition to differential gene expression, the read depth used in our RNA-seq analysis allowed analysis of isoform switches with functional consequences in order to investigate whether the genes in the critical interval play a regulative role in alternate splicing. Seven genes showed significantly different patterns of splicing in the PWS-cr mouse brain (Appendix 4), and there was no overlap between them and the DEGs. Among these was dual-specificity tyrosine-phosphorylation-regulated kinase (*Dyrk3*), which plays a role in the dissolution of stress granules and in the early secretory pathway (21). The DEGs and isoforms were pooled together for a gene ontology analysis, which revealed no enrichments.

PWS-IC differentially expressed genes and isoforms

In contrast to PWS-cr, analysis of PWS-IC mouse brain tissue RNA-sequencing data primarily clustered by genotype and revealed 59 significantly DEGs compared to their wild-type littermates (Figure 5a; Appendix 5). This included *Snhg14*, the non-coding RNA that host the *Snord116* and *Ipw*, but also, as expected, the other PWS genes such as *Snrpn, Necdin, Mkrn3,* and *Magel2*. Outside the PWS cluster there were other notable DEGs, including the circadian clock regulator *Per1* (z=-4.25, padj=0.012), Plp1 (z=6.11, padj<0.001) and insulin growth factor *Igf1* (z=4.164, padj=0.015).

Analysis of isoform switches identified 48 differentially spliced genes, and there was no overlap with the differentially expressed genes (Appendix 4). Among them was the GABA receptor subunit gene *Gabrg3*, which is associated with autism, and Angelman syndrome, a sister disorder to PWS caused by mutations in the same cluster of imprinted genes (22, 23). Differentially expressed genes and isoforms were pooled together for a gene ontology term analysis, which were enriched for molecular functions, cellular compartments and biological compartments relevant to oxygen transport (Appendix 6).

Comparing the combined DEG and isoform data for the two models directly shows that there were a greater number of differences in the PWS-IC mice overall (Figure 5b). As expected, a good proportion (~50%) of the changes seen in PWS-cr were shared with those found in PWS-IC (Figure 5b). However, this overlap was limited to DEGs, as there were no common differentially spliced genes in the PWS-cr and PWS-IC samples.

Enrichment of common genetic variants

We pooled the differentially expressed and differentially spliced gene sets (padj<0.05) together for each mouse model in order to look for enrichment of common genetic variants associated with schizophrenia or psychotic episodes. The analysis was performed using MAGMA's regression gene-set computation with the datasets from genome wide association studies (GWAS) that utilize the UK Biobank, ClozUK and CKDgen consortium databases respectively (24-26). The results showed no enrichment of genetic variants of interest in the PWS-cr samples (Figure 5c). In contrast, results from the PWS-IC model indicated an enrichment of SNPs common for the experience of 'any' and 'multiple' psychotic episodes (Appendix 7; β=0.215, p=0.023, β=0.271, p=0.007), but interestingly not for schizophrenia (Figure 5c; β=-0.206, p=0.913). As expected, neither DEG and isoform list was enriched for the negative control dataset, common genetic variants associated with chronic kidney disease (β=-0.161, p=0.909).

Figure 5. RNA-sequencing assay. Differential gene expression analysis (a) showed an overexpression of *Necdin* and *Mafa* in the PWS-cr tissue, as well as an expected loss of expression of the long non-coding RNA *Snhg14* that acts as a host for the critical interval. Among the differentially expressed genes in the PWS-IC tissue were the genes from the PWS locus, and notably the circadian clock regulator *Per1* and the insulin growth factor *Igf1*. Predicted genes of unknown function were excluded from the graph. The y axis on this graph (a) shows log2 fold change of expression, error bars indicate standard error of the mean. Differentially expressed genes and isoforms were pooled for enrichment analyses; PWS-cr and PWS-IC pooled genes had 6 genes in common (b). Gene set analysis of pooled differentially expressed genes and isoforms (c) showed a significant enrichment of common genetic variants associated with the experience of psychotic episodes, but not schizophrenia. GWAS summary data from a chronic kidney disease study was used as a negative control.

Discussion

Our aim in this study was to investigate the differential contribution of genes in the Prader-Willi syndrome imprinted interval to psychiatric illness using mouse models. We have previously demonstrated abnormalities in a number of behavioural and cognitive endophenotypes of relevance to psychiatric illness in a PWS imprinting centre deletion (PWS-IC) mouse model, one of the genotypes in people with PWS in which psychotic illness is most prevalent. Here we show that these same behavioural and cognitive measures are not altered in a mouse model for PWS that has a deletion limited to the PWS critical interval (PWS-cr mice). This suggests these behaviours are not a general feature of PWS models, but something specific to the PWS-IC model. We then used RNA-seq to examine the molecular bases of these differences in behaviour between the PWS-IC and PWS-cr

models. Although a number of interesting gene expression and alternate splicing events were detected in the brains of PWS-cr mice, a greater number were seen in the PWS-IC model. Moreover, these changes in gene expression and splicing in the PWS-IC brain were enriched for common genetic variants associated with the experience of psychotic episodes, but not those associated with schizophrenia. These findings shed light on the nature of psychotic illness in PWS and have implications for possible therapeutic strategies.

The behaviour of mice carrying a deletion of the PWS-cr has been extensively characterized with respect to various aspects of the core PWS phenotype including hunger and sleep (27-30), but behaviours relating to psychiatric illness have not been directly studied. Our previous behavioural studies of the PWS-IC model indicated reduced activity, abnormal sensory-motor gating, and deficits in attention (17). In contrast, here our studies of the PWS-cr mouse model suggests that deletion of the critical region has only very subtle effects on these same behavioural measures. Specifically, the PWS-cr mice showed no abnormalities in anxiety as measured by the EPM and OF tests, locomotor activity, sensory-motor gating as measured by PPI, or attention and impulsivity as measured by the 5-CSRTT. While we cannot rule out Type II error, we note that the PWS-cr behavioural cohort was considerably larger than that used for the PWS-IC mice. Furthermore, although conducted at separate times, critically the behavioural analysis of PWS-cr mice was conducted in an identical manner to that of the PWS-IC model, including the equipment, procedures, software, location (both Institute and rooms) and genetic background strain (F1 cross between C57Bl/6 and CD1).

Of all the behaviours assessed only ASR was altered in PWS-cr mice. However, in contrast to the enhanced startle response seen in PWS-IC mice, PWS-cr exhibited a decreased startle response. A reduced acoustic startle response could potentially be confounded by hearing loss and this cannot be completely ruled out here. However, this explanation would seem unlikely given the sensitivity of the response in PWS-cr mice to different startle pulse intensities (Supplementary figure 1), and that there are no records of hearing loss in any mouse models for PWS, or indeed in PWS patients. The combined phenotype of an increased startle response and deficient pre-pulse inhibition seen in the PWS-IC mouse model is indicative of psychotic-like behaviour (31, 32). Consequently, the lack of any PPI deficit in PWS-cr deletion model suggests that their reduced startle response is a distinct behavioural outcome. However, whilst increased ASR has been linked to stress, pain and fear (33-35), reduced startle response has been reported in individuals with major depressive disorder and anhedonia, as well as an effect of anxiolytic drugs (36-38). With this in mind, the reduced startle endophenotype seen in PWS-cr mice could fit the psychiatric profile of individuals with PWS that carry deletions spanning the critical interval, who are prone to depression and anxiety, but not psychosis (9-11). However, a broader variety of behavioural analysis is required to properly establish a depression phenotype in the PWS-cr mice.

The behavioural comparison of these two models indicates that the PWS-cr mice do not exhibit the same range of cognitive and psychotic illness endophenotypes as seen in the PWS-IC mice. We next explored the molecular bases for these behavioural differences using RNA-seq. Analysis of RNA-seq data comparing PWS-cr neonate brain with wild-type littermates revealed only very subtle changes of gene expression and alternative splicing, with seven differentially expressed genes (four predicted transcripts) and five differentially spliced transcripts. This may seem at odds with previous findings demonstrating differential expression of upwards of 200 genes in PWS-cr mice (39-41). However, these studies looked at gene expression in more focused tissues (e.g. hypothalamus) or cell types, whereas our study examined bulk whole brain, therefore obscuring some of the subtler region- or cell-specific changes, and revealing only the strongest effects across the whole brain. In addition to *Snord116* and *Ipw*, that are directly affected by the mutation (Supplementary figure 2), our data suggest that the PWS-cr deletion also leads to mis-expression of the PWS interval gene *Necdin,* which is an important regulator of neuronal outgrowth and differentiation (42, 43), loss of which is associated with motor deficits and enhanced learning and memory (44, 45). Whether the increased expression of *Necdin* is due to loss of suppression by *Snord116* and/or *Ipw*, in manner similar to the

regulatory loop between *Snord116* and *Snord115* (39) or the suppressive role of *IPW* on the maternally expressed genes from the *DLK1-DIO3* imprinted cluster (40), remains to be established. Nevertheless, it could be that this altered expression contributes to the learning and memory deficits seen in PWS-cr mice (46).

In contrast to the limited changes seen in the PWS-cr model, RNA-seq analysis of PWS-IC neonatal brain revealed a markedly larger number of DEGs and isoform changes. We suggest these gene expression changes may contribute to behavioural and cognitive deficits seen in the PWS-IC but not the PWS-cr mice and, by extension, may underpin the increased incidence of psychotic illness seen in individuals with imprinting centre deletions or mUPD (47). In line with this, as well as altered expression of a greater number of genes from within the PWS interval, there were changes in a number of other neurally important genes. These included the myelin protein proteolipid protein *Plp1*, the gamma-aminobutyric acid (GABA) A receptor subunit gamma 3 gene *Gabrg3*, and the circadian rhythm gene *Per1*.

In order to more formally assess the relevance of these gene expression and isoform changes to psychiatric illness we examined the DEGs and isoforms for enrichment of genetic variants relevant to psychosis. The GWAS data we used to look for genetic correlation between our samples with the occurrence of psychotic episodes was from a recent UK Biobank study that sampled individuals with a history of psychotic experiences including hallucinations and delusional ideation, but specifically without a diagnosis of schizophrenia (25). These authors found a shared genetic liability with schizophrenia from an external GWAS dataset (24), which we also used for our analysis here. Unsurprisingly given the limited number of changes, there was no enrichment in the gene expression changes seen in the PWS-cr neonate brain. However, there was an enrichment of genetic variants common to the experience of psychotic episodes in the gene expression changes seen in the PWS-IC neonate brain. The findings substantiate the relevance of the gene expression changes and behavioural endophenotype observed in the PWS-IC mouse model to the psychiatric profile of individuals with individuals with an IC deletion or a mUPD of chromosome 15. Strikingly, there was not an enrichment of genetic variants associated with schizophrenia. This finding provides a biological signal that confirms the observation that the psychotic illness seen in PWS is distinct from schizophrenia (13) and should not be treated in the same manner (16).

A closer inspection of the individual genes that carry genetic variants common for the occurrence of psychotic episodes shows that three of these genes are PEGs from within the PWS cluster itself, specifically *MAGEL2, MKRN3,* and *NECDIN.* It is likely that loss of expression of these genes in the PWS-IC mice brain is a strong, but importantly, not the sole driver of enrichment for the GWAS variants associated with multiple psychotic illness. This then suggests a subtly different model of vulnerability to psychotic illness in PWS. Previously a two-hit model was proposed, in which loss of PEG expression in PWS leads to basal neurodevelopmental problems and it is against this background that increased expression of MEGs (most probably *UBE3A*) in matDUP cases leads to psychotic illness (8, 48). However, our data indicate a two-hit model where loss of expression of paternally expressed imprinted genes not only causes neurodevelopmental problems but also predisposes to psychotic illness, and mis-expression of MEGs further exacerbates this vulnerability leading to the very high rates of psychotic illness seen in matDUP cases.

Overall, our findings suggest a cumulative effect of several of the genes from the PWS locus on the behavioural endophenotypes typical of Prader-Willi syndrome. The critical interval deletion, albeit important in the core phenotypes seen in PWS models, only induced very subtle behavioural and cognitive differences in the studies we conducted. By comparing the gene expression and isoform profile of the PWS-cr and PWS-IC lines were able to identify brain molecular changes of relevance to the psychiatric profile of PWS, underlined by the enrichment of genetic variants for psychotic episodes in the PWS-IC gene changes. The fact that genetic variants for schizophrenia showed no

enrichment reflects the distinct nature of psychotic illness in PWS and could have implications for therapeutic strategies used to treat psychosis in individuals with PWS.

Methods

Behavioural experiments

Animal husbandry

Due to the genomic imprinting of the 15q11.2-q13 locus, heterozygous mice carrying the deletion on the paternally inherited chromosome are expected to have a complete loss of expression of the critical interval. PWS-cr$^{m+/p-}$ mice (B6(Cg)-Snord116^{tm1Uta}/J) were purchased from The Jackson Laboratory. The females were mated with wild type c57BL/6 males in order to generate PWS-cr$^{m-/p+}$ mice, which carry the deletion on the silenced maternal copy of the critical interval mutation and are therefore expected not to exhibit any of the phenotypes associated with it. The behavioural cohort of 24 PWS-cr$^{m+/p-}$ (16 female, 8 male) and 31 wild type (17 female, 14 male) was produced by breeding PWS-cr$^{m-/p+}$ males with wild type CD1 females in order to exactly replicate the genetic background that we have used in a previous study, examining the behaviour of a PWS-IC mouse model (17). Weaning took place at approximately four weeks of age, when animals were housed in single sex and mixed genotype cages of 2-4 individuals. Ear tissue was collected for genotyping identification (Protocol in Appendix 3). Experimental testing began at approximately 8-9 weeks of age, and it took place in the order in which the tests are described below.

The animals were handled daily for two weeks prior to the start of experimentation in order to adapt them to being picked up. All mice were kept at a 12 h light/dark cycle, with the lights being switched on at 7:00 h every morning. Access to water and food was *ad libitum*, until two weeks before the start of the reward preference test, when the animals were gradually introduced to a water deprivation regime. During the first two days of this regime, the animals were given access to water for 4 hours per day. After that, the access to water was limited to 2 hours per day. The water deprivation regime continued all through the reward preference test and the 5-choice serial reaction time task. Water bottles were provided in the home-cages immediately after experimentation.

Elevated plus maze and open field

The elevated plus maze (EPM) and open field test (OF) were used to measure anxiety-related behaviours (49, 50). The EPM consisted of a plus shaped platform, with two enclosed arms (19x8x15 cm) and two open, exposed arms (19x8 cm). The maze was lifted 50 cm above the ground. The animals were placed in one of the enclosed arms of the maze at the beginning of the task, and allowed to freely explore for 5 min. The OF apparatus consisted of a square arena (75x75 cm) enclosed by walls (45 cm). The surface of the arena was divided into central zone (45x45 cm) and outer zone. The animals were placed into the outer zone of the arena and allowed to freely explore for 10 min. The movement of the animals in the EPM and OF was detected by a camera attached to a computer with Ethovision software (XT, Noldus, Netherlands). Time spent in movement during the trials was calculated post-hoc by dividing the distance by the velocity.

Locomotor activity

Spontaneous locomotor activity (LMA) of mice was explicitly measured in custom made chambers (21x36x20 cm) fitted with infrared beams, situated 3 cm from either end of the box and 1 cm from the floor of the box. The test was run in complete darkness in order to remove the anxiogenic effect of light. The mice were placed in the boxes and allowed to roam freely for 2 h, while the disturbance of the beams by their movement was recorded by an Acorn computer with an Arachnid software (Cambridge Cognition LTD). Number of beam breaks and the consecutive breaking of the two infrared beams (referred to as a "run") were considered the main measures of locomotor activity. The animals were assessed at the same time for two consecutive days, in order to assess habituation to a novel environment within and between sessions.

Acoustic startle and pre-pulse inhibition

Acoustic startle response (ASR) and pre-pulse inhibition (PPI) were used to test for phenotypes associated with psychotic and affective illness (31, 32). ASR and PPI were measured in a soundproofed SR LAB startle chamber with a speaker (San Diego Instruments), in which the mice were securely placed inside a plexiglass tube (3.5 cm dia). The trial commenced with 5 min of 70 dB white noise in order to allow for habituation to the environment. After habituation the mice were exposed to a range of acoustic stimuli, including pulse-alone stimuli at 120 dB and 105 dB above background noise (70dB) for 40 ms, then startle stimuli preceded by 20 ms pre-pulses of 4 dB ,8 dB and 16 dB, 80 ms prior to startle stimuli. All startle and pre-pulse stimuli were presented against a background noise of 70 dB. The startle movement in response to the noise was recorded with a piezoelectric accelerometer and normalized by body weight before analysis. PPI of the ASR was calculated as the percentage reduction in startle amplitude between responses to startle stimuli alone (averaged cross 13 trials) and pre-pulse trials (averaged for the 5 presentations at each pre-pulse amplitude.

Reward preference test

Prior to the 5-CSRTT and after a transition to a restricted home-cage water access, animals were gradually exposed to the substance (10% condensed milk in tap water) which was to be used as a reward for the tasks later on. The animals were placed individually in a cage without sawdust, for a 10 min session each day, over a 7-day period. Each cage contained two small plastic containers (~1 cm high x ~2 cm diameter) secured to the floor by Velcro, placed equidistant from the end wall, and evenly spaced across the width of the cage. On the first two days, both containers contained tap water. On the following four days, one container contained the condensed milk dilution and the other - water, with the positions alternating each day of testing. On the final day, both containers were filled with the condensed milk dilution. All consumption was measured in grams by weighing the containers before and after the task. The preference to the reward was calculated for days 3 to 6 as a percentage of total volume consumed.

5-choice serial reaction time task

The 5-choice serial reaction time task (5-CSRTT) is commonly used to measure visuo-spatial attention and impulsivity in rodent models of disease (51, 52). The 5-CSRTT was commenced on the day following the final day of the reward preference test and broadly followed the protocol described by

(53). For this task, mice were trained to respond to a brief visual stimulus appearing pseudorandomly at one of five locations, in order to gain a condensed milk reward. The variables analysed as indicative of performance were accuracy (ratio of correct to total trials), omissions (ratio of omitted trials to total trials), and premature responses (number of responses that took place before the stimulus was presented). Other variables such as completed trials (calculated as ratio out of 60), correct reaction time, reward collection latency, number of food magazine entries and total number of trial perseveration were also kept track of for further analysis. Subjects were trained to a set baseline performance observed at a stimulus duration of 0.8 s (>30 completed trials, >75% accuracy, and <25% omissions). Full details can be found in Supplementary information.

The conditions of the experiment were manipulated in order to study various aspects of attention and impulsivity. One of the manipulations entailed shortening the stimulus duration (SSD). During this session, stimuli of 0.6 s, 0.4 s and 0.2 s were interspersed in pseudorandom order along with the baseline stimulus duration of 0.8 s, in order to increase the difficulty of the task and to affect the accuracy of performance. In another manipulation lengthened it is (LITI) of 7 s, 9 s and 11 s were interspersed in pseudorandom order along with the baseline inter-trial interval of 5 s, in order to induce premature responses and test for impulsive behaviour.

Data analysis

All data were analysed using R Studio 1.1.383 (R Studio, Inc). Linear models (lm) and generalized linear models (glm) were built with genotype and sex as fixed effects. Since interactions between sex and genotype were not observed, the effect of sex is not reported in this paper. Logistic glm from the binomial family was used to analyse all proportional data. Mixed lms and glms with mouse ID as a random effect were built using the lme4 package (54), for data in which multiple measures were taken from each individual at different conditions. In the LMA mixed model, day of testing and time bins were added as fixed effects. In the ASR mixed model, startle trial number was added as fixed effect. In the PPI mixed model, the different pre-pulse decibels (4 dB, 8dB, 16 dB) were added as fixed effects. For the 5-CSRTT SSD and LITI manipulations, stimulus duration and inter-trial interval were also added as fixed effects. The p-values and chi-square values (χ^2) presented in the results were calculated using likelihood ratio tests run with the inbuilt ANOVA function of R and comparing the full statistical models to reduced models with genotype removed as a fixed factor.

RNA sequencing

Tissue collection, preparation and RNA sequencing

For the RNA sequencing experiments, whole brain neonatal tissue was taken from 6 PWS-cr mice (3 females and 3 males) and 6 of their wild type littermates (3 females and 3 males), as well as from 4 PWS-IC$^{m+/p-}$ mice (all female) and 6 of their wild type littermates (2 female, 4 males). The collected tissue was snap frozen in dry ice and stored at -80°C until further use. RNA was extracted using the Direct-zol RNA Miniprep kit. Libraries were prepared with the KAPA mRNA HyperPrep kit (Roche Sequencing solutions) from total RNA of RQN > 8.5 as assessed by the Fragment Analyser system (Agilent Technologies). 800 ng of total RNA input was used for each sample. The mRNA was fragmented at 94°C for 6 minutes to achieve a mean insert size of 200-300 bp. 8 cycles were used in the final amplification. A final bead clean-up step was added at the end of the protocol before the libraries were taken forward for NGS on an Illumina HiSeq4000 system.

RNA sequencing analysis

For analysis of differential gene expression, sequencing reads were mapped to the mouse genome (version mm10) using the STAR software package (55). The featureCounts software (56) was used to count the raw reads, which were subsequently analysed in R Studio 1.1.383 using the DESeq2 package (57), with genotype, sex and litter as fixed effects into the statistical model. Hierarchical clustering showed a female wild type outlier in both PWS-cr and PWS-IC samples, which were removed from further analysis (Supplementary figure 4). The UCSC genome browser was used for visualization of the data. Differential isoform usage was pseudo-aligned to the mouse transcriptome (mm10) with the Kallisto software (58) which was also used to quantify the raw reads. The Kallisto files were then analysed in R studio via DEXSeq with the IsoformSwitchAnalyzeR package (59-61), with genotype, sex and litter added as fixed effects into the statistical model. All p-values were adjusted using the Benjamini-Hochberg method. Prediction of protein domains was run on EBI's Pfam webserver, which performs biosequence analysis using hidden Markov models (62). The data from the PWS-cr and PWS-IC models was analysed separately since difference between the wild type littermates of the two groups did not allow for pooling of the data. Genes that were differentially expressed or exhibited differential usage of isoforms between genotypes (padj<0.05) were pooled together for functional enrichment analysis using the g:Profiler web server (63).

Common variant enrichment analysis

Genes that were differentially expressed or exhibited differential usage of isotopes between genotypes (padj<0.05) were pooled together for common variant enrichment analysis. The mouse gene IDs were converted to their homologous human gene IDs. The genes were run for gene set analysis through MAGMA along with GWAS summary statistics for schizophrenia, psychosis and chronic kidney disease (24-26). Gene locations build 37 and reference data from European population was downloaded from CNCR CTGLab (https://ctg.cncr.nl/software/magma).

DATA AVAILABILITY

All behavioural data (Appendices 1 and 2) and all RNA-seq data, including gene ontology and genetic enrichment data (Appendices 3 to 7) are publicly available at the following 'Open science framework' link: https://osf.io/wnx8r/

FUNDING

This work was supported by a UKRI Medical Research Council studentship to SZ (1976193) and UKRI Biotechnology and Biological Sciences Research Council research grant to ARI (BB/J016756/1).

Cassidy, S.B., Schwartz, S., Miller, J.L. and Driscoll, D.J. (2011) Prader-Willi syndrome. *Genetics In Medicine*, **14**, 10.
Angulo, M.A., Butler, M.G. and Cataletto, M.E. (2015) Prader-Willi syndrome: a review of clinical, genetic, and endocrine findings. *Journal of Endocrinological Investigation*, **38**, 1249-1263.
Whittington, J. and Holland, A. (2018) A review of psychiatric conceptions of mental and behavioural disorders in Prader-Willi syndrome. *Neurosci Biobehav Rev*, **95**, 396-405.
Reddy, L.A. and Pfeiffer, S.I. (2007) Behavioral and Emotional Symptoms of Children and Adolescents with Prader-Willi Syndrome. *Journal of Autism and Developmental Disorders*, **37**, 830-839.

Thuilleaux, D., Laurier, V., Copet, P., Tricot, J., Demeer, G., Mourre, F., Tauber, M. and Jauregi, J. (2018) A model to characterize psychopathological features in adults with Prader-Willi syndrome. *American Journal of Medical Genetics Part A*, **176**, 41--47.

Vogels, A., Hert, M.D., Descheemaeker, M.J., Govers, V., Devriendt, K., Legius, E., Prinzie, P. and Fryns, J.P. (2004) Psychotic disorders in Prader–Willi syndrome. *American Journal of Medical Genetics Part A*, **127A**, 238-243.

Sinnema, M., Boer, H., Collin, P., Maaskant, M.A., van Roozendaal, K.E.P., Schrander-Stumpel, C.T.R.M. and Curfs, L.M.G. (2011) Psychiatric illness in a cohort of adults with Prader-Willi syndrome. *Research in Developmental Disabilities*, **32**, 1729-1735.

McNamara, G.I. and Isles, A.R. (2013) Dosage-sensitivity of imprinted genes expressed in the brain: 15q11-q13 and neuropsychiatric illness. *Biochem Soc Trans*, **41**, 721-726.

Sahoo, T., del Gaudio, D., German, J.R., Shinawi, M., Peters, S.U., Person, R.E., Garnica, A., Cheung, S.W. and Beaudet, A.L. (2008) Prader-Willi phenotype caused by paternal deficiency for the HBII-85 C/D box small nucleolar RNA cluster. *Nature Genetics*, **40**, 719.

Fontana, P., Grasso, M., Acquaviva, F., Gennaro, E., Galli, M.L., Falco, M., Scarano, F., Scarano, G. and Lonardo, F. (2017) SNORD116 deletions cause Prader-Willi syndrome with a mild phenotype and macrocephaly. *Clinical Genetics*, **92**, 440-443.

Bieth, E., Eddiry, S., Gaston, V., Lorenzini, F., Buffet, A., Conte Auriol, F., Molinas, C., Cailley, D., Rooryck, C., Arveiler, B. *et al.* (2015) Highly restricted deletion of the SNORD116 region is implicated in Prader–Willi Syndrome. *European Journal of Human Genetics*, **23**, 252-255.

Yang, L., Zhan, G.-d., Ding, J.-j., Wang, H.-j., Ma, D., Huang, G.-y. and Zhou, W.-h. (2013) Psychiatric Illness and Intellectual Disability in the Prader–Willi Syndrome with Different Molecular Defects - A Meta Analysis. *PLOS ONE*, **8**, e72640.

Soni, S., Whittington, J., Holland, A.J., Webb, T., Maina, E.N., Boer, H. and Clarke, D. (2008) The phenomenology and diagnosis of psychiatric illness in people with Prader–Willi syndrome. *Psychological Medicine*, **38**, 1505-1514.

Boer, H., Holland, A., Whittington, J., Butler, J., Webb, T. and Clarke, D. (2002) Psychotic illness in people with Prader Willi syndrome due to chromosome 15 maternal uniparental disomy. *Lancet*, **359**, 135-136.

Isles, A.R., Ingason, A., Lowther, C., Walters, J., Gawlick, M., Stober, G., Rees, E., Martin, J., Little, R.B., Potter, H. *et al.* (2016) Parental Origin of Interstitial Duplications at 15q11.2-q13.3 in Schizophrenia and Neurodevelopmental Disorders. *PLoS Genet*, **12**, e1005993.

Soni, S., Whittington, J., Holland, A.J., Webb, T., Maina, E., Boer, H. and Clarke, D. (2007) The course and outcome of psychiatric illness in people with Prader-Willi syndrome: implications for management and treatment. *J Intellect Disabil Res*, **51**, 32-42.

Relkovic, D., Doe, C.M., Humby, T., Johnstone, K.A., Resnick, J.L., Holland, A.J., Hagan, J.J., Wilkinson, L.S. and Isles, A.R. (2010) Behavioural and cognitive abnormalities in an imprinting centre deletion mouse model for Prader–Willi syndrome. *European Journal of Neuroscience*, **31**, 156-164.

Doe, C., Relkovic, D., Garfield, A., Dalley, J., Theobald, D., Humby, T., Wilkinson, L. and Isles, A. (2009) Loss of the imprinted snoRNA mbii-52 leads to increased 5htr2c pre-RNA editing and altered 5HT2CR-mediated behaviour. *Human Molecular Genetics*, **18**, 2140-2148.

Mistry, S., Harrison, J.R., Smith, D.J., Escott-Price, V. and Zammit, S. (2018) The use of polygenic risk scores to identify phenotypes associated with genetic risk of schizophrenia: Systematic review. *Schizophrenia Research*, **197**, 2-8.

Schork, A.J., Won, H., Appadurai, V., Nudel, R., Gandal, M., Delaneau, O., Revsbech Christiansen, M., Hougaard, D.M., Bækved-Hansen, M., Bybjerg-Grauholm, J. *et al.* (2019) A genome-wide association study of shared risk across psychiatric disorders implicates gene regulation during fetal neurodevelopment. *Nature Neuroscience*, **22**, 353-361.

Wippich, F., Bodenmiller, B., Trajkovska, Maria G., Wanka, S., Aebersold, R. and Pelkmans, L. (2013) Dual Specificity Kinase DYRK3 Couples Stress Granule Condensation/Dissolution to mTORC1 Signaling. *Cell*, **152**, 791-805.

Wang, L., Li, J., Shuang, M., Lu, T., Wang, Z., Zhang, T., Yue, W., Jia, M., Ruan, Y., Liu, J. *et al.* (2018) Association study and mutation sequencing of genes on chromosome 15q11-q13 identified GABRG3 as a susceptibility gene for autism in Chinese Han population. *Translational Psychiatry*, **8**, 152.

Greger, V., Knoll, J.H.M., Woolf, E., Glatt, K., Tyndale, R.F., Delorey, T.M., Olsen, R.W., Tobin, A.J., Sikela, J.M., Nakatsu, Y. *et al.* (1995) The γ-aminobutyric acid receptor γ3 subunit gene (GABRG3) is tightly linked to the α5 subunit gene (GABRA5) on human chromosome 15q11–q13 and is transcribed in the same orientation. *Genomics*, **26**, 258-264.

Pardiñas, A.F., Holmans, P., Pocklington, A.J., Escott-Price, V., Ripke, S., Carrera, N., Legge, S.E., Bishop, S., Cameron, D., Hamshere, M.L. *et al.* (2018) Common schizophrenia alleles are enriched in mutation-intolerant genes and in regions under strong background selection. *Nature Genetics*, **50**, 381-389.

Legge, S.E., Jones, H.J., Kendall, K.M., Pardiñas, A.F., Menzies, G., Bracher-Smith, M., Escott-Price, V., Rees, E., Davis, K.A.S., Hotopf, M. *et al.* (2019) Association of Genetic Liability to Psychotic Experiences With Neuropsychotic Disorders and Traits. *JAMA Psychiatry*, **76**, 1256-1265.

Wuttke, M., Li, Y., Li, M., Sieber, K.B., Feitosa, M.F., Gorski, M., Tin, A., Wang, L., Chu, A.Y., Hoppmann, A. *et al.* (2019) A catalog of genetic loci associated with kidney function from analyses of a million individuals. *Nature Genetics*, **51**, 957-972.

Lassi, G., Maggi, S., Balzani, E., Cosentini, I., Garcia-Garcia, C. and Tucci, V. (2016) Working-for-Food Behaviors: A Preclinical Study in Prader-Willi Mutant Mice. *Genetics*, **204**, 1129-1138.

Lassi, G., Priano, L., Maggi, S., Garcia-Garcia, C., Balzani, E., El-Assawy, N., Pagani, M., Tinarelli, F., Giardino, D., Mauro, A. *et al.* (2016) Deletion of the Snord116/SNORD116 Alters Sleep in Mice and Patients with Prader-Willi Syndrome. *Sleep*, **39**, 637-644.

Ding, F., Li, H.H., Zhang, S., Solomon, N., Camper, S., Cohen, P. and Francke, U. (2008) SnoRNA *Snord116 (Pwcr1/MBII-85)* Deletion Causes Growth Deficiency and Hyperphagia in Mice. *Plos One*, **3**.

Qi, Y., Purtell, L., Fu, M., Lee, N.J., Aepler, J., Zhang, L., Loh, K., Enriquez, R.F., Baldock, P.A., Zolotukhin, S. *et al.* (2016) Snord116 is critical in the regulation of food intake and body weight. *Scientific Reports*, **6**, 18614.

Akdag, S.J., Nestor, P.G., O'Donnell, B.F., Niznikiewicz, M.A., Shenton, M.E. and McCarley, R.W. (2003) The startle reflex in schizophrenia: habituation and personality correlates. *Schizophrenia research*, **64**, 165-173.

Parwani, A., Duncan, E.J., Bartlett, E., Madonick, S.H., Efferen, T.R., Rajan, R., Sanfilipo, M., Chappell, P.B., Chakravorty, S., Gonzenbach, S. *et al.* (2000) Impaired prepulse inhibition of acoustic startle in schizophrenia. *Biological Psychiatry*, **47**, 662-669.

Sorenson, C.A. and Swerdlow, N.R. (1982) The effect of tail pinch on the acoustic startle response in rats. *Brain Research*, **247**, 105-113.

Hebb, A.L.O., Zacharko, R.M., Gauthier, M. and Drolet, G. (2003) Exposure of mice to a predator odor increases acoustic startle but does not disrupt the rewarding properties of VTA intracranial self-stimulation. *Brain Research*, **982**, 195-210.

Davis, M., Falls, W.A., Campeau, S. and Kim, M. (1993) Fear-potentiated startle: A neural and pharmacological analysis. *Behavioural Brain Research*, **58**, 175-198.

Kaviani, H., Gray, J.A., Checkley, S.A., Raven, P.W., Wilson, G.D. and Kumari, V. (2004) Affective modulation of the startle response in depression: influence of the severity of depression, anhedonia, and anxiety. *Journal of Affective Disorders*, **83**, 21-31.

Commissaris, R.L., Fomum, E.A. and Leavell, B.J. (2004) Effects of buspirone and alprazolam treatment on the startle-potentiated startle response. *Depression and Anxiety*, **19**, 146-151.

Riba, J., Rodríguez-Fornells, A., Urbano, G., Morte, A., Antonijoan, R. and Barbanoj, M. (2001) Differential effects of alprazolam on the baseline and fear-potentiated startle reflex in humans: a dose-response study. *Psychopharmacology (Berl)*, **157**, 358-367.

Falaleeva, M., Surface, J., Shen, M., de la Grange, P. and Stamm, S. (2015) SNORD116 and SNORD115 change expression of multiple genes and modify each other's activity. *Gene*, **572**, 266-273.

Stelzer, Y., Sagi, I., Yanuka, O., Eiges, R. and Benvenisty, N. (2014) The noncoding RNA IPW regulates the imprinted DLK1-DIO3 locus in an induced pluripotent stem cell model of Prader-Willi syndrome. *Nature genetics*, **46**, 551-557.

Polex-Wolf, J., Lam, B.Y.H., Larder, R., Tadross, J., Rimmington, D., Bosch, F., Cenzano, V.J., Ayuso, E., Ma, M.K.L., Rainbow, K. *et al.* (2018) Hypothalamic loss of Snord116 recapitulates the hyperphagia of Prader-Willi syndrome. *The Journal of Clinical Investigation*, **128**, 960-969.

Lee, S., Walker, C.L., Karten, B., Kuny, S.L., Tennese, A.A., O'Neill, M.A. and Wevrick, R. (2005) Essential role for the Prader–Willi syndrome protein necdin in axonal outgrowth. *Human Molecular Genetics*, **14**, 627-637.

Kuwajima, T., Nishimura, I. and Yoshikawa, K. (2006) Necdin Promotes GABAergic Neuron Differentiation in Cooperation with Dlx Homeodomain Proteins. *The Journal of Neuroscience*, **26**, 5383.

Muscatelli, F.o., Abrous, D.N., Massacrier, A., Boccaccio, I.n., Moal, M.L., Cau, P. and Cremer, H. (2000) Disruption of the mouse Necdin gene results in hypothalamic and behavioral alterations reminiscent of the human Prader–Willi syndrome. *Human Molecular Genetics*, **9**, 3101-3110.

Andrieu, D., Meziane, H., Marly, F., Angelats, C., Fernandez, P.-A. and Muscatelli, F. (2006) Sensory defects in Necdin deficient mice result from a loss of sensory neurons correlated within an increase of developmental programmed cell death. *BMC Developmental Biology*, **6**, 56.

Adhikari, A., Copping, N.A., Onaga, B., Pride, M.C., Coulson, R.L., Yang, M., Yasui, D.H., LaSalle, J.M. and Silverman, J.L. (2019) Cognitive deficits in the Snord116 deletion mouse model for Prader-Willi syndrome. *Neurobiology of learning and memory*, **165**, 106874-106874.

Boer, H., Holland, A., Whittington, J., Butler, J., Webb, T. and Clarke, D. (2002) Psychotic illness in people with Prader Willi syndrome due to chromosome 15 maternal uniparental disomy. *The Lancet*, **359**, 135-136.

Soni, S., Whittington, J., Holland, A.J., Webb, T., Maina, E.N., Boer, H. and Clarke, D. (2008) The phenomenology and diagnosis of psychiatric illness in people with Prader-Willi syndrome. *Psychol Med*, **38**, 1505-1514.

Walf, A.A. and Frye, C.A. (2007) The use of the elevated plus maze as an assay of anxiety-related behavior in rodents. *Nature Protocols*, **2**, 322-328.

Prut, L. and Belzung, C. (2003) The open field as a paradigm to measure the effects of drugs on anxiety-like behaviors: a review. *European Journal of Pharmacology*, **463**, 3-33.

Robbins, T. (2002) 5-choice serial reaction time task: behavioural pharmacology and functional neurochemistry. *Psychopharmacology*, **163**, 362:380.

Humby, T., Laird, F.M., Davies, W. and Wilkinson, L.S. (1999) Visuospatial attentional functioning in mice: interactions between cholinergic manipulations and genotype. *European Journal of Neuroscience*, **11**, 2813-2823.

Humby, T., Wilkinson, L. and Dawson, G. (2005) Assaying Aspects of Attention and Impulse Control in Mice Using the 5-Choice Serial Reaction Time Task. *Current Protocols in Neuroscience*, **31**, 8.5H.1-8.5H.15.

Bates, D., Maechler, M., Bolker, B. and Walker, S. (2015) Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, **67**, 1-48.

Dobin, A. and Gingeras, T.R. (2015) Mapping RNA-seq Reads with STAR. *Current protocols in bioinformatics*, **51**, 11.14.11-11.14.19.

Liao, Y., Smyth, G. and Shi, W. (2014) featureCounts: an Efficient General Purpose Program for Assigning Sequence Reads to Genomic Features *Bioinformatics*, **30**, 923-930.

Anders, S. and Huber, W. (2010) Differential expression analysis for sequence count data. *Genome Biology*, **11**, R106.

Bray, N.L., Pimentel, H., Melsted, P. and Pachter, L. (2016) Near-optimal probabilistic RNA-seq quantification. *Nature Biotechnology*, **34**, 525-527.

Vitting-Seerup, K. and Sandelin, A. (2017) The Landscape of Isoform Switches in Human Cancers. *Molecular Cancer Research*, **15**, 1206.

Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W. and Smyth, G.K. (2015) limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Research*, **43**, e47-e47.

Anders, S., Reyes, A. and Huber, W. (2012) Detecting differential usage of exons from RNA-seq data. *Genome Research*, **22**, 2008-2017.

Punta, M., Coggill, P.C., Eberhardt, R.Y., Mistry, J., Tate, J., Boursnell, C., Pang, N., Forslund, K., Ceric, G., Clements, J. *et al.* (2011) The Pfam protein families database. *Nucleic Acids Research*, **40**, D290-D301.

Raudvere, U., Kolberg, L., Kuzmin, I., Arak, T., Adler, P., Peterson, H. and Vilo, J. (2019) g:Profiler: a web server for functional enrichment analysis and conversions of gene lists (2019 update). *Nucleic Acids Research*, **47**, W191-W198.

Comparison of mouse models reveals the molecular bases for psychotic illness in Prader-Willi syndrome – Supplementary information

Simona Zahova[1], Trevor Humby[2], Jennifer R. Davies[1], Joanne E. Morgan[1], Anthony R. Isles[1]*

[1]MRC Centre for Neuropsychiatric Genetics and Genomics, School of Medicine, Cardiff University, Hadyn Ellis Building, Maindy Road, Cardiff, CF24 4HQ, U.K.; ZahovaS@cardiff.ac.uk

[2]School of Psychology, Cardiff University, Tower Building, 70 Park Place, Cardiff, CF10 3AT, U.K. HumbyT@cardiff.ac.uk

* Correspondence: Anthony R. Isles; IslesAR1@cardiff.ac.uk; MRC Centre for Neuropsychiatric Genetics and Genomics, School of Medicine, Cardiff University, Hadyn Ellis Building, Maindy Road, Cardiff, CF24 4HQ, U.K.

Supplementary Figure 1. Startle response to gradual increasing startle pulses (background noise 70dB) in PWS-cr and wild-type littermate mice. Data points are means for each group ±SEM.

Supplementary Figure 2. qPCR assay of the critical interval through stages E13.5, E18.5 and P0 of wild type mouse development. Error bars indicate 95% confidence intervals, the circles on the bars indicate mean average.

Supplementary Figure 3. UCSC browser track image of STAR generated transcript counts at the critical interval of PWS-cr mouse model and their wild type littermates shows the deletion in the PWS-cr mice spans all the copies of *Snord116,* as well as most of the exons of the *Ipw* non-coding RNA.

Supplementary Figure 4. Top 100 differentially expressed genes in the PWS-cr and PWS-IC mouse model samples. Hierarchical clustering heatmap in the PWS-cr mouse model showed no effect of genotype on gene expression. An outlier (far left) was excluded from further analysis. The PWS-IC samples, on the other hand, clustered by genotype and showed differences in gene expression between heterozygous and wild type. An outlier (far left) was excluded from further analysis.

Detailed 5-choice serial reaction time task methods

The task was carried out using a bank of eight 9-hole chambers (Campden Instruments, UK), each enclosed in a sound attenuating chamber. Each of the equidistantly spaced holes across the array could present a visual stimulus and also register a nose-poke response by the breaking of an infrared beam. For this task, 4 of the 9 holes in the array (number 2, 4, 6 and 8) were blocked by masks. On the far side of the nose-poke array there was a food magazine, to which the reward was delivered by a peristaltic pump, via silicone tubing. Entrances to the food magazine were also recorded by means of an infrared. The chambers could be illuminated by lights in their side walls ("house lights), although all sessions were conducted in the dark apart from time out periods.

The mice were given a single 5-CSRTT session each day, where a session lasted for either a maximum of 60 trials or 20 minutes. Each trial at baseline conditions consisted of a stimulus presentation, pseudorandomly presented to one of the 5 locations, and a 5 s limited-hold period during which the stimulus was extinguished but the subject could still make a response and a 5 s inter-trial interval (ITI). A correct response (i.e. a nose-poke where the stimulus was illuminated) was rewarded by a 70 µl delivery of the reward, whereas following an incorrect response or an omission (i.e. no response) a 5 s time out period was initiated consisting of illumination of the house light. Responding during the ITI was defined as a "premature response" and initiated a time out and restarted the current trial.

The 5-CSRTT was commenced on the day following the final day of the reward preference test and broadly followed the protocol described by (53). In the first stage of shaping, the mice were habituated to the chambers for half an hour, during which reward was presented to the food magazine every 30 s. Mice completed a minimum of 3 sessions and then if they made >35 food magazine entries in two consecutive sessions they were moved to the next stage of training. Mice remained at this stage until they met the same criteria.

In the next stage of training, the mice performed a simplified version of the 5-CSRTT whereby only the central of the five holes was illuminated, with a stimulus duration of 60 s (1-CSRT task). Once performance reached criteria (>30 completed trials, >75% accuracy, <25% trial omissions) for two consecutive days, mice were moved on to the 5-CSRTT proper, with an initial stimulus duration of 60 s. When mice met the above criteria, the stimulus duration was then decreased in the following way: 32 s, 16 s, 8 s, 4 s, 2 s, 1,8 s, 1.6 s, 1.4 s, 1.2 s, 1 s and 0.8 s. A stimulus duration of 0.8 s was considered baseline, and once reached, mice were kept on it for a minimum of five sessions and had to meet performance criteria for a minimum of two consecutive sessions, thus showing "stable" baseline performance before task manipulations were implemented to further study different aspects of attention and impulsivity. The average of the two final consecutive sessions before the first manipulation session was used to analyse performance at baseline conditions.