PRESENILIN 1 mutations causing early-onset familial Alzheimer's
disease or familial acne inversa differ in their effects on genes
facilitating energy metabolism and signal transduction

Karissa **Barthelson**[a*], Yang **Dong**[a], Morgan **Newman**[a] and Michael **Lardelli**[a]

[a] Alzheimer's Disease Genetics Laboratory, School of Biological Sciences, University of
Adelaide, North Terrace, Adelaide, SA 5005, Australia

* Corresponding author

Complete correspondence address: Room 1.24, Molecular Life Sciences Building. North

Terrace Campus. The University of Adelaide, SA 5005, Australia.

Telephone: 83134863

Email: karissa.barthelson@adelaide.edu.au

Running title: RNA-seq analysis of psen1 mutations

ABSTRACT

Background: The most common cause of early-onset familial Alzheimer's disease (EOfAD) is

mutations in *PRESENILIN 1* (*PSEN1*) allowing production of mRNAs encoding full-length, but

mutant, proteins. In contrast, a single known frameshift mutation in *PSEN1* causes familial

acne inversa (fAI) without EOfAD. The molecular consequences of heterozygosity for these

mutation types, and how they cause completely different diseases, remains largely

unexplored.

Objective: To analyse brain transcriptomes of young adult zebrafish to identify similarities

and differences in the effects of heterozygosity for *psen1* mutations causing EOfAD or fAI.

Methods: RNA sequencing was performed on mRNA isolated from the brains of a single

family of 6-month-old zebrafish siblings either wild type or possessing a single, heterozygous

EOfAD-like or fAI-like mutation in their endogenous *psen1* gene.

Results: Both mutations downregulate genes encoding ribosomal subunits, and upregulate

genes involved in inflammation. Genes involved in energy metabolism appeared significantly

affected only by the EOfAD-like mutation, while genes involved in Notch, Wnt and

neurotrophin signalling pathways appeared significantly affected only by the fAI-like

mutation. However, investigation of direct transcriptional targets of Notch signalling

revealed possible increases in γ-secretase activity due to heterozygosity for either *psen1*

mutation. Transcriptional adaptation due to the fAI-like frameshift mutation was evident.

Conclusion: We observed both similar and contrasting effects on brain transcriptomes of the

heterozygous EOfAD-like and fAI-like mutations. The contrasting effects may illuminate how

these mutation types cause distinct diseases.

INTRODUCTION

Cases of Alzheimer's disease (AD) can be classified by age of onset and mode of inheritance.

Dominant mutations in a small number of genes cause AD with an age of onset younger than

65 years (early onset familial AD, EOfAD). On a population basis, around 60% of the

mutations causing EOfAD occur in one gene, *PRESENILIN 1* (*PSEN1*) [1-4].

PSEN1 encodes a multi-pass transmembrane protein resident in the endoplasmic reticulum,

plasma membrane, endolysosomal pathway and other membranes [5, 6]. It has nine

recognised transmembrane domains [7]. A tenth transmembrane domain may exist when

PSEN1 protein is in its holoprotein state [7], before it undergoes autocatalytic

endoproteolysis to form the active catalytic core of γ-secretase [8], an enzyme complex

consisting of PSEN1 (or PSEN2), and the proteins NCSTN, PSENEN, and APH1A (or APH1B) [9,

10].

As a locus for genetic disease, *PSEN1* is truly remarkable both for the number of mutations

found there, and the variety of diseases these mutations cause. Mutations have been found

associated with Pick's disease [11], dilated cardiomyopathy [12] and acne inversa [13].

However, over 300 mutations of *PSEN1* are known to cause EOfAD

(www.alzforum.org/mutations/psen-1). In total, these mutations affect 161 codons of the

gene. Remarkably, the mutations are widely distributed in the PSEN1 coding sequence, but

are particularly common in the transmembrane domains. Only three regions of the PSEN1

protein are mostly devoid of EOfAD mutations: upstream of the first transmembrane

domain; a large part of the "cytosolic loop domain" (cytosolic loop 3); and the last two thirds

of the 9th transmembrane domain together with the lumenal C-terminus (see **Figure 1**).

The most common outcome of mutation of a protein sequence is either no effect or a

detrimental effect on the protein's evolved activity. Only rarely are mutations selectively

advantageous so that they enhance organismal survival and reproduction. The very large

number of EOfAD-causative mutations in *PSEN1* and their wide distribution in the protein

coding sequence is consistent with a loss-of-function. However, this cannot be a simple loss

of γ-secretase activity, as EOfAD-causative mutations have never been found in the genes

encoding the other components of γ-secretase complexes (other than less frequent

mutations in the *PSEN1* homologous gene, *PSEN2*, reviewed in [14]). Also, an *in vitro* analysis

of 138 EOfAD mutations of *PSEN1* published in 2017 by Sun et al. [15] found that

approximately 10% of these mutations actually increased γ-secretase activity.

Currently, the most commonly discussed hypothetical mechanism addressing how EOfAD

mutations of *PSEN1* cause disease is that these act through "qualitative changes" to γ-

secretase cleavage of the AMYLOID β A4 PRECURSOR PROTEIN (AβPP) to alter the length

distribution of the AMYLOID β (Aβ) peptides derived from it [16]. However, the

comprehensive study of Sun et al. revealed no consistency in the effects of *PSEN1* EOfAD

mutations on Aβ length distribution. The single consistent characteristic of all EOfAD

mutations in both *PSEN1* and *PSEN2* is that these permit production of transcripts with

coding sequences containing in-frame mutations, but terminated by the wild type stop

codons (i.e. they still permit production of a full length protein). This phenomenon was first

noted by De Strooper in 2007 [17] and described in detail by us in 2016 (the "reading frame

preservation rule" [18]). The universality of this rule, and that it reflects a critical feature of

the EOfAD-causative mechanism of *PSEN1* mutations, is shown by the fact that the

P242LfsX11 frameshift mutation of *PSEN1* (hereafter referred to as P242fs) causes a

completely different inherited disease, familial acne inversa (fAI, also known as hidradenitis

suppurativa), *without* EOfAD [13]. (Recently, a frameshift mutation in *PSEN1,* H21PfsX2, was

identified in an early-onset AD patient. However, whether the mutation is causative of

EOfAD mutation is still uncertain [19]. Questionable additional claims of EOfAD-causative

frameshift mutations in PSEN genes have been made and are reviewed in [18].) Critically, fAI

can also be caused by mutations in *NCSTN* and *PSENEN* [13], strongly supporting that this

disease is due to changes in γ-secretase activity.

Understanding the role of *PSENs* and their mutations is complicated by the partial functional

redundancy shared by *PSEN1* and *PSEN2* and their complex molecular biology. For example,

the PSEN1 holoprotein has been shown to have γ-secretase-independent activities required

for normal lysosomal acidification [20], can form multimers [21-24], and may interact with

the HIF1α protein [25-27] that is critical both for responses to hypoxia and for iron

homeostasis (reviewed in [28]). Additionally, within γ-secretase complexes, the PSENs act to

cleave at least 149 different substrates [29]. To simplify analysis, most previous investigation

of PSEN activity has involved inactivation (knock-out) of *PSEN1* and/or *PSEN2* in cells or

animals, and expression of only single forms of mutant PSEN (i.e. without simultaneous

expression of wild type forms). Forced expression of *PSEN* genes is also usually at non-

physiological levels which has unexpected regulatory feedback effects [30]. In humans,

investigating *PSEN*'s role in AD at the molecular level is restricted to post-mortem brain

tissues. However, these show substantially different patterns of gene expression compared

to the brains of people with mild cognitive impairment (MCI) or age-matched healthy

controls [31]. Since AD is thought to take decades to develop [32], we must understand the

pathological effects of EOfAD mutations in young adult brains if we wish to identify

preventative treatments. For this reason, we must model EOfAD mutations in animals.

The overwhelming majority of animal modelling of AD has utilised transgenic models

favoured for their apparent, partial reproduction of Aβ histopathology and easily discernible

cognitive disabilities. However, the relationship between Aβ histopathology and cognitive

change in these models is questionable [33]. Additionally, the most detailed form of

molecular phenotyping currently available, transcriptome analysis, shows little consistency

between the disturbed brain gene transcription of various transgenic models and limited

concordance between them and human sporadic AD brain transcriptomes [34]. "Knock-in"

mouse models of single EOfAD mutations (e.g. [35, 36]) make the fewest assumptions

regarding the pathological mechanism(s) of AD and most closely replicate the human EOfAD

genetic state (i.e. incorporating a single, dominant, endogenous mutation in the

heterozygous state). However, the brain transcriptomes of these mice have never been

analysed, and interest in them waned due to their lack of Aβ histopathology and mild

cognitive effects.

Analysis of mouse brain transcriptomes is complicated by strong effects on gene expression

of sex [37, 38] and, potentially, litter-of-origin (i.e. due to environmental and genotypic

variation) (K. Barthelson, unpublished results). In contrast, zebrafish brain transcriptomes

show only subtle influences of sex, and very large numbers of siblings can be generated from

single mating event, alleviating potential litter-of-origin issues [39-43]. In 2014, our

laboratory began a program of creating knock-in models of EOfAD-like (and non-EOfAD-like)

mutations in the zebrafish genes orthologous to *PSEN1*, *PSEN2*, and *SORL1*. In 2019 we

began publishing the results of transcriptome analyses of the young adult brains of these fish

[39-46] as we attempt to establish what effect(s) all the EOfAD mutations have in common

(and differentiate them from the non-EOfAD-like mutations).

Our previous analyses of an EOfAD-like mutation in the zebrafish *psen1* gene, Q96_K97del,

revealed very significant effects on the expression of genes involved in mitochondrial

function, lysosomal acidification, and iron homeostasis [45, 46]. Although Q96_K97del

follows the reading-frame preservation rule of EOfAD mutations in *PSEN1*, it is not an exact

equivalent of any human EOfAD-causative mutation. Consequently, in this study we aimed

to generate an additional, exactly equivalent, model of a human *PSEN1* EOfAD mutation. For

technical reasons, the T428 codon of zebrafish *psen1* (equivalent to the T440 codon of

human PSEN1) was predicted to be readily targetable using CRISPR-Cas9 technology, and we

subsequently deleted this codon in the zebrafish gene. This generated a zebrafish model of

the human EOfAD mutation $PSEN1^{T440del}$ [47]. This mutation was identified in a Japanese

man classified as displaying a mixed dementia phenotype (variant AD with spastic

paraparesis, Parkinson's disease and dementia with Lewy bodies). Also, to understand how

reading-frame preserving and frameshift mutations can cause completely different diseases

we generated a frameshift mutation in zebrafish *psen1*, $psen1^{W233fs}$, very similar to the fAI-

causative P242fs mutation of human *PSEN1*. We then performed an RNA-seq analysis with

high read depth and large sample numbers to compare the brain transcriptomes of fish from

a single family of young adult siblings heterozygous for either mutation or wild type. We

observed subtle, and mostly distinct, effects of the two mutations. In particular, changes in

the fAI-like brain transcriptomes implied significant effects on Notch, Wnt, neurotrophin,

and Toll-like receptor signalling, while changes in the EOfAD-like brain transcriptomes

implied effects on oxidative phosphorylation similar to those previously seen for EOfAD-like

mutations in *psen1* [45], *psen2* [40, 43], and *sorl1* [39, 41].

MATERIALS AND METHODS

Zebrafish husbandry and animal ethics

157 All zebrafish (Tübingen strain) used in this study were maintained in a recirculating water

158 system on a 14 hour light/10 hour dark cycle, fed dry food in the morning and live brine

159 shrimp in the afternoon. All zebrafish work was conducted under the auspices of the Animal

160 Ethics Committee (permit numbers S-2017-089 and S-2017-073) and the Institutional

161 Biosafety Committee of the University of Adelaide.

CRISPR-Cas9 genome editing

To mutate zebrafish *psen1,* we used the Alt-R® CRISPR-Cas9 system (Integrated DNA

Technologies, Coralville, IA, USA). To generate the T428del mutation (EOfAD-like) in exon 11

of *psen1*, we used a custom-designed crRNA recognising the sequence 5'

CTCCCCATCTCCATAACCTT 3' and a PAM of CGG. For the W233fs mutation (fAI-like), the

crRNA was designed to recognise the sequence 5' GATGAGCCATGCGGTCCACT 3' in exon 6

of *psen1,* with a PAM sequence of CGG. We aimed to generate exact equivalents of the

human P242fs mutation causing fAI, and the T440del mutation causing EOfAD by homology

directed repair (HDR). For the P242fs mutation, we used a plasmid DNA template as

described in [48] (synthesised by Biomatik, Kitchener, Ontario, Canada). For the T440del

mutation, we used an antisense, asymmetric single-stranded oligonucleotide with

phosphorothioate modifications (synthesised by Merck, Kenilworth, NJ, USA) as described in

[49] (HDR template DNA sequences are given in **Supplementary File 1.**)

Each crRNA was annealed with an equal amount of Alt-R® CRISPR-Cas9 tracrRNA (IDT) in

nuclease free duplex buffer (IDT) by heating at 95°C for 5 minutes, then allowed to cool to

room temperature, giving sgRNA solutions of 33 µM (assuming complete heteroduplex

formation of the RNA molecules). Then, 1 µL of the sgRNA solution was incubated with 1 µL

of Alt-R® S.p.Cas9 Nuclease 3NLS (IDT) at 64 µM at 37°C for 10 minutes to form

ribonucleoprotein (RNP) complexes. The final concentration for the linear ssODN for the

T428del mutation was 1 µM, and the final concentration of the plasmid DNA for the W233fs

mutation was 25 ng/µL. Approximately 2-5 nL of RNP complexes in solution with the

respective template DNAs were injected into Tübingen strain zebrafish embryos at the one

cell stage. The procedures followed for testing of the mutagenesis efficiencies of CRISPR-

Cas9 systems using allele-specific polymerase chain reactions and T7 endonuclease I assays,

and the breeding strategy to isolate the mutations of interest, are described in [41, 50].

RNA-seq raw data generation and processing

A

B

Figure 1: Experimental design. A Schematic of the human PSEN1 protein adapted from

https://www.alzforum.org/mutations/psen-1 with permission from FBRI LLC (Copyright © 1996–2020 FBRI LLC. All Rights

Reserved. Version 3.3 – 2020). Amino acid residues are colour-coded as to whether they are pathogenic for Alzheimer's

disease (red) or their pathogenicity is unclear (blue). The human mutation sites (P242 (fAI) and T440 (EOfAD)) are indicated by

black arrows. The site of the zebrafish W233fs-equivalent codon (W244) is shown by the red arrow. Note that the human T440

codon is equivalent to the zebrafish T428 codon. The residues equivalent to those deleted in the Q96_K97del mutation of

zebrafish *psen1* analysed previously are indicated by a red box. **B** A fish heterozygous for the W233fs mutation (fAI-like/+) was

mated with a fish heterozygous for the T428del mutation (EOfAD-like/+). The resulting family of fish contain genotypes fAI-like/+,

EOfAD-like/+, EOfAD-like/fAI-like and their wild type siblings. The pairwise comparisons performed in the RNA-seq experiment

are depicted. Since the EOfAD-like/fAI-like genotype is not representative of any human disease, it was not analysed.

We performed RNA-seq on a family of zebrafish as described in **Figure 1**. Total RNA (with

genomic DNA depleted by DNaseI treatment) was isolated from the brains of n = 4 fish per

genotype and sex as described in [41]. Then, 500 ng of total RNA ($RIN_e > 9$) was delivered to

the South Australian Genomics Centre (SAGC, Adelaide, Australia) for polyA+ library

construction (with unique molecular identifiers (UMIs)) and RNA-sequencing using the

Illumina Novaseq S1 2x100 SE platform.

The raw fastq files from SAGC were provided as 100 bp paired end reads as well as an index

file containing the UMIs for each read (over two Novaseq lanes which were subsequently

merged). The merged raw data was processed using a developed Nextflow [51] RNA-seq

workflow (see https://github.com/sagc-bioinformatics/sagc-rnaseq-nf). Briefly, UMIs were

added to headers of each read using *fastp* (v0.20.1). Alignment of the reads to the zebrafish

genome (GRCz11, Ensembl release 98) was performed using *STAR* (v2.5.3a). Then, reads

which contained the same UMI (i.e. PCR duplicates) were deduplicated using the *dedup*

function of *umi_tools* (version 1.0.1). Finally, the gene-level counts matrix was generated

using *featureCounts* from the *Subread* (version 2.0.1) package.

 Differential gene expression

Statistical analysis of the RNA-seq data was performed using *R* [52]. Since lowly expressed

genes are considered uninformative for differential expression analysis, we omitted genes

with less than 0.1 counts per million (CPM) (following the 10/minimum library size in millions

rule described in [53]). Library sizes after omitting the lowly expressed genes ranged

between 61 and 110 million reads. These were normalised using the trimmed mean of M-

values (TMM) method [54]. To test for differential expression of genes due to heterozygosity

for the T428del or W233fs mutation, we used a generalised linear model and likelihood ratio

tests using *edgeR* [55, 56]. A design matrix was specified with the wild type genotype as the

intercept, and the T428del/+ and W233fs/+ genotypes as the coefficients. We considered a

gene to be differentially expressed (DE) due to each *psen1* mutant genotype if the FDR

adjusted p-value was less than 0.05.

Enrichment analysis

We tested for over-representation of gene ontology (GO) terms within the DE gene lists

using *goseq* [57], using the average transcript length per gene to calculate the probability

weighting function (PWF). We considered a GO term to be significantly over-represented

within the DE gene lists relative to all detectable genes in the RNA-seq experiment if the

FDR-adjusted p-value generated by *goseq* was less than 0.05.

We also performed enrichment analysis on the entire list of detectable genes by calculating

the harmonic mean p-value from the raw p-values calculated from *fry* [58]*, camera* [59] and

fgsea [60, 61] as described in [41]. To test for changes to gene expression in a broad range of

biological processes, we used the KEGG [62] gene sets obtained from MSigDB [63] using the

msigdbr package [64]. We also used *msigdbr* to obtain gene sets which contain genes that

show changed expression in response to changes in the Notch signalling pathway

(*NGUYEN_NOTCH1_TARGETS_UP* and *NGUYEN_NOTCH1_TARGETS_DN*, *NOTCH_DN.V1_UP*,

NOTCH_DN.V1_DN and *RYAN_MANTLE_CELL_LYMPHOMA_NOTCH_DIRECT_UP*). We also

tested for evidence of possible iron dyshomeostasis using gene sets containing genes

encoding transcripts which contain iron-responsive elements in their untranslated regions

(described in [45]).

Comparison of the T428del and Q96_K97del mutations in psen1

Isolation of the zebrafish Q96_K97del mutation in zebrafish *psen1* and analysis of its effects

on zebrafish brain transcriptomes have been described previously [45, 46]. That dataset is

comprised of brain RNA-seq data for fish heterozygous for the Q96_K97del mutation and

their wild type siblings, at 6 months old (young adult) and 24 months old (aged), and under

normoxia or hypoxia treatment (n = 4 fish per genotype, age and treatment). In the analysis

presented here, we performed enrichment analysis using the methods described above on

the entire dataset, but presented the results for the pairwise comparison between 6-month-

old Q96_K97del/+ fish and wild type fish under normoxia.

To obtain a broader comparison on the effects of the Q96_K97del and T428del mutations,

we performed adaptive, elastic-net sparse PCA (AES-PCA) [65] as implemented in the

pathwayPCA package [66]. For this analysis, we utilised the HALLMARK [67] gene sets from

MSigDB to generate the pathway collection. The pathway principal components (PCs) were

calculated only on the gene expression data from samples heterozygous for an EOfAD-like

mutation (Q96_K97del or T428del) and their wild type siblings under normoxia at 6 months

of age. Then, the categorical effect of genotype was tested for association with the pathway

PCs using a permutation-based regression model as described in [66].

Data availability

The paired end fastq files and the output of *featureCounts* have been deposited in GEO

under accession number GSE164466. Code to reproduce this analysis can be found at

https://github.com/karissa-b/psen1_EOfAD_fAI_6m_RNA-seq.

RESULTS

Generation of an EOfAD-like and a fAI-like mutation in zebrafish psen1

An unsolved puzzle regarding the dominant EOfAD mutations of human *PSEN1* (and *PSEN2*)

is why these are consistently found to permit production of transcripts in which the reading

frame is preserved, while heterozygosity for mutations causing frameshifts (or deleting the

genes) does not cause EOfAD. To investigate this quandary in an *in vivo* model, we initially

aimed to generate mutations in zebrafish *psen1* which would be exact equivalents of the

T440del and P242fs mutations using homology directed repair (HDR). While screening for

the desired mutations, we identified the mutations W233fs and T428del, both likely

generated by the non-homologous end joining (NHEJ) pathway of DNA repair. T428del is a 3

nucleotide deletion which, nevertheless, produces a protein-level change exactly equivalent

to that observed for the human T440del mutation. Hereafter, for simplicity, we refer to the

T428del mutation as "EOfAD-like". W233fs is an indel mutation causing a frameshift in the

second codon downstream of the zebrafish *psen1* proline codon equivalent to human *PSEN1*

codon P242. This change is still within the short third lumenal loop of the Psen1 protein (see

Figure 1). Assuming no effect on splicing, the frameshift mutation results in a premature

stop codon 36 codons downstream of W233 (**Figure S1** in **Supplementary File 2**). Hereafter,

we refer to this mutation as "fAI-like". The alignment of the wild type and mutant sequences

in humans and zebrafish is shown in **Figure 2**.

Heterozygosity or homozygosity for neither the EOfAD-like nor the fAI-like mutation

produces any obvious morphological defects. However, this is unsurprising considering that

rare examples of humans homozygous for EOfAD mutations are known [68, 69] and loss of

PSEN1 γ-secretase activity is, apparently, compatible with viability in zebrafish [70] and rats

[71] (although not in mice [72]).

Figure 2: A Alignment of a region of wild type human (*Hs*) *PSEN1* exon 7, and the same region containing the human P242fs

(P242LfsX11) mutation, and the equivalent zebrafish (*Dr*) *psen1* exon 6 wild type and W233fs sequences. **B** Alignment of a

region of wild type human (*Hs*) *PSEN1* exon 12, and the same region containing the human T440del mutation, and the

equivalent zebrafish (*Dr*) *psen1* exon 11 wild type and T428del sequences.

Transcriptome analysis

To investigate global changes to the brain transcriptome due to heterozygosity for the

EOfAD-like or fAI-like mutations in *psen1,* we performed mRNA-seq on a family of fish as

described in **Figure 1**. The family of sibling fish were raised together in a single tank, thereby

reducing sources genetic and environmental variation between individuals and allowing

subtle changes to the transcriptome to be detected with minimal confounding effects.

To begin our exploration of the similarities and differences between the brain

transcriptomes of the mutant and wild type fish, we first performed principal component

analysis (PCA) on the gene level, log transformed counts per million (logCPM) of the

zebrafish RNA-seq samples. A plot of principal component 1 (PC1) against PC2 did not show

distinct clustering of samples by genotype or sex, indicating that these variables do not

result in stark changes to the brain transcriptome. This is consistent with our previous

observations of EOfAD-like mutations in other genes [39-41]. However, some separation of

the EOfAD-like and fAI-like samples is observed across PC2, indicating distinct, but subtle,

differences between these transcriptomes. Notably, the majority of the variation in this

dataset is not captured until PC6 (**Figure 3**).

Heterozygosity for the EOfAD-like or fAI-like mutations of psen1 causes only subtle effects on

gene expression

Which genes are dysregulated due to heterozygosity for the EOfAD-like or the fAI-like

mutations? To address this question, we performed differential gene expression analysis

using a generalised linear model and likelihood ratio tests with *edgeR*. We observed

statistical evidence for 13 genes as significantly differentially expressed (DE) due to

heterozygosity for the EOfAD-like mutation, and 5 genes due to the fAI-like mutation (**Figure
4**, **Supplementary Table 1**). Notably, *psen1* was the most significantly DE gene due to

heterozygosity for the fAI-like mutation (logFC = -0.8, FDR = 1.33e-78), consistent with the

observation that frame-shift mutations commonly induce nonsense-mediated mRNA decay

when they result in premature stop codons (reviewed in [73]). Total levels of *psen1*

transcripts were unchanged in EOfAD-like/+ brains (logFC = -0.0065, FDR = 1, **Figure S7** in

Supplementary File 2). No DE genes were found to be shared between the comparisons of

either form of heterozygous mutant to wild type, or were found to be significantly

overrepresented by any gene ontology (GO) terms by *goseq* (for the top 5 most significantly

over-represented GO terms in each comparison, see **Tables S1** and **S2** in **Supplementary File**

2). This is not unexpected due to the relatively low number of significantly DE genes

detected.

Figure 4: Differential expression analysis. **A** Mean-difference (MD) plots and **B** volcano plots of changes to gene expression in EOfAD-like/+ and fAI-like/+ mutant zebrafish brains. Note that the limits of the y-axis in **B** are restrained to between 0 and 10 for visualisation purposes. **C** Upset plot indicating the low number of genes which are significantly differentially expressed (DE) in either comparison.

 Significant differences in gene expression between the EOfAD-like and fAI-like mutants can be

detected at the pathway level.

Since very few DE genes were detected in each comparison of heterozygous *psen1* mutant

fish to their wild type siblings, we performed enrichment analysis on all detectable genes in

the RNA-seq experiment. Our method, inspired by the *EGSEA* framework [74], involves

calculation of the harmonic mean p-value [75] from the raw p-values of three different rank-

based gene set testing methods: *fry* [58], *camera* [59] and *GSEA* [60, 61]. Unlike *EGSEA*, we

use the harmonic mean p-value to combine the raw p-values, as the harmonic mean p-value

has been specifically shown to be robust for combining dependent p-values [75]. We

performed enrichment testing using the KEGG gene sets (describing 186 biological pathways

and processes) to obtain information on changes to activities for these pathways. We also

tested for evidence of iron dyshomeostasis using our recently defined sets of genes

containing iron-responsive elements (IREs) in the untranslated regions of their mRNAs [45].

We observed statistical evidence for 7 KEGG gene sets as significantly altered by

heterozygosity for the EOfAD-like mutation and 11 KEGG gene sets as significantly altered by

heterozygosity for the fAI-like mutation (**Figure 5,** full results of the raw p-values from each

algorithm as well as the harmonic mean p-value can be found in **Supplementary Table 2**).

Gene sets significantly altered in the brains of both forms of heterozygous mutant included

the KEGG gene sets for cytokine receptor interactions, Jak/Stat signalling, and encoding the

components of the ribosomal subunits. Inspection of the leading edge genes (which can be

interpreted as the core genes driving the enrichment of a gene set) showed that similar

genes were driving the enrichment of the gene sets for cytokine receptor interactions and

Jak/Stat signalling. Similar genes were also driving the enrichment of the *KEGG_RIBOSOME*

gene set in both heterozygous mutants. However, the magnitude of the logFC was greater in

the fAI-like/+ samples, suggesting a stronger effect (**Figure S4-S6** in **Supplementary File 2**).

Gene sets which were only altered significantly by heterozygosity for the EOfAD-like

mutation were involved in energy metabolism (*KEGG_PARKINSONS_DISEASE*,

KEGG_OXIDATIVE_PHOSPHORYLATION, *and KEGG_CITRATE_CYCLE_TCA_CYCLE*). Notably,

the KEGG gene sets for Parkinson's disease, Huntington's disease, and for oxidative

phosphorylation, share 55 leading-edge genes, implying that their enrichment is driven by,

essentially, the same gene expression signal (**Figure 5**). Conversely, the 10 KEGG gene sets

found to be altered significantly by heterozygosity for the fAI-like mutation appear to be

driven mostly by distinct gene expression signals. No IRE gene sets were observed

statistically to be altered in the brains of either mutant, suggesting that iron homeostasis is

unaffected (at least at 6 months of age). The changes to expression of genes within the KEGG

gene sets are likely not due to broad changes in cell-type proportions in the zebrafish brain

samples, since the expression of marker genes of neurons, astrocytes, oligodendrocytes and

microglia was similar in all samples (**Figure S13** in **Supplementary File 2**).

Figure 5: A KEGG gene sets with FDR-adjusted harmonic mean p-values of < 0.05 in *psen1* EOfAD-like/+ and fAI-like/+ mutant

brains. The colour of the cells indicates the level of significance (brighter colour indicates greater statistical significance, while

dark grey indicates the FDR-adjusted harmonic mean p-value > 0.05). The number within each cell is the FDR-adjusted

harmonic mean p-value. **B** Upset plot indicating the overlap of leading edge genes from the fgsea algorithm which drive the

enrichment of gene sets significantly altered in EOfAD-like/+ and **C** fAI-like/+ brains.

The EOfAD-like and fAI-like mutations alter expression of Notch signalling genes

Notch signalling plays a critical role in many cell differentiation events and is dependent on

PSEN's γ-secretase activity. Disturbance of Notch signalling due to decreased γ-secretase

activity has been suggested to contribute to the changes in skin histology of fAI, as Notch

signalling is required for normal epidermal maintenance ([76-78] and reviewed by [79]).

However, fAI has not been reported as associated with EOfAD, despite that the T440del

mutant form of *PSEN1* appears to have little intrinsic γ-secretase activity [15]. The expression

of genes involved in the KEGG gene set for the Notch signalling pathway was observed to be

highly significantly altered in the brains of fAI-like/+ mutants, but not of EOfAD/+ mutants,

implying that γ-secretase activity might only be affected significantly by the frameshift, fAI-

like mutation (**Figure 5**). However, inspection of the logFC of genes in the

KEGG_NOTCH_SIGNALING_PATHWAY gene set revealed similar patterns of changes to gene

expression in both mutants (**Figure 6**). Upregulation of the genes encoding the Notch and

Delta receptors is observed in both mutants compared to wild type. In fAI-like/+ brains, we

observe downregulation of the downstream transcriptional targets of the Notch intracellular

domain (NICD), implying decreased Notch signalling (and, likely, reduced γ-secretase

activity). Genes encoding repressors of Notch signalling are observed to be upregulated (i.e.

dyl and *numb*), reinforcing this interpretation.

Since the KEGG gene set for Notch signalling only contains two genes that are direct

transcriptional targets of the NICD, we investigated further whether Notch signalling is

perturbed in both mutants by analysis of gene sets from MSigDB containing information on

genes responsive to Notch signalling in different cell lines: *NGUYEN_NOTCH1_TARGETS_UP*;

NGUYEN_NOTCH1_TARGETS_DN; *NOTCH_DN.V1_UP*; *NOTCH_DN.V1_DN*; and

RYAN_MANTLE_CELL_LYMPHOMA_NOTCH_DIRECT_UP. The

NGUYEN_NOTCH1_TARGETS_UP and *_DOWN* gene sets consist of genes which have been

observed as up- or downregulated respectively in response to a constitutively active Notch

receptor in keratinocytes [80]. *The NOTCH_DN.V1_UP* and *_DN* gene sets contain genes

which are up- and down-regulated respectively in response to treatment with the γ-

secretase inhibitor DAPT in a T-cell acute lymphoblastic leukemia (T-ALL) cell line [81]. The

RYAN_MANTLE_CELL_LYMPHOMA_NOTCH_DIRECT_UP gene set contains genes showing

both increased expression upon rapid activation of Notch signalling by washout of the γ-

secretase inhibitor compound E, and evidence for a NICD binding site in the promotor by

ChIP-seq, in mantle cell lymphoma cell lines [82]. (Note that there is no equivalent

"*RYAN_MANTLE_CELL_LYMPHOMA*" gene set representing genes downregulated in

response to Notch signalling.) Of these 5 gene sets, statistical support was found only for

changes to the expression of genes in the

RYAN_MANTLE_CELL_LYMPHOMA_NOTCH_DIRECT_UP gene set, and this was found for

both the EOfAD-like (p=0.006) and the fAI-like (p=0.008) mutants. The leading edge genes

were mostly observed to be upregulated, which supports increases in Notch signalling

(implying increased γ-secretase activity). Transcriptional adaptation (previously known as

"genetic compensation") might contribute to the apparent increase in Notch signalling in the

frameshift, fAI-like/+ mutant brains via upregulated expression of the *psen1*-homologous

gene, *psen2* [83, 84]. Although no statistically significant differences in expression were

observed for *psen2* in the differential expression test using *edgeR* (see **Supplementary Table**

1), a trend towards upregulation in the fAI-like/+ mutants was observed following a simple

Student's t-test (p=0.074, **Figure 6D**). El-Brolosy et al. [83] showed that the wild type allele of

a mutated gene can also be upregulated by transcriptional adaptation (where the mutation

causes nonsense-medicated decay, NMD, of mutant transcripts). Inspection of the number

of reads aligning to the W233 mutation site across samples indicates that the expression of

the wild type *psen1* allele in fAI-like/+ brains appears to be greater than 50% of the

expression of the wild type *psen1* allele in wild type brains (p = 0.006), providing further

evidence for transcriptional adaptation due to the fAI-like mutation (**Figure S8** in

Supplementary File 2).

Together, these results suggest that Notch signalling and, by implication, γ-secretase activity,

may be enhanced in *psen1* mutant brains. However, future biochemical assays should be

performed to confirm this prediction.

Figure 6: A Pathview [85] visualisation of the changes to gene expression in the *KEGG_NOTCH_SIGNALING_PATHWAY*

gene set in EOfAD-like/+ mutants and **B** fAI-like/+ mutants. **C** The proportion of genes with increased expression (red, z > √2)

and decreased expression (blue, z < √2) in MSigDB gene sets for Notch signalling in EOfAD-like/+ and fAI-like/+ mutant brains.

Gene sets which contained a FDR-adjusted harmonic mean p-value (HMP) < 0.05 appear less transparent. The FDR adjust p-

values are also listed on the bars. **D** The expression of *psen2* is trending towards upregulation, particularly in fAI-like/+ mutants.

Here, p-values were determined by Student's unpaired t-tests. **E** Heatmap indicating the logFC values for genes in the

RYAN_MANTLE_CELL_LYMPHOMA_NOTCH_DIRECT_UP gene set. Genes are clustered based on their Euclidean distance,

and are labelled with green if they appear in the leading edge of the *fgsea* algorithm for each comparison of a *psen1*

heterozygous mutant with wild type.

The EOfAD-like mutation T428del has a milder phenotype than the previously studied

Q96_K97del EOfAD-like mutation of psen1

The T428del mutation of *psen1* is the first identified zebrafish mutation exactly equivalent,

(at the protein sequence level), to a characterised human EOfAD mutation. Therefore, we

sought to assess the consistency of its effects with those of a previously studied EOfAD-like

mutation, Q96_K97del, and to identify cellular processes affected in common by the two

mutations. The Q96_K97del mutation deletes two codons in the sequence encoding the first

lumenal loop of the Psen1 protein (see **Figure 1**). Comparison of transcriptomes from the 6-

month-old brains of Q96_K97del/+ and wild type siblings previously predicted changes to

expression of genes involved in energy metabolism, iron homeostasis and lysosomal

acidification [45, 46]. To compare which cellular processes are affected by heterozygosity for

the Q96_K97del mutation or the T428del mutation, we first performed enrichment analysis

on the RNA-seq data previously generated by our analysis of zebrafish heterozygous for the

Q96_K97del mutation relative to their wild type siblings. Here, we observed that

heterozygosity for the Q96_K97del mutation results in significant alterations in 7 KEGG gene

sets (at 6 months of age during normoxia, Figure 7A). We also found statistical evidence for

altered expression of genes possessing IREs in their 3' UTRs (see *IRE3_ALL* in **Figure 7A**),

consistent with our previous finding using a different method of enrichment analysis [45].

Gene sets affected in common between the two EOfAD-like mutations in *psen1* are involved

in energy metabolism and protein translation (**Figure 7A**). The expression of genes involved

in protein degradation, and of genes containing IREs in the 3' UTRs of their transcripts,

appeared significantly altered only by the Q96_K97del mutation (**Figure 7A**).

We also compared the effects of the two EOfAD-like mutations using adaptive, elastic-net

sparse PCA (AES-PCA) as implemented in the *pathwayPCA* package [66]. AES-PCA allows

reduction of data dimensionality and for the overall activity of predefined gene sets to be

observed in a sample-specific manner [65]. To obtain a global view of the changes to gene

expression between the two *psen1* EOfAD-like mutations over the two datasets, we utilised

the HALLMARK gene sets that encompass 50 distinct biological processes (rather than the

186 KEGG gene sets that share many genes).

The latent variables estimated by AES-PCA for the HALLMARK gene sets (i.e. the first

principal components) in each dataset did not show any significant association with *psen1*

genotype, suggesting that changes to gene expression (measured over entire brains) are too

subtle to be detected as statistically significant using this method. However, clustering of the

calculated PC1 values by AES-PCA for each HALLMARK gene set in each sample and dataset

revealed that samples in the Q96_K97del dataset clustered mostly according to genotype

(one wild type sample did not follow the trend), supporting that heterozygosity for the

Q96_K97del mutation does result in marked effects on gene expression for the HALLMARK

gene sets. Conversely, clustering of PC1 values in the T428del dataset resulted in two distinct

clusters of samples. However, samples did not group by genotype over the two clusters to

the same extent as seen for the Q96_K97del dataset. Intriguingly, the Q96_K97del dataset

had less sample numbers per genotype (n = 4), and did not have as great sequencing depth

as the current RNA-seq experiment. Therefore, this supports that heterozygosity for the

Q96_K97del mutation has more consistent (more severe) effects on young adult brain

transcriptomes than heterozygosity for the T428del mutation (**Figure 7 B,C**).

 Figure 7: A Comparison of KEGG and IRE gene sets significantly altered by the EOfAD-like mutations T428del and

Q96_K97del in 6-month-old zebrafish brains. Each cell is coloured according to statistical significance, and the FDR-adjusted

harmonic mean p-value is shown. Gene sets not significantly altered (FDR adjusted harmonic mean p-value > 0.05) in a

comparison between a *psen1* mutant zebrafish with their respective wild type siblings appear grey. **B** Principal component 1

(PC1) values for the HALLMARK gene sets as calculated by AES-PCA, clustered based on their Euclidean distance in

T428del/+ samples relative to their wild type siblings. **C** PC1 values for the HALLMARK gene sets as calculated by AES-PCA

clustered based on their Euclidean distance in Q96_K97del/+ samples relative to their wild type siblings at 6 months of age

under normal oxygen conditions.

DISCUSSION

In this study, we exploited transcriptome analysis of whole brains of young adult zebrafish

siblings, to detect differences in molecular state between the brains of fish heterozygous for

an EOfAD-like mutation or an fAI-like mutation of *psen1* compared to wild type *in vivo*. The

subtlety of the effects observed is consistent with that EOfAD is, despite its designation as

"early-onset", a disease affecting people overwhelmingly at ages older than 30 years [86].

The person reported to carry the T440del mutation of *PSEN1* showed cognitive decline at 41

years [47]. (Overall, EOfAD mutations in *PSEN1* show a median survival to onset of 45 years

[86]). In contrast, at 6 months of age, zebrafish are only recently sexually mature.

Nevertheless, since AD is thought to take decades to develop [32], it is these subtle, early

changes that we must target therapeutically if we wish to arrest the pathological processes

driving the progression to AD. As seen in all our previous analyses of EOfAD-like mutations

[39-41, 43, 45], changes in expression of genes involved in oxidative phosphorylation were

identified as significant. However, this was not the case for the fAI-like, frameshift mutation.

Therefore, oxidative phosphorylation changes appear to be an early signature cellular stress

of EOfAD. Changes to mitochondrial function have been observed in heterozygous *PSEN1*

mutant astrocytes [87], homozygous *PSEN1* mutant neurons [88], and in neurons

differentiated from human induced pluripotent stem cells (hIPSCs) from sporadic AD

patients [89], supporting our findings. However, such changes are not always observed [90],

possibly due to issues of experimental reproducibility between laboratories when working

with hIPSCs [91].

The EOfAD-like mutation also caused very statistically significant changes in the

KEGG_PARKINSONS_DISEASE gene set (that shares many genes with

KEGG_OXIDATIVE_PHOSPHORYLATION) and the person carrying the *PSEN1^{T440del}* mutation

modelled by zebrafish *psen1^{T428del}* initially showed symptoms of early onset parkinsonism at

34 years of age before those of cognitive decline at 41 years [47].

In contrast to this EOfAD-like mutation, the fAI-like mutation apparently caused very

statistically significant changes in Notch signalling and changes in other signal transduction

pathways such as those involving Wnt and neurotrophins, as might be expected from

changes in γ-secretase activity. Also notable was enrichment for the

KEGG_TOLL_LIKE_RECEPTOR_SIGNALLING_PATHWAY gene set since acne inversa is a chronic

inflammatory skin disorder and, in humans, increased expression of Toll-like receptor 2 has

been noted in acne inversa lesions [92].

Both the EOfAD-like and fAI-like mutations caused very statistically significant changes in the

gene sets *KEGG_CYTOKINE_CYTOKINE_RECEPTOR _INTERACTION* and *KEGG_RIBOSOME*. The

former gene set reflects that both mutations appear to affect inflammation that is a

characteristic of the pathologies of both EOfAD [93] and fAI (reviewed in [94]). Like oxidative

phosphorylation, we have also observed effects on ribosomal protein genes sets for every

EOfAD-like mutation we have studied [39-41, 43]. This may be due protein synthesis

consuming a large proportion of cells' energy budgets [95] and requiring amino acid

precursors that can be sourced from lysosomes. Recently, Bordi et al [96] noted that mTOR

is highly activated in fibroblasts from people with Down syndrome (DS, trisomy 21). DS

individuals commonly develop EOfAD due to overexpression of the AβPP gene (that is

resident on human chromosome 21). The consequent increased expression of AβPP's β-

CTF/C99 fragment (generated by β-secretase cleavage of AβPP without γ-secretase cleavage)

affects endolysosomal pathway acidification [97] in a similar manner to EOfAD mutations of

PSEN1 [98]. The mTOR protein is localised at lysosomes in the mTORC1 and mTORC2 protein

complexes (reviewed in [99, 100]) and monitors the energy and nutrient status of cells

(reviewed in [101]). It is important for regulating ribosomal activity, partly by regulating

transcription of ribosome components (reviewed in [102, 103]). Therefore, one explanation

for the consistent enrichment for transcripts of the *KEGG_RIBOSOME* gene set we see in

EOfAD mutant brains may be mTOR activation due to effects on lysosomal acidification

and/or the energy status of cells. We did not observe any significant changes to the

expression of genes involved in the mTOR signalling pathway in the analyses described in

this paper (the FDR-adjusted harmonic mean p-value for the

KEGG_MTOR_SIGNALING_PATHWAY was 0.7 for each comparison of the *psen1* mutant fish

to their wild type siblings, see **Supplementary Table 2**). However, these changes could be

undetectably subtle in young adult brains and/or occurring at the protein level and therefore

not observable in bulk RNA-seq data. (Statistically significant enrichment for genes in the

HALLMARK_PI3K_AKT_MTOR_SIGNALING gene set was seen previously for the normoxic, 6-

month-old brains of fish heterozygous for the more severe EOfAD-like mutation Q96_K97del

when compared to wild type siblings [45].)

While the brain transcriptome alterations caused by the EOfAD-like and fAI-like mutations

are subtle (as illustrated by the lack of tight clustering of samples in the principal component

analysis in **Figure 3,** and the low number of significantly differentially expressed genes), we

are reassured in their overall veracity by their similarity to the results of a parallel analysis of

sibling brain transcriptomes from 6-month-old zebrafish heterozygous for either a frameshift

or a frame-preserving mutation in the zebrafish *psen2* gene relative to wild type [40]. In that

similarly structured (but less statistically powered) experiment, only the frameshift mutation

significantly affected the *KEGG_NOTCH_SIGNALLING* gene set while only the frame-

preserving, EOfAD-like mutation significantly affected the

KEGG_OXIDATIVE_PHOSPHORYLATION gene set. Both *psen2* genotypes affected the

KEGG_RIBOSOME gene set, but in overall opposite directions (the frameshift mutation

largely upregulated these genes while the frame-preserving mutation did the opposite).

Transcriptome analysis can reveal a great deal of data on differences in gene transcript levels

between different genotypes or treatments. However, interpreting changes in cellular state

from this information is not straight forward. Are any changes seen direct molecular effects

of a mutation/treatment (e.g. the direct, downstream effects of a change in γ-secretase

activity) or homeostatic responses as cells/tissues adjust their internal states to promote

survival? For example, in the *KEGG_NOTCH_SIGNALING_PATHWAY* gene set shown in **Figure**

6B, more pathway components are upregulated than are downregulated. However, the

direct transcriptional targets of Notch signalling (*her4.2* and *heyl*) are downregulated, as

might be expected from reduced expression of wild type, catalytically-competent Psen1

protein. The upregulation of other components of the pathway may represent homeostatic

responses attempting to restore normal levels of Notch signalling.

Only two Notch downstream transcriptional target genes are described in the

KEGG_NOTCH_SIGNALING_PATHWAY gene set. Therefore, in an effort to assess more

generally the effects of the EOfAD-like and fAI-like mutations on γ-secretase activity, we also

analysed additional sets of genes previously identified (in various systems) as direct

transcriptional targets of γ-secretase-dependent signalling. One of these sets, encompassing

genes identified as Notch signalling targets by both γ-secretase inhibitor responses and

binding of the Notch intracellular domain to chromatin, revealed apparent upregulation of

Notch signalling in both the EOfAD-like and fAI-like heterozygous mutant brains relative to

wild type siblings. In both EOfAD-like/+ and fAI-like/+ mutant brains, the most highly ranked

genes in terms of differential expression tended to be upregulated (although some highly

ranked genes were downregulated in fAI-like/+ brains). The idea that putative low levels of a

form of Psen1 protein truncated in the third lumenal loop domain could increase Notch

signalling is not unexpected, as we previously observed an implied upregulation of Notch

signalling in zebrafish embryos with forced expression of the fAI-causative P242Lfs allele of

human *PSEN1* [104]. However, a widespread assumption within AD research is that EOfAD-

like mutations of *PSEN1* decrease γ-secretase cleavage of AβPP [105], possibly through a

dominant negative mechanism [21]. This assumption conflicts with the observation of Sun et

al. [15] that approximately 10% of the 138 EOfAD mutations of human *PSEN1* they studied

actually increased γ-secretase cleavage of AβPP's β-CTF/C99 fragment (in experiments

examining the activities of the mutant proteins in isolation from wild type protein). Zhou et

al. [106] also observed increased γ-secretase activity (cleavage of AβPP's β-CTF/C99

fragment) due to an EOfAD mutation of *PSEN1* (S365A, this replicated Sun et al.'s finding for

this mutation).

It is important to note that mutations of *PSEN1* need not cause similar effects on Notch and

AβPP cleavage [104, 107, 108]. The transmembrane domains of the Notch receptor and the

APP's C99 fragment have different conformations [109]. Therefore, changes in the

conformation of PSEN1 within γ-secretase due to a mutation may differentially affect Notch

and C99 cleavage. Indeed, in our previously mentioned study of forced expression of the

human *PSEN1^{P242Lfs}* allele in zebrafish embryos, increased apparent Notch signalling was

observed without change in AβPP processing [104]. Conversely, Zhang et al. [108] showed

that transgenic expression of an EOfAD mutation S169del in *PSEN1* under the control of the

Thy1 brain specific promotor altered the processing of AβPP *in vivo* without affecting Notch

signalling. Notably, both of these studies did not use the *PSEN1* gene's own promoter to

express mutant forms of this gene, and so the effects seen may be distorted by gene/protein

over-expression.

Unfortunately, the direct transcriptional targets of the intracellular domain of AβPP (AICD)

have not been characterised to the same extent as those of NICD (reviewed in [110]). This

constrains transcriptome analysis for detection of differential effects on γ-secretase cleavage

of AβPP caused by the EOfAD-like and fAI-like mutations. Future work should include further

investigation of how these mutations effect γ-secretase cleavage of AβPP *in vivo.*

In conclusion, we have performed the first direct comparison of an EOfAD-like and a fAI-like

mutation of *presenilin 1* in an *in vivo* model. Both forms of mutation cause apparent changes

in inflammation, downregulate expression of genes encoding the components of the

ribosome subunits, and potentially affect γ-secretase activity as supported by altered

expression of Notch signalling pathway transcriptional target genes. We see that changes to

mitochondrial function are a specific, common characteristic of EOfAD-like mutations while

the fAI-like mutation specifically affects important signal transduction pathways. These

differential effects on brain transcriptomes give insight into how reading-frame preserving

mutations in *PSEN1* cause EOfAD while frameshift mutations do not.

CONFLICT OF INTEREST/DISCLOSURE STATEMENT

The authors have no financial or non-financial competing interests to declare.

ACKNOWLEDGEMENTS

The authors would like to thank Dr. Nhi Hin for providing the Q96_K97del gene expression

values and the zebrafish IRE gene sets. We also would like to thank Dr. Jimmy Breen for his

assistance with using the Nextflow pipeline. The authors thank Dr Giuseppe Verdile for

critical reading of the manuscript.

This work was supported with supercomputing resources provided by the Phoenix HPC

service at the University of Adelaide and by grants GNT1061006 and GNT1126422 from the

National Health and Medical Research Council of Australia (NHMRC). KB was supported by

an Australian Government Research Training Program Scholarship and by funds from the

Carthew Family Charity Trust. YD was supported by an Adelaide Graduate Research

Scholarship from the University of Adelaide. MN was supported by funds from the grants

listed above. ML is an academic employee of the University of Adelaide.

REFERENCES

[1]   Janssen JC, Beck JA, Campbell TA, Dickinson A, Fox NC, Harvey RJ, Houlden H, Rossor
652   MN, Collinge J (2003) Early onset familial Alzheimer's disease: Mutation frequency in
653   31 families. *Neurology* **60**, 235-239.

654   [2]   Żekanowski C, Styczyńska M, Pepłońska B, Gabryelewicz T, Religa D, Ilkowski J,
655   Kijanowska-Haładyna B, Kotapka-Minc S, Mikkelsen S, Pfeffer A, Barczak A, Łuczywek
656   E, Wasiak B, Chodakowska-Żebrowska M, Gustaw K, Łączkowski J, Sobów T, Kuźnicki
657   J, Barcikowska M (2003) Mutations in presenilin 1, presenilin 2 and amyloid precursor
658   protein genes in patients with early-onset Alzheimer's disease in Poland.
659   *Experimental Neurology* **184**, 991-996.

660   [3]   Cruts M, van Duijn CM, Backhovens H, Van den Broeck M, Wehnert A, Serneels S,
661   Sherrington R, Hutton M, Hardy J, St George-Hyslop PH (1998) Estimation of the
662   genetic contribution of presenilin-1 and-2 mutations in a population-based study of
663   presenile Alzheimer disease. *Human molecular genetics* **7**, 43-51.

664   [4]   Kamimura K, Tanahashi H, Yamanaka H, Takahashi K, Asada T, Tabira T (1998) Familial
665   Alzheimer's disease genes in Japanese. *Journal of the neurological sciences* **160**, 76-
666   81.

667   [5]   Area-Gomez E, de Groof AJ, Boldogh I, Bird TD, Gibson GE, Koehler CM, Yu WH, Duff
668   KE, Yaffe MP, Pon LA, Schon EA (2009) Presenilins are enriched in endoplasmic
669   reticulum membranes associated with mitochondria. *American Journal of Pathology*
670   **175**, 1810-1816.

671   [6]   Schon EA, Area-Gomez E (2010) Is Alzheimer's disease a disorder of mitochondria-
672   associated membranes? *J Alzheimers Dis* **20 Suppl 2**, S281-292.

673   [7]   Laudon H, Hansson EM, Melén K, Bergman A, Farmery MR, Winblad B, Lendahl U, von
674   Heijne G, Näslund J (2005) A nine-transmembrane domain topology for presenilin 1. *J
675   Biol Chem* **280**, 35352-35360.

676   [8]   Fukumori A, Fluhrer R, Steiner H, Haass C (2010) Three-amino acid spacing of
677   presenilin endoproteolysis suggests a general stepwise cleavage of gamma-secretase-
678   mediated intramembrane proteolysis. *J Neurosci* **30**, 7853-7862.

[9]   Sun L, Zhao L, Yang G, Yan C, Zhou R, Zhou X, Xie T, Zhao Y, Wu S, Li X, Shi Y (2015)
Structural basis of human γ-secretase assembly. *Proceedings of the National*
Academy of Sciences **112**, 6003.

[10]   Wolfe MS (2020) Unraveling the complexity of γ-secretase. *Semin Cell Dev Biol* **105**,
3-11.

[11]   Dermaut B, Kumar-Singh S, Engelborghs S, Theuns J, Rademakers R, Saerens J, Pickut
BA, Peeters K, van den Broeck M, Vennekens K, Claes S, Cruts M, Cras P, Martin JJ,
Van Broeckhoven C, De Deyn PP (2004) A novel presenilin 1 mutation associated with
Pick's disease but not beta-amyloid plaques. *Ann Neurol* **55**, 617-626.

[12]   Li D, Parks SB, Kushner JD, Nauman D, Burgess D, Ludwigsen S, Partain J, Nixon RR,
Allen CN, Irwin RP, Jakobs PM, Litt M, Hershberger RE (2006) Mutations of presenilin
genes in dilated cardiomyopathy and heart failure. *Am J Hum Genet* **79**, 1030-1039.

[13]   Wang B, Yang W, Wen W, Sun J, Su B, Liu B, Ma D, Lv D, Wen Y, Qu T, Chen M, Sun M,
Shen Y, Zhang X (2010) -Secretase Gene Mutations in Familial Acne Inversa. **330**,
1065-1065.

[14]   Jiang H, Jayadev S, Lardelli M, Newman M (2018) A Review of the Familial Alzheimer's
Disease Locus PRESENILIN 2 and Its Relationship to PRESENILIN 1. *Journal of*
Alzheimer's Disease **66**, 1323-1339.

[15]   Sun L, Zhou R, Yang G, Shi Y (2017) Analysis of 138 pathogenic mutations in
presenilin-1 on the in vitro production of Aβ42 and Aβ40 peptides by γ-secretase.
Proceedings of the National Academy of Sciences **114**, E476-E485.

[16]   Szaruga M, Veugelen S, Benurwar M, Lismont S, Sepulveda-Falla D, Lleo A, Ryan NS,
Lashley T, Fox NC, Murayama S, Gijsen H, De Strooper B, Chávez-Gutiérrez L (2015)
Qualitative changes in human γ-secretase underlie familial Alzheimer's disease. *J Exp*
Med **212**, 2003-2013.

[17]   De Strooper B (2007) Loss-of-function presenilin mutations in Alzheimer disease.
Talking Point on the role of presenilin mutations in Alzheimer disease. *EMBO Rep* **8**,
141-146.

[18]   Jayne T, Newman M, Verdile G, Sutherland G, Munch G, Musgrave I, Moussavi Nik
SH, Lardelli M (2016) Evidence for and Against a Pathogenic Role of Reduced gamma-
Secretase Activity in Familial Alzheimer's Disease. *Journal of Alzheimer's Disease*.

[19]   Wong TH, Seelaar H, Melhem S, Rozemuller AJM, van Swieten JC (2020) Genetic
711   screening in early-onset Alzheimer's disease identified three novel presenilin
712   mutations. *Neurobiology of aging* **86**, 201.e209-201.e214.

713   [20]   Lee JH, McBrayer MK, Wolfe DM, Haslett LJ, Kumar A, Sato Y, Lie PP, Mohan P, Coffey
714   EE, Kompella U, Mitchell CH, Lloyd-Evans E, Nixon RA (2015) Presenilin 1 Maintains
715   Lysosomal Ca(2+) Homeostasis via TRPML1 by Regulating vATPase-Mediated
716   Lysosome Acidification. *Cell Report* **12**, 1430-1444.

717   [21]   Heilig EA, Gutti U, Tai T, Shen J, Kelleher RJ, 3rd (2013) Trans-dominant negative
718   effects of pathogenic PSEN1 mutations on γ-secretase activity and Aβ production. *J
719   Neurosci* **33**, 11606-11617.

720   [22]   Schroeter EH, Ilagan MXG, Brunkan AL, Hecimovic S, Li Y-m, Xu M, Lewis HD, Saxena
721   MT, De Strooper B, Coonrod A, Tomita T, Iwatsubo T, Moore CL, Goate A, Wolfe MS,
722   Shearman M, Kopan R (2003) A presenilin dimer at the core of the γ-secretase
723   enzyme: Insights from parallel analysis of Notch 1 and APP proteolysis. *Proceedings
724   of the National Academy of Sciences* **100**, 13075.

725   [23]   Brautigam H, Moreno CL, Steele JW, Bogush A, Dickstein DL, Kwok JBJ, Schofield PR,
726   Thinakaran G, Mathews PM, Hof PR, Gandy S, Ehrlich ME (2015) Physiologically
727   generated presenilin 1 lacking exon 8 fails to rescue brain PS1−/− phenotype and
728   forms complexes with wildtype PS1 and nicastrin. *Scientific Reports* **5**, 17042.

729   [24]   Li X, Dang S, Yan C, Gong X, Wang J, Shi Y (2013) Structure of a presenilin family
730   intramembrane aspartate protease. *Nature* **493**, 56-61.

731   [25]   De Gasperi R, Sosa MAG, Dracheva S, Elder GA (2010) Presenilin-1 regulates induction
732   of hypoxia inducible factor-1α: altered activation by a mutation associated with
733   familial Alzheimer's disease. *Molecular Neurodegeneration* **5**, 38-38.

734   [26]   Villa JC, Chiu D, Brandes AH, Escorcia FE, Villa CH, Maguire WF, Hu CJ, de Stanchina E,
735   Simon MC, Sisodia SS, Scheinberg DA, Li YM (2014) Nontranscriptional role of Hif-
736   1alpha in activation of gamma-secretase and notch signaling in breast cancer. *Cell
737   Replication* **8**, 1077-1092.

738   [27]   Newman M, Nik HM, Sutherland GT, Hin N, Kim WS, Halliday GM, Jayadev S, Smith C,
739   Laird AS, Lucas CW, Kittipassorn T, Peet DJ, Lardelli M (2020) Accelerated loss of
740   hypoxia response in zebrafish with familial Alzheimer's disease-like mutation of
741   presenilin 1. *Human Molecular Genetics* **29**, 2379-2394.

742   [28]   Lane DJR, Merlot AM, Huang MLH, Bae DH, Jansson PJ, Sahni S, Kalinowski DS,
743   Richardson DR (2015) Cellular iron uptake, trafficking and metabolism: Key molecules
744   and mechanisms and their roles in disease. *Biochimica et Biophysica Acta (BBA) -*
745   *Molecular Cell Research* **1853**, 1130-1144.

746   [29]   Güner G, Lichtenthaler SF (2020) The substrate repertoire of γ-secretase/presenilin.
747   *Seminars in Cell & Developmental Biology* **105**, 27-42.

748   [30]   Thinakaran G, Harris CL, Ratovitski T, Davenport F, Slunt HH, Price DL, Borchelt DR,
749   Sisodia SS (1997) Evidence that levels of presenilins (PS1 and PS2) are coordinately
750   regulated by competition for limiting cellular factors. *J Biol Chem* **272**, 28415-28422.

751   [31]   Berchtold NC, Sabbagh MN, Beach TG, Kim RC, Cribbs DH, Cotman CW (2014) Brain
752   gene expression patterns differentiate mild cognitive impairment from normal aged
753   and Alzheimer's disease. *Neurobiology of Aging* **35**, 1961-1972.

754   [32]   Villemagne VL, Burnham S, Bourgeat P, Brown B, Ellis KA, Salvado O, Szoeke C,
755   Macaulay SL, Martins R, Maruff P, Ames D, Rowe CC, Masters CL (2013) Amyloid β
756   deposition, neurodegeneration, and cognitive decline in sporadic Alzheimer's
757   disease: a prospective cohort study. *The Lancet Neurology* **12**, 357-367.

758   [33]   Foley AM, Ammar ZM, Lee RH, Mitchell CS (2015) Systematic review of the
759   relationship between amyloid-β levels and measures of transgenic mouse cognitive
760   deficit in Alzheimer's disease. *J Alzheimers Dis* **44**, 787-795.

761   [34]   Hargis KE, Blalock EM (2017) Transcriptional signatures of brain aging and
762   Alzheimer's disease: What are our rodent models telling us? *Behavioural Brain*
763   *Research* **322**, 311-328.

764   [35]   Guo Q, Fu W, Sopher BL, Miller MW, Ware CB, Martin GM, Mattson MP (1999)
765   Increased vulnerability of hippocampal neurons to excitotoxic necrosis in presenilin-1
766   mutant knock-in mice. *Nat Med* **5**, 101-106.

767   [36]   Kawasumi M, Chiba T, Yamada M, Miyamae-Kaneko M, Matsuoka M, Nakahara J,
768   Tomita T, Iwatsubo T, Kato S, Aiso S, Nishimoto I, Kouyama K (2004) Targeted
769   introduction of V642I mutation in amyloid precursor protein gene causes functional
770   abnormality resembling early stage of Alzheimer's disease in aged mice. *Eur J*
771   *Neurosci* **19**, 2826-2838.

772   [37]   Xu J, Burgoyne PS, Arnold AP (2002) Sex differences in sex chromosome gene
773   expression in mouse brain. *Human Molecular Genetics* **11**, 1409-1419.

[38]   Bundy JL, Vied C, Nowakowski RS (2017) Sex differences in the molecular signature of
the developing mouse hippocampus. *BMC Genomics* **18**, 237.

[39]   Barthelson K, Pederson S, Newman M, Lardelli M (2020) Transcriptome analysis of a
protein-truncating mutation in sortilin-related receptor 1 associated with early-onset
familial Alzheimer's disease indicates effects on mitochondrial and ribosome function
in young-adult zebrafish brains. *bioRxiv*, 2020.2009.2003.282277.

[40]   Barthelson K, Pederson SM, Newman M, Jiang H, Lardelli M (2020) Frameshift and
frame-preserving mutations in zebrafish presenilin 2 affect different cellular
functions in young adult brains. *bioRxiv*, 2020.2011.2021.392761.

[41]   Barthelson K, Pederson SM, Newman M, Lardelli M (2020) Brain transcriptome
analysis reveals subtle effects on mitochondrial function and iron homeostasis of
mutations in the SORL1 gene implicated in early onset familial Alzheimer's disease.
Molecular Brain **13**, 142.

[42]   Hin N, Newman M, Kaslin J, Douek AM, Lumsden A, Nik SHM, Dong Y, Zhou X-F,
Mañucat-Tan NB, Ludington A, Adelson DL, Pederson S, Lardelli M (2020) Accelerated
brain aging towards transcriptional inversion in a zebrafish model of the K115fs
mutation of human PSEN2. *PLOS ONE* **15**, e0227258.

[43]   Jiang H, Pederson SM, Newman M, Dong Y, Barthelson K, Lardelli M (2020)
Transcriptome analysis indicates dominant effects on ribosome and mitochondrial
function of a premature termination codon mutation in the zebrafish gene psen2.
PLOS ONE **15**, e0232559.

[44]   Dong Y, Newman M, Pederson S, Hin N, Lardelli M (2020) Transcriptome analyses of
7-day-old zebrafish larvae possessing a familial Alzheimer's disease-like mutation in
psen1 indicate effects on oxidative phosphorylation, mcm functions, and iron
homeostasis. *bioRxiv*, 2020.2005.2003.075424.

[45]   Hin N, Newman M, Pederson SM, Lardelli MM (2020) Iron Responsive Element (IRE)-
mediated responses to iron dyshomeostasis in Alzheimer's disease. *bioRxiv*,
2020.2005.2001.071498.

[46]   Newman M, Hin N, Pederson S, Lardelli M (2019) Brain transcriptome analysis of a
familial Alzheimer's disease-like mutation in the zebrafish presenilin 1 gene implies
effects on energy production. *Molecular Brain* **12**.

[47] Ishikawa A, Piao Y-S, Miyashita A, Kuwano R, Onodera O, Ohtake H, Suzuki M,
Nishizawa M, Takahashi H (2005) A mutant PSEN1 causes dementia with lewy bodies
and variant Alzheimer's disease. *Annals of Neurology* **57**, 429-434.

[48] Zhang Y, Zhang Z, Ge W (2018) An efficient platform for generating somatic point
mutations with germline transmission in the zebrafish by CRISPR/Cas9-mediated
gene editing. *J Biol Chem* **293**, 6611-6622.

[49] Prykhozhij SV, Fuller C, Steele SL, Veinotte CJ, Razaghi B, Robitaille JM, McMaster CR,
Shlien A, Malkin D, Berman JN (2018) Optimized knock-in of point mutations in
zebrafish using CRISPR/Cas9. *Nucleic Acids Research* **46**, e102-e102.

[50] Jiang H, Newman M, Lardelli M (2018) The zebrafish orthologue of familial
Alzheimer's disease gene PRESENILIN 2 is required for normal adult melanotic skin
pigmentation. *PLOS ONE* **13**, e0206155.

[51] Di Tommaso P, Chatzou M, Floden EW, Barja PP, Palumbo E, Notredame C (2017)
Nextflow enables reproducible computational workflows. *Nature Biotechnology* **35**,
316-319.

[52] Team RC (2019) R: A language and environment for statistical computing. *R
Foundation for Statistical Computing, Vienna, Austria*.

[53] Chen Y, Lun A, Smyth G (2016) From reads to genes to pathways: differential
expression analysis of RNA-Seq experiments using Rsubread and the edgeR quasi-
likelihood pipeline [version 2; peer review: 5 approved]. *F1000Research* **5**.

[54] Robinson MD, Oshlack A (2010) A scaling normalization method for differential
expression analysis of RNA-seq data. *Genome Biology* **11**, R25.

[55] McCarthy DJ, Chen Y, Smyth GK (2012) Differential expression analysis of multifactor
RNA-Seq experiments with respect to biological variation. *Nucleic Acids Research* **40**,
4288-4297.

[56] Robinson MD, McCarthy DJ, Smyth GK (2009) edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**,
139-140.

[57] Young MD, Wakefield MJ, Smyth GK, Oshlack A (2010) Gene ontology analysis for
RNA-seq: accounting for selection bias. *Genome Biology* **11**, R14.

[58]   Wu D, Lim E, Vaillant F, Asselin-Labat M-L, Visvader JE, Smyth GK (2010) ROAST:
836   rotation gene set tests for complex microarray experiments. *Bioinformatics* **26**, 2176-
837   2182.

838   [59]   Wu D, Smyth GK (2012) Camera: a competitive gene set test accounting for inter-
839   gene correlation. *Nucleic acids research* **40**, e133-e133.

840   [60]   Sergushichev AA (2016) An algorithm for fast preranked gene set enrichment analysis
841   using cumulative statistic calculation. *bioRxiv*, 060012.

842   [61]   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich
843   A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP (2005) Gene set enrichment analysis:
844   A knowledge-based approach for interpreting genome-wide expression profiles.
845   *Proceedings of the National Academy of Sciences* **102**, 15545.

846   [62]   Kanehisa M, Goto S (2000) KEGG: kyoto encyclopedia of genes and genomes. *Nucleic
847   acids research* **28**, 27-30.

848   [63]   Liberzon A (2014) A description of the Molecular Signatures Database (MSigDB) Web
849   site. *Methods Mol Biol* **1150**, 153-160.

850   [64]   Dolgalev I (2020), p. R package.

851   [65]   Chen X (2011) Adaptive elastic-net sparse principal component analysis for pathway
852   association testing. *Statistical applications in genetics and molecular biology* **10**, 48.

853   [66]   Odom GJ, Ban Y, Liu L, Sun X, Pico AR, Zhang B, Wang L, Chen X (2019) pathwayPCA:
854   an R package for integrative pathway analysis with modern PCA methodology and
855   gene selection. *bioRxiv*, 615435.

856   [67]   Liberzon A, Birger C, Thorvaldsdóttir H, Ghandi M, Mesirov Jill P, Tamayo P (2015)
857   The Molecular Signatures Database Hallmark Gene Set Collection. *Cell Systems* **1**,
858   417-425.

859   [68]   Kosik KS, Muñoz C, Lopez L, Arcila ML, García G, Madrigal L, Moreno S, Ríos Romenets
860   S, Lopez H, Gutierrez M, Langbaum JB, Cho W, Suliman S, Tariot PN, Ho C, Reiman
861   EM, Lopera F (2015) Homozygosity of the autosomal dominant Alzheimer disease
862   presenilin 1 E280A mutation. *Neurology* **84**, 206-208.

[69]   Parker J, Mozaffar T, Messmore A, Deignan JL, Kimonis VE, Ringman JM (2019)
Homozygosity for the A431E mutation in PSEN1 presenting with a relatively
aggressive phenotype. *Neurosci Lett* **699**, 195-198.

[70]   Sundvik M, Chen Y-C, Panula P (2013) Presenilin1 Regulates Histamine Neuron
Development and Behavior in Zebrafish, Danio rerio. *The*
Journal of Neuroscience **33**, 1589.

[71]   Tambini MD, D'Adamio L (2020) Knock-in rats with homozygous PSEN1(L435F)
Alzheimer mutation are viable and show selective γ-secretase activity loss causing
low Aβ40/42 and high Aβ43. *J Biol Chem* **295**, 7442-7451.

[72]   Xia D, Watanabe H, Wu B, Lee SH, Li Y, Tsvetkov E, Bolshakov VY, Shen J, Kelleher RJ,
3rd (2015) Presenilin-1 knockin mice reveal loss-of-function mechanism for familial
Alzheimer's disease. *Neuron* **85**, 967-981.

[73]   Nickless A, Bailis JM, You Z (2017) Control of gene expression through the nonsense-
mediated RNA decay pathway. *Cell & Bioscience* **7**, 26.

[74]   Alhamdoosh M, Law CW, Tian L, Sheridan JM, Ng M, Ritchie ME (2017) Easy and
efficient ensemble gene set testing with EGSEA. *F1000Research* **6**, 2010-2010.

[75]   Wilson DJ (2019) The harmonic mean p-value for combining dependent tests.
Proceedings of the National Academy of Sciences **116**, 1195.

[76]   Pan Y, Lin MH, Tian X, Cheng HT, Gridley T, Shen J, Kopan R (2004) gamma-secretase
functions through Notch signaling to maintain skin appendages but is not required
for their patterning or initial morphogenesis. *Dev Cell* **7**, 731-743.

[77]   Blanpain C, Lowry WE, Pasolli HA, Fuchs E (2006) Canonical notch signaling functions
as a commitment switch in the epidermal lineage. *Genes Dev* **20**, 3022-3035.

[78]   Kopan R, Ilagan MX (2009) The canonical Notch signaling pathway: unfolding the
activation mechanism. *Cell* **137**, 216-233.

[79]   Nowell C, Radtke F (2013) Cutaneous Notch signaling in health and disease. *Cold*
Spring Harbor perspectives in medicine **3**, a017772-a017772.

[80]   Nguyen B-C, Lefort K, Mandinova A, Antonini D, Devgan V, Della Gatta G, Koster MI,
Zhang Z, Wang J, di Vignano AT, Kitajewski J, Chiorino G, Roop DR, Missero C, Dotto

GP (2006) Cross-regulation between Notch and p63 in keratinocyte commitment to
differentiation. *Genes & Development* **20**, 1028-1042.

[81] Dohda T, Maljukova A, Liu L, Heyman M, Grandér D, Brodin D, Sangfelt O, Lendahl U
(2007) Notch signaling induces SKP2 expression and promotes reduction of p27Kip1
in T-cell acute lymphoblastic leukemia cell lines. *Exp Cell Res* **313**, 3141-3152.

[82] Ryan RJH, Petrovic J, Rausch DM, Zhou Y, Lareau CA, Kluk MJ, Christie AL, Lee WY,
Tarjan DR, Guo B, Donohue LKH, Gillespie SM, Nardi V, Hochberg EP, Blacklow SC,
Weinstock DM, Faryabi RB, Bernstein BE, Aster JC, Pear WS (2017) A B Cell Regulome
Links Notch to Downstream Oncogenic Pathways in Small B Cell Lymphomas. *Cell Rep*
21, 784-797.

[83] El-Brolosy MA, Kontarakis Z, Rossi A, Kuenne C, Günther S, Fukuda N, Kikhi K, Boezio
GLM, Takacs CM, Lai S-L, Fukuda R, Gerri C, Giraldez AJ, Stainier DYR (2019) Genetic
compensation triggered by mutant mRNA degradation. *Nature* **568**, 193-197.

[84] Rossi A, Kontarakis Z, Gerri C, Nolte H, Hölper S, Krüger M, Stainier DYR (2015)
Genetic compensation induced by deleterious mutations but not gene knockdowns.
Nature **524**, 230-233.

[85] Luo W, Pant G, Bhavnasi YK, Blanchard SG, Jr., Brouwer C (2017) Pathview Web: user
friendly pathway visualization and data integration. *Nucleic Acids Research* **45**, W501-
W508.

[86] Ryman DC, Acosta-Baena N, Aisen PS, Bird T, Danek A, Fox NC, Goate A, Frommelt P,
Ghetti B, Langbaum JBS, Lopera F, Martins R, Masters CL, Mayeux RP, McDade E,
Moreno S, Reiman EM, Ringman JM, Salloway S, Schofield PR, Sperling R, Tariot PN,
Xiong C, Morris JC, Bateman RJ (2014) Symptom onset in autosomal dominant
Alzheimer disease: A systematic review and meta-analysis. *Neurology* **83**, 253-260.

[87] Oksanen M, Petersen AJ, Naumenko N, Puttonen K, Lehtonen Š, Gubert Olivé M,
Shakirzyanova A, Leskelä S, Sarajärvi T, Viitanen M, Rinne JO, Hiltunen M, Haapasalo
A, Giniatullin R, Tavi P, Zhang S-C, Kanninen KM, Hämäläinen RH, Koistinaho J (2017)
PSEN1 Mutant iPSC-Derived Model Reveals Severe Astrocyte Pathology in
Alzheimer's Disease. *Stem Cell Reports* **9**, 1885-1897.

[88] Martín-Maestro P, Sproul A, Martinez H, Paquet D, Gerges M, Noggle S, Starkov AA
(2019) Autophagy Induction by Bexarotene Promotes Mitophagy in Presenilin 1
Familial Alzheimer's Disease iPSC-Derived Neural Stem Cells. *Molecular Neurobiology*
56, 8220-8236.

[89]   Birnbaum JH, Wanner D, Gietl AF, Saake A, Kündig TM, Hock C, Nitsch RM,
926   Tackenberg C (2018) Oxidative stress and altered mitochondrial protein expression in
927   the absence of amyloid-β and tau pathology in iPSC-derived neurons from sporadic
928   Alzheimer's disease patients. *Stem Cell Res* **27**, 121-130.

929   [90]   Kwart D, Gregg A, Scheckel C, Murphy EA, Paquet D, Duffield M, Fak J, Olsen O,
930   Darnell RB, Tessier-Lavigne M (2019) A Large Panel of Isogenic APP and PSEN1
931   Mutant Human iPSC Neurons Reveals Shared Endosomal Abnormalities Mediated by
932   APP β-CTFs, Not Aβ. *Neuron* **104**, 256-270.e255.

933   [91]   Volpato V, Smith J, Sandor C, Ried JS, Baud A, Handel A, Newey SE, Wessely F, Attar
934   M, Whiteley E, Chintawar S, Verheyen A, Barta T, Lako M, Armstrong L, Muschet C,
935   Artati A, Cusulin C, Christensen K, Patsch C, Sharma E, Nicod J, Brownjohn P, Stubbs
936   V, Heywood WE, Gissen P, De Filippis R, Janssen K, Reinhardt P, Adamski J, Royaux I,
937   Peeters PJ, Terstappen GC, Graf M, Livesey FJ, Akerman CJ, Mills K, Bowden R,
938   Nicholson G, Webber C, Cader MZ, Lakics V (2018) Reproducibility of Molecular
939   Phenotypes after Long-Term Differentiation to Human iPSC-Derived Neurons: A
940   Multi-Site Omics Study. *Stem Cell Reports* **11**, 897-911.

941   [92]   Hunger RE, Surovy AM, Hassan AS, Braathen LR, Yawalkar N (2008) Toll-like receptor
942   2 is highly expressed in lesions of acne inversa and colocalizes with C-type lectin
943   receptor. *British Journal of Dermatology* **158**, 691-697.

944   [93]   Otani K, Shichita T (2020) Cerebral sterile inflammation in neurodegenerative
945   diseases. *Inflammation and Regeneration* **40**, 28.

946   [94]   Zouboulis CC, Benhadou F, Byrd AS, Chandran NS, Giamarellos-Bourboulis EJ,
947   Fabbrocini G, Frew JW, Fujita H, González-López MA, Guillem P, Gulliver WPF,
948   Hamzavi I, Hayran Y, Hórvath B, Hüe S, Hunger RE, Ingram JR, Jemec GBE, Ju Q,
949   Kimball AB, Kirby JS, Konstantinou MP, Lowes MA, MacLeod AS, Martorell A,
950   Marzano AV, Matusiak Ł, Nassif A, Nikiphorou E, Nikolakis G, Nogueira da Costa A,
951   Okun MM, Orenstein LAV, Pascual JC, Paus R, Perin B, Prens EP, Röhn TA, Szegedi A,
952   Szepietowski JC, Tzellos T, Wang B, van der Zee HH (2020) What causes hidradenitis
953   suppurativa ?—15 years after. *Experimental Dermatology* **29**, 1154-1170.

954   [95]   Research IoMUCoMN (1999) The Energy Costs of Protein Metabolism: Lean and
955   Mean on Uncle Sam's Team In *The Role of Protein and Amino Acids in Sustaining and*
956   *Enhancing Performance* National Academies Press (US), Washington (DC).

957   [96]   Bordi M, Darji S, Sato Y, Mellén M, Berg MJ, Kumar A, Jiang Y, Nixon RA (2019) mTOR
958   hyperactivation in Down Syndrome underlies deficits in autophagy induction,
959   autophagosome formation, and mitophagy. *Cell Death & Disease* **10**, 563.

[97] Jiang Y, Sato Y, Im E, Berg M, Bordi M, Darji S, Kumar A, Mohan PS, Bandyopadhyay
U, Diaz A, Cuervo AM, Nixon RA (2019) Lysosomal Dysfunction in Down Syndrome Is
APP-Dependent and Mediated by APP-βCTF (C99). *The Journal of Neuroscience* **39**,
5255.

[98] Lee JH, Yu WH, Kumar A, Lee S, Mohan PS, Peterhoff CM, Wolfe DM, Martinez-
Vicente M, Massey AC, Sovak G, Uchiyama Y, Westaway D, Cuervo AM, Nixon RA
(2010) Lysosomal proteolysis and autophagy require presenilin 1 and are disrupted
by Alzheimer-related PS1 mutations. *Cell* **141**, 1146-1158.

[99] Linke M, Fritsch SD, Sukhbaatar N, Hengstschläger M, Weichhart T (2017) mTORC1
and mTORC2 as regulators of cell metabolism in immunity. *FEBS letters* **591**, 3089-
3103.

[100] Inpanathan S, Botelho RJ (2019) The Lysosome Signaling Platform: Adapting With the
Times. *Frontiers in Cell and Developmental Biology* **7**, 113.

[101] Bond P (2016) Regulation of mTORC1 by growth factors, energy status, amino acids
and mechanical stimuli at a glance. *J Int Soc Sports Nutr* **13**, 8.

[102] Loewith R, Hall MN (2011) Target of rapamycin (TOR) in nutrient signaling and growth
control. *Genetics* **189**, 1177-1201.

[103] Lempiäinen H, Shore D (2009) Growth control and ribosome biogenesis. *Current
Opinion in Cell Biology* **21**, 855-863.

[104] Newman M, Wilson L, Verdile G, Lim A, Khan I, Moussavi Nik SH, Pursglove S,
Chapman G, Martins RN, Lardelli M (2014) Differential, dominant activation and
inhibition of Notch signalling and APP cleavage by truncations of PSEN1 in human
disease. *Human Molecular Genetics* **23**, 602-617.

[105] Kelleher RJ, Shen J (2017) Presenilin-1 mutations and Alzheimer's disease.
Proceedings of the National Academy of Sciences **114**, 629.

[106] Zhou R, Yang G, Shi Y (2017) Dominant negative effect of the loss-of-function γ-
secretase mutants on the wild-type enzyme through heterooligomerization. *Proc Natl
Acad Sci U S A* **114**, 12731-12736.

[107] Chávez-Gutiérrez L, Bammens L, Benilova I, Vandersteen A, Benurwar M, Borgers M,
Lismont S, Zhou L, Van Cleynenbreugel S, Esselmann H, Wiltfang J, Serneels L, Karran
E, Gijsen H, Schymkowitz J, Rousseau F, Broersen K, De Strooper B (2012) The

mechanism of γ-Secretase dysfunction in familial Alzheimer disease. *The EMBO*
Journal **31**, 2261-2274.

[108]   Zhang S, Cai F, Wu Y, Bozorgmehr T, Wang Z, Zhang S, Huang D, Guo J, Shen L, Rankin
994   C, Tang B, Song W (2020) A presenilin-1 mutation causes Alzheimer disease without
995   affecting Notch signaling. *Molecular Psychiatry* **25**, 603-613.

996   [109]   Deatherage CL, Lu Z, Kroncke BM, Ma S, Smith JA, Voehler MW, McFeeters RL,
997   Sanders CR (2017) Structural and biochemical differences between the Notch and the
998   amyloid precursor protein transmembrane domains. *Science Advances* **3**, e1602794.

999   [110]   Bukhari H, Glotzbach A, Kolbe K, Leonhardt G, Loosse C, Müller T (2017) Small things
1000   matter: Implications of APP intracellular domain AICD nuclear signaling in the
1001   progression and pathogenesis of Alzheimer's disease. *Progress in Neurobiology* **156**,
1002   189-213.

1003