Fine mapping without phenotyping: Identification of selection targets in secondary Evolve

and Resequence experiments

Anna Maria Langmüller[1, 2], Marlies Dolezal[3], Christian Schlötterer[1, *]

[1] Institut für Populationsgenetik, Vetmeduni Vienna, Veterinärplatz 1, 1210 Vienna, Austria

[2] Vienna Graduate School of Population Genetics, Vetmeduni Vienna, Veterinärplatz 1, 1210

Vienna, Austria

[3] Plattform Bioinformatik und Biostatistik, Vetmeduni Vienna, Veterinärplatz 1, 1210 Vienna,

Austria

*Author of Correspondence:

Christian Schlötterer, Institut für Populationsgenetik, Vetmeduni Vienna, Veterinärplatz 1,

1210 Vienna, Austria

+43 1 25077-4301, +43 25077-4390

christian.schloetterer@vetmeduni.ac.at

Abstract

Evolve and Resequence (E&R) studies investigate the genomic selection response of populations in an Experimental Evolution (EE) setup. Despite the popularity of E&R, empirical studies typically suffer from an excess of candidate loci due to linkage, and single gene or SNP resolution is the exception rather than the rule. Recently, a secondary E&R design – where unevolved founder genotypes are added to a primary E&R study - has been suggested as promising experimental procedure to confirm putative selection targets. Furthermore, secondary E&R provides the opportunity to increase mapping resolution by allowing for additional recombination events, which separate the selection target from neutral hitchhikers. Here, we use computer simulations to assess the effect of crossing scheme, population size, experimental duration, and number of replicates on the power, and resolution of secondary E&R. We find that the crossing scheme, population size, and the experimental duration are crucial factors for the power and resolution of secondary E&R: a simple crossing scheme with few founder lines consistently outcompetes crossing schemes where evolved populations from a primary E&R experiment are mixed with a complex ancestral founder population. Regardless of the experimental design tested, a population size of at least 4,800 individuals, which is roughly 5 times larger than population sizes in typical E&R studies, is required to achieve a power of at least 75%. Our study provides an important step towards improved experimental designs aiming to characterize causative SNPs in EE studies.

Keywords

Experimental Evolution, secondary Evolve and Resequence, experimental design, *Drosophila*, fine mapping

Introduction

Deciphering the genetic architecture of adaptation is one of the longstanding goals in evolutionary biology. Experimental Evolution (EE) has become a popular approach to study adaptation in real time (1, 2). In contrast to natural populations EE offers the key advantage of replicating experiments under controlled laboratory conditions (3). Evolve and Resequence (E&R) (3–5) – a combination of EE with next-generation-sequencing – facilitates in-depth analysis of the genomic responses to selection, with the ultimate goal to identify and characterize individual adaptive loci.

E&R has already been successful in investigating genomic selection responses from standing genetic variation in adapting sexually reproducing organisms, such as chicken (6), yeast (7), *C. elegans* (8) and *Drosophila* (9–12). Despite its popularity, E&R typically suffers from an excess of candidates caused by linkage disequilibrium between true causative SNPs and neutral hitchhikers (13–15), which decreases the resolution of E&R studies and makes single gene resolution (9) the exception rather than the rule.

The problem of candidate excess in E&R studies has been approached from different angles. More refined statistical tests have been developed (16–19), and the combination of time-series data with replicate populations has been identified as particularly powerful (7, 20, 21). Organisms with a higher recombination rate and a lack of large segregating inversions that suppress recombination events have been suggested to be better suited for E&R studies (22). Computer simulations showed that the power of E&R studies can be significantly improved by increasing the number of replicate populations, the experimental duration, or by adjusting the applied selection regime (23–26).

Secondary E&R (27) has been recently introduced as a promising experimental follow-up

approach to validate selection signals of primary E&R. The basic idea of secondary E&R is that

putative selection targets determined in the primary E&R study should rise in frequency again

when exposed to the same environmental conditions. This experimental validation of

selection signals is especially attractive before starting the time-consuming functional

characterization of putatively selected alleles (e.g. based on the CRISPR/Cas technology (28)).

A hitherto under-explored potential of secondary E&R is that the additional recombination

events during the secondary E&R can be used to fine map selection signals of primary

experiments.

Here we suggest additional secondary E&R experimental designs and evaluate their power

and resolution to identify causative SNPs via extensive computer simulations. We use logistic

regression to assess which simulated experimental and population genetic parameters have

a significant effect on the success of secondary E&R. Selection coefficient, dominance

coefficient, and mean starting allele frequency of the selection target all have a significant

effect on the success of secondary E&R. However, crossing scheme and population size

emerge as the most influential parameters. We show that the population size of secondary

E&R experiments needs to be at least 5 times larger than currently used population sizes in

typical primary E&R studies with *Drosophila* to achieve a power above 75%. Furthermore, we

show that the crossing scheme is a crucial experimental parameter shaping the power of

secondary E&R - a simple crossing scheme with few founder lines results in higher power and

resolution compared to more complex crossing schemes.

Methods

<u>Outline of the simulation framework</u>

The founder lines used in our simulation study are a randomly chosen subset of 100

haplotypes from a panel of 189 sequenced *D. simulans* haplotypes originally collected in

Tallahassee, Florida, USA capturing the amount of standing genetic variation in a natural

Drosophila population (12, 29). In order to speed up the calculations, we only simulated

chromosome-arm 2L. Simulations with linkage were conducted with MimicrEE2 (v206) (30)

using the *D. simulans* recombination map (29). MimicrEE2 is a forward-simulation framework

for E&R studies that can simulate evolving experimental populations based on their haplotype

information and genome-wide recombination rates. We used the w-mode of MimicrEE2

which computes the fitness of individuals directly from the selection coefficients. We

simulated positive, additive selection for biallelic SNPs, with selection coefficients being

uniformly sampled between 0.07 and 0.1 for each positively selected SNP and tracked the

frequency of all SNPs over time (ranging from 479,507 to 970,466 SNPs, Table S1). We chose

to simulate rather strong selection reasoning that alleles with a high selection coefficient are

more likely to be experimentally tested. On the other side, strongly selected alleles result in

many neutral linked hitchhikers producing false positive signals that adversely impact the

mapping resolution (25). Selected SNPs were randomly chosen with an equal probability to

be either co-dominant (dominance coefficient h=0.5), or dominant (h=1). We did not consider

recessive loci (h=0), because we do not anticipate that fully recessive targets would result in

sufficiently large allele frequency changes to be detected in primary E&R experiments (24,

25). We rescaled allele counts for each biallelic position to a uniform read depth of 80.

<u>Experimental parameters</u>

The purpose of this study is to test the influence of different experimental parameters on the

power and resolution of secondary E&R. For this, we systematically varied the crossing

scheme, population size, experimental duration, and number of replicates to assess the effect

of these experimental parameters on the power and resolution of secondary E&R (Table 1).

We use the term "experimental design" to describe a distinct set of simulated experimental

parameters (*e.g.,* crossing scheme=1:1; population size=1,200 individuals; experimental

duration=60 generations; 5 replicates).

Crossing scheme

We simulated five different crossing schemes: (A) 1:1, (B) 1:few, (C) 1:many, (D) dil:st, and (E)

dil:mt (Figure 1). In the **1:1** crossing scheme (Figure 1A), two inbred lines are crossed at equal

proportions. One inbred focal line carries a single target of selection - known from a primary

E&R experiment – and is crossed with inbred non-focal lines, not carrying known adaptive

alleles. Replicates are created by crossing the same focal line to different non-focal lines

without known targets of selection. We used different non-focal lines as crossing partners to

account for the possibility that in real experiments these lines may contain unidentified

selected loci. By using different lines, the influence of selection targets present in a single line

will be outweighed by the focal locus, which is present in every replicate (Figure 1A).

In the **1:few** crossing scheme (Figure 1B) the focal line is crossed with a pool of five non-focal

lines that do not carry known beneficial alleles. The starting frequency of the focal line is 50

%, whereas each non-focal line has a starting frequency of 10 %. Each replicate consists of the

same focal/non-focal line mixture. In the **1:many** crossing scheme (Figure 1C), the focal line

is crossed with a pool of 99 non-focal lines. Replicates consist of the same mixture of lines,

and the starting frequency of the focal line is 50 %.

It has been recently suggested, that "diluting" evolved populations of a primary E&R

experiment with many ancestral genotypes of the very same primary E&R and exposing the

diluted populations to the same selection regime is a promising approach to validate selection

candidates (27). However, we lack a systematic power assessment of such experiments with

computer simulations. We thus included two "dilution" crossing schemes (**dil:mt, dil:st**) into

our analysis. To evaluate the power of dilution crossing schemes, it is important to first

simulate a primary E&R study. We chose to simulate a population consisting of 100 different

founder lines, a population size of 300 individuals, 60 generations of adaptation and one

replicate, which can then be diluted with genotypes of the ancestral population. SNPs in the

primary E&R were tested for allele frequency change with the χ^2 test.

The population of the **dil:st** crossing scheme carries only one beneficial SNP (**dil:st** = **dil**ution:

single **t**arget, Figure 1D). The starting allele frequency of this beneficial, focal SNP is sampled

from the empirical starting allele frequency distribution of putatively selected alleles from a

previous E&R study in which *D. simulans* populations adapted to a new temperature regime

(mean starting frequency = 0.1) (12). After simulating 60 generations of adaptation (primary

E&R), 50 % of the evolved population is replaced by flies of the ancestral founder population.

After this dilution step, the secondary E&R was simulated under the exact same selection

regime as in the primary E&R. A region with a strong selection signal from the primary E&R is

chosen for validation in the dilution crossing schemes. Hence, we investigated a 1 Mb

window, which is the previously reported median selected haplotype block length on

chromosome-arm 2L (12), around the SNP with the highest χ^2 test-statistic in the primary

E&R, in the secondary E&R.

The **dil:mt** (Figure 1E) crossing scheme has multiple selection targets (**dil:mt** = **dil**ution:

multiple **t**argets; 16 selection targets on chromosome-arm 2L (12)). Again, we investigate a 1

Mb window around the SNP with the highest χ^2 test statistic in the primary E&R, in the

secondary E&R. In case there are multiple selection targets in the 1 Mb window, we consider

the target that is closest to the SNP with the highest CMH test statistic in the 1Mb window of

the secondary E&R in our analysis.

171
Figure 1 Simulated crossing schemes. (A) **1:1** crossing scheme: Inbred flies with one target of selection (purple)
are crossed to inbred flies without known beneficial variants. The starting frequency of each genotype is 50 %.
In each replicate the line with the beneficial allele (focal line, purple) is crossed to a different line lacking
beneficial mutations (non-focal lines, different shades of grey). (B) **1:few** crossing scheme: The focal line is
crossed to a pool of flies with five different genotypes without known selection targets. The starting frequency
of the focal line is again 50 %. (C) **1:many** crossing scheme: The focal line is crossed to a pool of 99 lines without
known selection targets. (D) **dil:st** crossing scheme: 50 % of an evolved population originating from a primary
E&R is replaced by ancestral genotypes of the primary E&R. The entire population has only one single target of
selection (focal SNP, purple flies) (E) **dil:mt** crossing scheme: 50 % of an evolved population originating from a
primary E&R is replaced by ancestral genotypes of the primary E&R. The ancestral population carries 16 targets
of selection (flies carrying different beneficial SNPs are shown in purple, and green).

Population size

For two crossing schemes that are relatively easy to implement in empirical studies (1:1,

dil:mt) we performed simulations (experimental duration = 60 generations; 5 replicates) with

independently sampled selection targets for population sizes of 300; 1,200; 4,800; and 19,200

individuals per replicate to test for the effect of the population size on the power and

resolution of secondary E&R (Table 1). All other crossing schemes were evaluated at a

population size of 300 individuals.

Experimental duration

All crossing schemes were evaluated after 60 generations. To explore the possibility that

shorter experiments with less than 60 generations may already be sufficient to achieve

satisfactory power, we analyzed the simulations for 1:1 and dil:mt crossing schemes also

already after 20 generations (Table 1).

Number of replicates

To assess the impact of the number of replicates on power of secondary E&R, we conducted

for each crossing scheme 100 additional simulations (population size= 300 individuals;

experimental duration = 60 generations) with 30 replicates (Table 1).

Table 1 Simulation overview.

Crossing scheme	Population size	Generations	Replicates	Number of simulations
1:1	300; 1,200; 4,800; 19,200	60; 20	5/30	2,000/100
1:few	300	60	5/30	2,000/100
1:many	300	60; 20	5/30	2,000/100
dil:st	300	60	5/30	2,000/100
dil:mt	300; 1,200; 4,800; 19,200	60; 20	5/30	2,000/100

 <u>Statistical analysis</u>

All statistical analyses were performed using the R statistical computing environment (v3.5.3)

(31).

Power

For each simulation, we tested all SNPs on chromosome-arm 2L for allele frequency increase

between the start and the end of the simulated secondary E&R using the Cochran-Mantel-

Haenszel-test (CMH-test, implemented in the R package poolSeq (v0.3.2) (32)). The CMH-test

allows to test for independence of matched data – e.g. allele counts of replicated ancestral

and evolved populations (33). We ranked SNPs based on their CMH test statistic using the

dense ranking method in the R package data.table (v.1.12-8 (34)). In dense ranking, SNPs with

identical test statistics receive the same rank, and the following SNP is assigned the

immediately following rank.

Based on this ranking, we used two different approaches to classify simulations being either

successful, or unsuccessful. First, we considered a simulation to be only successful if the true

target of selection (*i.e.,* the focal SNP with a selection coefficient > 0) was the SNP with the

highest test statistic (success-A). In a second analysis step, we classified a simulation as

success, if the true target of selection was not more than 100 SNPs away from the SNP with

the highest test statistic (Figure 2, success-B). We acknowledge that success-B is dependent

on the maximum distance that we allow between the target of selection and the SNP with

the highest CMH test statistic (Figure 2). However, varying the maximum distance threshold

did not alter the relative performance of different experimental designs (data not shown).

The power of an experimental design is defined as the proportion of simulations that were

able to detect the true target of selection.

230
Figure 2 Schematic overview of the definition of success-B in a secondary Evolve and Resequence simulation. All
SNPs (neutral = circle, beneficial = purple diamond) are tested for an allele frequency change with the Cochran-
Mantel-Haenszel-test, and are ranked based on their test statistic (y-axis). If the true selection target (purple
diamond) is less than 100 SNPs away from the SNP with the highest test statistic, the simulation run is deemed
a success. In the example depicted, the distance in number of SNPs between the true target of selection, and
the SNP with the highest test statistic is 4, and the simulation is classified as success.

Resolution

For simulations where the selection target was detected (success-B), we determined the

resolution of fine mapping of the selection target by counting the number of SNPs between

the true selection target, and the SNP with the highest CMH test statistic.

Assessment of experimental and population genetic parameters

We used logistic regressions with a binomial error structure (ε) and a logit link function (35)

to test how different experimental and population genetic parameters affect success and

failure to identify targets of selection, i.e. success (Y) is treated as a binary response encoded

in 0 and 1 of secondary E&R. We fitted two different models with R function *glm* with μ being

the overall mean per model:

(1) $Y_{ijklmn} = \mu + cross_i + h_j + s_k + af_l + N_m + (cross{:}h)_{ij} + (h{:}af)_{jl} +$

$(cross{:}N)_{im} + (h{:}N)_{jm} + (cross{:}h{:}N)_{ijm} + \varepsilon_{ijklmn}$

(2) $Y_{ijklm} = \mu + cross_i + h_j + s_k + af_l + (cross{:}h)_{ij} + (h{:}af)_{jl} + \varepsilon_{ijklm}$

Model 1 (equation (1)) contained (i) five explanatory variables as main effects; crossing

scheme ($cross_i$), a fixed categorical effect with 2 levels - 1:1 and dil:mt; a fixed categorical

effect of dominance coefficient (h_j) with levels 0.5 and 1; selection coefficient (s_k) and mean

starting allele frequency (af_l) over replicated populations (both as continuous covariate); and

population size (N_m), a fixed categorical effect with 4 levels - 300; 1,200; 4,800; 19,200, (ii) an

interaction term between dominance coefficient and mean starting allele frequency

($(h{:}af)_{jl}$), (iii) and a triple interaction between crossing scheme, dominance coefficient, and

population size ($(cross{:}h{:}N)_{ijm}$), and all pairwise interaction terms of effects involved in the

triple interaction term ($(cross{:}h)_{ij}$; $(cross{:}N)_{im}$; $(h{:}N)_{jm}$). We included interaction terms

into the model that have population genetic interpretations.

Prior to model fitting, the two covariates selection coefficient and mean starting allele

frequency were multiplied by 100, and z-transformed to a mean of zero and standard

deviation of one for easier interpretable estimates (36). Data analyzed with model 1

contained 16,000 observations, namely 2,000 independent simulation runs for each crossing

scheme (1:1, dil:mt), and each of the four different population sizes (300; 1,200; 4,800;

19,200) (Table 1). To avoid potential bias, we randomly chose 4 sets of focal/non-focal

founder lines for the 1:1 crossing scheme and performed 500 simulation runs per set. The

chosen set of founder lines did not have a significant effect on the success of secondary E&R

and is thus not included in the final model (LRT full-null model comparison; data not shown).

Model 2 (equation (2)) contained (i) four explanatory variables; crossing scheme ($cross_i$), a

fixed categorical effect with 5 levels - 1:1; 1:few; 1:many; dil:st; dil:mt, dominance coefficient

(h_j), selection coefficient (s_k), and mean starting allele frequency (af_l) over replicated

populations as main effects, as described for model 1, (ii) an interaction term between

crossing scheme and dominance coefficient (($cross{:}h)_{ij}$), (iii) and an interaction term

between dominance coefficient and mean starting allele frequency (($h{:}af)_{jl}$),. We analyzed

10,000 samples (2,000 independent simulation runs for each crossing scheme) (Table 1). For

crossing schemes using only 6 different founder lines (1:1, 1:few), the simulation runs are

based on 4 randomly chosen sets of founder lines each (500 simulation runs per set). Again,

the set of founder lines does not have a significant effect on secondary E&R success and is

thus not included in the final model (LRT full-null model comparison; data not shown). We

performed additional analysis with 500 observations (100 samples per crossing scheme) to

determine the influence of the number of replicates on the power of secondary E&R (Table

1).

We performed all diagnostic checks required for logistic regression. Absence of collinearity

was confirmed by computing the generalized Variance Inflation Factors (37) using function *vif*

in R package *car* (v3.0-8 (38)). Model stability was checked with the R function *dfbeta*. For

visualization, linear predictors (LP) were back-transformed to success probabilities using the

inverse logit transformation: $p_{sucess} = \frac{e^{LP}}{1+e^{LP}}$. 95 % confidence intervals of the fitted values

were investigated with the R function *predict.glm*. Significance of single explanatory variables

was tested with a Type III Sum of Squares using function *Anova* in R package car (38).

Significance of explanatory variables including their modeled interactions was tested with a

likelihood ratio test (LRT) comparing the full model with a nested reduced model with the

same structure as the full model, but lacking the assessed explanatory variable (and its

interactions). Significance is declared at an alpha cut off-off of 5%. We used Nagelkerke's R^2

index (39) to calculate the improvement of each model parameter upon the prediction of a

null model.

<u>Data availability</u>

Information regarding the accessibility of raw sequence reads, phased haplotypes of the

ancestral *D. simulans* haplotypes as well as MimicrEE2 ready text files for the *D. simulans*

recombination map (file used in this project: Dsim_recombination_map_LOESS_100kb_1.txt)

can be found in (29). MimicrEE2 ready input files of the different experimental designs, the

simulated selection regimes, processed simulation results and all scripts that are necessary to

reproduce the results are available at SourceForge (link TBD).

Results

We used forward simulations to assess the influence of different experimental and population genetic parameters, more specifically crossing scheme, population size, dominance coefficient, selection strength, and mean starting allele frequency on the success to detect and fine map selection targets in secondary E&R experiments. We used a Cochran-Mantel-Haenszel (CMH) test to identify SNPs rising in allele frequency. The CMH test allows to test for independence of matched categorical data (33), and compares favorably to other statistical methods in reliably identifying possible targets of selection in E&R setups (40). A simulation run was considered successful, if the true target of selection was the SNP with the highest CMH test statistic (success-A). In a second analysis step, we considered simulation runs as successful, if the true target of selection was not more than 100 SNPs away from the SNP with the highest CMH test statistic (Figure 2; success-B). We used logistic regression to assess which experimental and population genetic parameters have a significant effect on the success of secondary E&R.

First, we evaluated two crossing schemes that can be easily implemented in empirical studies, more specifically the 1:1 (Figure 1A) and dil:mt (Figure 1E) crossing schemes. The 1:1 crossing scheme is solely based on founder lines (Figure 1A). Using information about the selected haplotype from a primary E&R study, it is possible to determine which founder lines carry a selection target (12), which requires them to be sequenced. Crossing a focal founder line with the selected haplotype with a non-focal line without known selection targets offers the advantage of reducing the number of selection targets dramatically. Using different non-focal lines without known strong selection targets in each replicate reduces the potential of confounding effects of unidentified selection targets in these lines. On the other hand, dil:mt

(Figure 1E) represents an entirely different approach. Dil:mt, which has multiple targets of

selection at different frequencies, is probably the simulated crossing scheme with the most

straight forward implementation (12, 27). It is based on a "dilution" approach, where evolved

individuals from a primary E&R experiment are crossed to the ancestral founder population

from the same primary E&R (27). In contrast to 1:1, a dil:mt crossing scheme requires both –

ancestral and evolved – populations of the primary E&R, but relatively limited information

about selection targets on individual founder lines.

To evaluate the 1:1 and dil:mt crossing scheme, we simulated secondary E&R consisting of 5

replicated populations with constant population size (300; 1,200; 4,800, or 19,200 individuals

per replicate) that evolve for 60 generations, and used logistic regression to assess the effects

of model parameters on secondary E&R success (Model 1, *Assessment of experimental and*

population genetic parameters in Material & Methods). While selection strength, dominance

coefficient, and the interaction between dominance coefficient and mean starting allele

frequency all have a significant effect on the power of secondary E&R (Table S2 (success-A),

Table S3 (success-B)), our analysis reveals that crossing scheme and population size have by

far the strongest influence on secondary E&R success. Both the crossing scheme (LRT full-null

model comparison (Model 1): $\chi^2 = 3510.1$, df = 8, p<0.001 (success-A); $\chi^2 = 3211.3$, df = 8,

p<0.001 (success-B)); and the population size (LRT full-null model comparison (Model 1): $\chi^2 =$

2633.4, df = 12, p<0.001 (success-A); $\chi^2 = 2142.9$, df = 12, p<0.001 (success-B)) have a

significant effect on the success of secondary E&R, and are the only parameters with a

Nagelkerke's R^2 index above 0.2 (Table 2).

While a 1:1 crossing scheme results in higher power values than the dil:mt crossing scheme

independently of the population size, the difference to the dil:mt crossing scheme is more

pronounced in bigger experimental populations (Figure 3). This does not hold only for the

power, but also for the resolution (= the distance in SNPs between the true target of selection,

and the SNP with the highest CMH test statistic, Figure 4). A potential disadvantage of the 1:1

crossing scheme is that the low number of different founder lines (n=6) causes more linkage

disequilibrium, indicated by the number of neighboring SNPs with identical test statistics

(=ties) and thus broadens signatures compared to dil:mt crossing scheme that has more

founder lines (n=100) (Figure S1). However, this is outweighed by superior power of the 1:1

crossing scheme at every population size investigated (Figure 3, Figure 4).

Table 2 Nagelkerke's R^2-index of each simulation parameter including its interaction in the logistic regression (2
crossing schemes (1:1; dil:mt), 4 population sizes (300; 1,200; 4,800; 19,200 individuals); Figure 3). If the reduced
model explains the data as well as a full model that includes the evaluated effect, Nagelkerke's R^2-index is 0.

effect	R^2 (success-A)	R^2 (success-B)
crossing scheme	0.305	0.264
dominance coefficient	0.077	0.069
selection coefficient	0.007	0.007
mean starting allele frequency	0.003	0.002
population size	0.204	0.224

379
Figure 3 Power of the 1:1 and dil:mt crossing scheme at different population sizes (2,000
simulations/experimental design). Bars show the power (*i.e.,* the proportion of successful simulations)
separately for each combination of crossing scheme (1:1, dil:mt), population size (300; 1,200; 4,800; 19,200
individuals), and dominance coefficient (additive in grey, dominant in white). The dots with error bars display
the estimate from the fitted model (Model 1) and its 95 % confidence interval. For the model fit, the selection
coefficient was fixed to its global average, and combination-specific average starting allele frequencies were
used. (A) shows the results for success-A (= selection target is the SNP with the highest Cochran-Mantel-Haenszel
(CMH) test statistic), (B) shows the results for success-B (= selection target is not more than 100 SNPs away from
the SNP with the highest CMH tests).

390
Figure 4 Resolution of the 1:1 and dil:mt crossing scheme at different population sizes. Proportion of simulations
(y-axis) that do not exceed a maximum distance in SNPs (x-axis) between the SNP with the highest Cochran-
Mantel-Haenszel test statistic and the true target of selection. Each panel shows the result for one simulated
population size.

<u>Experimental duration</u>

Our analysis has shown that after 60 generations of adaptation, only experimental designs

with a population size of at least 1,200 individuals exceed power values of 50 % (Figure 3).

The 1:1 crossing scheme clearly outperforms dil:mt regardless of the population size, and

reaches power above 75% only with a population size of at least 4,800 individuals. However,

the maintenance of 5 replicates for 60 generations is labor intensive and can be quite time

consuming for most sexual organisms. We explored the possibility that shorter experiments

with less than 60 generations may already be sufficient to achieve satisfactory power values.

We thus reanalyzed the power of the 1:1, and dil:mt crossing scheme after 20 generations of

adaptation (Table 1).

Consistent with previous computer simulation studies (23–25) and empirical results (41), we

observe reduced power for experimental designs with shorter experimental duration (Figure

S2). As observed after 60 generations, the crossing scheme (LRT full-null model comparison

(Model 1): $\chi^2 = 2672.7$, df = 8, p<0.001 (success-A); $\chi^2 = 2234.4$, df = 8, p<0.001 (success-

B)) and the population size (LRT full-null model comparison (Model 1): $\chi^2 = 3176.4$, df = 12,

p<0.001 (success-A); $\chi^2 = 3375.3$, df = 12, p<0.001 (success-B)) have the biggest effects on

the success of secondary E&R (Table S4 (success-A), Table S5 (success-B), Table S6). In contrast

to our analysis after 60 generations of adaptation, the dil:mt crossing scheme results in higher

power than the 1:1 crossing scheme for populations with the smallest simulated population

size (300 individuals) after 20 generations of adaptation (Figure S2). A possible explanation

for this is that the dil:mt crossing scheme has multiple beneficial targets. Because we simulate

additive selection, linked selection targets can act synergistically and increase the frequency

of the focal SNP over shorter time scales. This phenomenon of pronounced allele frequency

increase due to linked selection will be especially important if the population size is small (*i.e.,*

drift is not neglectable), and the experimental duration is short. This is reflected in a higher

Nagelkerke's R^2 index for population size in secondary E&R with an experimental duration of

20 generations compared to 60 generations of adaptation (Table S6, Table 2). With increasing

population size and thus, reduced genetic drift, the 1:1 crossing scheme outcompetes dil:mt,

similar to our analysis after 60 generations (Figure S2, Figure S3). Although shorter

experimental duration results in reduced power, our analysis after 20 generations of

adaptation highlights that if the maintenance of experimental populations for many

generations is not feasible, shorter experiments can achieve similar power if they are

maintained at larger population sizes (Figure 3, Figure S2).

<u>Alternative crossing schemes</u>

Since large population sizes can be challenging to maintain, we also evaluated additional

crossing schemes which may perform better even for smaller population sizes (Model 2,

Assessment of experimental and population genetic parameters in Material & Methods).

1:few (Figure 1B) is a modification of the 1:1 crossing scheme, which combines all non-focal

lines in each replicate - this provides a consistent genetic composition in each replicate. The

1:many crossing scheme (Figure 1C) uses 1 focal, and 99 non-focal lines without selection

target. We reasoned that the larger number of segregating variants (970,466 SNPs compared

to max 527,771 SNPs in 1:1 crossing scheme, Table S1) potentially provides a higher mapping

resolution. Finally, we modified the dilution crossing scheme by simulating only one selection

target in the founder population (dil:st, Figure 1D). Hence, no additional selection targets can

create potentially confounding selection signatures. For each experimental design, we

simulated a population size of 300 individuals, 60 generations of adaptation, and 5 replicates

(Table 1).

Regardless of the experimental design, the simulated focal selection targets experienced a

pronounced allele frequency increase (Figure S4). In "dilution" crossing schemes (dil:mt,

dil:st; Figure 1D-E) the frequency trajectories of the focal SNPs were highly variable because

of the heterogeneous starting frequency. The crossing schemes where a single focal line with

a starting frequency of 50 % carries the beneficial allele typically had a superior performance

than dilution crossing schemes (Figure 5). Notably, the significant influence of crossing

scheme on secondary E&R success (Table S7 (success-A), Table S8 (success-B), Table S9)

cannot be explained by a loss of the focal SNP in dilution crossing schemes, which occurred in

less than 1 % of the simulations.

We found that the dil:st crossing scheme is still inferior to the 1:1 crossing scheme for additive

loci (Figure 5). Also, the modifications of the 1:1 crossing scheme (1:few, 1:many) do not

provide a substantial improvement (Figure 5). Surprisingly, the 1:many crossing scheme

performed very poorly. This may be at least partly attributed to the larger number of SNPs/kb,

which will affect the SNP-based success rate (success-B). Given that the three additional

crossing schemes do not provide a clear advantage (Figure 5, Figure 6), and are not easier to

execute experimentally, we did not evaluate them with larger population sizes.

462
Figure 5 Power of five different crossing schemes (population size = 300 individuals; 2,000
simulations/experimental design). Bars show the power (*i.e.,* proportion of successful simulations) separately
for each combination of crossing scheme and dominance coefficient (additive in grey; dominant in white). The
dots with error bars display the estimate from the fitted model and its 95 % confidence interval. For the model
fit, the selection coefficient was fixed to its global average, and combination-specific average starting allele
frequencies were used. (A) shows the results for success-A (= selection target is the SNP with the highest
Cochran-Mantel-Haenszel (CMH) test statistic), (B) shows the results for success-B (= selection target is not more
than 100 SNPs away from the SNP with the highest CMH test statistic).

471
472
473

474
Figure 6 Resolution of five different crossing schemes. Proportion of simulations (y-axis) that do not exceed a
maximum distance in SNPs (x-axis) between the SNP with the highest Cochran-Mantel-Haenszel test statistic
and the true target of selection.

Number of replicates

Given the superior performance of experimental designs with larger population sizes (Figure

3), we were also interested whether more replicates with a smaller population size may

provide further improvements. We performed additional simulations for all five different

crossing schemes with a population size of 300 individuals per replicate, and 30 replicates

(Table 1). Consistent with other studies (23, 25), more replicates result in increased power

(Figure S5). The power of secondary E&R with 30 replicates (Table S10 (success-A), Table S11

(success-B)), is only significantly affected by the crossing scheme (LRT full-null model

comparison (Model 2): $\chi^2 = 37.744$, df = 8, p<0.001 (success-A); $\chi^2 = 16.692$, df = 8, p=0.033

(success-B); Table S12). The 1:1 crossing scheme with 30 replicates also has the highest

resolution (Figure S6A). As expected, the impact of linkage disequilibrium, indicated by the

number of ties, is dramatically reduced for all crossing schemes when more replicates are

simulated (Figure S6B compared to Figure S7). However, similar improvements as seen with

30 replicates can be achieved if 5 replicates of the 1:1 crossing scheme are used with a

population size of 1,200 individuals (Figure 3). Thus, the same power can be achieved while

maintaining about 33 % fewer individuals.

Dominance coefficient, Selection coefficient, and mean starting allele frequency

As expected, dominance coefficient (h), selection coefficient (s), and average starting allele

frequency (af) have a significant effect on the success of secondary E&R in all simulation

scenarios, except for data with 30 replicates. We observed that selection targets with a higher

selection coefficient and/or a higher mean starting allele frequency are easier to fine map.

The lack of significant effects of h, s, and af on secondary E&R success for the dataset with 30

replicates can probably be explained by the smaller number of conducted simulations (100

for 30 replicates and 2,000 for 5 replicates see Table 1). After crossing scheme and population

size, the dominance coefficient is the parameter with the highest Nagelkerke's R^2 index in our

analysis, with additive loci being easier to fine map. This is caused by the fact that

heterozygotes and target homozygotes with a dominant beneficial SNP have the same fitness,

resulting in less efficient selection that becomes especially apparent if the selected allele has

already reached a high frequency in the population, and non-selected allele homozygotes

become rare (indicated by a significant negative effect of the interaction term between

dominance coefficient and mean starting allele frequency in most of our statistical models).

Discussion

This work was inspired by the difficulty of most E&R studies with sexually reproducing

organisms to pinpoint selection targets, mostly due to numerous neutral hitchhikers resulting

in large haplotype blocks. Secondary E&R – a follow-up EE validating putative selection targets

of a primary E&R under an identical selection regime – has been recently suggested as

experimental approach for selection target confirmation (27). We used extensive computer

simulations to evaluate how experimental and population genetics parameter shape the

power and resolution of secondary E&R. As expected, dominance coefficient, selection

strength, and mean starting allele frequency of the selected target all have a significant effect

on the success of E&R, where dominant selection targets at high frequency are particularly

challenging to detect. However, population size and crossing scheme emerged as the most

influential parameters in our analysis.

The crossing scheme has a pronounced effect on secondary E&R success

We show that a simple crossing scheme, which only requires that (a subset of) the founder

lines are sequenced and that founder lines with and without the focal adaptive allele can be

distinguished, has the best power and resolution of the five crossing schemes tested. The 1:1

crossing scheme is particularly well-suited when many selection targets are detected in the

evolved populations of the primary E&R, because it uses only lines from the non-adapted

founder population. Ancestral haplotypes from the non-adapted founder population will

harbor on average less selection targets than evolved haplotypes that can acquire multiple

selection targets through recombination events during the primary E&R (42). Genotypes

harboring multiple selection targets make it challenging to fine-map single selection targets,

because selected alleles will not propagate independently from each other. Nevertheless,

highly polygenic architectures where selected haplotype blocks contain multiple, weakly

selected alleles are not well-suited for secondary E&R, because recombination will eventually

separate linked favorable alleles and no strongly selected allele can be identified.

Sufficient E&R power requires large population sizes

Our analyses also indicate that rather large population sizes are crucial for being able to

identify the causative variant with sufficient confidence using a secondary E&R approach. This

observation explains why an empirical secondary E&R study in *D. simulans* with a population

size of 1,250 flies per replicated population failed to narrow down a pronounced candidate

region from a primary E&R in a secondary E&R after 30 generations, and only confirmed the

presence of selected alleles (27).

Our results suggest that large population sizes become especially crucial for short-term

secondary E&R. The previously reported attenuated power of short-term E&R due to reduced

experimental duration (25) can be increased by a higher experimental population size. Our

analyses show that to achieve satisfactory power after only 20 generations requires

population sizes of tens of thousands of individuals– dimensions that are rather rare in

current EE designs with strictly sexually reproducing organisms, where population sizes are

mostly limited by the capacities of the lab. Similar to previous simulation studies (23, 25), we

demonstrated that increasing the number of replicates results in more powerful secondary

E&R. If maintaining large population sizes is not feasible, maintaining many replicates at small

population size can help to boost secondary E&R performance. Additionally, the loss of one

replicate in a highly replicated setup has a smaller impact than in an experimental setup with

very few replicates. However, we show that additional replication cannot completely

compensate the advantage of large population sizes in secondary E&R.

<u>Choice of model organism and selection regime</u>

Our simulations are parameterized for *D. simulans* because this species is better suited for

E&R studies than *D. melanogaster* (22). While other species, such as *S. cerevisiae* (7) or *C.*

remanei (43) have been used to study adaptive response from standing genetic variation in

outcrossing species, *Drosophila* is the most popular organism. Independent of the actual

experimental parameters used, we anticipate that our results can be generalized to other

species, but the availability of inbred founder lines is probably a more severe restriction,

which limits the wide-spread use of secondary E&R.

Finally, we simulated a rather simple selection scenario of directional selection, not

considering more complex scenarios that include pleiotropy and/or epistasis. While both are

very important factors that could affect the allele frequency changes in an E&R study, we

would like to point out that secondary E&R is designed to study loci, which experienced a

substantial allele frequency increase in a primary experiment, thus pleiotropic constraints are

not expected to be a major confounding factor. Epistasis, on the other hand may be more

pronounced in the 1:1 crossing scheme. It may be possible that the focal allele is only selected

in a subset of the replicates, but not in others-or the response may differ between replicates.

While this reduces the power of the secondary E&R to fine-map the selection target, such a

result may open the possibility to study the impact of epistatic effects. Since the founder lines

will be available, it will be possible to repeat the experiment with replication for each

genotype combination to assure that the heterogeneous response comes from epistatic

effects and is not linked to stochastic changes.

Acknowledgements

We thank the members of the Institute of Population Genetics for fruitful discussion and

support. Thanks to Roger Mundry who shared R scripts for the statistical analyses. Special

thanks to Rupert Mazzucco, Robert Kofler, Kathrin Otte and Christos Vlachos for their support

regarding the computer simulations. Special thanks to Claire Burny, and Robert Kofler for

helpful comments on earlier versions of the manuscript. This work was supported by the

Austrian Science Funds (FWF, W1225, P-32935), and by the European Research Council (ERC)

grant "ArchAdapt".

Authors Contribution

AML, MD, and CS designed the analysis. AML performed the forward simulations, and the

bioinformatic analysis. AML, and MD performed the statistical analysis. AML, MD, and CS

wrote the manuscript.

References

1. T. J. Kawecki, *et al.*, Experimental evolution. *Trends Ecol. Evol.* **27**, 547–560 (2012).

2. T. Garland, M. Rose, *Experimental Evolution: Concepts, Methods, and Applications of Selection Experiments* (2009).

3. C. Schlötterer, R. Kofler, E. Versace, R. Tobler, S. U. Franssen, Combining experimental evolution with next-generation sequencing: a powerful tool to study adaptation from standing genetic variation. *Heredity (Edinb).* **114**, 431–440 (2015).

4. T. L. Turner, A. D. Stewart, A. T. Fields, W. R. Rice, A. M. Tarone, Population-Based Resequencing of Experimentally Evolved Populations Reveals the Genetic Basis of Body Size Variation in Drosophila melanogaster. *PLoS Genet.* **7**, e1001336 (2011).

5. A. Long, G. Liti, A. Luptak, O. Tenaillon, Elucidating the molecular architecture of adaptation via evolve and resequence experiments. *Nat. Publ. Gr.* **16**, 567–582 (2015).

6. A. M. Johansson, M. E. Pettersson, P. B. Siegel, Ö. Carlborg, Genome-Wide Effects of Long-Term Divergent Selection. *PLOS Genet.* **6**, e1001188 (2010).

7. M. K. Burke, G. Liti, A. D. Long, Standing Genetic Variation Drives Repeatable Experimental Evolution in Outcrossing Populations of Saccharomyces cerevisiae. *Mol. Biol. Evol.* **31**, 3228–3239 (2014).

8. A. Papkou, *et al.*, The genomic basis of Red Queen dynamics during rapid reciprocal host–pathogen coevolution. *Proc. Natl. Acad. Sci.* **116**, 923–928 (2019).

9. N. E. Martins, *et al.*, Host adaptation to viruses relies on few genes with different cross-resistance properties. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 5938–43 (2014).

10. H. Teotónio, I. M. Chelo, M. Bradić, M. R. Rose, A. D. Long, Experimental evolution reveals natural selection on standing genetic variation. *Nat. Genet.* **41**, 251–257

627   (2009).

628   11.   S. C. Remolina, P. L. Chang, J. Leips, S. V Nuzhdin, K. A. Hughes, Genomic basis of

629   aging and life-history evolution in Drosophila melanogaster. *Evolution* **66**, 3390–3403

630   (2012).

631   12.   N. Barghi, *et al.*, Genetic redundancy fuels polygenic adaptation in Drosophila. *PLOS*

632   *Biol.* **17**, 1–31 (2019).

633   13.   R. Tobler, *et al.*, Massive Habitat-Specific Genomic Response in *D. melanogaster*

634   Populations during Experimental Evolution in Hot and Cold Environments. *Mol. Biol.*

635   *Evol.* **31**, 364–375 (2014).

636   14.   S. U. Franssen, V. Nolte, R. Tobler, C. Schlötterer, Patterns of Linkage Disequilibrium

637   and Long Range Hitchhiking in Evolving Experimental *Drosophila melanogaster*

638   Populations. *Mol. Biol. Evol.* **32**, 495–509 (2015).

639   15.   S. V Nuzhdin, T. L. Turner, Promises and limitations of hitchhiking mapping. *Curr.*

640   *Opin. Genet. Dev.* **23**, 694–699 (2013).

641   16.   A. Iranmehr, A. Akbari, C. Schlötterer, V. Bafna, CLEAR : Composition of Likelihoods

642   for Evolve and. *Genetics* **206**, 1011–1023 (2017).

643   17.   H. Topa, Á. Jónás, R. Kofler, C. Kosiol, A. Honkela, Gaussian process test for high-

644   throughput sequencing time series: Application to experimental evolution.

645   *Bioinformatics* **31**, 1762–1770 (2015).

646   18.   J. K. Kelly, K. A. Hughes, Pervasive Linked Selection and Intermediate-Frequency

647   Alleles Are Implicated in an Evolve-and-Resequencing Experiment of Drosophila

648   simulans . *Genetics* **211**, genetics.301824.2018 (2018).

649   19.   K. Spitzer, M. Pelizzola, A. Futschik, Modifying the Chi-square and the CMH test for

650   population genetic inference: Adapting to overdispersion. *Ann. Appl. Stat.* **14**, 202–

220 (2020).

20. G. I. Lang, *et al.*, Pervasive genetic hitchhiking and clonal interference in forty

evolving yeast populations. *Nature* **500**, 571–574 (2013).

21. N. Barghi, J. Hermisson, C. Schlötterer, Polygenic adaptation: a unifying framework to

understand positive selection. *Nat. Rev. Genet.* **21**, 769–781 (2020).

22. N. Barghi, R. Tobler, V. Nolte, C. Schlötterer, *Drosophila simulans* : A Species with

Improved Resolution in Evolve and Resequence Studies. *G3:*

Genes|Genomes|Genetics **7**, 2337–2343 (2017).

23. D. Kessner, J. Novembre, Power analysis of artificial selection experiments using

efficient whole genome simulation of quantitative traits. *Genetics* **199**, 991–1005

(2015).

24. J. G. Baldwin-Brown, A. D. Long, K. R. Thornton, The power to detect quantitative trait

loci using resequenced, experimentally evolved populations of diploid, sexual

organisms. *Mol. Biol. Evol.* **31**, 1040–1055 (2014).

25. R. Kofler, C. Schlötterer, A guide for the design of evolve and resequencing studies.

Mol. Biol. Evol. **31**, 474–483 (2014).

26. C. Vlachos, R. Kofler, Optimizing the Power to Identify the Genetic Basis of Complex

Traits with Evolve and Resequence Studies. *Mol. Biol. Evol.* **36**, 2890–2905 (2019).

27. C. Burny, *et al.*, Secondary Evolve and Resequencing: An Experimental Confirmation

of Putative Selection Targets without Phenotyping. *Genome Biol. Evol.* **12**, 151–159

(2020).

28. S. J. Gratz, *et al.*, Genome engineering of Drosophila with the CRISPR RNA-guided

Cas9 nuclease. *Genetics* **194**, 1029–1035 (2013).

29. J. M. Howie, R. Mazzucco, T. Taus, V. Nolte, C. Schlötterer, DNA motifs are not general

predictors of recombination in two drosophila sister species. *Genome Biol. Evol.* **11**,

1345–1357 (2019).

30. C. Vlachos, R. Kofler, MimicrEE2: Genome-wide forward simulations of Evolve and

Resequencing studies. *PLOS Comput. Biol.* **14**, e1006413 (2018).

31. R Core Team 3.5.3, A Language and Environment for Statistical Computing. *R Found.*

Stat. Comput. **2**, https://www.R-project.org (2019).

32. T. Taus, A. Futschik, C. Schlötterer, Quantifying selection with pool-seq time series

data. *Mol. Biol. Evol.* **34**, 3023–3034 (2017).

33. A. Agresti, M. Kateri, "Categorical Data Analysis" in *International Encyclopedia of*

Statistical Science, M. Lovric, Ed. (Springer Berlin Heidelberg, 2011), pp. 206–208.

34. M. Dowle, A. Srinivasan, data.table: Extension of `data.frame` (2019).

35. R. H. Baayen, *Analyzing Linguistic Data: A Practical Introduction to Statistics using R*

(Cambridge University Press, 2008) https:/doi.org/10.1017/CBO9780511801686.

36. H. Schielzeth, Simple means to improve the interpretability of regression coefficients.

Methods Ecol. Evol. **1**, 103–113 (2010).

37. J. Fox, G. Monette, Generalized Collinearity Diagnostics. *J. Am. Stat. Assoc.* **87**, 178–

183 (1992).

38. J. Fox, S. Weisberg, *An R Companion to Applied Regression*, Third (Sage, 2019).

39. N. Nagelkerke, A note on a general definition of the coefficient of determination.

Biometrika **78**, 691–692 (1991).

40. C. Vlachos, *et al.*, Benchmarking software tools for detecting and quantifying

selection in evolve and resequencing studies. *Genome Biol.* **20**, 169 (2019).

41. A. M. Langmüller, C. Schlötterer, Low concordance of short-term and long-term

selection responses in experimental Drosophila populations. *Mol. Ecol.* **29**, 3466–

3475 (2020).

42.   K. A. Otte, C. Schlötterer, Detecting selected haplotype blocks in evolve and

resequence experiments. *Mol. Ecol. Resour.* **21**, 93–109 (2021).

43.   H. Teotónio, S. Estes, P. C. Phillips, C. F. Baer, Experimental evolution with

Caenorhabditis nematodes. *Genetics* **206**, 691–716 (2017).

Supplementary Figures

Figure S1 Violin plots of the number of ties (y axis = $\log_{10}$ (number of ties + 1)) for simulations where the true
target of selection has the highest Cochran-Mantel-Haenszel (CMH) test statistic (success-A). The black
horizontal lines in the violin plots display the median number of observed ties. Each panel shows the results for
one particular simulated population size. Ties are defined as neighboring SNPs that have the same CMH test
statistic as the target of selection.

713
714

Figure S2 Power of the 1:1 and dil:mt crossing scheme at different population sizes (2,000
simulations/experimental design) after 20 generations of adaptation. Bars show the power (*i.e.,* proportion of
successful simulations) separately for each combination of crossing scheme (1:1, dil:mt), population size (300;
1,200; 4,800; 19,200 individuals), and dominance coefficient (additive in grey, dominant in white). The dots with
error bars display the model fit and its 95 % confidence interval. For the model fit, the selection coefficient was
fixed to its global average, and combination-specific average starting allele frequencies were used. A) shows the
results for success-A (= selection target is the SNP with the highest Cochran-Mantel-Haenszel (CMH) test
statistic) (B) shows the results for success-B (= selection target is not more than 100 SNPs away from the SNP
with the highest CMH test statistic).

727
Figure S3 Resolution for the 1:1 and dil:mt crossing scheme simulated with different population sizes after 20
generations of adaptation. (A) Proportion of simulations (y-axis) that do not exceed a maximum distance in SNPs
(x-axis) between the SNP with the highest Cochran-Mantel-Haenszel (CMH) test statistic and the true target of
selection (B) Violin plots of the number of ties (y axis = $\log_{10}$ (number of ties + 1)) for simulations where the true
target of selection has the highest CMH test statistic (success-A). The black horizontal lines in the violin plots
display the median number of observed ties. Ties are defined as neighboring SNPs that have the same CMH test
statistic as the target of selection.

736
Figure S4 Median allele frequency trajectory of focal targets in different experimental designs. For each single
simulation, the focal target frequency of a distinct generation was averaged over five replicates. Solid lines show
the median allele frequency trajectory over 2,000 independent simulations per experimental design. Dashed
lines show the 5 and 95 percentiles of the allele frequency trajectories. Each panel shows the median allele
frequency trajectory of one particular crossing scheme (1:1, dil:mt, 1:few, 1:many, dil:st) and population size
(300; 1,200; 4,800; 19,200 individuals) combination.

Figure S5 Power of five different crossing schemes (population size = 300 individuals; 30 replicates; 100
simulations/experimental design). Bars show the power (*i.e.* proportion of successful simulations) separately for
each combination of crossing scheme and dominance coefficient (additive in grey; dominant in white). The dots
with error bars display the estimate from the fitted model and its 95 % confidence interval. For the model fit,
the selection coefficient was fixed to its global average, and combination-specific average starting allele
frequencies were used. A) shows the results for success-A (= selection target is the SNP with the highest Cochran-
mantel-Haenszel (CMH) test statistic), (B) shows the results for success-B (= selection target is not more than
100 SNPs away from the SNP with the highest CMH test statistic).

752
753

Figure S6 Resolution for five different crossing schemes (population size = 300 individuals; 30 replicates; 100
simulations/experimental design). (A) Proportion of simulations (y-axis) that do not exceed a maximum distance
in SNPs (x-axis) between the SNP with the highest Cochran-Mantel-Haenszel (CMH) test statistic and the true
target of selection (B) Violin plots of the number of ties (y axis = $\log_{10}$ (number of ties + 1)) for simulations where
the true target of selection has the highest CMH test statistic (success-A). The black horizontal lines in the violin
plots display the median number of observed ties. Ties are defined as neighboring SNPs that have the same CMH
test statistic as the target of selection.

765
Figure S7 Violin plots of the number of ties (y axis = $\log_{10}$ (number of ties + 1)) for simulations where the true
target of selection has the highest Cochran-Mantel-Haenszel (CMH) test statistic (success-A, population size =
300 individuals; 5 replicates; 2,000 simulations/experimental design). The black horizontal lines in the violin plots
display the median number of observed ties. Ties are defined as neighboring SNPs that have the same CMH test
statistic as the target of selection.

Supplementary Tables

Table S1 Number of SNPs per crossing scheme on chromosome-arm 2L.

crossing scheme	median	range
1:1	523,008	479,507 – 527,771
1:few	526,510	522,470 – 530,230
1:many	970,466	970,466 – 970,466
dil:mt	970,466	970,466 – 970,566
dil:st	970,466	970,466 – 970,466

Table S2 Type III Sum of Squares of explanatory variables for the analysis of success in secondary Evolve and
Resequence studies after 60 generations of adaptation (success-A; Model 1). cross= crossing scheme; h=
dominance coefficient; s= selection coefficient; af= mean starting allele frequency of the focal target over 5
replicates; N= population size; coss:h = interaction term between crossing scheme and dominance coefficient;
h:af = interaction term between dominance coefficient and mean starting allele frequency; cross:N = interaction
term between crossing scheme and population size ; h:N = interaction term between dominance coefficient and
population size; cross:h:N = interaction term between crossing scheme, dominance coefficient, and population
size.
785

term	χ^2	df	P
cross	117.75	1	<0.001
h	113.81	1	<0.001
s [1]	67.38	1	<0.001
af [1]	0.10	1	0.751
N	1023.53	3	<0.001
cross:h	0.37	1	0.544
h:af	18.51	1	<0.001
cross:N	222.05	3	<0.001
h:N	3.05	3	0.384
cross:h:N	3.78	3	0.286

786
787 [1] multiplied by 100, and z-transformed to mean = 0, and sd =1; Mean/standard deviation of the original value
788 was 0.463/0.130 for the starting allele frequency, and 0.086/0.009 for the selection coefficient respectively

789
790
791
Table S3 Type III Sum of Squares of explanatory variables for the analysis of success in secondary Evolve and
Resequence studies after 60 generations of adaptation (success-B; Model 1). cross= crossing scheme; h=
dominance coefficient; s= selection coefficient; af= mean starting allele frequency of the focal target over 5
replicates; N= population size; coss:h = interaction term between crossing scheme and dominance coefficient;
h:af = interaction term between dominance coefficient and mean starting allele frequency; cross:N = interaction
term between crossing scheme and population size ; h:N = interaction term between dominance coefficient and
population size; cross:h:N = interaction term between crossing scheme, dominance coefficient, and population
size.
800

term	χ^2	df	P
cross	50.81	1	<0.001
h	115.74	1	<0.001
s [1]	73.39	1	<0.001
af [1]	1.95	1	0.162
N	1225.22	3	<0.001
cross:h	4.18	1	0.041
h:af	16.67	1	<0.001
cross:N	426.90	3	<0.001
h:N	6.03	3	0.110
cross:h:N	16.12	3	0.001

801
802 [1] multiplied by 100, and z-transformed to mean = 0, and sd =1; Mean/standard deviation of the original value
803 was 0.463/0.130 for the starting allele frequency, and 0.086/0.009 for the selection coefficient respectively
804

Table S4 Type III Sum of Squares of explanatory variables for the analysis of success in secondary Evolve and
Resequence studies after 20 generations of adaptation (success-A; Model 1). cross= crossing scheme; h=
dominance coefficient; s= selection coefficient; af= mean starting allele frequency of the focal target over 5
replicates; N= population size; coss:h = interaction term between crossing scheme and dominance coefficient;
h:af = interaction term between dominance coefficient and mean starting allele frequency; cross:N = interaction
term between crossing scheme and population size ; h:N = interaction term between dominance coefficient and
population size; cross:h:N = interaction term between crossing scheme, dominance coefficient, and population
size.

term	χ^2	df	P
cross	14.79	1	<0.001
h	6.07	1	0.0138
s [1]	18.21	1	<0.001
af [1]	30.24	1	<0.001
N	1714.41	3	<0.001
cross:h	0.01	1	0.942
h:af	0.02	1	0.898
cross:N	378.43	3	<0.001
h:N	0.64	3	0.887
cross:h:N	1.29	3	0.732

815
816 [1] multiplied by 100, and z-transformed to mean = 0, and sd =1; Mean/standard deviation of the original value
817 was 0.463/0.130 for the starting allele frequency, and 0.086/0.009 for the selection coefficient respectively

818
819
820
Table S5 Type III Sum of Squares of explanatory variables for the analysis of success in secondary Evolve and
Resequence studies after 20 generations of adaptation (success-B; Model 1). cross= crossing scheme; h=
dominance coefficient; s= selection coefficient; af= mean starting allele frequency of the focal target over 5
replicates; N= population size; coss:h = interaction term between crossing scheme and dominance coefficient;
h:af = interaction term between dominance coefficient and mean starting allele frequency; cross:N = interaction
term between crossing scheme and population size ; h:N = interaction term between dominance coefficient and
population size; cross:h:N = interaction term between crossing scheme, dominance coefficient, and population
size.

term	χ^2	df	P
cross	65.33	1	<0.001
h	11.80	1	<0.001
s [1]	58.96	1	<0.001
af [1]	60.39	1	<0.001
N	1789.63	3	<0.001
cross:h	1.60	1	0.206
h:af	0.36	1	0.551
cross:N	657.22	3	<0.001
h:N	1.88	3	0.598
cross:h:N	4.68	3	0.197

830
831 [1] multiplied by 100, and z-transformed to mean = 0, and sd =1; Mean/standard deviation of the original value
832 was 0.463/0.130 for the starting allele frequency, and 0.086/0.009 for the selection coefficient respectively

833
834

835 Table S6 Nagelkerke's R^2-index of each simulation parameter including its interaction in the logistic regression
836 (Model 1; 2 crossing schemes (1:1, dil:mt), 4 population sizes (300; 1,200; 4,800; 19,200 individuals), 20
837 generations, 5 replicates; Figure S2). If the reduced model explains the data equally well as a full model that
838 includes the evaluated effect, Nagelkerke's R^2-index is 0.

effect	R^2 (success-A)	R^2 (success-B)
crossing scheme	0.283	0.295
dominance	0.019	0.021
selection coefficient	0.002	0.007
starting allele frequency	0.005	0.009
population size	0.322	0.211

839
840

841 Table S7 Type III Sum of Squares of explanatory variables for the analysis of success in secondary Evolve and
842 Resequence studies after 60 generations of adaptation (success-A; Model 2). cross= crossing scheme; h=
843 dominance coefficient; s= selection coefficient; af= mean starting allele frequency of the focal target over 5
844 replicates; cross:h= interaction term between crossing scheme and dominance coefficient; h:af= interaction
845 term between dominance coefficient and mean starting allele frequency.

term	χ^2	df	P
cross	197.66	4	<0.001
h	115.11	1	<0.001
s [1]	76.20	1	<0.001
af [1]	11.94	1	<0.001
cross:h	77.19	4	<0.001
h:af	7.56	1	0.006

846
847 [1] multiplied by 100, and z-transformed to mean = 0, and sd =1; Mean/standard deviation of the original value
848 were 0.463/0.133 for the starting allele frequency, and 0.085/0.009 for the selection coefficient respectively

849
850
851
852 Table S8 Type III Sum of Squares of explanatory variables for the analysis of success in secondary Evolve and
853 Resequence studies after 60 generations of adaptation (success-B; Model 2). cross= crossing scheme; h=
854 dominance coefficient; s= selection coefficient; af= mean starting allele frequency of the focal target over 5
855 replicates; cross:h= interaction term between crossing scheme and dominance coefficient; h:af= interaction
856 term between dominance coefficient and mean starting allele frequency.

term	χ^2	df	P
cross	97.39	4	<0.001
h	116.20	1	<0.001
s [1]	72.09	1	<0.001
af [1]	19.69	1	<0.001
cross:h	97.2	4	<0.001
h:af	7.07	1	0.008

858
859 [1] multiplied by 100, and z-transformed to mean = 0, and sd =1; Mean/standard deviation of the original value
860 were 0.463/0.133 for the starting allele frequency, and 0.085/0.009 for the selection coefficient respectively

861
862
863
864 Table S9 Nagelkerke's R^2-index of each simulation parameter including its interaction in the logistic regression
865 (5 crossing schemes, population size = 300 individuals, experimental duration = 60 generations; 5 replicates per
866 simulation; Model 2). If the reduced model explains the data equally well as a full model that includes the
867 evaluated effect, Nagelkerke's R^2-index is 0.

effect	R^2 (success-A)	R^2 (success-B)
crossing scheme	0.061	0.033
dominance	0.095	0.076
selection coefficient	0.002	0.001
starting allele frequency	0.003	0.004

868 Table S10 Type III Sum of Squares of explanatory variables for the analysis of success in secondary Evolve and
869 Resequence studies after 60 generations of adaptation (success-A; Model 2). cross= crossing scheme; h=
870 dominance coefficient; s= selection coefficient; af= mean starting allele frequency of the focal target over 30
871 replicates; cross:h= interaction term between crossing scheme and dominance coefficient; h:af= interaction
872 term between dominance coefficient and mean starting allele frequency.
873

term	χ^2	df	P
cross	13.89	4	0.008
h	1.30	1	0.255
s [1]	2.74	1	0.098
af [1]	2.82	1	0.093
cross:h	5.95	4	0.203
h:af	1.72	1	0.189

874
875 [1] multiplied by 100, and z-transformed to mean = 0, and sd =1; Mean/standard deviation of the original value
876 were 0.463/0.133 for the starting allele frequency, and 0.085/0.009 for the selection coefficient, respectively
877
878
879 Table S11 Type III Sum of Squares of explanatory variables for the analysis of success in secondary Evolve and
880 Resequence studies after 60 generations of adaptation (success-B; Model 2). cross= crossing scheme; h=
881 dominance coefficient; s= selection coefficient; af= mean starting allele frequency of the focal target over 30
882 replicates; cross:h= interaction term between crossing scheme and dominance coefficient; h:af= interaction
883 term between dominance coefficient and mean starting allele frequency.
884

term	χ^2	df	P
cross	9.86	4	0.043
h	3.55	1	0.059
s [1]	3.65	1	0.056
af [1]	0.77	1	0.381
cross:h	5.41	4	0.247
h:af	1.01	1	0.315

885
886 [1] multiplied by 100, and z-transformed to mean = 0, and sd =1; Mean/standard deviation of the original value
887 were 0.463/0.133 for the starting allele frequency, and 0.085/0.009 for the selection coefficient, respectively
888
889
890 Table S12 Nagelkerke's R^2-index of each simulation parameter including its interaction in the logistic regression
891 (5 crossing schemes, population size = 300 individuals, experimental duration = 60 generations; 30 replicates
892 per simulation; Figure S5; Model 2). If the reduced model explains the data equally well as a full model that
893 includes the evaluated effect, Nagelkerke's R^2-index is 0.

effect	R^2 (success-A)	R^2 (success-B)
crossing scheme	0.116	0.046
dominance	0.046	0.034
selection coefficient	0.009	0.010
starting allele frequency	0.010	0.003