Fine mapping without phenotyping: Identification of selection targets in secondary Evolve

and Resequence experiments

Anna Maria Langmüller[1,2], Marlies Dolezal[3], Christian Schlötterer[1,*]

[1] Institut für Populationsgenetik, Vetmeduni Vienna, Veterinärplatz 1, 1210 Vienna, Austria

[2] Vienna Graduate School of Population Genetics, Vetmeduni Vienna, Veterinärplatz 1, 1210

Vienna, Austria

[3] Plattform Bioinformatik und Biostatistik, Vetmeduni Vienna, Veterinärplatz 1, 1210 Vienna,

Austria

*Author of Correspondence:

Christian Schlötterer, Institut für Populationsgenetik, Vetmeduni Vienna, Veterinärplatz 1,

1210 Vienna, Austria

+43 1 25077-4301, +43 1 25077-4390

christian.schloetterer@vetmeduni.ac.at

Abstract

Evolve and Resequence (E&R) studies investigate the genomic selection response of populations in an Experimental Evolution setup. Despite the popularity of E&R, empirical studies in sexually reproducing organisms typically suffer from an excess of candidate loci due to linkage disequilibrium, and single gene or SNP resolution is the exception rather than the rule. Recently, so-called "secondary E&R" has been suggested as promising experimental follow-up procedure to confirm putatively selected regions from a primary E&R study. Secondary E&R provides also the opportunity to increase mapping resolution by allowing for additional recombination events, which separate the selection target from neutral hitchhikers. Here, we use computer simulations to assess the effect of different crossing schemes, population size, experimental duration, and number of replicates on the power and resolution of secondary E&R. We find that the crossing scheme and population size are crucial factors determining power and resolution of secondary E&R: a simple crossing scheme with few founder lines consistently outcompetes crossing schemes where evolved populations from a primary E&R experiment are mixed with a complex ancestral founder population. Regardless of the experimental design tested, a population size of at least 4,800 individuals, which is roughly 5 times larger than population sizes in typical E&R studies, is required to achieve a power of at least 75%. Our study provides an important step towards improved experimental designs aiming to characterize causative SNPs in Experimental Evolution studies.

Keywords

Experimental Evolution, secondary Evolve and Resequence, experimental design, *Drosophila*, fine mapping

Significance

Despite the popularity of Evolve and Resequence (E&R) to investigate genomic selection responses, most studies that use sexually reproducing organisms have broad selection signatures and an excess of candidate loci due to linkage disequilibrium. In this study, we use computer simulations and statistical modelling to evaluate the effects of different experimental and population genetic parameters on the success of potential follow-up experiments (=secondary E&R) aiming to validate and fine-map selection signatures of primary studies. We found that a large population size in combination with a simple crossing scheme is key to the success of secondary E&R in *Drosophila*.

Introduction

Deciphering the genetic architecture of adaptation is one of the longstanding goals in evolutionary biology. Experimental Evolution (EE) has become a popular approach to study adaptation in real time (Garland & Rose 2009; Kawecki et al. 2012). In contrast to natural populations, EE offers the key advantage of replicating experiments under controlled laboratory conditions (Schlötterer et al. 2015). Evolve and Resequence (E&R) (Turner et al. 2011; Long et al. 2015; Schlötterer et al. 2015) – a combination of EE with Next Generation Sequencing – facilitates in-depth analysis of the genomic responses to selection, with the ultimate goal to identify and characterize individual adaptive loci.

E&R has already been successful in investigating genomic selection responses from standing genetic variation in adapting sexually reproducing organisms, such as chicken (Johansson et al. 2010), yeast (Burke et al. 2014), and *Drosophila* (Teotónio et al. 2009; Remolina et al. 2012; Martins et al. 2014; Barghi et al. 2019). Despite its popularity, E&R typically suffers from an excess of candidates caused by linkage disequilibrium between true

causative SNPs and neutral hitchhikers (Nuzhdin & Turner 2013; Tobler et al. 2014; Franssen et al. 2015), which decreases the resolution of E&R studies and makes single gene resolution (Martins et al. 2014) the exception rather than the rule.

The problem of candidate excess in E&R studies has been approached from different angles. More refined statistical tests have been developed (Topa et al. 2015; Iranmehr et al. 2017; Kelly & Hughes 2019; Spitzer et al. 2020), and the combination of time-series data with replicate populations has been identified as particularly powerful (Lang et al. 2013; Burke et al. 2014; Barghi et al. 2020). Organisms with a higher recombination rate and a lack of large segregating inversions that suppress recombination events have been suggested to be better suited for E&R studies (Barghi et al. 2017). Computer simulations showed that the power of E&R studies can be significantly improved by increasing the number of replicate populations, the experimental duration, or by adjusting the applied selection regime (Baldwin-Brown et al. 2014; Kofler & Schlötterer 2014; Kessner & Novembre 2015; Vlachos & Kofler 2019).

Burny et al. (2020) recently suggested an experimental follow-up procedure ("secondary E&R") to validate selection signals of primary E&R studies. The basic idea of secondary E&R is that putative selection targets determined in the primary E&R study should rise in frequency again when exposed to the same environmental conditions during an additional E&R conducted after the primary experiment (Figure 1A). This experimental validation of selection signals is especially attractive before starting the time-consuming functional characterization of putatively selected alleles (e.g. based on the CRISPR/Cas technology) (Gratz et al. 2013). A hitherto under-explored potential of secondary E&R is that the additional recombination events during the secondary E&R can be used to fine map selection signals of primary experiments. In addition to mixing evolved genotypes of a primary

E&R with non-adapted ancestral founder genotypes (coined "dilution" by Burny et al. (2020)),

we propose several different secondary E&R crossing schemes for validating and fine-mapping

of putative selection targets.

We evaluate the power and resolution of different secondary E&R designs to identify

causative SNPs via extensive computer simulations. We use logistic regression to assess which

simulated experimental and population genetic parameters have a significant effect on the

success of secondary E&R. Selection coefficient, dominance coefficient, and mean starting

allele frequency of the selection target all have a significant effect on the success of secondary

E&R. However, crossing scheme and population size emerge as the most influential parameters.

We show that the population size of secondary E&R experiments needs to be at least 5 times

larger than currently used population sizes in typical primary E&R studies with *Drosophila* to

achieve a power above 75%. Furthermore, we show that the crossing scheme is a crucial

experimental parameter shaping the power of secondary E&R - a simple crossing scheme with

few founder lines results in higher power and resolution compared to more complex crossing

schemes.

Material & Methods

<u>Outline of the simulation framework</u>

The non-adapted ancestral founder genotypes used in our simulation study are a randomly

chosen subset of 100 haplotypes from a panel of 189 sequenced *D. simulans* haplotypes

originally collected in Tallahassee (Florida, USA) capturing the amount of standing genetic

variation in a natural *Drosophila* population (Howie et al. 2019; Barghi et al. 2019). We use

the term "founder line" for an inbred isofemale line homozygous for one of these ancestral

haplotypes. In order to speed up the calculations, we only simulated chromosome-arm 2L.

Simulations with linkage were conducted with MimicrEE2 (v206) (Vlachos & Kofler 2018)

using the *D. simulans* recombination map (Howie et al. 2019). MimicrEE2 is a forward-

simulation framework for E&R studies that can simulate evolving experimental populations

based on their haplotype information and genome-wide recombination rates. We used the w-

mode of MimicrEE2 which computes the fitness of individuals directly from the selection

coefficients. If not stated otherwise, we simulated positive, additive selection for biallelic SNPs,

with selection coefficients being uniformly sampled between 0.07 and 0.1 for each positively

selected SNP and tracked the frequency of all SNPs over time (ranging from 479,507 to 970,466

SNPs, Table S1). We chose to simulate rather strong selection reasoning that alleles with a high

selection coefficient are more likely to be experimentally tested. On the other hand, strongly

selected alleles result in many neutral linked hitchhikers producing false positive signals that

adversely impact the mapping resolution (Kofler & Schlötterer 2014) - which requires follow-

up studies to identify the target of selection. If not stated otherwise, selected SNPs were

randomly chosen with an equal probability to be either co-dominant (dominance coefficient

$h=0.5$), or dominant ($h=1$). We did not consider recessive loci ($h=0$), because we do not

anticipate that fully recessive targets would result in sufficiently large allele frequency changes

to be detected in primary E&R experiments (Baldwin-Brown et al. 2014; Kofler & Schlötterer

2014). Similar to Baldwin-Brown et al. (2014), we did not model allele frequency estimation

errors caused for example by sequencing errors, limited read depth, read depth heterogeneity

across the chromosome, or the number of sequenced individuals. We used PoPoolation2 (Kofler

et al. 2011) to rescale allele counts for each biallelic position to a uniform read depth of 80.

Experimental parameters

The purpose of this study is to test the influence of different experimental parameters on the

power and resolution of secondary E&R. For this, we systematically varied the crossing

scheme, population size, experimental duration, and number of replicates to assess the effect of

these experimental parameters on the power and resolution of secondary E&R (Table 1).

We use the term "experimental design" to describe a distinct set of simulated experimental

parameters (e.g., crossing scheme=1:1_1f; population size=1,200 individuals; experimental

duration=60 generations; 5 replicates).

Crossing scheme

We simulated five different crossing schemes: 1:1 (versions 1f, 2f and 1f1nf), 1:few, 1:many,

dil:st, and dil:mt (Figure 1B-F). In the **1:1_1f** crossing scheme (Figure 1B), two inbred founder

lines are crossed at equal proportions. One inbred focal line carries a single target of selection

– known from a primary E&R experiment – and is crossed with inbred non-focal lines, not

carrying known adaptive alleles. We consider this a best-case scenario. Replicates are created

by crossing the same focal line to different non-focal lines without known targets of selection.

We used different non-focal lines as crossing partners to account for the possibility that in real

experiments these lines may contain unidentified selected loci. By using different lines, the

influence of selection targets present in a single non-focal line will be outweighed by the focal

locus, which is present in every replicate (Figure 1B). To further explore the influence of

unidentified selected loci, we simulated two more versions of the 1:1 crossing scheme with a

more realistic genetic architecture (Figure S1). In these versions, either the focal line itself, or

one of the non-focal lines carries one additional target of selection. We call these scenarios 2f,

for a total of 2 selected loci in the focal line (Figure S1B), and 1f1nf, for one selected locus in

the focal line and one selected locus in one of the non-focal lines (Figure S1C). For each

simulation, we sampled the selection coefficient of the additional selection target uniformly

between 0.07 and the selection coefficient of the selected SNP we consider for our analysis.

We simulated all possible combinations of dominance coefficients for the two selected SNPs

$(0.5 - 0.5; 0.5 - 1; 1 - 0.5; 1 - 1)$. We ran 500 simulations for each of the 4 combinations of

dominance coefficients for 2f and 1f1nf, respectively.

In the **1:few** crossing scheme (Figure 1C) the focal line is crossed with a pool of five

non-focal lines that do not carry known beneficial alleles. The starting frequency of the focal

line is 50 %, whereas each non-focal line has a starting frequency of 10 %. Each replicate

consists of the same focal/non-focal line mixture. In the **1:many** crossing scheme (Figure 1D),

the focal line is crossed with a pool of 99 non-focal lines. Replicates consist of the same mixture

of lines, and the starting frequency of the focal line is 50 %.

It has been recently suggested, that "diluting" evolved populations of a primary E&R

experiment with many non-adapted ancestral genotypes of the very same primary E&R and

exposing the diluted populations to the same selection regime is a promising approach to

validate selection candidates (Burny et al. 2020). However, we lack a systematic power

assessment of such experiments with computer simulations. We thus included two "dilution"

crossing schemes (**dil:mt, dil:st**) into our analysis. To evaluate the power of dilution crossing

schemes, it is important to first simulate a primary E&R study. We chose to simulate a

population consisting of 100 different founder lines, a population size of 300 individuals, 60

generations of adaptation and one replicate, which can then be diluted with non-adapted

ancestral genotypes. SNPs in the primary E&R were tested for allele frequency change with the

χ^2 test.

The population of the **dil:st** crossing scheme carries only one beneficial SNP (**dil:st** =

dilution: **s**ingle **t**arget, Figure 1E). The starting allele frequency of this beneficial, focal SNP is

sampled from the empirical starting allele frequency distribution of putatively selected alleles

from a previous E&R study in which *D. simulans* populations adapted to a new temperature

regime (mean starting frequency = 0.1) (Barghi et al. 2019). After simulating 60 generations of

adaptation (primary E&R), 50 % of the evolved population is replaced by flies of the non-

adapted ancestral founder population. After this dilution step, the secondary E&R was

simulated under the exact same selection regime as in the primary E&R. A region with a strong

selection signal from the primary E&R is chosen for validation in the dilution crossing schemes

(Burny et al. 2020). Hence, we investigated a 1 Mb window, which is the previously reported

median selected haplotype block length on chromosome-arm 2L (Barghi et al. 2019), around

the SNP with the highest χ^2 test-statistic in the primary E&R, in the secondary E&R.

The **dil:mt** (Figure 1F) crossing scheme has multiple selection targets (**dil:mt** =

dilution: **m**ultiple **t**argets; 16 selection targets on chromosome-arm 2L (Barghi et al. 2019)).

Again, we investigate a 1 Mb window around the SNP with the highest χ^2 test statistic in the

primary E&R, in the secondary E&R. In case of multiple selection targets in the 1 Mb window,

we consider the target that is closest to the SNP with the highest CMH test statistic in the 1Mb

window of the secondary E&R in our analysis.

Figure 1. Basic idea of secondary Evolve and Resequence (E&R) and simulated crossing schemes. (A) The basic idea of secondary E&R is that a putative selection target (purple) determined in a primary E&R study (left) should rise in frequency again when exposed to the same environment (i.e., selection regime), during secondary E&R (right). Additional recombination events during the secondary E&R allow to fine map selected regions of primary E&R experiments (*i.e.,* reduce the number hitchhikers indicated in light pink) The two arrows indicate that the secondary E&R is either started with evolved haplotypes (dashed arrow) or with specific founder lines of the primary E&R (dotted arrow). (B) **1:1_1f** crossing scheme: Inbred flies with one target of selection (purple) are crossed to inbred flies without known beneficial variants. The starting frequency of each genotype is 50 %. In each replicate the line with the beneficial allele (focal line, purple) is crossed to a different line lacking beneficial mutations (non-focal lines, different shades of grey). (C) **1:few** crossing scheme: The focal line is crossed to a pool of flies with five different genotypes without known selection targets. The starting frequency of the focal line is again 50 %. (D) **1:many** crossing scheme: The focal line is crossed to a pool of 99 lines without known selection targets. (E) **dil:st** crossing scheme: 50 % of an evolved population originating from a primary E&R is replaced by ancestral genotypes of the primary E&R. The entire population has only one single target of selection (focal SNP, purple flies) (F) **dil:mt** crossing scheme: 50 % of an evolved population originating from a primary E&R is replaced by ancestral genotypes of the primary E&R. The ancestral population carries 16 targets of selection (flies carrying different beneficial SNPs are shown in purple, and green).

Population size

For two crossing schemes that are relatively easy to implement in empirical studies 1:1 (1f, 2f

and 1f1nf) and dil:mt we performed simulations (experimental duration = 60 generations; 5

replicates) with independently sampled selection targets for population sizes of 300; 1,200;

4,800; and 19,200 individuals per replicate to test for the effect of the population size on the

power and resolution of secondary E&R (Table 1). All other crossing schemes were evaluated

at a population size of 300 individuals.

Experimental duration

234 All crossing schemes were evaluated after 60 generations. To explore the possibility that shorter

235 experiments with less than 60 generations may already be sufficient to achieve satisfactory

236 power, we analyzed the simulations for 1:1 (without 2f and 1f1nf) and dil:mt crossing schemes

237 also already after 20 generations (Table 1).

Number of replicates

240 To assess the impact of the number of replicates on power of secondary E&R, we conducted

241 for each crossing scheme (1:1 without 2f and 1f1nf) 100 additional simulations (population

242 size= 300 individuals; experimental duration = 60 generations) with 30 replicates (Table 1).

Table 1. Simulation overview. 1:1_1f, 1:few, 1:many, dil:st, dil:mt: For each selected SNP, selection coefficients were uniformly sampled between 0.07 and 0.1. Dominance coefficients were randomly chosen to be either 0.5 (co-dominant) or 1 (dominant). 1:1_2f, 1:1_1f1nf: For each simulation, the selection coefficient of the target of interest was uniformly sampled between 0.07 and 0.1. In contrast to 1:1_1f, we simulated an additional selection target, which was either located on the focal haplotype (1:1_2f) or one non-focal haplotype (1:1_1f1nf). The selection coefficient of the additional selection target was uniformly sampled between 0.07 and the selection coefficient of the target of interest. We simulated all possible combinations of dominance coefficients for the two selected SNPs (500 simulations per combination).

Crossing scheme	Population size	Generations	Replicates	Number of simulations
1:1_1f	300; 1,200; 4,800; 19,200	60; 20	5/30	2,000/100
1:1_2f	300; 1,200; 4,800; 19,200	60	5	2,000
1:1_1f1nf	300; 1,200; 4,800; 19,200	60	5	2,000
1:few	300	60	5/30	2,000/100
1:many	300	60; 20	5/30	2,000/100
dil:st	300	60	5/30	2,000/100
dil:mt	300; 1,200; 4,800; 19,200	60; 20	5/30	2,000/100

Statistical analysis

All statistical analyses were performed using the R statistical computing environment (v3.5.3) (R Core Team 3.5.3 2019).

Power

For each simulation, we tested all SNPs on chromosome-arm 2L for allele frequency increase between the start and the end of the simulated secondary E&R using the Cochran-Mantel-Haenszel-test (CMH-test, implemented in the R package poolSeq (v0.3.2) (Taus et al. 2017)). The CMH-test allows to test for independence of matched data – e.g. allele counts of replicated ancestral and evolved populations (Agresti & Kateri 2011). We ranked SNPs based on their CMH test statistic using the dense ranking method in the R package data.table (v.1.12-8)

(Dowle & Srinivasan 2019). In dense ranking, SNPs with identical test statistics receive the

same rank, and the following SNP is assigned the immediately following rank.

Based on this ranking, we used two different approaches to classify simulations being

either successful, or unsuccessful. First, we only considered a simulation to be successful if the

true target of selection (i.e., the focal SNP with a selection coefficient > 0) was the SNP with

the highest test statistic (success-A). In a second analysis step, we classified a simulation as

success, if the focal target of selection was not more than 100 SNPs away from the SNP with

the highest test statistic (success-B, Figure 2). We acknowledge that success-B depends on the

maximum distance allowed between the true target of selection and the SNP with the highest

CMH test statistic (Figure 2). However, varying the maximum distance threshold did not alter

the relative performance of different experimental designs (data not shown). The power of an

experimental design is defined as the proportion of simulations that were able to detect the true

target of selection.

Figure 2. Schematic overview of the definition of success-B in a secondary Evolve and Resequence simulation.
All SNPs (neutral = circle, beneficial = diamond) are tested for an allele frequency change with the Cochran-
Mantel-Haenszel-test, and are ranked based on their test statistic (y-axis). If the focal selection target (purple
diamond; one additional selected SNP is shown as green diamond - for crossing scheme 1:1_2f, 1:1_1fnf, and
dil:mt) is less than 100 SNPs away from the SNP with the highest test statistic, the simulation run is deemed a
success. In the example depicted, the distance in number of SNPs between the focal target of selection, and the
SNP with the highest test statistic is 4, and the simulation is classified as success.

Resolution

For simulations where the selection target was detected (success-B), we determined the

resolution of fine mapping of the selection target by counting the number of SNPs between the

true selection target, and the SNP with the highest CMH test statistic.

Assessment of experimental and population genetic parameters

We used logistic regressions with a binomial error structure (ε) and a logit link function

(Baayen 2008) to test how different experimental and population genetic parameters affect

success and failure to identify targets of selection, i.e. success (Y) is treated as a binary response

encoded in 0 (failure) and 1 (success) of secondary E&R.

We fitted three different models with R function *glm* with μ being the overall mean per

model: Model 1 includes only the two crossing schemes 1:1_1f and dil:mt for which we also

varied population size. Model 2 includes the three versions of crossing scheme 1:1 (1f, 2f and

1f1nf) at varying population sizes: 1f with only one positively selected SNP in the focal line;

version 2f with two selected SNPs in the focal line; and version 1f1nf with one selected SNP

in the focal line and one selected SNP in one non-focal line. Model 3 includes all 5 simulated

crossing schemes (1:1 without 2f and 1f1nf) at a constant population size of 300 individuals.

Prior to model fitting, the two covariates selection coefficient and mean starting allele

frequency (not applicable to model 2 as starting allele frequency is always 50%) were

multiplied by 100, and z-transformed to a mean of zero and standard deviation of one for easier

interpretable estimates (Schielzeth 2010).

(1) $Y_{ijklmn} = \mu + cross_i + h_j + s_k + af_l + N_m + (cross{:}h)_{ij} + (h{:}af)_{jl} +$

$(cross{:}N)_{im} + (h{:}N)_{jm} + (cross{:}h{:}N)_{ijm} + \varepsilon_{ijklmn}$

(2) $Y_{ijkmn} = \mu + architecture_i + h_j + s_k + N_m + (architecture{:}h)_{ij} +$

$(architecture{:}N)_{im} + (h{:}N)_{jm} + (architecture{:}h{:}N)_{ijm} + \varepsilon_{ijkmn}$

(3) $Y_{ijklm} = \mu + cross_i + h_j + s_k + af_l + (cross{:}h)_{ij} + (h{:}af)_{jl} + \varepsilon_{ijklm}$

Model 1 (equation (1)) contained (i) five explanatory variables as main effects; crossing

scheme ($cross_i$), a fixed categorical effect with 2 levels - 1:1_1f and dil:mt; a fixed categorical

effect of dominance coefficient (h_j) with levels 0.5 and 1; selection coefficient (s_k) and mean

starting allele frequency (af_l) over replicated populations (both as continuous covariate); and

population size (N_m), a fixed categorical effect with 4 levels - 300; 1,200; 4,800; 19,200, (ii) an interaction term between dominance coefficient and mean starting allele frequency (($h: af)_{jl}$), (iii) and a triple interaction between crossing scheme, dominance coefficient, and population size (($cross: h: N)_{ijm}$), and all pairwise interaction terms of effects involved in the triple interaction term (($cross: h)_{ij}$; ($cross: N)_{im}$; ($h: N)_{jm}$). We included interaction terms into the model that have population genetic interpretations.

Data analyzed with model 1 contained 16,000 observations, namely 2,000 independent simulation runs for each crossing scheme (1:1_1f, dil:mt), and each of the four different population sizes (300; 1,200; 4,800; 19,200) (Table 1). To avoid potential bias introduced by specific haplotypes being sampled, we randomly chose 4 sets of focal/non-focal founder lines for the 1:1_1f crossing scheme and performed 500 simulation runs per set. The chosen set of founder lines did not have a significant effect on the success of secondary E&R and is thus not included in the final model (likelihood ratio test (LRT) full-reduced model comparison; data not shown).

Model 2 (equation (2)) contained (i) four explanatory variables as main effects; a fixed categorical effect "architecture" ($architecture_i$) that describes the version of the 1:1 crossing scheme in combination with the dominance coefficient of the additional selected target (if present) resulting in 5 levels: 1f; 2f_h05; 2f_h1; 1f1nf_h05; 1f1nf_h. Model 2 further contained a fixed categorical effect of dominance coefficient for the focal SNP (h_j) with levels 0.5 and 1; selection coefficient (s_k) as continuous covariate; and population size (N_m), a fixed categorical effect with 4 levels: 300; 1,200; 4,800; 19,200, (ii) a triple interaction between architecture, dominance coefficient, and population size (($architecture: h: N)_{ijm}$), and all pairwise

interaction terms of effects involved in the triple interaction term $((architecture{:}h)_{ij};$

$(architecture{:}N)_{im}; (h{:}N)_{jm}).$

Data analyzed with model 2 contained 24,000 observations, namely 2,000 independent

simulation runs for each version of the 1:1 crossing scheme (1f, 2f, 1f1nf), and four different

population sizes (300; 1,200; 4,800; 19,200 individuals). Because model 1 showed that the

chosen set of founder lines did not have a significant effect on the power of secondary E&R,

we simulated only one set of focal/non-focal founder lines.

Model 3 (equation (3)) contained (i) four explanatory variables; crossing scheme $(cross_i)$,

a fixed categorical effect with 5 levels: 1:1_1f; 1:few; 1:many; dil:st; dil:mt, dominance

coefficient (h_j), selection coefficient (s_k), and mean starting allele frequency (af_l) over

replicated populations as main effects, as described for model 1, (ii) an interaction term between

crossing scheme and dominance coefficient $((cross{:}h)_{ij})$, (iii) and an interaction term between

dominance coefficient and mean starting allele frequency $((h{:}af)_{jl}).$

We analyzed 10,000 samples (2,000 independent simulation runs for each crossing scheme)

(

Table *1*). For crossing schemes using only few different founder lines (1:1_1f, 1:few), the

simulation runs are based on 4 randomly chosen sets of founder lines each (500 simulation runs

per set). As in model 1, the set of founder lines does not have a significant effect on secondary

E&R success and is thus not included in the final model (LRT full-reduced model comparison;

data not shown). We performed additional analysis with 500 observations (100 independent

simulation runs for each crossing scheme) to determine the influence of the number of replicates

on the power of secondary E&R (Table 1).

We performed all diagnostic checks required for logistic regression. Absence of collinearity

was confirmed by computing the generalized Variance Inflation Factors (Fox & Monette 1992)

using function *vif* in R package *car* (v3.0-8 (Fox & Weisberg 2019)). Model stability was

checked with the R function *dfbeta*. For visualization, linear predictors (LP) were back-

transformed to success probabilities using the inverse logit transformation: $p_{sucess} = \frac{e^{LP}}{1+e^{LP}}$. 95

% confidence intervals of the fitted values were investigated with the R function *predict.glm*.

Significance of single explanatory variables was tested with a Type II ANOVA using function

Anova in R package car (Fox & Weisberg 2019) and are provided in the Supplement.

Significance of explanatory variables including all their modeled interactions was tested with a

likelihood ratio test comparing the full model with a nested reduced model with the same

structure as the full model, but lacking the assessed explanatory variable (and its interactions).

Significance is declared at an alpha cut-off of 5%. We used Nagelkerke's R^2 index (Nagelkerke

1991) to calculate the improvement of each model parameter upon the prediction of a reduced

model.

We observed two cases where a combination of explanatory variables resulted in complete

separation of data points (Figure 5A (dominance coefficient: 0.5, population size: 1,200,

architecture: 2f_h05), Figure S8A (dominance coefficient: 0.5, crossing scheme: dil:mt)). To

obtain interpretable model estimates, we added one pseudo-observation with the missing

response (success) to the data for each of these two cases.

Data availability

Information regarding the accessibility of raw sequence reads, phased haplotypes of the

ancestral *D. simulans* haplotypes as well as MimicrEE2 ready text files for the *D. simulans*

recombination map (file used in this project: Dsim_recombination_map_LOESS_100kb_1.txt)

can be found in (Howie et al. 2019). MimicrEE2 ready input files of the different experimental

designs, the simulated selection regimes, processed simulation results and all scripts that are

necessary to reproduce the results are available at SourceForge

(https://sourceforge.net/projects/secondary-e-r-sim/files/).

Results

We used forward simulations to assess the influence of different experimental and population

genetic parameters, more specifically crossing scheme, population size, dominance coefficient,

selection strength, and mean starting allele frequency on the success to detect and fine map

selection targets in secondary E&R experiments (Figure 1A). We used a Cochran-Mantel-

Haenszel (CMH) test to identify SNPs rising in allele frequency (number of SNPs see Table

S1). The CMH test allows to test for independence of matched categorical data (Agresti &

Kateri 2011), and compares favorably to other statistical methods in reliably identifying

possible targets of selection in E&R setups (Vlachos et al. 2019). A simulation run was

considered successful, if the true target of selection was the SNP with the highest CMH test

statistic (success-A). In a second analysis step, we considered simulation runs as successful, if

the true target of selection was not more than 100 SNPs away from the SNP with the highest

CMH test statistic (success-B, Figure 2). We used logistic regression to assess which experimental and population genetic parameters have a significant effect on the success of secondary E&R.

First, we evaluated two crossing schemes that can be easily implemented in empirical studies, 1:1_1f (Figure 1B) and dil:mt (Figure 1F). The 1:1_1f crossing scheme (Figure 1B) is based on founder lines only (a founder line is an inbred isofemale line homozygous for one ancestral haplotype). Using information about the selected haplotype from a primary E&R study, it is possible to determine which founder lines carry a selection target (Barghi et al. 2019). Note, this requires focal founder lines to be sequenced. Crossing a focal founder line with the selected haplotype with a non-focal line without known selection targets offers the advantage of reducing the number of selection targets dramatically. Using different non-focal lines without known strong selection targets in each replicate reduces the potential of consistent confounding effects of unidentified selection targets outside the region of interest – a signal the CMH test is particularly sensitive to as it scans for consistent allele frequency changes across replicates.

Dil:mt (Figure 1F) represents an entirely different approach. Dil:mt has multiple targets of selection at different frequencies and is probably the simulated crossing scheme with the most straight forward empirical implementation (Barghi et al. 2019; Burny et al. 2020). It is based on a "dilution" approach, where evolved individuals from a primary E&R experiment are crossed to the non-adapted ancestral founder population from the same primary E&R (Burny et al. 2020). In contrast to 1:1_1f, a dil:mt crossing scheme requires both – ancestral and evolved – populations of the primary E&R, but relatively limited information about selection targets on individual founder lines.

To evaluate the 1:1_1f and dil:mt crossing scheme, we simulated secondary E&R consisting of 5 replicated populations with constant population size (300; 1,200; 4,800, or 19,200 individuals per replicate) that evolve for 60 generations. We simulated positively selected SNPs (selection coefficient is uniformly sampled between 0.07 and 0.1) that were randomly chosen with equal probability to be either co-dominant (h=0.5) or dominant (h=1). We used logistic regression to assess the effects of model parameters on secondary E&R success (Model 1, *Assessment of experimental and population genetic parameters* in Material & Methods). While selection strength (LRT full-reduced null model comparison (Model 1): $\chi^2 = 67.4$, df = 1, p<0.001 (success-A); $\chi^2 = 73.4$, df = 1, p<0.001 (success-B)), dominance coefficient (LRT full-reduced model comparison (Model 1): $\chi^2 = 733.4$, df = 9, p<0.001 (success-A); $\chi^2 = 714.5$, df = 9, p<0.001 (success-B)), and mean starting allele frequency (LRT full-reduced model comparison (Model 1): $\chi^2 = 22.4$, df = 2, p<0.001 (success-A); $\chi^2 = 17.4$, df = 2, p<0.001 (success-B)) all have a significant effect on the power of secondary E&R (Table S2), our analysis reveals that crossing scheme and population size have by far the strongest influence on secondary E&R success. Both the crossing scheme (LRT full-reduced model comparison (Model 1): $\chi^2 = 3510.1$, df = 8, p<0.001 (success-A); $\chi^2 = 3211.3$, df = 8, p<0.001 (success-B)); and the population size (LRT full-reduced model comparison (Model 1): $\chi^2 = 2142.9$, df = 12, p<0.001 (success-A); $\chi^2 = 2633.4$, df = 12, p<0.001 (success-B)) have a significant effect on the success of secondary E&R, and are the only parameters with a Nagelkerke's R^2 index above 0.2 (Table S3).

While a 1:1_1f crossing scheme results in higher power values than the dil:mt crossing scheme independently of the population size, the difference to the dil:mt crossing scheme is more pronounced in larger experimental populations (Figure 3). This does not hold only for the

power, but also for the resolution (= the distance in SNPs between the true target of selection,

and the SNP with the highest CMH test statistic, Figure 4). A potential disadvantage of the

1:1_1f crossing scheme is that the low number of different founder lines (n=6, Figure 1B)

causes more linkage disequilibrium, indicated by the number of neighboring SNPs with

identical test statistics (=ties) and thus broadens signatures compared to dil:mt crossing scheme

that has more founder lines (n=100) (Figure S2). However, this is outweighed by superior

power of the 1:1_1f crossing scheme at every population size investigated (Figure 3-4).

Figure 3. Power of the 1:1_1f and dil:mt crossing scheme at different population sizes (2,000
simulations/experimental design). Bars show the power (i.e., the proportion of successful simulations) separately
for each combination of crossing scheme (1:1_1f, dil:mt), population size (300; 1,200; 4,800; 19,200 individuals),
and dominance coefficient (additive in grey, dominant in white). The dots with error bars display the estimate from
the fitted model (Model 1) and its 95 % confidence interval. For the model fit, the selection coefficient was fixed
to its global average, and combination-specific average starting allele frequencies were used. (A) shows the results
for success-A (= selection target is the SNP with the highest Cochran-Mantel-Haenszel (CMH) test statistic), (B)
shows the results for success-B (= selection target is not more than 100 SNPs away from the SNP with the highest
CMH test statistic).

Figure 4. Resolution of the 1:1_f and dil:mt crossing scheme at different population sizes. Proportion of simulations (y-axis) that do not exceed a maximum distance in SNPs (x-axis) between the SNP with the highest Cochran-Mantel-Haenszel test statistic and the true target of selection. Each panel shows the result for one simulated population size.

Experimental duration

Our analysis showed that after 60 generations of adaptation, only experimental designs with a population size of at least 1,200 individuals have more power than 50 % (Figure 3). The 1:1_1f crossing scheme clearly outperforms dil:mt regardless of the population size, and reaches power above 75% only with a population size of at least 4,800 individuals. However, the maintenance of 5 replicates for 60 generations is labor intensive and can be quite time consuming for most sexual organisms. We explored the possibility that shorter experiments with less than 60 generations may already be sufficient to achieve satisfactory power values. We thus reanalyzed the power of the 1:1_1f, and dil:mt crossing scheme after 20 generations of adaptation (Table 1).

Consistent with previous computer simulation studies (Baldwin-Brown et al. 2014;

Kofler & Schlötterer 2014; Kessner & Novembre 2015) and empirical results (Langmüller &

Schlötterer 2020), we observe reduced power for experimental designs with shorter

experimental duration (Figure S3). Consistent with the results after 60 generations, the crossing

scheme (LRT full-reduced model comparison (Model 1): $\chi^2 = 2672.7$, df $= 8$, p<0.001

(success-A); $\chi^2 = 2234.4$, df $= 8$, p<0.001 (success-B)) and the population size (LRT full-

reduced model comparison (Model 1): $\chi^2 = 3176.4$, df $= 12$, p<0.001 (success-A); $\chi^2 =$

3375.3, df $= 12$, p<0.001 (success-B)) have the biggest effects on the success of secondary

E&R (TableS2-3).

In contrast to our analysis after 60 generations of adaptation, the dil:mt crossing scheme

results in higher power than the 1:1_1f crossing scheme for populations with the smallest

simulated population size (300 individuals) after 20 generations of adaptation (Figure S4). A

possible explanation for this is that the dil:mt crossing scheme has multiple beneficial targets.

Because we simulate additive selection, linked selection targets can act synergistically and

increase the frequency of the focal SNP over shorter time scales. This phenomenon of

pronounced allele frequency increase due to linked selection will be especially important if the

population size is small (i.e., drift is not neglectable), and the experimental duration is short.

This is reflected in a higher Nagelkerke's R^2 index for population size in secondary E&R with

an experimental duration of 20 generations compared to 60 generations of adaptation (Table

S3).

With increasing population size and thus, reduced genetic drift, the 1:1_1f crossing

scheme outcompetes dil:mt, as already seen in our analysis after 60 generations (Figure S3-4).

Although shorter experimental duration reduces power, our analysis after 20 generations of

adaptation highlights that if the maintenance of experimental populations for many generations

is not feasible, shorter experiments can achieve similar power if they are maintained at larger

population sizes (Figure 3, Figure S3).

<u>Additional target of selection in the 1:1 crossing scheme</u>

In contrast to dil:mt, the 1:1_1f crossing scheme harbors only one target of selection. We

simulated two additional versions of the 1:1 crossing scheme to investigate how one additional

target of selection influences the power of secondary E&R (Figure S1; Model 2, *Assessment of*

experimental and population genetic parameters in Material & Methods). We observed that

one additional beneficial SNP has a significant effect on the power of secondary E&R using

the 1:1 crossing scheme (LRT full-reduced model comparison (Model 2): $\chi^2 = 1727.7$, df =

32, p<0.001 (success-A); $\chi^2 = 2588.42$, df =32, p<0.001 (success-B); Table S4-5). This

significant effect is mainly driven by one scenario: when the focal line harbors two targets of

selection with the target of interest being dominant and the additional target being co-dominant

(Figure 5) the power to detect the target of interest is close to zero. The reason is that with a

starting frequency of 50 %, co-dominant alleles respond more to strong selection than dominant

alleles, because non-favored alleles are masked by high frequency dominant alleles (Figure S5).

This differential behavior is particularly pronounced for high allele frequencies. Note that our

definitions of success do not use prior information on the location of the focal SNP of interest.

We propose that a substantial fraction of the power can be recovered if the analysis is restricted

to the approximate location of the focal target determined in the primary E&R study. On the

other hand, for additive targets of interest one additional target of selection hardly reduces the

power of the 1:1 crossing scheme, regardless of the dominance coefficient of the additional

target and whether the additional target is positioned on the focal or one non-focal haplotype

(Figure 5). Overall, our results show that the 1:1 crossing scheme still outperforms dil:mt even

550 in the presence of one additional target of selection (Figure 3, Figure 5). For the remaining

551 analysis we will focus on a 1:1_1f crossing scheme with only one target of selection.

553 **Figure 5.** Power of the different 1:1 crossing schemes at different population sizes (Model 2) (2,000
554 simulations/experimental design) after 60 generations of adaptation. Bars show the power (i.e., proportion of
555 successful simulations) separately for each combination of 1:1 crossing scheme version with the dominance
556 coefficient of the additional target of selection: 1f, only one target of selection on the focal haplotype;
557 2f_h05/2f_h1, 2 targets of selection on the focal haplotype; 1f1nf_h05/1f1nf_h1, 1 target of selection on the focal
558 haplotype and one on one non-focal haplotype. Columns depict different population sizes (300; 1,200; 4,800;
559 19,200 individuals), and rows show different dominance coefficients of the target of interest (additive in top row,
560 dominant in bottom row). The dots with error bars display the model fit (Model 2) and its 95 % confidence interval.
561 For the model fit, the selection coefficient was fixed to its global average. (A) shows the results for success-A (=
562 selection target is the SNP with the highest Cochran-Mantel-Haenszel (CMH) test statistic) (B) shows the results
563 for success-B (= selection target is not more than 100 SNPs away from the SNP with the highest CMH test
564 statistic).

Alternative crossing schemes

567 Since large population sizes can be challenging to maintain, we also evaluated additional

568 crossing schemes which may perform better even for smaller population sizes (Model 3,

569 *Assessment of experimental and population genetic parameters* in Material & Methods). 1:few

570 (Figure 1C) is a modification of the 1:1_1f crossing scheme, which combines all non-focal lines

571 in each replicate - this provides a consistent genetic composition in each replicate. The 1:many

crossing scheme (Figure 1D) uses 1 focal, and 99 non-focal lines without selection target. We reasoned that the larger number of segregating variants (970,466 SNPs compared to max 527,771 SNPs in 1:1 crossing scheme, Table S1) potentially provides a higher mapping resolution. Finally, we modified the dilution crossing scheme by simulating only one selection target in the founder population (dil:st, Figure 1E). Hence, no additional selection targets can create potentially confounding selection signatures. For each experimental design, we simulated a population size of 300 individuals, 60 generations of adaptation, and 5 replicates (Table1).

Regardless of the experimental design, the simulated focal selection targets experienced a pronounced allele frequency increase (Figure S6). In "dilution" crossing schemes (dil:st, dil:mt; Figure 1E-F) the frequency trajectories of the focal SNPs were highly variable because of the heterogeneous starting frequency. The crossing schemes where a single focal line with a starting frequency of 50 % carries the beneficial allele typically had a superior performance than dilution crossing schemes (Figure 6). Notably, the significant influence of crossing scheme on secondary E&R success (LRT full-reduced model comparison (Model 3): $\chi^2 = 295.9$, df $=$ 8, p<0.001 (success-A); $\chi^2 = 190.23$, df $=8$, p<0.001 (success-B)) (Table S6-7) cannot be explained by a loss of the focal SNP in dilution crossing schemes, which occurred in less than 1 % of the simulations.

We found that the dil:st crossing scheme is still inferior to the 1:1_1f crossing scheme for additive loci (Figure 6). Also, the modifications of the 1:1_1f crossing scheme (1:few, 1:many) do not provide a substantial improvement (Figure 6). Surprisingly, the 1:many crossing scheme performed very poorly. This may be at least partly attributed to the larger number of SNPs/kb, which will affect the SNP-based success rate (success-B, Figure 2). Given that the

three additional crossing schemes do not provide a clear advantage (Figure 6-7, Figure S7), and

are not easier to execute experimentally, we did not evaluate them with larger population sizes.

Figure 6. Power of five different crossing schemes (population size = 300 individuals; 2,000
simulations/experimental design). Bars show the power (i.e., proportion of successful simulations) separately for
each combination of crossing scheme and dominance coefficient (additive in grey; dominant in white). The dots
with error bars display the estimate from the fitted model (Model 3) and its 95 % confidence interval. For the
model fit, the selection coefficient was fixed to its global average, and combination-specific average starting allele
frequencies were used. (A) shows the results for success-A (= selection target is the SNP with the highest Cochran-
Mantel-Haenszel (CMH) test statistic), (B) shows the results for success-B (= selection target is not more than 100
SNPs away from the SNP with the highest CMH test statistic).

Figure 7. Resolution of five different crossing schemes. Proportion of simulations (y-axis) that do not exceed a

maximum distance in SNPs (x-axis) between the SNP with the highest Cochran-Mantel-Haenszel test statistic and

the true target of selection.

<u>Number of replicates</u>

Given the superior performance of experimental designs with larger population sizes (Figure

3), we were also interested whether more replicates with a smaller population size may provide

further improvements. We performed additional simulations for all five different crossing

schemes with a population size of 300 individuals per replicate, and 30 replicates (Table 1).

Consistent with other simulation studies (Kofler & Schlötterer 2014; Kessner & Novembre

2015), more replicates result in increased power (Figure S8). The power of secondary E&R

with 30 replicates (Table S6) is only significantly affected by the crossing scheme considering

all modeled interactions (LRT full-reduced model comparison (Model 3): $\chi^2 = 31.467$, df $= 8$,

p<0.001 (success-A); $\chi^2 = 16.692$, df $= 8$, p=0.033 (success-B); Table S7). The 1:1_1f

crossing scheme with 30 replicates also has the highest resolution (Figure S9A). As expected,

the impact of linkage disequilibrium, indicated by the number of ties, is dramatically reduced

for all crossing schemes when more replicates are simulated (Figure S9B compared to Figure

S7). However, similar improvements as seen with 30 replicates can be achieved if 5 replicates

of the 1:1_1f crossing scheme are used with a population size of 1,200 individuals (Figure 3).

Thus, the same power can be achieved while maintaining about 33 % fewer individuals.

<u>Dominance coefficient, Selection coefficient, and mean starting allele frequency</u>

As expected, dominance coefficient, selection coefficient, and average starting allele frequency

have a significant effect on the success of secondary E&R in all simulation scenarios, except

for data with 30 replicates. We observed that selection targets with a higher selection coefficient

and/or a higher mean starting allele frequency are easier to fine map. The lack of significant

effects of these parameters on secondary E&R success for the dataset with 30 replicates can

probably be explained by the smaller number of conducted simulations (100 for 30 replicates

and 2,000 for 5 replicates, Table 1). After crossing scheme and population size, the dominance

coefficient is the parameter with the highest Nagelkerke's R^2 index in our analysis, with

additive loci being easier to fine map. This is caused by the fact that heterozygotes and target

homozygotes with a dominant beneficial SNP have the same fitness, resulting in less efficient

selection that becomes especially apparent if the selected allele has already reached a high

frequency in the population, and non-selected allele homozygotes become rare (indicated by a

significant negative effect of the interaction term between dominance coefficient and mean

starting allele frequency in most of our statistical models).

Discussion

This work was inspired by the difficulty of most E&R studies with sexually reproducing

organisms to pinpoint selection targets, mostly due to numerous neutral hitchhikers resulting in

large haplotype blocks. Secondary E&R – a follow-up EE validating putative selection targets

of a primary E&R under an identical selection regime – has been recently suggested as

experimental approach for selection target confirmation (Burny et al. 2020). We used extensive

computer simulations to evaluate how experimental and population genetics parameter shape

the power and resolution of secondary E&R. As expected, dominance coefficient, selection

strength, and mean starting allele frequency of the selected target all have a significant effect

on the success of E&R, where dominant selection targets at high frequency are particularly

challenging to detect. However, population size and crossing scheme emerged as the most

influential parameters in our analysis.

<u>The crossing scheme has a pronounced effect on secondary E&R success</u>

We show that a simple crossing scheme, which only requires that (a subset of) the founder lines

are sequenced and that founder lines with and without the selection target of interest can be

distinguished, has the best power and resolution of the five crossing schemes tested. The 1:1

crossing scheme is particularly well-suited when many selection targets are detected in the

evolved populations of the primary E&R, because it uses only a subset of the lines from the

non-adapted ancestral founder population of the primary E&R study. This reduces potential

confounding effects between the selection target of interest and other adaptive loci in two ways.

First, non-adapted ancestral founder haplotypes will harbor on average less selection targets

than evolved haplotypes that can acquire multiple selection targets through recombination

events during the primary E&R (Otte & Schlötterer 2021). Because evolved haplotypes will

often harbor multiple selection targets, beneficial alleles will not propagate independently from

each other, which makes it challenging to fine-map single selection targets. Second, the total

number of potential beneficial alleles in a 1:1 crossing scheme is deliberately reduced by

picking only a subset of the founder lines of the primary E&R, which facilitates fine mapping

of one particular selection target of interest with secondary E&R.

We would like to point out that in our study "crossing schemes" are defined such that

they do not only differ in the actual crossing procedure, but also in the number and starting

allele frequency distribution of simulated selection targets (e.g., 1:1_1f vs. dil:mt). Additional

beneficial alleles can without doubt have an impact on the power of secondary E&R. While the

1:1 crossing scheme still outperforms dil:mt in the presence of one additional selection target,

the relative performance of crossing schemes might change given an even more complex

underlying genetic architecture. Furthermore, we focused in this study on the assessment of

experimental and population genetic parameters given a directional selection regime. Future

research on the potential of secondary E&R to fine map selection targets under a

complex/polygenic genetic architecture is needed, such as for example a quantitative trait under

stabilizing selection that experienced a recent shift in trait optimum.

<u>Sufficient E&R power requires large population sizes</u>

Our analyses also indicate that rather large population sizes are crucial to identify the causative

variant with sufficient confidence using a secondary E&R approach. This observation explains

why an empirical secondary E&R study in *D. simulans* with a population size of 1,250 flies per

replicated population failed to narrow down a pronounced candidate region from a primary

E&R in a secondary E&R after 30 generations, and only confirmed the presence of selected

alleles (Burny et al. 2020).

Our results suggest that large population sizes become especially crucial for short-term

secondary E&R. The low power of short-term E&R with reduced experimental duration (Kofler

& Schlötterer 2014) can be improved by a larger experimental population size. Our analyses

show that to achieve satisfactory power after only 20 generations requires population sizes of

tens of thousands of individuals– dimensions that are rather rare in current EE designs with

strictly sexually reproducing organisms, where population sizes are mostly limited by the

capacities to conduct such large experiments. Similar to previous simulation studies (Kofler &

Schlötterer 2014; Kessner & Novembre 2015), we demonstrated that increasing the number of

replicates results in more powerful secondary E&R. If large population sizes are not feasible,

maintaining many replicates at small population size can help to boost secondary E&R

performance. Additionally, the loss of one replicate in a highly replicated setup has a smaller

impact than in an experimental setup with very few replicates. However, we show that

additional replication cannot completely compensate the advantage of large population sizes in

secondary E&R. Based on this observation, we conclude that independent of the crossing

scheme, the power of secondary E&R will benefit from larger population sizes. Furthermore,

the resolution will significantly increase with more replicates – a result that we anticipate can

be generalized to different underlying genetic architectures as well as (model) organisms.

<u>Allele frequency estimation errors and read depth</u>

The results of this study are based on true allele frequencies estimated without error - a rather

optimistic assumption that will not hold for empirical data. In empirical E&R studies, numerous

factors influence the accuracy of allele frequency estimates, such as the rate of sequencing

errors, genome-wide read depth heterogeneity, and the average read depth (Baldwin-Brown et

al. 2014; Kofler & Schlötterer 2014; Tilk et al. 2019).

Individual whole-genome sequencing for entire populations becomes quickly

prohibitive with increasing sample size. Sequencing pools of individuals (Pool-Seq)

(Schlötterer et al. 2014) provides a cost-effective approach that allows to robustly estimate

allele frequency estimates and has become the method of choice for most E&R studies (Turner

et al. 2011; Schlötterer et al. 2015). Since typical Pool-Seq studies combine all individuals from

a given generation to reduce the sampling error (Schlötterer et al. 2014), the read depth of Pool-

Seq studies will affect all experimental designs to the same extent as long as the coverage is

considerably lower than the pool size.

<u>Choice of model organism</u>

Our simulations are parameterized for *D. simulans* because this species is better suited for E&R

studies than *D. melanogaster* (Barghi et al. 2017). While other species, such as *S. cerevisiae*

(Burke et al. 2014) or *C. remanei* (Castillo et al. 2015) have been used to study adaptive

response from standing genetic variation in outcrossing species, *Drosophila* is currently the

most popular organism. Nevertheless, we anticipate that the results of this study can be

generalized to other species, but the availability of sequenced inbred founder lines is probably

a more severe restriction, which could limit the widespread use of secondary E&R.

<u>Selection regime</u>

Our simulations are targeted for selection regimes with a moderate number of selection targets,

with rather strong effects, as seen in empirical E&R studies using *Drosophila* (e.g. Mallard et

al. 2018; Barghi et al. 2019; Michalak et al. 2019). Since strongly selected haplotype blocks

typically have low starting frequencies, we consider the 1:1 crossing scheme a very realistic

case, as most of the selection targets will be found only in a few ancestral founder lines. As

outlined above, evolved haplotypes in dil:mt on the other hand will most likely carry multiple

beneficial alleles that have recombined over the course of the primary E&R, which makes it

challenging to fine-map one distinct selection target of interest as our results suggest.

Furthermore, we assumed the rather simple selection scenario of directional selection.

More complex scenarios that include pleiotropy and/or epistasis were not considered. While

both are very important factors that could affect the allele frequency changes in an E&R study,

we would like to point out that secondary E&R is designed to study loci, which experienced a

substantial allele frequency increase in a primary experiment, thus pleiotropic constraints are

not expected to be a major confounding factor. Epistasis, on the other hand may be more

pronounced in the 1:1 crossing scheme. It may be possible that the focal allele is only selected

in a subset of the replicates, but not in others-or the response may differ between replicates.

While this reduces the power of secondary E&R to fine-map the selection target, such a result

may open the possibility to study the impact of epistatic effects. Since the founder lines will be

available, it will be possible to repeat the experiment with replication for each genotype

combination to assure that the heterogeneous response comes from epistatic effects and is not

linked to stochastic changes.

Finally, it is important to keep in mind that the secondary E&R designs discussed here

are not tailored to study highly polygenic traits, because in this study we were assuming that

recombination facilitates the identification of a selection target with a major effect. For a highly

polygenic trait, selected haplotype blocks identified in the primary E&R experiment (with

multiple selected loci) would be broken down during the secondary E&R, which reduces their

selective advantage. To what extent this different behavior in secondary E&R experiments can

be used to distinguish these different architectures requires further work.

Acknowledgements

We thank the members of the Institute of Population Genetics for fruitful discussion and

support. Thanks to Roger Mundry who shared R scripts for the statistical analyses. Special

thanks to Rupert Mazzucco, Robert Kofler, Kathrin Otte, and Christos Vlachos for their support

regarding the computer simulations. Special thanks to Claire Burny and Robert Kofler for

helpful comments on earlier versions of the manuscript. Last, but not least we are thankful to

the anonymous reviewers for their constructive feedback. This work was supported by the

Austrian Science Funds (FWF) [grant numbers W1225, P-32935]; and by the European

Research Council (ERC) [grant ArchAdapt].

Authors Contribution

AML, MD, and CS designed the analysis. AML performed the forward simulations, and the

bioinformatic analysis. AML, and MD performed the statistical analysis. AML, MD, and CS

wrote the manuscript.

References

Agresti A, Kateri M. 2011. Categorical Data Analysis. In: International Encyclopedia of

Statistical Science. Lovric, M, editor. Springer Berlin Heidelberg: Berlin, Heidelberg pp.

206–208. doi: 10.1007/978-3-642-04898-2_161.

Baayen RH. 2008. *Analyzing Linguistic Data: A Practical Introduction to Statistics using R*.

Cambridge University Press doi: 10.1017/CBO9780511801686.

Baldwin-Brown JG, Long AD, Thornton KR. 2014. The Power to Detect Quantitative Trait

Loci Using Resequenced, Experimentally Evolved Populations of Diploid, Sexual Organisms.

Mol. Biol. Evol. 31:1040–1055. doi: 10.1093/molbev/msu048.

Barghi N et al. 2019. Genetic redundancy fuels polygenic adaptation in *Drosophila*. PLOS

Biol. 17:1–31. doi: 10.1371/journal.pbio.3000128.

Barghi N, Hermisson J, Schlötterer C. 2020. Polygenic adaptation: a unifying framework to

understand positive selection. Nat. Rev. Genet. 21:769–781. doi:

https://doi.org/10.1038/s41576-020-0250-z.

Barghi N, Tobler R, Nolte V, Schlötterer C. 2017. *Drosophila simulans* : A Species with

Improved Resolution in Evolve and Resequence Studies. G3 Genes|Genomes|Genetics.

7:2337–2343. doi: 10.1534/g3.117.043349.

Burke MK, Liti G, Long AD. 2014. Standing Genetic Variation Drives Repeatable

Experimental Evolution in Outcrossing Populations of *Saccharomyces cerevisiae*. Mol. Biol.

Evol. 31:3228–3239. doi: 10.1093/molbev/msu256.

Burny C et al. 2020. Secondary Evolve and Resequencing: An Experimental Confirmation of

Putative Selection Targets without Phenotyping. Genome Biol. Evol. 12:151–159. doi:

10.1093/gbe/evaa036.

Castillo DM, Burger MK, Lively CM, Delph LF. 2015. Experimental evolution: Assortative

mating and sexual selection, independent of local adaptation, lead to reproductive isolation in

the nematode Caenorhabditis remanei. Evolution. 69:3141–3155. doi: 10.1111/evo.12815.

Dowle M, Srinivasan A. 2019. data.table: Extension of `data.frame`. https://cran.r-

project.org/package=data.table.

Fox J, Monette G. 1992. Generalized Collinearity Diagnostics. J. Am. Stat. Assoc. 87:178–

183. http://www.jstor.org/stable/2290467.

Fox J, Weisberg S. 2019. *An R Companion to Applied Regression*. Third. Sage

https://socialsciences.mcmaster.ca/jfox/Books/Companion/.

Franssen SU, Nolte V, Tobler R, Schlötterer C. 2015. Patterns of Linkage Disequilibrium and

Long Range Hitchhiking in Evolving Experimental *Drosophila melanogaster* Populations.

Mol. Biol. Evol. 32:495–509. doi: 10.1093/molbev/msu320.

Garland T, Rose M. 2009. *Experimental Evolution: Concepts, Methods, and Applications of*

Selection Experiments.

Gratz SJ et al. 2013. Genome engineering of *Drosophila* with the CRISPR RNA-guided Cas9

nuclease. Genetics. 194:1029–1035. doi: 10.1534/genetics.113.152710.

Howie JM, Mazzucco R, Taus T, Nolte V, Schlötterer C. 2019. DNA Motifs Are Not General

Predictors of Recombination in Two *Drosophila* Sister Species. Genome Biol. Evol.

11:1345–1357. doi: 10.1093/gbe/evz082.

Iranmehr A, Akbari A, Schlötterer C, Bafna V. 2017. CLEAR: Composition of Likelihoods

for Evolve and Resequence Experiments. Genetics. 206:1011–1023. doi:

10.1534/genetics.116.197566.

Johansson AM, Pettersson ME, Siegel PB, Carlborg Ö. 2010. Genome-Wide Effects of Long-

Term Divergent Selection. PLOS Genet. 6:e1001188. doi: 10.1371/journal.pgen.1001188.

Kawecki TJ et al. 2012. Experimental evolution. Trends Ecol. Evol. 27:547–560. doi:

10.1016/j.tree.2012.06.001.

Kelly JK, Hughes KA. 2019. Pervasive Linked Selection and Intermediate-Frequency Alleles

Are Implicated in an Evolve-and-Resequencing Experiment of *Drosophila simulans*.

Genetics. 211:943–961. doi: 10.1534/genetics.118.301824.

Kessner D, Novembre J. 2015. Power analysis of artificial selection experiments using

efficient whole genome simulation of quantitative traits. Genetics. 199:991–1005. doi:

10.1534/genetics.115.175075.

Kofler R, Pandey RV, Schlötterer C. 2011. PoPoolation2: identifying differentiation between

populations using sequencing of pooled DNA samples (Pool-Seq). Bioinformatics. 27:3435–

3436. doi: 10.1093/bioinformatics/btr589.

Kofler R, Schlötterer C. 2014. A guide for the design of evolve and resequencing studies.

Mol. Biol. Evol. 31:474–483. doi: 10.1093/molbev/mst221.

Lang GI et al. 2013. Pervasive genetic hitchhiking and clonal interference in forty evolving

yeast populations. Nature. 500:571–574. doi: 10.1038/nature12344.

Langmüller AM, Schlötterer C. 2020. Low concordance of short-term and long-term selection

responses in experimental Drosophila populations. Mol. Ecol. 29:3466–3475. doi:

10.1111/mec.15579.

Long A, Liti G, Luptak A, Tenaillon O. 2015. Elucidating the molecular architecture of

adaptation via evolve and resequence experiments. Nat. Publ. Gr. 16:567–582. doi:

10.1038/nrg3937.

Mallard F, Nolte V, Tobler R, Kapun M, Schlötterer C. 2018. A simple genetic basis of

adaptation to a novel thermal environment results in complex metabolic rewiring in

Drosophila. Genome Biol. 19:119. doi: 10.1186/s13059-018-1503-4.

Martins NE et al. 2014. Host adaptation to viruses relies on few genes with different cross-

resistance properties. Proc. Natl. Acad. Sci. U. S. A. 111:5938–43. doi:

10.1073/pnas.1400378111.

Michalak P, Kang L, Schou MF, Garner HR, Loeschcke V. 2019. Genomic signatures of

experimental adaptive radiation in *Drosophila*. Mol. Ecol. 28:600–614. doi:

https://doi.org/10.1111/mec.14917.

Nagelkerke N. 1991. A note on a general definition of the coefficient of determination.

Biometrika. 78:691–692.

Nuzhdin S V, Turner TL. 2013. Promises and limitations of hitchhiking mapping. Curr. Opin.

Genet. Dev. 23:694–699. doi: 10.1016/j.gde.2013.10.002.

Otte KA, Schlötterer C. 2021. Detecting selected haplotype blocks in evolve and resequence

experiments. Mol. Ecol. Resour. 21:93–109. doi: 10.1111/1755-0998.13244.

R Core Team 3.5.3. 2019. A Language and Environment for Statistical Computing. R Found.

Stat. Comput. 2:https://www.R-project.org. http://www.r-project.org.

Remolina SC, Chang PL, Leips J, Nuzhdin S V, Hughes KA. 2012. Genomic basis of aging

and life-history evolution in *Drosophila melanogaster*. Evolution. 66:3390–3403. doi:

10.1111/j.1558-5646.2012.01710.x.

Schielzeth H. 2010. Simple means to improve the interpretability of regression coefficients.

Methods Ecol. Evol. 1:103–113. doi: 10.1111/j.2041-210X.2010.00012.x.

Schlötterer C, Kofler R, Versace E, Tobler R, Franssen SU. 2015. Combining experimental

evolution with next-generation sequencing: a powerful tool to study adaptation from standing

genetic variation. Heredity (Edinb). 114:431–440. doi: 10.1038/hdy.2014.86.

Schlötterer C, Tobler R, Kofler R, Nolte V. 2014. Sequencing pools of individuals — mining

genome-wide polymorphism data without big funding. Nat. Rev. Genet. 15:749–763. doi:

10.1038/nrg3803.

Spitzer K, Pelizzola M, Futschik A. 2020. Modifying the Chi-square and the CMH test for

population genetic inference: Adapting to overdispersion. Ann. Appl. Stat. 14:202–220. doi:

10.1214/19-AOAS1301.

Taus T, Futschik A, Schlötterer C. 2017. Quantifying selection with pool-seq time series data.

Mol. Biol. Evol. 34:3023–3034. doi: 10.1093/molbev/msx225MBE.

Teotónio H, Chelo IM, Bradić M, Rose MR, Long AD. 2009. Experimental evolution reveals

natural selection on standing genetic variation. Nat. Genet. 41:251–257. doi: 10.1038/ng.289.

Tilk S et al. 2019. Accurate Allele Frequencies from Ultra-low Coverage Pool-Seq Samples

in Evolve-and-Resequence Experiments. G3 Genes|Genomes|Genetics. 9:4159–4168. doi:

10.1534/g3.119.400755.

Tobler R et al. 2014. Massive Habitat-Specific Genomic Response in *D. melanogaster*

Populations during Experimental Evolution in Hot and Cold Environments. Mol. Biol. Evol.

31:364–375. doi: 10.1093/molbev/mst205.

Topa H, Jónás Á, Kofler R, Kosiol C, Honkela A. 2015. Gaussian process test for high-

throughput sequencing time series: Application to experimental evolution. Bioinformatics.

31:1762–1770. doi: 10.1093/bioinformatics/btv014.

Turner TL, Stewart AD, Fields AT, Rice WR, Tarone AM. 2011. Population-Based

Resequencing of Experimentally Evolved Populations Reveals the Genetic Basis of Body

Size Variation in *Drosophila melanogaster*. PLoS Genet. 7:e1001336. doi:

10.1371/journal.pgen.1001336.

Vlachos C et al. 2019. Benchmarking software tools for detecting and quantifying selection in

evolve and resequencing studies. Genome Biol. 20:169. doi: 10.1186/s13059-019-1770-8.

Vlachos C, Kofler R. 2018. MimicrEE2: Genome-wide forward simulations of Evolve and

Resequencing studies. PLOS Comput. Biol. 14:e1006413.

https://doi.org/10.1371/journal.pcbi.1006413.

Vlachos C, Kofler R. 2019. Optimizing the Power to Identify the Genetic Basis of Complex

Traits with Evolve and Resequence Studies. Mol. Biol. Evol. 36:2890–2905. doi:

10.1093/molbev/msz183.

Supplementary Figures

Figure S1. Simulated versions of the 1:1 crossing scheme.

Figure S2. Number of ties in the 1:1_1f and dil:mt crossing schemes.

Figure S3. Power of the 1:1_1f and dil:mt crossing schemes after 20 generations of adaptation.

Figure S4. Resolution of the 1:1_1f and dil:mt crossing schemes after 20 generations of

adaptation.

Figure S5. Allele frequency trajectories of selected SNPs in 1:1_2f.

Figure S6. Allele frequency trajectories of focal SNPs in 5 different crossing schemes.

Figure S7. Number of ties in 5 different crossing schemes.

Figure S8. Power of 5 different crossing schemes with 30 replicates per simulation.

Figure S9. Resolution of 5 different crossing schemes with 30 replicates per simulation.

Figure S1. Simulated versions of the 1:1 crossing scheme. (A) Default version of the 1:1 crossing scheme (=**1f**

for 1 focal SNP). Inbred flies with one target of selection (purple) are crossed to inbred flies without known

beneficial variants. The starting frequency of each genotype is 50%. In each replicate the line with the beneficial

allele (focal line, purple) is crossed to a different line lacking beneficial mutations (non-focal lines are colored in

different shades of grey). Figure S1A is equal to Figure 1B in the main manuscript. (B) 2 focal SNPs (=**2f**) version

of the 1:1 crossing scheme. The focal line carries two beneficial variants: the focal SNP we aim to fine map (purple)

and one additional target of selection (green). (C) 1 focal, 1 non-focal SNP (=**1f1nf**) version of the 1:1 crossing

scheme: One non-focal line carries on additional beneficial variant (green).

Figure S2. Violin plots of the number of ties (y axis = log$_{10}$ (number of ties + 1)) for simulations where the true target of selection has the highest Cochran-Mantel-Haenszel (CMH) test statistic (success-A). The black horizontal lines in the violin plots display the median number of observed ties. Each panel shows the results for one particular simulated population size. Ties are defined as neighboring SNPs that have the same CMH test statistic as the target of selection.

Figure S3. Power of the 1:1_1f and dil:mt crossing scheme at different population sizes (2,000

simulations/experimental design) after 20 generations of adaptation. Bars show the power (i.e., proportion of

successful simulations) separately for each combination of crossing scheme (1:1_1f, dil:mt), population size (300;

1,200; 4,800; 19,200 individuals), and dominance coefficient (additive in grey, dominant in white). The dots with

error bars display the model fit (Model 1) and its 95 % confidence interval. For the model fit, the selection

coefficient was fixed to its global average, and combination-specific average starting allele frequencies were used.

(A) shows the results for success-A (= selection target is the SNP with the highest Cochran-Mantel-Haenszel

(CMH) test statistic) (B) shows the results for success-B (= selection target is not more than 100 SNPs away from

the SNP with the highest CMH test statistic).

Figure S4. Resolution for the 1:1_1f and dil:mt crossing scheme simulated with different population sizes after

20 generations of adaptation. (A) Proportion of simulations (y-axis) that do not exceed a maximum distance in

SNPs (x-axis) between the SNP with the highest Cochran-Mantel-Haenszel (CMH) test statistic and the true target

of selection (B) Violin plots of the number of ties (y axis = log10 (number of ties + 1)) for simulations where the

true target of selection has the highest CMH test statistic (success-A). The black horizontal lines in the violin plots

display the median number of observed ties. Ties are defined as neighboring SNPs that have the same CMH test

statistic as the target of selection.

Figure S5. Median allele frequency trajectory of focal (red) and additional targets (grey) in 1:1_2f. For each single simulation, the target frequencies of a distinct generation were averaged over five replicates. Solid lines show the median allele frequency trajectory over 2,000 independent simulations per experimental design. Dashed lines show the 5 and 95 percentiles of the allele frequency trajectories. Each panel shows the median allele frequency trajectory of one particular dominance coefficient (focal target; additional target) and population size (300; 1,200; 4,800; 19,200 individuals) combination.

Figure S6. Median allele frequency trajectory of focal targets in different experimental designs. For each single

simulation, the focal target frequency of a distinct generation was averaged over five replicates. Solid lines show

the median allele frequency trajectory over 2,000 independent simulations per experimental design. Dashed lines

show the 5 and 95 percentiles of the allele frequency trajectories. Each panel shows the median allele frequency

trajectory of one particular crossing scheme (1:1_1f, dil:mt, 1:few, 1:many, dil:st) and population size (300; 1,200;

4,800; 19,200 individuals) combination.

Figure S7. Violin plots of the number of ties (y axis = log10 (number of ties + 1)) for simulations where the true

target of selection has the highest Cochran-Mantel-Haenszel (CMH) test statistic (success-A, population size =

300 individuals; 5 replicates; 2,000 simulations/experimental design). The black horizontal lines in the violin plots

display the median number of observed ties. Ties are defined as neighboring SNPs that have the same CMH test

statistic as the target of selection.

Figure S8. Power of five different crossing schemes (population size = 300 individuals; 30 replicates; 100 simulations/experimental design). Bars show the power (i.e., proportion of successful simulations) separately for each combination of crossing scheme and dominance coefficient (additive in grey; dominant in white). The dots with error bars display the estimate from the fitted model (Model 3) and its 95 % confidence interval. For the model fit, the selection coefficient was fixed to its global average, and combination-specific average starting allele frequencies were used. (A) shows the results for success-A (= selection target is the SNP with the highest Cochran-mantel-Haenszel (CMH) test statistic), (B) shows the results for success-B (= selection target is not more than 100 SNPs away from the SNP with the highest CMH test statistic).

Figure S9. Resolution for five different crossing schemes (population size = 300 individuals; 30 replicates; 100 simulations/experimental design). (A) Proportion of simulations (y-axis) that do not exceed a maximum distance in SNPs (x-axis) between the SNP with the highest Cochran-Mantel-Haenszel (CMH) test statistic and the true target of selection (B) Violin plots of the number of ties (y axis = $\log_{10}$ (number of ties + 1)) for simulations where the true target of selection has the highest CMH test statistic (success-A). The black horizontal lines in the violin plots display the median number of observed ties. Ties are defined as neighboring SNPs that have the same CMH test statistic as the target of selection.

Supplementary Tables

Table S1. Number of SNPs per crossing scheme.

Table S2. Type II ANOVA (Model 1).

Table S3. Nagelkerke's R^2-index (Model 1).

Table S4. Type II ANOVA (Model 2).

Table S5. Nagelkerke's R^2-index (Model 2).

Table S6. Type II ANOVA (Model 3).

Table S7. Nagelkerke's R^2-index (Model 3).

Table S1. Number of SNPs per crossing scheme on chromosome-arm 2L.

crossing scheme	median	range
1:1	523,008	479,507 – 527,771
1:few	526,510	522,470 – 530,230
1:many	970,466	970,466 – 970,466
dil:mt	970,466	970,466 – 970,466
dil:st	970,466	970,466 – 970,466

Table S2. Type II ANOVA of explanatory variables for the analysis of success in secondary Evolve and Resequence studies with model 1 after 60 and 20 generations of adaptation. cross= crossing scheme; h= dominance coefficient; s= selection coefficient; af= mean starting allele frequency of the focal target over 5 replicates; N= population size; cross:h = interaction term between crossing scheme and dominance coefficient; h:af = interaction term between dominance coefficient and mean starting allele frequency; cross:N = interaction term between crossing scheme and population size ; h:N = interaction term between dominance coefficient and population size; cross:h:N = interaction term between crossing scheme, dominance coefficient, and population size.

generations	term	df	χ^2	P	χ^2	P
			success-A		success-B	
60	cross*	1	3188.7	<0.001	2514.70	<0.001
	h*	1	687.5	<0.001	652.41	<0.001
	s [1]	1	67.4	<0.001	73.39	<0.001
	af*[1]	1	3.9	0.048	0.69	0.407
	N*	3	1791.3	<0.001	1898.77	<0.001
	cross:h	1	2.1	0.145	8.24	0.004
	h:af	1	18.5	<0.001	16.67	<0.001
	cross:N	3	315.5	<0.001	677.98	<0.001
	h:N	3	6.8	0.079	2.70	0.440
	cross:h:N	3	3.8	0.286	16.12	0.001
20	cross*	1	2079.94	<0.001	1056.48	<0.001
	h*	1	139.04	<0.001	181.36	<0.001
	s [1]	1	18.21	<0.001	58.96	<0.001
	af* [1]	1	39.91	<0.001	77.13	<0.001
	N*	3	2589.34	<0.001	2204.36	<0.001
	cross:h	1	3.73	0.053	1.15	0.284
	h:af	1	0.02	0.898	0.36	0.551
	cross:N	3	573.81	<0.001	1156.81	<0.001
	h:N	3	0.58	0.900	0.87	0.833
	cross:h:N	3	1.29	0.732	4.68	0.197

* In type II ANOVA, main effects are not corrected for the interaction part of the model, while interaction terms are corrected for the main effects in the model.

[1] multiplied by 100, and z-transformed to mean = 0, and sd =1; Mean/standard deviation of the original value was 0.463/0.130 for the starting allele frequency, and 0.086/0.009 for the selection coefficient respectively

106 **Table S3.** Nagelkerke's R^2-index of each explanatory variable including all its interactions in the logistic

107 regression (Model 1). If the reduced model explains the data as well as a full model that includes the evaluated

108 effect, Nagelkerke's R^2-index is 0.

generations	effect	R^2 success-A	R^2 success-B
60	crossing scheme	0.305	0.264
	dominance coefficient	0.077	0.069
	selection coefficient	0.007	0.007
	mean starting allele frequency	0.003	0.002
	population size	0.204	0.224
20	crossing scheme	0.283	0.211
	dominance coefficient	0.019	0.021
	selection coefficient	0.002	0.007
	mean starting allele frequency	0.005	0.009
	population size	0.322	0.295

Table S4. Type II ANOVA of explanatory variables for the analysis of success in secondary Evolve and

Resequence studies with model 2. architecture= version of the 1:1 crossing scheme in combination with the

dominance coefficient of the additional target; h= dominance coefficient of the selection target of interest; s=

selection coefficient; N= population size; architecture:h = interaction term between architecture and dominance

coefficient of the selection target of interest; architecture:N = interaction term between architecture and population

size ; h:N = interaction term between dominance coefficient of the selection target of interest and population size;

architecture:h:N = interaction term between architecture, dominance coefficient of the selection target of interest,

and population size.

term	df	χ^2	P	χ^2	P
		success-A		success-B	
architecture*	4	855.4	<0.001	1509.9	<0.001
h*	1	2742.4	<0.001	2755.9	<0.001
s [1]	1	181.9	<0.001	137.5	<0.001
N*	3	5230.6	<0.001	6661.9	<0.001
architecture:h	4	697.8	<0.001	708.6	<0.001
architecture:N	12	17.5	0.132	42.7	<0.001
h:N	3	12.3	0.006	3.0	0.399
architecture:h:N	12	27.9	0.006	29.8	0.003

* In type II ANOVA, main effects are not corrected for the interaction part of the model, while interaction terms

are corrected for the main effects in the model.

[1] multiplied by 100, and z-transformed to mean = 0, and sd =1; Mean/standard deviation of the original value

was 0.085/0.009.

Table S5. Nagelkerke's R^2-index of each explanatory variable including all its interactions in the logistic

regression (Model 2). If the reduced model explains the data as well as a full model that includes the evaluated

effect, Nagelkerke's R^2-index is 0.

effect	R^2 success-A	R^2 success-B
architecture	0.107	0.158
dominance coefficient	0.201	0.210
selection coefficient	0.012	0.009
population size	0.285	0.348

126 **Table S6.** Type II ANOVA of explanatory variables for the analysis of success in secondary Evolve and

127 Resequence studies with model 3. cross= crossing scheme; h= dominance coefficient; s= selection coefficient; af=

128 mean starting allele frequency of the focal target over 5 replicates; cross:h= interaction term between crossing

129 scheme and dominance coefficient; h:af= interaction term between dominance coefficient and mean starting allele

130 frequency.

generations	term	df	χ^2	P	χ^2	P
			success-A		success-B	
5	cross*	4	218.65	<0.001	93.01	<0.001
	h*	1	371.35	<0.001	321.13	<0.001
	s [1]	1	76.20	<0.001	72.09	<0.001
	af* [1]	1	5.55	0.019	12.66	<0.001
	cross:h	4	77.19	<0.001	97.21	<0.001
	h:af	1	7.56	0.006	7.07	0.008
30	cross*	4	28.79	<0.001	11.28	0.024
	h*	1	5.13	0.024	4.40	0.036
	s [1]	1	2.63	0.105	3.65	0.056
	af* [1]	1	1.34	0.247	0.13	0.716
	cross:h	4	2.68	0.612	5.41	0.247
	h:af	1	1.73	0.189	1.01	0.315

* In type II ANOVA, main effects are not corrected for the interaction part of the model, while interaction terms

are corrected for the main effects in the model.

[1] multiplied by 100, and z-transformed to mean = 0, and sd =1; Mean/standard deviation of the original value

were 0.463/0.133 for the starting allele frequency, and 0.085/0.009 for the selection coefficient respectively.

Table S7. Nagelkerke's R^2-index of each explanatory variable including all its interactions in the logistic

regression (Model 3). If the reduced model explains the data equally well as a full model that includes the evaluated

effect, Nagelkerke's R^2-index is 0.

replicates	effect	R^2 success-A	R^2 success-B
5	crossing scheme	0.061	0.033
	dominance	0.095	0.076
	selection coefficient	0.016	0.013
	starting allele frequency	0.003	0.004
30	crossing scheme	0.097	0.046
	dominance	0.034	0.034
	selection coefficient	0.009	0.010
	starting allele frequency	0.010	0.003