Title: Genome-wide, genic, and site-specific changes associated with the
convergent evolution of sociality in spiders

Authors: Chao Tong[a,*], Leticia Avilés[b], Linda S. Rayor[c], Alexander S. Mikheyev[d,e], Timothy A.
Linksvayer[a,f,*]

Affiliations:

[a] Department of Biology, University of Pennsylvania, Philadelphia, PA, 19104, USA

[b] Department of Zoology, University of British Columbia, Vancouver, British Columbia V6T 1Z4, Canada

[c] Department of Entomology, Cornell University, Ithaca, New York 14853, USA

[d] Ecology and Evolution Unit, Okinawa Institute of Science and Technology, Okinawa, 904-0495, Japan

[e] Evolutionary genomics group, Research School of Biology, Australian National University, Canberra,
0200, Australia

[f] Department of Biological Sciences, Texas Tech University, Lubbock, TX, 79409, USA

[*] Correspondence: Chao Tong, Timothy A. Linksvayer

Email: tongchao1990@gmail.com (C.T.), tlinksvayer@gmail.com (T.A.L.)

C.T. 0000-0001-5202-5507; L.A. 0000-0002-9609-2952; L.S.R. 0000-0002-7132-6281; A.S.M. 0000-
0003-4369-1019; T.A.L. 0000-0001-7034-1546

Classification

Major – Biological Sciences; Minor – Evolution

Keywords

comparative genomics; convergent evolution; sociality; spiders

Author Contributions

CT and TAL conceived and designed the study. LA and LSR provided spider samples and expertise.
ASM provided reagents and sequencing support. CT generated libraries for RNA sequencing. CT and
TAL performed all transcriptomic and comparative genomic analyses and created all figures. CT and TAL
wrote the paper, with input from all authors.

This PDF file includes:
Main Text
Figures 1 to 4
33

Abstract

Sociality is a striking phenotypic innovation that independently evolved dozens of times across animals.

Sociogenomic approaches have begun to elucidate the molecular underpinnings of social life in social

insects and vertebrates, but the degree to which the convergent evolution of sociality involves convergent

molecular evolution remains controversial and largely unknown. Spiders are a powerful system for

identifying the genomic causes and consequences of social life because sociality is estimated to have

independently evolved 15 times, and each origin likely occurred recently, within the past few million years.

To determine if there are statistically supported genomic signatures of protein-coding sequence evolution

associated with the convergent evolution of sociality in spiders, we compared the genomes or

transcriptomes of 24 spider species that vary in social organization and represent at least seven

independent origins of sociality. We identified hundreds of genes that experienced shifts in patterns of

molecular evolution during the convergent evolution of sociality, and these genes were enriched for

several annotated functions, including neural function, neurogenesis, and behavior, as well as immune

function, growth, and metabolism. We also found evidence that directional selection for specific

substitutions repeatedly occurred in social species for several genes, in particular the calcium channel

gene *TPC1*. Finally, supporting previous results, we found elevated genome-wide rates of molecular

evolution in social species, resulting mainly from relaxation of selection. Altogether, we identify genome-

wide, genic, and site-specific changes that repeatedly occurred during the evolution of sociality in spiders.

Significance Statement

The transition from solitary to social life is a major transition in evolution that repeatedly occurred, but the

genetic underpinnings are largely unknown. To identify genomic changes associated with sociality, we

compared the genomes of 24 spider species, representing seven recent and independent origins of

sociality. We identified hundreds of genes and many functional classes of genes that tended to

experience shifts in molecular evolution, including genes that have previously been implicated in animal

social behavior and human behavioral disorders. Our study shows that while the precise genetic details

vary, the repeated evolution of sociality in spiders predictably leaves genome-wide signatures and

involves sets of genes with conserved functions that may play general roles in the evolution of social

behavior.

Main Text

Introduction

A longstanding goal of biology is to identify the genetic causes of major phenotypic innovations such as

multicellularity, sociality, and flight (1). When individuals within a population vary for the trait of interest,

identifying associations between genetic variants and traits of interest is relatively straightforward with

experimental mapping populations or genome-wide association studies (2). However, major phenotypic

innovations often only differ between divergent lineages. This makes identifying their genetic

underpinnings more challenging because of the difficulty in distinguishing the relevant genetic differences

from those unrelated to the phenotype of interest (3,4).

Increasingly, comparative genomic approaches have been used to elucidate the genomic underpinnings

of phenotypic innovations, in particular when phenotypes of interest repeatedly evolve in different

evolutionary lineages, so-called convergent phenotypic evolution (5-9). If specific genetic changes and

genomic signatures are common across independent origins of a phenotype of interest, comparative

genomic approaches can provide a powerful means to identify genetic changes that are repeatedly

involved, and thus likely to be especially important in phenotypic innovation. At the same time, these

studies can begin to answer a second complementary longstanding question: to what degree are the

genetic changes underlying independent phenotypic innovation the same, and more generally, how

repeatable is the evolutionary process (10,11)?

Sociality is an exemplar phenotypic innovation that convergently evolved repeatedly and sporadically

across vertebrates, insects, spiders, and crustaceans (12). Sociality and social behavior more generally

are hypothesized to evolve via changes in a set of deeply conserved genes (13-17). However, the degree

to which convergent social evolution in diverse lineages consistently involves evolutionary changes in the

expression or protein-coding sequence of the same or similar genes remains largely unclear. Many

transcriptomic studies have emphasized significantly overlapping sets of genes or molecular pathways

underlying the expression of key social phenotypes in different lineages (18-21), but other transcriptomic

studies emphasize the importance of lineage-specific genes, or genes with lineage-specific expression

(22-24). Social insect studies searching for patterns of molecular evolution associated with the

convergent evolution of sociality have also found differing results. For example, a study comparing

population genomic signatures of selection in two bee and one wasp species, altogether representing two

origins of sociality, found significant overlap in genes experiencing positive selection among all three

species (25). In contrast, the largest comparative genomic study of sociality to date, a comparative

genomic study of ten bee species representing two independent origins and two independent

elaborations of eusociality, found low and insignificant overlap for rapidly evolving genes associated with

social evolution in each lineage (26). The authors concluded that the evolution of sociality in each lineage
involved distinct evolutionary routes (26).

One difficulty with comparative genomic studies of sociality is that in most lineages there are relatively few
origins of sociality, which typically occurred many millions of years ago, making the genomic signatures of
the evolution of sociality difficult to detect (4). For example, social hymenopteran insects (wasps, bees,
and ants), and in particular the bees, are often described as ideal study systems because they include the
full range of social complexity and multiple origins of sociality (26, 27). However, there are only an
estimated total 2-3 independent origins of sociality within the bees, and 7-8 across all social
hymenopterans, and these origins occurred between 20 and 100 million years ago (4, 26, 27). Other
groups with many independent origins of sociality may complement studies in social insects and help to
further identify common genomic signatures of sociality.

Like the social insects, spiders exhibit the full range of social complexity and multiple independent origins
of sociality (28-30). Sociality evolved independently in spiders an estimated 15-16 times (31, 32) and
each independent origin is thought to be relatively recent, at most a few million years ago (33, 34).
Spiders are classified into several categories of social organization, including solitary, subsocial,
prolonged subsocial, and social: solitary species live in individual nests, having dispersed from the egg
sac soon after hatching (28, 32); subsocial spiders have nests that contain a single mother and up to a
few dozen offspring, which may remain together for several instars before dispersing to initiate their own
nests (28, 29); prolonged subsocial spiders form colonies containing a single mother and multiple cohorts
of offspring that remain in their natal nest until late adolescence or sexual maturity before dispersing
independently to found their own colony (29); and social spiders (i.e. non-territorial permanent-social
species) form colonies that contain multiple adult females and offspring that remain in the natal nest
through maturity, mating with each other to produce new generations that reoccupy the natal nest (28). In
social and subsocial species, individuals cooperate in building and maintaining the communal nest,
capture prey cooperatively, and share their food. In social species, individuals additionally exhibit
communal brood care. Social spider species have additional distinct features, including colonies that grow
over multiple generations and in some species can reach colony sizes of tens of thousands of individuals,
female-biased sex ratios, high reproductive skew, and high rates of inbreeding (30, 35).

Spider sociality has a twiggy phylogenetic distribution, where the closest relatives of most social species
are not social. Several previous studies have compared pairs of social species and their closest nonsocial
(i.e. subsocial) relatives within a genus in order to identify genetic consequences of spider sociality and
associated shifts in breeding system from outbreeding to inbreeding (33, 36-39). These studies, together

with a preliminary comparative genomic study including two social species from one genus and several

solitary species (40), identified putative genomic signatures of the evolution of sociality in spiders,

including elevated genome-wide rates of molecular evolution likely caused by relaxed purifying selection

(36-40).

Here, we use a comparative genomic approach to determine whether there are common genomic

signatures across repeated origins of spider sociality that can provide insight into the genomic causes

and consequences of social life. We used 24 total spider species in four lineages that range in social

organization and represent an estimated seven independent origins of sociality (34, 41-43). We focused

our comparisons on testing whether the convergent evolution of sociality in spiders is associated with the

following signatures of convergent molecular evolution: (i) convergent genome-wide patterns of molecular

evolution; (ii) convergent shifts in evolutionary rates for specific genes; and (iii) convergent amino acid

substitutions in specific genes.

Results

Spider samples

We constructed transcriptomes in parallel from 16 ethanol-stored specimens from species in the genera

Anelosimus, *Theridion* (Theriidiade), *Stegodyphus* (Eresidae), and members of the family Sparassidae (SI

Appendix, Table S1). We complemented these new transcriptomes with an additional two available

genomes and six transcriptomes from another eight related species (38, 44-48) (SI Appendix, Table S2).

Altogether, our study included 24 total spider species across the range of spider sociality: eight social

species, one prolonged subsocial species, five subsocial species, and ten solitary species. Since we were

interested in identifying genomic changes associated with sociality, we compared the eight social species

(hereafter labeled as "social") with the 16 species in the remaining categories (hereafter all grouped

together as "nonsocial") (Fig. 1A).

Transcriptome assembly, ortholog identification, and species phylogeny

Following *de novo* assembly and annotation, each of 22 spider transcriptomes had an average of 33,870

transcripts representing the complete or partial protein-coding regions of genes (SI Appendix, Table S2).

Of the 1,066 core arthropod orthologous genes in the BUSCO database (49), 85.2% to 92.1% were

present in the 22 spider transcriptome assemblies (SI Appendix, Table S2), indicating that all

transcriptomes were relatively complete. We also included three available spider genomes from two

social *Stegodyphus* species and a solitary outgroup, *Acanthoscurria geniculata* (SI Appendix, Table S2).

We used a reciprocal hit search strategy with Orthofinder (50) and OrthoDB (https://www.orthodb.org/),

and identified a total of 7,580 orthologous gene groups (OGGs) shared by all 25 spider species, and

1,618 core single-copy orthologs across all species (SI Appendix, Table S2). Finally, we used coalescent

single-copy orthologs to reconstruct a species phylogeny (Fig. 1A, SI Appendix, Fig. S1) that was strongly

supported (bootstrap value = 100) for all nodes, and was mostly consistent with previously-published

phylogenies (34, 41–43, 51) (Fig. S1).

174   Social spiders exhibit convergent genome-wide patterns of accelerated molecular evolution that is

175   caused mainly by the relaxation of selection

As a first test of signatures of convergent molecular evolution, we asked whether social branches had

different rates of genome-wide molecular evolution (i.e. dN/dS) compared to the nonsocial branches. We

estimated the genome-wide ratio of nonsynonymous to synonymous substitution (dN/dS) across the

spider phylogeny, using CodeML in PAML 4.7a (52) with concatenated 1,618 single-copy ortholog data.

Social branches had higher genome-wide dN/dS that estimated by HyPhy 2.5 (53) than nonsocial

branches (likelihood ratio test (LRT), $P < 0.00001$) (Fig. 1B).

Elevated dN/dS can be caused by increased positive selection, relaxed purifying selection, or a

combination of both. RELAX (54) quantifies the degree to which shifts in the distribution of dN/dS across

individual genes or whole genomes are caused by overall *relaxation* of selection (i.e. weakening of both

purifying selection and positive selection, towards neutrality) versus overall *intensification* of selection (i.e.

strengthening of both purifying selection and positive selection, away from neutrality). Specifically, RELAX

models the distribution of three categories of dN/dS (i.e. positive selection, neutral evolution, purifying

selection) across a phylogeny and compares the distributions for foreground (i.e. social) branches to

background (i.e. nonsocial) branches and estimates a parameter K that indicates overall relaxation (K < 1)

or intensification (K > 1). We found that the shifts in genome-wide dN/dS between social and nonsocial

branches were caused mainly by relaxation of both purifying and positive selection (K = 0.80, LRT, $P <$

0.001) (Fig. 1C).

195   Many genes experience convergent patterns of accelerated or decelerated evolution in social

196   spiders

Next we asked if there was evidence for convergent molecular evolution at the gene level. First we used

RELAX (54) to identify genes that experienced relaxed or intensified selection in all social branches

relative to all nonsocial branches. Consistent with the genome-wide RELAX results emphasizing

relaxation of selection, out of 5,243 single-copy orthologs included in the RELAX analysis, more genes

showed evidence of relaxation of selection (598 genes, K < 1, LRT *P* < 0.05, Fig. 2A and 2B, SI Appendix,
Table S3) compared to intensification of selection (275 genes, K > 1, LRT *P* < 0.05, Fig. 2A & 2C, SI
Appendix, Table S3). Genes experiencing significant relaxation of selection in social branches were
enriched for 57 gene ontology (GO) terms, including many related to neural function, neurogenesis, and
cell growth (Fig. 2D, SI Appendix, Table S4), such as neuron projection extension (GO:1990138),
regulation of nervous system process (GO:0031644), regulation of neuron differentiation (GO:0045664),
regulation of extent of cell growth (GO:0061387), and regulation of cell morphogenesis (GO:0022604). In
addition, genes experiencing intensification of selection in social branches were enriched for 71 GO terms,
including terms relating to energy metabolic, immune response, nervous system development,
reproduction and chemosensory behavior (Fig. 2E, SI Appendix, Table S5), such as NADH
dehydrogenase complex assembly (GO:0010257), regulation of protein catabolic process (GO:0042176),
defense response to virus (GO:0051607), synapse organization (GO:0050808), peripheral nervous
system development (GO:0007422), gonad development (GO:0008406), development of primary sexual
characteristics (GO:0045137), chemosensory behavior (GO:0007635) and olfactory behavior
(GO:0042048). The pattern of relaxation of selection in social relative to nonsocial branches can be
caused by relaxation of purifying selection, relaxation of positive selection, or both, and similarly,
intensification of selection can be caused by intensification of purifying selection, intensification of positive
selection, or both. Indeed, inspection of the RELAX results for individual genes showed that genes
experienced relaxation of selection because of either mainly relaxed purifying selection (Fig. 2D), mainly
relaxed positive selection (Fig. 2B), or both (Fig. 2B). Similarly, genes experienced intensification of
selection either because of intensified purifying selection (Fig. 2C), intensified positive selection (Fig. 2C),
or both (Fig. 2C).

Further, we applied the relative evolutionary rate (RER) test implemented in RERconverge (55) that is
specifically designed to test for signatures of convergent molecular evolution underlying convergent
phenotypic evolution. This test examines the rate of protein sequence evolution on every branch across
the phylogeny for each gene, standardized to the distribution of rates across all genes. Genes for which
these standardized rates (RER) are consistently either higher or lower in foreground branches (defined
here as the tips for each social species as well as the branch leading to the common ancestor of *A.*
eximus and *A. domingo*, which may have also been social (42, 51); see Figure 1A) compared to
background branches (i.e. all remaining branches) as experiencing accelerated molecular evolution in
social species or in nonsocial species. We used a phylogenetically restricted permutation strategy,
dubbed permulations, to assess the statistical significance of genes and enriched GO terms (9, 56). This
permulation approach combines phylogenetic simulations and permutations to construct null expectations
for *P* values, and has been shown to be unbiased and more conservative than other approaches such as
permutations alone (9, 56).

We ran 10,000 permulations and identified genes and enriched GO terms (Fig. 3A) that experienced significant convergent shifts in RERs in social compared to nonsocial branches. Specifically, out of 7,345 single-copy orthologs included in REconverge analysis, we identified 170 genes experiencing significantly accelerated molecular evolution in social branches, namely rapidly evolving genes in social spiders (permulation $P < 0.05$) (Fig. 3A and 3B, SI Appendix, Table S6) and we identified 39 genes experiencing significantly decelerated molecular evolution in social branches, namely rapidly evolving genes in nonsocial spiders (permulation $P < 0.05$) (Fig. 3A & 3X, SI Appendix, Table S6). We found 46 GO terms significantly associated with social acceleration (permulation $P < 0.05$), mainly related to nervous system development and neural function (Fig. 3D), such as cerebral cortex development (GO:0021987), neurotransmitter secretion (GO:0007269), memory (GO:0007613) and visual learning (GO:0008542) (SI Appendix, Table S7). We found 22 GO terms significantly associated with social deceleration (permulation $P < 0.05$) (Fig. 3E), mainly associated with metabolism and DNA repair processes, such as carbohydrate metabolic process (GO:0005975), ATP metabolic process (GO:0046034), DNA repair (GO:0006281) and DNA replication checkpoint (GO:0000076) (SI Appendix, Table S8).

To clarify further what patterns of selection contribute to the acceleration and deceleration of protein evolution in social branches, we examined the overlap of lists of genes identified by RERconverge and RELAX. The lists of social-accelerated genes and genes experiencing relaxed selection in social branches overlapped more than expected by chance (permutation $P = 0.00017$; Fig. S2), indicating that elevated rates of protein evolution for some genes in social branches is caused primarily by relaxation of purifying selection. Similarly, lists of social-decelerated genes and genes experiencing intensified selection in social branches were significantly overlapping (permutation $P = 0.015$; SI Appendix, Fig. S2), indicating that reduced rates of protein evolution for some genes in social branches is caused primarily by intensified purifying selection.

Social spiders harbor convergent amino acid substitutions

We asked if there were any amino acid substitutions consistently associated with the evolution of sociality in spiders. We used FADE in HyPhy 2.5 (53) to identify sites experiencing directional selection towards specific amino acids in social compared to nonsocial branches. After filtering sites that were also identified by FADE as experiencing directional selection in nonsocial relative to social branches, we found 907 sites in 169 genes with evidence (Bayes Factor > 100) for directional selection in social relative to background branches. Since we were most interested in identifying sites that showed a repeated pattern of substitution across the social branches, we further focused on sites that were inferred by FADE to have experienced the same substitution at least four separate times in social branches, resulting in a list of 90

sites in 19 genes (SI Appendix, Table S9). We inspected protein alignments for these 19 genes and found
that there were no substitutions that occurred in all social branches and no nonsocial branches. However,
site 64 in the ion channel gene, two pore calcium channel protein 1 (*TPC1*) had seven identical inferred
substitutions from valine to glycine in six social spider species as well as a single nonsocial spider, the
prolonged subsocial *Delena cancerides*, and another two substitutions from valine to alanine in two social
Anelosimus species, *A. oritoyacu* and *A. guacamayos*, which are thought to have independently evolved
sociality (Fig. 4A). Besides *D. cancerides*, all other nonsocial species had the presumed ancestral valine.

As two substitutions at site 64 repeatedly occurred in *TPC1* in social spider branches (Fig. 4A), we asked
if these amino acid changes are predicted to have a functional effect. We generated a three-dimensional
protein structure homology model of TPC1, consisting of 12 transmembrane helices connected by loop
regions, forming a transporter protein-like structure (Fig. 4B). The substrate transportation cavity is
surrounded by four transmembrane helices: TM2, TM3, TM4, TM6, TM7 and TM9 (Fig. 4C and 4D). Site
64 is located on the TM4 domain facing the ion channel and close to the periplasm, indicating that change
from valine to glycine or alanine at this site (Fig. 4E) might affect the activity of ion transport.

Discussion

We used a comparative genomic approach with 24 spider species, representing seven independent and
likely recent origins of sociality, in order to answer a longstanding question: what genetic changes
underpin the convergent evolution of sociality? We found the following genomic signatures of convergent
phenotypic evolution: (i) Supporting previous results in spiders and insects, we found elevated genome-
wide rates of molecular evolution in social species compared to nonsocial species, resulting mainly from
relaxation of selection, which is an expected genomic consequence of living in inbred social groups with
reduced effective population size. (ii) We identified hundreds of genes that experienced shifts in patterns
of molecular evolution during the convergent evolution of sociality. These genes were enriched for Gene
Ontology (GO) terms related to neural function, neurogenesis, and behavior, as well as immune function,
growth, and metabolism, indicating that changes in these functions are likely important during or following
the transition to social life. (iii) We found evidence for specific substitutions that tended to occur in social
species, with substitutions at one site in the calcium channel gene *TPC1* found almost exclusively in
social species. Overall, our results suggest that patterns of natural selection that lead to, and result from
the repeated evolution of sociality in spiders, tend to act on sets of genes with similar functions and also
repeatedly lead to genome-wide signatures.

Consistent with several previous studies in social spiders (33, 36-39) and social insects (26, 57, 58), we

found that social branches tend to experience elevated genome-wide rates of molecular evolution (i.e.

dN/dS) compared to nonsocial branches. Such elevated genome-wide rates of molecular evolution are

hypothesized to be caused by reduced effective population size in social species, which results in

relaxation of purifying selection and positive selection experienced by genes (33, 36-39). In social spiders,

factors causing this reduced effective population size include inbreeding, female-biased sex ratios, and

reproductive skew. Supporting this hypothesis, we found that the genome-wide pattern was primarily

driven by the relaxation of both purifying selection and positive selection (Fig. 1C), so that genes in social

branches tend to experience more relaxed, neutral evolution when compared to genes in nonsocial

branches. This genome-wide pattern is likely to be a longer-term *consequence* of the switch from

outbreeding to inbreeding, and to female-biased sex ratios, which likely occurred after the origin of

sociality. There was also variation for genome-wide dN/dS that did not depend on sociality (Fig. 1A),

which might be explained by lineage-specific differences in effective population size or other factors.

Similarly, in a previous study in hymenopteran insects, social branches experienced elevated genome-

wide dN/dS compared to nonsocial branches, but most variation existed between lineages, with bees --

regardless social organization -- having the highest dN/dS (58).

We identified approximately 200 genes that tended to experience convergent shifts in evolutionary rates

during the transition to social life (Fig. 2A). Such gene-wide shifts could be caused by changes in the

patterns of purifying selection, positive selection or both (Fig. 2D). We separately identified hundreds of

genes that experienced an average relaxation or intensification of selection during the transition to social

life (Fig. 2C). The list of genes experiencing *elevated* rates of protein evolution significantly overlapped

with the list of genes experiencing *relaxation* of selection (Fig. S2), indicating that the pattern of gene-

wide elevated protein evolution that we detected is often caused by the relaxation of purifying selection

during the transition to sociality. Similarly, the list of genes experiencing *decreased* rates of protein

evolution significantly overlapped with the list of genes experiencing *intensification* of selection (Fig. S2),

indicating that gene-wide decreased protein evolution is largely caused by intensified purifying selection

during the transition to sociality.

The genes that we identified as experiencing convergent patterns of molecular evolution were not

randomly distributed with respect to annotated gene function, as would be expected if the gene-wide

patterns were driven only by genome-wide relaxation of selection associated with reduced effective

population size. Instead, these genes were enriched for many specific annotated functional categories,

notably those related to neural function, neurogenesis, and behavior, as well as immune function,

metabolism, development, and reproduction (Fig. 2 & 3). We can infer that genes and molecular

pathways with these functions likely played important roles in the evolution of sociality, either in initial

genetic changes directly associated with the evolution of social traits and the transition to social life, or

genetic changes that were later consequences of social life or the elaboration of social phenotypes.

Genes that experienced shifts in patterns of molecular evolution during the transition to sociality were

consistently enriched for functions relating to neural function, neurogenesis, and behavior (SI Appendix,

Table S4, S5, S6), indicating that changes in these functions were likely particularly important. Genes

experiencing relaxation of selection in social branches were enriched for many GO terms associated with

neural function or neural development, including neurotransmitter receptor transport and regulation of

neuron differentiation (Fig. 2B). Similarly, genes tending to experience elevated evolutionary rates in

social branches were enriched with functional annotations such as cerebral development,

neurotransmitter secretion, visual learning, and memory (Fig. 3B). Genes experiencing intensified

selection were also enriched for synapse organization and peripheral nervous system development, as

well as behavior-related terms such as olfactory behavior and negative regulation of locomotion (Fig. 2C).

A fundamental phenotypic difference between social and nonsocial species is the extent of social

interactions with conspecifics, which would have required increased behavioral tolerance towards

conspecifics as an important preliminary change necessary for the evolution of cooperative group living in

social spiders (35, 59, 60) and other animals (61). Furthermore, shifts in behavioral development and the

timing of social behaviors, reproductive behavior, and dispersal behaviors, are also hypothesized to be

critical in the evolution of sociality in spiders (30, 35) and insects (15). Finally, changes in the size of

certain brain regions and sensory organs are hypothesized to be important for detecting and processing

social signals and social information (62). The changes that we detected in genes associated with neural

function, neurogenesis, and behavior may underlie changes in these important behavioral and neural

phenotypes.

Genes involved in neurotransmitter secretion and neurotransmitter receptor transport stand out as

interesting candidates influencing the evolution of spider social behavior because many genes annotated

with these functions experienced significant shifts in evolutionary rates in social spiders and have also

previously been implicated in the regulation of social behaviors in other animals (SI Appendix, Table S6).

For example, the gene *adenosine A$_{2A}$ receptor* (*adora2a*), which experienced relaxed selection in social

spiders, regulates glutamate and dopamine release and influences motivation for social interactions in

mice (63, 64), and variation in *adora2a* has been associated with mood and anxiety disorders in humans

(65, 66). The gene *dystrophin, isoforms A/C/F/G/H* (*dys*), which experienced accelerated evolution in

social spiders, affects social behavior, communication, and synaptic plasticity in mice (67, 68), and

mutations in *dys* are associated with autism and intellectual disability in humans (68). The gene *osmotic*

avoidance abnormal protein 3 (*osm-3*), which experienced relaxed selection in social spiders, regulates

social feeding and aggregation behavior in nematodes (69). The gene *calsyntenin-1* (*clstn1*), which

experienced relaxed selection and accelerated evolution in social spiders, regulates synapse formation
and synapse plasticity in mice (70), is essential for chemotaxis learning in nematodes (71), and is a
candidate for Fragile X syndrome and autism in humans (70, 72). Finally, the genes *synaptosomal-*
associated protein 25kDa (*snap-25*), *synaptobrevin*, and *syntaxin-1A* (*stx1a*) interact to regulate the
secretion of neurotransmitters and neuromodulators (73). Each of these has been implicated in the
expression of social behavior in various animals (74-76), and also implicated in abnormal behavioral
phenotypes in humans, including autism and schizophrenia (77, 78). Each of these experienced
significant relaxed or accelerated rates of molecular evolution in social spiders, except for *stx1a*
(RERconverge permulation P = 0.06). Further studies will be necessary to validate the function and
mechanism of these candidates influencing spider social behavior.

Besides critical changes in behavioral and neural phenotypes, transitions to social living are also
associated with several other phenotypic changes. Social living is hypothesized to be associated with
increased disease pressure, although researchers have found mixed support for genetic changes in
immune-related genes in social insects compared to solitary insects (79-81). Genes experiencing
intensified selection in social spiders were enriched for several immune-related GO terms such as
defense response to virus and defense response to other organisms, which is consistent with social
spiders experiencing increased disease pressure. Transitions to social living in spiders are also thought to
be associated with shifts in development, including increasing paedomorphosis, and a trend towards
smaller brood sizes (35) and smaller body size (82). Genes experiencing relaxed selection in social
spiders were enriched for GO terms including regulation of cell size and developmental growth, which
could underlie these phenotypic changes. Finally, another phenotypic consequence of transitions to
sociality in spiders, which may occur over longer time scales, is the change of breeding system from
outbreeding to inbreeding and the evolution of female-biased sex ratios (30, 35). Genes experiencing
intensified selection were enriched for GO terms including gonad development and male mating behavior,
which could be associated with these shifts in the breeding system. Metabolism-related genes and gene
functions have also frequently been implicated in the expression of various types of social behavior in
many animals (20, 83), and comparative genomic studies in social insects have identified metabolism-
associated genes as experiencing accelerated evolution in eusocial bees (84) and ants (85). Consistent
with these previous studies, we found that genes experiencing accelerated evolution and intensified
selection in spiders were also enriched for metabolism-related functions (e.g., the GO terms ATP
metabolic process and regulation of cellular protein catabolic process).

In addition to genome-wide and gene-wide signatures, we also identified some signatures of site-specific
convergent molecular evolution. Specifically, 90 sites in 19 genes showed evidence for directional
selection for specific amino acids in social relative to nonsocial species, and genes with such signatures

of specific substitutions were enriched for several GO terms related to nervous system development and

synapse organization. However, inspection of the amino acid alignments for these genes showed that

amino acid substitutions that occurred repeatedly in social branches also tended to be found in one or

more nonsocial branches. An amino acid substitution at position 64 in the calcium channel gene *two pore*

calcium channel protein 1 (*TPC1*) was found in all eight social species and just one nonsocial species.

Interestingly, the only nonsocial species that also had a substitution at this position was *Delena*

cancerides, which stands as being relatively more socially complex: *D. cancerides* is classified as

prolonged subsocial because individuals spend most of their lives living in social groups with their mother

and siblings, while the other nonsocial species in our study are classified as solitary or subsocial.

Homology modeling of the protein structure of *TPC1* indicated that position 64 is located in a

transmembrane domain facing the ion channel and close to the periplasm and may affect ion transport

activity. Ion channel genes are widely expressed in the nervous system and influence a variety of

behavioral phenotypes (86), including social behaviors such as depression-related behavior (87),

acquired fear-related behavior (88), and courtship behavior (89). Our finding suggests that the amino acid

substitution at position 64 in TPC1 across social species might contribute to a key social trait, and this

gene and specific position is an interesting candidate for future functional tests.

We used phylogenetic comparative approaches to identify genome-wide, gene-wide, and site-specific

signatures of convergent molecular evolution. We identified hundreds of genes that tended to experience

different rates of molecular evolution in the replicate social branches compared to background nonsocial

branches, and these genes were enriched for certain functions. However, we did not identify genes or

sites within genes that *always* experienced evolutionary shifts in each of the seven branches that

independently evolved sociality. For example, inspection of patterns of gene-wide relative evolutionary

rates (e.g., Fig. 3D, 3E) shows that genes with a relatively strong association between social status and

evolutionary rate still showed a lot of variation within each category. This means that we identified certain

genes and functions that tended to experience shifts in evolutionary rates, but the exact details differed

somewhat between the seven independent origins of sociality. Thus, specific substitutions or changes in

specific genes are not required for the evolution of spider sociality, but changes do tend to occur in

certain sets of genes or molecular pathways. Put another way, the strong statistical signatures across

seven independent origins of sociality indicate that social spiders tend to travel similar evolutionary paths

in terms of the functional genetic changes and genome-wide consequences, but the precise genetic

details do vary. In practice, these results also emphasize that for complex phenotypes such as sociality,

datasets representing many independent origins may be necessary to have the statistical power to detect

these signatures. When samples representing multiple independent origins are not available, lineage-

specific processes such as lineage-specific adaptations or historical contingency can be confounded with

genetic changes that are actually associated with the phenotype of interest (3, 4, 40, 90).

Our study adds to the growing list of studies that use comparative genomic approaches to elucidate the

molecular underpinnings of convergent evolution and the degree to which similar phenotypic outcomes

result from the same or similar genetic changes (5-9). Collectively, these studies show that some

convergent phenotypes, especially those involving relatively simple physiological adaptation, such as the

evolution of antibiotic resistance, pesticide resistance, or resistance to neurotoxins (91), often predictably

involve the same substitutions or changes in one family of genes. Other phenotypic changes that are

relatively more complex, such as the evolution of sociality, still have identifiable genomic signatures

across independent origins, but these signatures are not as highly predictable in the sense that they

always involve the same substitution or changes in the same genes, but instead involve changes in

genes with similar functions.

Finally, our study emphasizes the potential for ethanol-preserved specimens to be used in transcriptome-

based comparative genomics research. Increasingly, ethanol-preserved specimens from collections are

being used for DNA-based phylogenomic and comparative genomic studies (92), but RNA-based studies

have usually been restricted to samples collected in liquid nitrogen or RNA later, and stored at -80 C (93).

Our study builds on previous studies that successfully extracted RNA from ethanol-stored samples and

subsequently conducted transcriptome-based phylogenomics (94, 95). We extracted RNA from spider

specimens that had been preserved in ethanol at room temperature for up to 11 years (SI Appendix,

Table S1), and generated relatively high quality RNA sequence data for subsequent comparative

molecular evolution analyses (SI Appendix, Table S2). Collectively, our study highlights the heretofore

unappreciated potential of transcriptomics using ethanol-preserved specimens for comparative genomics

and phylogenomics analyses, especially in species with large genomes, such as spiders, where

sequencing and assembling whole genomes remains challenging.

Materials and Methods

Sample collection

We collected spider specimens of 16 species that range in social organization, including social, prolonged

subsocial, subsocial and solitary, and stored the samples in 95% ethanol (Decon Labs, USA). Specifically,

this set of spider specimens includes five social and three subsocial species from the genus *Anelosimus*,

one social and four solitary species from the genus *Theridion*, and one prolonged subsocial and two

solitary species from the family Sparassidae (Fig.1A, SI Appendix, Table S1). Full details of this as well as

other materials and methods are provided in the SI Appendix.

Acknowledgements

We would like to thank Drs. Sarah Rose, Marc Milne, and Jonathan Pruitt for kindly providing ethanol-preserved spider specimens. This work was funded by the National Institutes of Health Grant GM115509 to TAL.

References

1. A. Wagner, The molecular origins of evolutionary innovations. *Trends Genet.* 27, 397–410 (2011).
2. T. F. C. Mackay, E. A. Stone, J. F. Ayroles, The genetics of quantitative traits: challenges and prospects. *Nat. Rev. Genet.* 10, 565–577 (2009).
3. W. Enard, The Molecular Basis of Human Brain Evolution. *Curr. Biol.* 26, R1109–R1117 (2016).
4. T. A. Linksvayer, B. R. Johnson, Re-thinking the social ladder approach for elucidating the evolution and molecular basis of insect societies. *Curr Opin Insect Sci* 34, 123–129 (2019).
5. Y. Zhen, M. L. Aardema, E. M. Medina, M. Schumer, P. Andolfatto, Parallel molecular evolution in an herbivore community. *Science* 337, 1634–1637 (2012).
6. J. Parker, et al., Genome-wide signatures of convergent evolution in echolocating mammals. *Nature* 502, 228–231 (2013).
7. R. Partha, et al., Subterranean mammals show convergent regression in ocular genes and enhancers, along with adaptation to tunneling. *Elife* 6 (2017).
8. C. R. Feldman, E. D. Brodie Jr, E. D. Brodie 3rd, M. E. Pfrender, Constraint shapes convergence in tetrodotoxin-resistant sodium channels of snakes. *Proc. Natl. Acad. Sci. U. S. A.* 109, 4556–4561 (2012).
9. A. Kowalczyk, R. Partha, N. L. Clark, M. Chikina, Pan-mammalian analysis of molecular constraints underlying extended lifespan. *Elife* 9 (2020).
10. E. B. Rosenblum, C. E. Parent, E. E. Brandt, The molecular basis of phenotypic convergence. *Annu. Rev. Ecol. Evol. Syst.* 45, 203-226 (2014).
11. D. I. Bolnick, R. D. H. Barrett, K. B. Oke, D. J. Rennison, Y. E. Stuart, (Non) parallel evolution. *Annu. Rev. Ecol. Evol. Syst.* 49, 303–330 (2018).
12. D. R. Rubenstein, P. Abbot, Comparative Social Evolution (Cambridge University Press, 2017).
13. A. L. Toth, G. E. Robinson, Evo-devo and the evolution of social behavior. *Trends Genet.* 23, 334–341 (2007).
14. G. V. Amdam, A. Csondes, M. K. Fondrk, R. E. Page Jr, Complex social behaviour derived from maternal reproductive traits. *Nature* 439, 76–78 (2006).
15. T. A. Linksvayer, M. J. Wade, The evolutionary origin and elaboration of sociality in the aculeate Hymenoptera: maternal effects, sib-social effects, and heterochrony. *Q. Rev. Biol.* 80, 317–336 (2005).
16. L. A. O'Connell, H. A. Hofmann, The vertebrate mesolimbic reward system and social behavior network: a comparative synthesis. *J. Comp. Neurol.* 519, 3599–3639 (2011).
17. G. E. Robinson, C. M. Grozinger, C. W. Whitfield, Sociogenomics: social life in molecular terms. *Nat. Rev. Genet.* 6, 257–270 (2005).

18. A. J. Berens, J. H. Hunt, A. L. Toth, Comparative transcriptomics of convergent evolution:
different genes but conserved pathways underlie caste phenotypes across lineages of eusocial
insects. *Mol. Biol. Evol.* 32, 690–703 (2015).

19. R. L. Young, et al., Conserved transcriptomic profiles underpin monogamy across vertebrates.
Proc. Natl. Acad. Sci. U. S. A. 116, 1331–1336 (2019).

20. C. C. Rittschof, et al., Neuromolecular responses to social challenge: common mechanisms
across mouse, stickleback fish, and honey bee. *Proc. Natl. Acad. Sci. U. S. A.* 111, 17929–17934
(2014).

21. M. R. Warner, L. Qiu, M. J. Holmes, A. S. Mikheyev, T. A. Linksvayer, Convergent eusocial
evolution is based on a shared reproductive groundplan plus lineage-specific plastic genes. *Nat.*
Commun. 10, 2651 (2019).

22. B. R. Johnson, N. D. Tsutsui, Taxonomically restricted genes are associated with the evolution of
sociality in the honey bee. *BMC Genomics* 12, 164 (2011).

23. W. C. Jasper, et al., Large-scale coding sequence change underlies the evolution of
postdevelopmental novelty in honey bees. *Mol. Biol. Evol.* 33, 1379 (2016).

24. S. Sumner, The importance of genomic novelty in social evolution. *Mol. Ecol.* 23, 26–28 (2014).

25. K. A. Dogantzis, et al., Insects with similar social complexity show convergent patterns of
adaptive molecular evolution. *Sci. Rep.* 8, 10388 (2018).

26. K. M. Kapheim, et al., Genomic signatures of evolutionary transitions from solitary to group living.
Science 348, 1139–1143 (2015).

27. S. D. Kocher, R. J. Paxton, Comparative methods offer powerful insights into social evolution in
bees. *Apidologie* 45, 289–305 (2014).

28. L. Avilés, J. Guevara, "Sociality in Spiders" in Comparative Social Evolution, R. Rubenstein, P.
Abbot, Eds. (Cambridge University Press, 2017), pp. 188–223.

29. E. C. Yip, L. S. Rayor, Maternal care and subsocial behaviour in spiders. Biol. Rev. Camb. Philos.
Soc. 89, 427–449 (2014).

30. Y. Lubin, T. Bilde, "The Evolution of Sociality in Spiders" in Advances in the Study of Behavior,
(Academic Press, 2007), pp. 83–145.

31. I. Agnarsson, L. Avilés, J. A. Coddington, W. P. Maddison, Sociality in theridiid spiders: repeated
origins of an evolutionary dead end. *Evolution* 60, 2342–2351 (2006).

32. L. Avilés, "Social Spiders" in Encyclopedia of Social Insects, C. K. Starr, Ed. (Springer, 2020), pp.
1–10.

33. I. Agnarsson, L. Avilés, W. P. Maddison, Loss of genetic variability in social spiders: genetic and
phylogenetic consequences of population subdivision and inbreeding. *J. Evol. Biol.* 26, 27–37
(2013).

34. J. Johannesen, Y. Lubin, D. R. Smith, T. Bilde, J. M. Schneider, The age and evolution of
sociality in Stegodyphus spiders: a molecular phylogenetic perspective. *Proc. Biol. Sci.* 274, 231–
237 (2007).

35. L. Avilés, 23• Causes and consequences of cooperation and permanent-sociality in spiders. The
evolution of social behavior in insects and arachnids, 476–498 (1997).

36. V. Settepani, et al., Evolution of sociality in spiders leads to depleted genomic diversity at both
population and species levels. *Mol. Ecol.* 26, 4197–4210 (2017).

37. V. Settepani, J. Bechsgaard, T. Bilde, Phylogenetic analysis suggests that sociality is associated
with reduced effectiveness of selection. *Ecol. Evol.* 6, 469–477 (2016).

38. J. Bechsgaard, et al., Evidence for faster X chromosome evolution in spiders. *Mol. Biol. Evol.* 36,
1281–1293 (2019).

39. T. M. Mattila, J. S. Bechsgaard, T. T. Hansen, M. H. Schierup, T. Bilde, Orthologous genes
identified by transcriptome sequencing in the spider genus Stegodyphus. *BMC Genomics* 13, 70
(2012).

40. C. Tong, G. M. Najm, N. Pinter-Wollman, J. N. Pruitt, T. A. Linksvayer, Comparative Genomics
Identifies Putative Signatures of Sociality in Spiders. *Genome Biol. Evol.* 12, 122–133 (2020).

41. M. A. Arnedo, I. Agnarsson, R. G. Gillespie, Molecular insights into the phylogenetic structure of
the spider genus Theridion (Araneae, Theridiidae) and the origin of the Hawaiian Theridion-like
fauna. *Zool. Scr.* 36, 337–352 (2007).

42. Y. Luo, et al., Global Diversification of Anelosimus Spiders Driven by Long Distance Overwater
Dispersal and Neogene Climate Oscillations. *Syst. Biol.* 69 (6), 1122-1136 (2020).

43. I. Agnarsson, L. S. Rayor, A molecular phylogeny of the Australian huntsman spiders
(Sparassidae, Deleninae): implications for taxonomy and social behaviour. *Mol. Phylogenet. Evol.*
69, 895–905 (2013).

44. K. W. Sanggaard, et al., Spider genomes provide insight into composition and evolution of venom
and silk. *Nat. Commun.* 5, 3765 (2014).

45. P. J. P. Croucher, M. S. Brewer, C. J. Winchell, G. S. Oxford, R. G. Gillespie, De novo
characterization of the gene-rich transcriptomes of two color-polymorphic spiders, Theridion
grallator and T. californicum (Araneae: Theridiidae), with special reference to pigment genes.
BMC Genomics 14, 862 (2013).

46. L. Shao, S. Li, Early Cretaceous greenhouse pumped higher taxa diversification in spiders. *Mol.*
Phylogenet. Evol. 127, 146–155 (2018).

47. R. Fernández, et al., Phylogenomics, Diversification Dynamics, and Comparative Transcriptomics
across the Spider Tree of Life. *Curr. Biol.* 28, 2190–2193 (2018).

48. S. Liu, A. Aageaard, J. Bechsgaard, T. Bilde, DNA Methylation Patterns in the Social Spider,
Stegodyphus dumicola. *Genes* 10 (2019).

49. F. A. Simão, R. M. Waterhouse, P. Ioannidis, E. V. Kriventseva, E. M. Zdobnov, BUSCO:
assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics 31, 3210–3212 (2015).

50. D. M. Emms, S. Kelly, OrthoFinder: solving fundamental biases in whole genome comparisons
dramatically improves orthogroup inference accuracy. *Genome Biol.* 16, 157 (2015).

51. I. Agnarsson, W. P. Maddison, L. Avilés, The phylogeny of the social Anelosimus spiders
(Araneae: Theridiidae) inferred from six molecular loci and morphology. *Mol. Phylogenet. Evol.*
43, 833–851 (2007).

52. Z. Yang, PAML 4: phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* 24, 1586–1591
(2007).

53. S. L. Kosakovsky Pond, et al., HyPhy 2.5—A Customizable Platform for Evolutionary Hypothesis
Testing Using Phylogenies. *Mol. Biol. Evol.* 37, 295–299 (2020).

54. J. O. Wertheim, B. Murrell, M. D. Smith, S. L. Kosakovsky Pond, K. Scheffler, RELAX: detecting
relaxed selection in a phylogenetic framework. *Mol. Biol. Evol.* 32, 820–832 (2015).

55. A. Kowalczyk, et al., RERconverge: an R package for associating evolutionary rates with
convergent traits. *Bioinformatics* 35, 4815–4817 (2019).

56. E. Saputra, A. Kowalczyk, L. Cusick, N. Clark, M. Chikina, Phylogenetic Permulations: a
statistically rigorous approach to measure confidence in associations between phenotypes and
genetic elements in a phylogenetic context. Cold Spring Harbor Laboratory, 2020.10.14.338608
(2020).

57. J. Romiguier, et al., Population genomics of eusocial insects: the costs of a vertebrate-like
effective population size. *J. Evol. Biol.* 27, 593–603 (2014).

58. A. Weyna, J. Romiguier, Relaxation of purifying selection suggests low effective population size
in eusocial Hymenoptera and pollinating bees. *bioRxiv,* 2020.04.14.038893 (2020).

59. C. Viera, I. Agnarsson, "Parental Care and Sociality" in Behaviour and Ecology of Spiders:
Contributions from the Neotropical Region, C. Viera, M. O. Gonzaga, Eds. (Springer International
Publishing, 2017), pp. 351–381.

60. E. J. Kullmann, Evolution of social behavior in spiders (Araneae; Eresidae and Theridiidae). Am.
Zool. 12, 419–426 (1972).

61. N. Lin, C. D. Michener, Evolution of Sociality in Insects. *Q. Rev. Biol.* 47, 131–159 (1972).

62. B. Wittwer, et al., Solitary bees reduce investment in communication compared with their social
relatives. *Proc. Natl. Acad. Sci. U. S. A.* 114, 6569–6574 (2017).

63. L. López-Cruz, et al., Adenosine A2A receptor deletion affects social behaviors and anxiety in
mice: Involvement of anterior cingulate cortex and amygdala. *Behav. Brain Res.* 321, 8–17
(2017).

64. L. López-Cruz, et al., Ethanol and Caffeine Effects on Social Interaction and Recognition in Mice:
Involvement of Adenosine A2A and A1 Receptors. *Front. Behav. Neurosci.* 10, 206 (2016).

65. S. Oliveira, et al., Impact of genetic variations in ADORA2A gene on depression and symptoms: a
cross-sectional population-based study. *Purinergic Signal.* 15, 37–44 (2019).

66. D. van Calker, K. Biber, K. Domschke, T. Serchov, The role of adenosine receptors in mood and
anxiety disorders. *J. Neurochem.* 151, 11–27 (2019).

67. R. Miranda, et al., Altered social behavior and ultrasonic communication in the dystrophin-
deficient mdx mouse model of Duchenne muscular dystrophy. *Mol. Autism* 6, 60 (2015).

68. F. Daoud, et al., Role of mental retardation-associated dystrophin-gene product Dp71 in
excitatory synapse organization, synaptic plasticity and behavioral functions. *PLoS One* 4, e6574
(2008).

69. M. de Bono, D. M. Tobin, M. W. Davis, L. Avery, C. I. Bargmann, Social feeding in Caenorhabditis
elegans is induced by neurons that detect aversive stimuli. *Nature* 419, 899–903 (2002).

70. K. Cheng, et al., Calsyntenin-1 Negatively Regulates ICAM5 Accumulation in Postsynaptic
Membrane and Influences Dendritic Spine Maturation in a Mouse Model of Fragile X Syndrome.
Front. Neurosci. 13, 1098 (2019).

71. D. D. Ikeda, et al., CASY-1, an ortholog of calsyntenins/alcadeins, is essential for learning in
Caenorhabditis elegans. *Proc. Natl. Acad. Sci. U. S. A.* 105, 5260–5265 (2008).

72. S. V. Ranneva, K. S. Pavlov, A. V. Gromova, T. G. Amstislavskaya, T. V. Lipina, Features of
emotional and social behavioral phenotypes of calsyntenin2 knockout mice. *Behav. Brain Res.*
332, 343–354 (2017).

73. N. A. Ramakrishnan, M. J. Drescher, D. G. Drescher, Direct interaction of otoferlin with syntaxin
1A, SNAP-25, and the L-type voltage-gated calcium channel Cav1.3. *J. Biol. Chem.* 284, 1364–
1372 (2009).

74. I. Corradini, et al., Epileptiform activity and cognitive deficits in SNAP-25+/- mice are normalized
by antiepileptic drugs. *Cereb. Cortex* 24, 364–376 (2014).

75. T. Fujiwara, M. Sanada, T. Kofuji, K. Akagawa, Unusual social behavior in HPC-1/syntaxin1A
knockout mice is caused by disruption of the oxytocinergic neural system. *J. Neurochem.* 138,
117–123 (2016).

76. S. D. Kocher, et al., The genetic basis of a social polymorphism in halictid bees. *Nat. Commun.* 9,
4338 (2018).

77. D. Braida, et al., Association between SNAP-25 gene polymorphisms and cognition in autism:
functional consequences and potential therapeutic strategies. *Transl. Psychiatry* 5, e500 (2015).

78. I. Corradini, C. Verderio, M. Sala, M. C. Wilson, M. Matteoli, SNAP-25 in neuropsychiatric
disorders. *Ann. N. Y. Acad. Sci.* 1152, 93–99 (2009).

79. K. Meusemann, J. Korb, M. Schughart, F. Staubach, No evidence for single-copy immune-gene
specific signals of selection in termites. *Front. Ecol. Evol.* 8 (2020).

80. L. Viljakainen, et al., Rapid evolution of immune proteins in social insects. *Mol. Biol. Evol.* 26,
1791–1801 (2009).

81. S. Otani, N. Bos, S. H. Yek, Transitional Complexity of Social Insect Immunity. *Frontiers in
Ecology and Evolution* 4, 69 (2016).

82. L. Grinsted, et al., Prey to predator body size ratio in the evolution of cooperative hunting—a
social spider test case. *Dev. Genes Evol.* 230, 173–184 (2020).

83. C. C. Rittschof, G. E. Robinson, Behavioral Genetic Toolkits: Toward the Evolutionary Origins of
Complex Phenotypes. *Curr. Top. Dev. Biol.* 119, 157–204 (2016).

84. S. H. Woodard, et al., Genes involved in convergent evolution of eusociality in bees. *Proc. Natl.
Acad. Sci. U. S. A.* 108, 7472–7477 (2011).

85. J. Roux, et al., Patterns of positive selection in seven ant genomes. *Mol. Biol. Evol.* 31, 1661–
1685 (2014).

86. N. S. Atkinson, et al., Molecular separation of two behavioral phenotypes by a mutation affecting
the promoters of a Ca-activated K channel. *J. Neurosci.* 20, 2988–2993 (2000).

87. M. W. Coryell, et al., Acid-sensing ion channel-1a in the amygdala, a novel therapeutic target in
depression-related behavior. *J. Neurosci.* 29, 5381–5388 (2009).

88. J. A. Wemmie, et al., Overexpression of acid-sensing ion channel 1a in transgenic mice increases
acquired fear-related behavior. *Proc. Natl. Acad. Sci. U. S. A.* 101, 3621–3626 (2004).

89. Y. Ding, A. Berrocal, T. Morita, K. D. Longden, D. L. Stern, Natural courtship song variation
caused by an intronic retroelement in an ion channel gene. *Nature* 536, 329–332 (2016).

90. F. Delsuc, M.-K. Tilak, Naked but not Hairless: the pitfalls of analyses of molecular adaptation
based on few genome sequence comparisons. *Genome Biol. Evol.* 7, 768–774 (2015).

91. J. W. McGlothlin, et al., Historical Contingency in a Multigene Family Facilitates Adaptive
Evolution of Toxin Resistance. *Curr. Biol.* 26, 1616–1621 (2016).

92. A. E. Z. Short, T. Dikow, C. S. Moreau, Entomological Collections in the Age of Big Data. *Annu.
Rev. Entomol.* 63, 513–530 (2018).

93. Z. Wang, M. Gerstein, M. Snyder, RNA-Seq: a revolutionary tool for transcriptomics. *Nat. Rev.*
Genet. 10, 57–63 (2009).

94. H. M. Gough, J. M. Allen, E. F. A. Toussaint, C. G. Storer, A. Y. Kawahara, Transcriptomics
illuminate the phylogenetic backbone of tiger beetles. *Biol. J. Linn. Soc. Lond.* 129, 740–751
(2020).

95. A. L. Bazinet, M. P. Cummings, K. T. Mitter, C. W. Mitter, Can RNA-Seq resolve the rapid
radiation of advanced moths and butterflies (Hexapoda: Lepidoptera: Apoditrysia)? An
exploratory study. *PLoS One* 8, e82615 (2013).

    Figures

701
Figure 1. Spiders experienced elevated genome-wide molecular evolution during the transition to sociality, driven by the relaxation of purifying and positive selection. (A) phylogram showing the phylogenetic relationships among the 24 species included in the study. Branch color shows estimated genome-wide rate of molecular evolution (dN/dS) and branch length is proportional to sequence divergence. Social and nonsocial spider taxa are indicated by orange or sky blue dots, respectively. X's indicate independent origins of sociality as inferred in previous studies (31, 32). (B) barplot depicting the genome-wide dN/dS ratios for social and nonsocial branches, estimated by HyPhy. ** $P□<□0.001$ (likelihood ratio test, LRT). (C) plot showing the pattern of selection experienced across sites in the genomes of social and nonsocial spiders, estimated with the Partitioned Descriptive model in RELAX (54). The distribution of dN/dS across sites in the genomes are illustrated by three categories of dN/dS for social (orange) and nonsocial (sky blue) branches. The vertical dashed line at dN/dS = 1 represents neutral evolution, bars at dN/dS > 1 represent sites experiencing positive selection, and bars at dN/dS <1 represent sites experiencing purifying selection. The arrows show the direction of change in dN/dS between nonsocial and social branches, demonstrating relaxation of purifying and positive selection associated with the transition to sociality. K < 1 indicates genome-wide relaxation of selection.
717

A

Genes and GO terms experiencing relaxation and intensification in social branches

Genes

N=598   N=275

☐ Relaxation ☐ Intensification

Enriched GO terms

N=57   N=71

☐ Relaxation ☐ Intensification

B

Ontology for genes experiencing relaxation of selection in social branches

neural function and neurogenesis

negative regulation of axonogenesis
astrocyte differentiation
excitatory postsynaptic potential
negative regulation of neuron projection development
chemical synaptic transmission, postsynaptic
regulation of axonogenesis
regulation of neuron projection development
neuron projection extension
regulation of axon extension
regulation of nervous system process
axon extension
neurotransmitter receptor transport
regulation of neuron differentiation

cell growth

negative regulation of cellular component organization
regulation of cell morphogenesis involved in differentiation
negative regulation of cell projection organization
regulation of extent of cell growth
regulation of cell morphogenesis
negative regulation of cell morphogenesisinvolved in differentiation
developmental growth

-LOG10(P-value)

C

Ontology for genes experiencing intensification of selection in social branches

energy metabolism

protein insertion into mitochondrial membrane
regulation of ubiquitin-dependent protein catabolic process
establishment of protein localization to mitochondrial membrane
regulation of cellular protein catabolic process
NADH dehydrogenase complex assembly

immune response

response to biotic stimulus
response to external biotic stimulus
response to other organism
response to virus
defense response to virus
defense response to other organism

neurogenesis

synapse organization
peripheral nervous system development

behavior

olfactory behavior
male mating behavior
chemosensory behavior
negative regulation of locomotion
regulation of locomotion

reproduction

gonad development
development of primary sexual characteristics

-LOG10(P-value)

D   Examples of genes experiencing relaxation in social branches

relaxed purifying selection
PGGT1B   K = 0.97 P = 0.0001
Social Nonsocial
Proportion of sites dN/dS

relaxed positive selection
NINJ1   K = 0.58 P = 0.0012
Social Nonsocial
Proportion of sites dN/dS

relaxed purifying and positive selection
AGRN   K = 0.63 P = 0.0043
Social Nonsocial
Proportion of sites dN/dS

E   Examples of genes experiencing intensification in social branches

intensified purifying selection
CenG1A   K = 1.46 P = 0.0099
Social Nonsocial
Proportion of sites dN/dS

intensified positive selection
MRRF   K = 1.84 P = 0.0037
Social Nonsocial
Proportion of sites dN/dS

intensified purifying and positive selection
PHRF1   K = 1.28 P = 0.0013
Social Nonsocial
Proportion of sites dN/dS

Figure 2. RELAX identifies a substantial set of genes that tend to experience relaxation or
intensification of selection during the transition to sociality. (A) the number of protein-coding genes
and enriched GO terms experiencing relaxation or intensification of selection. Genes or enriched GO
terms are depicted in chalk yellow and light grey, respectively. (B) many enriched GO terms for genes
experiencing relaxation of selection were associated with neural function, neurogenesis, and cell growth.
20 out of a total of 57 significantly ($P < 0.05$, Fisher's exact test) enriched GO terms (biological process,
BP) are shown. (C) enriched GO terms (BP) for genes experiencing intensification of selection were
associated with energy metabolism, immune response, nervous system development, reproduction and
behavior. 20 out of a total of 71 significantly enriched GO terms are shown. (D) examples of genes that
experienced relaxation of selection in social branches caused by relaxed purifying selection, relaxed
positive selection, or both. (E) examples of genes that experienced intensification of selection in selection
caused by intensified purifying selection, intensified positive selection, or both.
731

**Figure 3. RERconverge identifies over 200 genes that tended to experience shifts in gene-wide
rates of molecular evolution during the transition to sociality.** (A) the number of genes and GO terms
experiencing acceleration or deceleration. Genes or enriched GO terms are depicted in green and light
grey, respectively. (B) 8 out of a total of 46 significantly (permulation *P* value < 0.05) enriched GO terms
(biological process, BP) in social branches, such as main categories of neural function and behavior. (C)
8 out of a total of 22 significantly decelerated GO terms (BP) in social branches, including main
categories of metabolism and DNA repair functions. (D) an example of an accelerated gene in social
branches, the nervous system development related gene, Integrin alpha-PS2 (ITA2). Social spider taxa
are depicted in orange, nonsocial taxa are in sky blue. (E) an example of a decelerated gene in social
branches, Kinesin-like protein KIF2A (KIF2A), a microtubule-associated motor protein with effects on
neural development and function.

746
Figure 4. Amino acid substitutions in an ion channel gene occur almost exclusively in social
branches and are predicted to cause protein structural changes. (A) Amino acid substitutions at
position 64 of the ion channel gene Two pore calcium channel protein 1 (TPC1) were found in all social
species and only a single nonsocial species, the prolonged subsocial *D.* cancerides. Triangles indicate a
substitution from valine to glycine and squares represent a substitution of valine to alanine. Social spider
species are depicted in orange, and nonsocial spider taxa in sky blue. X's on the phylogenetic tree
represents the independent origins of sociality. (B) Schematic diagram depicting the topology and domain
arrangement in TPC1. Periplasmic or cytoplasmic regions (dark line between boxes) and the 12
transmembrane (TM) domains (boxes) of TPC1 are shown. The substitutions V64G and V64A are located
in the TM4 domain. (C) Ribbon representation of the protein structure of spider TPC1 viewed from within
the membrane. The amino acid residue V64 is located at the center of TPC1, close to the periplasm side.
(D) Periplasm (top) view of the homology structure of TPC1, showing the location of the substitutions at
position 64. The 12 transmembrane alpha helices are shown as ribbons. (E) Residue V64 (blue color)
and substitutions V64G (orange) and V64A (green) are shown as stick representations.