Games of risk and reward in carnivore communities

Joel Ruprecht[a,1], Charlotte E. Eriksson[a,2], Tavis D. Forrester[b,2], Derek B. Spitz[c], Darren A. Clark[b], Mike J. Wisdom[d], Marcus Bianco[b], Mary M. Rowland[d], Joshua B. Smith[b], Bruce K. Johnson[b], and Taal Levi[a].

[a]Department of Fisheries and Wildlife, Oregon State University, [b]Oregon Department of Fish and Wildlife, [c]Environmental Studies Department, University of California, Santa Cruz, [d]USDA Forest Service Pacific Northwest Research Station

[1]To whom correspondence may be addressed.

[2]These authors contributed equally to this work.

Email: ruprechtjoel@gmail.com

Author Contributions: All authors designed the research and edited the manuscript. J.R. led the writing. J.R., T.D.F., M.B., and J.B.S. collected data. C.E.E. conducted DNA metabarcoding for diet. J.R., T.L. and D.B.S. analyzed the data.

Competing Interest Statement: The authors declare no competing interest.

Classification: Biological Sciences, Ecology.

Keywords: Carnivore, facilitation, mesopredator release, scavenging, suppression.

This PDF file includes:

> Main Text
> Figures 1 to 6
> Tables 1 to 1

Abstract

Mesopredator release theory suggests that dominant predators suppress subordinate carnivores
and ultimately shape community dynamics, but the assumption that subordinate species are only
negatively affected ignores the possibility of facilitation through scavenging. We examined the
interplay within a carnivore community consisting of cougars, coyotes, black bears, and bobcats
using contemporaneous Global Positioning System telemetry data from 51 individuals, diet
analysis from 972 DNA-metabarcoded scats, and data from 128 physical investigations of cougar
kill sites, 28 of which were monitored with remote cameras. Resource provisioning from
competitively-dominant cougars to coyotes through scavenging was so prolific as to be an
overwhelming determinant of coyote behavior, space use, and resource acquisition. This was
evident via strong attraction of coyotes to cougar kill sites, frequent scavenging of cougar-killed
prey, and coyote diets that nearly matched cougars in the magnitude of ungulate consumption.
Yet coyotes were often killed by cougars and used space to minimize encounters, complicating
the fitness benefits gained from scavenging. We estimated that 23% (95% CI: 8–55%) of the
coyote population in our study area was killed by cougars annually suggesting that coyote
interactions with cougars are a complex behavioral game of risk and reward. In contrast, we
found no indication that bobcat space use or diet was influenced by cougars. Black bears avoided
cougars, but there was no evidence of attraction to cougar kill sites, and much lower levels of
ungulate consumption and carcass visitation than for coyotes. Interspecific interactions among
carnivores are multifaceted encompassing both suppression and facilitation.

Significance Statement

An incomplete understanding of the total influence top predators exert on subordinate species
hinders our ability to anticipate the effects that changing carnivore populations will have in
ecological communities. Here we show that cougars are the architects of a complex behavioral
game of risk and reward, as subordinate or co-occurring carnivores are both provisioned and
preyed on by the top predators. Each co-occurring carnivore species considered here employed a
different strategy to approach the risk-reward tradeoff suggesting there are multiple viable
solutions to the game. By not considering the multitude of effects top predators have on other
carnivores, we are missing important linkages in terrestrial food webs.

30
31
32 **Main Text**
33
34 **Introduction**
35
The mesopredator release hypothesis posits a trophic cascade wherein larger-bodied
and competitively-dominant predators suppress subordinate predators with consequences for
prey populations (1-4). Such suppression is often the result of interspecific killing which may be
so pervasive as to account for one- to two-thirds of the total mortality of terrestrial carnivores (5-
7). Subordinate species may respond to this risk by altering their space use or activity patterns to
minimize the probability of encounters with dominant predators (8, 9), but nonetheless, the
antagonism between predator species can result in reductions in density or complete exclusion of
subordinate carnivores from certain habitats or regions (4, 10, 11).
The assumption that only negative interactions influence carnivore community structure is
likely overly simplistic because scavenging of dominant predator kills can also provision
subordinate species (7, 12, 13). While the prey killed by dominant predators may represent a
'fatal attraction' that leads mesopredators to their death, the energetic rewards from scavenging
in some cases may outweigh the risks (14). For example, if the risk of scavenging is the same as
hunting alternative prey but yields a higher probability of finding food, the fitness-maximizing
decision for the subordinate or co-occurring species should be to scavenge (Fig. 1; *SI Appendix*,

Text S1). But as risks associated with scavenging increase, the optimal choice becomes a more
nuanced function of the relative risk versus reward and depends on the energetic state of the
potential scavenger (Fig. 1a; *SI Appendix*, Text S1). And, of course, the scavenger must make
this assessment under imperfect knowledge of the risk involved. Thus, the decision for a
subordinate species to scavenge from a dominant species represents a complex behavioral
game of risk and reward (Fig. 1b). How the costs and benefits of interactions with dominant
predators influence the overall fitness of subordinate or co-occurring species has rarely been
examined.
Understanding the intraguild dynamics of carnivores is critical to anticipating the
ecosystem-level consequences of changing predator populations, but establishing the relative
prevalence of suppression and facilitation among carnivores has remained intractable. Strong
inference about the magnitude and direction of behavioral and population-level interactions
among carnivores requires (1) quantifying the intraguild predation rate on subordinate or co-
occurring carnivores, (2) understanding the extent to which subordinate or co-occurring
carnivores utilize carrion from dominant predator kills, and (3) evaluating whether dominant
predators or their kills influence the movements or space use of subordinate or co-occurring
carnivores. Spatial avoidance of dominant predators and high rates of intraguild predation would
suggest other carnivores are suppressed. By contrast, high utilization of dominant predator kills
by subordinates would support the facilitation hypothesis. We note that while the literature often
refers to the benefit conferred by scavenging as facilitation (7, 12, 14), this term implies no harm
to the dominant species. If carrion is lost due to scavenging (15, 16), the phenomenon may be
more appropriately termed parasitism, or more specifically, kleptoparasitism.
Here we quantify the influence of cougars (*Puma concolor*) on coyotes (*Canis latrans*),
black bears (*Ursus americanus*), and bobcats (*Lynx rufus*). Cougars are widely considered
dominant to coyotes and bobcats, and although previous researchers have suggested cougars
are subordinate to black bears, the available literature suggests that cougars kill bears more often
than the reverse (17). We therefore consider bears and cougars to be co-occurring and not
necessarily fitting within a clear dominance hierarchy. We leveraged four empirical datasets
including contemporaneous Global Positioning System (GPS) telemetry data of 51 individuals
across the four species of carnivores, diet analysis from 972 DNA-metabarcoded carnivore scats,
and data from 128 physical investigations of cougar kill sites, 28 of which were monitored for
scavenging via remote cameras to evaluate how sympatric carnivores are influenced by cougars.
We hypothesized that black bears, bobcats and coyotes could be unaffected by cougars (null
hypothesis), negatively influenced by cougars (suppression hypothesis), positively influenced by
cougars (facilitation hypothesis), or both positively and negatively affected by cougars (provision-
predation hypothesis) (Fig. 1c). We used contemporaneous GPS data to infer whether each
carnivore species altered movements due to the proximity of a cougar or cougar kill site. We
hypothesized that carnivore species attracted to cougar kill sites would have a high frequency of
carrion in their diet—suggesting facilitation from cougars—but would also be killed by cougars at
higher rates than species that avoided cougar kill sites, indicating suppression. Together, these
approaches provided a robust evaluation of the fine-scale behavioral interactions and facilitation
versus suppression tradeoff among species within carnivore communities, which will be
increasingly important as predator populations recover across portions of their former range and
are extirpated in others.
95
96 **Materials and Methods**
97
Data collection. We collected data on four sympatric carnivore species (cougars, coyotes, black
bears, and bobcats) in and adjacent to the Starkey Experimental Forest and Range in the Blue
Mountains of northeastern Oregon, USA (Fig. 2 and *SI Appendix*, Text S2) between 2016 and
2020. We tagged 17 cougars (8M, 9F), 17 coyotes (10M, 7F), 11 black bears (7M, 4F), and 6
bobcats (3M, 3F) with GPS collars (Vectronic Aerospace or Lotek Wireless, Inc.) programmed to
record positions every 2 or 3 hours (*SI Appendix*, Text S3). We used conservation detection dogs
(18) to locate scats (*SI Appendix*, Text S4) and genetically confirmed 774 coyote scats, 85 black
bear scats, 96 bobcat scats, and 17 cougar scats (19) that were DNA-metabarcoded for diet

analysis (*SI Appendix*, Text S5). We considered the frequency of occurrence of ungulates (i.e.,
deer spp. [*Odocoileus hemionus* and *O. virginianus*] and elk [*Cervus canadensis*]) in scats of
subordinate carnivores as a qualitative but imperfect proxy for scavenging (14). Although each of
these carnivore species can kill deer of all ages, elk consumption by coyotes and bobcats in
particular is likely exclusively from scavenging because these species rarely kill elk calves in the
region (20). In addition to inference from scat analysis, we further quantified cougar diets by
conducting ground searches of clusters of cougar GPS locations indicating potential kill sites of
tagged individuals (*SI Appendix*, Text S6) (21). At 28 of the 128 confirmed cougar kill sites, we
deployed remote cameras (Bushnell Trophy Cam Aggressor, Browning Dark Ops HD Pro, and
Reconyx HyperFire 2) to estimate scavenging rates by the other carnivores. We separately tallied
the number of daily visits made by coyotes, bears and bobcats to cougar kill sites that were
detected on camera, where a visit was defined as a detection separated by at least 30 minutes
from another detection of a given species. We used negative binomial regression to model the
number of daily visits to the carcass by a given species as:
120
   number of daily visits = cougar present + ln(time since kill) + prey size + kill ID
121
122 where cougar present (1/0) indicates whether the cougar was present at the kill that day, time
123 since kill is the number of days since the kill was made, prey size (1/0) indicates whether the prey
124 was large (adult ungulate) or not (young of the year ungulate), and kill ID is a random intercept
125 identifying a given kill site.
126
Calculating Intraguild Predation Rate. At times, the prey items found at cougar kill sites were
other carnivores which allowed us to calculate the proportion of cougar kills representing
intraguild predation. Combining these data with information on contemporaneously-estimated
carnivore densities (22) and cougar kill rates (21) from a recent study in the area allowed us to
calculate intraguild predation rates, i.e., the proportion of the co-occurring carnivore population
dying annually due to cougar predation. We calculated the intraguild predation rate on species i
(P_i) as the number of individuals from species i killed per unit time (K_i) divided by the population
density of species i (D_i) where K_i is the product of cougar population density (D_c) cougar kill rate
(kills/year), R, and proportion of kills corresponding to species i (F_i):
136

$$P_i = \frac{K_i}{D_i} = \frac{D_c \times R \times F_i}{D_i}$$

137
138 We quantified uncertainty in P_i by propagating error inherent in each parameter using Markov
139 Chain Monte Carlo simulations (*SI Appendix*, Text S7).
140
Step selection functions. We fit integrated step-selection functions (hereafter *SSFs*) (23, 24) to
GPS collar data to estimate how coyote, bobcat, and bear movements were influenced by
landscape variables, cougar proximity, or cougar kill site proximity. We included movement
attributes as parameters in each model to reduce bias arising from sampling control locations (25)
using a gamma distribution to characterize the natural log of step lengths and a von Mises
distribution to characterize the cosine of turning angles (24, 26). We generated 20 random steps
for each observed step (27). To ensure that territoriality did not bias inference, we did not allow
the endpoints of random steps to fall outside a 1,000 m buffer of the 99% minimum convex
polygon of each animal's territory. We did this to prevent detecting spurious avoidance behavior
due to a random step occurring in an area for which the animal did not have uninhibited access
due to territoriality.
We sought to identify whether landscape features and the proximity of a cougar
(hereafter, 'distance to cougar' or D2C) or cougar kill site (hereafter, 'distance to kill' or D2K)
influenced the movements of coyotes, bobcats, and bears. We calculated distance to cougar as
the natural log-transformed Euclidean distance between each focal individual (coyote, bobcat, or
bear) and the nearest GPS-collared cougar at that time to determine whether the subsequent

movements of the animal were toward or away from the cougar. To restrict encounters to only
those in which the cougar could plausibly be detected by the subordinate species, we only
allowed cougars within 1,000 m of the focal bear, bobcat, or coyote to influence the movements
of the other carnivores by using a binary indicator for whether a cougar was present (hereafter,
'cougar present' or CP) (1/0) at each GPS fix. We cast 'distance to cougar' as an interaction term
with 'cougar present' such that the term was nullified when the nearest known cougar was >
1,000 m away. We used a buffer of 1,000 m because that is the distance elk alter behavior in
response to wolves (28) and has also been used to determine interaction distances between
cheetahs (29). Further, it is implausible that species could detect cougars at distances greatly
exceeding this value. However, to ensure this cutoff did not change our inference about the
direction or magnitude of the effect of cougars, we replicated the analyses using additional cutoff
values of 500 and 1,500 m. If GPS fixes between cougars and the other carnivores were not
contemporaneous (i.e. falling on the same hour plus a tolerance of 300 seconds) due to
misalignment or missed fixes, we imputed the location of the cougar using a correlated random
walk (30). We allowed cougar locations to be imputed if a single GPS fix was missed but we
censored location data for any gaps exceeding 6 hours.
We used this same approach to assess how carnivores responded to cougar kill sites by
including a term for 'distance to kill,' calculated as the natural log of the Euclidean distance
between each GPS position of a focal individual and the nearest cougar kill site. Potential kill sites
were identified from clusters of sequential cougar GPS locations indicating reduced movement
and were either confirmed via physical investigations or identified as probable kills using a
predictive model (21) (*SI Appendix*, Text S6). We paired 'distance to kill' with a binary indicator
variable describing whether there was a known or predicted cougar kill present (hereafter, 'kill
present' or KP) (1/0) within a 3,000 m buffer of the focal individual. When there was not a kill
within the buffer, the indicator variable took the value of zero and the whole term became null. If
there was more than one kill within this buffer for a given individual at a given time, we considered
only the distance to the most recent kill. Our assumption that carnivores could detect a carcass
up to distances of 3,000 m was conservative; for example, one GPS-collared coyote in our study
exhibited two bouts of directed travel of 2.2 km and 3.6 km outside its territory to feed on an elk
carcass suggesting an ability to detect carrion from these distances (*SI Appendix*, Fig. S1).
Further, previous studies report that coyotes can travel between 12.2 and 20.5 km to reach
carcasses of domestic animals (31, 32), while arctic foxes routinely detect marine mammal
carcasses at distances of at least 10 km (33). While less is known about olfaction distances in
bears, it is reported that polar bears (*Ursus maritimus*) can detect seal breathing holes at
distances of 3 km (34). While the available literature suggests 3,000 m is an appropriate value for
a cutoff, we replicated each analysis using cutoffs of 2,000 m and 4,000 m to ensure inference
was not an artifact of the buffer chosen.
To avoid data contamination due to old carcasses that no longer provided scavenging
opportunities, we classified all kills occurring within 30 days of the focal fix as present by
censoring kills exceeding this threshold. To quantify how movements of carnivores toward or
away from the kill may change as the carcass aged, we created another variable cast as an
interaction between the kill present indicator variable (KP), the natural log of the distance to kill
(D2K), and the natural log of the time elapsed since the kill (hereafter, 'time since kill' or TSK).
This interaction allowed the magnitude of avoidance or attraction to change as a function of the
age of the carcass.
It is certain that at times additional cougars were present in the study area not monitored
with GPS collars, so the inferences made about effects of cougars and their kill sites on the other
carnivore species should be considered conservative. Thus, with complete information on all
cougar and kill site locations, the D2C and D2K variables may have had larger effect sizes and/or
smaller standard errors than reported here.
Before including cougar variables, we first fit models containing only landscape, and
movement variable (hereafter, *habitat model*) according to

$w(x) = exp(\boldsymbol{\beta} \times landscape\ variables\ + \beta_{turn\ angle} \times cosine(turn\ angle) + \beta_{step\ length} \times ln(step\ length))$

210
where 'landscape variables' refer to the continuous variables *ln*(distance to open road),
ln(distance to perennial water source), canopy cover, topographic ruggedness, and the factor
variable potential vegetation type. *A priori*, we expected each of these landscape variables to
influence resource selection of each of the species considered, so for simplicity we did not
conduct model selection on subsets of these variables and instead retained them all.
We then added the cougar and kill site terms and used Akaike's Information Criterion
(AIC) (35) to assess whether this full model (hereafter, *habitat + cougar model*) outperformed the
habitat-only model:
219

$$w(x) = \exp(\boldsymbol{\beta} \times \textit{landscape variables} + \beta_{D2C} \times CP \times D2C + \beta_{D2K} \times KP \times D2K + \beta_{D2K \times TSK} \times KP \times D2K \times TSK + \beta_{turn\ angle} \times cosi$$

220
221 By setting $\beta_{D2K} \times \mathrm{KP} \times \mathrm{D2K} + \beta_{D2K \times TSK} \times \mathrm{KP} \times \mathrm{D2K} \times \mathrm{TSK}$ equal to zero and solving for
222 TSK, we estimated the time until the effect of the kill becomes null. Assuming a kill site is initially
223 a source of attraction that diminishes in time, this is estimated by
224

$$\exp\left(-\frac{\beta_{D2K}}{\beta_{D2K \times TSK}}\right)$$

225
226 We used the 'amt' package (26) in Program R 3.6.1 (R Development Core Team 2019) to
227 format data, generate random steps, and fit models.
228
Evaluating Weight of Evidence for Facilitation, Suppression, and Null Hypotheses. We
leveraged multiple independent datasets on the four carnivore species using a weight of evidence
approach to determine whether cougars suppress and/or facilitate black bears, bobcats, and
coyotes. *A priori*, we established the following criteria: 1) spatial avoidance of cougars would be
evidence of suppression, attraction to cougars would be evidence of facilitation, and indifference
toward cougars would support the null hypothesis, 2) spatial avoidance of cougar kill sites would
be evidence of suppression, attraction to kill sites would be evidence of facilitation, and
indifference toward kill sites would support the null hypothesis, 3) a higher frequency of ungulates
found in scats of subordinate or co-occurring predators (a proxy for scavenging in our system) in
similar proportions to those observed in cougar scats would suggest a higher degree of facilitation
than would a lower frequency of ungulates found in scats, and no ungulates found in scats would
support the null hypothesis, 4) higher scavenging rates would suggest a greater degree of
facilitation than would lower scavenging rates, and the absence of scavenging would support the
null hypothesis, and 5) any amount of intraguild predation by cougars on other carnivores would
be evidence of suppression, whereas complete absence of intraguild predation would support the
null hypothesis.
245
246 **Results**
247
Step selection functions. Coyotes, black bears, and bobcats each responded to a subset of
landscape variables; most notably, bears and bobcats strongly selected for high canopy cover
while coyotes avoided high canopy cover (Table 1 and SI Appendix; Fig. S2). Including the three
cougar parameters (CP × D2C, KP × D2K, and KP × D2K × TSK) substantially improved model fit
for bears (ΔAIC = 10.36) and particularly for coyotes (ΔAIC = 23.96), but the habitat-only model
outperformed the habitat + cougar model for bobcats (ΔAIC = 2.69) (SI Appendix, Table S1).
Notably, coyotes exhibited a strong attraction to cougar kill sites (β = -0.26, p < 0.001) which
attenuated as the carcass aged (β = 0.071, p < 0.001; Fig. 3 and Table 1) and resulted in
approximately 39 days of selection for cougar kill sites. Post-hoc analyses revealed that attraction
of coyotes to cougar kill sites was driven primarily by resident male coyotes (SI Appendix, Text
S9–S10, Table S2–S3). Coyotes avoided cougars (β = 0.20, p = 0.014; Table 1), but there was
no evidence to suggest attraction to kill sites was diminished if a cougar was present at the kill,
indicating coyotes may disregard the risk of cougars when a food reward is present (SI Appendix,

Text S11, Table S4). Bears were indifferent to kill sites (β = 0.01, p = 0.93) but actively avoided
cougars (β = 0.55, p $\leq$ 0.001; Table 1). Bobcats were indifferent to both cougars (β = 0.26, p =
0.33) and their kill sites (β = 0.11, p = 0.54; Table 1). For all species, the choice of buffer for
determining whether a cougar (CP) or kill (KP) was present did not appreciably change inference
about the direction or magnitude of these effects except that coyote avoidance of cougars was
only evident when buffers 1,000 m or less were used (SI Appendix, Text S12, Table S5–S10).
267
Diet analysis. DNA metabarcoding of carnivore scats revealed a variable frequency of
occurrence of deer and elk across species. In coyotes, elk and deer were present in 58% and
12% of scats, respectively (Fig. 4b). In bears, elk was found in 29% of scats and deer in 8% of
scats (Fig. 4b). Zero bobcat scats contained elk and 8% contained deer (Fig. 4b). For cougars,
61% of scats contained elk and 22% contained deer. Investigations of 128 cougar kill sites with
confirmed prey revealed that elk represented 64% and deer 16% of the prey items killed by
cougars (Figs. 4a, 4b).
275
Scavenging Rates. We monitored 28 cougar kill sites containing ungulate prey using remote
cameras to estimate scavenging rates. Cameras were placed an average of 4.4 (range: 0–16)
days after the kill was made and were operational an average of 17.4 (range: 1–52) days. We
documented coyotes present at 89% of carcasses, bears at 50% of the carcasses outside the
hibernation period (N = 22), and zero bobcats were detected at cougar kills (Fig. 4c, 4d) during
the period cameras were active. For coyotes, the negative binomial regression model predicted
that one day after a large ungulate was killed, the carcass would receive 1.8 visits by coyotes but
that the number of visits would be reduced by 27% with every additional ln(day) (β = –0.31, p =
0.067; Fig. 5a). The coyote visitation rate was expected to be 160% higher for a large ungulate
than a juvenile ungulate (β = 0.97, p = 0.059) and was not influenced by cougar presence (β =
0.43, p = 0.26). The bear model predicted 0.36 visits by bears the day after a large ungulate was
killed and that every additional ln(day) would reduce the number of visits by 52% (β = –0.73, p =
0.051; Fig. 5b). The number of daily bear visits was not influenced by the size of prey (β = 0.39, p
= 0.50) nor whether a cougar was present (β = –0.12, p = 0.89). The intercepts were –0.81 (p =
0.22) and –1.32 (p = 0.29) for the coyote and bear models, respectively.
291
Intraguild Predation. Physical investigations of cougar kill sites indicated that coyotes
represented 7.0% (9 of 128; 95% CI: 3.3–11.1%) of cougar kills (Fig. 4e, f). In eight cases
coyotes were the only prey item found and were consumed by cougars, and in one case a dead
coyote was found in conjunction with another prey item. In the latter scenario, the coyote was not
consumed but exhibited trauma consistent with cougar predation and was presumably killed as it
attempted to scavenge from the kill. One transient GPS-collared coyote in our sample was killed
and consumed by an uncollared cougar outside the study area (Fig. 4e) and in another instance,
a GPS-collared cougar killed and consumed a GPS-collared coyote in the study area. With all
sources of uncertainty propagated, we estimated that annually, cougars kill approximately 23.0%
(95% CI: 8.4–54.7%) of the coyote population according to:
302

$$Intraguild\ Predation\ rate = \frac{(2.2\ cougars \cdot 100\ km^{-2}) \times (54.5\ kills \cdot cougar^{-1} \cdot year^{-1}) \times (0.07\ dead\ coyotes \cdot kill^{-1})}{33.9\ coyotes \cdot 100\ km^{-2}}$$

303
Weight of Evidence for Facilitation, Suppression, and Null Hypotheses. We found evidence
of both facilitation and suppression of other carnivore species by cougars (Fig. 6). Most notably,
coyotes exhibited facilitation in three of the five measures evaluated (statistically significant
attraction to kill sites, frequent carrion in diet, and high scavenging rates) and indication of
suppression in two measures (the presence of intraguild predation and avoidance of cougars).
Bears showed evidence of facilitation in two measures (moderate scavenging rates and moderate
levels of carrion in diet), suppression in one measure (statistically significant avoidance of
cougars), and no evidence in two measures (absence of statistically significant attraction or
avoidance of kill sites, absence of intraguild predation). We found no evidence that cougars

influenced bobcats in any of the five measures we evaluated, fully supporting the null hypothesis
that cougars neither suppress nor facilitate bobcats.
315
316 **Discussion**
317
We combined contemporaneous GPS-tracking of four carnivore species, prey
identification at cougar kill sites, diet analysis with DNA metabarcoding, and remote-camera
monitoring of cougar kills to assess the relative strength of suppressive and facilitative forces
between a competitively dominant predator and three subordinate or co-occurring predators. Our
strongest evidence of facilitation suggested that resource provisioning from cougars to coyotes
through scavenging was so ubiquitous as to be an overwhelming determinant of coyote behavior,
space use, and resource acquisition. This finding was demonstrated through strong attraction of
coyotes to cougar kill sites, coyotes scavenging nearly all carcasses at cougar kill sites that we
monitored, and coyote diets that nearly matched cougars in the magnitude and composition of
ungulate consumption. However, we also show that cougars readily killed coyotes as prey and to
a much lesser degree from competitive interactions at kill sites. These previously unquantified
interactions highlight a complex game of risk and reward as coyotes seek to maximize their
fitness by scavenging while simultaneously evading cougars (7, 36).
Selective forces operating on coyote behavior influence their acceptance of risk while
acquiring resources at cougar kills. By observing dead coyotes at 7% of cougar kill sites, we
found the risk cougars pose to coyotes to be substantial. We estimated that each cougar killed
approximately 3.8 coyotes annually, though we note previous studies in the region have reported
less coyote predation (21, 37) so this value may partially be the result of a small sample size.
Nonetheless, given a cougar density of 2.2 per 100 km^2 in our study area (22), this suggests that
approximately 8.4 coyotes are killed per 100 km^2 per year. With a coyote density of 33.9 per 100
km^2, this level of mortality reflects 23.0% (95% CI: 8.4–54.5% when all sources of uncertainty are
propagated) of coyotes killed by cougars annually. Thus, cougar predation is a very significant
mortality source to coyotes, but coyotes nevertheless readily accept this risk to forage extensively
on cougar kills. This risk may not be evenly shared by all coyotes, however, because many (6 of
9) of the coyotes killed by cougars were young (< 2 years old), indicating that breeding coyotes
may gain the most benefit from scavenging while younger and dispersing coyotes may bear most
of the intraguild predation cost. Therefore, cougar predation on coyotes is probably compensatory
to some degree although we cannot confirm this with the available data.
Only 1 of 9 coyotes killed by cougars occurred at a kill site. While this is mildly suggestive
of increased risk while scavenging because coyotes spend substantially less than 1/9 of their
daily activity budget (2.67 hours daily) scavenging, this small sample size precludes statistical
determination of how much, or if, predation risk per unit time is higher while actively scavenging.
If risk from cougars is not higher at kill sites than elsewhere on the landscape, then scavenging
becomes an even more appealing prospect (Fig. 1a). While heightened risk at cougar kill sites is
intuitive, this may not be the case for several reasons. First, coyotes may exhibit heightened
vigilance levels when scavenging if dominant predators are perceived to be near (38). Second,
assuming cougars are satiated after feeding on a kill, they may have little motivation to kill
coyotes if their primary reason for doing so is for consumptive purposes. Determining whether
cougars kill coyotes for an immediate energetic gain or the long-term benefit of removing a
competitor should be the focus of further study. Two lines of evidence suggest coyotes do not
reduce their use of kill sites if cougars are present at the kill; terms accounting for this were not
significant in either the regression model of kill site visits from camera data nor a post-hoc step-
selection function with an added term indicating whether a cougar was on a kill in question (*SI
Appendix*, Text S11).
Not all species showed such a strong influence from cougars. Black bears were
indifferent to cougar kill sites as determined from step-selection functions using GPS data which
was surprising given substantial bear scavenging of cougar prey has been previously
documented in other studies (15, 16, 20, 39). Not only did step selection functions indicate no
attraction toward cougar kills, scat analysis revealed deer and elk constituted only a moderate
percentage of the diets of black bears. Because bears readily kill and consume deer and elk

neonates (40), a substantial proportion of ungulates in their diet may have come from direct
predation. A lower rate of bear scavenging at cougar kills is further supported by camera data
indicating that bears were less frequent visitors to cougar kills than were coyotes. The lack of
attraction to kill sites as measured by GPS data suggests that the bears that did visit cougar-killed
prey may have encountered them opportunistically, in contrast to coyotes which at times made
long and directed movements toward kill sites. Further, bears may not have frequently overlapped
with cougar kills for several reasons. First, bears exhibit seasonal movements following
ephemeral resource pulses and in this region, these seasonal resources are often located in
habitats not frequently used by cougars. Second, the time when bears are active (i.e. not
denning) generally coincides with time periods in which cougars kill smaller prey (21) further
limiting the potential for scavenging, particularly if coyotes arrive first. The selection coefficient for
distance to kill still was not statistically significant when male and female bears were modeled
separately; however, the direction of the coefficient suggested male bears tended to be slightly
attracted to kill sites while female bears tended to avoid them (*SI Appendix*, Text S13, Table
S11). This accords with previous studies of brown bears wherein females with cubs had a lower
propensity to scavenge than males (41).
While bears neither selected for nor avoided cougar kills, step selection functions
indicated that black bears did actively avoid cougars. This was unexpected because black bears
often usurp cougar prey at kill sites in other areas (15, 16). However, two recent studies on
cougar prey selection in an adjacent study area reported finding bears as prey items (21, 37) and
other literature reports more instances of cougars killing black bears than the converse (17). Our
results suggest that black bears in our system view cougars as sufficiently threatening that they
utilize the landscape to reduce the risk of confrontation. When modeled separately, both male
and female bears exhibited avoidance of cougars (*SI Appendix*, Text S13, Table S11). Because
body size is a major determinant of intraguild conflict (6), it is possible that larger bears (either
larger individuals or larger bear species) would be less risk averse toward cougars, but we did not
have the data to assess this.
Previous research indicates that intraguild predation is most likely to operate between
carnivore species that have intermediate body size differences, broad dietary overlap, and that
are taxonomically related (5, 6). Despite their taxonomic relatedness to cougars, we found no
evidence suggesting that bobcats were attracted to cougar kill sites or otherwise influenced by
cougars (42). Previous studies have found variable levels of bobcat scavenging (15, 39), but our
three data sources indicated very clearly that bobcats in this system are not scavenging from
cougars. We found zero instances of elk in any of 96 bobcat scats and zero observations of
bobcats at cougar kill sites. While 8% of bobcat scats contained deer, this could be explained by
direct predation because bobcats kill deer in this system (43). By foregoing energetically
profitable cougar kills, bobcats in this system are solving the game of risk and reward quite
differently than are coyotes. This is expected when the availability of non-scavenged foods is
higher, or body condition is high, thus obviating the need to incur risk, or if predation risk at kills is
higher, which might be the case for solitary bobcats relative to group-living coyotes (Fig. 1a).
While we have assessed behavioral interactions among a dominance hierarchy of extant
carnivores, gray wolves (*Canis lupus*) are now recolonizing much of western North America
including our study area. At the time of data collection, wolves were only a minor presence on the
landscape and very rarely killed or scavenged prey in our study area (ODFW, unpublished data).
But as wolves become more prominent on the landscape, the interplay between carnivores in this
system is likely to change. For example, wolves are likely to deposit carrion in more open
grassland environments than do cougars. The larger viewsheds in such vegetation types may be
attractive to coyotes seeking scavenging opportunities (44), but the increased post-encounter
predation risk by a wolf pack versus a solitary cougar may shift coyote behavior toward increased
avoidance of wolf kills. Future work might assess whether coyotes perceive the risk-reward
landscape with wolves differently than that with cougars by monitoring selection for, and
scavenging at, both cougar and wolf kills.
The extent to which scavenging by co-occurring carnivores affects cougars is not clear.
Despite cougars being an average of 4–5 times larger than coyotes in this area, we observed at
least one instance via remote camera monitoring of kill sites where coyotes displaced a cougar

from its kill. The cougar involved had dependent young that were likely nearby, which may have
made it especially vulnerable or responsive to the mesopredators. A question with broader
ecological significance, however, is whether cougars increase their kill rate due to lost feeding
opportunities caused by scavengers. While others have shown that black bears often usurp
cougar kills which may lead to an increase in kill rate (15, 16), there are many confounding
factors such as spatiotemporal variation in prey size, composition, or density, seasonality of kills,
carrion decomposition rates, etc. to consider this a generality for all scavengers. However, our
camera monitoring of carcasses showed that coyotes often arrived and began scavenging before
cougars abandoned their kills, so any loss of carrion should technically be considered
kleptoparasitism and not facilitation *per se*.
433   To conclude, it is probably a gross oversimplification to propose that species interactions
within terrestrial carnivore communities have either strictly positive, negative, or even neutral
consequences (7). We demonstrated that coyotes, and a lesser extent bears, were confronted by
the opposing effects of both facilitation (i.e. food provisioning) and suppression (i.e. death), yet
bobcats were seemingly unaffected by a dominant predator. Scavenging carrion from dominant
predators incurs both risk and reward, and thus the willingness to engage in this activity must be
mediated by behavior. But why would coyotes, bears, and bobcats approach the risk-reward
tradeoff so differently? One possible explanation is that the selective pressures favoring one risk
aversion strategy over another operated differently by species. As the efficiency of acquiring
alternative prey increases, scavenging becomes less advantageous (Fig. 1a) and this should
cause species with higher proficiency finding alternative prey to scavenge less. The ability to
perceive risk likely differs by species as well, and regardless of whether these are innate or
learned behaviors, our results suggest that the propensity to meet energetic demands by
scavenging is variable across species, with neither strategy appearing to have an obvious fitness
benefit over the other. This suggests that there are likely to be multiple viable strategies to solve
this complex behavioral game.
449
450   **Acknowledgments**
451

452   We are grateful to the Oregon Department of Fish and Wildlife, the USDA Forest Service Pacific
453   Northwest Research Station, Oregon State University, and the Wildlife Restoration Act for funding
454   and/or logistical support. We thank a multitude of technicians and volunteers who assisted with
455   fieldwork and photo tagging.

456
457
References

1.   K. R. Crooks, M. E. Soulé, Mesopredator release and avifaunal extinctions in a
fragmented system. *Nature* **400**, 563-566 (1999).
2.   S. E. Henke, F. C. Bryant, Effects of coyote removal on the faunal community in western
Texas. *The Journal of Wildlife Management*, 1066-1081 (1999).
3.   L. R. Prugh *et al.*, The rise of the mesopredator. *Bioscience* **59**, 779-791 (2009).
4.   T. Levi, C. C. Wilmers, Wolves–coyotes–foxes: a cascade among carnivores. *Ecology*
93, 921-929 (2012).
5.   F. Palomares, T. M. Caro, Interspecific killing among mammalian carnivores. *The*
American Naturalist **153**, 492-508 (1999).
6.   E. Donadio, S. W. Buskirk, Diet, morphology, and interspecific killing in *Carnivora*. *The*
American Naturalist **167**, 524-536 (2006).
7.   L. R. Prugh, K. J. Sivy, Enemies with benefits: integrating positive and negative
interactions among terrestrial carnivores. *Ecology Letters* **23**, 902-918 (2020).
8.   S. M. Durant, Living with the enemy: avoidance of hyenas and lions by cheetahs in the
Serengeti. *Behavioral Ecology* **11**, 624-632 (2000).
9.   E. Dröge, S. Creel, M. S. Becker, J. M'soka, Spatial and temporal avoidance of risk within
a large carnivore guild. *Ecology and Evolution* **7**, 189-199 (2017).

10. D. J. Harrison, J. A. Bissonette, J. A. Sherburne, Spatial relationships between coyotes
and red foxes in eastern Maine. *The Journal of Wildlife Management*, 181-185 (1989).
11. J. D. Linnell, O. Strand, Interference interactions, co-existence and conservation of
mammalian carnivores. *Diversity and Distributions* **6**, 169-176 (2000).
12. K. J. Sivy, C. B. Pozzanghera, K. E. Colson, M. A. Mumma, L. R. Prugh, Apex predators
and the facilitation of resource partitioning among mesopredators. *Oikos* **127**, 607-621
(2018).
13. C. C. Wilmers, R. L. Crabtree, D. W. Smith, K. M. Murphy, W. M. Getz, Trophic facilitation
by introduced top predators: grey wolf subsidies to scavengers in Yellowstone National
Park. *Journal of Animal Ecology* **72**, 909-916 (2003).
14. K. J. Sivy, C. B. Pozzanghera, J. B. Grace, L. R. Prugh, Fatal attraction? Intraguild
facilitation and suppression among predators. *The American Naturalist* **190**, 663-679
(2017).
15. M. L. Allen, L. M. Elbroch, C. C. Wilmers, H. U. Wittmer, The comparative effects of large
carnivores on the acquisition of carrion by scavengers. *The American Naturalist* **185**,
822-833 (2015).
16. L. M. Elbroch, P. E. Lendrum, M. L. Allen, H. U. Wittmer, Nowhere to hide: pumas, black
bears, and competition refuges. *Behavioral Ecology* **26**, 247-254 (2015).
17. L. M. Elbroch, A. Kusler, Are pumas subordinate carnivores, and does it matter? *PeerJ* **6**,
e4293 (2018).
18. S. K. Wasser *et al.*, Scat detection dogs in wildlife research and management: application
to grizzly and black bears in the Yellowhead Ecosystem, Alberta, Canada. *Canadian
Journal of Zoology* **82**, 475-492 (2004).
19. C. E. Eriksson, J. Ruprecht, T. Levi, More affordable and effective noninvasive single
nucleotide polymorphism genotyping using high-throughput amplicon sequencing.
Molecular Ecology Resources **20**, 1505-1516 (2020).
20. B. K. Johnson *et al.*, Roles of maternal condition and predation in survival of juvenile elk
in Oregon. *Wildlife Monographs* **201**, 3-60 (2019).
21. D. A. Clark, G. A. Davidson, B. K. Johnson, R. G. Anthony, Cougar kill rates and prey
selection in a multiple-prey system in northeast Oregon. *The Journal of Wildlife
Management* **78**, 1161-1176 (2014).
22. J. Ruprecht *et al.*, Evaluating and integrating spatial capture-recapture models with data
of variable individual identifiability. *BioRxiv* [Preprint] (2020).
https://www.biorxiv.org/content/10.1101/2020.03.27.010850v2 (accessed 26 January
2021).
23. D. Fortin *et al.*, Wolves influence elk movements: behavior shapes a trophic cascade in
Yellowstone National Park. *Ecology* **86**, 1320-1330 (2005).
24. T. Avgar, J. R. Potts, M. A. Lewis, M. S. Boyce, Integrated step selection analysis:
bridging the gap between resource selection and animal movement. *Methods in Ecology
and Evolution* **7**, 619-630 (2016).
25. J. D. Forester, H. K. Im, P. J. Rathouz, Accounting for animal movement in estimation of
resource selection functions: sampling and data analysis. *Ecology* **90**, 3554-3565 (2009).
26. J. Signer, J. Fieberg, T. Avgar, Animal movement tools (amt): R package for managing
tracking data and conducting habitat selection analyses. *Ecology and Evolution* **9**, 880-
890 (2019).
27. G. Latombe, D. Fortin, L. Parrott, Spatio-temporal dynamics in the response of woodland
caribou and moose to the passage of grey wolf. *Journal of Animal Ecology* **83**, 185-198
(2014).
28. A. D. Middleton *et al.*, Linking anti-predator behaviour to prey demography reveals limited
risk effects of an actively hunting large carnivore. *Ecology Letters* **16**, 1023-1030 (2013).
29. F. Broekhuis, E. K. Madsen, K. Keiwua, D. W. Macdonald, Using GPS collars to
investigate the frequency and behavioural outcomes of intraspecific interactions among
carnivores: A case study of male cheetahs in the Maasai Mara, Kenya. *PloS One* **14**,
e0213910 (2019).

30. B. T. McClintock, T. Michelot, momentuHMM: R package for generalized hidden Markov
models of animal movement. *Methods in Ecology and Evolution* **9**, 1518-1530 (2018).
31. D. A. Danner, N. S. Smith, Coyote home range, movement, and relative abundance near
a cattle feedyard. *The Journal of Wildlife Management* **44**, 484-487 (1980).
32. J. F. Kamler, W. B. Ballard, R. L. Gilliland, K. Mote, Coyote (Canis latrans) movements
relative to cattle (*Bos taurus*) carcass areas. *Western North American Naturalist*, 53-58
(2004).
33. S. Lai, J. Bêty, D. Berteaux, Spatio–temporal hotspots of satellite–tracked arctic foxes
reveal a large detection range in a mammalian predator. *Movement Ecology* **3**, 37 (2015).
34. R. R. Togunov, A. E. Derocher, N. J. Lunn, Windscapes and olfactory foraging in a large
carnivore. *Scientific Reports* **7**, 46332 (2017).
35. K. P. Burnham, D. R. Anderson, *Model Selection and Multimodel Inference: A practical
information-theoretic approach* (Springer, 2002).
36. P. J. Mahoney, "Spatial ecology of coyotes and cougars: understanding the influence of
multiple prey on the spatial interactions of two predators," PhD dissertation, Utah State
University, Logan Utah, USA (2017).
37. E. K. Orning, "Carnivore Competition: Spatial and Dietary Implications of Gray Wolf
Recolonization for Cougars in Northeast Oregon," PhD dissertation, Oregon State
University, Corvallis, Oregon, USA (2019).
38. J. Young, P. J. Mahoney (2017) "Foraging behavior of coyotes under intraguild predation
risk by cougars: an experimental approach" in *Proceedings of the 12th Mountain Lion
Workshop*, C. R. McLaughlin, M. Viera, Eds. (Western Association of Fish and Wildlife
Agencies, Estes Park, Colorado, USA, 2017), p. 232.
39. M. L. Allen, L. M. Elbroch, C. C. Wilmers, H. U. Wittmer, Trophic facilitation or limitation?
Comparative effects of pumas and black bears on the scavenger community. *PLoS One*
9 (2014).
40. P. Zager, J. Beecham, The role of American black bears and brown bears as predators
on ungulates in North America. *Ursus* **17**, 95-108 (2006).
41. A. Ordiz *et al.*, Individual Variation in Predatory Behavior, Scavenging and Seasonal Prey
Availability as Potential Drivers of Coexistence between Wolves and Bears. *Diversity* **12**,
(2020).
42. C. Hass, Competition and coexistence in sympatric bobcats and pumas. *Journal of
Zoology* **278**, 174-180 (2009).
43. N. J. Jackson, K. M. Stewart, M. J. Wisdom, D. A. Clark, M. M. Rowland, Demographic
performance of a large herbivore: effects of winter nutrition and weather. *Ecosphere* **12**,
e03328 (2021).
44. E. M. Gese, R. L. Ruff, R. L. Crabtree, Foraging ecology of coyotes (*Canis latrans*): the
influence of extrinsic factors and a dominance hierarchy. *Canadian Journal of Zoology*
74, 769-783 (1996).

569
570

Figures and Tables

573
Figure 1. a) Predictions from a two patch dynamic state variable model used as an illustrative
example. The conceptual model predicts the fitness-maximizing decision where the choices are
to scavenge (patch 1, tan shading) or forego the scavenging opportunity and hunt alternative prey
(patch 2, blue shading). The optimal decision in this theoretical model depends on the risk,
quantified as daily probability of death, and reward, quantified as probability of finding food. The
fitness-maximizing decision depends not only on risk and reward of each patch, but also the
energetic state of the scavenger (y-axis) and the week within an annual cycle (x-axis). b)
Interaction between coyotes and a cougar at a kill site monitored by remote camera. c) Possible
influences of dominant predators in a carnivore community: 1) dominant predators do not
influence other carnivores (null hypothesis), 2) dominant predators negatively influence other
carnivores through interference competition or interspecific killing (suppression hypothesis), 3)
dominant predators positively influence other carnivores due to resource provisioning via
scavenging (facilitation hypothesis), and 4) dominant predators both positively and negatively
influence other carnivores (provision-predation hypothesis).
588
589

590
591 **Figure 2.** Study area. Left panel: Known and predicted cougar kill sites (blue points) and 95%
592 kernel density home range estimates (colored polygons) for collared cougars. Right panels, top to
593 bottom: coyote, black bear and bobcat 95% kernel density home range estimates (colored
594 polygons). All panels: Starkey Experimental Forest and Range (black polygon).
595
596

597
Figure 3. Attraction of coyotes to cougar kill sites. The first column presents marginal response
plots for step selection functions showing the relative probability of selection as a function of the
distance to the kill site. The second column presents predictive maps showing the relative
probability of selecting for areas on the landscape as a function of the distance to the nearest kill
site (black triangle). Warm colors indicate selection and cool colors indicate avoidance. In both
columns, each row shows the predicted response at 1, 15, and 30 days after the kill was made.
604
605

606
Figure 4. Left panel: Remote camera photos of a) cougar feeding on an adult female elk, c)
coyotes scavenging a cougar kill, e): coyote killed by cougar. Right panel: plots showing b)
percentages of deer and elk in carnivore diets determined from DNA-metabarcoding of scats
(except for cougars in which diet estimates came from kill site investigations), d) scavenging rates
of carnivores at cougar kill sites documented by remote cameras, and f) the estimated proportion
of the population killed annually by cougars.
613

614
Figure 5. Remote-camera based estimates of daily visits to cougar kill sites by coyotes (panel a)
and black bears (panel b). The thick line displays the predicted number of visits for coyotes (solid
blue line) and bears (dotted tan line). Shaded regions indicate 95% confidence interval for the
predictions. The vertical bars display the mean number of visits for every day since the kill was
made, across all kill sites monitored (N = 28). The predictions assume that the kill was a large
ungulate (i.e., the binary factor variable for prey size = 1) and that a cougar was present (i.e., the
binary factor variable for cougar present = 1).
622

	GPS collars		Scats	Cameras	Clusters
	Cougar presence	Kill site presence	Diet	Carcass Visitations	Intraguild predation
	Suppression	Facilitation	Facilitation	Facilitation	Suppression
	Suppression	Null	Facilitation	Facilitation	Null
	Null	Null	Null	Null	Null

Figure 6. Weight of evidence for support of suppression, facilitation, and null hypotheses regarding the influence of a dominant predator, cougars, on subordinate or co-occurring carnivores: coyotes, bears, and bobcats.

628 **Table 1.** Parameter estimates for landscape and cougar variables influencing the relative
629 probability of selection in step-selection functions in coyotes, black bears, and bobcats. PVT =
630 potential vegetation type, CP = cougar present, D2C = distance to nearest cougar, KP = kill
631 present, D2K = distance to nearest kill, and TSK = time since kill.
632

	Coyote			Bear			Bobcat		
	β	*SE*	*P*	*β*	*SE*	*P*	*β*	*SE*	*P*
CP × D2C[1]	0.20	0.08	0.01	0.55	0.15	<0.01	0.26	0.26	0.33
KP × D2K[1]	-0.26	0.05	<0.01	0.01	0.10	0.93	0.11	0.17	0.54
KP × D2K[1] × TSK[1]	0.07	0.02	<0.01	-0.02	0.04	0.59	-0.08	0.07	0.26
Canopy Cover	-0.13	0.01	<0.01	0.36	0.01	<0.01	0.32	0.01	<0.01
Distance to Road[1]	0.09	0.01	<0.01	0.13	0.01	<0.01	0.10	0.02	<0.01
Distance to Water[1]	0.02	0.01	0.01	-0.01	0.01	0.44	0.01	0.02	0.45
Ruggedness	-0.06	0.01	<0.01	0.16	0.01	<0.01	0.02	0.02	0.23
PVT, Wet Forest[2]	-0.09	0.02	<0.01	0.42	0.03	<0.01	0.98	0.07	<0.01
PVT, Dry Forest[2]	-0.08	0.02	<0.01	0.26	0.03	<0.01	0.71	0.07	<0.01
PVT, Other[2]	-0.25	0.15	0.09	1.39	0.11	<0.01	1.26	0.31	<0.01
Step Length[1]	0.03	<0.01	<0.01	0.04	0.01	<0.01	0.05	0.01	<0.01
Turning Angle[3]	-0.09	0.01	<0.01	0.04	0.00	<0.01	0.22	0.02	<0.01

633 [1]Indicates the variable was natural log transformed; [2]the reference category was Grassland;
634 [3]indicates the cosine of the variable was used.