TITLE PAGE

Title: Water levels primarily drive variation in photosynthesis and nutrient use of scrub

Red Mangroves in the southeastern Florida Everglades

Authors: J. Aaron Hogan[1*], Edward Castañeda-Moya[2], Lukas Lamb-Wotton[1], Tiffany

Troxler[2,3], Christopher Baraloto[1]

Author Affiliations: [1] Department of Biological Sciences, Florida International University,

Miami, FL USA; [2] Institute of Environment, Florida International University, Miami, FL

USA, [3] Department of Earth and Environment, Florida International University, Miami,

FL, USA.

* Corresponding Author: J. Aaron Hogan, email: jhogan@fiu.edu

ORCiDs:

J. Aaron Hogan: 0000-0001-9806-3074; Edward Castañeda-Moya: 0000-0001-7759-4351;

Lukas Lamb-Wotton: 0000-0002-0712-9005, Troxler: 0000-0007-7100-7742, Baraloto: 0000-

0001-7322-8581

Authorship statement: The conception and design of the study and the acquisition of data

were conducted by JAH, EC-M, and LL-W. Data analyses were performed by JAH and EC-M.

Interpretation of the data was carried out by JAH, EC-M, LL-W, TT, and CB. The manuscript

was drafted by JAH with the help of EC-M. All authors have contributed to the final version of

the manuscript and have approved its submission to the journal.

Data deposition: The data collected and used in this study have been archived and are

available through the Environmental Data Initiative data repository:

Hogan, J.A., E. Castaneda, and L. Lamb-Wotton. 2021. FCE LTER Taylor Slough/Panhandle-7

Site Scrub Red Mangrove (*Rhizophora mangle*) Leaf Gas Exchange Data, Florida, USA from

January-December 2019 ver 1. Environmental Data Initiative.

https://doi.org/10.6073/pasta/27f6332609eb1ef6d398c7855855f2e3 (Accessed 2021-01-26).

Note: this data archive also contains R code for running the linear mixed effects models using

the archived datasets.

ABSTRACT

We investigated how differences in mangrove island micro-elevation (i.e., habitat: center vs.

edge) affect tree physiology in a scrub mangrove forest of the southeastern Everglades. We

measured leaf gas exchange rates of scrub *Rhizophora mangle* trees monthly during 2019,

hypothesizing that CO_2 assimilation (A_{net}) and stomatal conductance (g_{sw}) would decline with

increases in water level and salinity, with larger differences at mangrove island edges than

centers, where physiological stress is greatest. Water levels relative to the soil surface varied

between 0 and 60 cm, rising during the wet season (May-October) relative to the dry season

(November-April). Porewater salinity ranged from 15 to 30 ppt, being higher at mangrove island

edges compared to centers. A_{net} maximized at 15.1 μmol m^{-2} $s^{-1,}$ and g_{sw} was typically <0.2 mol

m^{-2} s^{-1}, both of which were greater in the dry than the wet season and greater at island centers

than edges with the magnitude to seasonal variation being roughly equal to variation among

habitats. After accounting for season and habitat, water level had a positive effect on A_{net} in

both seasons, but no effect on g_{sw}. Our findings suggest that water levels are a main driver of

variation of leaf gas exchange rates in scrub mangroves of the Florida Everglades, while also

constraining A_{net} more than g_{sw}, and that the interaction between permanent flooding and habitat

varies with season as physiological stress is alleviated at higher-elevation mangrove island

center habitats during the dry season. Additionally, habitat heterogeneity leads to differences in

nutrient and water acquisition and use between trees growing in island centers versus edges,

creating distinct physiological controls on leaf photosynthesis, which could ultimately affect

carbon flux dynamics of scrub mangrove forests across the Everglades landscape.

Keywords: Scrub mangroves, Florida Coastal Everglades, photosynthesis, porewater salinity,

hydroperiod, *Rhizophora mangle*.

INTRODUCTION

Global climate change is affecting coastal mangrove ecosystems in an unprecedented manner, principally through increased flooding and saltwater intrusion (Pezeshki et al. 1990a, Yu et al. 2020). Increases in flooding severity and salinity due to sea-level rise (SLR) have the potential to push ecosystems to degraded alternative stable states, where biogeochemical cycles (e.g., carbon sequestration and storage potential) are impaired (Neubauer et al., 2013, Tully et al. 2019, Yu et al. 2020). Mangrove wetlands are particularly susceptible to SLR because of their position at the boundary between terrestrial and marine ecosystems (Field 1995, Ellison and Farnsworth 1997). Mangrove species have developed large variation in key life-history traits, such rates of photosynthesis, water-use and nutrient-use efficiencies, and growth rates and biomass allocation ratios in response to the interactions among resources (e.g., light and nutrients), regulators (e.g., salinity, sulfides), and hydroperiod gradients (e.g., frequency, depth, and duration of inundation – Twilley and Rivera-Monroy 2005, Alongi 2008, Twilley and Rivera-Monroy 2009, Castañeda-Moya et al. 2013). Due to such physiological flexibility and the large carbon sequestration and storage capacity of mangroves across a variety of geomorphic settings (e.g., karstic vs. deltaic; Mcleod et al. 2011, Murdiyarso et al. 2015, Lovelock et al. 2017, Rovai et al. 2018), there is an increasing need to strengthen our understanding of the effects of SLR and saltwater intrusion on mangrove tree physiology to assess trajectories of ecosystem structure and function in response to global change drivers.

Scrub mangrove forests, dominated by *Rhizophora mangle,* are common in Caribbean karstic environments (Cintron et al. 1978, Lugo and Snedaker 1974). The stunted physiognomy (i.e., reduced growth and development) of scrub mangroves results from severe nutrient (e.g., phosphorus, P) limitation, prolonged or permanent inundation with little tidal influence, and seasonal water stress (Feller 1995, Koch and Snedaker 1997, Cheeseman and Lovelock 2004, Medina et al. 2010). Scrub mangrove forests develop distinct landscape patterning, forming mangrove island clusters with higher elevations than their surrounding shallow open-water ponds and channels (Figure 1A). Soil elevation differences are maintained through the vertical accumulation of biogenic material via root biomass and production, leaf litter accretion and wood deposition (McKee et al. 2007, McKee 2011, Krauss et al. 2014). For example, in scrub mangrove islands of the southeastern Florida Everglades, island center habitats have 66% more root biomass and 52% more root production than island edges (Castañeda-Moya et al. 2011), which leads to spatial differences in soil elevation among island habitats. These differences in soil elevation interact with environmental gradients (e.g., hydroperiod, salinity)

along the intertidal zone in complex ways to affect mangrove physiology (e.g., rates of net CO_2

assimilation – A_{net}, growth rates, or sap flux) at variable scales (Medina and Francisco 1997,

Twilley et al. 1998, Medina et al. 2010, Twilley et al. 2017).

Hydrological dynamics (e.g., depth and duration of inundation) can cause mangrove

physiological stress, which cascades to affect carbon cycle dynamics and other biogeochemical

processes across spatial and temporal scales (Medina 1999, Castañeda-Moya et al. 2013,

Twilley et al. 2017, 2019). Although mangrove species can tolerate flooded conditions, they are

still susceptible to damage if plants become completely submerged for days to weeks (Wanless

1998, Mendelssohn and McKee 2000, McKee 2011). Inundation stress typically decreases

rates of leaf gas exchange (e.g., A_{net}, transpiration) and tree growth in mangroves (He et al.

2007, Cardona-Olarte et al. 2013). For example, greenhouse studies have revealed a 20%

reduction in maximum A_{net} when mangrove seedlings and saplings were subjected to short-term

intermittent seawater flooding (6 to 22 days, Krauss et al. 2006). Mangrove leaf gas exchange

is further affected by how seawater flooding interacts with fresh water and nutrient inputs

(Wolanski 1992). For instance, a reduction in stomatal conductance (g_{sw}) and leaf water

potential in *Bruguiera gymnorrhiza* seedlings occurred when exposed to prolonged flooding for

up to 80 days with 33% seawater compared to the control plants; however, seedlings flooded

with fresh water for 80 days showed an increase in both parameters (Naidoo 1983). In contrast,

seedlings of *Avicennia germinans* and *Laguncularia racemosa* exposed to permanent flooding

with 23% seawater showed no change in g_{sw}, A_{net}, or intrinsic water use efficiency (*wue*), but

had reduced leaf area (Krauss et al. 2006). Hydrologic conditions can further negatively

influence mangrove physiology through the interaction with soil phytotoxins (i.e., sulfides),

produced as by-products of low oxygen availability and soil redox conditions due to permanent

flooding, which can potentially depress water and nutrient uptake, and affect rates of leaf gas

exchange (Nickerson and Thibodeau 1985, McKee 1993, Ball 1996, Pezeshki and DeLaune

2012, Lamers et al. 2013). Regarding permanently inundated scrub mangroves, such as those

in the southeastern Florida Everglades, it is not entirely understood how seasonal dynamics

interact with inundation levels to influence leaf and forest carbon uptake dynamics.

Mangroves are highly adapted to tolerate salt stress, yet salinity has the greatest impact

on forest productivity, tree growth rates, and rates of leaf gas exchange. The negative effects of

increasing salinity are most evident along steep salinity gradients (i.e., those >30 ppt),

particularly in dry environments (Lugo and Snedaker 1974, Cintron et al. 1978, Medina and

Francisco 1997, Reef and Lovelock 2015). Salt stress variably affects mangrove tree

physiology, depending on species-specific salt tolerance levels and mechanisms for processing salt (Parida and Jha 2010, Reef and Lovelock 2015). For example, $R.$ mangle naturally inhabits Neotropical environments with salinities from near zero to around 35 ppt but may also be found in dry coastal environments with salinities up to 50-60 ppt (Cintron et al. 1978, Cardona-Olarte et al. 2006). All mangroves have the ability to exclude salt through the roots; however, $R.$ mangle is a highly-efficient salt excluder because its roots largely prevent salt from entering the plant. Additionally, $R.$ mangle lacks the excretory glands that other mangrove species (e.g., $L.$ racemosa) use to excrete salt once it has entered the plant. As such, the xylem of $R.$ mangle is 100 times less saline than seawater (Scholander et al. 1962, Scholander 1968, Medina and Francisco 1997, Tomlinson 2016) because of the Casparian strip (Lawton et al. 1981) and ultrafiltration by cell membranes in the thick aerenchyma and cortical layers of its root tissues (Field 1984, Werner and Stelzer 1990). However, some salt still enters the plant through the roots, which has a deleterious effect on the physiology of $Rhizophora$ trees, causing decreases in growth and A_{net} rates, and water and nutrient use efficiencies (Ball 1988, Clough and Sim 1989, Lugo et al. 2007, Medina et al. 2010, Cardona-Olarte et al. 2013).

Mangrove A_{net} varies widely with the environment (e.g., water and salinity levels) and nutrient availability. A_{net} for $R.$ mangle maximizes around 20 µmol m^{-2} s^{-1} (Golley et al. 1962, Bjorkman et al. 1988, Lin and Sternberg 1992, Lovelock and Feller 2003, Lugo et al. 2007, Ball 2009); however, A_{net} for scrub mangroves is lower, generally ranging from <5 µmol m^{-2} s^{-1} (Golley et al. 1962, Cheeseman et al. 1997, Cheeseman and Lovelock 2004) to roughly 13 µmol m^{-2} s^{-1} (Lugo et al. 2007, Barr et al. 2009). A field study from Jobos Bay in southern Puerto Rico demonstrated a significant decrease in $R.$ mangle A_{net} (from 12.7 to 7.9 µmol m^{-2} s^{-1}) and g_{sw} (from 0.28 to 0.19 mol m^{-2} s^{-1}) when comparing fringe habitats at 35 ppt salinity to inland salt flat habitats at 80 ppt (Lugo et al. 2007). Reductions in A_{net} and g_{sw} were accompanied by changes in leaf morphology (i.e., smaller specific leaf area, SLA), reduced nutrient-use efficiency, and increased nutrient resorption, which demonstrates how environmental effects on mangrove physiology can have consequences for ecosystem functioning. Thus, increasing salinity decreases A_{net} and g_{sw} and increases intrinsic water use efficiency (wue, defined as A_{net}/g_{sw}) in mangroves, with $Rhizophora$ species exemplifying these trends (Ball 2009, Clough and Sim 1989). Moreover, the high salt tolerance of mangrove species leads to strong stomatal control, which creates dynamics between A_{net} and water use which depend on relative reduction in transpiration rates versus the degree to which leaves are biochemically limited to fix carbon (e.g., via RUBISCO carboxylation efficiency versus RUBP regeneration) at low stomatal conductance (Sobrado 2000, Lovelock and Feller 2003, Ball 2009). For instance, $R.$ mangle

has succulent leaves with lower *wue* compared to more salt-tolerant species (i.e., *A. germinans*

or *L. racemosa*); however, *R. mangle* has greater efficiency of water transport in stems than

more salt-tolerant species (Sobrado 2000). Thus, when considering the effects of salinity on

leaf gas exchange rates, plant water use must be considered in concert because both A_{net} and

g_{sw} decline in a similar fashion with increasing salinity, effectively creating co-limitation of

photosynthesis at moderate to high salinities (Ball 2009).

159   In mangrove wetlands of the Florida Everglades, variation in environmental gradients

including hydroperiod (e.g., duration of inundation) and soil P availability control mangrove

forest structure and function (e.g., litterfall production) across the coastal landscape (Chen and

Twilley 1999; Castañeda-Moya et al. 2011, 2013). However, it is not completely understood

how the interaction between water level dynamics and salinity affect *in situ* rates of mangrove

leaf gas exchange in this region. Experimental evidence using *R. mangle* seedlings from south

Florida showed that inundation created a greater degree of physiological stress than did salinity

levels; however, salinity accelerated the adverse effects of inundation stress on leaf function

over time (Cardona-Olarte et al. 2013). In contrast, other studies have reported no clear effect

of water levels or flooding duration on rates of mangrove gas exchange, although inundation

duration decreased variability in leaf gas exchange measurements (Hoppe-Speer et al. 2011).

Using Florida mangroves, Krauss et al. (2006) found that short-term intermittent flooding

decreased rates of leaf gas exchange relative to unflooded or permanently flooded greenhouse-

grown seedlings, but that for *in situ* established *R. mangle* saplings growing along a natural tidal

inundation gradient along Shark River in the southwestern Everglades, flooding led to increases

in A_{net} and *wue*. Permanent flooding leads to decreases in A_{net} and g_{sw} rates in most wetland

plants (Kozlowski 1997); however, how inundation dynamics interact with salinity along the

intertidal zone to influence mangrove physiology at different spatial and temporal scales in south

Florida mangroves remains largely unknown. Further, global change-driven SLR and saltwater

intrusion in South Florida coupled to large-scale freshwater diversion have accelerated

mangrove encroachment into inland freshwater wetlands over the past 60 years (Ross et al.

2000). As sea levels continue to rise, it is imperative to further our understanding of the

magnitude of the effects of inundation and salinity on mangrove physiology and subsequent

ecosystem functioning (e.g., carbon flux) in the region.

183   Here, we present a comprehensive, one-year analysis of the spatial and seasonal

effects of salinity (surface and porewater) and water levels on rates of leaf gas exchange of *R.*

mangle scrub mangroves in southeastern Florida Everglades. We focused our sampling on

mangrove islands with noticeable micro-elevational differences to understand the magnitude of

influence of water levels and salinity on *R. mangle* tree physiology between mangrove island

center and edge habitats. We addressed the following questions: (1) how do rates of leaf gas

exchange (e.g., A_{net}, g_{sw}) vary with mangrove island micro-elevation (center vs. edge habitats)?

(2) how does leaf gas exchange respond to seasonal changes in salinity and water levels? (3)

how do water- and nutrient-use efficiencies of *R. mangle* leaves vary between mangrove island

center and edge habitats. We hypothesized that A_{net} would be greater for *R. mangle* leaves

located in higher elevation center habitats relative to lower elevation mangrove edges. We also

expected that A_{net} should vary little with season (i.e., <2 μmol CO_2 m^{-2} s^{-1}) and that seasonal

variation in g_{sw} would be less than variation in A_{net}, relative to the range of variability among

leaves because of strong control on g_{sw} and a potential for decoupling to some degree between

g_{sw} and A_{net}. Moreover, given that scrub mangroves in Taylor River basin are strongly limited by

phosphorus (i.e., soil N:P = 102-109 – Castañeda-Moya et al. 2013), *R. mangle* plants should

have high rates of P resorption. Finally, we predicted that mangrove island centers function at a

higher physiological level (i.e., with greater rates of A_{net} and g_{sw}) due to lower levels of

inundation and salt stress. Thus, mangrove trees in island centers should have greater *wue*

(Ball 2009) and higher relative rates of nutrient resorption (Lugo et al. 2007, Medina et al. 2010)

than trees at island edges.

METHODS

Study site

This study was conducted in the southeastern region of Everglades National Park in a

mangrove site known as Taylor Slough/Panhandle-7 (TS/Ph-7: 25.197°N, 80.642°W, Figure 1B),

one of the six mangrove sites established in 2000 as part of the Florida Coastal Everglades

Long-Term Ecological Research (FCE-LTER) program (Childers 2006; http://fcelter.fiu.edu).

TS/Ph-7 is located approximately 1.5 km inland from Florida Bay in the downstream section of

the Taylor River. Mangroves zones at TS/Ph-7 are dominated by scrub *R. mangle* L. trees, with

clusters of *L. racemosa* L. and *Conocarpus erectus* L. – a mangrove associate, intermixed with

low densities of freshwater grasses *Cladium jamaicense* (Crantz) Kük and *Eleocharis cellulosa*

Torr. (Loveless 1959). Mangrove tree heights reach 1.5 to 2 m (Ewe et al. 2006).

The substrate at this site is organic mangrove peat soil (~1 m depth) overlying the karstic

bedrock (depth ~1.5-2 m, Table 1; Castañeda-Moya et al. 2011, Ewe et al. 2006). Surface (0-

45 cm depth) soils at TS/Ph-7 have high organic matter content (71%), low bulk density (0.16 g

cm^{-3}), low total nitrogen (TN, 2.5 mg cm^{-3}) and low total phosphorus (TP, 0.06 mg cm^{-3})

concentrations, resulting in a highly P-limited environment with soil N:P ratios of about 102

(Castañeda-Moya et al. 2013). Mangrove zones in Taylor River are permanently flooded for

most of the year, with an average annual flooding duration averaging 360 d yr^{-1} from 2001 to

2005. This results in anoxic soil conditions and buildup of porewater sulfide (range: 0.5-2 mM)

throughout the year that constrains mangrove growth (Castañeda-Moya et al. 2011, 2013). The

tidal effect is negligible in Taylor River, and water flow and hydrology are determined by

seasonal precipitation, upland runoff, and wind (Michot et al. 2011, Sutula et al. 2001). The

interaction between low P fertility and permanent flooding conditions results in the formation of

scrub forests with restricted tree height and aboveground productivity, and high root biomass

allocation and high root:shoot ratios compared to riverine mangrove forests along Shark River

estuary in southwestern FCE (Ewe et al. 2006, Castañeda-Moya et al. 2011, 2013).

South Florida has a subtropical savanna climate per the Köppen climate classification,

where the average air temperature is between 20 and 30°C and relative humidity is high (70-

80%). Rainfall and evapotranspiration vary interannually and average 1500 and 1300 mm $year^{-1}$,

respectively (Abiy et al. 2019). In the Everglades, 60% of the precipitation occurs during the

wet season, and only 25% during the dry season (Duever et al. 1994). Analysis of long-term

(110-year) rainfall trends for South Florida has shown that the annual hydrologic regime can be

divided into two seasons: a wet season from May to October and a dry season from November

to April (Abiy et al. 2019). For the 2019 calendar year, temperature and relative humidity data

were collected from an eddy covariance flux tower installed at TS/Ph-7 and operational since

December 2016. Rainfall data were collected from a nearby meteorological station (station

name: "Taylor_River_at_mouth") managed by the US Geological Survey as a part of the

Everglades Depth Estimation Network (https://sofia.usgs.gov/eden).

Experimental design

Eight distinct mangrove islands of similar size (3-5 m in diameter) were selected for

repeated measurements of leaf photosynthesis and physicochemical variables from January to

December 2019. Mangrove islands were selected within previously-established permanent

vegetation plots (two 20×20 m plots) based on their location relative to the shoreline (i.e., Taylor

River), with four islands located in the fringe mangrove zone (~50-60 m from the edge) and four

islands located in the interior forest (~100-110 m inland; Figure 1B). Mangrove islands with

distinct micro-elevational gradients were selected, having higher soil elevation center habitats

and lower elevation edge habitats. Mangrove islands are surrounded by open water ponds

(Figure 1A) and remain flooded for most of the year, except the center island habitats during the
dry season (Castañeda-Moya et al. 2011, 2013, Figure S1).

Within each mangrove island, a higher-elevation center and a lower-elevation edge
habitat were each permanently marked with an aluminum rod exposed 1.5 m above the soil
surface and buried approximately 1 m below the soil surface. Soil surface elevation was
measured for all mangrove islands at both habitats, in addition to six measurements in the
adjacent shallow ponds surrounding mangrove islands. Measurements were taken using real-
time kinematics referenced to the 1988 North American Vertical Datum (NAVD88) with a
Trimble R8 global navigation satellite system receiver (Trimble; Sunnyvale, CA, USA), which
has a horizontal accuracy of ±1 cm and vertical accuracy of ±2 cm.

Water level and salinity measurements

Water levels relative to the soil surface were measured monthly with a meter stick at
each of the permanent aluminum rods established at all island habitats. Continuous
measurements of water levels relative to the soil surface were recorded for the duration of the
2019 calendar year (see Figure S2 for details). Continuous data were used to confirm trends in
water levels and porewater salinity measurements made by hand across islands. We use the
measurements taken by hand at each island as the predictors in our models of leaf gas
exchange. A porewater sample was collected at 30 cm depth at each habitat using a 60 ml
syringe attached to a stopcock and a rigid tubing probe (3/16" Ø). Porewater temperature and
salinity were measured using a handheld YSI conductivity-salinity-temperature meter (model
Pro 30, YSI Inc., Yellow Springs, OH, USA). A sample of surface water (when present) was
also collected at each island habitat to measure salinity and temperature.

Leaf gas exchange measurements

Photosynthetic gas exchange measurements of *R. mangle* leaves were conducted once
a month (9:00 AM to 1:00 PM, typically during sunny days) at eight scrub mangrove islands
from January to December 2019 using a Li-COR Li-6800 portable photosynthesis system (Li-
COR Inc., Lincoln, NE, USA). At each island habitat (center vs. edge), five mature green leaves
were randomly selected from top mangrove branches. Fully developed and healthy (i.e.,
without herbivory) green leaves from the second-most distal pair of leaves on the leaf node
were chosen. The Li-6800 was clamped onto each leaf and held until machine stability was
reached (which typically happened in 2-3 minutes), wherein data points were logged.

The environmental configuration of the Li-6800 was: flow rate of 600 µmol s^{-1}, 50-70%

relative humidity of the incoming air (slightly drier than ambient air to prevent condensation in

the instrument), 400 µmol mol^{-1} CO_2 concentration, and light level of 1000 µmol m^{-2} s^{-1}, which

was determined to be non-limiting and similar to ambient environmental conditions. We used

five stability criteria, which were all assessed over a 15 s interval: the slope of A_{net} being <1

µmol m^{-2} s^{-1}, the slope of the concentration of intracellular CO_2 (c_i, which is a calculated

parameter using the difference in CO_2 concentrations between IRGAs in the Li-6800) being < 5

µmol mol^{-1}, the slope of g_{sw} being < 0.5 mol m^{-2} s^{-1}, the slope of the transpiration rate (E) being

less than 1 mol m^{-2} s^{-1}, and the slope of the difference in air water vapor concentration between

the sample and reference IRGA (ΔH_2O) being less than 1 mmol mol^{-1}. All five stability criteria

were met before logging data. Air temperature within the leaf chamber was not controlled, but

allowed to vary with the ambient conditions at the site, ranging from 26.1 to 32.0°C. Thus, leaf

temperatures ranged from 25.85 - 32.44 °C, averaging 29.41 ± 0.08 °C, in the wet season, and

ranged from 25.06 - 28.72 °C, averaging 26.79 ± 0.04 °C, in the dry season. We calculated

intrinsic *wue* as the ratio of leaf net CO_2 uptake to leaf gas exchange (i.e., $A_{net}/g_{sw}/1000$, where

we divide by 1000 to get *wue* in mmol mol^{-1}).

Measurement of leaf functional traits, nutrient content, and isotopic signatures

During the monthly photosynthesis measurements in February, May, August, and

November, measured mature green leaves (n = 5 per habitat, 40 in total) were collected at half

of the islands (four of the eight islands with two per location) for determination of leaf functional

traits and total carbon (TC), nitrogen (TN), and phosphorus (TP) content. Leaves were

numbered, placed in a sealed, moist bag to prevent water loss, and transported to the

laboratory in a cooler with ice for further analyses. Five senescent (i.e., yellowing) leaves were

also collected from trees in the same islands at the same time to determine carbon and nutrient

content. At the laboratory, leaves were removed from bags, wiped dry, and immediately

weighed to obtain leaf fresh mass. Green leaves were then scanned at high resolution and

oven-dried for at least 72 hours at 60°C to constant weight before recording their dry mass.

Leaf area was measured using ImageJ (Schneider et al. 2012). Leaf dry mass was recorded

and used to calculate leaf dry matter content (LDMC) as the ratio of the dry leaf mass (in mg) to

its fresh mass (in g, mg g^{-1}), percent leaf water content (1000-LDMC; %), and SLA, the ratio of

leaf dry weight to leaf area (cm^2 g^{-1}). These methods followed Cornelissen et al. (2003).

For nutrient analyses, composite leaf samples containing the five leaves from each

island habitat per collection were ground into a fine powder using a vibrating ball mill

(Pulversette 0, Frtisch GmbH, Idar-Oberstein, Germany). Green and senescent leaf samples

were stored in scintillation vials at room temperature and analyzed separately. Leaf TC and TN

content were determined for each sample with a NA1500 elemental analyzer (Fisons

Instruments Inc., Danvers, MA, USA). TP was extracted using an acid-digest (HCl) extraction,

and concentrations of soluble reactive P were determined by colorimetric analysis (Methods

365.4 and 365.2, US EPA 1983). Leaf C and N bulk isotopic signatures (δ^{13}C, δ^{15}N) were

analyzed on a Thermo Scientific Delta V Plus CF-IRMS coupled to an 1108 elemental analyzer

via a ConFlo IV interface (Thermo Fisher Scientific, Waltham, MA, USA). All C and N analyses

were conducted at the Southeast Environmental Research Center Analysis Laboratory (SERC-

NAL). SERC-NAL follows strict internal and external QA assurance practices and is NELAC

Certified for non-potable-water-General Chemistry under State Lab ID E76930.

Using leaf carbon isotope fractionation values, we calculated the concentration of

intracellular CO_2 and plant water use efficiency integrated over the lifespan of the leaf tissue

samples (i.e., intrinsic water use efficiency, *WUE*) via methods described by O'Leary (1988) and

Marshall et al. (2007) (and outlined in Lambers et al. 2008). We used an ambient concentration

of atmospheric CO_2 of 408 µmol mol^{-1} for our calculations, which is a conservative estimate for

the 2019 calendar year and was indicative of the atmospheric conditions at the site based on

IRGA measurements from an eddy covariance tower at the site. Thus, the equation used to

calculate c_i and *WUE* from carbon isotope data were: c_i = ((-8.5 - δ^{13}C - 4.4) ÷ 22.6) × 408), and

WUE = (408 × (1 - c_i ÷ 408)) ÷ 1.6, where c_i is the value derived from the previous equation.

Additionally, the following equation was used to calculate resorption of N and P using green (G)

and senescent (S) leaf nutrient content: Relative resorption (%) = ((G - S) ÷ G × 100) (Pugnaire

and Chapin 1993).

Statistical analyses

Repeated measures analysis of variance (ANOVA) was used to test for differences in

water level, surface water salinity, and porewater salinity among locations (fringe and interior),

island habitats (center and edge), and season (wet and dry), as well as for the interaction

between these effects and season, which was used as the repeated measure. For the repeated

measures ANOVA, islands were nested within locations and treated as experimental units. All

effects were considered fixed, except for when testing for significant differences in habitat,

which included location as a random effect to account for the nested structure of the sampling

scheme. One-way ANOVAs were used to test for differences in soil surface elevation among

locations and habitats, and the interaction between them. Two-way ANOVAs were carried out

for all leaf functional traits and nutrient concentrations, where comparisons were made across

all habitat and season combinations. Tukey HSD post-hoc tests were used to identify significant

pairwise comparisons when ANOVAs indicated statistical differences. Repeated measures

ANOVAs were performed using PROC MIXED (SAS Institute, Cary, NC, USA) and the one-way

and two-way ANOVAs were performed in R v3.5.1 (R Core Team 2018).

We constructed linear mixed-effects models (with a Gaussian error distribution and

identity link function) to address our research questions. Island habitat and season were

included as fixed effects in the models to address questions 1 and 2, respectively, with water

levels and porewater salinity being also included as the continuous covariates to parse out their

marginal effects. We couple inference from these models to leaf nutrient analyses and our

measurements of the hydrological environment to inform about nutrient and water use of *R.*

mangle (question 3). Prior to model fitting, response variables were confirmed to meet the

assumptions of data normality. Four separate models were constructed, one for each of four

gas exchange variables of interest, A_{net}, g_{sw}, c_i, and *wue*. For each model, fixed effects for

season (wet and dry), habitat (center and edge), porewater salinity and water level were

considered, including interaction terms for water level and porewater salinity with season. In all

models, random intercept terms were considered for location (i.e., fringe vs. interior), islands,

and islands nested within location. Random slopes were explored but determined to not

improve model fits. The best-fit models were determined via stepwise model comparison using

AIC based on backward selecting random effects then backward selecting fixed effects, as

implemented with the 'lmerStep' function in the lmerTest R package (Kuznetsova et al. 2017).

The best fit models included a random intercept term for islands, which helped remove

variability in the data because of the sampling design. Random effects for location were

insignificant, signifying that most of the random variance in the gas exchange data was among

islands, which we consider as the experimental unit in all mixed-effects models. The mixed

effect models were fit using restricted maximum likelihood estimates via the lme4 R package

(Bates et al. 2015). Models were evaluated using model predicting, tabling, and plotting

functions from the sjPlot R package (Lüdecke 2018). All analyses were complete in R v3.5.1 (R

Core Team 2018).

RESULTS

Mangrove island micro-elevational differences and ecohydrology

Soil surface elevation (measured in relation to the NAVD88 datum) significantly declined

from mangrove island center to edge habitats from -0.14 ± 0.1 m at island centers to -0.4 ± 0.02

m at island edges, a mean difference of about 30 cm ($F_{1,20}$ = 108.42, p < .001; Table 1). Water

levels relative to the soil surface were significantly higher in the edge than in center habitats

($F_{1,178}$ = 178.33, p < .001), measuring on average 36.9 ± 1.4 cm in edge habitats, and 12.8 ± 1.2

cm in mangrove island centers (Table 2). We recorded water levels of 0 cm (i.e., non-inundated

habitats) in 10% of our measurements, and those were exclusive to mangrove island centers

during the dry season (Figure S1C). There was a significant effect of season ($F_{1,178}$ = 11.11, p

< .001) on water levels, where they increased from 17.05 ± 1.5 cm in the dry season to 30.4 ±

1.6 cm in the wet season (Table 2, Figure S1C).

Continuous water level data recorded at the fringe and interior mangrove zones

indicated a similar flooding trends between locations, with lower water levels during the dry

season and higher water levels in the wet season, up to 40-47 cm above the soil surface in both

locations (Figure S2). Water levels at the interior mangrove forest always remained higher than

those registered in the fringe mangrove zone (Figure S2A). Porewater salinity was significantly

different between habitats ($F_{1,178}$ = 91.45, p < .001) and seasons ($F_{1,178}$ = 17.87, p < .001), with

lower salinity values in the center (21.5 ± 0.3) of the islands relative to the edge (25.1 ± 0.3)

habitats, and slightly lower porewater salinity during the dry season (22.5 ± 0.4) than in the wet

season (24.1 ± 0.3; Table 2, Figure S1D). There was no significant interaction ($F_{1,178}$ = 0.26,

p > .05) between island habitats and seasons, indicating that the variation in porewater salinity

between habitats is independent of seasonality (Table 2). Surface water salinity was not

significantly different among center and edge habitats ($F_{1,163}$ = 2.36, p > .05), but increased

significantly from the dry to the wet season ($F_{1,163}$ = 8.97, p < .01, Table 2). There was also a

significant habitat-season interaction for surface water salinity, but a Tukey post-hoc HSD test

indicated that only island center habitats in the dry season were different from all other pairwise

comparisons (Table 2).

Rates of leaf gas exchange and their relationships to the hydrological environment

A_{net} measurements ranged from 0.1 to 15.1 µmol m^{-2} s^{-1}, with 90% of the observations

recorded between 2 and 14 µmol m^{-2} s^{-1} (see Figure S3). g_{sw} values were low, ranging from

<0.01 to 0.27 and averaging 0.1 mol m^{-2} s^{-1}(see Figure S3). Associated c_i values ranged from

40 to 377 and averaged 242 µmol mol^{-1}, with 98% of them being greater than 150 µmol mol^{-1}.

Lastly, measured rates of *wue* varied between >0.01 and 0.21 mmol CO_2 mol H_2O^{-1}, being

normally distributed about a mean value of 0.09 mmol mol^{-1}.

The linear mixed-effects model for A_{net} included fixed effects for island habitat, porewater

salinity, water level, season, and an interaction term for water level with season (Figure S4 &

414 Table S4). There was substantial variation in A_{net} rates among leaves (σ^2 of about 6 µmol m^{-2} s$^-$

415 ^{1}), and the random variation among islands was about 0.02 µmol m^{-2} s^{-1} (see Table S4). All

416 fixed effects were statistically significant ($p < .05$), except the interaction term, which was

417 marginally significant ($p = .05$) but greatly improved model fit. Mangrove edge habitats reduced

418 A_{net} by over 2.5 µmol m^{-2} s^{-1} relative to mangrove island centers (Figure 2). Seasonality had a

419 comparable negative effect, leading to an average decrease in A_{net} of just over 2 µmol m^{-2} s^{-1} in

420 the wet season relative to the dry season (Figure 2). After accounting for variation in the data

421 because of habitat and season, the marginal effects of water level and porewater salinity were

422 positive, albeit weak, leading to increases in A_{net} of <0.1 µmol m^{-2} s^{-1} per cm increase in water

423 level (Figure 3) or per ppt increase in porewater salinity (Figure 4). Therefore, as water levels

424 increased, A_{net} increased, with increases being consistent across habitats (Figure 3); a similar

425 pattern was observed in relation to soil porewater salinity, although the magnitude of increase in

426 A_{net} was smaller (Figure 4). These relationships of A_{net} with water level variability were

427 consistent across seasons, although rates of A_{net} were depressed during the wet season (Figure

428 2). The mixed-effects model for A_{net} fit satisfactorily for these types of linear mixed effects

429 models modeling leaf-gas exchange data using environmental predictors, explaining 24% of the

430 variation in the data, 22% of which was explained by ecohydrological data (i.e., fixed effects)

431 (Table S4).

432   g_{sw} was modeled using an identical mixed-effects model to that of A_{net} (Figure S5 &

433 Table S5). Generally, rates of g_{sw} were low, with 98% of g_{sw} measurements being <0.2 mmol

434 mol^{-1}. Random variance in g_{sw} among islands was negligible, being <0.01 mmol mol^{-1}. Leaf g_{sw}

435 in edge habitats was statistically lower than that of mangrove island centers ($p < .001$), being

436 depressed by about 0.02 mmol mol^{-1} (Figure 2). Water levels did not affect rates of g_{sw} ($p > .05$,

437 Figure 3, Table S5), and soil porewater salinity had a marginal effect ($p = .07$) on g_{sw}, where

438 conductance increased slightly at high salinities, after accounting for the effects of other

439 environmental variables in the model (Figure 4). The effect of season on rates of g_{sw} was

440 significant in the model, with the wet season leading to a 0.05 mmol mol^{-1} decrease in

441 conductance (Figure 2) and the interaction between water levels and season being statistically

442 significant (Figure 3). Overall, the mixed-effects model for g_{sw} did not fit the data as well as the

443 model for A_{net}, in that the model only explained about 12% of the variability in the data, with 9%

444 of its explanatory power coming from the environmental predictors (Table S5).

445   Although the model selection approach was the same as the other mixed-effects models,

446 the best-fit model for c_i was different from the models for A_{net} and g_{sw}. The model did not

include a fixed effect for soil porewater salinity (which was dropped out of the model in the

model selection procedure) but included all the same fixed effects as the models for A_{net} and g_{sw},

which were all statistically significant ($p < .001$), and a random intercept term for islands (Table

S6). Mangrove island edge habitats had consistently higher c_i values than island centers, being

about 27 μmol mol^{-1} greater (19 to 35, 95% CI; Figure 2). The marginal effect of season alone

was similar in magnitude to that of habitat; the wet season led to a decrease in c_i of 24 μmol

mol^{-1} (14 to 34, 95% CI) relative to the dry season (Figure 2, Table S6). Water levels, by

themselves (again, the marginal effect), led to a slight decrease in c_i but had a positive

interaction with season, indicating that the relative decrease in c_i due to increasing water levels

was suppressed during the wet season (Figure 3). The random intercept term in the model (for

islands) explained a considerable amount of variation in the data ($\sigma^2 = 128$ μmol mol^{-1}, with

$\tau_{island} = 66$ μmol mol^{-1}). The mixed-effect model for c_i fit the poorest of all four models,

explaining just under 12% of the variance in c_i, about 9% of which was explained by data from

the hydrological environment (Table S6).

Lastly, we modeled *wue* using a similar mixed-effects model to that of g_{sw}. In the model

for *wue*, all fixed effects were statistically significant ($p < .001$); however, the fixed effects were

more subtle in magnitude. Similar to the model for c_i, porewater salinity was not included in the

best-fit model. *wue* values were normally distributed about a mean value of 0.09 μmol mol^{-1},

with 83% of the data having values between 0.05 and 0.15 μmol mol^{-1}. Mangrove island edge

habitats had lower *wue* by 0.01 μmol mol^{-1} than island centers (Figure 2). The marginal effect of

water level, although being statistically significant in the model, was negligible; however, the wet

season caused an increase in *wue* by 0.02 μmol mol^{-1} relative to the dry season, with the

interaction between water level and season being slightly negative (Figure 2, Figure 3).

Random variation in *wue* structured across the eight mangrove islands measured was

minuscule, being < 0.01 μmol mol^{-1}. The model fit for the *wue* mixed-effects model was

comparable to and slightly better than the model for c_i, with fixed effects explaining just over 12%

of the variance in the data, about 9% of which was explained using the environmental predictors

(Table S7).

Rhizophora mangle leaf functional traits, nutrient content, and isotopic signatures

Leaf SLA values did not vary significantly between seasons ($F_{1,155} = 0.46$, $p > .05$) and

island habitats ($F_{1,155} = 3.07$, $p > .05$, Table 3), despite having some variation in SLA with

average values ranging from 29 to 40 cm^2 g^{-1}. Similarly, leaf water content was not significantly

different between all season-habitat combinations ($F_{1,155} = 0.32$, $p > .05$), despite a statistically

significant effect of season alone ($F_{1,155}$ = 9.10, p < .01), where leaf water content was greater in

the dry season (65.6 ± 0.3%) relative to the wet season (63.8 ± 0.4%, Table 3).

Leaf TC content ranged from 400 to 450 mg g^{-1} (Table 3) and was not different between

seasons ($F_{1,155}$ = 1.10, p > .05), habitats ($F_{1,12}$ = 0.10, ns), or their interaction ($F_{1,12}$ = 1.77,

p > .05). Leaf TN concentrations were higher in the dry season compared to the wet season

($F_{1,12}$ = 11.95, p < .01) and ranged from 8-10 mg g^{-1} (Table 3). There was no significant

difference ($F_{1,12}$ = 11.86, p > .05) in leaf TN between habitats nor a significant interaction ($F_{1,12}$ =

0.11, ns) between seasons and habitats (Table 3). Leaf TP content did vary between seasons

($F_{1,28}$ = 22.89, p < .001) and habitats ($F_{1,28}$ = 8.97, p > .05), but the interaction effect was not

significant ($F_{1,28}$ = 0.002, ns). Overall, mean leaf TP values ranged from 0.42 to 0.55 mg g^{-1}

across seasons and habitats, with higher concentrations during the dry season than in the wet

season and higher leaf tissue TP values in the island center habitats compared to edge habitats

(Table 3). Mean N resorption for *R. mangle* leaves was slightly similar across seasons and

habitats and ranged from 60 to 63% (Table 3). In contrast, P resorption of leaf tissue had a

broad range compared to that of N, ranging from 73.2 ± 6.2% (center, wet season) to 78.6 ± 0.2%

(edge, dry season) across seasons and habitats. Overall, P resorption of *R. mangle* leaves was

higher in the edge habitats relative to the center during both seasons (Table 3).

Patterns in green leaf carbon isotope signatures (δ^{13}C) mirrored those of leaf TN and TP

concentrations. Carbon isotopic fractionation was more negative during the wet season than in

the dry season ($F_{1,12}$ = 18.88, p < .01, Table 3, Figure S8), with no statistical difference between

habitats ($F_{1,12}$ = 1.17, p > .05). Green leaves bulk δ^{13}C values ranged from -25.9 to -25.1‰

across seasons and habitats (Table 3). Physiologically, the differences in carbon isotopic

fractionation were estimated to result in a maximum difference of about 10 µmol mol^{-1} c_i

between the center and edge habitats and a difference of 5 to 15 µmol mol^{-1} c_i within habitats

($F_{1,12}$ = 1.71, p > .05) because of seasonality ($F_{1,12}$ = 18.88, p < .01). These differences resulted

in slightly greater, but not statistically different, c_i values in mangrove island centers than in

edge habitats in the wet season; however, the opposite pattern was found during the dry season,

with c_i being about 10 µmol mol^{-1} greater in island edge habitats relative to their centers (Table

3). The interaction between season and habitat was marginally significant ($F_{1,12}$ = 4.53, p

= .055). Intrinsic water-use efficiency (*WUE*) was calculated from leaf δ^{13}C values; accordingly,

WUE was greatest in mangrove island centers during the dry season relative to all habitat-

season combinations. Additionally, *WUE* was significantly lower in the wet season than the dry

season ($F_{1,12}$ = 18.88, p < .01, Table 3). Mean leaf bulk δ^{15}N values were significantly ($F_{1,12}$ =

19.66, $p < .001$) higher in the center habitats (-0.60 ± 0.66‰) relative to the edge (-4.79 ±
0.66‰), but there was no difference between seasons ($F_{1,12} = 0.17$, $n.s.$) and no interaction
between season and island habitat ($F_{1,12} = 0.58$, $n.s.$, Figure S8A, Table 3).

DISCUSSION

Our first research question asked how mangrove island habitat affects rates of leaf gas
exchange. We can confirm our hypothesis that photosynthetic rates and stomatal conductance
are greater at island centers than edges (Figure 2). However, contrary to our expectation,
habitat driven-variation in rates of leaf gas exchange was roughly equal to seasonal variation,
with no apparent decoupling between A_{net} and g_{sw} (Figure 2). Our second research question
asked whether water levels or salinity exerted a greater effect on mangrove leaf gas exchange
in situ. Porewater salinities at TS/Ph-7 are relatively low (i.e., between 15 and 30 ppt) relative to
the levels of salinity at which leaf gas exchange rates of *R. mangle* are negatively affected (i.e.,
salinities > 35 ppt), and did not vary considerably over time. Therefore, we conclude that
inundation stress is the main driver of variation in *R. mangle* leaf gas exchange rates. Lastly,
we predicted that general physiological stress would be lower at island centers than island edge
habitats, leading to increased *wue* and higher rates of nutrient resorption. Indeed, intrinsic *wue*
was greater at island centers than edge habitats, with results being consistent across gas
exchange-measured *wue* and isotope-derived *WUE*. Water levels modulated leaf intrinsic water
use efficiency (Figure 3, Table 2), especially in the dry season. Patterns of nutrient resorption
were less clear but seem to indicate differences in leaf N and P concentrations on mangrove
island centers versus edges, illustrating differences in water and nutrient use of *R. mangle*
among habitats.

The effect of mangrove island habitat on leaf gas exchange

Our results showed significant differences in soil elevation of about 30 cm between
mangrove island habitats (Table 1), which had an apparent effect on *R. mangle* leaf gas
exchange rates (Figures 2, 3, 4). The soil elevation gradient at our study site is driven by
differences in mangrove root biomass and productivity between center and edge island habitats,
with higher total root biomass and productivity (top 0-90 cm of soil) observed in center habitats
compared to the edge habitats (Castañeda-Moya et al. 2011). Along this elevation gradient, we
measured clear differences in A_{net} and g_{sw} (Figure 2). A_{net} was nearly 3 µmol m^{-2} s^{-1} (or 20%)
greater at mangrove island centers than edges, and g_{sw} was >0.1 mol m^{-2} s^{-1} (or >37%) higher;
these differences were attributable to mangrove island habitat alone, after accounting for
variation explained by water level, salinity, or seasonality (i.e., marginal differences).

Associated c_i concentrations were about 30 µmol mol^{-1} (or 12%) lower and *wue* was >0.01

mmol mol^{-1} (or about 10%) greater at island centers than at island edges (Figure 3).

Thus, these findings support our first hypothesis about the effect of habitat micro-

elevation (center vs. edge) on A_{net}, with overall greater leaf gas exchange rates at mangrove

island centers compared to their edges. Interestingly, the effect of habitat on *R. mangle* leaf gas

exchange rates was similar in magnitude to the effect of season (Figure 3). The magnitude of

variation in A_{net} that we report in this study is slightly larger than the magnitude of variation

reported by Lin and Sternberg (1992) who found that A_{net} varied up to 2 µmol m^{-2} s^{-1} between

scrub and fringe *R. mangle* trees in the nearby Florida Keys. Furthermore, our A_{net}

measurements with average values between 5.7 µmol m^{-2} s^{-1} (edge habitat, wet season) and

10.2 µmol m^{-2} s^{-1} (center habitat, dry season, Figure 2), are within the range of values reported

for *R. mangle* interior scrub (5.3 µmol m^{-2} s^{-1}) and fringe (10 µmol m^{-2} s^{-1}) mangroves along a

distinct zonation pattern in the intertidal zone at Twin Cays, Belize (Cheeseman and Lovelock

2004). Island center habitats may also support greater access to mixed soil-groundwater

sources in the dry season and higher leaf gas exchange rates as compared with edge habitats

(Ewe et al. 2007). These results demonstrate the effect that higher elevation center habitats at

TS/Ph-7 have on alleviating inundation stress, which pervades scrub mangrove physiology,

making trees growing in center habitats in the dry season physiologically comparable to fringe

mangroves. Certainly, the stress relief is short lived when water levels rise in the wet season

(Table 2, Figure S1), and rates of leaf gas exchange are depressed once more (Figure 2. Figure

S3).

Seasonal signals in R. mangle *physiology with implications for ecosystem functioning*

We found that A_{net} varied over 2.5 µmol m^{-2} s^{-1} (17%), and g_{sw} varied about 0.03 mol m^{-2}

s^{-1} (11%) within habitats between the wet and dry seasons (Figure 3). Despite differences in

A_{net} and g_{sw} between seasons, we found no statistical differences in c_i and *wue* between

seasons, although there was some variation (Figure 3). These differences point to habitat-

specific optimization of the diffusion and uptake of CO_2 into (i.e., c_i) and the movement of water

vapor out of (i.e., *wue*) leaves (Cardona-Olarte et al. 2006, Barr et al. 2009, Reef and Lovelock

2015, Lopes et al. 2019). As precipitation and freshwater flow increased during the wet season,

water levels increased, and mangrove island centers experienced greater inundation levels

(Figures S1 & S2), resulting in decreased A_{net} and g_{sw} (Figure 3). A similar reduction in A_{net} and

g_{sw} was measured in mangrove island edge habitats during the wet season (Figures 3 & 4).

Although A_{net} was depressed in the wet season, the effect of inundation levels on reducing A_{net}

was consistent across seasons (Figure 4). g_{sw} showed a similar pattern to A_{net}, being highest in

mangrove island centers during the dry season (Figure 3). The effect of water levels on g_{sw},

however, resulted in increased g_{sw} in the wet season, an effect which was tempered during the

dry season (Figure 4).

In the Florida Everglades, irradiance peaks in April and May (Barr et al. 2009), and

rainfall and temperature reach maxima in June, July, and August (Figure 1 A, B). Thus,

photosynthetic demand for water is likely highest at the end of the dry season in April and

beginning of the wet season in May. During this time, we measured lower water levels and

porewater salinity values relative to the wet season. Barr et al. (2009) recorded earlier diurnal

and more considerable reductions in g_{sw} during late May versus July or August for mangroves at

Key Largo. Additionally, greatest A_{net} rates for tall fringe mangroves in the southeastern

Everglades occur from March to May (Barr et al. 2009). The difference in surface water and

porewater salinity (*Δsw-pw*) can be used as a proxy for tree transpiration (Reef and Lovelock

2015). Average *Δsw-pw* measured 10.7 and 8.1 ppt at mangrove island edges in the wet and

dry seasons, respectively, whereas it measured 7.8 and -0.3 in mangrove island centers in the

wet and dry seasons, respectively. Indeed, transpiration was highest in March and April (Figure

S3). Thus, photosynthetic demand for water is higher in the dry season in mangrove island

centers relative to edges or to either habitat in the wet season. The drying of the soils at slightly

higher elevation island center habitats in this scrub mangrove forest likely increases A_{net}.

Therefore, the seasonal variation in hydrology, mainly reductions in water levels and porewater

salinity during the dry season, albeit coupled with an increase in surface water salinity in this

study (Table 2), likely have key consequences for mangrove forest carbon fluxes at greater

spatial scales. Potentially drying of soils could also lead to an increase in ecosystem respiration

(Chambers et al. 2014). As such, future research could look at soil metabolic dynamics (e.g.,

soil respiration, microbial C and N, or changes in microbial communities) with hydrology and

season, which may show unique responses in this scrub *R. mangle* forest (Lovelock 2008,

Chambers et al. 2014).

The effect of salinity and water level on R. mangle *leaf gas exchange*

During 2019, the hydrological environment (Figure 1C,D) at our study site was

seasonally dynamic (Table 1) and tended to mirror patterns in local rainfall (Figure 1B). Water

level and porewater salinity both increased during the wet season (Table 2, Figure S1) from the

beginning of the rainy season in May through November. This likely led to increased water

column stratification via an increase in the freshwater lens (Hughes et al. 1998, Uncles et al.

1992). Indeed, the difference in surface water and porewater salinity increased in the wet

season, with surface water salinities decreasing, despite a slight increase in porewater salinities

(Table 1). When data were grouped by season, edge habitats were slightly more saline (about

4 ppt on average) than mangrove edges (Table 1), and there were no clear differences between

fringe and interior scrub mangrove zones (Figures S1 & S2). Comparing these changes in the

hydrological environment with previous years, long-term water level and porewater salinity data

at this site show that water level usually increases and porewater salinity usually decreases in

the wet season relative to the dry season (Castañeda-Moya et al. 2013). We measured the

opposite trend in 2019 with slight differences between seasons, likely because it was a wet year.

Total rainfall for 2019 (929 mm yr^{-1}; Figure S1) was 10% greater than total for rainfall 2018 (859

mm yr^{-1}; https://sofia.usgs.gov/eden).

Rates of mangrove leaf gas exchange (i.e., A_{net} and g_{sw}) typically decrease with

porewater salinity, especially along strong salinity gradients in the environment (i.e.,

gradients >30 ppt, Ball 2009, Clough and Sim 1989, Lugo et al. 2007). Porewater salinity was

only included in the linear mixed-effects models for A_{net} and g_{sw}, and its marginal effect was

minimal, slightly increasing A_{net} by <0.1 µmol m^{-2} s^{-1} per ppt increase in porewater salinity. Like

the effect of porewater salinity on A_{net}, the effect of porewater salinity on g_{sw} was small in

magnitude and consistent across seasons and mangrove island habitats but was not statistically

significant (Figure 4). The minimal influence of porewater salinity on leaf gas exchange is likely

due to the small seasonal and spatial variations that we observed in our study. Differences

were not large, maximizing at 16.4 ppt and averaging 5.2 ppt, especially when considering that

R. mangle frequently occupies natural habitats that have salinities greater than seawater (Reef

and Lovelock 2015), potentially up to 50-60 ppt (Cintron et al. 1978). At our study site, variation

in porewater salinity from long-term monitoring data (2001-2020) have shown similar

magnitudes of relatively minor variation in porewater salinity, with overall mean values ranging

from 19-22, and rarely exceeding 30 ppt (Castañeda-Moya et al. 2013). Additionally, long-term

variation in porewater salinity (<30 ppt) across the FCE mangrove sites (Shark and Taylor River

sites) is below the critical value of 65 ppt that influences forest structure and productivity across

the FCE landscape (Castañeda-Moya et al. 2013). Thus, the limited effects of salinity in our

linear mixed effects models, are likely broadly indicative of relatively weak salinity effects in both

scrub and tall *R. mangle* dominated forests of the Everglades . This is a significant finding given

that these scrub forests are adapted to relatively low salinities, and if salinity increases greatly

due to SLR and saltwater intrusion in the region, they will likely experience more stress

conditions affecting their physiological performance as observed in other studies in the

neotropics (Lugo et al. 2007).

The effects of inundation on *R. mangle* photosynthesis can be difficult to separate from

the effects of salinity, however the linear mixed modeling approach we used permitted doing so.

We found that the intermittent flooding conditions of mangrove island centers which averaged

water levels of 10-15 cm above the soil surface, allowed greater A_{net} and g_{sw} than permanently

flooded mangrove island edges, which averaged water levels of 30-40 cm. This indicates that

the hydrological regime in center habitats allows for mangrove soils to repeatedly flood and

desiccate, which may help the species to maintain optimal stem water potentials and g_{sw} (Ball

2009, Reef and Lovelock 2015). In typical greenhouse experiments where mangrove seedlings

are grown, inundation alone has little effect on photosynthetic rates or biomass production

(Pezeshki et al. 1990b, Hoppe-Speer et al. 2011). Inundation may sometimes lead to increases

in rates of leaf gas exchange over the short term, and often interacts with salinity over time to

reduce A_{net}, g_{sw} and growth rates (e.g., Cardona-Olarte et al. 2013). Thus, water levels and

flooding duration are key drivers controlling A_{net} in mangroves, and mangroves seem to

physiologically optimize to water levels. For instance, findings from a long-term greenhouse

inundation study by Farnsworth and Ellison (1996) exemplify how short-term responses of *R.*

mangle to inundation differ from longer-term responses. Over several years, high inundation

levels led to steady declines in A_{net} of up 25% for a given g_{sw} and decreases in growth rates.

Results of the high-water level (30-40 cm above soil surface) treatment were similar to those of

the low water level (10-15 cm) treatment, suggesting that *R. mangle* physiology is optimized at

inundation levels that reach just a few centimeters above the soil surface at high water level

(Ellison and Farnsworth 1997). Further research could assess the coupled effect of the depth

and duration of flooding with salinity in the FCE by characterizing photosynthetic rates across a

landscape-scale gradient that encompassed multiple sites and a broad range of salinity.

Although initial increases in *R. mangle* g_{sw} can result from short term inundation (Krauss

et al. 2006, Hoope-Speer et al. 2011), especially at low salinities (Pezeshki et al. 1990b),

several studies have linked stomatal closure to longer-term inundation (Kozlowski 1997, Ellison

and Farnsworth 1997). We measured depressed g_{sw} during the wet season and in mangrove

island edge habitats relative to centers; however, this was not attributable to water levels after

accounting for variation in seasonality and habitat, in that gsw increased with increasing water

levels during the wet season . Our measurements of g_{sw} were consistent with those reported in

other studies from across a range of inundation levels (Clough and Sim 1989, Lin and Sternberg

1992, Ellison and Farnsworth 1997, Krauss et al. 2006, Lugo et al. 2007, Barr et al. 2009),

supporting the understanding that *R. mangle* leaves limit g_{sw} in response to flooding to optimize

c_i for carbon gain without losing unnecessary amounts of water, but that g_{sw} can increase with

freshwater inputs, resulting in decrease in c_i, as A_{net} increases (Figure 3), likely because of

faster Calvin cycle reactions. What is interesting is that the scrub *R. mangle* leaves of TS/Ph-7

operate with low c_i concentrations (range = 220-260 µmol mol^{-1})., which suggests pervasive

inundation stress.

R. mangle nutrient use at TS/Ph-7

We found little variation (60-63%) in N resorption efficiencies for *R. mangle* leaves

across scrub mangrove island habitats. In contrast, higher overall efficiencies of P resorption

(73-79%) of leaf tissue were measured across habitats, with higher P resorption in mangrove

island edge habitats relative to centers, suggesting higher P availability in island centers. Our

findings are roughly comparable to N and P resorption efficiencies for *R. mangle* in the control

plots of scrub-dominated forests in Panama (~50%, and 80%, respectively; Lovelock et al.

2004). Resorption of nutrients from senescent leaves prior to leaf fall is a within-stand nutrient

recycling mechanism that may reduce nutrient losses via tidal export in coastal systems

(Vitousek 1982, Aerts and Chapin 2002). Like other tropical trees, mangroves exhibit several

physiological mechanisms that reduce nutrient losses via tidal exchange, including resorption of

N and P prior to leaf abscission, which can lead to increased availability of limiting nutrients and

ultimately change nutrient use and conservation patterns (Twilley et al. 1986, Alongi et al. 1992,

Feller et al. 2003a, 2003b).

Our findings suggest that *R. mangle* conserves P better than N in this P-limited

environment and indicate that canopy N and P resorption efficiency at TS/Ph-7 potentially

results from the differential partitioning of these nutrients among leaf stages and within the soil.

Mangrove species prioritize resorption of nutrients that are limited in the soil, and it has been

suggested that plants growing in nutrient-poor environments resorb a higher proportion of

nutrients, potentially decreasing nutrient loss by efficient nutrient recycling (Chapin and

Moilanen 1991). At our study site, low soil TP concentrations probably determine the higher

recycling efficiency of P relative to N. Indeed, soil TP concentrations in the upper 50 cm of soil

at TS/Ph-7 (0.06 ± 0.004 mg cm^{-3}) are three times lower than soils at the mouth of Shark River

estuary (SRS-6) dominated by fertile well-developed tall riverine mangroves, resulting in

extreme P limitation with average soil N:P ratios of 102 ± 6 (Castañeda-Moya et al. 2013).

Therefore, the non-homogeneous distribution of essential nutrients within mangrove habitats

creates distinct gradients and hot spots along the intertidal zone, influencing the efficiency of

internal nutrient recycling. This is supported by observations that nutrient resorption efficiencies

in mangroves vary with nutrient availability, e.g., via nutrient addition (Feller et al. 1999, Feller et

al. 2003b) or along natural fertility gradients (Medina et al. 2010). Such variation in nutrient

availability and resorption efficiencies within mangrove trees likely scales with variation in

productivity (e.g., litterfall) and carbon residence times (e.g., soil and biomass dynamics) of

mangrove forests.

Foliar δ^{15}N values integrate long-term processes of N sources because isotopic

fractionation against the heavier isotope (i.e., ^{15}N) occurs during N transformations and

interactions between biotic (e.g., mycorrhizal fungi, or bacteria) and biogeochemical (e.g.,

nitrification, denitrification) nutrient cycling processes (Garten 1993). In our study site, patterns

of δ^{15}N in *R. mangle* leaves differed drastically between mangrove habitats, with values around

-4 to -5‰ for mangrove edge habitats and between 0 and -1‰ for island centers, indicating

lower ^{15}N discrimination in island center habitats (Table 3, Figure S8A). These δ^{15}N values are

considerably more depleted than the *R. mangle* leaf δ^{15}N values reported for riverine

mangroves along Shark River estuary (He et al. 2020), where values were negatively correlated

with distance inland from the mouth of the estuary with more enriched leaves occurring near the

mouth of Shark River (4‰) relative to upstream (0.4‰) regions. Reported δ^{15}N values for *R.*

mangle leaves across different ecotypes in the neotropics range from 0 to -11‰, with more

negative values for scrub mangrove forests (-5 to -10‰) than for fringe mangroves (0-7‰; Reis

et al. 2017a). Similarly, Medina et al. (2010) showed that leaves from interior scrub mangrove

communities had more negative δ^{15}N values than tall fringe mangroves in eastern Puerto Rico (-

12‰ vs. 0‰, respectively). Those δ^{15}N values were more negative than those reported for

scrub *R. mangle* forests in Florida (Fry et al. 2000), Belize (McKee et al. 2002, Wooller et al.

2003, Fogel et al. 2008), or Brazil (Reis et al. 2017b).

Patterns of foliar δ^{15}N between mangrove ecotypes can be discerned using *in situ* leaf

nutrient content. For example, a direct relationship between ^{15}N discrimination and leaf N:P

ratios of *R. mangle* leaves previously reported for the six FCE mangrove sites, including our

study site, indicates that leaf N:P ratios accounted for 70% of the variability in ^{15}N discrimination

(Mancera-Pineda et al. 2009). Thus, foliar ^{15}N composition can reflect *in situ* N status and

differences in plant N use. Hypoxic conditions in the soil may inhibit denitrification and ammonia

volatilization, two processes that enrich the soil substrate in ^{15}N (Craine et al. 2015). Therefore,

the substrate should be less enriched at mangrove island edges relative to their centers,

because of interactions with the soil and the open water channels, which can alleviate hypoxia.

Thus, it appears that more negative $\delta^{15}N$ values in the edge habitats may be associated with

lower inorganic N (i.e., porewater ammonium) use by edge mangrove trees compared to those

in center island habitats (Fry et al. 2000). However, our results contrast slightly with those of

Mancera-Pineda et al. (2009), who reported mean $\delta^{15}N$ values of +3 from 65 mature leaves

collected in 2001 at our study site. We argue that differences in $\delta^{15}N$ values between the two

studies could be attributed to the location where leaves were collected during the 2009 study,

concluding that it is very likely that Mancera-Pineda et al. only collected leaves from the center

of mangrove islands, avoiding edge habitats. Taking this into consideration, mangrove island

centers potentially may have even more positive $\delta^{15}N$ signatures than we found, illustrating that

in the center of mangrove islands, N is taken up by roots in inorganic soluble forms (e.g.,

porewater ammonium, nitrate) readily available for plant uptake, and not biotically via root

symbionts. Another potential explanation of why $\delta^{15}N$ values were more negative at mangrove

island edges than in their centers is because lateral surface roots of *R. mangle* can extend into

open water where they associate with symbiotic biofilms (i.e., algae and aquatic bacteria) that

facilitate N acquisition from open water (Potts 1979). A significant source of isotopic

discrimination occurs during N transfer between belowground symbionts (e.g., mycorrhizal fungi

or bacteria) and plant roots during processes such as nitrification, denitrification, and ammonia

volatilization. The lighter isotope ^{14}N reacts faster than ^{15}N (i.e., is preferentially given to the

host plant by the symbiont), so that plant tissues are depleted, while substrates are enriched

(Högberg 1997, Robinson 2001). Indeed, at our study site, we have observed several long,

absorptive, fine lateral root systems that protrude from the edge of mangrove islands into the

open water ponds where biofilms colonize them.

Mangrove trees can potentially adapt to nutrient shortage or localized nutrient

deficiencies in the soil by patterns of nutrient use. This plant strategy may maximize the

efficiency of capturing limiting resources essential for growth (e.g., N, P) from soil or

surrounding open water areas in nutrient-poor environments such as Taylor River, as proposed

by the optimal plant allocation theory (Chapin et al. 1987, Gleeson and Tilman 1992). We

observed a slight decrease in foliar $\delta^{15}N$ during the wet season (Figure S8A, Table 3), as water

levels and porewater salinity increased, suggesting that N-acquisition by *R. mangle* via algal

biofilms may be slightly greater in the dry season than in the wet season. Lastly, highly

depleted (i.e., negative) N isotopic values in leaf tissues is characteristic of tropical wetlands

with P limitation because P-limitation increases N fractionation, especially in flooded wetlands

with limited P pedogenesis (McKee et al. 2002, Troxler 2007, Medina et al. 2008), as is the case

with the scrub *R. mangle* forest at TS/Ph-7 where the main source of P is brackish groundwater

discharge (Price et al 2006). Soil total P concentrations in top 10 cm of the peat soils at this site

have measured 0.055 (± 0.01) mg cm^{-3}, with atomic N:P ratios of roughly 72 (± 2) (Mancera-

Pineda et al 2009), which is considerably lower than soils of most mangrove forests globally, but

consistent with mangrove forests in karstic environments (Rovai et al 2018).

CONCLUSIONS

Habitat heterogeneity, resulting from micro-elevational differences in mangrove tree

locations on islands within the open water-mangrove island forest landscape, drives variation in

scrub *R. mangle* leaf physiological performance. Mangrove island edge habitats experience

greater and more-prolonged inundation than island centers in a seasonal dynamic, which leads

to reductions in g_{sw}, reduced A_{net}, and lower intrinsic *wue*. Conversely, mangrove island center

habitats are alleviated from inundation stress in the dry season, leading to increases in A_{net} and

g_{sw}. Interestingly, c_i levels increase with increasing water levels because inundation likely slows

not only g_{sw}, but the entire biochemical process of CO_2 assimilation, including mesophyll and

lower level (i.e., cell wall, plasma membrane, cytosol) conductance. Additionally, differences in

nutrient acquisition and use patterns among scrub *R. mangle* trees growing at island edges vs.

centers affect leaf nutrient status and photosynthetic potential.

The findings from this study indicate that it is the interaction of inundation stress with

mangrove island micro-elevational habitat in the flooded scrub mangroves of the southeastern

Florida Everglades that principally alters tree water and nutrient use dynamics, which appear to

cascade to affect leaf gas exchange rates through their effects on g_{sw}. Reductions in A_{net}

interact with the salinity of the water that inundates scrub *R. mangle* trees, in theory, because

g_{sw} rates are low and primarily respond to water loss from leaves rather than carbon gain. In

our field measurements, however, we find that prolonged inundation more than porewater

salinity drives reduction in A_{net} because the hydrological regime at Taylor River is characterized

by long hydroperiods and minor fluctuations in salinity throughout the year (Figure S2). At the

forest level, such physiological differences in scrub mangrove functioning with habitat and the

hydrological environment can help to parameterize demographic and carbon flux models to

forecast ecosystem trajectories in response to the impacts of SLR and saltwater intrusion in this

coastal region. This is particularly significant given the current freshwater restoration efforts in

the Everglades and the associated uncertainties of water management and projected climate

change scenarios on the spatial distribution of mangrove forests.

ACKNOWLEDGEMENTS

This research was supported by the Department of Interior – National Park Service (Cooperative Agreement #P16AC00032) and the Florida Coastal Everglades Long-Term Ecological Research (FCE-LTER) program funded by the National Science Foundation (Grant #DEB-1832229). We thank Mandy Kuhn, Austin Pezoldt, and Ralph Diaz-Hung for field assistance. We thank the Everglades National Park for granting research permits and the Florida Bay Interagency Science Center-Everglades National Park (FBISC-ENP) for logistic support during the study. We thank Dr. Zoran Bursac from the center for Statistical Consulting and Collaboration at FIU for statistical advice. We thank Prof. Ernesto Medina and two anonymous reviewers for constructive comments on an earlier version of this manuscript. This is contribution number XX from the Southeast Environmental Research Center in the Institute of Environment at Florida International University.

REFERENCES

Abiy AZ, Melesse AM, Abtew W, Whitman D (2019) Rainfall trend and variability in Southeast Florida: Implications for freshwater availability in the Everglades. PLoS One 14:e0212008.

Aerts R, Chapin FS. (2002) The mineral nutrition of wild plants revisited: a re-evaluation of processes and patterns. Adv Ecol Res 30:1-67.

Alongi DM (2008) Mangrove forests: Resilience, protection from tsunamis, and responses to global climate change. Estuarine Coastal Shelf Sci 76:1-13.

Alongi DM, Boto KG, Robertson AI (1992) Nitrogen and phosphorus cycles. In: Robertson AI, Alongi DM (eds) Tropical Mangrove Ecosystems. American Geophysical Union, Washington, DC.

Ball MC (2009) Photosynthesis in mangroves. Wetlands Australia 6:12-22.

Ball MC (1988) Ecophysiology of mangroves. Trees-Struct Funct 2:129-142.

Ball MC (1996) Comparative ecophysiology of mangrove forest and tropical lowland moist rainforest. In: Mulkey SS, Chazdon RL, Smith AP (eds) Tropical forest plant ecophysiology Springer, Boston, Massachusetts, pp 461-496.

Barr JG, Fuentes JD, Engel V, Zieman JC (2009) Physiological responses of red mangroves to the climate in the Florida Everglades. J Geophys Res Biogeosci 114:G02008.

Bates D, Mächler M, Bolker B, Walker S. (2015) Fitting linear fixed-effects models using lme4. J Stat Softw 67:1-48.

Bjorkman O, Demmig B, Andrews TJ (1988) Mangrove photosynthesis: response to high-irradiance stress. Funct Plant Bio 15:43-61.

Cardona-Olarte P, Krauss KW, Twilley RR (2013) Leaf gas exchange and nutrient use efficiency help explain the distribution of two Neotropical mangroves under contrasting flooding and salinity. Int J For Res 2013:1-10. doi:10.1155/2013/524625

Cardona-Olarte P, Twilley RR, Krauss KW, Rivera-Monroy VH (2006) Responses of neotropical mangrove seedlings grown in monoculture and mixed culture under treatments of hydroperiod and salinity. Hydrobiologia 569:325-341.

Castañeda-Moya E, Twilley RR, Rivera-Monroy VH, Marx BD, Coronado-Molina C, Ewe SML (2011) Patterns of root dynamics in mangrove forests along environmental gradients in the Florida Coastal Everglades, USA. Ecosystems 14:1178-1195.

Castañeda-Moya E, Twilley RR, Rivera-Monroy VH (2013) Allocation of biomass and net primary productivity of mangrove forests along environmental gradients in the Florida Coastal Everglades, USA. For Ecol Manage 307:226-241.

Chambers LG, Davis SE, Troxler T, Boyer JN, Downey-Wall A, Scinto LJ (2014) Biogeochemical effects of simulated sea level rise on carbon loss in an Everglades mangrove peat soil. Hydrobiologia 726:195-211.

Chapin FS, Bloom AJ, Field CB, Waring RH (1987) Plant responses to multiple environmental factors. BioScience 37:49-57.

Chapin FS, Moilanen L (1991) Nutritional controls over nitrogen and phosphorus resorption from Alaskan birch leaves. Ecology 72:709-715.

Cheeseman J, Herendeen L, Cheeseman A, Clough B (1997) Photosynthesis and photoprotection in mangroves under field conditions. Plant Cell Environ 20:579-588.

Cheeseman J, Lovelock C (2004) Photosynthetic characteristics of dwarf and fringe Rhizophora mangle L. in a Belizean mangrove. Plant Cell Environ 27:769-780.

Chen R, Twilley RR (1999) A simulation model of organic matter and nutrient accumulation in mangrove wetland soils. Biogeochemistry. 44:93-118.

Childers DL (2006) A synthesis of long-term research by the Florida Coastal Everglades LTER Program. Hydrobiologia. 569:531-544.

Cintron G, Lugo AE, Douglas D, Pool J, Morris G (1978) Mangroves of Arid Environments in
Puerto Rico and Adjacent Islands. Biotropica 10:110-121.
Clough B, Sim R (1989) Changes in gas exchange characteristics and water use efficiency of
mangroves in response to salinity and vapour pressure deficit. Oecologia 79:38-44.
Cornelissen JH, Lavorel S, Garnier E, Díaz S, Buchmann N, Gurvich DE, Reich PB, Ter Steege
H, Morgan HD, Van Der Heijden MG, Pausas JG (2003) A handbook of protocols for
standardised and easy measurement of plant functional traits worldwide. Aust J Bot
51:335-80.
Craine JM, Brooksheir ENJ, Cramer MD, Hasselquist NJ, Koba K, Marin-Spiotta E, Wang L
(2015) Ecological interpreations of nitrogen isotope ratios of terrestrial plans and soils
Plant Soil 396:1-26.
Duever MJ, Meeder JF, Meeder LC, McCollom JM (1994) The climate of South Florida and its
role in shaping the Everglades ecosystem. In: Davis SM,Ogden JC (eds) Everglades:
The Ecosystem and Its Restoration. St. Lucie Press, Delray Beach, Florida, pp 225-248.
Ellison AM, Farnsworth EJ (1997) Simulated sea level change alters anatomy, physiology,
growth, and reproduction of red mangrove (*Rhizophora mangle* L.). Oecologia 112:435-
446.
Ewe SM, Gaiser EE, Childers DL, Iwaniec D, Rivera-Monroy VH, Twilley RR (2006) Spatial and
temporal patterns of aboveground net primary productivity (ANPP) along two freshwater-
estuarine transects in the Florida Coastal Everglades. Hydrobiologia 569:459-474.
Ewe SM, Sternberg LSL, Childers DL. 2007. Seasonal plant water uptake patterns in the saline
southeast Everglades ecotone. Oecologia 152: 607-616.
Farnsworth EJ, Ellison AM (1996) Sun-shade adaptability of the red mangrove, *Rhizophora*
mangle (Rhizophoraceae): Changes through ontogeny at several levels of biological
organization. Amer J Bot 83:1131-1143.
Feller IC (1995) Effects of nutrient enrichment on growth and herbivory of dwarf red mangrove
(Rhizophora mangle). Ecol Monogr 65:477-505.
Feller IC, Whigham DF, O'Neil JP, McKee KL (1999) Effects of nutrient enrichement on within-
stand cycling in a mangrove forest. Ecology. 80:2193-2205.
Feller IC, McKee KL, Whigham DF, O'Neill JP (2003a) Nitrogen vs. phosphorus limitation across
an ecotonal gradient in a mangrove forest. Biogeochemistry 62:145-175.
Feller IC, Whigham DF, McKee KL, Lovelock CE (2003b) Nitrogen limitation of growth and
nutrient dynamics in a disturbed mangrove forest, Indian River Lagoon, Florida.
Oecologia 134:405-414.
Field CD (1984) Ions in mangroves. In: Teas HJ (ed) Physiology and management of
mangroves. Springer, Dordrecht, Netherlands, pp 43-48.
Field CD (1995) Impact of expected climate change on mangroves. Hydrobiologia 295:75-81.
Fogel M, Wooller M, Cheeseman J, Smallwood B, Roberts Q, Romero I, Jacobsen Meyers M
(2008) Unusually negative nitrogen isotopic compositions (δ^{15}N) of mangroves and
lichens in an oligotrophic, microbially-influenced ecosystem. Biogeosci Discuss 5:1693-
1704.
Fry B, Bern A, Ross M, Meeder J (2000) δ^{15}N studies of nitrogen use by the red mangrove,
Rhizophora mangle L. in south Florida. Estuarine Coastal Shelf Sci 50:291-296.
Garten CTJ (1993) Variation in foliar ^{15}N abundance and the availability of soil nitrogen on
Walker Branch watershed. Ecology 74:2098-2113.
Gleeson SK, Tilman D (1992) Plant allocation and the multiple limitation hypothesis. Am Nat
139:1322-1343.
Golley F, Odum HT, Wilson RF (1962) The structure and metabolism of a Puerto Rican red
mangrove forest in May. Ecology 43:9-19.

He B, Lai T, Fan H, Wang W, Zheng H (2007) Comparison of flooding-tolerance in four
mangrove species in a diurnal tidal zone in the Beibu Gulf. Estuarine Coastal Shelf Sci
74:254-262.
He D, Rivera-Monroy VH, Jaffé R, Zhao X (2020) Mangrove leaf species-specific isotopic
signatures along a salinity and phosphorus soil fertility gradients in a subtropical estuary.
Estuarine Coastal Shelf Sci 106768.
Högberg P (1997) ^{15}N natural abundance in soil-plant systems. New Phytol 137:179-203.
Hoppe-Speer SC, Adams JB, Rajkaran A, Bailey D (2011) The response of the red mangrove
Rhizophora mucronata Lam. to salinity and inundation in South Africa. Aquat Bot 95:71-
76.
Hughes CE, Binning P, Willgoose GR (1998) Characterisation of the hydrology of an estuarine
wetland. J Hydrol 211:34-49.
Koch MS, Snedaker SC (1997) Factors influencing *Rhizophora mangle* L. seedling development
in Everglades carbonate soils. Aquat Bot 59:87-98.
Kozlowski TT (1997) Responses of woody plants to flooding and salinity. Tree Physiol 17:490-
490.
Krauss KW, McKee KL, Lovelock CE, Cahoon DR, Saintilan N, Reef R, Chen L (2014) How
mangrove forests adjust to rising sea level. New Phytol 202:19-34.
Krauss KW, Twilley RR, Doyle TW, Gardiner ES (2006) Leaf gas exchange characteristics of
three neotropical mangrove species in response to varying hydroperiod. Tree Physiol
26:959-968.
Kuznetsova A, Brockhoff PB, Christensen RHB (2017) lmerTest Package: Tests in Linear Mixed
Effects Models. J Stat Softw 82:13.
Lambers H, Chapin FS, Pons TL (eds) (2008) Plant physiological ecology. Springer, New York,
New York.
Lamers LPM, Govers LL, Janssen ICJM, Geurts JJM, Van der Welle MEW, Van Katwijk MM,
Van del Heide T, Roelofs JGM, Smolders AJP (2013) Sulfide as a soil phytotoxin - a
review. Front Plant Sci 4:1-14.
Lawton JR, Todd ANN, Naidoo DK (1981) Preliminary inverstigations into the structure of the
roots of mangroves, *Avicennia marina* and *Brugeria gymnorrhiza*, in relation to ion
uptake. New Phytol 88:713-722.
Lin G, Sternberg LSL (1992) Comparative study of water uptake and photosynthetic gas
exchange between scrub and fringe red mangroves, *Rhizophora mangle* L. Oecologia
90:399-403.
Lopes D, Tognella M, Falqueto A, Soares M (2019) Salinity variation effects on photosynthetic
responses of the mangrove species *Rhizophora mangle* L. growing in natural habitats.
Photosynthetica 57:1142-1155.
Loveless CM (1959) A study of the vegetation in the Florida Everglades. Ecology 40:2-9.
Lovelock CE (2008) Soil respiration and belowground carbon allocation in mangrove forests.
Ecosystems 11:342-354.
Lovelock CE, Feller I (2003) Photosynthetic performance and resource utilization of two
mangrove species coexisting in hypersaline scrub forest. Oecologia 134:455-462.
Lovelock CE, Feller IC, McKee KL, Engelbrecht BM, Ball MC (2004) The effect of nutrient
enrichment on growth, photosynthesis and hydraulic conductance of dwarf mangroves in
Panama. Funct Ecol 18:25-33.
Lovelock CE, Atwood T, Baldock J, Duarte CM, Hickey S, Lavery P, Masque P, Macreadie PI,
Ricart AM, Serrano O, Steven A (2017) Assesing the risk of carbon dioxide emissions
from blue carbon ecosystems. Front Ecol Environ 15:257-265.
Lüdecke D (2018) sjPlot: Data visualization for statistics in social science. R package version 2.
https://cran.r-project.org/web/packages/sjPlot/

Lugo AE, Medina E, Cuevas E, Cintrón G, Nieves ENL, Novelli YS (2007) Ecophysiology of a
971   mangrove forest in Jobos Bay, Puerto Rico. Carib J Sci 43:200-219.
972   Lugo AE, Snedaker SC (1974) The ecology of mangroves. Ann Rev Ecol System 5:39-64.
973   Mancera-Pineda JE, Twilley RR, Rivera-Monroy VH (2009) Carbon (δ^{13}C) and nitrogen (δ^{15}N)
974   isotopic discrimination in mangroves in Florida coastal Everglades as a function of
975   enviromnetal stress. Contrib Mar Sci 38:109-129.
976   Marshall JD, Brooks JR, Lajtha K (2007) Stable isotopes in ecology and environmental science.
977   In: Michener R, Lajtha K (eds) Sources of variation in the stable isotopic composition of
978   plants .Blackwell, Singapore, pp 22-60.
979   McKee KL (1993) Soil physicochemical patterns and mangrove species distribution-reciprocal
980   effects? J Ecol 81:477-487.
981   McKee KL (2011) Biophysical controls on accretion and elevation change in Caribbean
982   mangrove ecosystems. Estuarine Coastal Shelf Sci 91:475-483.
983   McKee KL, Cahoon DR, Feller IC (2007) Caribbean mangroves adjust to rising sea level
984   through biotic controls on change in soil elevation. Global Ecol Biogeogr 16:545-556.
985   McKee KL, Feller IC, Popp M, Wanek W (2002) Mangrove isotopic (δ^{15}N and δ^{13}C) fractionation
986   across a nitrogen vs. phosphorus limitation gradient. Ecology 83:1065-1075.
987   Mcleod E, Chmura GL, Bouillon S, Salm R, Bjork M, Duarte CM, Lovelock CE, Schlesinger WH,
988   Silliman BR (2011) A blueprint for blue carbon: Toward an improved understanding of
989   the role of vegetated coastal habitats in sequestering CO_2. Front Ecol Environ 9:552-560.
990   Medina E (1999) Mangrove physiology: the challenge of salt, heat, and light stress under
991   recurrent flooding. In: Yañez-Arancibia A, Lara-Domínguez L (eds) Ecosistemas de
992   manglar en América tropical. Instituto de Ecología A.C. México, UICN/ORMA, Costa
993   Rica, NOAA/NMFS Silver Spring, MD USA. pp 109-126.
994   Medina E, Cuevas E, Lugo AE (2010) Nutrient relations of dwarf *Rhizophora mangle* L.
995   mangroves on peat in eastern Puerto Rico. Plant ecolog 207:13-24.
996   Medina E, Francisco M (1997) Osmolality and δ^{13}C of leaf tissues of mangrove species from
997   environments of contrasting rainfall and salinity. Estuarine Coastal Shelf Sci 45:337-344.
998   Medina E, Francisco M, Quilice A (2008) Isotopic signatures and nutrient relations of plants
999   inhabiting brackish wetlands in the northeastern coastal plain of Venezuela. Wetlands
1000   Ecol Manage 16:51.
1001   Mendelssohn I, McKee K (2000) Saltmarshes and mangroves. In: Barbour MG, Billings WD
1002   (eds) North American terrestrial vegetation. Cambridge University Press, New York, NY,
1003   pp 501-536.
1004   Michot B, Meselhe EA, Rivera-Monroy VH, Coronado-Molina C, Twilley RR (2011) A tidal creek
1005   water budget: estimation of groundwater discharge and overland flow using hydrologic
1006   modeling in the southern Everglades. Estuarine Coastal Shelf Sci 93:438-448.
1007   Murdiyarso D, Purbopuspito J, Kauffman JB, Warren MW, Sasmito SD, Donato DC, Manuri S,
1008   Krisnawati H, Taberima S, Kurnianto S (2015) The potential of Indonesian mangrove
1009   forests for global climate change mitigation. Nat Clim Change 5:1089-1092.
1010   Naidoo G (1983) Effects of flooding on leaf water potential and stomatal resistance in *Bruguiera*
1011   gymnorrhiza (L.) LAM. New Phytol 93:369-373.
1012   Neubauer SC, Franklin RB, Berrier DJ (2013) Saltwater intrusion into tidal freshwater marshes
1013   alters the biogeochemical processing of organic carbon. Biogeosciences Discuss
1014   10:10685-10720.
1015   Nickerson NH, Thibodeau FR (1985) Association between porewater sulfide concentrations and
1016   distribution of mangroves. Biogeochemistry 1:183-192.
1017   O'Leary MH (1988) Carbon isotopes in photosynthesis. BioScience. 38:328-336.
1018   Parida AK, Jha B (2010) Salt tolerance mechanisms in mangroves: a review. Trees-Struct Funct
1019   24:199-217.

Pezeshki S, DeLaune R, Patrick Jr W (1990a) Flooding and saltwater intrusion: Potential effects on survival and productivity of wetland forests along the U.S. Gulf Coast. For Ecol Manage 33:287-301.

Pezeshki S, DeLaune R, Patrick Jr W (1990b) Differential response of selected mangroves to soil flooding and salinity: gas exchange and biomass partitioning. Can J For Res 20:869-874.

Pezeshki SR, DeLaune RD (2012) Soil oxidation-reduction in wetlands and its impact on plant functioning. Biology 1:196-221.

Potts M (1979) Nitrogen fixation (acetylene reduction) associated with communities of heterocystous and non-heterocystous blue-green algae in mangrove forests of Sinai. Oecologia 39:359-373.

Pugnaire FI, Chapin FS (1993) Controls over nutrient resorption from leaves of evergreen Mediterranean species. Ecology 4:124-129.

Price RM, PK Swart, Fourqurean JW (2006) Coastal groundwater discharge - an additional source of phosphorus for the oligotrophic wetlands of the Everglades. Hydrobiologia 569: 23-36.

R Core Team (2018) R: a language and environment for statiscal computing. R foundation for staistical computing. Vienna, Austria. www.r-project.org

Reef R, Lovelock CE (2015) Regulation of water balance in mangroves. Ann Bot 115:385-395.

Reis CRG, Nardoto GB, Oliveira RS (2017a) Global overview on nitrogen dynamics in mangroves and consequences of increasing nitrogen availability for these systems. Plant Soil 410:1-19.

Reis CRG, Nardoto GB, Rochelle ALC, Vieira SA, Oliveira RS (2017b) Nitrogen dynamics in subtropical fringe and basin mangrove forests inferred from stable isotopes. Oecologia 183:841-848.

Robinson D (2001) δ^{15}N as an integrator of the nitrogen cycle. Trends Ecol Evol 16:153-162.

R Core Team (2018) R: a language and environment for statiscal computing. R foundation for staistical computing. Vienna, Austria. www.r-project.org

Ross MS, Meeder JF, Sah JP, Ruiz PL, Telesnicki GJ (2000) The southeast saline Everglades revisited: 50 years of coastal vegetation change. J Veg Sci 11:101-112.

Rovai AS, Twilley RR, Castañeda-Moya E, Riul P, Cifuentes-Lara M, Manrow-Villalobos M, Horta PA, Simonassi JC, Fonseca AL, Pagliosa PR (2018) Global controls of carbon storage in mangrove soils. Nat Clim Change 8:534-538.

Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image analysis. Nat Methods 9:671-675.

Scholander P (1968) How mangroves desalinate seawater. Physiol Plant 21:251-261.

Scholander P, Hammel H, Hemmingsen E, Garey W (1962) Salt balance in mangroves. Plant Physiol 37:722.

Sobrado M (2000) Relation of water transport to leaf gas exchange properties in three mangrove species. Tree-Struct Funct 14:258-262.

Sutula M, Day JW, Cable JE, Rudnick D (2001) Hydrological and nutrient budgets of freshwater and estuarine wetlands of Taylor Slough in southern Everglades, Florida (U.S.A.). Biogeochemistry 56:287-310.

Tully K, Gedan K, Epanchin-Niell R, Strong A, Bernhardt ES, BenDor T, Mitchell M, Kominoski J, Jordan TE, Neubauer SC, Weston NB (2019) The invisible flood: The chemistry, ecology, and social implications of coastal saltwater intrusion. BioScience 69:368-378.

Tomlinson PB (2016) The botany of mangroves. Cambridge University Press, Cambridge, Massachusetts.

Troxler TG (2007) Patterns of phosphorus, nitrogen and δ^{15}N along a peat development gradient in a coastal mire, Panama. J Trop Ecology 23:683-691.

Twilley RR, Castañeda-Moya E, Rivera-Monroy VH, Rovai A (2017) Productivity and carbon
dynamics in mangrove wetlands. In: Rivera-Monroy VH, Lee SY, Kristensen E, Twilley
RR (eds) Mangrove ecosystems: A global biogeographic perspective. Springer
International, pp 113-162.
Twilley RR, Lugo AE, Patterson-Zucca C (1986) Litter production and turnover in basin
mangrove forests in southwest Florida. Ecology 67:670-683.
Twilley RR, Rivera-Monroy V (2005) Developing performance measures of mangrove wetlands
using simulation models of hydrology, nutrient biogeochemistry, and community
dynamics. J Coast Res 40:79-93.
Twilley RR, Rivera-Monroy V (2009) Ecogeomorphic models of nutrient biogeochemistry for
mangrove wetlands. In: Coastal Wetlands: An integrated ecosystem approach (eds)
Perillo GME, Wolanski E, Cahoon DR and Brinson MM. Elsevier, pp 641-683.
Twilley RR, Rivera-Monroy V, Rovai AS, Castañeda-Moya E, Davis SE, III. (2019) Mangrove
biogeochemistry at local to global scales using ecogeomorphic approaches. In: Perillo
GME, Wolanski E, Cahoon DR, Hopkinson CS (eds) Coastal Wetlands: An Integrated
Ecosystem Approach.Elsevier, pp 717-785.
Twilley RR, Rivera-Monroy VH, Chen R, Botero L (1998) Adapting and ecological mangrove
model to simulate trajectories in restoration ecology. Mar. Poll. Bull. 37:404-419.
Uncles R, Gong W-K, Ong J-E (1992) Intratidal fluctuations in stratification within a mangrove
estuary. In: Martens E, Jaccarini V (eds) The Ecology of Mangrove and Related
Ecosystems. Springer, pp 163-171.
Vitousek P (1982) Nutrient cycling and nutrient use efficiency. Am Nat 119:553-572.
Wanless H (1998) Mangroves, hurricanes, and sea level rise. South Florida Study Group, The
Conservancy, Naples, Florida
Werner A, Stelzer R (1990) Physiological responses of the mangrove *Rhizophora mangle* grown
in the absence and presence of NaCl. Plant Cell Environ 13:243-255.
Wolanski E (1992) Hydrodynamics of mangrove swamps and their coastal waters.
Hydrobiologia 247:141-161.
Wooller M, Smallwood B, Scharler U, Jacobson M, Fogel M (2003) A taphonomic study of δ13C
and δ15N values in Rhizophora mangle leaves for a multi-proxy approach to mangrove
palaeoecology. Org Geochem 34:1259-1275.
Yu M, Rivera-Ocasio E, Heartsill-Scalley T, Davila-Casanova D, Rios-López N, Gao Q. (2019)
Landscape-level consequences of rising sea-level on coastal wetlands: saltwater
intrusion drives displacement and mortality in the twenty-first century. Wetlands 39:1343-
1355.
1105

 TABLES

Table 1. Mean (± 1 SE) soil surface elevation, bedrock elevation, and soil depth, for open water, mangrove island edge and center habitats in the fringe and interior scrub mangrove areas at TS/Ph-7 in southeastern Florida Coastal Everglades. Elevation measurements are referenced to the North American Vertical Datum 1988 (NAVD88). Letters denote statistically different groupings via Tukey HSD post-hoc test ($p < .05$).

Mangrove Location	Habitat	Soil Surface Elevation (NAVD88, m)	Bedrock Elevation (NAVD88, m)	Soil depth (m)
Fringe	Open water	-0.63 ± 0.05^C	-1.84 ± 0.05^A	1.22 ± 0.05^C
	Island edge	-0.41 ± 0.03^B	-1.83 ± 0.02^A	1.43 ± 0.05^{BC}
	Island center	-0.14 ± 0.02^A	-1.86 ± 0.06^A	1.72 ± 0.05^A
Interior	Open water	-0.84 ± 0.02^D	-2.01 ± 0.15^A	1.18 ± 0.14^C
	Island edge	-0.49 ± 0.03^B	-1.78 ± 0.04^A	1.39 ± 0.04^C
	Island center	-0.15 ± 0.02^A	-1.77 ± 0.03^A	1.62 ± 0.03^{AB}

1113 **Table 2.** Seasonal variation in water levels, surface water and porewater salinity measured in
1114 mangrove island habitats at scrub *R. mangle* dominated mangroves at TS/Ph-7 in southeastern
1115 Florida Everglades. Means (± 1 SE) with different letters within each column denoting
1116 significant differences among groups (Tukey HSD post hoc, $p < .05$).

1117

Season	Habitat	Water level (cm)	Surface water salinity (ppt)	Porewater salinity (ppt)
Dry	Edge	33.5 ± 1.9^A	16.11 ± 1.16^A	24.22 ± 0.44^A
	Center	10.1 ± 1.9^B	21.14 ± 1.51^B	20.83 ± 0.44^B
Wet	Edge	40.2 ± 1.9^C	15.31 ± 1.16^A	26.00 ± 0.44^C
	Center	15.5 ± 1.9^B	14.41 ± 1.47^A	22.22 ± 0.44^D

Table 3. Leaf functional traits, carbon and nutrient contents and N:P ratios, nitrogen and
phosphorus resorption efficiencies and bulk isotopic signatures, and intrinsic intracellular CO_2
concentrations (c_i) and intrinsic water use-efficiency (*WUE*) (calculated from ^{13}C fractionation)
for scrub *R. mangle* leaves collected at mangrove island habitats at TS/Ph-7 during the dry and
wet seasons of 2019. Means (± 1 SE) with different letters across each row denoting
significantly different groups (Tukey HSD test, $p < .05$).

Leaf Trait	Dry season		Wet season	
	Edge	Center	Edge	Center
Leaf dry mass (g)	0.64 ± 0.02^A	0.65 ± 0.02^A	0.69 ± 0.02^A	0.70 ± 0.02^A
Leaf area (cm^2)	24.9 ± 0.7^A	25.5 ± 0.7^A	26.0 ± 1.4^A	27.3 ± 0.9^A
SLA ($cm^2 g^{-1}$)	28.97 ± 0.66^A	39.79 ± 0.68^A	37.33 ± 0.40^A	40.04 ± 1.71^A
LWC (%)	65.7 ± 0.4^A	65.4 ± 0.5^A	63.6 ± 0.5^A	64.0 ± 0.8^A
Total C ($mg g^{-1}$)	450.9 ± 5.7^A	441.4 ± 1.0^A	428.7 ± 17.7^A	444.0 ± 1.0^A
Total N ($mg g^{-1}$)	9.8 ± 0.2^{AB}	10.2 ± 0.3^A	8.4 ± 0.5^B	9.0 ± 0.3^{AB}
Total P ($mg g^{-1}$)	0.50 ± 0.01^{AB}	0.55 ± 0.04^A	0.42 ± 0.01^B	0.46 ± 0.02^{AB}
Leaf N_{area} ($mg cm^{-2}$)	0.258 ± 0.012	0.264 ± 0.019	0.263 ± 0.035	0.306 ± 0.020
Leaf P_{area} ($mg cm^{-2}$)	0.013 ± 0.001	0.014 ± 0.002	0.013 ± 0.001	0.016 ± 0.001
Atomic N:P	41.7 ± 1.9	40.0 ± 0.1	43.5 ± 3.7	42.5 ± 2.1
N resorption (%)	60.0 ± 0.4	62.8 ± 0.5	60.8 ± 2.8	62.9 ± 1.5
P resorption (%)	78.6 ± 0.2	74.3 ± 3.2	75.5 ± 0.4	73.2 ± 6.2
$\delta^{13}C$ (‰)	-25.5 ± 0.1^{AB}	-25.1 ± 0.1^A	-25.8 ± 0.1^B	-25.9 ± 0.2^{AB}
c_i ($\mu mol mol^{-1}$)	228.1 ± 2.2^{AB}	219.9 ± 1.8^B	233.3 ± 1.1^A	235.3 ± 3.6^A
WUE ($mmol mol^{-1}$)	0.1124 ± 0.0014^{AB}	0.1175 ± 0.0012^B	0.1092 ± 0.0007^A	0.1080 ± 0.0022^A
$\delta^{15}N$ (‰)	-5.3 ± 0.5^B	-0.4 ± 0.4^A	-4.2 ± 1.0^{AB}	-0.8 ± 1.5^A

1124

 FIGURES

1126

Figure 1. A) Photograph of TS/Ph-7, showing scrub *R. mangle* tree islands which characterize
the study site. Mangrove canopy heights are approximately 1.5-2 m tall, facilitating canopy
measurements of leaf physiology. Boardwalk (1.3 m height) is pictured for reference. **B)** Aerial
view (Google Earth) of mangrove islands measured for this study within TS/Ph-7, near the
mouth of the Taylor River in southeastern Florida Coastal Everglades, USA. The inset shows
the location of TS/Ph-7 within the boundary of Everglades National Park. Colors indicate scrub
mangroves and fringe and interior zones relative to the shoreline (i.e., Taylor River). Symbols
denoted paired higher-elevation, center and lower-elevation, edge habitats for each mangrove
island (squares and triangles, respectively).

1136

1139   **Figure 2.** Predicted marginal mean (±95% confidence intervals) values of photosynthesis (A_{net}),
1140   stomatal conductance (g_{sw}), the concentration of intracellular CO_2 (c_i), and instantaneous water
1141   use efficiency (*wue*) by mangrove island habitat and season. The dry season is from November
1142   to April, and the wet season is from May to October. See supplemental material for complete
1143   model summaries.

1144

Figure 3. The effect of water level on leaf photosynthesis (A_{net}) and stomatal conductance (g_{sw}), the concentration of intracellular CO_2 (c_i), and instantaneous water use efficiency (*wue*) by season. Lines are habitat-specific predicted mean marginal mean values (± 95% confidence intervals) from linear mixed-effects models.

1149

1150

1151 **Figure 4.** The effect of soil porewater salinity on leaf photosynthesis (A_{net}) and stomatal
1152 conductance (g_{sw}) by season. Porewater salinity was not included in the best-fit models for c_i or
1153 *wue*. Lines are predicted mean marginal effects from linear mixed-effects models ± 95%
1154 confidence intervals (colored by island habitat).