TITLE PAGE

Title: Water levels primarily drive variation in photosynthesis and nutrient use of scrub

Red Mangroves in the southeastern Florida Everglades

Authors: J. Aaron Hogan[1*], Edward Castañeda-Moya[2], Lukas Lamb-Wotton[1], Tiffany

Troxler[2,3], Christopher Baraloto[1]

Author Affiliations: [1] Department of Biological Sciences, Florida International University,

Miami, FL USA; [2] Institute of Environment, Florida International University, Miami, FL

USA, [3] Department of Earth and Environment, Florida International University, Miami,

FL, USA.

* Corresponding Author: J. Aaron Hogan, email: jhogan@fiu.edu

ORCiDs:

J. Aaron Hogan: 0000-0001-9806-3074; Edward Castañeda-Moya: 0000-0001-7759-4351;

Lukas Lamb-Wotton: 0000-0002-0712-9005, Troxler: 0000-0007-7100-7742, Baraloto: 0000-

0001-7322-8581

Authorship statement: The conception and design of the study and the acquisition of data

were done by JAH, EC-M, and LL-W. JAH and EC-M performed data analyses. Interpretation

of the data was carried out by JAH, EC-M, LL-W, TT, and CB. JAH drafted the manuscript. All

authors have contributed to the final version of the manuscript and have approved its

submission to the journal.

Data deposition: The data collected and used in this study have been archived and are

available through the Environmental Data Initiative data repository:

Hogan, J.A., E. Castaneda, and L. Lamb-Wotton. 2021. FCE LTER Taylor Slough/Panhandle-7

Site Scrub Red Mangrove (*Rhizophora mangle*) Leaf Gas Exchange Data, Florida, USA from

January-December 2019 ver 1. Environmental Data Initiative.

https://doi.org/10.6073/pasta/27f6332609eb1ef6d398c7855855f2e3 (Accessed 2021-01-26).

Note: this data archive also contains R code for running the linear mixed-effects models using

the archived datasets.

Running head: mangrove photosynthesis varies with island-habitat

ABSTRACT

We investigated how mangrove-island micro-elevation (i.e., habitat: center vs. edge) affects tree physiology in a scrub mangrove forest of the southeastern Everglades. We measured leaf gas exchange rates of scrub *Rhizophora mangle* trees monthly during 2019, hypothesizing that CO_2 assimilation (A_{net}) and stomatal conductance (g_{sw}) would decline with increasing water levels and salinity, expecting more-considerable differences at mangrove-island edges than centers, where physiological stress is greatest. Water levels varied between 0 and 60 cm from the soil surface, rising during the wet season (May-October) relative to the dry season (November-April). Porewater salinity ranged from 15 to 30 ppt, being higher at mangrove-island edges than centers. A_{net} maximized at 15.1 µmol m^{-2} s$^{-1,}$ and g_{sw} was typically <0.2 mol m^{-2} s^{-1}, both of which were greater in the dry than the wet season and greater at island centers than edges, with seasonal variability being roughly equal to variation between habitats. After accounting for season and habitat, water level positively affected A_{net} in both seasons but did not affect g_{sw}. Our findings suggest that inundation stress (i.e., water level) is the primary driver of variation in leaf gas exchange rates of scrub mangroves in the Florida Everglades, while also constraining A_{net} more than g_{sw}. The interaction between inundation stress due to permanent flooding and habitat varies with season as physiological stress is alleviated at higher-elevation mangrove-island center habitats during the dry season. Freshwater inflows during the wet season, increase water levels and inundation stress at higher-elevation mangrove-island centers, but also potentially alleviate salt and sulfide stress in soils. Thus, habitat heterogeneity leads to differences in nutrient and water acquisition and use between trees growing in island centers versus edges, creating distinct physiological controls on photosynthesis, which likely affect carbon flux dynamics of scrub mangroves in the Everglades.

Keywords: Scrub mangroves, Florida Coastal Everglades, photosynthesis, porewater salinity, water levels, inundation, *Rhizophora mangle*.

INTRODUCTION

Global climate change is affecting coastal mangrove ecosystems unprecedentedly, principally through increased flooding and saltwater intrusion (Pezeshki et al. 1990a, Yu et al. 2020). Increases in flooding severity and salinity due to sea-level rise (SLR) have the potential to push ecosystems to degraded alternative stable states, where biogeochemical cycles (e.g., carbon sequestration and storage potential) are impaired (Neubauer et al. 2013, Tully et al. 2019, Yu et al. 2020). Mangrove wetlands are particularly susceptible to SLR because of their position between terrestrial and marine ecosystems (Field 1995, Ellison and Farnsworth 1997). Mangrove species have developed considerable variation in crucial life-history traits, such as rates of photosynthesis, water- and nutrient-use efficiencies, growth rates, and biomass allocation ratios in response to the interactions among resources (e.g., light and nutrients), regulators (e.g., salinity, sulfides), and inundation – Twilley and Rivera-Monroy 2005, Alongi 2008, Twilley and Rivera-Monroy 2009, Castañeda-Moya et al. 2013). Due to such physiological flexibility and the significant carbon sequestration and storage capacity of mangroves across a variety of geomorphic settings (e.g., karstic vs. deltaic; Mcleod et al. 2011, Murdiyarso et al. 2015, Lovelock et al. 2017, Rovai et al. 2018), there is an increasing need to strengthen our understanding of the effects of SLR and saltwater intrusion on mangrove tree physiology to assess trajectories of ecosystem structure and function in response to global change drivers.

Scrub mangrove forests, dominated by *Rhizophora mangle,* are typical in Caribbean karstic environments (Cintron et al. 1978, Lugo and Snedaker 1974). The stunted physiognomy (i.e., reduced growth and development) of scrub mangroves results from severe nutrient (e.g., phosphorus, P) limitation, prolonged or permanent inundation with little tidal influence, and seasonal water stress (Feller 1995, Koch and Snedaker 1997, Cheeseman and Lovelock 2004, Medina et al. 2010). Scrub mangrove forests develop distinct landscape patterning, forming mangrove-island clusters with higher-elevations than their surrounding shallow open-water ponds and channels (Figure 1A). Soil elevation gradients result from differences in root biomass stocks and production, leaf litter accretion, and wood deposition (McKee et al. 2007, McKee 2011, Krauss et al. 2014). For example, in scrub mangrove-islands of the southeastern Florida Everglades, island center habitats have 66% more root biomass and 52% more root production than island edges (Castañeda-Moya et al. 2011), which leads to spatial differences in soil elevation among island habitats. These differences in soil elevation interact with environmental gradients (e.g., hydroperiod, salinity) along the intertidal zone in complex ways to

affect mangrove physiology (e.g., rates of net CO_2 assimilation – A_{net}, growth rates, or sap flux)

at variable scales (Medina and Francisco 1997, Twilley et al. 1998, Medina et al. 2010, Twilley

et al. 2017).

Hydrological dynamics (e.g., depth and duration of inundation) can cause mangrove

physiological stress, which cascades to affect carbon cycle dynamics and other biogeochemical

processes across spatial and temporal scales (Medina 1999, Castañeda-Moya et al. 2013,

Twilley et al. 2017, 2019). Although mangrove species can tolerate flooded conditions, they are

still susceptible to damage if plants become entirely submerged for days to weeks (Wanless

1998, Mendelssohn and McKee 2000, McKee 2011). Inundation stress typically decreases

rates of leaf gas exchange (e.g., A_{net}, transpiration) and tree growth in mangroves (He et al.

2007, Cardona-Olarte et al. 2013). For example, greenhouse studies have revealed a 20%

reduction in maximum A_{net} when mangrove seedlings and saplings were subjected to short-term

intermittent seawater flooding (6 to 22 days, Krauss et al. 2006). Mangrove leaf gas exchange

is further affected by how seawater flooding interacts with fresh water and nutrient inputs

(Wolanski 1992). For instance, a reduction in stomatal conductance (g_{sw}) and leaf water

potential in *Bruguiera gymnorrhiza* seedlings occurred when exposed to prolonged flooding for

up to 80 days with 33% seawater compared to the control plants; however, seedlings flooded

with fresh water for 80 days showed an increase in both parameters (Naidoo 1983). In contrast,

seedlings of *Avicennia germinans* and *Laguncularia racemosa* exposed to permanent flooding

with 23% seawater showed no change in g_{sw}, A_{net}, or intrinsic water use efficiency (*wue*) but had

reduced leaf area (Krauss et al. 2006). Hydrologic conditions can further negatively influence

mangrove physiology through the interaction with soil phytotoxins (i.e., sulfides), produced as

by-products of low oxygen availability and soil redox conditions due to permanent flooding,

which can potentially depress water and nutrient uptake and affect rates of leaf gas exchange

(Nickerson and Thibodeau 1985, McKee 1993, Ball 1996, Pezeshki and DeLaune 2012, Lamers

et al. 2013). Regarding permanently inundated scrub mangroves, such as those in the

southeastern Florida Everglades, how seasonal dynamics interact with inundation levels to

influence leaf and forest carbon uptake dynamics is not entirely understood.

Mangroves are highly adapted to tolerate salt stress, yet salinity has the most significant

impact on forest productivity, tree growth rates, and rates of leaf gas exchange. The adverse

effects of increasing salinity are most evident along steep salinity gradients (i.e., those >30 ppt),

particularly in dry environments (Lugo and Snedaker 1974, Cintron et al. 1978, Medina and

Francisco 1997, Reef and Lovelock 2015). Salt stress variably affects mangrove tree

physiology, depending on species-specific salt tolerance levels and mechanisms for processing

salt (Parida and Jha 2010, Reef and Lovelock 2015). For example, *R. mangle* naturally inhabits

Neotropical environments with salinities from near zero to around 35 ppt but may also be found

in dry coastal environments with salinities up to 50-60 ppt (Cintron et al. 1978, Cardona-Olarte

et al. 2006). All mangroves can exclude salt through the roots; however, *R. mangle* is a highly

efficient salt excluder because its roots essentially prevent salt from entering the plant.

Additionally, *R. mangle* lacks the excretory glands that other mangrove species (e.g., *L.*

racemosa) use to excrete salt once it has entered the plant. As such, the xylem of *R. mangle* is

100 times less saline than seawater (Scholander et al. 1962, Scholander 1968, Medina and

Francisco 1997, Tomlinson 2016) because of the Casparian strip (Lawton et al. 1981) and

ultrafiltration by cell membranes in the thick aerenchyma and cortical layers of its root tissues

(Field 1984, Werner and Stelzer 1990). However, some salt still enters the plant through the

roots, which has a deleterious effect on the physiology of *Rhizophora* trees, causing decreases

in growth and A_{net} rates, and water and nutrient use efficiencies (Ball 1988, Clough and Sim

1989, Lugo et al. 2007, Medina et al. 2010, Cardona-Olarte et al. 2013).

Mangrove A_{net} varies widely with the environment (e.g., water and salinity levels) and

nutrient availability. A_{net} for *R. mangle* maximizes around 20 µmol m^{-2} s^{-1} (Golley et al. 1962,

Bjorkman et al. 1988, Lin and Sternberg 1992, Lovelock and Feller 2003, Lugo et al. 2007, Ball

2009); however, A_{net} for scrub mangroves is lower, generally ranging from <5 µmol m^{-2} s^{-1}

(Golley et al. 1962, Cheeseman et al. 1997, Cheeseman and Lovelock 2004) to roughly 13 µmol

m^{-2} s^{-1} (Lugo et al. 2007, Barr et al. 2009). A field study from Jobos Bay in southern Puerto Rico

demonstrated a significant decrease in *R. mangle* A_{net} (from 12.7 to 7.9 µmol m^{-2} s^{-1}) and g_{sw}

(from 0.28 to 0.19 mol m^{-2} s^{-1}) when comparing fringe habitats at 35 ppt salinity to inland salt flat

habitats at 80 ppt (Lugo et al. 2007). Reductions in A_{net} and g_{sw} were accompanied by changes

in leaf morphology (i.e., smaller specific leaf area, SLA), reduced nutrient-use efficiency, and

increased nutrient resorption, demonstrating how environmental effects on mangrove

physiology can have consequences for within plant nutrient dynamics, and thus ecosystem

functioning. Therefore, increasing salinity decreases A_{net} and g_{sw} and increases intrinsic water

use efficiency (*wue*, defined as A_{net}/g_{sw}) in mangroves, with *Rhizophora* species exemplifying

these trends (Ball 2009, Clough and Sim 1989). Moreover, the high salt tolerance of mangrove

species leads to strong stomatal control, which creates dynamics between A_{net} and water use

which depend on relative reduction in transpiration rates versus the degree to which leaves are

biochemically limited to fix carbon (e.g., via RUBISCO carboxylation efficiency versus RUBP

regeneration) at low stomatal conductance (Sobrado 2000, Lovelock and Feller 2003, Ball 2009).

For instance, *R. mangle* has succulent leaves with lower *wue* than more salt-tolerant species
(i.e., *A. germinans* or *L. racemosa*); however, *R. mangle* has greater water transport efficiency
in stems than more salt-tolerant species (Sobrado 2000). Thus, when considering the effects of
salinity on leaf gas exchange rates, plant water use must be considered in concert because
both A_{net} and g_{sw} decline similarly with increasing salinity, effectively creating co-limitation of
photosynthesis at moderate to high salinities (Ball 2009).

In mangrove wetlands of the Florida Everglades, variation in environmental gradients,
including hydroperiod (e.g., duration of inundation) and soil P availability, control mangrove
forest structure and function (e.g., biomass and litterfall production) across the coastal
landscape (Chen and Twilley 1999; Castañeda-Moya et al. 2011, 2013). However, how the
interaction between water level dynamics and salinity affects mangrove leaf gas exchange rates
in situ is not entirely understood. Experimental evidence using *R. mangle* seedlings from south
Florida showed that inundation created a greater degree of physiological stress than salinity
levels; however, salinity accelerated the adverse effects of inundation stress on leaf function
over time (Cardona-Olarte et al. 2013). In contrast, other studies have reported no apparent
effect of water levels or flooding duration on rates of mangrove gas exchange, although
inundation duration decreased variability in leaf gas exchange measurements (Hoppe-Speer et
al. 2011). Using Florida mangroves, Krauss et al. (2006) found that short-term intermittent
flooding decreased rates of leaf gas exchange relative to unflooded or permanently flooded
greenhouse-grown seedlings, but that for *in situ* established *R. mangle* saplings growing along a
natural tidal inundation gradient along Shark River in the southwestern Everglades, flooding led
to increases in A_{net} and *wue*. Permanent flooding leads to decreases in A_{net} and g_{sw} rates in
most wetland plants (Kozlowski 1997); however, how inundation dynamics interact with salinity
along the intertidal zone to influence mangrove physiology at different spatial and temporal
scales in south Florida mangroves remains largely unknown. Further, global change-driven
SLR and saltwater intrusion in South Florida coupled to large-scale freshwater diversion have
accelerated mangrove encroachment into inland freshwater wetlands over the past 60 years
(Ross et al. 2000). As SLR continues, it is imperative to quantify the relative effects of
inundation and salinity on mangrove physiology and subsequent ecosystem functioning (e.g.,
carbon flux) in the region.

Here, we present a comprehensive, one-year analysis of the spatial and seasonal
effects of salinity (surface and porewater) and water levels on leaf gas exchange rates of *R.*
mangle scrub mangroves in southeastern Florida Everglades. We focused our sampling on

mangrove-islands with noticeable micro-elevational differences to understand the magnitude of

influence of water levels and salinity on *R. mangle* tree physiology between mangrove-island

center and edge habitats. We addressed the following questions: (1) how do rates of leaf gas

exchange (e.g., A_{net}, g_{sw}) vary with mangrove-island micro-elevation (center vs. edge habitats)?

(2) how does leaf gas exchange respond to seasonal changes in salinity and water levels? (3)

how do water- and nutrient-use efficiencies of *R. mangle* leaves vary between mangrove-island

center and edge habitats? We hypothesized that A_{net} would be greater for *R. mangle* leaves in

higher-elevation center habitats than lower-elevation edges. We also expected that A_{net} should

vary little with season and that seasonal variation in g_{sw} would be less than variation in A_{net},

relative to the range of variability among leaves because of strong control on g_{sw} and potential

for decoupling to some degree between g_{sw} and A_{net}. Moreover, given that scrub mangroves in

Taylor River basin are strongly limited by phosphorus (i.e., soil N:P = 102-109 – Castañeda-

Moya et al. 2013), *R. mangle* plants should have high rates of P resorption. Finally, we

predicted that trees within mangrove-island centers function at a higher physiological level (i.e.,

with greater rates of A_{net} and less of a relative reduction in g_{sw}) due to lower levels of inundation

and salt stress, and should, therefore, have greater *wue* (Ball 2009) and higher relative rates of

nutrient resorption (Lugo et al. 2007, Medina et al. 2010) than trees at island edges. However,

the magnitude of the reduction in g_{sw} relative to A_{net}, because of inundation stress at mangrove-

island edge habitats should drive patters in *wue.*

METHODS

Study site

This study was conducted in the southeastern region of Everglades National Park in a

mangrove site known as Taylor Slough/Panhandle-7 (TS/Ph-7: 25.197°N, 80.642°W, Figure 1B),

one of the six mangrove sites established in 2000 as part of the Florida Coastal Everglades

Long-Term Ecological Research (FCE-LTER) program (Childers 2006; http://fcelter.fiu.edu).

TS/Ph-7 is located approximately 1.5 km inland from Florida Bay in the downstream section of

the Taylor River. Mangroves zones at TS/Ph-7 are dominated by scrub *R. mangle* L. trees, with

clusters of *L. racemosa* L. and *Conocarpus erectus* L. – a mangrove associate, intermixed with

low densities of freshwater grasses *Cladium jamaicense* (Crantz) Kük and *Eleocharis cellulosa*

Torr. (Loveless 1959). Mangrove tree heights reach 1.5 to 2 m (Ewe et al. 2006, Figure 1A).

The substrate at this site is organic mangrove peat soil (~1 m depth) overlying the karstic

bedrock (depth ~1.5-2 m, Table 1; Castañeda-Moya et al. 2011, Ewe et al. 2006). Surface (0-

45 cm depth) soils at TS/Ph-7 have high organic matter content (71%), low bulk density (0.16 g

cm^{-3}), low total nitrogen (TN, 2.5 mg cm^{-3}), and low total phosphorus (TP, 0.06 mg cm^{-3})

concentrations, resulting in a highly P-limited environment with soil N:P ratios of about 102

(Castañeda-Moya et al. 2013). Mangrove zones in Taylor River are permanently flooded for

most of the year, with an annual flooding duration averaging 360 d yr^{-1} from 2001 to 2005. The

permanent flooding results in anoxic soil conditions and buildup of porewater sulfide (range: 0.5-

2 mM) throughout the year that constrains mangrove growth (Castañeda-Moya et al. 2011,

2013). The tidal effect is negligible in Taylor River, and water flow and hydrology are

determined by seasonal precipitation, upland runoff, and wind (Michot et al. 2011, Sutula et al.

2001). The interaction between low P fertility and permanent flooding conditions results in the

formation of scrub forests with restricted tree height and aboveground productivity, high root

biomass allocation and high root: shoot ratios compared to riverine mangrove forests along

Shark River estuary in southwestern FCE (Ewe et al. 2006, Castañeda-Moya et al. 2011, 2013).

South Florida has a subtropical savanna climate per the Köppen climate classification,

where the average air temperature is between 20 and 30°C and relative humidity is high (70-

80%). Rainfall and evapotranspiration vary interannually and average 1500 and 1300 mm year^{-1}

^{1}, respectively (Abiy et al. 2019). In the Everglades, 60% of the precipitation occurs during the

wet season, and only 25% during the dry season (Duever et al. 1994). Analysis of long-term

(110-year) rainfall trends for South Florida has shown that the annual hydrologic regime can be

divided into two seasons: a wet season from May to October and a dry season from November

to April (Abiy et al. 2019). For the 2019 calendar year, temperature and relative humidity data

were collected from an eddy covariance flux tower installed at TS/Ph-7. Rainfall data were

collected from a nearby meteorological station (station name: "Taylor_River_at_mouth")

managed by the US Geological Survey as a part of the Everglades Depth Estimation Network

(https://sofia.usgs.gov/eden).

Experimental design

Eight distinct mangrove-islands of similar size (3-5 m in diameter) were selected for

repeated measurements of leaf photosynthesis and physicochemical variables from January to

December 2019. Mangrove-islands were selected within previously-established permanent

vegetation plots (two 20×20 m plots) based on their location relative to the shoreline (i.e., Taylor

River), with four islands located in the fringe mangrove zone (~50-60 m from the edge) and four

islands located in the interior forest (~100-110 m inland; Figure 1B). Mangrove-islands with

distinct micro-elevational gradients were selected, having higher soil elevation center habitats

and lower-elevation edge habitats. Mangrove-islands are surrounded by open water ponds

(Figure 1A) and remain flooded for most of the year, except the center island habitats during the
dry season (Castañeda-Moya et al. 2011, 2013, Figure S1).

Within each mangrove-island, a higher-elevation center and a lower-elevation edge
habitat were each permanently marked with an aluminum rod. At these locations, soil surface
elevation was measured for all mangrove-islands at both habitats, in addition to six
measurements in the adjacent shallow ponds surrounding mangrove-islands. Measurements
were taken using real-time kinematics referenced to the 1988 North American Vertical Datum
(NAVD88) with a Trimble R8 global navigation satellite system receiver (Trimble; Sunnyvale, CA,
USA), which has a horizontal accuracy of ±1 cm and vertical accuracy of ±2 cm.

Water level and salinity measurements

Water levels relative to the soil surface were measured monthly with a meter stick at the
permanent aluminum rods established at all island habitats. Continuous measurements of water
levels relative to the soil surface were recorded for the duration of the 2019 calendar year (see
Figure S2 for details). Continuous data were used to confirm trends in water levels and
porewater salinity measurements made by hand across islands. We use the measurements
taken by hand at each island as the predictors in our models of leaf gas exchange. A porewater
sample was collected at 30 cm depth at each habitat using a 60 ml syringe attached to a
stopcock and a rigid tubing probe (3/16" Ø). Porewater temperature and salinity were measured
using a handheld YSI conductivity-salinity-temperature meter (model Pro 30, YSI Inc., Yellow
Springs, OH, USA). A sample of surface water (when present) was also collected at each
island habitat to measure salinity and temperature.

Leaf gas exchange measurements

Photosynthetic gas exchange measurements of *R. mangle* leaves were conducted once
a month (9:00 AM to 1:00 PM, typically during sunny days) at eight scrub mangrove-islands
from January to December 2019 using a Li-COR Li-6800 portable photosynthesis system (Li-
COR Inc., Lincoln, NE, USA). At each island habitat (center vs. edge), five mature green leaves
were randomly selected from top mangrove branches. Fully developed and healthy (i.e.,
without herbivory) green leaves from the second-most distal pair of leaves on the leaf node
were chosen. The Li-6800 was clamped onto each leaf and held until machine stability was
reached (which typically happened in 2-3 minutes), wherein data points were logged.

The environmental configuration of the Li-6800 was: flow rate of 600 µmol s^{-1}, 50-70%
relative humidity of the incoming air (slightly drier than ambient air to prevent condensation in

the instrument), 400 μmol mol^{-1} CO_2 concentration, and light level of 1000 μmol m^{-2} s^{-1}, which

was determined to be non-limiting and similar to ambient environmental conditions. We used

five stability criteria, which were all assessed over a 15 s interval: the slope of A_{net} being <1

μmol m^{-2} s^{-1}, the slope of the concentration of intracellular CO_2 (c_i, which is a calculated

parameter using the difference in CO_2 concentrations between IRGAs) being <5 μmol mol^{-1}, the

slope of g_{sw} being < 0.5 mol m^{-2} s^{-1}, the slope of the transpiration rate (E) being <1 mol m^{-2} s^{-1},

and the slope of the difference in air-water vapor concentration between the sample and

reference IRGA (ΔH_2O) being <1 mmol mol^{-1}. All five stability criteria were met before logging

data. Air temperature within the leaf chamber was not controlled but allowed to vary with the

ambient conditions at the site, ranging from 26.1 to 32.0°C. Thus, leaf temperatures ranged

from 25.85 - 32.44 °C, averaging 29.41 ± 0.08 °C, in the wet season, and ranged from 25.06 -

28.72 °C, averaging 26.79 ± 0.04 °C, in the dry season. We calculated intrinsic *wue* as the ratio

of leaf net CO_2 uptake to leaf gas exchange (i.e., A_{net}/g_{sw}/1000, where we divide by 1000 to get

wue in mmol mol^{-1}).

Measurement of leaf functional traits, nutrient content, and isotopic signatures

During the monthly photosynthesis measurements in February, May, August, and

November measured mature green leaves (n = 5 per habitat, 40 in total) were collected at half

of the islands (four of the eight islands with two per location) for determination of leaf functional

traits and total carbon (TC), nitrogen (TN), and phosphorus (TP) content. Leaves were

numbered, placed in a sealed, moist bag to prevent water loss, and transported to the

laboratory in a cooler with ice for further analyses. Five senescent (i.e., yellowing) leaves were

also collected from trees in the same islands at the same time to determine carbon and nutrient

content. Leaves were removed from bags, wiped dry, and immediately weighed to obtain leaf

fresh mass at the laboratory. Green leaves were then scanned at high resolution and oven-

dried for at least 72 hours at 60°C to constant weight before recording their dry mass. Leaf area

was measured using ImageJ (Schneider et al. 2012). Leaf dry mass was recorded and used to

calculate leaf dry matter content (LDMC) as the ratio of the dry leaf mass (in mg) to its fresh

mass (in g, mg g^{-1}), percent leaf water content (1000-LDMC; %), and SLA, the ratio of leaf dry

weight to leaf area (cm^2 g^{-1}). These methods followed Cornelissen et al. (2003).

For nutrient analyses, composite leaf samples containing the five leaves from each

island habitat per collection were ground into a fine powder using a vibrating ball mill

(Pulversette 0, Frtisch GmbH, Idar-Oberstein, Germany). Green and senescent leaf samples

were stored in scintillation vials at room temperature and analyzed separately. Leaf TC and TN

content were determined with a Carlo-Erba NA-1500 elemental analyzer (Fisons Instruments

Inc., Danvers, MA, USA). TP was extracted using an acid-digest (HCl) extraction, and

concentrations of soluble reactive P were determined by colorimetric analysis (Methods 365.4

and 365.2, US EPA 1983). Leaf C and N bulk isotopic signatures (δ^{13}C, δ^{15}N) were analyzed on

a Thermo Scientific Delta V Plus CF-IRMS coupled to an 1108 elemental analyzer via a ConFlo

IV interface (Thermo Fisher Scientific, Waltham, MA, USA). All C and N analyses were

conducted at the Southeast Environmental Research Center Analysis Laboratory.

Using leaf carbon isotope fractionation values, we calculated the concentration of

intracellular CO_2 and plant water use efficiency integrated over the lifespan of the leaf tissue

samples (i.e., intrinsic water use efficiency, *WUE*) via methods described by O'Leary (1988) and

Marshall et al. (2007) (and outlined in Lambers et al. 2008). We used an ambient concentration

of atmospheric CO_2 of 408 µmol mol^{-1} for our calculations, which is a conservative estimate for

the 2019 calendar year and indicative of the site's atmospheric conditions based on IRGA

measurements from an eddy covariance tower at the site. Thus, the equation used to calculate

c_i and *WUE* from carbon isotope data were: $c_i = ((-8.5 - \delta^{13}C - 4.4) \div 22.6 \times 408)$, and *WUE* =

$(408 \times (1 - c_i \div 408)) \div 1.6$, where c_i is the value derived from the previous equation.

Additionally, the following equation was used to calculate the resorption of N and P using green

(G) and senescent (S) leaf nutrient content: Relative resorption (%) = $((G - S) \div G \times 100)$

(Pugnaire and Chapin 1993).

Statistical analyses

Repeated measures analysis of variance (ANOVA) was used to test for differences in

water level, surface water salinity, and porewater salinity among locations (fringe and interior),

island habitats (center and edge), and season (wet and dry), as well as for the interaction

between these effects and season, which was used as the repeated measure. For the repeated

measures ANOVA, islands were nested within locations and treated as experimental units. All

effects were considered fixed, except for when testing for significant differences in habitat,

which included location as a random effect to account for the nested structure of the sampling

scheme. One-way ANOVAs were used to test for differences in soil surface elevation among

locations and habitats and their interaction. Two-way ANOVAs were carried out for all leaf

functional traits and nutrient concentrations, making comparisons across all habitat and season

combinations. Tukey HSD posthoc tests were used to identify significant pairwise comparisons

when ANOVAs indicated statistical differences. Repeated measures ANOVAs were performed

using PROC MIXED (SAS Institute, Cary, NC, USA), and the one-way and two-way ANOVAs

were performed in R v3.5.1 (R Core Team 2018).

We constructed linear mixed-effects models (with a Gaussian error distribution and

identity link function) to address our research questions. Island habitat and season were

included as fixed effects in the models to address questions 1 and 2, respectively, with water

levels and porewater salinity being also included as the continuous covariates to parse out their

marginal effects. We couple inference from these models to leaf nutrient analyses and our

measurements of the hydrological environment to inform about nutrient and water use of *R.*

mangle (question 3). Before model fitting, response variables were confirmed to meet the

assumptions of data normality. Four separate models were constructed for each of four gas

exchange variables of interest: A_{net}, g_{sw}, c_i, and *wue*. For each model, fixed effects for season

(wet and dry), habitat (center and edge), porewater salinity, and water level were considered,

including interaction terms for water level and porewater salinity with season. All models

considered random intercept terms for location (i.e., fringe vs. interior), islands, and islands

nested within location. Random slopes were explored but determined not to improve model fits.

The best-fit models were determined via stepwise model comparison using AIC based on

backward selecting random effects then backward selecting fixed effects, as implemented with

the 'lmerStep' function in the lmerTest R package (Kuznetsova et al. 2017). The best fit models

included a random intercept term for islands, which helped remove variability in the data

because of the sampling design. Random effects for location were insignificant, signifying that

most of the random variance in the gas exchange data was among islands, which we consider

as the experimental unit in all mixed-effects models. The mixed-effect models were fit using

restricted maximum likelihood estimates via the lme4 R package (Bates et al. 2015). Models

were evaluated using model predicting, tabling, and plotting functions from the sjPlot R package

(Lüdecke 2018). All analyses were complete in R v3.5.1 (R Core Team 2018).

RESULTS

Mangrove-island micro-elevational differences and ecohydrology

Soil surface elevation (measured in relation to the NAVD88 datum) significantly declined

from mangrove-island center to edge habitats from -0.14 ± 0.1 m at island centers to -0.4 ± 0.02

m at island edges, a mean difference of about 30 cm ($F_{1,20}$ = 108.42, p < .001; Table 1). Water

levels relative to the soil surface were significantly higher in edge than in center habitats ($F_{1,178}$ =

178.33, p < .001), measuring on average 36.9 ± 1.4 cm in edge habitats, and 12.8 ± 1.2 cm in

mangrove-island centers (Table 2). We recorded water levels of 0 cm (i.e., non-inundated

habitats) in 10% of our measurements, and those were exclusive to mangrove-island centers

during the dry season (Figure S1C). There was a significant effect of season ($F_{1,178} = 11.11$, p

< .001) on water levels, where they increased from 17.05 ± 1.5 cm in the dry season to 30.4 ±

1.6 cm in the wet season (Table 2, Figure S1C).

Continuous water level data recorded at the fringe and interior mangrove zones

indicated similar flooding trends between locations, with lower water levels during the dry

season and higher water levels in the wet season, up to 40-47 cm above the soil surface in both

locations (Figure S2). Water levels at the interior mangrove forest always remained higher than

those registered in the fringe mangrove zone (Figure S2A). Porewater salinity was significantly

different between habitats ($F_{1,178} = 91.45$, $p < .001$) and seasons ($F_{1,178} = 17.87$, $p < .001$), with

lower salinity values in the center (21.5 ± 0.3) of the islands relative to the edge (25.1 ± 0.3)

habitats, and slightly lower porewater salinity during the dry season (22.5 ± 0.4) than in the wet

season (24.1 ± 0.3; Table 2, Figure S1D). There was no significant interaction ($F_{1,178} = 0.26$,

$p > .05$) between island habitats and seasons, indicating that the variation in porewater salinity

between habitats is independent of seasonality (Table 2). Surface water salinity was not

significantly different among center and edge habitats ($F_{1,163} = 2.36$, $p > .05$), but increased

significantly from the dry to the wet season ($F_{1,163} = 8.97$, $p < .01$, Table 2). There was also a

Tukey posthoc HSD test indicated that only island center habitats in the dry season differed

from all other pairwise comparisons (Table 2).

Rates of leaf gas exchange and their relationships to the hydrological environment

A_{net} measurements ranged from 0.1 to 15.1 µmol m^{-2} s^{-1}, with 90% of the observations

recorded between 2 and 14 µmol m^{-2} s^{-1} (see Figure S3). g_{sw} values were low, ranging from

<0.01 to 0.27 and averaging 0.1 mol m^{-2} s^{-1} (see Figure S3). Associated c_i values ranged from

40 to 377 and averaged 242 µmol mol^{-1}, with 98% of them being greater than 150 µmol mol^{-1}.

Lastly, measured rates of *wue* varied between >0.01 and 0.21 mmol CO_2 mol H_2O^{-1}, being

normally-distributed about a mean value of 0.09 mmol mol^{-1}.

The linear mixed-effects model for A_{net} included fixed effects for island habitat, porewater

salinity, water level, season, and an interaction term for water level with season (Figure S4 &

Table S4). There was substantial variation in A_{net} rates among leaves (σ^2 of about 6 µmol m^{-2} s^{-}

^{1}), and the random variation among islands was about 0.02 µmol m^{-2} s^{-1} (see Table S4). All

fixed effects were statistically significant ($p < .05$), except the interaction term, which was

marginally significant ($p = .05$) but greatly improved model fit. Mangrove edge habitats reduced

A_{net} by over 2.5 µmol m^{-2} s^{-1} relative to mangrove-island centers (Figure 2). Seasonality had a

comparable negative effect, leading to an average decrease in A_{net} of just over 2 µmol m^{-2} s^{-1} in

the wet season relative to the dry season (Figure 2). After accounting for variation in the data

because of habitat and season, the marginal effects of water level and porewater salinity were

positive, albeit weak, leading to increases in A_{net} of <0.1 µmol m^{-2} s^{-1} per cm increase in water

level (Figure 3) or per ppt increase in porewater salinity (Figure 4). Therefore, A_{net} increased as

water levels increased, with increases consistent across habitats (Figure 3); a similar pattern

was observed concerning soil porewater salinity, although the magnitude of increase in A_{net} was

smaller (Figure 4). These relationships of A_{net} with water level variability were consistent across

seasons, although rates of A_{net} were depressed during the wet season (Figure 2). The mixed-

effects model for A_{net} fit satisfactorily for these types of linear mixed-effects models modeling

leaf-gas exchange data using environmental predictors, explaining 24% of the variation in the

data, 22% of which was explained by ecohydrological data (i.e., fixed effects) (Table S4).

g_{sw} was modeled using an identical mixed-effects model as was used for A_{net} (Figure S5

& Table S5). Generally, rates of g_{sw} were low, with 98% of g_{sw} measurements being <0.2 mmol

mol^{-1}. Random variance in g_{sw} among islands was negligible, being <0.01 mmol mol^{-1}. Leaf g_{sw}

in edge habitats was statistically lower than that of mangrove-island centers ($p < .001$), being

depressed by about 0.02 mmol mol^{-1} (Figure 2). Water levels did not affect rates of g_{sw} ($p > .05$,

Figure 3, Table S5), and soil porewater salinity had a marginal effect ($p = .07$) on g_{sw}, where

conductance increased slightly at high salinities, after accounting for the effects of other

environmental variables in the model (Figure 4). The effect of season on rates of g_{sw} was

significant in the model, with the wet season leading to a 0.05 mmol mol^{-1} decrease in

conductance (Figure 2) and the interaction between water levels and season being statistically

significant (Figure 3). Overall, the mixed-effects model for g_{sw} did not fit the data as well as the

model for A_{net}. The model only explained about 12% of the variability in the data, with 9% of its

explanatory power coming from the environmental predictors (Table S5).

Although the model selection approach was the same as the other mixed-effects models,

the best-fit model for c_i differed from the models for A_{net} and g_{sw}. The model did not include a

fixed effect for soil porewater salinity (which dropped out of the model in the model selection

procedure) but included all the same fixed effects as the models for A_{net} and g_{sw}, which were all

statistically significant ($p < .001$), and a random intercept term for islands (Table S6).

Mangrove-island edge habitats had consistently higher c_i values than island centers, being

about 27 µmol mol^{-1} greater (19 to 35, 95% CI; Figure 2). The marginal effect of season alone

was similar in magnitude to that of habitat; the wet season led to a decrease in c_i of 24 µmol

mol^{-1} (14 to 34, 95% CI) relative to the dry season (Figure 2, Table S6). Water levels, by

themselves (again, the marginal effect), led to a slight decrease in c_i but had a positive

interaction with season, indicating that the relative decrease in c_i due to increasing water levels

was suppressed during the wet season (Figure 3). The random intercept term in the model (for

islands) explained a considerable amount of variation in the data (σ^2 = 128 µmol mol^{-1}, with

τ_{island} = 66 µmol mol^{-1}). The mixed-effect model for c_i fit the poorest of all four models,

explaining just under 12% of the variance in c_i, about 9% of which was explained by data from

the hydrological environment (Table S6).

Lastly, we modeled *wue* using a similar mixed-effects model to that of g_{sw}. In the model

for *wue*, all fixed effects were statistically significant ($p < .001$); however, the fixed effects were

more-subtle in magnitude. Similar to the model for c_i, porewater salinity was not included in the

best-fit model. *wue* values were normally-distributed about a mean value of 0.09 µmol mol^{-1},

with 83% of the data having values between 0.05 and 0.15 µmol mol^{-1}. Mangrove-island edge

habitats had lower *wue* by 0.01 µmol mol^{-1} than island centers (Figure 2). The marginal effect of

water level, although being statistically significant in the model, was negligible; however, the wet

season caused an increase in *wue* by 0.02 µmol mol^{-1} relative to the dry season, with the

interaction between water level and season being slightly negative (Figure 2, Figure 3).

Random variation in *wue* structured across the eight mangrove-islands was minuscule, being <

0.01 µmol mol^{-1}. Thus, the model fit for *wue* was comparable to, and slightly better than, the

model for c_i, with fixed effects explaining just over 12% of the variance in the data, about 9% of

which was explained using the environmental predictors (Table S7).

Rhizophora mangle leaf functional traits, nutrient content, and isotopic signatures

Leaf SLA values did not vary significantly between seasons ($F_{1,155}$ = 0.46, $p > .05$) and

island habitats ($F_{1,155}$ = 3.07, $p > .05$, Table 3), despite having some variation in SLA with

average values ranging from 29 to 40 cm^2 g^{-1}. Similarly, leaf water content was not significantly

different between all season-habitat combinations ($F_{1,155}$ = 0.32, $p > .05$), despite a statistically

significant effect of season alone ($F_{1,155}$ = 9.10, $p < .01$), where leaf water content was greater in

the dry season (65.6 ± 0.3%) relative to the wet season (63.8 ± 0.4%, Table 3).

Leaf TC content ranged from 400 to 450 mg g^{-1} (Table 3) and was not different between

seasons ($F_{1,12}$ = 1.10, $p > .05$), habitats ($F_{1,12}$ = 0.10, $p > .05$), or their interaction ($F_{1,12}$ = 1.77,

$p > .05$). Leaf TN concentrations were higher in the dry season compared to the wet season

($F_{1,12}$ = 11.95, $p < .01$) and ranged from 8-10 mg g^{-1} (Table 3). There was neither a significant

difference in leaf TN between habitats ($F_{1,12}$ = 1.86, $p > .05$), nor a significant interaction

between seasons and habitats ($F_{1,12}$ = 0.11, p > .05, Table 3). Leaf TP content did vary

significantly between seasons ($F_{1,12}$ = 15.05, p < .01) and had marginally-significant difference

between habitats ($F_{1,12}$ = 4.55, p = .054), but the interaction effect was not significant ($F_{1,12}$ =

0.08, p > .05). Overall, mean leaf TP values ranged from 0.42 to 0.55 mg g^{-1} across seasons

and habitats, with higher concentrations during the dry season than in the wet season and

higher leaf tissue TP values in the island center habitats compared to edge habitats (Table 3,

Figure 5). Mean N resorption for *R. mangle* leaves was slightly similar across seasons and

habitats and ranged from 60 to 63% (Table 3, Figure 5). In contrast, P resorption of leaf tissue

had a broad range compared to that of N, ranging from 73.2 ± 6.2% (center, wet season) to 78.6

± 0.2% (edge, dry season) across seasons and habitats. Overall, P resorption of *R. mangle*

leaves was higher in the edge habitats relative to the center during both seasons (Table 3).

Patterns in green leaf carbon isotope signatures (δ^{13}C) mirrored those of leaf TN and TP

concentrations. Carbon isotopic fractionation was more negative during the wet season than in

the dry season ($F_{1,12}$ = 18.88, p < .01, Table 3, Figure 5), with no statistical difference between

habitats ($F_{1,12}$ = 1.17, p > .05). Green leaves bulk δ^{13}C values ranged from -25.9 to -25.1‰

across seasons and habitats (Table 3, Figure 5). Physiologically, the differences in carbon

isotopic fractionation were estimated to result in a maximum difference of about 10 μmol mol^{-1} c_i

between the center and edge habitats and a difference of 5 to 15 μmol mol^{-1} c_i within habitats

($F_{1,12}$ = 1.71, p > .05) because of seasonality ($F_{1,12}$ = 18.88, p < .01). These differences resulted

in slightly greater, but not statistically different, c_i values in mangrove-island centers than in

edge habitats in the wet season; however, the opposite pattern was found during the dry season,

with c_i being about 10 μmol mol^{-1} greater in island edge habitats relative to their centers (Table

3). The interaction between season and habitat was marginally significant ($F_{1,12}$ = 4.53, p

= .055). Intrinsic water-use efficiency (*WUE*) was calculated from leaf δ^{13}C values; accordingly,

WUE was greatest in mangrove-island centers during the dry season relative to all habitat-

season combinations. Additionally, *WUE* was significantly lower in the wet season than the dry

season ($F_{1,12}$ = 18.88, p < .01, Table 3). Mean leaf bulk δ^{15}N values were significantly ($F_{1,12}$ =

19.66, p < .001) higher in the center habitats (-0.60 ± 0.66‰) relative to the edge (-4.79 ±

0.66‰), but there was no difference between seasons ($F_{1,12}$ = 0.17, p > .05) and no interaction

between season and island habitat ($F_{1,12}$ = 0.58, p > .05, Table 3, Figure 5A).

DISCUSSION

Our first research question asked how mangrove-island habitat affects rates of leaf gas

exchange. We can confirm our hypothesis that photosynthetic rates and stomatal conductances

are greater at island centers than edges (Figure 2). However, contrary to our expectation,

habitat driven-variation in leaf gas exchange rates was roughly equal to seasonal variation, with

no apparent decoupling between A_{net} and g_{sw} (Figure 2). Our second research question asked

whether water levels or salinity exerted a more-substantial effect on mangrove leaf gas

exchange *in situ*. Porewater salinities at TS/Ph-7 are relatively low (i.e., between 15 and 30 ppt),

compared to the levels of salinity at which leaf gas exchange rates of *R. mangle* are negatively

affected (i.e., salinities > 35 ppt) and did not vary considerably over time (Figures S1 & S2).

Therefore, we conclude that inundation stress is the primary driver of variation in *R. mangle* leaf

gas exchange rates. Lastly, we predicted that general physiological stress would be lower at

island centers than island edge habitats, leading to increased *wue* and higher rates of nutrient

resorption at centers relative to edges. Indeed, intrinsic *wue* was greater at island centers than

edge habitats, with results being consistent across gas exchange-measured *wue* and isotope-

derived *WUE.* In addition, *w*ater levels modulated leaf intrinsic water use efficiency (Figure 3,

Table 2), especially in the dry season. Patterns of nutrient resorption were less clear but

seemed to indicate differences in leaf N and P concentrations on mangrove-island centers

versus edges, illustrating differences in water and nutrient use of *R. mangle* among habitats,

which likely drive variation in leaf gas exchange rates.

The effect of mangrove-island habitat on leaf gas exchange

Our results showed significant differences in soil elevation of about 30 cm between

mangrove-island habitats (Table 1), which affected *R. mangle* leaf gas exchange rates (Figures

2, 3, 4). The soil elevation gradient at our study site is driven by differences in mangrove root

biomass and productivity between center and edge island habitats, with higher total root

biomass and productivity (top 0-90 cm of soil) observed in center habitats compared to the edge

habitats (Castañeda-Moya et al. 2011). Along this micro-elevation gradient, we measured clear

differences in A_{net} and g_{sw} (Figure 2). A_{net} was nearly 3 µmol m^{-2} s^{-1} (or 20%) greater at

mangrove-island centers than edges, and g_{sw} was >0.1 mol m^{-2} s^{-1} (or >37%) higher; these

differences were attributable to mangrove-island habitat alone, after accounting for variation

explained by water level, salinity, or seasonality (i.e., marginal differences). Associated c_i

concentrations were about 30 µmol mol^{-1} (or 12%) lower, and *wue* was >0.01 mmol mol^{-1} (or

about 10%) greater at island centers than at island edges (Figure 3).

Thus, these findings support our first hypothesis about the effect of habitat micro-

elevation (center vs. edge) on A_{net}, with overall greater leaf gas exchange rates at mangrove-

island centers compared to their edges. Interestingly, the effect of habitat on *R. mangle* leaf gas

exchange rates was similar in magnitude to the effect of season (Figure 3). The magnitude of

variation in A_{net} that we report in this study is slightly larger than the magnitude of variation

reported by Lin and Sternberg (1992), who found that A_{net} varied up to 2 µmol m^{-2} s^{-1} between

scrub and fringe *R. mangle* trees in the nearby Florida Keys. Furthermore, our A_{net}

measurements with average values between 5.7 µmol m^{-2} s^{-1} (edge habitat, wet season) and

10.2 µmol m^{-2} s^{-1} (center habitat, dry season, Figure 2), are within the range of values reported

for *R. mangle* interior scrub (5.3 µmol m^{-2} s^{-1}) and fringe (10 µmol m^{-2} s^{-1}) mangroves along a

distinct zonation pattern in the intertidal zone at Twin Cays, Belize (Cheeseman and Lovelock

2004). Island center habitats may also support greater access to mixed soil-groundwater

sources in the dry season facilitating higher leaf gas exchange rates because of increased

freshwater availability, or a reduction in the energy demand for processing saline water (Ewe et

al. 2007). Our results demonstrate the effect that higher-elevation center habitats at TS/Ph-7

have on alleviating inundation stress, which pervades scrub mangrove physiology, making trees

growing in center habitats in the dry season physiologically comparable to fringe mangroves.

Certainly, the stress relief is short-lived when water levels rise in the wet season (Table 2,

Figure S1), and leaf gas exchange rates are depressed once more (Figure 2. Figure S3).

Seasonal signals in R. mangle *physiology with implications for ecosystem functioning*

We found that A_{net} varied over 2.5 µmol m^{-2} s^{-1} (17%), and g_{sw} varied about 0.03 mol m^{-2}

s^{-1} (11%) within habitats between the wet and dry seasons (Figure 3). Despite differences in

A_{net} and g_{sw} between seasons, we found no statistical differences in c_i and *wue* between

seasons, although there was some variation (Figure 3). These differences point to habitat-

specific optimization of the diffusion and uptake of CO_2 into (i.e., c_i) and the movement of water

vapor out of (i.e., *wue*) leaves (Cardona-Olarte et al. 2006, Barr et al. 2009, Reef and Lovelock

2015, Lopes et al. 2019). As precipitation and freshwater flow increased during the wet season,

water levels increased, and mangrove-island centers experienced greater inundation levels

(Figures S1 & S2), resulting in decreased A_{net} and g_{sw} (Figure 3). A similar reduction in A_{net} and

g_{sw} was measured in mangrove-island edge habitats during the wet season (Figures 3 & 4).

Although A_{net} was depressed in the wet season, the effect of inundation levels on reducing A_{net}

was consistent across seasons (Figure 4). g_{sw} showed a similar pattern to A_{net}, being highest in

mangrove-island centers during the dry season (Figure 3). However, the effect of water levels

on g_{sw} resulted in increased g_{sw} in the wet season, an effect which was tempered during the dry

season (Figure 4).

In the Florida Everglades, irradiance peaks in April and May (Barr et al. 2009), and
rainfall and temperature reach maxima in June, July, and August (Figure 1 A, B). Thus,
photosynthetic demand for water is likely highest from April to May, at the end of the dry season
and the beginning of the wet season. During this time, we measured lower water levels and
porewater salinities relative to the peak wet season. Barr et al. (2009) recorded earlier diurnal
and more considerable reductions in g_{sw} during late May versus July or August for mangroves at
Key Largo, evidencing the greatest water-limitation on photosynthesis at the end of the dry
season. Additionally, the greatest A_{net} rates for tall fringe mangroves in the southeastern
Everglades occur during the dry season from March to May (Barr et al. 2009). The difference in
surface water and porewater salinity ($\Delta sw\text{-}pw$) can be used as a proxy for tree transpiration
(Reef and Lovelock 2015). Average $\Delta sw\text{-}pw$ measured 10.7 and 8.1 ppt at mangrove-island
edges in the wet and dry seasons, respectively, whereas it measured 7.8 and -0.3 in mangrove-
island centers in the wet and dry seasons, respectively. Indeed, transpiration was highest in
March and April (Figure S3). Thus, photosynthetic demand for water is higher in the dry season
in mangrove-island centers relative to edges or either habitat in the wet season. The drying of
the soils at slightly higher-elevation island center habitats in this scrub mangrove forest likely
facilitates increases A_{net}. Therefore, the seasonal variation in hydrology, mainly reductions in
water levels and porewater salinity during the dry season, albeit coupled with an increase in
surface water salinity in this study (Table 2), likely have critical consequences for mangrove
forest carbon fluxes at greater spatial scales. Potentially drying soils could also lead to an
increase in ecosystem respiration (Chambers et al. 2014), or non-stomatal derived CO_2 uptake
(Reef and Lovelock 2015). Future research could look at soil metabolic dynamics (e.g., soil
respiration, microbial C and N, or changes in microbial communities) with hydrology and season,
which may show unique responses in this scrub *R. mangle* forest (Lovelock 2008, Chambers et
al. 2014).

The effect of salinity and water level on R. mangle *leaf gas exchange*

During 2019, the hydrological environment (Figure 1C, D) at our study site was
seasonally dynamic (Table 2) and tended to mirror patterns in local rainfall (Figure 1B). Water
levels and porewater salinity both increased during the wet season (Table 2, Figure S1) from
the beginning of the rainy season in May through November. This likely led to increased water
column stratification via a larger freshwater lens (Hughes et al. 1998, Uncles et al. 1992).
Indeed, the difference in surface water and porewater salinity increased in the wet season, with
surface water salinities decreasing, despite a slight increase in porewater salinities (Table 2,

Figure S1). When data were grouped by season, edge habitats were slightly more saline (about

4 ppt on average) than mangrove centers (Table 2), and there were no apparent differences

between fringe and interior scrub mangrove zones (Figures S1 & S2). Comparing these

changes in the hydrological environment with previous years, long-term water level and

porewater salinity data at this site show that water level usually increases and porewater salinity

usually decreases in the wet season relative to the dry season (Castañeda-Moya et al. 2013).

We measured the opposite trend in 2019 with slight differences between seasons, likely

because it was a wet year. Total rainfall for 2019 (929 mm yr^{-1}; Figure S1) was 10% greater

than the total for rainfall 2018 (859 mm yr^{-1}; https://sofia.usgs.gov/eden).

Rates of mangrove leaf gas exchange (i.e., A_{net} and g_{sw}) typically decrease with

porewater salinity, especially along strong salinity gradients in the environment (i.e.,

gradients >30 ppt, Ball 2009, Clough and Sim 1989, Lugo et al. 2007). Porewater salinity was

only included in the linear mixed-effects models for A_{net} and g_{sw}, and its marginal effect was

minimal, slightly increasing A_{net} by <0.1 µmol m^{-2} s^{-1} per ppt increase in porewater salinity. Like

the effect of porewater salinity on A_{net}, the effect of porewater salinity on g_{sw} was small in

magnitude and consistent across seasons and mangrove-island habitats but was not statistically

significant (Figure 4). The minimal influence of porewater salinity on leaf gas exchange is likely

due to the minor seasonal and spatial variation in salinity that we measured during 2019.

Differences were not large, maximizing at 16.4 ppt and averaging 5.2 ppt, especially when

considering that *R. mangle* frequently occupies natural habitats with salinities greater than

seawater (Reef and Lovelock 2015), potentially up to 50-60 ppt (Cintron et al. 1978). At our

study site, variation in porewater salinity from long-term monitoring data (2001-2020) has shown

similar magnitudes of relatively-minor variation in porewater salinity, with overall mean values

ranging from 19-22 and rarely exceeding 30 ppt (Castañeda-Moya et al. 2013). Additionally,

long-term variation in porewater salinity (<30 ppt) across the FCE mangrove sites (Shark and

Taylor River sites) is below the critical value of 65 ppt that influences forest structure and

productivity across the FCE landscape (Castañeda-Moya et al. 2013). Thus, the limited effect

of salinity in our linear mixed-effects models is likely broadly indicative of relatively weak salinity

effects in both scrub and tall *R. mangle*-dominated forests of the Everglades. This is a

significant finding, given that these scrub forests are adapted to relatively low salinities. If

salinity increases greatly due to SLR and saltwater intrusion in the region, they will likely

experience more stressful conditions with could diminish their physiological performance, as

observed in other studies in the neotropics (Lugo et al. 2007).

The effects of inundation on *R. mangle* photosynthesis can be difficult to separate from

the effects of salinity; however, the linear mixed modeling approach we used permitted doing so.

We found that the intermittent flooding conditions of mangrove-island centers that averaged 10-

15 cm above the soil surface allowed greater A_{net} and g_{sw} than permanently flooded mangrove-

island edges, which averaged 30-40 cm water levels. This indicates that the hydrological

regime in center habitats allows mangrove soils to repeatedly flood and desiccate, which may

help the species maintain optimal stem water potentials and g_{sw} (Ball 2009, Reef and Lovelock

2015). In typical greenhouse experiments where mangrove seedlings are grown, inundation

alone has little effect on photosynthetic rates or biomass production (Pezeshki et al. 1990b,

Hoppe-Speer et al. 2011). However, inundation may sometimes lead to increases in leaf gas

exchange rates over the short term and often interacts with salinity over time to reduce A_{net}, $g_{sw,}$

and growth rates (e.g., Cardona-Olarte et al. 2013). Thus, water levels and flooding duration

are key drivers controlling A_{net} in mangroves, and mangroves seem to optimize photosynthesis

to water levels physiologically. For instance, findings from a long-term greenhouse inundation

study by Farnsworth and Ellison (1996) exemplify how short-term responses of *R. mangle* to

inundation differ from longer-term responses. Over several years, high inundation levels led to

steady declines in A_{net} of up 25% for a given g_{sw} and decreases in growth rates. Results of the

high-water level (30-40 cm above soil surface) treatment were similar to those of the low water

level (10-15 cm) treatment, suggesting that *R. mangle* physiology is optimized at inundation

levels that reach just a few centimeters above the soil surface at high water level (Ellison and

Farnsworth 1997). Further research could assess the coupled effect of the depth and duration

of flooding with salinity in the FCE by characterizing photosynthetic rates across a landscape-

scale gradient that encompassed multiple sites and a broad range of salinity.

Although initial increases in *R. mangle* g_{sw} can result from short term inundation (Krauss

et al. 2006, Hoope-Speer et al. 2011), especially at low salinities (Pezeshki et al. 1990b),

several studies have linked stomatal closure to longer-term inundation (Kozlowski 1997, Ellison

and Farnsworth 1997). We measured depressed g_{sw} during the wet season and in mangrove-

island edge habitats relative to centers; however, this was not attributable to water levels after

accounting for variation in seasonality and habitat, in that g_{sw} increased with increasing water

levels during the wet season. Our measurements of g_{sw} were consistent with those reported in

other studies from across a range of inundation levels (Clough and Sim 1989, Lin and Sternberg

1992, Ellison and Farnsworth 1997, Krauss et al. 2006, Lugo et al. 2007, Barr et al. 2009),

supporting the understanding that *R. mangle* leaves limit g_{sw} in response to flooding. Limits on

g_{sw} seek to optimize c_i for carbon gain without losing unnecessary amounts of water, but our

findings show that g_{sw} can increase with freshwater inputs, resulting in a decrease in c_i as A_{net}

increases (Figure 3), likely because of faster Calvin cycle reactions (see supplemental

information Figure S9). Interestingly, the scrub *R. mangle* leaves of TS/Ph-7 operate with low c_i

concentrations (range = 220-260 µmol mol^{-1}), which suggests pervasive inundation stress.

Such pervasive inundation stress likely leads to water and nutrient (i.e., leaf N and Rubisco)-

stressed photosynthesis, which decreases max A_{net} at mangrove-island edges by maximum

rates of carboxylation, especially in the wet season.

R. mangle nutrient use at TS/Ph-7

We found little variation (60-63%) in N resorption efficiencies for *R. mangle* leaves

across scrub mangrove-island habitats (Figure 5). In contrast, higher overall efficiencies of P

resorption (73-79%) of leaf tissue were measured across habitats, with higher P resorption in

mangrove-island edge habitats relative to centers, suggesting higher P availability in island

centers. Our findings are roughly comparable to N and P resorption efficiencies for *R. mangle*

in the control plots of scrub-dominated forests in Panama (~50%, and 80%, respectively;

Lovelock et al. 2004). Resorption of nutrients from senescent leaves before leaf fall is a within-

stand nutrient recycling mechanism that may reduce nutrient losses via tidal export in coastal

systems (Vitousek 1982, Aerts and Chapin 2002). Like other tropical trees, mangroves exhibit

several physiological mechanisms that reduce nutrient losses via tidal exchange, including

resorption of N and P before leaf abscission, which can lead to increased availability of limiting

nutrients and ultimately change nutrient use and conservation patterns (Twilley et al. 1986,

Alongi et al. 1992, Feller et al. 2003a, 2003b).

Our findings suggest that *R. mangle* conserves P better than N in this P-limited

environment and indicate that canopy N and P resorption efficiency at TS/Ph-7 potentially

results from the differential acquisition of these nutrients from the soil between habitat, and

variation in use of these nutrients among leaf stages. Mangrove species prioritize resorption of

nutrients that are limited in the soil, and it has been suggested that plants growing in nutrient-

poor environments resorb a higher proportion of nutrients, potentially decreasing nutrient loss by

efficient nutrient recycling (Chapin and Moilanen 1991). At our study site, low soil TP

concentrations probably determine the higher recycling efficiency of P relative to N. Indeed, soil

TP concentrations in the upper 50 cm of soil at TS/Ph-7 (0.06 ± 0.004 mg cm^{-3}) are three times

lower than soils at the mouth of Shark River estuary (SRS-6), which is dominated by fertile well-

developed tall riverine mangroves. Such low TP concentrations result in extreme P limitation at

TS/Ph-7 with average soil N:P ratios of 102 ± 6 (Castañeda-Moya et al. 2013). Therefore, the

non-homogeneous distribution of essential nutrients within mangrove habitats creates distinct

gradients and hot spots along the intertidal zone, influencing the efficiency of internal nutrient

recycling. This is supported by observations that nutrient resorption efficiencies in mangroves

vary with nutrient availability, e.g., via nutrient addition (Feller et al. 1999, Feller et al. 2003b) or

along natural fertility gradients (Medina et al. 2010). Such variation in nutrient availability and

resorption efficiencies within mangrove trees likely scales with variation in photosynthesis and

productivity (e.g., growth, litterfall) and carbon residence times (e.g., soil and biomass dynamics)

of mangrove forests.

Foliar δ^{15}N values integrate long-term processes of N sources because isotopic

fractionation against the heavier isotope (i.e., ^{15}N) occurs during N transformations and

interactions between biotic (e.g., mycorrhizal fungi, or bacteria) and biogeochemical (e.g.,

nitrification, denitrification) nutrient cycling processes (Garten 1993). In our study site, patterns

of δ^{15}N in *R. mangle* leaves differed drastically between mangrove habitats, with values around

-4 to -5‰ for mangrove edge habitats and between 0 and -1‰ for island centers, indicating

lower ^{15}N discrimination in island center habitats (Table 3, Figure 5A). These δ^{15}N values are

considerably more depleted than the *R. mangle* leaf δ^{15}N values reported for riverine

mangroves along Shark River estuary (He et al. 2020), where values were negatively correlated

with distance inland from the mouth of the estuary with more enriched leaves occurring near the

mouth of Shark River (4‰) relative to upstream (0.4‰) regions. Reported δ^{15}N values for *R.*

mangle leaves across different ecotypes in the neotropics range from 0 to -11‰, with more

negative values for scrub mangrove forests (-5 to -10‰) than for fringe mangroves (0-7‰; Reis

et al. 2017a). Similarly, Medina et al. (2010) showed that leaves from interior scrub mangrove

communities had more negative δ^{15}N values than tall fringe mangroves in eastern Puerto Rico (-

12‰ vs. 0‰, respectively). Those δ^{15}N values were more negative than those reported for

scrub *R. mangle* forests in Florida (Fry et al. 2000), Belize (McKee et al. 2002, Wooller et al.

2003, Fogel et al. 2008), or Brazil (Reis et al. 2017b).

Patterns of foliar δ^{15}N between mangrove ecotypes can be discerned using *in situ* leaf

nutrient content. For example, a direct relationship between ^{15}N discrimination and leaf N:P

ratios of *R. mangle* leaves previously reported for the six FCE mangrove sites, including our

study site, indicates that leaf N:P ratios accounted for 70% of the variability in ^{15}N discrimination

(Mancera-Pineda et al. 2009). Thus, foliar ^{15}N composition can reflect *in situ* leaf N-status and

differences in plant N-use. Hypoxic conditions in the soil may inhibit denitrification and

ammonia volatilization, two processes that enrich the soil substrate in ^{15}N (Craine et al. 2015).

Therefore, the substrate should be less enriched at mangrove-island edges relative to their

centers, because of interactions with the soil and the open water channels, which can alleviate

hypoxia. Thus, it appears that more negative δ^{15}N values in the edge habitats may be

associated with lower inorganic N (i.e., porewater ammonium) use by edge mangrove trees

compared to those in center island habitats (Fry et al. 2000). However, our results contrast

slightly with those of Mancera-Pineda et al. (2009), who reported mean δ^{15}N values of +3 from

65 mature leaves collected in 2001 at our study site. We argue that differences in δ^{15}N values

between the two studies could be attributed to the location where leaves were collected during

the 2009 study, concluding that it is very likely that Mancera-Pineda et al. only collected leaves

from the center of mangrove-islands, avoiding edge habitats. Taking this into consideration,

mangrove-island centers potentially may have even more-positive δ^{15}N signatures than we

found, illustrating that in the center of mangrove-islands, N is taken up by roots in inorganic

soluble forms (e.g., porewater ammonium, nitrate) and not biotically via root symbionts.

Another potential explanation of why δ^{15}N values were more negative at mangrove-

island edges than in their centers is because lateral surface roots of *R. mangle* can extend into

open water where they associate with symbiotic biofilms (i.e., algae and aquatic bacteria) that

facilitate N acquisition from open water (Potts 1979). A significant source of isotopic

discrimination occurs during N transfer between belowground symbionts (e.g., mycorrhizal fungi

or bacteria) and plant roots during nitrification, denitrification, and ammonia volatilization. The

lighter isotope ^{14}N reacts faster than ^{15}N (i.e., it is preferentially given to the host plant by the

symbiont) so that plant tissues are depleted while substrates are enriched (Högberg 1997,

Robinson 2001). Indeed, at our study site, we observed several long, absorptive, fine lateral

root systems that protruded from the edge of mangrove-islands into the open water ponds,

which were colonized by algal biofilms. Mangrove trees can potentially adapt to nutrient

shortage or localized nutrient deficiencies in the soil by altering patterns of nutrient use. This

plant strategy may maximize the efficiency of capturing limiting resources essential for growth

(e.g., N, P) from soil or surrounding open water areas in nutrient-poor environments such as

Taylor River, as proposed by the optimal plant allocation theory (Chapin et al. 1987, Gleeson

and Tilman 1992). We observed a slight decrease in foliar δ^{15}N during the wet season (Figure

5A, Table 3) as water levels and porewater salinity increased, suggesting that N-acquisition by

R. mangle via algal biofilms may be slightly greater in the dry season than in the wet season.

Lastly, highly depleted (i.e., negative) N-isotope values in leaf tissues are characteristic of

tropical wetlands with P limitation because P limitation increases N fractionation, especially in

flooded wetlands with limited P pedogenesis (McKee et al. 2002, Troxler 2007, Medina et al.

2008). This is likely the case with the scrub *R. mangle* forest at TS/Ph-7, where the main

source of P is brackish groundwater discharge (Price et al. 2006). Soil total P concentrations in

the top 10 cm of the peat soils at this site have measured 0.055 (± 0.01) mg cm^{-3}, with atomic

N:P ratios of roughly 72 (± 2) (Mancera-Pineda et al. 2009), which is considerably lower than

soils of most mangrove forests globally, but consistent with mangrove forests in karstic

environments (Rovai et al. 2018).

CONCLUSIONS

Habitat heterogeneity, resulting from micro-elevational differences in mangrove tree

locations on islands within the open water, mangrove-island forest landscape, drives variation in

scrub *R. mangle* leaf physiological performance. In particular, mangrove-island edge habitats

experience greater and more-prolonged inundation than island centers in a seasonal dynamic,

which leads to reductions in g_{sw}, reduced A_{net}, and lower intrinsic *wue*. Conversely, mangrove-

island center habitats are alleviated from inundation stress in the dry season, leading to

increases in A_{net} and g_{sw}. Interestingly, c_i levels increase with increasing water levels because

inundation likely slows not only g_{sw}, but the entire biochemical process of CO_2 assimilation,

including mesophyll and lower level (i.e., cell wall, plasma membrane, cytosol) conductance.

Additionally, differences in nutrient acquisition and use patterns among scrub *R. mangle* trees

growing at island edges vs. centers affect leaf-nutrient status and photosynthetic potential.

The findings from this study indicate that it is the interaction of inundation stress with

mangrove-island micro-elevational habitat in the flooded scrub mangroves of the southeastern

Florida Everglades that principally alters tree water and nutrient-use dynamics, which appear to

cascade to affect leaf gas exchange rates through their effects on g_{sw}. Reductions in A_{net}

interact with the salinity of the water that inundates scrub *R. mangle* trees, in theory, because

g_{sw} rates are low and primarily respond to water loss from leaves rather than carbon gain (see

supplemental section on *R. mangle* CO_2 assimilation and stomatal behavior). In our field

measurements, however, we found that prolonged inundation more than porewater salinity

drives reduction in A_{net} because the hydrological regime at Taylor River is characterized by long

hydroperiods and minor fluctuations in salinity throughout the year (Figure S2). At the forest

level, such physiological differences in scrub mangrove functioning with habitat and hydrological

environment can help inform ecosystem carbon cycle models and mangrove forest responses to

SLR and saltwater intrusion.

ACKNOWLEDGEMENTS

This research was supported by the Department of Interior – National Park Service

(Cooperative Agreement #P16AC00032) and the Florida Coastal Everglades Long-Term

Ecological Research (FCE-LTER) program funded by the National Science Foundation (Grant

#DEB-1832229). We thank Mandy Kuhn, Austin Pezoldt, and Ralph Diaz-Hung for field

assistance. We thank the Everglades National Park for granting research permits and the

Florida Bay Interagency Science Center-Everglades National Park (FBISC-ENP) for logistic

support during the study. We acknowledge Dr. Zoran Bursac from the Center for Statistical

Consulting and Collaboration at FIU for statistical advice. Finally, we thank Prof. Ernesto Medina

and two anonymous reviewers for constructive comments on an earlier version of this

manuscript. This is contribution number 1040 from the Southeast Environmental Research

Center in the Institute of Environment at Florida International University.

REFERENCES

Abiy AZ, Melesse AM, Abtew W, Whitman D (2019) Rainfall trend and variability in Southeast Florida: Implications for freshwater availability in the Everglades. PLoS One 14:e0212008.

Aerts R, Chapin FS. (2002) The mineral nutrition of wild plants revisited: a re-evaluation of processes and patterns. Adv Ecol Res 30:1-67.

Alongi DM (2008) Mangrove forests: Resilience, protection from tsunamis, and responses to global climate change. Estuarine Coastal Shelf Sci 76:1-13.

Alongi DM, Boto KG, Robertson AI (1992) Nitrogen and phosphorus cycles. In: Robertson AI, Alongi DM (eds) Tropical Mangrove Ecosystems. American Geophysical Union, Washington, DC.

Ball MC (2009) Photosynthesis in mangroves. Wetlands Australia 6:12-22.

Ball MC (1988) Ecophysiology of mangroves. Trees-Struct Funct 2:129-142.

Ball MC (1996) Comparative ecophysiology of mangrove forest and tropical lowland moist rainforest. In: Mulkey SS, Chazdon RL, Smith AP (eds) Tropical forest plant ecophysiology Springer, Boston, Massachusetts, pp 461-496.

Barr JG, Fuentes JD, Engel V, Zieman JC (2009) Physiological responses of red mangroves to the climate in the Florida Everglades. J Geophys Res Biogeosci 114:G02008.

Bates D, Mächler M, Bolker B, Walker S. (2015) Fitting linear fixed-effects models using lme4. J Stat Softw 67:1-48.

Bjorkman O, Demmig B, Andrews TJ (1988) Mangrove photosynthesis: response to high-irradiance stress. Funct Plant Bio 15:43-61.

Cardona-Olarte P, Krauss KW, Twilley RR (2013) Leaf gas exchange and nutrient use efficiency help explain the distribution of two Neotropical mangroves under contrasting flooding and salinity. Int J For Res 2013:1-10. doi:10.1155/2013/524625

Cardona-Olarte P, Twilley RR, Krauss KW, Rivera-Monroy VH (2006) Responses of neotropical mangrove seedlings grown in monoculture and mixed culture under treatments of hydroperiod and salinity. Hydrobiologia 569:325-341.

Castañeda-Moya E, Twilley RR, Rivera-Monroy VH, Marx BD, Coronado-Molina C, Ewe SML (2011) Patterns of root dynamics in mangrove forests along environmental gradients in the Florida Coastal Everglades, USA. Ecosystems 14:1178-1195.

Castañeda-Moya E, Twilley RR, Rivera-Monroy VH (2013) Allocation of biomass and net primary productivity of mangrove forests along environmental gradients in the Florida Coastal Everglades, USA. For Ecol Manage 307:226-241.

Chambers LG, Davis SE, Troxler T, Boyer JN, Downey-Wall A, Scinto LJ (2014) Biogeochemical effects of simulated sea level rise on carbon loss in an Everglades mangrove peat soil. Hydrobiologia 726:195-211.

Chapin FS, Bloom AJ, Field CB, Waring RH (1987) Plant responses to multiple environmental factors. BioScience 37:49-57.

Chapin FS, Moilanen L (1991) Nutritional controls over nitrogen and phosphorus resorption from Alaskan birch leaves. Ecology 72:709-715.

Cheeseman J, Herendeen L, Cheeseman A, Clough B (1997) Photosynthesis and photoprotection in mangroves under field conditions. Plant Cell Environ 20:579-588.

Cheeseman J, Lovelock C (2004) Photosynthetic characteristics of dwarf and fringe Rhizophora mangle L. in a Belizean mangrove. Plant Cell Environ 27:769-780.

Chen R, Twilley RR (1999) A simulation model of organic matter and nutrient accumulation in mangrove wetland soils. Biogeochemistry. 44:93-118.

Childers DL (2006) A synthesis of long-term research by the Florida Coastal Everglades LTER Program. Hydrobiologia. 569:531-544.

Cintron G, Lugo AE, Douglas D, Pool J, Morris G (1978) Mangroves of Arid Environments in
Puerto Rico and Adjacent Islands. Biotropica 10:110-121.
Clough B, Sim R (1989) Changes in gas exchange characteristics and water use efficiency of
mangroves in response to salinity and vapour pressure deficit. Oecologia 79:38-44.
Cornelissen JH, Lavorel S, Garnier E, Díaz S, Buchmann N, Gurvich DE, Reich PB, Ter Steege
H, Morgan HD, Van Der Heijden MG, Pausas JG (2003) A handbook of protocols for
standardised and easy measurement of plant functional traits worldwide. Aust J Bot
51:335-80.
Craine JM, Brooksheir ENJ, Cramer MD, Hasselquist NJ, Koba K, Marin-Spiotta E, Wang L
(2015) Ecological interpreations of nitrogen isotope ratios of terrestrial plans and soils
Plant Soil 396:1-26.
Duever MJ, Meeder JF, Meeder LC, McCollom JM (1994) The climate of South Florida and its
role in shaping the Everglades ecosystem. In: Davis SM,Ogden JC (eds) Everglades:
The Ecosystem and Its Restoration. St. Lucie Press, Delray Beach, Florida, pp 225-248.
Ellison AM, Farnsworth EJ (1997) Simulated sea level change alters anatomy, physiology,
growth, and reproduction of red mangrove (*Rhizophora mangle* L.). Oecologia 112:435-
446.
Ewe SM, Gaiser EE, Childers DL, Iwaniec D, Rivera-Monroy VH, Twilley RR (2006) Spatial and
temporal patterns of aboveground net primary productivity (ANPP) along two freshwater-
estuarine transects in the Florida Coastal Everglades. Hydrobiologia 569:459-474.
Ewe SM, Sternberg LSL, Childers DL. 2007. Seasonal plant water uptake patterns in the saline
southeast Everglades ecotone. Oecologia 152: 607-616.
Farnsworth EJ, Ellison AM (1996) Sun-shade adaptability of the red mangrove, *Rhizophora
mangle* (Rhizophoraceae): Changes through ontogeny at several levels of biological
organization. Amer J Bot 83:1131-1143.
Feller IC (1995) Effects of nutrient enrichment on growth and herbivory of dwarf red mangrove
(Rhizophora mangle). Ecol Monogr 65:477-505.
Feller IC, Whigham DF, O'Neil JP, McKee KL (1999) Effects of nutrient enrichment on within-
stand cycling in a mangrove forest. Ecology. 80:2193-2205.
Feller IC, McKee KL, Whigham DF, O'Neill JP (2003a) Nitrogen vs. phosphorus limitation across
an ecotonal gradient in a mangrove forest. Biogeochemistry 62:145-175.
Feller IC, Whigham DF, McKee KL, Lovelock CE (2003b) Nitrogen limitation of growth and
nutrient dynamics in a disturbed mangrove forest, Indian River Lagoon, Florida.
Oecologia 134:405-414.
Field CD (1984) Ions in mangroves. In: Teas HJ (ed) Physiology and management of
mangroves. Springer, Dordrecht, Netherlands, pp 43-48.
Field CD (1995) Impact of expected climate change on mangroves. Hydrobiologia 295:75-81.
Fogel M, Wooller M, Cheeseman J, Smallwood B, Roberts Q, Romero I, Jacobsen Meyers M
(2008) Unusually negative nitrogen isotopic compositions (δ^{15}N) of mangroves and
lichens in an oligotrophic, microbially-influenced ecosystem. Biogeosci Discuss 5:1693-
1704.
Fry B, Bern A, Ross M, Meeder J (2000) δ^{15}N studies of nitrogen use by the red mangrove,
Rhizophora mangle L. in south Florida. Estuarine Coastal Shelf Sci 50:291-296.
Garten CTJ (1993) Variation in foliar ^{15}N abundance and the availability of soil nitrogen on
Walker Branch watershed. Ecology 74:2098-2113.
Gleeson SK, Tilman D (1992) Plant allocation and the multiple limitation hypothesis. Am Nat
139:1322-1343.
Golley F, Odum HT, Wilson RF (1962) The structure and metabolism of a Puerto Rican red
mangrove forest in May. Ecology 43:9-19.

He B, Lai T, Fan H, Wang W, Zheng H (2007) Comparison of flooding-tolerance in four
mangrove species in a diurnal tidal zone in the Beibu Gulf. Estuarine Coastal Shelf Sci
74:254-262.
He D, Rivera-Monroy VH, Jaffé R, Zhao X (2020) Mangrove leaf species-specific isotopic
signatures along a salinity and phosphorus soil fertility gradients in a subtropical estuary.
Estuarine Coastal Shelf Sci 106768.
Högberg P (1997) ^{15}N natural abundance in soil-plant systems. New Phytol 137:179-203.
Hoppe-Speer SC, Adams JB, Rajkaran A, Bailey D (2011) The response of the red mangrove
Rhizophora mucronata Lam. to salinity and inundation in South Africa. Aquat Bot 95:71-
76.
Hughes CE, Binning P, Willgoose GR (1998) Characterisation of the hydrology of an estuarine
wetland. J Hydrol 211:34-49.
Koch MS, Snedaker SC (1997) Factors influencing *Rhizophora mangle* L. seedling development
in Everglades carbonate soils. Aquat Bot 59:87-98.
Kozlowski TT (1997) Responses of woody plants to flooding and salinity. Tree Physiol 17:490-
490.
Krauss KW, McKee KL, Lovelock CE, Cahoon DR, Saintilan N, Reef R, Chen L (2014) How
mangrove forests adjust to rising sea level. New Phytol 202:19-34.
Krauss KW, Twilley RR, Doyle TW, Gardiner ES (2006) Leaf gas exchange characteristics of
three neotropical mangrove species in response to varying hydroperiod. Tree Physiol
26:959-968.
Kuznetsova A, Brockhoff PB, Christensen RHB (2017) lmerTest Package: Tests in Linear Mixed
Effects Models. J Stat Softw 82:13.
Lambers H, Chapin FS, Pons TL (eds) (2008) Plant physiological ecology. Springer, New York,
New York.
Lamers LPM, Govers LL, Janssen ICJM, Geurts JJM, Van der Welle MEW, Van Katwijk MM,
Van del Heide T, Roelofs JGM, Smolders AJP (2013) Sulfide as a soil phytotoxin - a
review. Front Plant Sci 4:1-14.
Lawton JR, Todd ANN, Naidoo DK (1981) Preliminary inverstigations into the structure of the
roots of mangroves, *Avicennia marina* and *Brugeria gymnorrhiza*, in relation to ion
uptake. New Phytol 88:713-722.
Lin G, Sternberg LSL (1992) Comparative study of water uptake and photosynthetic gas
exchange between scrub and fringe red mangroves, *Rhizophora mangle* L. Oecologia
90:399-403.
Lopes D, Tognella M, Falqueto A, Soares M (2019) Salinity variation effects on photosynthetic
responses of the mangrove species *Rhizophora mangle* L. growing in natural habitats.
Photosynthetica 57:1142-1155.
Loveless CM (1959) A study of the vegetation in the Florida Everglades. Ecology 40:2-9.
Lovelock CE (2008) Soil respiration and belowground carbon allocation in mangrove forests.
Ecosystems 11:342-354.
Lovelock CE, Feller I (2003) Photosynthetic performance and resource utilization of two
mangrove species coexisting in hypersaline scrub forest. Oecologia 134:455-462.
Lovelock CE, Feller IC, McKee KL, Engelbrecht BM, Ball MC (2004) The effect of nutrient
enrichment on growth, photosynthesis and hydraulic conductance of dwarf mangroves in
Panama. Funct Ecol 18:25-33.
Lovelock CE, Atwood T, Baldock J, Duarte CM, Hickey S, Lavery P, Masque P, Macreadie PI,
Ricart AM, Serrano O, Steven A (2017) Assesing the risk of carbon dioxide emissions
from blue carbon ecosystems. Front Ecol Environ 15:257-265.
Lüdecke D (2018) sjPlot: Data visualization for statistics in social science. R package version 2.
https://cran.r-project.org/web/packages/sjPlot/

Lugo AE, Medina E, Cuevas E, Cintrón G, Nieves ENL, Novelli YS (2007) Ecophysiology of a
mangrove forest in Jobos Bay, Puerto Rico. Carib J Sci 43:200-219.
Lugo AE, Snedaker SC (1974) The ecology of mangroves. Ann Rev Ecol System 5:39-64.
Mancera-Pineda JE, Twilley RR, Rivera-Monroy VH (2009) Carbon ($\delta^{13}C$) and nitrogen ($\delta^{15}N$)
isotopic discrimination in mangroves in Florida coastal Everglades as a function of
enviromnetal stress. Contrib Mar Sci 38:109-129.
Marshall JD, Brooks JR, Lajtha K (2007) Stable isotopes in ecology and environmental science.
In: Michener R, Lajtha K (eds) Sources of variation in the stable isotopic composition of
plants .Blackwell, Singapore, pp 22-60.
McKee KL (1993) Soil physicochemical patterns and mangrove species distribution-reciprocal
effects? J Ecol 81:477-487.
McKee KL (2011) Biophysical controls on accretion and elevation change in Caribbean
mangrove ecosystems. Estuarine Coastal Shelf Sci 91:475-483.
McKee KL, Cahoon DR, Feller IC (2007) Caribbean mangroves adjust to rising sea level
through biotic controls on change in soil elevation. Global Ecol Biogeogr 16:545-556.
McKee KL, Feller IC, Popp M, Wanek W (2002) Mangrove isotopic ($\delta^{15}N$ and $\delta^{13}C$) fractionation
across a nitrogen vs. phosphorus limitation gradient. Ecology 83:1065-1075.
Mcleod E, Chmura GL, Bouillon S, Salm R, Bjork M, Duarte CM, Lovelock CE, Schlesinger WH,
Silliman BR (2011) A blueprint for blue carbon: Toward an improved understanding of
the role of vegetated coastal habitats in sequestering CO_2. Front Ecol Environ 9:552-560.
Medina E (1999) Mangrove physiology: the challenge of salt, heat, and light stress under
recurrent flooding. In: Yañez-Arancibia A, Lara-Domínguez L (eds) Ecosistemas de
manglar en América tropical. Instituto de Ecología A.C. México, UICN/ORMA, Costa
Rica, NOAA/NMFS Silver Spring, MD USA. pp 109-126.
Medina E, Cuevas E, Lugo AE (2010) Nutrient relations of dwarf *Rhizophora mangle* L.
mangroves on peat in eastern Puerto Rico. Plant ecolog 207:13-24.
Medina E, Francisco M (1997) Osmolality and $\delta^{13}C$ of leaf tissues of mangrove species from
environments of contrasting rainfall and salinity. Estuarine Coastal Shelf Sci 45:337-344.
Medina E, Francisco M, Quilice A (2008) Isotopic signatures and nutrient relations of plants
inhabiting brackish wetlands in the northeastern coastal plain of Venezuela. Wetlands
Ecol Manage 16:51.
Mendelssohn I, McKee K (2000) Saltmarshes and mangroves. In: Barbour MG, Billings WD
(eds) North American terrestrial vegetation. Cambridge University Press, New York, NY,
pp 501-536.
Michot B, Meselhe EA, Rivera-Monroy VH, Coronado-Molina C, Twilley RR (2011) A tidal creek
water budget: estimation of groundwater discharge and overland flow using hydrologic
modeling in the southern Everglades. Estuarine Coastal Shelf Sci 93:438-448.
Murdiyarso D, Purbopuspito J, Kauffman JB, Warren MW, Sasmito SD, Donato DC, Manuri S,
Krisnawati H, Taberima S, Kurnianto S (2015) The potential of Indonesian mangrove
forests for global climate change mitigation. Nat Clim Change 5:1089-1092.
Naidoo G (1983) Effects of flooding on leaf water potential and stomatal resistance in *Bruguiera*
gymnorrhiza (L.) LAM. New Phytol 93:369-373.
Neubauer SC, Franklin RB, Berrier DJ (2013) Saltwater intrusion into tidal freshwater marshes
alters the biogeochemical processing of organic carbon. Biogeosciences Discuss
10:10685-10720.
Nickerson NH, Thibodeau FR (1985) Association between porewater sulfide concentrations and
distribution of mangroves. Biogeochemistry 1:183-192.
O'Leary MH (1988) Carbon isotopes in photosynthesis. BioScience. 38:328-336.
Parida AK, Jha B (2010) Salt tolerance mechanisms in mangroves: a review. Trees-Struct Funct
24:199-217.

Pezeshki S, DeLaune R, Patrick Jr W (1990a) Flooding and saltwater intrusion: Potential effects
on survival and productivity of wetland forests along the U.S. Gulf Coast. For Ecol
Manage 33:287-301.
Pezeshki S, DeLaune R, Patrick Jr W (1990b) Differential response of selected mangroves to
soil flooding and salinity: gas exchange and biomass partitioning. Can J For Res 20:869-
874.
Pezeshki SR, DeLaune RD (2012) Soil oxidation-reduction in wetlands and its impact on plant
functioning. Biology 1:196-221.
Potts M (1979) Nitrogen fixation (acetylene reduction) associated with communities of
heterocystous and non-heterocystous blue-green algae in mangrove forests of Sinai.
Oecologia 39:359-373.
Pugnaire FI, Chapin FS (1993) Controls over nutrient resorption from leaves of evergreen
Mediterranean species. Ecology 4:124-129.
Price RM, PK Swart, Fourqurean JW (2006) Coastal groundwater discharge - an additional
source of phosphorus for the oligotrophic wetlands of the Everglades. Hydrobiologia 569:
23-36.
R Core Team (2018) R: a language and environment for statiscal computing. R foundation for
staistical computing. Vienna, Austria. www.r-project.org
Reef R, Lovelock CE (2015) Regulation of water balance in mangroves. Ann Bot 115:385-395.
Reis CRG, Nardoto GB, Oliveira RS (2017a) Global overview on nitrogen dynamics in
mangroves and consequences of increasing nitrogen availability for these systems.
Plant Soil 410:1-19.
Reis CRG, Nardoto GB, Rochelle ALC, Vieira SA, Oliveira RS (2017b) Nitrogen dynamics in
subtropical fringe and basin mangrove forests inferred from stable isotopes. Oecologia
183:841-848.
Robinson D (2001) $\delta^{15}N$ as an integrator of the nitrogen cycle. Trends Ecol Evol 16:153-162.
R Core Team (2018) R: a language and environment for statiscal computing. R foundation for
staistical computing. Vienna, Austria. www.r-project.org
Ross MS, Meeder JF, Sah JP, Ruiz PL, Telesnicki GJ (2000) The southeast saline Everglades
revisited: 50 years of coastal vegetation change. J Veg Sci 11:101-112.
Rovai AS, Twilley RR, Castañeda-Moya E, Riul P, Cifuentes-Lara M, Manrow-Villalobos M,
Horta PA, Simonassi JC, Fonseca AL, Pagliosa PR (2018) Global controls of carbon
storage in mangrove soils. Nat Clim Change 8:534-538.
Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image
analysis. Nat Methods 9:671-675.
Scholander P (1968) How mangroves desalinate seawater. Physiol Plant 21:251-261.
Scholander P, Hammel H, Hemmingsen E, Garey W (1962) Salt balance in mangroves. Plant
Physiol 37:722.
Sobrado M (2000) Relation of water transport to leaf gas exchange properties in three
mangrove species. Tree-Struct Funct 14:258-262.
Sutula M, Day JW, Cable JE, Rudnick D (2001) Hydrological and nutrient budgets of freshwater
and estuarine wetlands of Taylor Slough in southern Everglades, Florida (U.S.A.).
Biogeochemistry 56:287-310.
Tully K, Gedan K, Epanchin-Niell R, Strong A, Bernhardt ES, BenDor T, Mitchell M, Kominoski J,
Jordan TE, Neubauer SC, Weston NB (2019) The invisible flood: The chemistry, ecology,
and social implications of coastal saltwater intrusion. BioScience 69:368-378.
Tomlinson PB (2016) The botany of mangroves. Cambridge University Press, Cambridge,
Massachusetts.
Troxler TG (2007) Patterns of phosphorus, nitrogen and $\delta^{15}N$ along a peat development
gradient in a coastal mire, Panama. J Trop Ecology 23:683-691.

Twilley RR, Castañeda-Moya E, Rivera-Monroy VH, Rovai A (2017) Productivity and carbon
1074   dynamics in mangrove wetlands. In: Rivera-Monroy VH, Lee SY, Kristensen E, Twilley
1075   RR (eds) Mangrove ecosystems: A global biogeographic perspective. Springer
1076   International, pp 113-162.
1077   Twilley RR, Lugo AE, Patterson-Zucca C (1986) Litter production and turnover in basin
1078   mangrove forests in southwest Florida. Ecology 67:670-683.
1079   Twilley RR, Rivera-Monroy V (2005) Developing performance measures of mangrove wetlands
1080   using simulation models of hydrology, nutrient biogeochemistry, and community
1081   dynamics. J Coast Res 40:79-93.
1082   Twilley RR, Rivera-Monroy V (2009) Ecogeomorphic models of nutrient biogeochemistry for
1083   mangrove wetlands. In: Coastal Wetlands: An integrated ecosystem approach (eds)
1084   Perillo GME, Wolanski E, Cahoon DR and Brinson MM. Elsevier, pp 641-683.
1085   Twilley RR, Rivera-Monroy V, Rovai AS, Castañeda-Moya E, Davis SE, III. (2019) Mangrove
1086   biogeochemistry at local to global scales using ecogeomorphic approaches. In: Perillo
1087   GME, Wolanski E, Cahoon DR, Hopkinson CS (eds) Coastal Wetlands: An Integrated
1088   Ecosystem Approach.Elsevier, pp 717-785.
1089   Twilley RR, Rivera-Monroy VH, Chen R, Botero L (1998) Adapting and ecological mangrove
1090   model to simulate trajectories in restoration ecology. Mar. Poll. Bull. 37:404-419.
1091   Uncles R, Gong W-K, Ong J-E (1992) Intratidal fluctuations in stratification within a mangrove
1092   estuary. In: Martens E, Jaccarini V (eds) The Ecology of Mangrove and Related
1093   Ecosystems. Springer, pp 163-171.
1094   Vitousek P (1982) Nutrient cycling and nutrient use efficiency. Am Nat 119:553-572.
1095   Wanless H (1998) Mangroves, hurricanes, and sea level rise. South Florida Study Group, The
1096   Conservancy, Naples, Florida
1097   Werner A, Stelzer R (1990) Physiological responses of the mangrove *Rhizophora mangle* grown
1098   in the absence and presence of NaCl. Plant Cell Environ 13:243-255.
1099   Wolanski E (1992) Hydrodynamics of mangrove swamps and their coastal waters.
1100   Hydrobiologia 247:141-161.
1101   Wooller M, Smallwood B, Scharler U, Jacobson M, Fogel M (2003) A taphonomic study of δ13C
1102   and δ15N values in Rhizophora mangle leaves for a multi-proxy approach to mangrove
1103   palaeoecology. Org Geochem 34:1259-1275.
1104   Yu M, Rivera-Ocasio E, Heartsill-Scalley T, Davila-Casanova D, Rios-López N, Gao Q. (2019)
1105   Landscape-level consequences of rising sea-level on coastal wetlands: saltwater
1106   intrusion drives displacement and mortality in the twenty-first century. Wetlands 39:1343-
1107   1355.
1108

 TABLES

1110 **Table 1.** Mean (± 1 SE) soil surface elevation, bedrock elevation, and soil depth, for open water,
1111 mangrove-island edge and center habitats in the fringe and interior scrub mangrove areas at
1112 TS/Ph-7 in southeastern Florida Coastal Everglades. Elevation measurements are referenced
1113 to the North American Vertical Datum 1988 (NAVD88). Letters denote statistically different
1114 groupings via Tukey HSD post-hoc test ($p < .05$).

1115

Mangrove Location	Habitat	Soil Surface Elevation (NAVD88, m)	Bedrock Elevation (NAVD88, m)	Soil depth (m)
Fringe	Open water	-0.63 ± 0.05^{C}	-1.84 ± 0.05^{A}	1.22 ± 0.05^{C}
	Island edge	-0.41 ± 0.03^{B}	-1.83 ± 0.02^{A}	1.43 ± 0.05^{BC}
	Island center	-0.14 ± 0.02^{A}	-1.86 ± 0.06^{A}	1.72 ± 0.05^{A}
Interior	Open water	-0.84 ± 0.02^{D}	-2.01 ± 0.15^{A}	1.18 ± 0.14^{C}
	Island edge	-0.49 ± 0.03^{B}	-1.78 ± 0.04^{A}	1.39 ± 0.04^{C}
	Island center	-0.15 ± 0.02^{A}	-1.77 ± 0.03^{A}	1.62 ± 0.03^{AB}

1116 **Table 2.** Seasonal variation in water levels, surface water and porewater salinity measured in
1117 mangrove-island habitats at scrub *R. mangle* dominated mangroves at TS/Ph-7 in southeastern
1118 Florida Everglades. Means (± 1 SE) with different letters within each column denoting
1119 significant differences among groups (Tukey HSD post hoc, $p < .05$).

1120

Season	Habitat	Water level (cm)	Surface water salinity (ppt)	Porewater salinity (ppt)
Dry	Edge	33.5 ± 1.9^A	16.11 ± 1.16^A	24.22 ± 0.44^A
	Center	10.1 ± 1.9^B	21.14 ± 1.51^B	20.83 ± 0.44^B
Wet	Edge	40.2 ± 1.9^C	15.31 ± 1.16^A	26.00 ± 0.44^C
	Center	15.5 ± 1.9^B	14.41 ± 1.47^A	22.22 ± 0.44^D

Table 3. Leaf functional traits, carbon and nutrient contents and N:P ratios, nitrogen and
phosphorus resorption efficiencies, bulk isotopic signatures, intrinsic intracellular CO_2
concentrations (c_i) and intrinsic water use-efficiency (*WUE*) (calculated from [13]C fractionation)
for scrub *R. mangle* leaves collected from mangrove-island habitats at TS/Ph-7 during the dry
and wet seasons of 2019. Means (± 1 SE), with different letters across each row denoting
significantly different groups (Tukey HSD test, *p* < .05).

Leaf Trait	Dry season		Wet season	
	Edge	Center	Edge	Center
Leaf dry mass (g)	0.64 ± 0.02^A	0.65 ± 0.02^A	0.69 ± 0.02^A	0.70 ± 0.02^A
Leaf area (cm^2)	24.9 ± 0.7^A	25.5 ± 0.7^A	26.0 ± 1.4^A	27.3 ± 0.9^A
SLA ($cm^2\ g^{-1}$)	28.97 ± 0.66^A	39.79 ± 0.68^A	37.33 ± 0.40^A	40.04 ± 1.71^A
LWC (%)	65.7 ± 0.4^A	65.4 ± 0.5^A	63.6 ± 0.5^A	64.0 ± 0.8^A
Total C ($mg\ g^{-1}$)	450.9 ± 5.7^A	441.4 ± 1.0^A	428.7 ± 17.7^A	444.0 ± 1.0^A
Total N ($mg\ g^{-1}$)	9.8 ± 0.2^{AB}	10.2 ± 0.3^A	8.4 ± 0.5^B	9.0 ± 0.3^{AB}
Total P ($mg\ g^{-1}$)	0.50 ± 0.01^{AB}	0.55 ± 0.04^A	0.42 ± 0.01^B	0.46 ± 0.02^{AB}
Leaf N_{area} ($mg\ cm^{-2}$)	0.258 ± 0.012	0.264 ± 0.019	0.263 ± 0.035	0.306 ± 0.020
Leaf P_{area} ($mg\ cm^{-2}$)	0.013 ± 0.001	0.014 ± 0.002	0.013 ± 0.001	0.016 ± 0.001
Atomic N:P	41.7 ± 1.9	40.0 ± 0.1	43.5 ± 3.7	42.5 ± 2.1
N resorption (%)	60.0 ± 0.4	62.8 ± 0.5	60.8 ± 2.8	62.9 ± 1.5
P resorption (%)	78.6 ± 0.2	74.3 ± 3.2	75.5 ± 0.4	73.2 ± 6.2
$\delta^{13}C$ (‰)	-25.5 ± 0.1^{AB}	-25.1 ± 0.1^A	-25.8 ± 0.1^B	-25.9 ± 0.2^{AB}
c_i ($\mu mol\ mol^{-1}$)	228.1 ± 2.2^{AB}	219.9 ± 1.8^B	233.3 ± 1.1^A	235.3 ± 3.6^A
WUE ($mmol\ mol^{-1}$)	0.1124 ± 0.0014^{AB}	0.1175 ± 0.0012^B	0.1092 ± 0.0007^A	0.1080 ± 0.0022^A
$\delta^{15}N$ (‰)	-5.3 ± 0.5^B	-0.4 ± 0.4^A	-4.2 ± 1.0^{AB}	-0.8 ± 1.5^A

1127

FIGURES LEGENDS

Figure 1. A) Photograph of TS/Ph-7 shows scrub R. mangle tree islands that characterize the study site. Mangrove canopy heights are approximately 1.5-2 m tall, facilitating canopy measurements of leaf physiology. Boardwalk (1.3 m height) is pictured for reference. **B)** Aerial view (Google Earth) of mangrove-islands measured for this study within TS/Ph-7, near the mouth of the Taylor River in southeastern Florida Coastal Everglades, USA. The inset shows the location of TS/Ph-7 within the boundary of Everglades National Park. Colors indicate scrub mangroves and fringe (white) and interior zones (yellow) relative to Taylor River. Symbols denoted paired higher-elevation center and lower-elevation edge habitats for each mangrove-island (squares and triangles, respectively).

Figure 2. Predicted marginal mean (±95% confidence intervals) values of photosynthesis (A_{net}), stomatal conductance (g_{sw}), the concentration of intracellular CO_2 (c_i), and intrinsic water use efficiency (*wue*) by mangrove-island habitat and season. The dry season is from November to April, and the wet season is from May to October. See supplemental material for complete model summaries.

Figure 3. The effect of water level on leaf photosynthesis (A_{net}) and stomatal conductance (g_{sw}), the concentration of intracellular CO_2 (c_i), and intrinsic water use efficiency (*wue*) by season. Lines are habitat-specific predicted mean marginal mean values (± 95% confidence intervals) from linear mixed-effects models.

Figure 4. The effect of soil porewater salinity on leaf photosynthesis (A_{net}) and stomatal conductance (g_{sw}) by season. Porewater salinity was not included in the best-fit models for c_i or *wue*. Lines are predicted mean marginal effects from linear mixed-effects models ± 95% confidence intervals (colored by island habitat).

Figure 5. Mean (± 1 SE) leaf isotopic signatures and nutrient resorption efficiency by island habitat and season combination. **A)** the relationship between $\delta^{15}N$ and $\delta^{13}C$ in *R. mangle* leaves, **B)** the relationship between N resorption efficiency and $\delta^{13}C$ for *R. mangle* leaves, and **C)** the relationship between P resorption efficiency and $\delta^{13}C$ for R. mangle leaves. Error bars colors denote island habitats, while point symbols and colors show seasons.