1 **High contrast, moving targets in an emerging target paradigm promote fast visuomotor**
2 **responses during visually guided reaching**
3
4
5 Abbreviated title: Moving targets promote fast visuomotor responses
6

7 Rebecca A. Kozak[1,4], Brian D. Corneil[1,2,3,4]
8
9 [1]Graduate Program in Neuroscience, Western University, London, On, Canada, N6A 5B7
10 [2]Department of Psychology, Western University, London, Ontario, Canada, N6A 5B7
11 [3]Department of Physiology and Pharmacology, Western University, London, Ontario, Canada,
12 N6A 5B7
13 [4]Robarts Research Institute, 1151 Richmond St. N, London, Ontario, Canada, N6A 5B7
14
15 *Corresponding Author: Brian D. Corneil, bcorneil@uwo.ca

16
17
Number of pages: 39
Number of figures: 10
Number of tables: 0
Abstract word count: 235
Introduction word count: 763
Discussion word count: 1985
24

25 Conflicts of interest: The authors declare no competing financial interests
26 Acknowledgements: We thank Dr. Andrew Pruszynski for use of his Kinarm exoskeleton for
27 Experiment 1.
28
This work is supported by a Discovery Grant to BDC from the Natural Sciences and Engineering
Research Council of Canada (NSERC; RGPIN 311680) and an Operating Grant to BDC from the
Canadian Institutes of Health Research (CIHR; MOP-93796). RAK was supported by an Ontario
Graduate Scholarship. The apparatus used in Experiment 2 was purchased using funds from the
Canadian Foundation for Innovation. Additional support came from the Canada First Research
Excellence Fund (BrainsCAN).

Keywords: visually-guided reaches, EMG, SLR, moving targets, contrast, reaction time, RT

Abstract

Humans have a remarkable capacity to rapidly interact with the surrounding environment, often

by transforming visual input into motor output on a moment-to-moment basis. But what visual

features promote the shortest-latency reach responses? To address this question, we had human

subjects perform visually guided reaches to moving targets varied by speed (experiment 1), or

speed and contrast (experiment 2) in an emerging target paradigm, which has recently been

shown to robustly elicit fast visuomotor responses. Our analysis focused on stimulus-locked

responses (SLRs) on upper limb muscles. SLRs represent the first wave of muscle recruitment

tied to visual target onset, appearing within <100 ms. Across 32 subjects studied in both

experiments, 97% expressed SLRs in the emerging target paradigm. In comparison, 69% of these

subjects expressed SLRs in a visually-guided reach paradigm. Within the emerging target

paradigm, we found that target speed impacted SLR magnitude, whereas target contrast impacted

SLR latency and magnitude. Thus, high contrast, faster-moving targets in the emerging target

paradigm robustly recruited the circuitry mediating the most rapid visuomotor transformations

for reaching, and such responses were associated with shorter latency RTs. Our results support

the hypothesis that a subcortical pathway originating in the superior colliculus may be involved

in the earliest wave of muscle recruitment following visual stimulus presentation. In scenarios

requiring expedited responses, cortical areas may serve to prime this subcortical pathway, and

elaborate subsequent phases of muscle recruitment following the SLR.

Significance Statement

Humans have a remarkable capacity, when necessary, to rapidly transform vision into action. But

how does the brain do this? Here, by studying human subjects reaching to suddenly-appearing

targets, we find that the earliest visually-guided actions are produced in response to high-

contrast, moving targets. A millisecond-resolution examination of upper limb muscle recruitment

shows that motor output can begin within less than 100 ms of target presentation. We surmise

that this earliest recruitment arises from a phylogenetically-conserved brainstem circuit

originating in the superior colliculus. Rather than being directly involved in the earliest phase of

visuomotor actions, cortical areas may prime this brainstem circuit to produce initial muscle

recruitment, and then elaborate subsequent phases of recruitment when time is of the essence.

Introduction

To perform a visually-guided reach, for example to pick-up a baseball off a table, visual information such as the shape and size of the ball is transformed into motor commands which are communicated to the motor periphery. However, if the ball is thrown to us, time is of the essence and we must perform a rapid reach in order to catch the ball. In such situations, the latency of the response depends on the visual attributes of the stimulus; for example, earlier responses are elicited by larger or higher contrast targets, or stimuli defined by lower spatial frequencies (Veerman et al., 2008; Wood et al., 2015; Kozak et al., 2019). Other work has suggested that the most rapid visuomotor transformation for reaching is mediated by a tecto-reticulo-spinal pathway, which lies in parallel to the well-studied corticospinal pathway (Day and Lyon, 2000; Pruszynski et al., 2010; Gu et al., 2016; Glover and Baker, 2019).

In order to better understand the most rapid visuomotor transformation at the limb, we and others have recorded electromyographic (EMG) activity from upper limb muscles during rapid visually-guided actions. Doing so has led to the identification of stimulus-locked responses (SLRs; also termed rapid visuomotor responses by (Glover and Baker, 2019) or express responses by (Contemori et al., 2020)), which are the first waves of muscle activity influenced by the visual stimulus. SLRs evolve within < 100 ms of stimulus onset, preceding some response latencies in visual and motor cortices (Maunsell and Gibson, 1992; Schmolesky et al., 1998; Lara et al., 2018). In contrast, visual responses latencies in the superior colliculus (SC; from which the tecto-reticulo-spinal track originates) precede the SLR (Stuphorn et al., 1999; Rezvani and Corneil, 2008), which makes the SC a potential candidate for mediating the SLR. Further, high contrast (Li and Basso, 2008; Marino et al., 2012; Wood et al., 2015) or low spatial frequency (Chen et al., 2018; Kozak et al., 2019) targets elicit larger-magnitude and shorter-latency visual

responses in the SC and SLRs. Robust SC responses are also elicited by moving visual targets

(Marrocco and Li, 1977); however, a systematic investigation of target motion on the SLR has

not been performed.

Previous research investigating the most rapid visuomotor transformations involved in

visually guided reaching has typically relied on targets that either suddenly appear (a *static target*

paradigm; (Pruszynski et al., 2010)), or are displaced during a reach (*target jump paradigm*;

(Soechting and Lacquaniti, 1983; Day and Lyon, 2000)). Recently, we have presented an

emerging target paradigm, where a target travels down an inverted 'Y' path before disappearing

and then emerging from behind an occluder (Kozak et al., 2020). Benefits of the emerging target

paradigm include the stability of the arm when the target emerges (which is not the case in the

target jump paradigm), and the generation of robust fast visuomotor responses in almost every

subject (which is not the case in the static target paradigm). In Kozak et al. (2020), while only 3

of 5 subjects expressed an SLR in the static target paradigm, all 5 subjects expressed an SLR in

the emerging target paradigm, with the SLR being four to five times larger in the emerging target

paradigm. The efficacy of the emerging target paradigm in generating SLRs has been replicated

in a larger sample of subjects using the emerging target paradigm, wherein 20 of 21 subjects

expressed an SLR (Contemori et al., 2020). Thus, essentially every subject has the circuitry to

generate fast visuomotor responses; the failure to do so in past studies arose from the reliance on

suboptimal stimuli and/or tasks.

The goal of the current study is to systematically investigate how target speed and

contrast impact fast visuomotor responses in the emerging target paradigm. Based on the

hypothesis that transient visual responses in the SC initiate the SLR, larger and more robust

SLRs should accompany the presentation of fast moving, high-contrast targets. Accordingly, we

tested the influence of targets varied by speed (experiment 1) or speed and contrast (experiment

2) on the SLR, in both static and emerging target paradigms. A secondary goal is to better

understand how the emerging target paradigm promotes the generation of SLRs. Hence, we also

presented non-moving stimuli in the context of both the static target and emerging target

paradigms in Experiment 1. In such cases, the retinal stimulus is exactly the same, meaning that

any differences in the SLR arise from differences related to the behavioural task. Overall, we

find that high contrast, fast moving targets in the emerging target paradigm robustly recruits the

circuitry mediating the most rapid visuomotor transformations for reaching movements.

Materials and methods

Subjects

A total of 34 subjects (18 females, 16 males; mean age: 24.2 years SD: 5.6) completed at least

one of the two experiments. Two subjects completed both experiments. All subjects provided

informed consent, were paid for their participation, and were free to withdraw from the

experiment at any time. All but 1 subject was right-handed. All subjects had normal or corrected-

to-normal vision, with no current visual, neurological, or musculoskeletal disorders. All

procedures were approved by the Health Science Research Ethics Board at the University of

Western Ontario. Two subjects (1 from each experiment) were excluded due to low quality EMG

recordings.

Apparatus

Subjects were seated in either a KINARM robotic Exoskeleton (experiment 1) or a KINARM

End-Point Lab (experiment 2) (KINARM Technologies, Kingston, ON, Canada), and performed

reaching movements with their right arm. Both platforms relied on the same experimental

software. Visual stimuli were projected onto an upward facing mirror from either a downward

facing monitor in experiment 1 (LG 47LS35A; size: 47" resolution: 1920x1080 pixels

(Exoskeleton)) or custom built in projector in experiment 2 (PROPixx projector by VPixx, Saint-

Bruno, QC, Canada; custom integrated into the KINARM End-Point Lab). In both experiments,

a shield below the mirror occluded direct vision of the hand, but real-time hand position was

represented on the monitor via a real-time cursor projected onto the screen. A photodiode was

used to indicate the precise time of visual stimulus presentation, and all kinematic and

electromyographic (EMG) data were aligned to photodiode onset. Eye movements were not

measured.

Experimental design:

In experiment 1 and 2, we used an emerging target paradigm as previously described in ((Kozak

et al., 2020) see **Figure 1**). Briefly, a target travels down an inverted 'y' path, disappears behind

an occluder before reappearing, continuing to move, and evoking a reach from a subject. In both

experiments, we also collected data in a *static task*, where subjects performed visually guided

reaches towards stationary targets in a separate block of trials. This allowed us to directly

compare SLRs in the emerging target paradigm to those obtained with a static task typically used

to evoke the SLR (Pruszynski et al., 2010; Wood et al., 2015; Gu et al., 2016; Kozak et al.,

2019).

Experiment 1: effects of speed.

In the static task in Experiment 1, subjects ($n = 16$) performed visually guided reaches from a

stationary arm position to static targets appearing pseudorandomly to either the right or left of

the start position. Throughout this task, a constant torque of 2 Nm was applied to the shoulder,

increasing tonic activity on the muscle of interest. Doing so allows for the SLR to be expressed

as either an increase or decrease in muscle activity following target appearance to the left or

right, respectively. Each trial was initiated when a subject moved the real time cursor (RTC)

representing their hand into a start position (T0). T0 was 1.5 cm in diameter, located

approximately 45 cm in front of the subject relative to midline, and changed to grey once the

RTC was in the start position. If the subject exited the start position before completion of a

constant hold period (1s), the trial was reset, and T0 changed back to white. Subjects were

instructed to also look at T0 before the onset of the target (T1). After the hold period, a white T1

was presented on the black background 17 cm to the left or right of the start position (~17.5

degrees of visual angle). A 17 cm eccentricity was chosen as it resembled the interception point

of the emerging target paradigm. Subjects were instructed to reach towards the peripheral target

as quickly as possible, and the trial ended when the RTC made contact with the center of T1. T0

remained visible as a grey target, changing back to white once the trial ended. Subjects

completed 1 block of 220 trials (110 trials towards each target), before they performed the

emerging target paradigm. In contrast to previous static tasks which used a randomized hold

delay for T0 (Gu et al., 2016; Gu et al., 2018), we chose to have a constant hold delay (1s), and

allowed T0 to remain visible during the reach. These conditions more closely resembled the

conditions in the emerging target task which allowed for a more direct comparison between a

static task and an emerging target task.

After completing the static task, the same 16 subjects performed visually guided reaches

towards moving targets that emerged below an occluder in the emerging target paradigm (See

Figure 1 for protocol). The emerging target paradigm involves a target starting at the top of an

inverted y (**Figure 1a**), moving towards an occluder and disappearing behind it (**Figure 1b**), then

emerging below the occluder at one of two outlets. Subjects were instructed to reach and

intercept the target after it emerged from behind the occluder (**Figure 1c**) (Kozak et al., 2020). In

experiment 1, we varied the speed of the moving target. The start position, RTC, hold period,

background loading torque, and target radius were the same as in the static task.

We implemented three target speeds in our paradigm (10, 15 or 20 cm/s), translating to

initial velocities of approximately 10, 15 or 20 retinal degrees per second, respectively. These

speeds were chosen to span a range from 'slow' to 'fast' while still permitting target interception

of the fastest target. The duration that T1 was visible above the occluder was constant (.5s)

regardless of target speed; as targets travelled at different speeds, this meant that they started at

different heights above the occluder. The occluder was 30 cm wide by 15 cm high, and the center

of the occluder was located ~63 cm away from the subject. Behind the occluder, the target

moved at a constant speed of (15 cm/s) regardless of initial or final target speed. Thus, the target

consistently disappeared behind the occluder for 1s on trials with a moving target. The target

only became visible once it had fully moved past the occluder (i.e., the target was presented in its

entirety, rather than sliding past the occluder to avoid a half-moon stimulus). After emergence,

the target stayed visible and usually continued down the inverted y path. To control for eye

position, subjects were instructed to foveate a notch located on the bottom of the occluder

halfway between the exit points, while the moving target was occluded. The trial ended as soon

as the RTC contacted the peripheral target, or if the target passed the bottom of the screen.

We also included a static T1 within the emerging target paradigm, where T1 began

moving at a speed of 20 cm/s but appeared ~ 17 cm to the left or right of the start position and

did not move; we term this the 0 cm/s target. Thus, the initial conditions where the target drops

209 into the occluder were the same for the 0 cm/s and 20 cm/s targets. On the 0 cm/s trial, the trial

210 ended when the subjects made contact with the target. Due to an oversight, the target disappeared

211 behind the barrier for 900 ms on 0 cm/s trials, rather than 1000 ms as on moving target trials. If

212 the slightly shorter time the target was obscured behind the barrier on 0 cm/s trials were to be

213 relevant, then results by Contemori and colleagues (2020) would suggest that SLR should be

214 weaker on such trials given less temporal certainty. Instead, as we will show, we observed robust

215 SLRs to both moving and non-moving targets in the emerging target task. Further, no subject

216 reported perceiving any difference in the time the target was behind the occluder on trials with a

217 moving versus a non-moving target.

218  All targets in experiment 1 were white (110 cd/m^2) presented against a black (.6 cd/m^2)

219 background (contrast ratio: 183:1). There were 8 unique trial conditions (4 target speeds 0, 10, 15

220 and 20 cm/s presented to the left and right). Subjects completed 2 blocks of 480 trials, with each

221 block containing 60 repetitions of each unique trial condition presented pseudorandomly without

222 replacement, yielding a total of 120 trials for each unique trial condition.

224 *Experiment 2: effects of contrast and speed*

225 Experiment 2 was designed to investigate the interaction between target speed and contrast on

226 the SLR. This experiment was conducted in a different apparatus, the KINARM End-Point

227 system. As in experiment 1, two tasks were run in separate blocks.

228  In the *static task* in experiment 2, subjects ($n = 18$) performed visually guided reaches

229 initiated from a stationary position, towards static left or right targets. As in experiment 1, each

230 trial was initiated when a subject moved the real time cursor (RTC) representing the subjects

231 hand position into a start position (T0). We used a constant torque throughout the experiment

(2Nm down, 5Nm right) applied to increase the baseline activity of the muscle of interest. This

torque is different than experiment 1, due to a different apparatus and subject posture. T0 was

aligned with the subject's midline (0, -42 cm relative to the midpoint between the robot

manipulanda; radius 1 cm). T0 changed from white to grey once the RTC was moved into it. The

trial was reset if the subject exited T0 before completion of the hold period (randomized time

between 1000 to 2000 ms). Our use of a randomized hold time in experiment 2 resembles that

used in the work by Wood and colleagues (2015), which also manipulated stimulus contrast.

Recall that the static task in experiment 1 used a fixed hold time of 1 s, which matched the time

the target disappeared behind the barrier in the emerging target paradigm in experiment 1.

Subjects were instructed to foveate T0 before the onset of the target (T1). After the hold period,

T1 (target with a 1 cm radius) appeared 10 cm to the left or right (~15.4 degrees of visual angle).

Subjects were instructed to reach towards the peripheral target as quickly as possible, and the

trial ended as soon as RTC made contact with the peripheral target. There were 4 unique trial

conditions; a high contrast white (275 cd/m^2: contrast ratio 793.8:1) or low contrast grey (18

cd/m^2: contrast ratio 50:1) target presented against a black (.36 cd/m^2) background to either the

left or right side. T0 remained visible as a grey target, changing back to white once the trial

ended. Subjects completed 1 block of 400 pseudorandomized trials, yielding a total of 100 trials

for each unique trial condition.

After completing the static task, the same 18 subjects completed the emerging target

paradigm. As in experiment 1, subjects performed visually guided reaches initiated from a stable

posture towards moving targets that emerged suddenly beneath an occluder. Throughout the

experiment, the same torque was applied as used in the static task. Each trial was initiated when

a subject moved the RTC into a start position (T0), and the trial was reset if the subject exited T0

before completion of the hold period (1s). In experiment 2, we used a two-by-two design, where

we modulated speed (including a 'slow' and a 'fast' target (10 cm/s, 20 cm/s; translating to

approximately 10 and 20 degrees of visual angle); and contrast (the 'low' and 'high' contrasts).

In experiment 2, the occluder was 30cm by 15cm, centered at the midpoint between the

manipulanda (0, - 29 cm). To control for eye position, subjects were instructed to foveate a notch

located on the bottom of the occluder halfway between the exit points, while T1 was occluded.

As in experiment 1, we controlled for the amount of time that T1 was visible before being

obscured by the occluder (.5s) by starting T1 at different heights above the occluder for different

target speeds. The time T1 disappeared behind the occluder in experiment 2 is shorter (0.5 s)

than in experiment 1 (1 s). T1 was made visible only after it had fully emerged beneath the

occluder, avoiding an initial half-moon stimulus. T1 then continued on the inverted y path. The

trial ended as soon as the RTC made contact with T1 or if the target passed the bottom of the

screen. There were 8 unique trial conditions (2 speeds, 2 contrasts, leftward or rightward targets).

The same contrasts were used from the static task. Subjects completed 2 blocks of 400

pseudorandomized trials, yielding a total of 100 trials for each unique trial condition.

Data acquisition and analysis

Surface electromyographic (EMG) activity was recorded from the right pectoralis muscle with

double-differential surface electrodes (Delsys Inc. Bagnoli-8 system, Boston, MA USA). As

stated in (Kozak et al., 2020), hair or excess adipose may compromise the EMG signal;

therefore, we targeted either the medial or lateral portion of the clavicular head, or the sternal

portion, depending on such idiosyncratic factors. We placed two electrodes on each subject, and

during post-hoc analysis we used the recording exhibiting the higher level of EMG activity.

EMG signals were sampled at 1 kHz, amplified by 1000, full-wave rectified off-line, and

smoothed with a 7-point smoothing function. We excluded two subjects with poor quality muscle

recordings, which was defined as an increase in peak EMG activity above baseline of less than

40 μV during leftward reaching movements.

Kinematic data were sampled at 1 kHz by the Kinarm data acquisition system in both

experiments. Reaction time (RT) was calculated as time from peripheral target (T1) appearance

(measured by the photodiode) to reach initiation. Reach initiation in both experiments was

defined as 8% of the peak tangential velocity. We excluded trials with RTs less than 120 ms due

to presumed anticipation, and also excluded trials with RTs exceeding 500 ms due to presumed

inattentiveness.

As described previously (Kozak et al., 2020), we used a receiver-operating characteristic

(ROC) analysis to define the presence and latency of the SLR. Briefly, the time-series ROC

analysis indicates the likelihood of discriminating stimulus presentation based on EMG alone. As

the SLR is the first wave of muscle activity influenced by the visual stimulus, we rely on this

method to discriminate the moment a visual stimulus influences EMG activity. The ROC value

(i.e., the area under the ROC curve) was calculated and plotted at every time sample (1ms) from

100 ms before to 300 ms after target presentation (an example may be seen in **Figure 2d** overlaid

with mean +/- SE of EMG activity for the data shown in **Figure 2a**). A ROC value of .5

indicates chance discrimination, whereas values of 1 or 0 indicate perfectly correct or incorrect

discrimination relative to target presentation, respectively. Discrimination thresholds were set to

0.6, and the discrimination latency was determined as the first of 8 of 10 consecutive points that

exceeded this value.

To define the presence of an SLR, an RT split analysis was used (as described in: (Wood

et al., 2015; Kozak et al., 2020)). The purpose of the RT split analysis is to empirically test

whether EMG activity is more likely to evolve with stimulus presentation, or arm movement

over a large number of trials. This method is used as an additional conservative approach to

relying on pooled EMG, which may be unfairly biased towards a subset of the shortest latency

trials. To perform the RT split analysis, separate time series ROC analyses were conducted on

early and late RT trials (purple or green highlighted trials in **Figure 2a**; see also mean EMG

traces in **Figure 2b**), defined relative to mean RT. We then plotted the mean RT for the early or

late RT subsets versus the ROC discrimination time, and then connected these two points with a

line (**Figure 2c**). The slope of this line is used to determine whether the initial change in EMG

activity is more locked to stimulus onset (slope $\geq$ 67.5 deg.) or not (slope < 67.5 deg.). The

example shown in **Figure 2** illustrates an example where an SLR was detected (termed SLR+;

versus SLR-).

We used the same criterion for determining latency and magnitude of the SLR as used in

(Kozak et al., 2020). For SLR+ observations, SLR latency was defined as the first point where

the time-series ROC constructed from all trials (regardless of RT) surpassed .6 for 8 out of 10

trials (see **Figure 2d**). SLR magnitude was defined as the area between leftward and rightward

mean EMG traces for an interval spanning from SLR latency to 30 ms later.

Statistical analyses

Statistical analyses were performed in MATLAB (version R2016a, The MathWorks, Inc.,

Natick, Massachusetts, United States). Chi-squared tests were used to determine differences in

the proportion of subjects exhibiting an SLR (termed SLR prevalence). To compare SLR

latencies and magnitudes, we used a one-way ANOVA for experiment 1 and a 2 by 2 ANOVA

for experiment 2. Post-hoc pairwise comparisons relied on the Tukey-Kramer critical value, as

well as paired and unpaired t-tests when appropriate. We relied on non-normalized EMG for our

magnitude analysis. When comparing SLR magnitudes between subjects, all SLR+ values were

normalized to a maximum SLR magnitude within participants. For example, if a subject

exhibited SLRs in 5 conditions, the largest SLR condition for that subject is given a value of 1

and all other magnitudes are presented as a fraction of 1. When comparing SLR magnitudes to

recruitment associated with voluntary movement, we divided the magnitudes attained from the

SLR epoch (non-normalized) by the sum of the movement-aligned activity with a 30 ms window

surrounding peak EMG activity. Correlational analyses were performed on latency, magnitude

and RT measures on a subject-by-subject basis.

Results

An emerging target paradigm increases SLR prevalence

The emerging target paradigm is highly effective at eliciting stimulus-locked responses (SLRs)

and other fast visuomotor responses, such as short-latency reach reaction times (RTs)(Kozak et

al., 2020). Here we sought to investigate the features of the stimulus which best elicit SLRs by

altering the speed (experiment 1) or the speed and contrast (experiment 2) of the target. We first

determined how effective a given target condition was at eliciting an SLR.

All 32 subjects exhibited an SLR in at least one condition in both experiment 1 and 2

(**Figure 3;** recall 2 subjects performed in both experiments); as can be seen in this figure, each

subject had at least one diamond above the dotted line, indicating a detectable SLR. Previous

studies have reported that less than 65% of subjects exhibit an SLR to static high contrast targets

(Wood et al., 2015). In our sample, we found that 31 (97%) exhibited an SLR to a moving target

condition, whereas only 22 (68.7%) exhibited an SLR in the static target task. These differences

in detection are statistically significant (chi-squared test: p= 0.0189, $\chi 2$= 5.51, df= 1), and

reaffirm previous observations that subjects not exhibiting an SLR to static targets almost always

exhibit an SLR to targets in the emerging target task (Kozak et al., 2020).

We further investigated whether SLR prevalence was influenced by target speed and/or

contrast. In experiment 1, the target could move at three different speeds; 'slow' (10 cm/s),

'medium' (15 cm/s), or 'fast' (20cm/s). We found that stimulus speed did not influence the

probability of detecting an SLR (10 v. 15 v. 20 cm/s- chi-squared, all p> .05). Therefore,

although subjects are more likely to exhibit SLRs in response to moving versus static targets, the

speed at which the target moves does not increase the likelihood of detection.

Recall that in experiment 1, we presented a static target either in the emerging target task

(i.e., the 0 cm/s target, which was intermixed with other moving targets), or in a standard static

task used previously (S1). We found that all 16 subjects (100%) exhibited an SLR to a static

target in the emerging paradigm, but only 9 of these subjects (~56%) exhibited an SLR in the

standard, static task (S1). This difference in prevalence was statistically significant (chi-squared

test: p= 0.00275, $\chi 2$= 8.96, df=1), indicating that an emerging target paradigm promotes the

expression of an SLR regardless of whether the target is moving or not.

In experiment 2, we investigated whether the contrast and speed of a target impacted SLR

expression. As shown in experiment 1, we confirmed that target speed does not impact SLR

prevalence, therefore we grouped together both fast- and slow- moving target conditions and

determined that a high contrast moving target is more likely to evoke the SLR than a low

contrast moving target (chi-squared test: p= 0.00075, $\chi 2$= 11.361, df=1). In fact, low contrast

moving targets were not significantly better at evoking SLRs than static targets (slow + fast low

contrast moving targets versus high + low contrast static targets: chi-squared test: p>.05, χ^2=

.056, df=1). These results demonstrate that high contrast, moving targets are better at evoking

SLRs than static or low contrast targets, furthermore, low contrast targets are equally as effective

as static targets.

Next, we sought to characterize the latency and magnitude of the SLR. As can be seen in

representative data from experiment 1 (**Figure 4**) and experiment 2 (**Figure 5**), various SLR

features changed with different target conditions. In these plots, the SLR appears as a vertical

banding of EMG activity, due to EMG changes occurring around the same time after target

emergence on every trial (grey boxes, two left columns in **Figure 4** and **Figure 5**). **Figure 4** and

5 also shows the mean +/- SE EMG traces and time-series ROC plots (3rd column), and a

depiction of how the presence of an SLR is detected (see methods- RT split analysis; slopes

>67.5 indicate an SLR is present). In **Figure 4**, the representative subject from experiment 1

exhibited SLRs in all conditions; note the vertical banding in the color plots, and the slopes of

the lines in the rightmost plots. SLRs in this subject began ~90 ms after target emergence in all

conditions, regardless of target speed, or whether a stationary stimulus was presented in the

emerging target task or static task. In contrast, the representative subject from experiment 2

(**Figure 5**) did not exhibit an SLR in the static task (rightmost columns, slope <67.5) but did

exhibit an SLR to both high and low contrast moving targets in the emerging target task. Data

from this subject illustrates an example of an SLR following target appearance in the emerging

target but not static task (compare first two rows to final four rows). In the following sections,

we further investigate the impact of varying target speed and contrast on the latency and

magnitude of the SLR.

Experiment 1: Faster moving targets elicit larger magnitude SLRs that evolve at similar

latencies

We have previously reported that SLRs may differ in magnitude (Gu et al., 2016) and/or latency

(Kozak et al., 2019) depending on task context and stimulus properties, respectively. We

examined the data from experiment 1 to investigate the effect of target speed on SLR latency and

magnitude. There were 5 different conditions associated with experiment 1: either a static target

appearing in the static task (S1), or static (0cm/s), 'slow' (10 cm/s), 'medium' (15 cm/s) or 'fast'

(20 cm/s) targets in the emerging paradigm. Intriguingly, as seen in (**Figure 6a**), there were no

differences in SLR latency in any of the conditions (one-way ANOVA, F(4,53) = .58, p = .68).

These results are also apparent in the individual subject data shown in **Figure 4**; as can be seen

in the trial-by-trial colour plots on the left-hand side, irrespective of the condition, the banding of

activity defining the SLR evolves consistently at ~90 ms.

Although it is clear that the latency for the SLR is not affected by target speed, we

observed clear evidence that SLR magnitude is impacted by target speed (**Figure 6b**). From the

left side of **Figure 6b** (S1) to the right side of the subplot (fastest moving targets), magnitude

values are increased for moving targets (one-way ANOVA (all conditions), F(4, 53)= 15.15, p=

2.53e-8), with all emerging targets evoking larger magnitude SLRs than static targets (all

p<.0005). Indeed, SLRs evoked by the fastest moving target in the emerging target paradigm

were approximately twice as large as SLRs evoked by the target in the static paradigm (S1 versus

20 cm/s; unpaired t-test; t= -7.78, p= 3.61e-7, df= 18). Even when removing the influence of

both static targets from our analysis (s1 and 0cm/s), we found that faster moving targets in the

emerging target paradigm promote larger magnitude SLRs (one-way ANOVA (10, 15 and 20

cm/s speeds), F(2, 32)= 4.59, *p*= .0183).

While SLR magnitude tends to increase for faster moving targets, **Figure 6b** shows that

SLR magnitudes were comparable for the fastest (20 cm/s) and the non-moving (0 cm/s) target in

the emerging target paradigm. Recall as well that the 0cm/s target presented in the emerging

target paradigm evoked SLRs more often than targets in the static paradigm. These results (large

magnitudes and high prevalence) were very interesting to us given that in both conditions the

stimulus presented to the visual system is the same (blue rectangle in **Figure 6**). A direct

comparison showed that 0 cm/s targets in the emerging target paradigm evoked larger magnitude

SLRs than target in the static paradigm (unpaired t-test; t= -6.1, p= 3.02e-6, df= 23). Thus, the

behavioural context in which a static target is presented influences both the prevalence and

magnitude of the SLR, but not the latency.

We were also interested in how SLR magnitude related to the maximal movement

aligned activity. Therefore, we took the magnitude of the EMG activity in SLR epoch and

compared it to movement aligned magnitude for a 30 ms window around the peak (this is the

same duration used for the SLR magnitude). As shown in **Figure 6c**, we observed substantial

variability in how SLR magnitudes compared with the peak EMG activity aligned to movement

onset, with SLRs being anywhere between ~10-90% as large. We did not detect any differences

in this scaled measure of SLR magnitude across conditions (one-way ANOVA, F(4,53)=.62, *p* =

.65). On average, SLRs tended to be ~50% as large as the activity aligned to movement onset.

Despite such large variability, the SLR magnitudes reported here are significantly larger than any

reported previously (Pruszynski et al., 2010; Gu et al., 2016; Kozak et al., 2019).

Overall, when using targets moving at different speeds in an emerging target paradigm,

all moving targets are equally capable of eliciting SLRs and do so at similar latencies. However,

faster moving targets are associated with larger magnitude SLRs. An emerging target paradigm

was also more effective than a static target paradigm at eliciting more prevalent and larger

magnitude SLRs. Static targets presented in the context of an emerging target paradigm,

intermixed with moving targets, were equally effective at eliciting large magnitude and prevalent

SLR responses.

Experiment 2: Higher contrast targets evoke shorter latency and larger magnitude SLRs

The most intriguing finding in experiment 1 was that the latency of the SLR did not depend on

the speed of the target; all targets (whether statically presented or the fastest moving) evoked

SLRs at the same latencies. This was intriguing given that previous studies which systematically

varied visual features (contrast, spatial frequency) found that lower magnitudes accompany

longer-latency SLRs (Wood et al., 2015; Kozak et al., 2019). However, the results from

experiment 1 indicate that manipulations of stimulus features, and specifically stimulus velocity,

can influence SLR magnitude without influencing SLR latency. Therefore, we further probed

this finding by introducing targets varied by contrast, which have been shown to alter SLR

latency for static targets (Wood et al., 2015). In particular, we examined if SLR latency and

magnitude would negatively correlate across the speed of motion of lower-contrast targets. These

lower contrast targets would avoid possible 'floor effects' on SLR latency, which may have been

caused by the use of salient, high contrast moving targets in experiment 1. In experiment 2, we

examined the SLR in response to high and low contrast targets moving at a 'fast' (20 cm/s) or

'slow' (10 cm/s) speed.

We first examined the effect of target contrast on SLR latency. We found that the

contrast, but not the speed of the moving target, impacted the SLR latency (2 x 2 ANOVA,

$F(1,47) = 72.85$, $p = 4.1e-11$). High contrast targets evoked shorter latency SLRs by

approximately 14ms compared to low contrast targets (**Figure 7a**; $p = 2.07e-8$). This trend is

consistent with previous results (Wood et al., 2015). We also found that targets with the same

contrast, regardless of the speed they were moving, evoked SLRs of similar latencies ($p > .05$;

Figure 7a no differences within green points or blue points), which confirms results from

experiment 1. Thus, the same contrast targets evoke similar latency SLRs, regardless of how fast

they are moving. Thus, the results from experiment 1 were not simply due to a floor effect on

SLR latency, beyond which the SLR latency could not be further lowered.

In contrast to experiment 1, static targets in experiment 2 tended to be longer latency

compared to emerging targets, (**Figure 7a**; unpaired t-test; $t(18) = -3.27$, $p = .004$), but were

similar latencies to those reported previously (Wood et al., 2015). We contend that the most

likely reason for longer latency SLRs in the experiment 2 static target task is due to differences

in experimental set up between experiment 1 and 2, related to randomized hold time (see

discussion).

We next examined the influence of target contrast and speed on SLR magnitude (**Figure**

7b). Here we found that both the speed and the contrast impacts SLR magnitude (2 x 2 ANOVA,

speed: $F(1,47)=67.8$ $p = 1.1e-10$; contrast: $F(1,47)=11.7$ $p = .001$). Higher contrast targets

evoked larger magnitude SLRs compared to lower contrast targets (**Figure 7b**, all blue points

lower than all green points; unpaired t-test; t= 2.7, p= .01, df= 49). We also found that faster

moving targets evoked larger magnitude SLRs compared to slower moving targets (all second

points within a condition are higher than first points; unpaired t-test; t= -7.8, p= 4.1e-10, df= 49).

This supports the results from experiment 1, where faster moving targets evoked larger

magnitude SLRs. We did not observe interaction effects between target speed and contrast (2 x 2

ANOVA, p=.87). We also did not observe any differences in magnitude for static targets. Taken

together, higher contrast and faster moving targets evoked shorter latency and larger magnitude

SLRs, compared to low contrast or static targets.

487   When examining SLR magnitudes in relation to the magnitude of reach-related activity in

experiment 2 (**Figure 7c**), we also observed large variability as seen in experiment 1 (**Figure**

6c). However, in experiment 2, SLR magnitudes in the emerging target paradigm tended to be

even larger than in experiment 1. SLR magnitudes for moving targets reached ~75% of the level

of reach-related activity (~20% higher than experiment 1; compare **Figures 6c** versus **7c**), with a

few values approaching 150%. An example of such a phenomenon is shown in **Figure 8**. As seen

in **Figure 8a**, the differences between red lines defining the SLR epoch highlighted by the grey

box, are approximately equal to, and slightly greater than the differences between lines of

movement aligned activity shown in **Figure 8b** (movement aligned magnitude epoch highlighted

by grey box). This figure demonstrates that SLR magnitude may be on par, or even greater than,

movement-aligned activity. We consider that the most likely reason for larger SLR magnitudes

in experiment 2 is due to differences in the apparatus used (see discussion). **Figure 8** also

illustrates the variability in the appearance of the SLR across subjects. For example. while the

SLR in **Figure 8** is crisp and distinct from subsequent recruitment, those seen in previous figures

(e.g., see mean EMG activity in **Figures** 4 and **5**) show a 'smoothing' of fast visuomotor activity

into voluntary activity.

503   To summarize, the results from experiment 2 concurred with the results from experiment

1 in that SLR latency was not influenced by target speed, regardless of target contrast. A target

of the same contrast will elicit SLRs of the same latency, regardless of how fast the target is

moving.

The SLR and its relationship to reaching behaviour

So far, we have shown that 97% of tested subjects were capable of generating an SLR towards

emerging targets, and furthermore that visual features such as speed and contrast of a target alter

the latency and magnitude of the SLR. We now turn to the relationship between the latency and

magnitude of the SLR, and how these features impact ensuing RT.

In both experiments SLR latency and magnitude were negatively correlated with each

other (**Figure 9a** exp. 1; R=-.48, p=.00012; **Figure 9d** exp. 2; R=-.41, p=.00038), indicating that

larger magnitude SLRs are associated with shorter latency SLRs. Recall that there were no

significant latency differences for any target conditions found in experiment 1. This negative

correlation indicates that short latency SLRs, even with small variability and a narrow range of

latencies (all SLR latencies between ~90-110 ms **Figure 9a**), still tend to be associated with

larger magnitude SLRs.

When examining how the SLR relates to RT, we found a positive correlation between

latency and RT in both experiments (experiment 1- **Figure 9b**; R=.36, p=.0047; experiment 2-

Figure 9e; R=.67, p=8.4e-11); indicating that longer SLR latencies were associated with longer

latency RTs. Conversely, there was a strong negative correlation between magnitude and RT for

both experiments (experiment 1- **Figure 9c**; R=-.75, p=8.69e-12; experiment 2- **Figure 9f**; R=-

.40, p=.0005), indicating that decreasing SLR magnitudes are correlated with increasing RT

latency. Thus, in conditions where stimulus appearance evoked larger and earlier SLRs, subjects

generally produced shorter RTs.

These findings raise the question of how much RTs change in accordance with the

changes in SLR latency and magnitude. In both experiments, we saw a significant reduction in

RT latency for faster moving, or higher contrast targets, compared to static targets (see **Figure**

10). In experiment 1, we found that faster moving targets were associated with shorter latency

RTs (one-way ANOVA, $F(4, 79)= 2.71$, $p= .036$). RTs decreased by an average of ~20 ms for

static versus fast targets (static versus fastest; 250ms+/-25 versus 231ms+/- 16; paired t-test; t=

3.25, p=.005, df= 15) (see **Figure 4** for an example; note the leftward shift of white boxes

indicating RTs in the first row versus the last row). In experiment 2, we found that both target

speed (two-by-two ANOVA, $F(1, 107)= 17.33$, $p= 6.5e-5$) and contrast (two-by-two ANOVA,

$F(2, 107)= 103.07$, $p= 3.25e-25$) impacted RT. RTs significantly decreased in experiment 2 by

an average of 62 ms for high contrast moving targets (static versus fastest; 243ms+/-26 versus

181ms+/- 9; paired t-test; t= 10.2, p=5.7e-9, df= 17), and 57 ms for low contrast moving targets

(static versus fastest; 253ms+/-26 versus 196ms+/- 11; paired t-test; t= 9.7, p=2.2e-8, df= 17). Of

great interest, the RTs shown in **Figure 10** strongly resemble our manipulated SLR

characteristics depending on the experiment, for both magnitude (compare **Figure 6b** to

experiment 1 in **Figure 10)**, and latency (compare **Figure 7a** to experiment 2 in **Figure 10).**

Taken together, these results indicate that strongly evoking a fast visuomotor system (indicated

by short latency and large magnitude SLRs) is correlated to decreasing RT. RTs may be

expedited by as much as 62 ms in an emerging target paradigm with high contrast fast moving

targets, compared to static targets.

Recall in experiment 1 that we presented two different static conditions, one in the

context of an emerging target paradigm (0 cm/s), and one in the standard static task (S1). The 0

cm/s target evoked larger and more prevalent SLRs compared to S1 targets presented out of the

emerging target paradigm. Therefore, we also investigated whether differences in SLRs related

to differences in RT. Averaged across subjects, RTs in the 0cm/s condition were not different

than RTs in the S1 condition (250 +/- 22 ms versus 250 +/- 25 ms; n.s.). Thus, despite

differences in SLR magnitudes between static targets, there was little difference in RTs. Taken

together, these results indicate that fast visuomotor relates to, but is not fully predictive of, the

ensuing RT.

Interestingly, we noticed in **Figure 6a** and **Figure 7a**, that there were latency differences

between experiments 1 and 2, despite both using 'high contrast' targets (unpaired t-test; t= 7.99,

p= 2.35e-8, df= 25). We found that experiment 2 targets were associated with shorter latency

SLRs by approximately 15 ms (99 +/-5 ms versus 84 +/- 4ms). However, when we compared

SLR latencies between experiment 1 and low contrast fast moving targets in experiment 2, we

did not find any differences (unpaired t-test; t= 1.2, p= .22, df= 25). Therefore, its possible that

the reason for this RT difference is due to differences in monitors, where the 'high contrast'

target in experiment 1 is similar to the low contrast target in experiment 2 (see discussion).

Here we have demonstrated that fast moving, and high contrast targets presented in an

emerging target paradigm tended to elicit the most prevalent, shortest latency and largest

magnitude SLRs, which were correlated with shorter latency RTs. We found that RTs decreased

by as much as 62 ms when using these targets. These results indicate that the neural circuitry

mediating a fast visuomotor response are best evoked by a faster moving, higher contrast targets.

Discussion

The brain has a remarkable capacity to rapidly transform vision into action, allowing us to

initiate reaches towards a target within fractions of a second. Previous research has shown that

some visual attributes, such as stimuli composed of lower spatial frequencies or higher contrast,

can elicit earlier reaching responses than others (Veerman et al., 2008; Wood et al., 2015; Kozak

et al., 2019). Here, by using a newly-described emerging target paradigm (Kozak et al., 2020),

we show that high-contrast, moving stimuli are particularly capable of eliciting fast visuomotor

responses such as SLRs and expedited reach reaction times. Further, as the same subjects

performed both the novel emerging target paradigm and a visually-guided reach paradigm used

previously, we can better understand the sensory and cognitive factors promoting fast

visuomotor responses. Finally, by studying a variety of stimuli in the emerging target paradigm,

we can gain further insights into the relationship between the earliest phase of stimulus-related

EMG recruitment, subsequent phases of EMG recruitment leading up to movement onset, and

the RT of the movement itself. Together, our results contribute to a growing understanding of

the interplay between the cortical and subcortical substrates mediating visually-guided reaches in

conditions requiring expedited responses.

Comparison to past studies

Previous research on rapid visually-guided reaches have typically used either target-jump

paradigm where a visual target is displaced during an arm movement (Soechting and Lacquaniti,

1983; Day and Lyon, 2000), or a static target reaching paradigm where a suddenly appearing

target evokes reaches from a stationary arm position (Pruszynski et al., 2010). While useful, such

approaches have limitations: the response of interest evolves during an on-going movement in

target-jump experiments, or not all subjects exhibit SLRs in a typical static target paradigm, even

with intramuscular electrodes (Pruszynski et al., 2010; Gu et al., 2016). In contrast, in the

emerging target paradigm, the arm is stationary at the time of target emergence and evokes

highly prevalent fast visuomotor responses that can be recorded even with surface electrodes

(Contemori et al., 2020; Kozak et al., 2020).

The results in experiment 1 show that much larger SLRs are provoked by the appearance

of a stationary target in the emerging target paradigm rather than the static task (blue rectangle,

Figure 6, approximately 2 times larger). Larger magnitude SLRs likely relate to the increased

proportion of subjects expressing an SLR to a stationary target in the emerging target paradigm

rather than static task (100% vs 56%). We speculate that a signal of implied motion of the target

behind the barrier is central to the increased SLR magnitude in the emerging target paradigm

(Kozak et al., 2020). Recent work has shown that the certainty of the time of target emergence

also facilitates the expression of more prevalent and larger SLRs (Contemori et al., 2020). Note

that temporally predictable targets still have to be presented within the emerging target paradigm,

as smaller magnitude and less prevalent SLRs are elicited in static tasks that had high temporal

certainty via a predictable gap between the offset of a central hold stimulus and target

presentation (Pruszynski et al., 2010; Wood et al., 2015).

Past work has shown that stimuli composed of progressively higher contrast (Wood et al.,

2015) or lower spatial frequencies (Kozak et al., 2019) elicit progressively larger magnitude and

lower latency SLRs. Indeed, higher contrast moving stimuli evoked larger and earlier SLRs in

experiment 2 (**Figure 7**). Somewhat surprisingly, this inverse relationship between SLR latency

and magnitude was not observed in the manipulation of the velocity of moving stimuli. Instead,

SLR latency remained invariant while SLR magnitude was greatest for the fastest moving

stimulus; this was observed both in experiment 1 (**Figure 6**) and in experiment 2 for stimuli of

similar contrast (**Figure 7**). Thus, at least across the parameters explored in this study, SLR

magnitude is impacted by both stimulus contrast and velocity, whereas SLR latency is impacted

by stimulus contrast but not velocity.

We speculate that this dissociation between SLR latency and magnitude for stimulus

velocity may relate to the displacement of the stimulus across sequential display frames, with

larger displacements increasing the vigor of sensory transients that remain timed to the onset of

the first frame. Future studies should examine the impact of subsequent display frames on the

SLR, given that Contemori and colleagues (2020) found that a stimulus flashed for a single

frame (~16 ms) provokes earlier and larger SLRs than moving targets; it is possible that flashed

stimuli may prove to be a particularly potent means of evoking vigorous sensory transients from

the retina.

Relating the SLR to subsequent RTs

SLRs represent the first change in muscle recruitment following visual stimulus presentation. As

such, they demarcate the completion of the most rapid sensory-to-motor transformation at the

upper limb. Past reports show that larger magnitude SLRs precede shorter latency movements on

a trial-by-trial basis (Pruszynski et al., 2010; Gu et al., 2016; Contemori et al., 2020), and here

we show, when pooling across all conditions in which subjects exhibited an SLR, that larger

magnitude and earlier SLRs precede shorter reach RTs (**Figure 9;** see also Kozak et al., 2019).

These correlative relationships are consistent with a functional contribution of the SLR to the

forces necessary to overcome the limb's inertia. However, we are mindful that the SLR is

followed by sustained levels of muscle recruitment preceding the onset of the reaching

movement, which also contribute to movement initiation. Indeed, comparing SLR latency and

magnitude to reach RT in experiments 1 and 2 leads to some interesting insights. For example,

while nearly identical SLRs were observed in the emerging target task in experiment 1 to the 0

deg/s and 20 deg/s stimulus (**Figure 6**), the reach RTs to the 20 deg/s stimulus were ~20 ms

shorter (**Figure 10**). Conversely, while nearly identical reach RTs in experiment 2 were observed

to fast moving low-contrast versus slow moving high-contrast targets (**Figure 10**), the SLRs to

these stimuli were different, being larger magnitude but starting later for low-contrast, fast

moving targets (**Figure 7**). These distinctions between SLR parameters and RTs likely attest to

continued processing of the visual stimulus, and the contribution of such processing to the RT.

The judicious selection of stimulus parameters, e.g., to elicit more robust SLRs but longer RTs,

may aid the study of the fast visuomotor responses by reducing the overlap between the SLR and

activity aligned to movement onset.

Plausible neural substrates

The superior colliculus (SC) has been implicated as a potential candidate mediating the SLR

(Corneil et al., 2004; Pruszynski et al., 2010). The SC is a midbrain structure whose intermediate

and deep layers (SCi) interface with premotor structures for eye, head, and upper limb

movements within the reticular formation (Corneil and Munoz, 2014), whereas superficial layers

respond to visual input. Our results add further circumstantial evidence for a role for the SC in

the SLR. Previous work established that both the SLR and visual responses in the SC

preferentially respond to low spatial frequency (Chen et al., 2018; Kozak et al., 2019) and high

contrast targets (Li and Basso, 2008; Marino et al., 2015; Wood et al., 2015). Moving targets also

evoke robust SC responses (Marrocco and Li, 1977), and we have shown here that moving

targets also evoke robust SLRs. Importantly, the target speeds used in the current study resemble

those used in the Marrocco and Li (1977) study, which evoked robust responses in the

intermediate and deep layers of the SC where visuomotor and motor neurons activated before

arm movements reside (Werner et al., 1997; Stuphorn et al., 1999; Philipp and Hoffmann, 2014)

(Stuphorn et al., 1999). The hypothesis that the SC mediates the SLR requires further testing

with casual manipulation in tasks that promote SLR expression. Work reporting that SCi

inactivation impacted the selection of reach targets but not reach kinematics (Song et al., 2011)

employed paradigms that would not promote SLR expression.

An open question is how visual input from target motion accesses the SC. Visual

information may access the superficial SC directly via a retinotectal route (summarized in: (May,

2006)), or indirectly via a retino-geniculo-striate loop through cortex, and then on to the SC

(Schiller et al., 1979). The SC also receives projections from higher order cortical centers such as

MT (an area which responds to target motion (Nhan and Callaway, 2012)); such signals may

provide the SC with a pre-target, priming signal related to implied motion that potentiates SLR

expression.

Methodological considerations

All of our subjects exhibited an SLR in at least one stimulus condition. This replicates the high

prevalence of SLR detection in the emerging target paradigm (Contemori et al., 2020; Kozak et

al., 2020), and affirms the contention that most if not all subjects are capable of exhibiting SLRs

in an appropriate paradigm. However, subjects did not necessarily exhibit an SLR to every

stimulus in the emerging target paradigm, even for high-contrast, fast moving stimuli (**Figure 2**).

We speculate that this result may be due, at least in part, to how we detect the presence of the

SLR with the RT split analysis, which allows for a direct comparison of SLR prevalence to those

reported elsewhere which also used this method (e.g.,(Kozak et al., 2019, 2020; Contemori et al.,

2020)). The RT split analysis is somewhat conservative in that it requires an SLR to both short-

and long-latency trials. SLR- observations could therefore include those situations where an SLR

precedes the short- but not long-RT trials (see top two rows of **Figure 5**, which appear to exhibit

SLRs for the shorter-than-average RTs). The RT split analysis is also more suited to paradigms

engendering a larger degree of RT variance. As noted previously (Kozak et al., 2020), one

potential complication of the emerging target paradigm is that it may shorten RTs too much, not

only reducing RT variance but also leading to a potential overlap of SLR with activity linked to

movement onset. As noted above, judicious stimulus selection may help avoid such overlap, but

we also expect the continued evolution of SLR detection methods suited to a particular

paradigm.

We noted large differences in the latencies of fast visuomotor responses across

experiments, with SLRs evolving ~15 ms earlier and reach RTs being ~50 ms shorter in

experiment 2 compared to experiment 1. The most likely reason for these differences is the

contrast of the stimuli, as the VPIXX projector used in experiment 2 has a higher level of

contrast (275 cd/m2) than the standard monitor used in experiment 1 (110 cd/m2). Consistent

with this, the SLR latencies from the high-contrast target in Experiment 1 (**Figure 6b**) resemble

those of the low contrast target in experiment 2 (**Figure 7a**). The large difference in reach RTs

across experiments, even to targets associated with similar latency SLRs, may relate to postural

differences in the exoskeleton-based Kinarm robot in experiment 1 versus the endpoint-based

Kinarm robot in experiment 2.

Conclusions

The work presented here demonstrates that all subjects can generate SLRs, but are more likely to

present one in an emerging target paradigm. As a consequence, reliable SLRs can be recorded

even with surface EMG, and here we use this paradigm to show that targets moving at faster

speeds elicit larger magnitude SLRs, whereas higher contrast targets decrease SLR latency.

Larger magnitude and shorter-latency SLRs tend to proceed expedited RTs, although RTs are

also influenced by continued stimulus processing after the SLR. As the SLR is the first wave of

muscle recruitment tied to visual target onset, muscle recruitment during this interval may

provide a unique window into the contribution of the tecto-reticulospinal signaling that is distinct

from subsequent phases of target-related recruitment of corticospinal or cortico-reticulo-spinal

circuits. Rather than being directly involved in the SLR, cortical areas may serve to prime this

phylogenetically-conserved circuit to produce the first wave of muscle recruitment in situations

when time is of the essence, preceding subsequent waves of EMG recruitment that arise

following more detailed, cortically-related processing of the sensory signal.

References

Chen C-Y, Sonnenberg L, Weller S, Witschel T, Hafed ZM (2018) Spatial frequency sensitivity

in macaque midbrain. Nat Commun 9:2852.

Contemori S, Loeb GE, Corneil BD, Wallis G, Carroll TJ (2020) The influence of temporal

predictability on express visuomotor responses. J Neurophysiol.

Corneil BD, Munoz DP (2014) Overt Responses during Covert Orienting. Neuron 82:1230–

1243.

Corneil BD, Olivier E, Munoz DP (2004) Visual responses on neck muscles reveal selective

gating that prevents express saccades. Neuron 42:831–841.

Day BL, Lyon IN (2000) Voluntary modification of automatic arm movements evoked by

motion of a visual target. Exp Brain Res 130:159–168.

Glover IS, Baker SN (2019) Multimodal stimuli modulate rapid visual responses during

reaching. J Neurophysiol 122:1894–1908.

Gu C, Pruszynski JA, Gribble PL, Corneil BD (2018) Done in 100 ms: path-dependent

visuomotor transformation in the human upper limb. J Neurophysiol 119:1319–1328.

Gu C, Wood DK, Gribble PL, Corneil BD (2016) A Trial-by-Trial Window into Sensorimotor

Transformations in the Human Motor Periphery. J Neurosci 36:8273–8282.

Kozak RA, Cecala AL, Corneil BD (2020) An Emerging Target Paradigm to Evoke Fast

Visuomotor Responses on Human Upper Limb Muscles. J Vis Exp:e61428.

Kozak RA, Kreyenmeier P, Gu C, Johnston K, Corneil BD (2019) Stimulus-Locked Responses

on Human Upper Limb Muscles and Corrective Reaches Are Preferentially Evoked by Low

Spatial Frequencies. eneuro 6:ENEURO.0301-19.2019.

Lara AH, Elsayed GF, Zimnik AJ, Cunningham JP, Churchland MM (2018) Conservation of

preparatory neural events in monkey motor cortex regardless of how movement is initiated.

Elife 7.

Li X, Basso MA (2008) Preparing to Move Increases the Sensitivity of Superior Colliculus

Neurons. J Neurosci 28:4561–4577.

Marino RA et al. (2015) Linking express saccade occurance to stimulus properties and

sensorimotor integration in the superior colliculus. J Neurophysiol 114:879–892.

Marino RA, Levy R, Boehnke S, White BJ, Itti L, Munoz DP (2012) Linking visual response

properties in the superior colliculus to saccade behavior. Eur J Neurosci 35:1738–1752.

Marrocco RT, Li RH (1977) Monkey superior colliculus: properties of single cells and their

afferent inputs. J Neurophysiol 40:844–60.

Maunsell JH, Gibson JR (1992) Visual response latencies in striate cortex of the macaque

monkey. J Neurophysiol 68:1332–1344.

May PJ (2006) The mammalian superior colliculus: Laminar structure and connections. Prog

Brain Res.

Nhan HL, Callaway EM (2012) Morphology of superior colliculus- and middle temporal area-

projecting neurons in primate primary visual cortex. J Comp Neurol.

Philipp R, Hoffmann K-P (2014) Arm movements induced by electrical microstimulation in the

superior colliculus of the macaque monkey. J Neurosci 34:3350–3363.

Pruszynski AJ, King GL, Boisse L, Scott SH, Randall Flanagan J, Munoz DP (2010) Stimulus-

locked responses on human arm muscles reveal a rapid neural pathway linking visual input

to arm motor output. Eur J Neurosci 32:1049–1057.

Rezvani S, Corneil BD (2008) Recruitment of a head-turning synergy by low-frequency activity

in the primate superior colliculus. J Neurophysiol 100:397–411.

Schiller PH, Malpeli JG, Schein SJ (1979) Composition of geniculostriate input to superior

colliculus of the rhesus monkey. J Neurophysiol 42:1124–33.

Schmolesky MT, Wang Y, Hanes DP, Thompson KG, Leutgeb S, Schall JD, Leventhal AG

(1998) Signal timing across the macaque visual system. J Neurophysiol 79:3272–3278.

Soechting JF, Lacquaniti F (1983) Modification of trajectory of a pointing movement in response

to a change in target location. J Neurophysiol 49:548–564.

Song J-H, Rafal RD, McPeek RM (2011) Deficits in reach target selection during inactivation of

the midbrain superior colliculus. Proc Natl Acad Sci 108:E1433–E1440.

Stuphorn V, Hoffmann KP, Miller LE (1999) Correlation of primate superior colliculus and

reticular formation discharge with proximal limb muscle activity. J Neurophysiol 81:1978–

1982.

Veerman MM, Brenner E, Smeets JBJ (2008) The latency for correcting a movement depends on

the visual attribute that defines the target. Exp Brain Res 187:219–228.

Werner W, Dannenberg S, Hoffmann K-P (1997) Arm-movement-related neurons in the primate

superior colliculus and underlying reticular formation: comparison of neuronal activity with

EMGs of muscles of the shoulder, arm and trunk during reaching. Exp Brain Res 115:191–

205.

Wood DK, Gu C, Corneil BD, Gribble PL, Goodale MA (2015) Transient visual responses reset

the phase of low-frequency oscillations in the skeletomotor periphery. Eur J Neurosci

42:1919–1932.

Figure legends

Figure 1. The emerging target paradigm (Kozak et al., 2020). a) A target (T1) appears above an

occluder (grey box) and the subject the real-time cursor (RTC; white dot below occluder that

represents hand position), into the start position (T0; grey dot below the occluder). b) Once the

RTC is in the start position, T1 moves towards the subject along the inverted y path and then

disappears behind the occluder. c) The subject makes a visually-guided reach to T1 as soon as it

emerges at the right or left outlet below the occluder.

Figure 2. Detection of SLRs. a) Exemplar trial-by-trial recruitment of activity from right

pectoralis muscle during leftward and rightward reaches from a subject with a detectable SLR.

Each row is a different trial, with the intensity of color conveying degree of recruitment. Trials

sorted by RT (white boxes) and aligned to stimulus onset. The SLR (highlighted by asterisk)

appears as a vertical banding of increased or decreased recruitment aligned to left or right

stimulus presentation, respectively, rather than movement onset. b) Mean +/- SE of EMG activity

for the early- (purple) and late- (green) RT subsets based on mean RT. c) In the RT split analysis,

separate time-series ROCs were run for the early and late subsets, and then the mean RT for each

subset is plotted as a function of the discrimination time. The slope of the line connecting these

points determines whether the divergence of EMG activity is more time-locked to stimulus or

movement onset. d) For SLR+ observations, all trials were pooled together regardless of RT to

determine the SLR latency (derived from the time-series ROC) and magnitude (derived from the

mean EMG waveformes; see methods). Grey rectangle in b) and d) depict the SLR interval.

Figure 3. Prevalence of SLRs across the population. Slope of the relationship between

discrimination time and average RT for early and late RT groups (e.g. see slope in Figure 2c)

Each data point represents a unique subject for a particular condition. Slopes are capped at 90

degrees and a dotted black line is drawn at 67.5 deg. A data point above this line represents a

SLR+ observation, whereas a data point below the line represents SLR- observations. a)

Experiment 1. Black and grey diamonds represent static conditions in the static and emerging

target paradigms, respectively. Blue diamonds represent moving targets in emerging target

paradigm, with slower moving targets are represented by darker shades. b) Experiment 2. Purple

diamonds represent high contrast targets; green diamonds represent lower contrast targets. Static

and slower moving targets are represented by darker shades. 2 subjects completed both

experiments; subjects 2 and 16 from experiment 1 are subjects 18 and 12 from experiment 2,

respectively.

Figure 4. Data from representative subject in experiment 1, using the same format as in figure 2.

From left to right, columns shown trial-by-trial EMG activity for left and right targets sorted by

RT, mean +/- SE EMG activity for left and right targets with overlaid time-series ROC, and the

RT split analysis for data shown in the corresponding row (the number in each panel is the

slope).

Figure 5. Data from representative subject in experiment 2,using the same format as figure 4.

Figure 6. Summary of results from experiment 1, showing SLR latency (a), SLR magnitude (b,

normalized within each subject to the largest SLR across all conditions, and SLR magnitude

normalized to the maximum of movement-aligned EMG activity. In c), each asterisk represents

data from an individual subject. S1 indicates the results from the static task, which presents the

same retinal stimulus to the subject as the 0 cm/s target in the emerging target paradigm . Error

bars show SE.. * indicates significance (see results text for more details).

Figure 7. Summary of results from experiment 2 (same format as Fig. 6), with SLR latency (a),

SLR magnitude normalized to the within-subject largest SLR (b), and SLR magnitude

normalized relative to maximum of movement-aligned activity. All values presented as mean +/-

standard error. * indicates significance. HC= high contrast LC= low contrast.

Figure 8. Example subject whose SLR magnitude exceeds the maximum level of EMG activity

aligned to target onset. Data, shown as mean +/- SE, are aligned to stimulus onset in a), or

movement onset in b). SLR magnitude is measured as area between left and right reaches for the

30 ms interval after the SLR latency (grey box, running from 73-103 ms relative to stimulus

onset). + 30 ms. For movement-aligned activity, magnitude is measured as area 15 ms before and

15 ms after RT onset (grey box in b).

Figure 9. Correlations for SLR latency, magnitude and associated reaction time in experiment 1

(top row) or 2 (bottom row). a, d) SLR latency and magnitude correlations. b, d) SLR latency

and RT correlations. c, f) SLR magnitude and RT correlations. All correlations shown are

significant. Each blue circle represents data from an SLR+ subject in a unique experimental

condition. Solid red lines indicates first order fit of the data.

Figure 10. Summary of RTs in experiment 1 (a) and 2 (b), following the formats in figures 6 and

7.

a

Stabilize hand at start position

b

target moves down path and disappears
behind occluder

c

subject reaches to intercept target
from left or right outlet

Figure 1- Kozak and Corneil 2021

Figure 2- Kozak and Corneil 2021

Figure 3- Kozak and Corneil 2021

Figure 4- Kozak and Corneil 2021

Figure 5- Kozak and Corneil 2021

Figure 6- Kozak and Corneil 2021

Figure 7- Kozak and Corneil 2021

a

b

Figure 8- Kozak and Corneil 2021

Figure 9- Kozak and Corneil 2021

Figure 10- Kozak and Corneil 2021