1 **High contrast, moving targets in an emerging target paradigm promote fast visuomotor**
2 **responses during visually guided reaching**
3
4
5 Abbreviated title: Moving targets promote fast visuomotor responses
6

7 Rebecca A. Kozak[1,4], Brian D. Corneil[1,2,3,4]
8
9 [1]Graduate Program in Neuroscience, Western University, London, On, Canada, N6A 5B7
10 [2]Department of Psychology, Western University, London, Ontario, Canada, N6A 5B7
11 [3]Department of Physiology and Pharmacology, Western University, London, Ontario, Canada,
12 N6A 5B7
13 [4]Robarts Research Institute, 1151 Richmond St. N, London, Ontario, Canada, N6A 5B7
14
15 *Corresponding Author: Brian D. Corneil, bcorneil@uwo.ca

16
17
Number of pages: 40
Number of figures: 10
Number of tables: 0
Abstract word count: 250
Introduction word count: 691
Discussion word count: 2340
24

25 Conflicts of interest: The authors declare no competing financial interests
26 Acknowledgements: We thank Dr. Andrew Pruszynski for use of his Kinarm exoskeleton for
27 Experiment 1.
28
This work is supported by a Discovery Grant to BDC from the Natural Sciences and Engineering
Research Council of Canada (NSERC; RGPIN 311680) and an Operating Grant to BDC from the
Canadian Institutes of Health Research (CIHR; MOP-93796). RAK was supported by an Ontario
Graduate Scholarship. The apparatus used in Experiment 2 was purchased using funds from the
Canadian Foundation for Innovation. Additional support came from the Canada First Research
Excellence Fund (BrainsCAN).

35
36 **Keywords:** visually-guided reaches, EMG, SLR, moving targets, contrast, reaction time, RT

Abstract

Humans have a remarkable capacity to rapidly interact with the surrounding environment, often by transforming visual input into motor output on a moment-to-moment basis. But what visual features promote rapid reaching? High contrast, fast-moving targets elicit strong responses in the superior colliculus (SC), a structure associated with express saccades and implicated in rapid electromyographic (EMG) responses on upper limb muscles. To test the influence of stimulus properties on rapid reaches, we had human subjects perform visually guided reaches to moving targets varied by speed (experiment 1) or speed and contrast (experiment 2), in an emerging target paradigm which has recently been shown to robustly elicit fast visuomotor responses. Our analysis focused on stimulus-locked responses (SLRs) on upper limb muscles. SLRs appear within <100 ms of target presentation, and as the first wave of muscle recruitment, they have been hypothesized to arise from the SC. Across 32 subjects studied in both experiments, 97% expressed SLRs in the emerging target paradigm, whereas only 69% expressed SLRs in an immediate response paradigm towards static targets. Faster moving targets (experiment 1) evoked large magnitude SLRs, while high contrast fast moving targets (experiment 2) evoked short latency, large magnitude SLRs. In some instances, SLR magnitude exceeded the magnitude of movement aligned activity. Both large magnitude and short latency SLRs were correlated with short latency reach reaction times. Our results support the hypothesis that, in scenarios requiring expedited responses, a subcortical pathway originating in the SC elicits the earliest wave of muscle recruitment, expediting reaction times.

New & Noteworthy (72 words)

60 How does the brain rapidly transform vision into action? Here, by recording upper limb muscle

61 activity, we find that high contrast and fast moving targets are highly effective at evoking rapid

62 visually guided reaches. We surmise that a brainstem circuit originating in the superior colliculus

63 contributes to the most rapid reaching responses. When time is of the essence, cortical areas may

64 serve to prime this circuit, and elaborate subsequent phases of recruitment.

Introduction

To perform a visually-guided reach, for example to pick-up a baseball, visual information such as the shape and size of the ball is transformed into motor commands which are communicated to the motor periphery. However, if the ball is thrown to us, time is of the essence and we must perform a rapid reach in order to catch the ball. In such situations, the latency of the response depends on the visual attributes of the stimulus; for example, earlier responses are elicited by larger size, higher contrast, or lower spatial frequency targets (Veerman et al., 2008; Wood et al., 2015; Kozak et al., 2019). Other work has suggested that the most rapid visuomotor transformation for reaching is mediated by a tecto-reticulo-spinal pathway, which lies in parallel with the better studied corticospinal pathway (Day and Lyon, 2000; Pruszynski et al., 2010; Gu et al., 2016; Glover and Baker, 2019).

To better understand the most rapid visuomotor transformations for limb control, we and others have recorded electromyographic (EMG) activity from upper limb muscles during rapid visually-guided reaches. Doing so has led to the identification of stimulus-locked responses (SLRs) which are the first wave of muscle activity influenced by the visual stimulus (SLRs are also termed rapid visuomotor responses by (Glover and Baker, 2019) or express responses by (Contemori et al., 2021)). SLRs evolve within < 100 ms of stimulus onset, which is before some response latencies in visual and motor cortices (Maunsell and Gibson, 1992; Schmolesky et al., 1998; Lara et al., 2018), but after response latencies in the superior colliculus (SC) from which the tecto-reticulo-spinal track originates (Stuphorn et al., 1999; Rezvani and Corneil, 2008). Further, responses in the SC and SLRs exhibit similar stimulus preferences, in that both are preferentially evoked by high contrast (Li and Basso, 2008; Marino et al., 2012; Wood et al.,

2015) or low spatial frequency (Chen et al., 2018; Kozak et al., 2019) stimuli. Such observations

are consistent with the hypothesis that the SC mediates the SLR.

Robust SC responses are also elicited by moving visual targets (Marrocco and Li, 1977),

however, a systematic investigation of target motion on the SLR has not been performed.

Previous reports examining short latency reaching responses have typically relied on targets that

either suddenly appear (a *static target paradigm*; (Pruszynski et al., 2010)), or are displaced

during a reach (*target jump paradigm*; (Soechting and Lacquaniti, 1983; Day and Lyon, 2000)).

Recently, we have presented an *emerging target paradigm*, where a target travels down an

inverted 'Y' path before disappearing and then emerging from behind an occluder (Kozak et al.,

2020). Subjects are instructed to reach and intercept the target when it emerges from behind the

occluder. Benefits of the emerging target paradigm include the stability of the arm when the

target emerges (which is not the case in the target jump paradigm), and the generation of robust

fast visuomotor responses in almost every subject (which is not the case in the static target

paradigm). In Kozak et al. (2020), all subjects exhibited at SLR in the emerging target paradigm

with magnitudes five times larger than in the static paradigm. The efficacy of the emerging target

paradigm in generating SLRs has been replicated in a larger sample of subjects, wherein 20 of 21

subjects expressed an SLR (Contemori et al., 2021). Thus, essentially every subject has the

circuitry to generate fast visuomotor responses; the failure to do so in past studies arose from the

reliance on suboptimal stimuli and/or tasks.

The main goal of the current study is to systematically investigate how target speed and

contrast impact fast visuomotor responses in the emerging target paradigm. We hypothesize that

the SC mediates rapid visuomotor responses on the upper limb, and therefore predict that high

contrast fast moving targets will elicit more prevalent, shorter latency, and larger magnitude

SLRs. A secondary goal is to better understand how the emerging target paradigm promotes the

generation of SLRs, by comparing such SLRs to those in the static target paradigm. Overall, we

find that fast moving, high-contrast, targets in the emerging target paradigm robustly recruits the

circuitry mediating the most rapid visuomotor transformations for reaching movements.

Materials and methods

Subjects

A total of 34 subjects (18 females, 16 males; mean age: 24.2 years SD: 5.6) completed at least

one of the two experiments. Two subjects completed both experiments. All subjects provided

informed consent, were paid for their participation, and were free to withdraw from the

experiment at any time. All but 1 subject was right-handed. All subjects had normal or corrected-

to-normal vision, with no current visual, neurological, or musculoskeletal disorders. All

procedures were approved by the Health Science Research Ethics Board at the University of

Western Ontario. Two subjects (1 from each experiment) were excluded due to low quality EMG

recordings (see below).

Apparatus

Subjects were seated in either a KINARM robotic Exoskeleton (experiment 1) or a KINARM

End-Point Lab (experiment 2) (KINARM Technologies, Kingston, ON, Canada), and performed

reaching movements with their right arm. Both platforms relied on the same experimental

software. Visual stimuli were projected onto an upward facing mirror from a downward facing

monitor in experiment 1 (LG 47LS35A; size: 47" resolution: 1920x1080 pixels (Exoskeleton)) or

a custom built in projector in experiment 2 (PROPixx projector by VPixx, Saint-Bruno, QC,

Canada; custom integrated into the KINARM End-Point Lab). Each platform has a different

distance from the eyes to the target. We positioned visual stimuli at different locations in each

platform so that targets were presented at similar retinal eccentricities and velocities. In both

platforms, a shield below the mirror occluded direct vision of the hand, but hand position was

represented on the monitor in real-time via a *real-time cursor* (RTC) projected onto the screen. A

photodiode was used to indicate the precise time of visual stimulus presentation, and all

kinematic and electromyographic (EMG) data were aligned to photodiode onset. A loading force

was provided in both experiments to increase the baseline activity of the muscle of interest, so

that the SLR could be expressed as either an increase or decrease in muscle activity following

leftward or rightward target presentation, respectively. This loading force was different in

Experiment 1 and 2 given that these experiments were conducted on different platforms.

Experimental design:

In experiment 1 and 2, we used an emerging target paradigm as previously described in

((Kozak et al., 2020) see **Figure 1**). The emerging target paradigm involves a target starting at

the top of an inverted y (**Figure 1a**), moving towards an occluder and disappearing behind it

(**Figure 1b**), then emerging below the occluder at one of two outlets. Subjects were instructed to

reach and intercept the target after it emerged from behind the occlude as quickly and accurately

as possible (**Figure 1c**) (Kozak et al., 2020). During target occlusion, subjects were instructed to

fixate on a notch at the bottom of the occluder, midway between the inverted Y outlets. The

target only became visible once it had fully moved past the occluder (i.e., the target was

presented in its entirety, rather than sliding past the occluder to avoid a half-moon stimulus). The

target then continued to move down the inverted 'Y' path. The trial ended as soon as the RTC

contacted the peripheral target, or if the target passed the bottom of the screen. In both

experiments, we also collected data in a *static task*, where subjects performed visually guided

reaches as quickly and accurately as possible towards stationary targets in a separate block of

trials. Doing so allowed us to directly compare SLRs in the emerging target paradigm to those

obtained with a static task typically used to evoke the SLR (Pruszynski et al., 2010; Wood et al.,

2015; Gu et al., 2016; Kozak et al., 2019). Kinematic and electromyographic (EMG) data are

aligned to target emergence below the occluder in the emerging target task, or relative to

peripheral target presentation in the static task. Eye movements were not measured.

Experiment 1: effects of speed.

Static task

In the static task in Experiment 1, subjects (*n* = 16) performed visually guided reaches from a

stationary arm position to static targets appearing pseudorandomly to either the right or left of

the start position. Throughout this task, a constant torque of 2 Nm was applied to the shoulder,

increasing tonic activity on the muscle of interest. Each trial was initiated when a subject moved

the RTC representing their hand into a white target representing the start position (T0). T0 was

1.5 cm in diameter, located approximately 45 cm in front of the subject relative to midline, and

changed to grey once the RTC was in T0. A target (T1) was presented after a 1s hold period. If

the subject exited the start position before completion of this hold period, the trial was reset, and

T0 changed back to white. Subjects were instructed to also look at T0 before the onset of the

target (T1). After the hold period, a white T1 was presented on the black background 17 cm to

the left or right of the start position (~17.5 degrees of visual angle). A 17 cm eccentricity was

chosen as it resembled the interception point of the emerging target paradigm. The trial ended

when the RTC made contact with the center of T1. T0 remained visible as a grey target,

changing back to white once the trial ended. Subjects completed 1 block of 220 trials (110 trials

towards each target), before they performed the emerging target paradigm.

Emerging target paradigm

After completing the static task, the same 16 subjects performed visually guided reaches towards

moving targets in the emerging target paradigm (See **Figure 1** for protocol). The start position,

RTC, hold period, background loading torque, and target radius were the same as in the static

task.

We implemented three target speeds in our paradigm (10, 15 or 20 cm/s), translating to

initial velocities of approximately 10, 15 or 20 retinal degrees per second, respectively. These

speeds were chosen to span a range from 'slow' to 'fast' while still permitting target interception

of the fastest target. To control for the time that a participant has to prepare for an upcoming

reach, the duration that T1 was visible above the occluder was constant (.5s) regardless of target

speed. As targets travelled at different speeds, this meant that they started at different heights

above the occluder. The occluder was 30 cm wide by 15 cm high, and the center of the occluder

was located ~63 cm away from the subject. Behind the occluder, the target moved at a constant

speed of (15 cm/s) regardless of initial or final target speed. Thus, the target consistently

disappeared behind the occluder for 1s on all trials.

We also included a static 0cm/s target within the emerging target paradigm, where a

target began moving at a speed of 20 cm/s, disappeared behind the occluder, and then appeared ~

17 cm to the left or right of the start position and did not move. Thus, the initial conditions where

the target drops into the occluder were the same for the 0 cm/s and 20 cm/s targets. On the 0

cm/s trial, the trial ended when the subjects made contact with the target. The target disappeared

behind the occluder for 900 ms on 0 cm/s trials, rather than 1000 ms as on moving target trials,

due to an oversight. Therefore, a target appeared below the occluder 1.5 s after initiation for

moving targets, and 1.4 s for static targets. Contemori and colleagues (2021) recently showed

that SLRs are promoted by certainty of the time of target appearance, which would have

predicted that weaker SLRs would have been observed for 0 cm/s targets. This indicates that our

results would be biased against robust SLRs in 0cm/s conditions, which we did not observe.

All targets in experiment 1 were white (110 cd/m^2) presented against a black ($.6 \text{ cd/m}^2$)

background (contrast ratio: 183:1). There were 8 unique trial conditions (4 target speeds 0, 10, 15

and 20 cm/s presented to the left and right). Subjects completed 2 blocks of 480 trials, with each

block containing 60 repetitions of each unique trial condition presented pseudorandomly without

replacement, yielding a total of 120 trials for each unique trial condition.

Experiment 2: effects of contrast and speed

Experiment 2 was designed to investigate the interaction between target speed and contrast on

the SLR. This experiment was conducted in a different apparatus than experiment 1, in

aKINARM End-Point system. As in experiment 1, we ran a static task followed by an emerging

target paradigm.

Static task

In the *static task* in experiment 2, subjects ($n = 18$) performed visually guided reaches initiated

from a stationary position, towards static left or right targets. As in experiment 1, each trial was

initiated when a subject moved the RTC representing hand position into a start position (T0).

Due to different apparatuses being used in experiment 1 and 2, we note relevant differences

below. A constant torque was applied through the handle throughout the experiment (2Nm down,

5Nm right) to increase the baseline activity of the muscle of interest. T0 was aligned with the

subject's midline (0, -42 cm relative to the midpoint between the robot manipulanda; radius 1

cm). T0 changed from white to grey once the RTC was moved into it. The trial was reset if the

subject exited T0 before completion of the hold period (randomized time between 1000 to 2000

ms). Our use of a randomized hold time in experiment 2 resembles that used in the work by

Wood and colleagues (2015), which also manipulated stimulus contrast. Subjects were instructed

to foveate T0 before presentation of the target (T1). After the hold period, T1 (target with a 1 cm

radius) appeared 10 cm to the left or right (~15.4 degrees of visual angle). The trial ended as

soon as RTC made contact with the peripheral target. There were 4 unique trial conditions; a

high contrast white (275 cd/m^2: contrast ratio 793.8:1) or low contrast grey (18 cd/m^2: contrast

ratio 50:1) target presented against a black (.36 cd/m^2) background to either the left or right side.

T0 remained visible as a grey target, changing back to white once the trial ended. Subjects

completed 1 block of 400 pseudorandomized trials, yielding a total of 100 trials for each unique

trial condition.

Emerging target paradigm

After completing the static task, the same 18 subjects completed the emerging target paradigm.

As in experiment 1, subjects performed visually guided reaches initiated from a stable posture

towards moving targets that emerged suddenly beneath an occluder. Throughout the experiment,

the same torque was applied as used in the static task. In experiment 2, we used a two-by-two

design, where we modulated speed (including a 'slow' and a 'fast' target (10 cm/s, 20 cm/s;

translating to approximately 10 and 20 degrees of visual angle); and contrast (the 'low' and

'high' contrasts). In experiment 2, the occluder was 30cm by 15cm, centered at the midpoint

between the manipulanda (0, - 29 cm). As in experiment 1, we controlled for the amount of time

that T1 was visible before being obscured by the occluder (.5s). The time T1 disappeared behind

the occluder in experiment 2 is shorter (0.5 s) than in experiment 1 (1 s). There were 8 unique

trial conditions (2 speeds, 2 contrasts, leftward or rightward targets). The same contrasts were

used from the static task. Subjects completed 2 blocks of 400 pseudorandomized trials, yielding a

total of 100 trials for each unique trial condition.

Data acquisition and analysis

Surface electromyographic (EMG) activity was recorded from the right pectoralis muscle with

double-differential surface electrodes (Delsys Inc. Bagnoli-8 system, Boston, MA USA). We

placed one surface electrode on the sternal portion of the muscle, and another surface electrode

on the clavicular head of the muscle. Further details regarding electrode placement can be found

in our previous methods paper (Kozak et al., 2020). Our rationale for placing two electrodes is as

a backup, given the potential of losing the adhesion of surface electrodes during a long

experiment. Off-line, we analyzed the EMG recording with the higher signal to noise ratio. EMG

signals were sampled at 1 kHz, amplified by 1000, full-wave rectified off-line, and smoothed

with a 7-point smoothing function. We excluded two subjects with poor quality muscle

recordings, which was defined as an increase in peak EMG activity above baseline of less than

40 µV during leftward reaching movements.

Kinematic data were sampled at 1 kHz by the Kinarm data acquisition system in both

experiments. Reaction time (RT) was calculated as time from sudden target onset in the static

target task, or target emergence below the occluder in the emerging target paradigm (both

measured by the photodiode) to reach initiation. Reach initiation in both experiments was

defined as 8% of the peak tangential velocity. We excluded trials with RTs less than 120 ms due

to presumed anticipation, and also excluded trials with RTs exceeding 500 ms due to presumed

inattentiveness. Overall we retained more than 93% of all trials.

As described previously (Kozak et al., 2020), we used a receiver-operating characteristic

(ROC) analysis to define the presence and latency of the SLR. Briefly, the time-series ROC

analysis indicates the likelihood of discriminating stimulus presentation based on EMG alone. As

the SLR is the first wave of muscle activity influenced by the visual stimulus, we rely on this

method to discriminate the moment a visual stimulus influences EMG activity. The ROC value

(i.e., the area under the ROC curve) was calculated and plotted at every time sample (1ms) from

100 ms before to 300 ms after target presentation (an example may be seen in **Figure 2d** overlaid

with mean +/- SE of EMG activity for the data shown in **Figure 2a**). A ROC value of .5

indicates chance discrimination, whereas values of 1 or 0 indicate perfectly correct or incorrect

discrimination relative to target presentation, respectively. Discrimination thresholds were set to

0.6, and the discrimination latency was determined as the first of 8 of 10 consecutive points that

exceeded this value.

To define the presence of an SLR, an RT split analysis was used (as described in: (Wood

et al., 2015; Kozak et al., 2020)). The purpose of the RT split analysis is to empirically test

whether EMG activity is more likely to evolve with stimulus presentation, or arm movement

over a large number of trials. This method is used as an additional conservative approach to

relying on pooled EMG, which may be unfairly biased towards a subset of the shortest latency

trials. To perform the RT split analysis, separate time series ROC analyses were conducted on

early and late RT trials (purple or green highlighted trials in **Figure 2a**; see also mean EMG

traces in **Figure 2b**), defined relative to mean RT. We then plotted the mean RT for the early or

late RT subsets versus the ROC discrimination time, and then connected these two points with a

line (**Figure 2c**). The slope of this line is used to determine whether the initial change in EMG

activity is more locked to stimulus onset (slope $\geq$ 67.5 deg.) or not (slope < 67.5 deg.). The

example shown in **Figure 2** illustrates an example where an SLR was detected (termed SLR+;

versus SLR-).

We used the same criterion for determining latency and magnitude of the SLR as used in

(Kozak et al., 2020). For SLR+ observations, SLR latency was defined as the first point where

the time-series ROC constructed from all trials (regardless of RT) surpassed .6 for 8 out of 10

trials (see **Figure 2d**). SLR magnitude was defined as the area between leftward and rightward

mean EMG traces for an interval spanning from SLR latency to 30 ms later. When comparing

SLR magnitudes between subjects, all SLR+ values were first normalized to the maximum

within-subject SLR magnitude. For example, if a subject exhibited SLRs in 5 conditions, the

largest SLR condition for that subject is given a value of 1 and all other magnitudes are

presented as a fraction of 1. When comparing SLR magnitudes to recruitment associated with

voluntary movement, we divided the magnitudes attained from the SLR epoch by the sum of the

movement-aligned activity with a 30 ms window surrounding peak EMG activity.

Statistical analyses

Statistical analyses were performed in MATLAB (version R2016a, The MathWorks, Inc.,

Natick, Massachusetts, United States). When comparing SLR prevalence across conditions, chi-

squared tests were used to determine differences in the proportion of subjects exhibiting an SLR.

To compare SLR latencies and magnitudes, as well as reaction times across conditions, we used

a one-way ANOVA for experiment 1 and a 2 by 2 ANOVA for experiment 2. Post-hoc pairwise

comparisons relied on the Tukey-Kramer critical value, as well as paired and unpaired t-tests

when appropriate. Correlational analyses were performed on latency, magnitude and RT

measures on a subject-by-subject basis.

Results

Previous work has shown that the emerging target paradigm is highly effective at eliciting rapid

visuomotor responses, such as stimulus-locked responses (SLRs) and short-latency reach

reaction times (RTs) (Kozak et al., 2020; Contemori et al., 2021). Here, we sought to investigate

the features of the stimulus which best elicit SLRs by altering the speed (experiment 1) or the

speed and contrast (experiment 2) of the target, and compared these results to static tasks. We

first confirmed that all 32 subjects exhibited an SLR in at least one condition in both experiment

1 and 2 in either the static task or emerging target paradigm (**Figure 3;** each subject had at least

one diamond above the dotted line, indicating a detectable SLR). Significantly more subjects

exhibited an SLR in the emerging target paradigm compared to the static target task (31 (97%) vs

22 (69%) SLR+ subjects, respectively; chi-squared test: p= 0.0189, $\chi2$= 5.51, df= 1). Our low

SLR detection rates in the static task is consistent with previous reports (Wood et al., 2015; Gu et

al., 2016; Pruszynski et al., 2018).

Experiment 1: Faster moving targets elicit larger magnitude SLRs

In experiment 1, we investigated whether target speed influences the SLR in an emerging target

paradigm. Data from a representative subject is shown in **Figure 4.** In the plots depicting single-

trial EMG activity, the SLR appears as a vertical band of EMG activity due to changes occurring

around the same time after target emergence on every trial (grey boxes, two left columns in

Figure 4). The SLR differs from the subsequent phases of EMG activity tied to movement onset

(indicated by white boxes demarcating sorted reaction time). **Figure 4** also shows the mean

EMG traces, time-series ROC plots, and a depiction of how an SLR is detected (see methods- RT

split analysis; slopes >67.5 indicate an SLR is present). In this subject, the SLR began ~90 ms

after target emergence in all conditions, regardless of target speed, or whether a stationary

stimulus was presented in the emerging target paradigm or static task. Further, while SLR

latency did not appear to change between conditions, it appears that SLR magnitude increased

for faster moving targets (e.g., compare the intensity of yellows for the 0 vs 20 cm/s targets in

the left column of **Figure 4**, or the peak of mean EMG activity in the SLR interval in the 3rd

column).

In experiment 1, we first quantified SLR prevalence (the proportion of subjects

expressing an SLR) towards targets moving at different speeds. Every subject had an SLR in at

least one moving target condition (**Figure 3**). We found no difference in SLR prevalence across

conditions (10 v. 15 v. 20 cm/s- chi-squared, all p> .05). Recall in experiment 1 that we also

presented a static target either in the emerging target paradigm (i.e., the 0 cm/s target, which was

intermixed with other moving targets), or in a static task (S1). We found that all 16 subjects

(100%) exhibited an SLR to a 0cm/s target, but only 9 of these subjects (~56%) exhibited an

SLR in the standard, static task (S1). This difference in prevalence was statistically significant

(chi-squared test: p= 0.00275, $\chi 2$= 8.96, df=1), indicating that an emerging target paradigm

promotes the expression of an SLR regardless of whether the target is moving or not.

We next examined the latency and magnitude of the SLR. Intriguingly, and reaffirming

the impression from the individual data in **Figure** 4, we found no differences in SLR latency

across any of the conditions (**Figure 5a;** one-way ANOVA, $F_{(4,53)} = .58$, $p = .68$). Thus,

regardless of target speed, we found that the latency of the SLR remained ~90 ms. In contrast,

we found that target speed impacted SLR magnitude (**Figure 5b)**. Moving from left to right in

Figure 5b, we found that SLR magnitude increased for faster moving targets (one-way ANOVA

across all conditions, $F_{(4, 53)} = 15.15$, $p = 2.53e-8$; all emerging targets evoked larger magnitude

SLRs than static targets; all $p < .0005$). Indeed, SLRs evoked by the fastest moving target in the

emerging target paradigm were approximately twice as large as SLRs evoked by the target in the

static paradigm (S1 versus 20 cm/s; unpaired t-test; $t = -7.78$, $p = 3.61e-7$, $df = 18$). Even when

removing the influence of both static targets from our analysis (s1 and 0cm/s), we found that

faster moving targets in the emerging target paradigm promoted larger magnitude SLRs (one-

way ANOVA (10, 15 and 20 cm/s speeds), $F_{(2, 32)} = 4.59$, $p = .0183$).

In **Figure 5b** (blue rectangle), we noticed differences between the SLR magnitude in the

static paradigm (S1) versus that driven by a non-moving (0 cm/s) stimulus in the emerging target

paradigm. In these cases, the stimulus presented to the visual system is the same, however SLRs

in the emerging target paradigm were almost twice as large (unpaired t-test; $t = -6.1$, $p = 3.02e-6$,

$df = 23$). Recall that the SLR prevalence was also higher to the 0 cm/s target in the emerging

target paradigm. Thus, depending on the context in which stimuli are presented, they may evoke

different SLRs.

Finally, we were interested in how SLR magnitude in the emerging target task related to

the maximal movement aligned activity. Previous reports using static tasks observed relatively

modest SLR magnitudes which were ~10-20% as large as muscle activity just prior to movement

onset (Pruszynski et al., 2010; Gu et al., 2016; Kozak et al., 2019). As shown in **Figure 5c**, SLRs

could be ~50% (range: 10-90%) as large as the peak of such movement-aligned activity. This is a

notable increase in magnitude compared to previous reports. When measured this way, we did

not observe any tendency for this scaled measure of SLR magnitude to vary across condition

(one-way ANOVA, $F(4,53)=.62$, $p = .65$).

Overall, targets presented in the emerging target paradigm readily evoked SLRs,

regardless of whether they were moving or not. SLRs evoked at similar latencies, however faster

moving targets evoked larger magnitude SLRs that occasionally approached levels of movement-

aligned activity. We also found that presentation of static targets in the emerging target versus

static paradigm evoked larger and more prevalent SLRs, attesting to the influence of task context

in favoring SLR expression.

Experiment 2: High-contrast, fast moving targets evoke shorter latency, larger magnitude SLRs

The most intriguing finding in experiment 1 was that all targets evoked similar latency SLRs.

Conversely, all previous reports manipulating visual features found that larger SLR magnitudes

were associated with shorter latencies (Wood et al., 2015; Kozak et al., 2019). Lower contrast

targets evoke longer latency SLRs (Wood et al., 2015). Thus, to further investigate the

relationship between SLR latency and magnitude, in experiment 2 we examined the SLR in

response to high and low contrast targets moving at a 'fast' (20 cm/s) or 'slow' (10 cm/s) speed.

Data from a representative subject in experiment 2 is shown in **Figure 6** (same format as

Figure 4). This figure illustrates EMG data from both static and emerging target paradigms

(compare first two rows to final four rows). In this subject, SLRs in the emerging target

paradigm evolved at shorter latencies and attained larger magnitudes in response to high contrast

408	targets (e.g., compare the timing and intensity of EMG activity in the grey box in color plots, or
409	the timing and peak of mean EMG activity in the SLR interval in the 3rd column). As expected,
410	in this subject high contrast targets appear to evoke shorter latency and larger magnitude SLRs
411	than low contrast moving targets. We therefore quantified differences in SLR prevalence, latency
412	and magnitude across the population.
413	As in experiment 1, all subjects exhibited an SLR in at least 1 condition (indicated by a
414	diamond above the dotted line in **Figure 3**). In the emerging target paradigm, we found that high
415	contrast targets were more likely to evoke SLRs than low contrast targets (chi-squared test: p=
416	0.00075, $\chi2$= 11.361, df=1). In fact, low contrast moving targets were no better at evoking SLRs
417	than static targets (slow + fast low contrast moving targets versus high + low contrast static
418	targets: chi-squared test: p>.05, $\chi2$= .056, df=1). Consistent with results from experiment 1,
419	target speed did not impact SLR prevalence (chi-squared, all p>.5).
420	We next examined SLR latency, and found that target contrast, but not target speed,
421	impacted SLR latency (**Figure 7a**; 2 x 2 ANOVA, $F(1,47) = 72.85$, $p = 4.1e-11$; note that there
422	is no difference within green or blue data points). High contrast targets evoked shorter latency
423	SLRs than low contrast targets (**Figure 7a**; $p = 2.07e-8$, mean within-subject difference ~14 ms).
424	These results confirm our findings from experiment 1, indicating that target speed does not
425	impact latency.
426	In contrast to experiment 1, we noticed that SLRs evoked in the static task had longer
427	latencies than those in the emerging target paradigm (**Figure 7a;** unpaired t-test; $t(18) = -3.27$, p
428	= .004; compare high and low contrast purple data with respective green and blue points). This
429	differed from what we saw in experiment 1, where SLR latencies did not change. We suspect
430	that these differences may relate to the predictability of the time of target appearance (Contemori

et al., 2021). Recall that the time of target appearance was fully predictable in experiment 1, but

randomized in experiment 2 in order to match a previous study (Wood et al., 2015). These results

indicate that emerging targets evoke shorter latency SLRs compared to static targets, if the

timing of target appearance is unpredictable.

We next examined the influence of target contrast and speed on SLR magnitude (**Figure**

7b). Here, we found that both target contrast and speed impacted SLR magnitude (2 x 2

ANOVA, speed: $F(1,47)=67.8$ p = 1.1e-10; contrast: $F(1,47)=11.7$ p = .001). When matched for

speed, higher contrast targets evoked larger magnitude SLRs (**Figure 7b**, all blue points lower

than all green points; unpaired t-test; t= 2.7, p= .01, df= 49). We also found that faster moving

targets evoked larger magnitude SLRs (unpaired t-test; t= -7.8, p= 4.1e-10, df= 49). This

supports the results from experiment 1, where faster moving targets evoked larger magnitude

SLRs. We did not observe interaction effects between target speed and contrast (2 x 2 ANOVA,

p=.87). We also did not observe any differences in magnitude for static targets.

Next, we compared SLR magnitude to movement-aligned activity (**Figure 7c**) and as

seen in experiment 1, observed large variability (**Figure 5c**). However, SLR magnitudes tended

to be larger than in experiment 1, reaching ~75% of movement-aligned activity (~20% higher

than experiment 1; compare **Figures 5c** versus **7c**). In a few cases, SLRs exceeded movement-

aligned activity, in the most extreme case approaching 150%. An example of this is shown in

Figure 8, plotting EMG relative to either stimulus onset (illustrating the SLR; **Figure 8a**) or

movement onset (illustrating reach-aligned activity; **Figure 8b**). This figure indicates that

recruitment during the SLR interval can exceed movement-aligned activity. **Figure 8a** also

provides an example of an SLR that is crisp and distinct from subsequent recruitment, in contrast

to the SLRs shown in **Figures 4** and **6**, which blend more smoothly into the ensuing phase of

reach-related activity.

To summarize, the results from experiment 2 confirmed and extended those from

experiment 1, in showing that SLR latency was not influenced by target speed, regardless of

target contrast. High contrast, fast moving targets improve SLR expression compared to static

targets. Furthermore, we report that SLR magnitude may rival, and even exceed activity attained

just before movement onset.

The SLR and its relationship to reaching behaviour

We further explored the relationship between SLR latency, magnitude, and the ensuing RT. In

both experiments, SLR latency and magnitude were negatively correlated with each other

(**Figure 9a** exp. 1; R=-.48, p=.00012; **Figure 9d** exp. 2; R=-.41, p=.00038), indicating that larger

magnitude SLRs are associated with shorter latency SLRs. Recall that there were no significant

latency differences for any target conditions found in experiment 1. This negative correlation

indicates that short latency SLRs, even with small variability and a narrow range of latencies (all

SLR latencies between ~90-110 ms **Figure 9a**), still tend to be associated with larger magnitude

SLRs.

When examining how the SLR relates to RT, we found a positive correlation between

latency and RT in both experiments (experiment 1- **Figure 9b**; R=.36, p=.0047; experiment 2-

Figure 9e; R=.67, p=8.4e-11); shorter latency SLRs were correlated with more rapid RTs.

Conversely, there was a strong negative correlation between SLR magnitude and RT for both

experiments (experiment 1- **Figure 9c**; R=-.75, p=8.69e-12; experiment 2- **Figure 9f**; R=-.40,

p=.0005), indicating that large magnitude SLRs were associated with more rapid RTs. Thus,

robust SLRs are correlated with expedited RTs.

Across both experiments, we found that faster moving (experiment 1) or fast moving high

contrast targets (experiment 2) were associated with more rapid RTs (see **Figure 10**). In

experiment 1, target speed influenced RTs (one-way ANOVA, $F(4, 79)= 2.71$, $p= .036$), with

RTs decreasing by ~20 ms (static versus fastest; 250ms+/-25 versus 231ms+/- 16; paired t-test;

t= 3.25, p=.005, df= 15). This trend may also be seen in **Figure 4,** which demonstrates a leftward

shift of RTs when moving from upper to lower color plots. In experiment 2, both target speed

(two-by-two ANOVA, $F(1, 107)= 17.33$, $p= 6.5e-5$) and contrast (two-by-two ANOVA, $F(2,$

$107)= 103.07$, $p= 3.25e-25$) influenced RTs. RTs decreased by ~62 ms for high contrast targets

(static versus fastest; 243ms+/-26 versus 181ms+/- 9; paired t-test; t= 10.2, p=5.7e-9, df= 17) or

~57 ms for low contrast targets (static versus fastest; 253ms+/-26 versus 196ms+/- 11; paired t-

test; t= 9.7, p=2.2e-8, df= 17). Importantly, changes in RTs (**Figure 10**) resembled changes in

the latency and magnitude of the SLR (e.g., compare Figure 10a to Figure 5, and Figure 10b to

Figure 7). These results indicate a strong relationship between robust SLRs and rapid RTs.

When further examining **Figure 10**, we noticed RT differences between experiments 1

and 2 (~50 ms), despite both using 'high contrast' fast moving targets. These same differences

were present in the SLR (**Figure 5a** and **Figure 7a**; unpaired t-test; t= 7.99, p= 2.35e-8, df= 25;

~15 ms). Recall that experiment 1 and 2 were performed in different apparatuses, with different

monitors. We suspect he 'high contrast' target in experiment 1 is similar to the low contrast

target in experiment 2 (see discussion). In support of this, we found no difference between

latencies for low contrast targets in experiment 2 and latencies in experiment 1 (unpaired t-test;

t= 1.2, p= .22, df= 25).

Recall in experiment 1 that we presented two different static conditions, one in the

standard static task (S1), and one in the context of an emerging target paradigm (0 cm/s) which

produced larger and more prevalent SLRs. When examining RTs between these conditions, we

observed similar RTs in both tasks (250 +/- 22 ms versus 250 +/- 25 ms; n.s.). Thus, despite

differences in SLRs, we observed no differences in RTs. Given the correlation of SLRs to rapid

RTs (see above), this result indicates that factors which impact the SLR do not always impact RT

in the same way.

To summarize, robust SLRs are correlated with rapid RTs. We found that RTs toward

high contrast, fast moving targets decreased by as much as 62 ms. However, we observed

conditions where equivalent SLRs preceded different RTs, as well as conditions where different

SLRs preceded equivalent RTs. Thus, although robust SLRs promote shorter-latency RTs, RTs

are also influenced by periods of muscle recruitment after the SLR.

Discussion

The brain has a remarkable capacity to rapidly transform vision into action, allowing us to

initiate reaches towards a target within fractions of a second. Previous research has shown that

some visual stimuli, such those composed of lower spatial frequencies or higher contrast, elicit

earlier reaching responses than others (Veerman et al., 2008; Wood et al., 2015; Kozak et al.,

2019). Here, by using a new emerging target paradigm (Kozak et al., 2020), we show that high-

contrast, moving stimuli are particularly capable of eliciting fast visuomotor responses such as

SLRs and expedited reach reaction times. Further, since subjects performed both the emerging

target paradigm and a visually-guided reach paradigm used previously, we can better understand

the sensory and cognitive factors promoting fast visuomotor responses. Finally, by studying a

variety of stimuli in the emerging target paradigm, we can gain further insights into the

relationship between the earliest phase of stimulus-related EMG recruitment, subsequent phases

of EMG recruitment leading up to movement onset, and the RT of the movement itself.

Together, our results contribute to a growing understanding of the interplay between the cortical

and subcortical substrates mediating visually-guided reaches in conditions requiring expedited

responses.

Comparison to past studies

Previous research on rapid visually-guided reaches have typically used either target-jump

paradigm where a visual target is displaced during an arm movement (Soechting and Lacquaniti,

1983; Day and Lyon, 2000), or a static target reaching paradigm where a suddenly appearing

target evokes reaches from a stationary arm position (Pruszynski et al., 2010). While useful, such

approaches have limitations: the response of interest evolves during an on-going movement in

target-jump experiments, or not all subjects exhibit SLRs in a typical static target paradigm even

with intramuscular electrodes (Pruszynski et al., 2010; Gu et al., 2016). In contrast, in the

emerging target paradigm, the arm is stationary at the time of target emergence and prevalent fast

visuomotor responses can be recorded even with surface electrodes (Contemori et al., 2021;

Kozak et al., 2020).

The results in experiment 1 show that larger SLRs are provoked by the appearance of a

stationary target in the emerging target paradigm rather than the static task (blue rectangle,

Figure 5, ~2 times larger). Larger magnitude SLRs likely relate to the increased proportion of

detectable SLRs in the emerging target paradigm rather than static task (100% vs. 56%). We

speculate that the increase in SLR magnitude and prevalence in the emerging target paradigm is

due to a combination of a signal of implied motion of the target behind the barrier (Kozak et al.,

2020) and certainty of target appearance (Contemori et al. 2021). Implied motion is critical, as a

recent study presenting only moving targets did not report SLRs (Lara et al., 2018). Further,

temporally predictable targets still have to be presented within the emerging target paradigm, as

smaller magnitude and less prevalent SLRs are elicited in the static tasks used here despite a

predictable gap between the offset of a central hold stimulus and target presentation (Pruszynski

et al., 2010; Wood et al., 2015).

Past work has shown that stimuli composed of progressively higher contrast (Wood et al.,

2015) or lower spatial frequencies (Kozak et al., 2019) elicit progressively larger magnitude and

lower latency SLRs. Somewhat surprisingly, manipulating stimulus velocity did not elicit this

inverse relationship between SLR latency and magnitude. Instead, SLR latency remained

invariant while SLR magnitude was greatest for the fastest moving stimulus; this was observed

both in experiment 1 (**Figure 5**) and in experiment 2 for stimuli of similar contrast (**Figure 7**).

This finding demonstrates that the latency and magnitude of the SLR are dissociable. We

speculate that this dissociation relates to the displacement of the stimulus across sequential

display frames, with larger displacements increasing the vigor of sensory transients that remain

timed to the onset of the first frame. Future studies should examine the impact of subsequent

display frames on the SLR, given that Contemori and colleagues (2021) found that a stimulus

flashed for a single frame (~16 ms) provokes earlier and larger SLRs than moving targets; it is

possible that flashed stimuli may provoke particularly vigorous sensory transients from the

retina.

Our observations of scaled EMG responses for faster moving targets resembles the

phenomenon of speed coupling in the interception timing literature, where participants adjust

hand velocity to match target velocity (Savelsbergh et al., 1992; van Donkelaar et al., 1992;

Carnahan and McFadyen, 1996; Brouwer et al., 2000, 2002, 2003). Speed coupling may reflect

the temporal accuracy required to intercept targets moving at different speeds, rather than direct

coupling to the speed of the target (Brouwer et al., 2000, 2002; Tresilian and Lonergan, 2002;

Caljouw et al., 2004). Aspects of our results may be consistent with speed coupling, particularly

since target motion before disappearance cues the subject about upcoming target velocity. On the

other hand, the extremely short latency of the SLR, regardless of target speed, is consistent with it

arising from a feed-forward response. Future work should examine SLR magnitude across target

velocity in the absence of any predictability, and compare such SLRs to subsequent recruitment

preceding the RT.

Relating the SLR to subsequent RTs

SLRs represent the first change in muscle recruitment following visual stimulus presentation. As

such, they demarcate the completion of the most rapid sensory-to-motor transformation for

visually-guided reaching. Past reports show that larger magnitude SLRs precede shorter latency

movements on a trial-by-trial basis (Pruszynski et al., 2010; Gu et al., 2016; Contemori et al.,

2021), and here we show, when pooling across all conditions in which subjects exhibited an

SLR, that larger magnitude and earlier SLRs precede shorter reach RTs (**Figure 9;** see also

Kozak et al., 2019). These correlative relationships, are consistent with a functional contribution

of the SLR to the forces necessary to overcome the limb's inertia. However, the SLR is often

followed by sustained levels of muscle recruitment preceding the RT. Consequently, muscle

recruitment during the SLR interval is not fully predictive of the RT. Nearly identical SLRs can

precede different RTs (e.g., in Experiment 1, different RTs toward 0 deg/s and 20 deg/s stimuli

despite similar SLRs), or different SLRs can precede the same RTs (e.g., in Experiment 2,

similar RTs toward fast moving low-contrast or slow moving high-contrast stimuli despite

different SLRs). These distinctions between SLRs and RTs attest to continued recruitment after

the SLR, and the contribution of such recruitment to overcoming the limb's inertia. Judicious

selection of stimulus parameters, e.g., to elicit robust SLRs but longer RTs, may aid the study of

the fast visuomotor responses by reducing the overlap between the SLR and activity aligned to

movement onset.

Previous reports of SLRs emphasized their relatively small magnitude, reaching ~10-20%

of the volitional recruitment achieved before movement onset. Such modest levels of recruitment

during the SLR interval can induce subtle motion toward the target, even if a subsequent

movement is withheld or proceeds in a different direction (Gu et al., 2016; Atsma et al., 2018). A

remarkable aspect of the current results is that we occasionally observed SLR recruitment that

equaled if not exceeded volitional, movement-related activity (**Figure 8**). As mentioned above,

increased SLRs magnitude likely relates to the increased propensity of detecting an SLR.

Further, for many subjects, the shortest RTs occurred essentially right after the SLR interval

(e.g., see shortest RT trials for the 20 cm/s targets in **Figures 4** and **6;** trials in lowest rows). In

such trials, EMG recruitment during the SLR interval is clearly sufficient to overcome the limb's

inertia to produce a detectable RT. On such trials, any distinction between the SLR and

subsequent volitional recruitment is blurred; phases of recruitment that are separated for longer-

RT movements are essentially indistinguishable for shorter-RT movements. We are struck by the

resemblance of the single-trial EMG data shown in **Figures 2, 4, and 6** to raster plots of activity

of the superior colliculus during express and non-express saccades (Edelman and Keller, 1996;

Dorris et al., 1997; Sparks et al., 2000). For shorter latency movements, be they express saccades

or short-RT movements in the emerging target paradigm, phases of recruitment tied to stimulus

onset merge into those phases of recruitment loosely termed 'volitional' for longer-latency

movements.

Plausible neural substrates

The superior colliculus (SC) has been implicated as a potential candidate mediating the SLR

(Corneil et al., 2004; Pruszynski et al., 2010). The SC is a midbrain structure whose intermediate

and deep layers (SCi) interface with premotor structures for eye, head, and upper limb

movements within the reticular formation (Corneil and Munoz, 2014), whereas superficial layers

respond to visual input. Our results add further circumstantial evidence for a role for the SC in

the SLR. Previous work established that both the SLR and visual responses in the SC

preferentially respond to low spatial frequency (Chen et al., 2018; Kozak et al., 2019) and high

contrast targets (Li and Basso, 2008; Marino et al., 2015; Wood et al., 2015). Moving targets also

evoke robust SC responses (Marrocco and Li, 1977), and we have shown here that moving

targets also evoke robust SLRs. Importantly, the target speeds used in the current study resemble

those used in the Marrocco and Li (1977) study, which evoked robust responses in the

intermediate and deep layers of the SC where visuomotor and motor neurons activated before

arm movements reside (Werner et al., 1997; Stuphorn et al., 1999; Philipp and Hoffmann, 2014).

The hypothesis that the SC mediates the SLR requires further testing with casual manipulation in

tasks that promote SLR expression. Work reporting that SCi inactivation impacted the selection

of reach targets but not reach kinematics (Song et al., 2011) employed paradigms that would not

promote SLR expression.

An open question is how visual input from target motion accesses the SC, and how

features of the emerging target task potentiate SLRs. Visual information may access the

superficial SC directly via a retinotectal route (summarized in: (May, 2006)), or indirectly via a

retino-geniculo-striate loop through cortex, and then on to the SC (Schiller et al., 1979). The SC

also receives projections from higher-order cortical centers such as MT ((Fries, 1984; Nhan and

Callaway, 2012); an area which responds to target motion). As MT response latencies to visual

stimuli are ~60-70 ms in NHPs (Schmolesky et al., 1998; Smith et al., 2005), which are

concurrent with SLR latencies on NHP neck muscles (Corneil et al., 2004), motion processing in

MT may not be directly involved in relaying visual transients that evoke the SLR. Instead, as

implied motion strongly activates MT (Krekelberg et al., 2005), top-down signals from MT may

strongly activate pre-target activity in the SC, increasing the vigor of the SC's response when the

target appears beneath the occluder, and subsequently increasing the magnitude of the SLR.

Methodological considerations

All of our subjects exhibited an SLR in at least one stimulus condition. This replicates the high

prevalence of SLR detection in the emerging target paradigm (Kozak et al., 2020; Contemori et

al., 2021), and affirms the contention that most if not all subjects are capable of exhibiting SLRs

in an appropriate paradigm. However, subjects did not necessarily exhibit an SLR to every

stimulus in the emerging target paradigm, even for high-contrast, fast moving stimuli (**Figure 2**).

We speculate that this result may be due, at least in part, to how we detect the presence of the

SLR with the RT split analysis, which allows for a direct comparison of SLR prevalence to those

reported elsewhere which also used this method (e.g.,(Kozak et al., 2019, 2020; Contemori et al.,

2021)). The RT split analysis is somewhat conservative in that it requires an SLR to both short-

and long-latency trials. SLR- observations could therefore include those situations where an SLR

precedes the short- but not long-RT trials (see top two rows of **Figure 4**, which appear to exhibit

SLRs for the shorter-than-average RTs). The RT split analysis is also more suited to paradigms

engendering a larger degree of RT variance. As noted previously (Kozak et al., 2020), one

potential complication of the emerging target paradigm is that it may shorten RTs too much, not

only reducing RT variance but also leading to a potential overlap of SLR with activity linked to

movement onset. As noted above, judicious stimulus selection may help avoid such overlap, but

we also expect the continued evolution of SLR detection methods suited to a particular

paradigm.

We noted large differences in the latencies of fast visuomotor responses across

experiments, with SLRs evolving ~15 ms earlier and reach RTs being ~50 ms shorter in

experiment 2 compared to experiment 1. The most likely reason for these differences is the

contrast of the stimuli, as the VPIXX projector used in experiment 2 has a higher level of

contrast (275 cd/m2) than the standard monitor used in experiment 1 (110 cd/m2). Consistent

with this, the SLR latencies from the high-contrast target in Experiment 1 (**Figure 5b**) resemble

those of the low contrast target in experiment 2 (**Figure 7a**). The large difference in reach RTs

across experiments, even to targets associated with similar latency SLRs, may also relate to

postural differences in the exoskeleton-based Kinarm robot in experiment 1 versus the endpoint-

based Kinarm robot in experiment 2.

Conclusions

The work presented here demonstrates that all subjects can generate SLRs, but are more likely to

do so in an emerging target paradigm. As a consequence, reliable SLRs can be recorded even

with surface EMG, and here we use this paradigm to show that targets moving at faster speeds

elicit larger magnitude SLRs, whereas higher contrast targets decrease SLR latency. Larger

magnitude and shorter-latency SLRs tend to precede expedited RTs. As the SLR is the first wave

of muscle recruitment tied to visual target onset, muscle recruitment during this interval may

provide a unique window into the contribution of the tecto-reticulospinal signaling that is distinct

from subsequent phases of target-related recruitment of corticospinal or cortico-reticulo-spinal

circuits. Rather than being directly involved in the SLR, cortical areas such as MT may serve to

prime this circuit to produce the first wave of muscle recruitment in situations when time is of

the essence.

References

Atsma J, Maij F, Gu C, Medendorp WP, Corneil BD (2018) Active Braking of Whole-Arm

692   Reaching Movements Provides Single-Trial Neuromuscular Measures of Movement

693   Cancellation. J Neurosci 38:4367–4382.

694   Brouwer A-M, Brenner E, Smeets JBJ (2002) Hitting moving objects: is target speed used in

695   guiding the hand? Exp Brain Res 143:198–211.

696   Brouwer A-M, Middelburg T, Smeets JBJ, Brenner E (2003) Hitting moving targets: a

697   dissociation between the use of the target's speed and direciton of motion. Exp Brain Res

698   152:368–375.

699   Brouwer AM, Brenner E, Smeets JBJ (2000) Hitting moving objects. The dependency of hand

700   velocity on the speed of the target. Exp Brain Res 133:242–248.

701   Caljouw S, van der Kamp J, Savelsbergh GJP (2004) Chapter 19 The fallacious assumption of

702   time-to-contact perception in the regulation of catching and hitting. In: Advances in

703   Psychology, pp 443–474.

704   Carnahan H, McFadyen BJ (1996) Visuomotor control when reaching toward and grasping

705   moving targets. Acta Psychol (Amst) 92:17–32.

706   Chen C-Y, Sonnenberg L, Weller S, Witschel T, Hafed ZM (2018) Spatial frequency sensitivity

707   in macaque midbrain. Nat Commun 9:2852.

708   Contemori S, Loeb GE, Corneil BD, Wallis G, Carroll TJ (2021) The influence of temporal

709   predictability on express visuomotor responses. J Neurophysiol 125:731–747.

710   Corneil BD, Munoz DP (2014) Overt Responses during Covert Orienting. Neuron 82:1230–

711   1243.

Corneil BD, Olivier E, Munoz DP (2004) Visual responses on neck muscles reveal selective

gating that prevents express saccades. Neuron 42:831–841.

Day BL, Lyon IN (2000) Voluntary modification of automatic arm movements evoked by

motion of a visual target. Exp Brain Res 130:159–168.

Dorris MC, Paré M, Munoz DP (1997) Neuronal activity in monkey superior colliculus related to

the initiation of saccadic eye movements. J Neurosci 17:8566–8579.

Edelman JA, Keller EL (1996) Activity of visuomotor burst neurons in the superior colliculus

accompanying express saccades. J Neurophysiol 76:908–926.

Fries W (1984) Cortical projections to the superior colliculus in the macaque monkey: a

retrograde study using horseradish peroxidase. J Comp Neurol 230:55–76.

Glover IS, Baker SN (2019) Multimodal stimuli modulate rapid visual responses during

reaching. J Neurophysiol 122:1894–1908.

Gu C, Wood DK, Gribble PL, Corneil BD (2016) A Trial-by-Trial Window into Sensorimotor

Transformations in the Human Motor Periphery. J Neurosci 36:8273–8282.

Kozak RA, Cecala AL, Corneil BD (2020) An Emerging Target Paradigm to Evoke Fast

Visuomotor Responses on Human Upper Limb Muscles. J Vis Exp:e61428.

Kozak RA, Kreyenmeier P, Gu C, Johnston K, Corneil BD (2019) Stimulus-Locked Responses

on Human Upper Limb Muscles and Corrective Reaches Are Preferentially Evoked by Low

Spatial Frequencies. eneuro 6:ENEURO.0301-19.2019.

Krekelberg B, Vatakis A, Kourtzi Z (2005) Implied motion from form in the human visual

cortex. J Neurophysiol 94:4373–4386.

Lara AH, Elsayed GF, Zimnik AJ, Cunningham JP, Churchland MM (2018) Conservation of

preparatory neural events in monkey motor cortex regardless of how movement is initiated.

eLife 7:e31826.

Li X, Basso MA (2008) Preparing to Move Increases the Sensitivity of Superior Colliculus

Neurons. J Neurosci 28:4561–4577.

Marino RA et al. (2015) Linking express saccade occurance to stimulus properties and

sensorimotor integration in the superior colliculus. J Neurophysiol 114:879–892.

Marino RA, Levy R, Boehnke S, White BJ, Itti L, Munoz DP (2012) Linking visual response

properties in the superior colliculus to saccade behavior. Eur J Neurosci 35:1738–1752.

Marrocco RT, Li RH (1977) Monkey superior colliculus: properties of single cells and their

afferent inputs. J Neurophysiol 40:844–60.

Maunsell JH, Gibson JR (1992) Visual response latencies in striate cortex of the macaque

monkey. J Neurophysiol 68:1332–1344.

May PJ (2006) The mammalian superior colliculus: laminar structure and connections. In:

Progress in Brain Research, pp 321–378.

Nhan HL, Callaway EM (2012) Morphology of superior colliculus- and middle temporal area-

projecting neurons in primate primary visual cortex. J Comp Neurol 520:52–80.

Philipp R, Hoffmann K-P (2014) Arm movements induced by electrical microstimulation in the

superior colliculus of the macaque monkey. J Neurosci 34:3350–3363.

Pruszynski AJ, King GL, Boisse L, Scott SH, Randall Flanagan J, Munoz DP (2010) Stimulus-

locked responses on human arm muscles reveal a rapid neural pathway linking visual input

to arm motor output. Eur J Neurosci 32:1049–1057.

Pruszynski JA, Gribble PL, Corneil BD (2018) Done in 100 ms: path-dependent visuomotor

transformation in the human upper limb. J Neurophysiol 119:1319–1328.

Rezvani S, Corneil BD (2008) Recruitment of a head-turning synergy by low-frequency activity

in the primate superior colliculus. J Neurophysiol 100:397–411.

Savelsbergh GJP, Whiting HTA, Burden AM, Bartlett RM (1992) The role of predictive visual

temporal information in the coordination of muscle activity in catching. Exp Brain Res

89:223–228.

Schiller PH, Malpeli JG, Schein SJ (1979) Composition of geniculostriate input to superior

colliculus of the rhesus monkey. J Neurophysiol 42:1124–33.

Schmolesky MT, Wang Y, Hanes DP, Thompson KG, Leutgeb S, Schall JD, Leventhal AG

(1998) Signal timing across the macaque visual system. J Neurophysiol 79:3272–3278.

Smith MA, Majaj NJ, Movshon JA (2005) Dynamics of motion signaling by neurons in macaque

area MT. Nat Neurosci 8:220–228.

Soechting JF, Lacquaniti F (1983) Modification of trajectory of a pointing movement in response

to a change in target location. J Neurophysiol 49:548–564.

Song J-H, Rafal RD, McPeek RM (2011) Deficits in reach target selection during inactivation of

the midbrain superior colliculus. Proc Natl Acad Sci 108:E1433–E1440.

Sparks D, Rohrer WH, Zhang Y (2000) The role of the superior colliculus in saccade initiation: a

study of express saccades and the gap effect. Vision Res 40:2763–2777.

Stuphorn V, Hoffmann KP, Miller LE (1999) Correlation of primate superior colliculus and

reticular formation discharge with proximal limb muscle activity. J Neurophysiol 81:1978–

1982.

Tresilian J, Lonergan A (2002) Intercepting a moving target: effects of temporal precision

constraints and movement amplitude. Exp Brain Res 142:193–207.

van Donkelaar P, Lee RG, Gellman RS (1992) Control strategies in directing the hand to moving

targets. Exp brain Res 91:151–161.

Veerman MM, Brenner E, Smeets JBJ (2008) The latency for correcting a movement depends on

the visual attribute that defines the target. Exp Brain Res 187:219–228.

Werner W, Dannenberg S, Hoffmann K-P (1997) Arm-movement-related neurons in the primate

superior colliculus and underlying reticular formation: comparison of neuronal activity with

EMGs of muscles of the shoulder, arm and trunk during reaching. Exp Brain Res 115:191–

205.

Wood DK, Gu C, Corneil BD, Gribble PL, Goodale MA (2015) Transient visual responses reset

the phase of low-frequency oscillations in the skeletomotor periphery. Eur J Neurosci

42:1919–1932.

Figure legends

Figure 1. The emerging target paradigm (Kozak et al., 2020). a) A target (T1) appears above an

occluder (grey box) and the subject moves the real-time cursor (RTC; white dot below occluder

that represents hand position), into the start position (T0; grey dot below the occluder). b) Once

the RTC is in the start position, T1 moves towards the subject along the inverted y path and then

disappears behind the occluder. c) The subject makes a visually-guided reach to T1 as soon as it

emerges at the right or left outlet below the occluder.

Figure 2. Detection of SLRs. a) Exemplar trial-by-trial recruitment of activity from right

pectoralis muscle during leftward and rightward reaches from a subject with a detectable SLR.

Each row is a different trial, with the intensity of color conveying degree of recruitment. Trials

sorted by RT (white boxes) and aligned to stimulus onset. The SLR (highlighted by asterisk)

appears as a vertical banding of increased or decreased recruitment aligned to left or right

stimulus presentation, respectively, rather than movement onset. b) Mean +/- SE of EMG activity

for the early- (purple) and late- (green) RT subsets based on mean RT. c) In the RT split analysis,

separate time-series ROCs were run for the early and late subsets, and then the mean RT for each

subset is plotted as a function of the discrimination time. The slope of the line connecting these

points determines whether the divergence of EMG activity is more time-locked to stimulus or

movement onset. d) For SLR+ observations, all trials were pooled together regardless of RT to

determine the SLR latency (derived from the time-series ROC) and magnitude (derived from the

mean EMG waveforms; see methods). Grey rectangle in b) and d) depict the SLR interval.

Figure 3. Prevalence of SLRs across the population. Slope of the relationship between

discrimination time and average RT for early and late RT groups (e.g. see slope in Figure 2c)

Each data point represents a unique subject for a particular condition. Slopes are capped at 90

degrees and a dotted black line is drawn at 67.5 deg. A data point above this line represents a

SLR+ observation, whereas a data point below the line represents SLR- observations. a)

Experiment 1. Black and grey diamonds represent static conditions in the static and emerging

target paradigms, respectively. Blue diamonds represent moving targets in emerging target

paradigm, with slower moving targets are represented by darker shades. b) Experiment 2. Purple

diamonds represent high contrast targets; green diamonds represent lower contrast targets. Static

and slower moving targets are represented by darker shades. 2 subjects completed both

experiments; subjects 2 and 16 from experiment 1 are subjects 18 and 12 from experiment 2,

respectively.

Figure 4. Data from a representative subject in experiment 1, using the same format as in figure

2. From left to right, columns shown trial-by-trial EMG activity for left and right targets sorted

by RT, mean +/- SE EMG activity for left and right targets with overlaid time-series ROC, and

the RT split analysis for data shown in the corresponding row (the number in each panel is the

slope).

Figure 5. Summary of results from experiment 1, showing (a) SLR latency, (b) SLR magnitude

normalized within each subject to the largest SLR across all conditions, and (c) SLR magnitude

normalized to the maximum of movement-aligned EMG activity. In c), each asterisk represents

data from an individual subject. S1 indicates the results from the static task, which presents the

same retinal stimulus to the subject as the 0 cm/s target in the emerging target paradigm. Error

bars show SE. * indicates significance (see results text for more details).

Figure 6. Data from representative subject in experiment 2,using the same format as figure 4.

Figure 7. Summary of results from experiment 2 (same format as Fig. 5), with SLR latency (a),

SLR magnitude normalized to the within-subject largest SLR (b), and SLR magnitude

normalized relative to maximum of movement-aligned activity (c). All values presented as mean

+/- standard error. * indicates significance. HC = high contrast. LC = low contrast.

Figure 8. Example subject whose SLR magnitude exceeds the level of EMG activity aligned to

target onset. Data, shown as mean +/- SE, are aligned to stimulus onset in a), or movement onset

in b). SLR magnitude is measured as area between left and right reaches for the 30 ms interval

after the SLR latency (grey box, running from 73-103 ms relative to stimulus onset). For

movement-aligned activity, magnitude is measured as area 15 ms before and 15 ms after RT

onset (grey box in b).

Figure 9. Correlations for SLR latency, magnitude and associated reaction time in experiment 1

(top row) or 2 (bottom row). a, d) SLR latency and magnitude correlations. b, d) SLR latency

and RT correlations. c, f) SLR magnitude and RT correlations. All correlations shown are

significant. Each blue circle represents data from an SLR+ subject in a unique experimental

condition. Solid red lines indicate first order fit of the data.

Figure 10. Summary of RTs in experiment 1 (a) and 2 (b), following the formats in figures 5 and

7.

a Stabilize hand at start position

b target moves down path and disappears behind occluder

c subject reaches to intercept target from left or right outlet

Figure 1- Kozak and Corneil 2021

Figure 2- Kozak and Corneil 2021

Figure 3- Kozak and Corneil 2021

Figure 4- Kozak and Corneil 2021

Figure 5- Kozak and Corneil 2021

Figure 6- Kozak and Corneil 2021

Figure 7- Kozak and Corneil 2021

Figure 8- Kozak and Corneil 2021

Figure 9- Kozak and Corneil 2021

Figure 10- Kozak and Corneil 2021