1
2
3 **Crop modeling defines opportunities and challenges for drought escape, water capture, and**
4 **yield increase using chilling-tolerant sorghum**
5

6 Rubí Raymundo[1], Sarah Sexton-Bowser[2], Ignacio A. Ciampitti[2], and Geoffrey Morris[1]*
7
8 [1] Department of Soil and Crop Science, Colorado State University, Fort Collins, CO 80523 USA
9 [2] Department of Agronomy, Kansas State University, Manhattan, KS, 66506 USA
10
11 * Corresponding author: Geoff.Morris@colostate.edu

12
13 Submission date: 05/04/2021
14
15

ABSTRACT

Many crop species, particularly those of tropical origin, are chilling sensitive so improved chilling tolerance can enhance production of these crops in temperate regions. For the cereal crop sorghum (*Sorghum bicolor L.*) early planting and chilling tolerance have been investigated for >50 years, but the potential value or tradeoffs of this genotype × management change has not been formally evaluated with modeling. To assess the potential of early-planted chilling-tolerant grain sorghum in the central US sorghum belt, we conducted CERES-Sorghum simulations and characterized scenarios under which this change would be expected to enhance (or diminish) drought escape, water capture, and yield. We conducted crop growth modeling for full- and short-season hybrids under rainfed systems that were simulated to be planted in very early (April), early (May 15), and normal (June 15) planting dates over 1986–2015 in four locations in Kansas representative of the central US sorghum belt. Simulations indicated that very early planting will generally lead to lower initial soil moisture, longer growing periods, and higher evapotranspiration. Very early planting is expected to extend the growing period by 20% for short- or full-season hybrids, reduce evaporation during fallow periods, and increase plant transpiration in the two-thirds of years with the highest precipitation (mean > 428 mm), leading to 11% and 7% increase grain yield for short- and full-season hybrids, respectively. Thus, in this major sorghum growing region very early and early planting could reduce risks of terminal droughts, extend seasons, and increase rotation options, suggesting that further development of chilling tolerant hybrids is warranted.

Keywords: Stress tolerance, crop growth model, crop adaptation, plant breeding, agronomy.

INTRODUCTION

Sustainable agriculture in water-limited environments requires cropping systems with low water requirements and relatively high productivity per unit of water used (Tilman et al., 2002; Zhang et al., 2018). Breeding programs generally rely on multi-environment trials in the target environment to assess the performance of genotypes and evaluate novel traits. However, this approach has limitations: (i) resources to deploy such trials are often limited, (ii) it is difficult to reproduce and/or interpret findings from year to year due to spatio-temporal variability of environmental conditions, and (iii) in most of the cases the novel trait is not yet available in commercial varieties, so cannot be studied in relevant genetic backgrounds (Lenaerts et al., 2019). In this context, crop growth models complement field experiments (van Ittersum et al., 2003; Challinor et al., 2018) to help breeding programs evaluate the potential value of novel traits within an existing or novel cropping system (Aggarwal et al., 1997; Chenu et al., 2017; Cooper et al., 2002). Models can be used to evaluate the potential benefits and tradeoffs of combining new genotypes and new management approaches in the target production environments, by accounting for interactions of genotype, environment and management (G×E×M) (Chenu et al., 2013; Kholová et al., 2014; Singh et al., 2014).

Sorghum (*Sorghum bicolor* L. Moench) is one of the most drought resilient crops, grown worldwide for grain, forage, and biomass (Adegbeye et al., 2020). The crop originated in the warm semi-arid tropics of Africa (Doggett and Majisu, 1968) and diffused widely, including to semi-arid and sub-humid temperate regions of the world (Smith and Frederiksen, 2000). However, like many tropical-origin crops, sorghum is sensitive to chilling (0-16 °C) (Lyons, 1973; Taylor and Rowley, 1971). Chilling sensitivity may be an ancestral trait stemming from sorghum's origin and adaptation in warm climates, or a derived trait originating inadvertently during selection for non tannin and/or semi-dwarf traits (Marla et al., 2019). Following Vavilov's phytogeographic approach, seedling chilling tolerance was identified in high-latitude Asian (Stickler et al., 1962) and high-altitude African (Singh, 1985) landraces, and more recently the trait was genetically mapped (Burow et al., 2011; Chopra et al., 2017; Marla et al., 2019). However, this trait has remained a breeding target for >50 years and has not been deployed in commercial grain sorghum hybrids. Research and development on chilling-tolerant sorghum remains ongoing in Europe (Schaffasz et al., 2019), Asia (Mocoeur et al., 2015), Australia (Wylie, 2008), and the Americas (Marla et al., 2019).

In temperate production regions, chilling temperatures restrict the sorghum growing season from late spring to fall (Ercoli et al., 2004). In the semi-arid region of the US sorghum belt (Laingen, 2015), a commercial hybrid with early-chilling tolerance trait has the potential to change the agricultural landscape by shifting planting dates from late spring to early spring to hypothetically position the crop in better environments to ensure early establishment (Burow et al., 2011; Franks et al., 2006; Kapanigowda et al., 2013). In the US, the state of Kansas leads grain sorghum production with 40% of national production on 2.7 million acres, mostly under rainfed conditions (>90%) (Laingen, 2015). Early crop establishment could potentially extend the crop growing period, synchronizing crop water demand and soil water supply during the

grain filling to avoid terminal droughts. Furthermore, early planting practices can facilitate
integrating the crop into a double crop rotation to maximize farm productivity (Burow et al.,
2011). For these reasons, it is believed that the early-season chilling tolerance trait would make
the crop more competitive for farmers and an alternative option to outperform traditional crops
with high water demand and less adapted to dryland systems (Bhattarai et al., 2019).

Ongoing efforts to improve chilling tolerance in sorghum (Burke et al., 2019; Chopra et
al., 2017; Marla et al., 2019; Ostmeyer et al., 2020) are based on the hypothesis that early
planting of a sorghum hybrid with chilling tolerance provides a commercially relevant increase
in grain yield in the US by extending the season and avoiding late season drought. Due to the
lack of chilling tolerant commercial hybrids this hypothesis cannot be directly tested under field
conditions. However, this hypothesis can be formally evaluated with crop modeling. In this study
we used the CERES-Sorghum growth model (Jones et al., 2003; White et al., 2015) to test these
hypotheses *in silico*, and quantify the expected agronomic impact of an early-chilling tolerant
sorghum hybrids in the northern part of the U.S. sorghum belt. We find support for several of the
proposed benefits of early-planted chilling-tolerant sorghum, along with some unanticipated
tradeoffs that must be accounted for in the overall G×E×M strategy for this cropping system.

MATERIALS AND METHODS
The sorghum cropping system
Kansas has a precipitation gradient decreasing from east to west (Lin et al., 2017) (Figure 1a),
with Koppen-Geiger climate classification of continental-humid in the east and semi-arid climate
in the west (Pražnikar, 2017). This gradient in environmental conditions has substantially
influenced the management of this rainfed crop such as plant density and varietal selection.
Recommended plant density declines from east to west, and later maturity hybrids are
recommended in the east while earlier maturity hybrids are recommended in the west
(Roozeboom and Fjell, 1998). Given sorghum's chilling sensitivity and the common practice of
prioritizing planting of other crops (i.e maize), sorghum is one of the later-planted crops in the
system and consequently the crop can not take advantage of the cumulative growing degrees
units or thermal time accumulated in early spring (Figure 1b). The sorghum growing season
ranges from early-May to early-July and the main planting season concentrates in the second
week of June (Figure 1c) (Shroyer et al., 1998). Otherwise, maize is planted from early-April to
mid-June and the main planting season concentrates in mid-May (Figure 1c). Therefore,
introgressing chilling tolerance in sorghum would allow the crop to be planted in early spring.

Overview of the model
CERES-Sorghum (White et al., 2015) belongs to the family of crop models available in the
Decision Support Systems for Agro technology Transfer-Crop Simulation Model (DSSAT-CSM)
(Hoogenboom et al., 2019; Jones et al., 2003). The model simulates the daily dynamics of plant
growth, phenology development and partitioning affected by stress factors (White et al., 2015).
The daily growth is simulated via the potential carbon assimilation (PCARB) that is calculated
using the radiation use efficiency (RUE, g MJ^{-1}), photosynthetically active radiation (PAR, MJ
m^{-1} day^{-1}) and the fraction of PAR intercepted ($1 - exp^{(-k * LAI)}$) by the crop (Ritchie et al., 1998),

where k (unitless) represents the extinction coefficient and LAI (m_{leaf}^2 m_{ground}^{-2}) is the leaf area
index. The model simulates daily water stress factors that penalize the daily carbon assimilation
(SWFAC) and plant growth (TURFAC) (White et al., 2015). The model requires cultivar-
specific genotype parameters (G), environmental information (E), and crop management (M)
practices, which are described in detail below.

Genotype parameters

In the model, G, a set of genotype-specific parameters, represents differences among hybrids
(White et al., 2015) that affect phenology development and crop growth. In this study, the
difference between early and late season hybrids is represented by the parameter potential
duration from emergence to end of juvenile phase (P1) (Table 1). In the model the phenology
development is a function of temperature and daylength. The model simulates germination,
emergence, end of the juvenile phase, panicle initiation, end of flag leaf expansion, anthesis,
beginning of grain filling and physiological maturity. Each phenology stage is the result of the
summation of daily thermal time (DTT) calculated via the formula

$$DTT = (Tx + Tn) \times 2^{-1} - Tbase$$

where Tbase is the minimum temperature for thermal time accumulation, set to 8 °C (White et
al., 2015), and Tx and Tn are daily maximum and minimum temperature, respectively. Note,
given that the DTT uses a Tbase of 8 °C and does not implement a chilling damage routine (i.e.
there is no penalty to growth or development for temperatures below Tbase) we interpret
CERES-Sorghum as modeling a chilling-tolerant sorghum genotype by default. The summation
of DTT to reach a phenology stage is indicated in Table 1 (e.g. P1: Potential duration from
emergence to end of juvenile phase). The only phenology stage affected by both temperature and
daylength is the duration from the end of juvenile stage to panicle initiation, represented by
parameters P2R and P2O (Table 1). The plant leaf growth and panicle partitioning are
determined by parameter G1 and G2, respectively.

Environmental data

In the model, E comprises daily weather data and multilayer soil profile parameters. Daily
weather data, including precipitation (mm), solar radiation (MJ m^{-2} day^{-1}), and daily maximum
(°C) and minimum temperature (°C) was obtained for agronomic research sites (Manhattan,
Hays, Garden City, and Colby) from Kansas Mesonet from 1986 to 2015 (Kansas Mesonet,
2020). Soil profile information was downloaded from the web soil survey (Soil Survey Staff,
2020) and detailed soil characteristics such as soil texture (%), bulk density (g ml^{-1}), organic
carbon (%), pH, wilting point (LL), field capacity (DUL), saturation (SAT), and soil root growth
factor (SRGF) for each site are presented in Table S1. The model computes the differences
between DUL and LL, and between SAT and DUL, to predict the daily soil extractable water and
the water drained in each soil layer, respectively. And, the SRGF represents the physical or
chemical constraints in each soil profile.

Management practices

The model requires realistic agronomic management (M) data such as planting date, planting density, row spacing, fertilization, and water supply. In this study, the crop was simulated to be planted each year in April 15 (very early), May 15 (early), and June 15 (normal) at a row distance of 76 cm, a planting depth of 3 cm, and nitrogen fertilizer rates of 190 kg ha^{-1}. Plant density was 14 plants m^{-2} in Manhattan, 8 plants m^{-2} in Hays, and 6 plants m^{-2} in both Garden City and Colby. Given that sorghum grows mostly under dryland systems across Kansas, simulations were performed under rainfed conditions.

Seasonal simulation settings

As outlined above, information on G, E, and M were defined for each research site. Simulations under rainfed conditions started on the first day of each year with 80% of soil moisture, 3 Mg ha^{-1} of surface residue, and 2 Mg ha^{-1} of root residue. Evapotranspiration was calculated with the Priestley-Taylor/Ritchie formula (Priestley and Taylor, 1972). Soil water infiltration was computed with the capacity approach method (Ritchie et al., 1998) and soil evaporation was estimated with the Suleiman-Ritchie method (Suleiman and Ritchie, 2003). Dynamics of carbon and nitrogen were simulated with the CENTURY model.

Evaluation of model performance and the effect of Tbase on days to emergence

For normal planting dates (June), as shown in Figure 2, CERES-Sorghum has shown satisfactory predictions for anthesis, grain yield, and evapotranspiration across Kansas (Araya et al., 2017; Staggenborg and Vanderlip, 2005; White et al., 2015) using a Tbase of 8 °C. To demonstrate that simulations in early spring represent a chilling-tolerant sorghum, the model accuracy in predicting days to emergence for a chilling-tolerant genotype (Kaoliang), simulations were conducted to match field experiments using a Tbase of 6 °C, 8 °C, and 10 °C. In these scenarios, we expect that seedling emergence for simulations with a Tbase of 8 °C will be closer to field data. Simulations were set up for field experiments that were planted in April, May, and June in 2016–2018 at three sites in Kansas (Marla et al., 2019). For these simulations, cultivar specific-parameters (G) were obtained for full-season hybrids (80 days to anthesis and 140 days to physiological maturity) (Araya et al., 2018). The model accuracy was quantified using the root mean square error (RMSE) that determines the distance of model prediction from a perfect prediction (Wallach et al., 2014).

Unlike commercial grain sorghum hybrids with semi-dwarf stature and high sink capacity, kaoliang sorghum has undesirable traits for this cropping system such as tall plants and low harvest index (Marla et al., 2019). Due to these differences, the direct comparison of observed and simulated grain weight for these known chilling tolerant genotypes is not possible. To ensure that a Tbase of 8 °C represents a sorghum tolerant to chilling temperatures, when planted early, a sensitivity analysis was conducted for a Tbase ranging from 6 °C to 10 °C. Under this assumption, in the simulations the crop will require less time for seedling emergence in early planting dates. For sensitive analysis, simulations were conducted under well-watered conditions to remove the confounding effect of soil moisture on emergence. In our simulations

the genotype used across all planting dates have the same cultivar parameters (Table 1) so the
effect of planting date can be isolated.

Statistical analysis and interpretation

Statistical analyses were performed in the R statistical environment (R Core Team, 2017) with
lmer (Bates et al., 2015) for mixed linear models and visualized using ggplot (Villanueva and
Chen, 2019). Analysis scripts are provided in Supplemental File 1 and data are provided in
Supplemental File 2. The observed annual precipitation, averaged across locations, was used to
classify years into three "precipitation level" quantiles: "dry" years (161–428 mm), "moderate"
years (428–580 mm), and "wet" years (580–1380 mm). Note, this classification is not intended
as a formal envirotyping of water deficit scenarios (Chenu et al., 2013) but to reflect the type of
informal classification used by growers and agronomic practitioners in the region (Baeumler and
Gupta, 2020). The effects of planting date and the interaction of planting date and precipitation
levels on all variables were tested using analysis of variance (ANOVA), considering planting
date, precipitation level, and hybrid type as fixed factors, and location and year as random
factors. Tukey test for the interaction of planting date and precipitation levels was performed
when the F value was below an $\alpha < 0.05$ significance threshold. Daily transpiration and water
stress events were assigned to vegetative and reproductive stages. Daily soil evaporation was
aggregated into fallow and growing periods. Soil evaporation was identical from January 1 to
April 15 across all three planting dates, and not modeled from the date of the latest harvest
(typically October) to December 31, so these periods were ignored.

RESULTS

Model performance for days to emergence

The CERES-Sorghum model performance for days to emergence using a Tbase of 6 °C, 8 °C,
and 10 °C was evaluated by comparing observed and simulated days to emergence for
experiments conducted on different planting dates in Kansas . Results indicated a RMSE of 2.2,
1.7, and 2.0 days for a Tbase of 6 °C, 8 °C and 10 °C, respectively (Figure 2a). For six
experiments planted in April, observed days to seedling emergence ranged from 13 to 19 days,
while simulated days to emergence, with a Tbase of 8 °C, ranged from 11 to 18 days (Figure 2a).
For two experiments planted in May, observed days to emergence were 5 and 7 and the
corresponding simulated values, with a Tbase of 8 °C, were 6 and 8 days, respectively (Figure
2a).

Seasonal simulations for days to emergence for a Tbase ranging from 6 °C to 10 °C in the
study site (Figure 2b) indicated that, for instance, in northwest location (Colby) an increase in 1
°C for Tbase delays the crop emergence around 1.5, 0.6, and 0.2 days for planting dates in mid-
April, mid-May and mid-June, respectively. Across locations, an increase in 1 °C delays the crop
emergence in around 1.3, 0.5, and 0.2 for planting dates in very early, early, and normal planting
dates, respectively. Overall, these results suggest that a lower Tbase reduces the time required for
emergence, particularly when the crop is simulated to be planted in early spring (April 15).

Therefore, as outlined above, simulations in early spring using a Tbase of 8 °C do represent days
to emergence for kaoliang, a chilling tolerant genotype, when planted in early spring.

Soil moisture at planting

At each location and year, simulations started on January 1 to accrue soil moisture in the soil
profile as a function of daily precipitation (Figure 3). In this growing region, high variability of
monthly precipitation is observed from May to August (Figure 3a). The direct impact of
precipitation in soil moisture is illustrated in Figure 3b and 3c, where the soil moisture or
extractable soil water on each planting date varied across locations, with less soil moisture in
western locations than in eastern locations and less soil moisture in very early and early planting
(April 15, May 15) dates than in normal planting dates (June 15). In general, less moisture
accrued prior to planting for very early planting dates (158 ∓35 mm) compared to early (168
∓40 mm) and normal planting dates (185 ∓46 mm).

Growing season length

In each year, the phenology development and corresponding environmental conditions were
classified into days from planting to seedling emergence, vegetative growth (days from seedling
emergence to anthesis), and reproductive growth (days from anthesis to physiological maturity)
(Figure 4, Table S2). Simulations indicated that shifting planting dates from early summer to
early spring slowed down the rate of seedling emergence. For instance, in very early, early, and
normal planting dates seedlings are expected to emerge 16, 10, and 6 days after planting,
respectively. The number of days to seedling emergence and vegetative growth were greater in
western regions than in eastern regions (Figure 4a and 4b). For both full- and short-season
hybrids, very early planting dates extended the vegetative stage and shortened the reproductive
stage. By contrast, normal planting dates shortened the vegetative stage and extended the
reproductive stage (Figure 4, Table S2). For instance, a full-season hybrid that was simulated to
be planted in early spring (April 15) and early summer (June 15) completed the vegetative
growth in 79 and 63 days and reproductive growth in 44 and 48 days, respectively (Table S2).

Temperatures experienced by growth stage

Very early, early and normal planting scenarios led to contrasting temperature profiles during the
growing period (Figure 4d-f), as summarized by the maximum (Tmax) and minimum (Tmin)
temperatures experienced by the crop at the given stage (Table S2). When a full-season hybrid
was simulated to be planted very early (mid-April), it experienced a temperature trend from low
(vegetative stage) to high (reproductive stage). By contrast, when planted in mid-June, it
experienced a temperature trend from high (vegetative stage) to low (reproductive stage).

In general, across all three planting dates most of the precipitation (from 63–65%)
occurred during the vegetative growth (Table S2). On average, in very early planting dates the
crop experienced slightly higher seasonal precipitation compared to normal planting (347 mm vs.
263 mm for full-season hybrids; 310 mm vs. 240 mm for short-season hybrids). In western sites,
temperatures were lower than in the eastern sites. For instance, given normal planting, during
reproductive stages the minimum temperature in the northwestern site (Colby) was 11 °C and in
the eastern site (Manhattan) was 14 °C.

Soil evaporation in fallow and growing periods

Daily soil evaporation was aggregated into fallow and growing periods and compared across planting date and hybrid type. Simulations of a full-season hybrid indicate that on average across all planting dates, 30% of soil evaporation occurred in fallow periods and 70% during growing periods. Most of the soil evaporation during the growing period occurred before anthesis (53% of total soil evaporation). For all precipitation levels (dry, moderate, and wet years), very early planting dates significantly reduced the total soil evaporation compared with normal planting dates (Figure 5a, upper panel). Likewise, for all precipitation levels, very early planting dates reduced the soil evaporation during fallow periods (Figure 5a, lower panel, upper bars light colors) but increased the soil evaporation during the growing periods (Figure 5a, lower panel, lower bars and dark colors).

Plant transpiration

Daily transpiration for simulated full- and short-season hybrids was aggregated into vegetative and reproductive growth for each planting date in each year (Table S3). As would be expected, transpiration was substantially higher in the wet years (289 mm) compared to moderate (212 mm) or dry years (128 mm) across all planting dates. Simulations for a full-season hybrid indicated higher average total transpiration or water capture for very early planting dates (236 mm) than early (232 mm) or normal planting dates (211 mm). In dry years, total transpiration was similar in all planting dates (Figure 5b, upper panel). In both, moderate and wet years, very early and early planting dates significantly increased plant transpiration compared with late planting dates (Figure 5b, upper panel).

During vegetative growth, no significant differences in transpiration were found among planting dates in all precipitation levels (Figure 5b, lower panel, upper bars and light color). During reproductive growth, discrepancies in transpiration among planting dates were negligible only in dry years. However, these discrepancies in transpiration among planting dates were consistent, in both, moderate and wet years (Figure 5b, lower panel, lower bars and dark colors). Overall for a full-season hybrid, in dry years, transpiration during reproductive growth in very early and normal planting dates averaged 27% and 21%, respectively. By contrast, in wet years, transpiration during reproductive growth was greater, averaging 44% and 42% of transpiration for very early and normal planting, respectively. This suggests that very early and early planting can lead to drought escape under favorable scenarios.

Water stress events

CERES-Sorghum has two water stress factors whose values range from 0 (no stress) to 1 (stress). In this study a water stress event is defined when the minimum value between water deficit factor for photosynthesis (SWFAC) or water deficit factor for development (TURFAC) was higher than 0.6. Next, the number of water stress events were aggregated into vegetative and reproductive growth and are presented in Table S4. Simulations for a full-season hybrid indicate that on average the number of days under water stress during the growing period in very early, early, and normal planting dates was 31, 32 and 28, respectively. Similarly, on average across all

planting dates the number of water stress events during the growing period was higher in dry

years (39), followed by moderate (30) and wet years (12).

The number of days under water stress in different planting dates and precipitation levels

during the growing period for a full-season hybrid is provided in Figure 5c. During the growing

period (Figure 5c, upper panel) and during vegetative stage (Figure 5c, upper bars and light

color) the frequency of water stress events among planting dates was negligible in all

precipitation levels. During the reproductive stage the frequency of water stress events was

significantly higher in very early and early planting dates in all precipitation levels (Figure 5c,

lower bars and dark color).

Grain yield

CERES-Sorghum simulates the plant-soil-water dynamics on a daily time step that finally is

translated into grain yield. To evaluate the effect of planting date on grain yield, we have

conducted simulations for full- and short-season hybrids (Figure 6 and Table S5). The simulated

grain yield over 1986–2015 averaged 4 Mg ha^{-1} across locations, planting dates, and hybrids, and

ranged from 0 to 12 Mg ha^{-1}. Zero yield was obtained in 1989 in the northwest site (Colby) for

all planting dates due to exceptionally low annual precipitation (161 mm) and in 2004 in the

southwest site (Garden City) due to low precipitation in very early spring. In general, grain yield

varied significantly among planting dates ($p < 0.001$), precipitation levels ($p < 0.001$) and

hybrids ($p < 0.001$).

Low grain yield was obtained in dry years, as expected, and differences in grain yield

among planting dates were negligible. For instance, in dry years, the simulated yield for full-

season hybrid was 1.6 Mg ha^{-1}, not significantly different ($p > 0.05$) from yields of either very

early, early or normal planting dates. In wet years, by contrast, yields were higher for very early

and early planting versus normal planting. For instance, in wet years very early planting dates

averaged 7.4 Mg ha^{-1} for a full-season hybrid, 13% greater than the mean for normal planting

dates (6.5 Mg ha^{-1}) .

Overall, interactions between planting dates and hybrids were not significant (Table S5),

indicating similar trends for both maturing genotypes with very early and early planting dates

with yield increase in environments from moderate to high precipitation. Nevertheless, grain

yield across locations for full- and short-season hybrids in contrasting sites (Table S6) indicated

that short-season hybrids can contribute a slight yield increase for normal planting dates. For

instance, in the northwestern site (Colby) under normal planting, the grain yield of a short-season

hybrid was 2.9 Mg ha^{-1} and for a full-season hybrid was 2.6 Mg ha^{-1}. Overall, results indicate

that very early and early planting of chilling-tolerant sorghum hybrids can potentially increase

grain yield in 66% of the years.

DISCUSSION

Validity of the modeling approach for the target cropping system

After more than 50 years of breeding for the early-planted chilling tolerance trait in grain

sorghum, this study provides insights on the impact of this G × E × M intervention at crop scale,

narrowing the bridge between genotype to phenotype. To this end, this study used an agricultural

systems tool, the CERES-Sorghum model, because it is one of the most widely used sorghum
models (Adam et al., 2018; Amouzou et al., 2019; Folliard et al., 2004; Fu et al., 2016; Kothari et
al., 2020; Pachta, 2007; Singh et al., 2014), and its accuracy has been established for simulations
of grain yield in Kansas for normal planting dates (Araya et al., 2017; Staggenborg and
Vanderlip, 2005; White et al., 2015). Further, the model accuracy in predicting days of
emergence for early and normal planting dates (Figure 2a) and the sensitivity analysis for Tbase
(Figure 2b) suggests that the CERES-Sorghum model, which uses a Tbase of 8 °C, is likely
appropriate to forecast the effects of variables analyzed in this study.

Still, the congruence between observed and simulated values for soil moisture,
evapotranspiration, soil evaporation, and transpiration for the CERES-Sorghum model has not
yet been demonstrated. Crop model applications rely on rigorous model testing or model
comparison between observed versus simulated variables (Wallach et al., 2014). Other models
such as APSIM-Maize and APSIM-Soybean simulated field observations of soil moisture for
maize and soybean (Ebrahimi-Mollabashi et al., 2019; Yang et al., 2018). Likewise, observed
seasonal evapotranspiration for sweet sorghum and maize matched those values simulated with
crop models CERES-Sorghum and CERES-Maize (Araya et al., 2017; DeJonge et al., 2012;
Lopez et al., 2017). Commonly, in sorghum and other cereal crops, crop model evaluation is
conducted for grain yield and biomass components (Kassie et al., 2016; Yakoub et al., 2017).
Consequently, field experimentation to evaluate water budgets are needed for grain sorghum.

Some aspects of farm conditions not considered by the model may affect the
interpretation of our findings. For instance, simulated fallow periods before planting disregard
weed growth pressure that can deplete soil moisture and nutrients. Studies in Kansas indicate that
in dry years, green fallow or a crop can deplete soil moisture decreasing the productivity of the
subsequent winter wheat season (Holman et al., 2016; Schlegel et al., 2017; Schlegel and Havlin,
1997). Note, the fallow period before planting, which allows weed growth, was longer for
normal planting dates versus the earlier planting dates. For this reason, it is possible that the
model overestimates expected soil moisture available at this planting date.

Potential benefits of early-planted chilling tolerant sorghum
In the US maize and sorghum belts, optimum planting windows for maize and soybean vary
between mid-April to late-May (Baum et al., 2019; Long et al., 2017; Mourtzinis et al., 2017;
Staggenborg et al., 1999a); therefore, these optimum planting windows would be suitable for
sorghum. Indeed, simulations show that bringing forward the sorghum planting date in mid-April
(very early) and mid-May (early) in years with moderate to high precipitation (top two-thirds of
years based on annual precipitation) can lead to a yield increase on average by 12% for a full-
season hybrid and and 7% for a short-season hybrid compared to normal planting dates (Table
S5). Higher sorghum yields were reported in Kansas in planting dates in early June (Ciampitti et
al., 2019). By contrast, when shifting planting dates from June to May, a yield increase (4–35%)
was reported in four experiments while a yield decline (-69%) was reported in one experiment
(Maiga, 2012). Those experimental findings are in line with our simulations and indicate that
early chilling tolerance can increase yields under favorable scenarios (Figure 6). Given that

yields in very early and early planting dates were not substantially different (Figure 6, Table S5),
it is likely that the optimum planting window for chilling tolerant sorghum assuming an adequate
early stand establishment ranges from mid-April to mid-May, similar to those of maize and
soybean.
396   In cereals, drought escape can be achieved through short-season varieties that flower
earlier (Blum, 2011; Shavrukov et al., 2017), and in sorghum drought escape is a proposed
benefit of early planting and chilling tolerance (Burow et al., 2011; Franks et al., 2006).
However, our simulations suggest that the probability of water stress during grain filling would
actually be expected to increase slightly for planting dates in early spring (Figure 5c). The results
to some extent support that it is possible to avoid the terminal drought by increasing plant
transpiration or water capture in years with moderate to high precipitation (Figure 5b). In our
simulations, the reduction of soil evaporation during the fallow portion of the season can not
directly be capitalized into grain yield, but this water budget can hypothetically benefit a double
crop rotation (Burow et al., 2011) or be available in the soil profile for the subsequent cropping
season (Figure S2). In dryland systems of western Kansas, wheat-sorghum is a common crop
rotation due to benefits in grain yield and soil water moisture (Schlegel and Havlin, 1997). Using
an early-planted chilling-tolerant grain sorghum in this rotation can potentially further improve
grain yield and water productivity for both crops. Additional simulations of crop rotations would
be needed to assess these benefits.
Expected tradeoffs for early and normal planting dates
Environmental conditions dictate the rate of phenology development. Planting dates in early
spring extended the duration of the growing season (Figure 4, Table S2), especially days for
seedling emergence and days to anthesis, as previously predicted (Burow et al., 2011) and
observed (Kapanigowda et al., 2013). The delay in seedling emergence in very early spring
(Figure 4) was additionally associated with low levels of soil moisture (Figure 3). Although,
studies on chilling tolerance in field conditions (Maulana and Tesso, 2013; Ostmeyer et al.,
2020) do not report the effect of soil moisture on germination and seedling emergence. Soil
moisture deficit and low temperature both impair germination and delay emergence in sorghum
(Evans et al., 1961), but their interactions are not well understood. Consequently, our study
suggests that hybrids with chilling tolerance should be planted under optimum levels of initial
soil moisture to ensure early crop establishment, as recommended for other crops such as maize
(Lu et al., 2017).
424   Sorghum grain yield is affected when the crop experiences extreme temperatures, above
38 °C and below 11 °C, during grain filling (Singh et al., 2015; Staggenborg et al., 1999b). In
our simulations, very early planting exposed the crop to moderately high temperatures (<38 °C)
during grain filling. By contrast, normal planting exposed the crop to temperatures below 10 °C
during grain filling in 50% of the years in the northwest site (Colby) (Figure 4f and Table S6).
Risk of freezing temperatures for normal planting dates were reported in studies conducted in
Kansas, Colorado and Texas (Baumhardt and Howell, 2006; McMaster et al., 2016; Staggenborg
et al., 1999b; Staggenborg and Vanderlip, 1996). Furthermore, in Colorado short-season hybrids

can avoid freezing temperatures during grain filling for planting dates in late spring (McMaster
et al., 2016).
Climate change, which was not accounted for in our simulations, may affect the potential
benefits and tradeoffs for very early and early planting of chilling-tolerant sorghum. For
instance, climate time series analysis in Kansas indicates that the frost risk events have become
less frequent in the last decade (Lin et al., 2017) and global temperatures are expected to rise
another 0.5–2.5 °C by mid-century (IPCC, 2013). In future scenarios it is possible that the risk of
heat stress during flowering can increase for very early planting, while the frequency of frost
damage may not be important for planting dates in early summer. Alternatively, very early
planting of chilling-tolerant hybrids could provide heat stress escape. Simulations under future
climate scenarios would be needed to resolve these potential tradeoffs.
A potential risk to very early, early and normal plantings was the low grain yield in years
with low precipitation (Figure 6), due to the low soil moisture (Figure 3) and high frequency of
daily droughts during pre-flowering and post-flowering stages (Figure 5c). This effect was
reported in other crops. For instance, in Iowa, locations exposed to earlier water stress exhibited
lower soybean yield (Irmak et al., 2002), while across the US midwest, locations with low soil
moisture imposed a greater penalty on maize yield (Seifert et al., 2017). In Texas, a simulation
study indicated that a short-season hybrid is better adapted to dryland systems when the crop is
planted in June (Baumhardt and Howell, 2006). Similar results were obtained for the
northwestern site (Colby), where a short-season hybrid yielded higher than a full-season hybrid
in planting dates in late spring (Table S6). Overall, low early-season water budgets as a result of
low precipitation can override the value of very early planting.

Next steps for models of chilling tolerance to guide crop improvement
In breeding and genetics studies, chilling tolerance is most often characterized through seedling
emergence rate and seedling vigor scores, usually based on visual evaluation (Burow et al., 2011;
Franks et al., 2006; Marla et al., 2019; Parra-Londono et al., 2018). By contrast, crop models
simulate days to emergence, seedling biomass weight, and leaf area. CERES-Sorghum,
developed for normal conditions, does not model any detrimental effect of chilling temperatures
with respect to seedling emergence or leaf damage. Consequently, the model can not simulate the
grain yield for a genotype which was damaged by chilling temperatures.
Future work to extend a chilling sensitive routine could entail penalizing the development
rate, growth rate, or leaf area. The development rate can be reduced by shifting the base
temperature (e.g. from 8 °C to 10 °C) and varying the duration from planting to emergence. For
instance, currently thermal time from planting to germination (50 degree days) and the coleoptile
extension rate (0.1 °C day cm^{-1}) are fixed parameters. Chilling tolerant genotypes can attain 50%
seedling emergence two days earlier than their susceptible counterparts (Stickler et al., 1962),
suggesting that there is variability for base temperature, thermal time from planting to
germination, and/or coleoptile extension rate. Otherwise, differences for base temperature (0–9.8
°C) were recently reported for Ethiopian sorghum germplasm, which is believed to harbor
adaptations to chilling-prone high-altitude regions (Tirfessa et al., 2020).

In this study, simulations were aggregated into three precipitation levels using a simple
quantile classification. Although the frequency of water stress corresponded well to each
precipitation level (Figure 5c) the daily water stress can be used to characterize drought patterns
more precisely. For instance, the APSIM-Wheat and the SSM model were used to classify and
quantify drought patterns across the Australian wheat belt (Chenu et al., 2013) and the US maize
belt (Messina et al., 2015), respectively. Thus, the characterization of drought patterns (i.e.
envirotyping) with CERES-Sorghum, in these locations and across the US sorghum belt would
guide crop improvement programs to focus on traits better-suited for target environments.

Overall, through years of field experimentation, research on early season chilling
tolerance pointed out the potential outcomes of a sorghum hybrid with this trait. This study has
shown that a crop simulation model can translate the genetic value of this breeding trait in terms
of grain yield by integrating environmental conditions (i.e. the impact of annual variability) and
realistic agronomic management. Furthermore, this study tested hypotheses regarding benefits of
early chilling tolerance proving that it is difficult to informally predict the state of any variable
(i.e. evapotranspiration) at crop system scale, without formal modeling. Thus, our study
underscores the value of crop modeling to complement and guide plant breeding and genetics.

ACKNOWLEDGEMENTS
This study was supported by funding from the Foundation for Food and Agriculture Research -
Seeding Solution "CA18-SS-0000000094 – Bridging the Genome-to-Phenome Breeding Gap for
Water-Efficient Crop Yields (G2P Bridge)" to G.P.M. and I.C; the Kansas Department of
Agriculture "Collaborative Sorghum Investment Program Water Optimized Sorghum for
Kansas" to G.P.M; and the Kansas Grain Sorghum Commission.

AUTHOR CONTRIBUTIONS
Designed the research: GPM, RR, IC. Performed research: RR. Analyzed data: RR, SSB. Wrote
the paper: RR, GPM.
497

REFERENCES

Adam, M., Dzotsi, K.A., Hoogenboom, G., Traoré, P.C.S., Porter, C.H., Rattunde, H.F.W., Nebie, B., Leiser, W.L., Weltzien, E., Jones, J.W., 2018. Modelling varietal differences in response to phosphorus in West African sorghum. European Journal of Agronomy, Recent advances in crop modelling to support sustainable agricultural production and food security under global change 100, 35–43. https://doi.org/10.1016/j.eja.2018.04.001

Adegbeye, M.J., Ravi Kanth Reddy, P., Obaisi, A.I., Elghandour, M.M.M.Y., Oyebamiji, K.J., Salem, A.Z.M., Morakinyo-Fasipe, O.T., Cipriano-Salazar, M., Camacho-Díaz, L.M., 2020. Sustainable agriculture options for production, greenhouse gasses and pollution alleviation, and nutrient recycling in emerging and transitional nations - An overview. Journal of Cleaner Production 242, 118319. https://doi.org/10.1016/j.jclepro.2019.118319

Aggarwal, P.K., Kropff, M.J., Cassman, K.G., ten Berge, H.F.M., 1997. Simulating genotypic strategies for increasing rice yield potential in irrigated, tropical environments. Field Crops Research, Special Issue: Applications of Rice Modelling 51, 5–17. https://doi.org/10.1016/S0378-4290(96)01044-1

Amouzou, K.A., Lamers, J.P.A., Naab, J.B., Borgemeister, C., Vlek, P.L.G., Becker, M., 2019. Climate change impact on water- and nitrogen-use efficiencies and yields of maize and sorghum in the northern Benin dry savanna, West Africa. Field Crop. Res. 235, 104–117. https://doi.org/10.1016/j.fcr.2019.02.021

Araya, A., Kisekka, I., Gowda, P.H., Prasad, P.V.V., 2018. Grain sorghum production functions under different irrigation capacities. Agricultural Water Management 203, 261–271. https://doi.org/10.1016/j.agwat.2018.03.010

Araya, A., Kisekka, I., Gowda, P.H., Prasad, P.V.V., 2017. Evaluation of water-limited cropping systems in a semi-arid climate using DSSAT-CSM. Agricultural Systems 150, 86–98. https://doi.org/10.1016/j.agsy.2016.10.007

Baeumler, N.W., Gupta, S.C., 2020. Precipitation as the Primary Driver of Variability in River Nitrogen Loads in the Midwest United States. JAWRA Journal of the American Water Resources Association 56, 113–133. https://doi.org/10.1111/1752-1688.12809

Bates, D., Maechler, M., Bolker, B.M., Walker, S.C., 2015. Fitting Linear Mixed-Effects Models Using lme4. J. Stat. Softw. 67, 1–48. https://doi.org/10.18637/jss.v067.i01

Baum, M.E., Archontoulis, S.V., Licht, M.A., 2019. Planting Date, Hybrid Maturity, and Weather Effects on Maize Yield and Crop Stage. Agronomy Journal 111, 303–313. https://doi.org/10.2134/agronj2018.04.0297

Baumhardt, R.L., Howell, T.A., 2006. Seeding Practices, Cultivar Maturity, and Irrigation Effects on Simulated Grain Sorghum Yield. Agronomy Journal; Madison 98, 462–470.

Bhattarai, B., Singh, S., West, C.P., Saini, R., 2019. Forage Potential of Pearl Millet and Forage Sorghum Alternatives to Corn under the Water-Limiting Conditions of the Texas High Plains: A Review. Crop, Forage & Turfgrass Management 5. https://doi.org/10.2134/cftm2019.08.0058

Blum, A., 2011. Drought Resistance and Its Improvement, Plant Breeding for Water-Limited Environments. Springer-Verlag Berlin, Berlin. https://doi.org/10.1007/978-1-4419-7491-4_3

Burke, J.J., Emendack, Y., Hayes, C., Xin, Z., Burow, G., 2019. Registration of Five Grain Sorghum Lines with Early-Season Seedling Cold Tolerance. Journal of Plant Registrations 13, 422–426. https://doi.org/10.3198/jpr2019.02.0010crg

Burow, G., Burke, J.J., Xin, Z., Franks, C.D., 2011. Genetic dissection of early-season cold
tolerance in sorghum (Sorghum bicolor (L.) Moench). Mol Breeding 28, 391–402.
https://doi.org/10.1007/s11032-010-9491-4
Challinor, A.J., Müller, C., Asseng, S., Deva, C., Nicklin, K.J., Wallach, D., Vanuytrecht, E.,
Whitfield, S., Ramirez-Villegas, J., Koehler, A.-K., 2018. Improving the use of crop
models for risk assessment and climate change adaptation. Agricultural Systems 159,
296–306. https://doi.org/10.1016/j.agsy.2017.07.010
Chenu, K., Deihimfard, R., Chapman, S.C., 2013. Large-scale characterization of drought
pattern: a continent-wide modelling approach applied to the Australian wheatbelt –
spatial and temporal trends. New Phytologist 198, 801–820.
https://doi.org/10.1111/nph.12192
Chenu, K., Porter, J.R., Martre, P., Basso, B., Chapman, S.C., Ewert, F., Bindi, M., Asseng, S.,
2017. Contribution of Crop Models to Adaptation in Wheat. Trends in Plant Science 22,
472–490. https://doi.org/10.1016/j.tplants.2017.02.003
Chopra, R., Burow, G., Burke, J.J., Gladman, N., Xin, Z., 2017. Genome-wide association
analysis of seedling traits in diverse Sorghum germplasm under thermal stress. BMC
Plant Biology 17. http://dx.doi.org.ezproxy2.library.colostate.edu/10.1186/s12870-016-
0966-2
Ciampitti, I.A., Prasad, P.V.V., Schlegel, A.J., Haag, L., Schnell, R.W., Arnall, B., Lofton, J.,
2019. Genotype × Environment × Management Interactions: US Sorghum Cropping
Systems, in: Sorghum. John Wiley & Sons, Ltd, pp. 277–296.
https://doi.org/10.2134/agronmonogr58.c13
Cooper, M., Chapman, S.C., Podlich, D.W., Hammer, G.L., 2002. The GP Problem: Quantifying
Gene-to-Phenotype Relationships. In Silico Biology 2, 151–164.
DeJonge, K.C., Ascough, J.C., Andales, A.A., Hansen, N.C., Garcia, L.A., Arabi, M., 2012.
Improving evapotranspiration simulations in the CERES-Maize model under limited
irrigation. Agricultural Water Management 115, 92–103.
https://doi.org/10.1016/j.agwat.2012.08.013
Doggett, H.J., Majisu, B.N., 1968. Disruptive selection in crop development. Heredity 23, 1–23.
https://doi.org/10.1038/hdy.1968.1
Ebrahimi-Mollabashi, E., Huth, N.I., Holzworth, D.P., Ordóñez, R.A., Hatfield, J.L., Huber, I.,
Castellano, M.J., Archontoulis, S.V., 2019. Enhancing APSIM to simulate excessive
moisture effects on root growth. Field Crops Research 236, 58–67.
https://doi.org/10.1016/j.fcr.2019.03.014
Ercoli, L., Mariotti, M., Masoni, A., Arduini, I., 2004. Growth responses of sorghum plants to
chilling temperature and duration of exposure. European Journal of Agronomy 21, 93–
103. https://doi.org/10.1016/S1161-0301(03)00093-5
Evans, W.F., Stickler, F.C., Laude, H.H., 1961. Sorghum Seed Germination as Affected by
Moisture and Temperature. Transactions of the Kansas Academy of Science (1903-) 64,
210–217. https://doi.org/10.2307/3626708
Folliard, A., Traore, P.C.S., Vaksmann, M., Kouressy, M., 2004. Modeling of sorghum response
to photoperiod: a threshold-hyperbolic approach. Field Crop. Res. 89, 59–70.
https://doi.org/10.1016/j.fcr.2004.01.006
Franks, C.D., Burow, G.B., Burke, J.J., 2006. A Comparison of U.S. and Chinese Sorghum
Germplasm for Early Season Cold Tolerance. Crop Science 46, 1371–1376.
https://doi.org/10.2135/cropsci2005.08-0279

Fu, T., Ko, J., Wall, G.W., Pinter, P.J., Kimball, B.A., Ottman, M.J., Kim, H.-Y., 2016.
Simulation of climate change impacts on grain sorghum production grown under free air
CO2 enrichment. Int. Agrophys. 30, 311–322. https://doi.org/10.1515/intag-2016-0007
Holman, J., Roberts, T., Maxwell, S., 2016. Fallow Replacement Crop (Cover Crops, Annual
Forages, and Short-Season Grain Crops) Effects on Wheat and Grain Sorghum Yields.
Kansas Agricultural Experiment Station Research Reports 2.
https://doi.org/10.4148/2378-5977.1247
Hoogenboom, G., Porter, C.H., Boote, K.J., Shelia, V., Wilkens, P.W., Singh, U., White, J.W.,
Asseng, S., Lizaso, J.I., Moreno, L.P., Pavan, W., Ogoshi, R., Hunt, L.A., Tsuji, G.Y.,
Jones, J.W., 2019. The DSSAT crop modeling ecosystem, in: Advances in Crop
Modeling for Sustainable Agriculture. Burleigh Dodds Science Publishing, Cambridge,
United Kingdom, pp. 173–216.
IPCC, 2013. Summary for Policymakers. Climate Change 2013: The Physical Science Basis.
Contribution of Working Group I to the Fifth Assessment Report of the
Intergovernmental Panel on Climate Change.
https://doi.org/10.1017/CBO9781107415324.004
Irmak, A., Batchelor, W.D., Jones, J.W., Irmak, S., Paz, J.O., Beck, H.W., Egeh, M., 2002.
Relationship between plant available soil water and yield for explaining soybean yield
variability. Appl. Eng. Agric. 18, 471–482.
Jones, J.W., Hoogenboom, G., Porter, C.H., Boote, K.J., Batchelor, W.D., Hunt, L.A., Wilkens,
P.W., Singh, U., Gijsman, A.J., Ritchie, J.T., 2003. The DSSAT cropping system model.
European Journal of Agronomy, Modelling Cropping Systems: Science, Software and
Applications 18, 235–265. https://doi.org/10.1016/S1161-0301(02)00107-7
Kansas Mesonet, 2020. Kansas Mesonet [WWW Document]. URL https://mesonet.k-state.edu/
(accessed 1.6.21).
Kapanigowda, M.H., Perumal, R., Aiken, R.M., Herald, T.J., Bean, S.R., Little, C.R., 2013.
Analyses of sorghum [Sorghum bicolor (L.) Moench] lines and hybrids in response to
early-season planting and cool conditions. Can. J. Plant Sci. 93, 773–784.
https://doi.org/10.4141/cjps2012-311
Kassie, B.T., Asseng, S., Porter, C.H., Royce, F.S., 2016. Performance of DSSAT-Nwheat across
a wide range of current and future growing conditions. European Journal of Agronomy
81, 27–36. https://doi.org/10.1016/j.eja.2016.08.012
Kholová, J., Murugesan, T., Kaliamoorthy, S., Malayee, S., Baddam, R., Hammer, G.L.,
McLean, G., Deshpande, S., Hash, C.T., Craufurd, P.Q., Vadez, V., 2014. Modelling the
effect of plant water use traits on yield and stay-green expression in sorghum. Functional
Plant Biol. 41, 1019–1034. https://doi.org/10.1071/FP13355
Kothari, K., Ale, S., Bordovsky, J.P., Porter, D.O., Munster, C.L., Hoogenboom, G., 2020.
Potential benefits of genotype-based adaptation strategies for grain sorghum production
in the Texas High Plains under climate change. European Journal of Agronomy 117,
126037. https://doi.org/10.1016/j.eja.2020.126037
Laingen, C., 2015. A Spatiotemporal Analysis of Sorghum in the United States. Papers in
Applied Geography 1, 307–311. https://doi.org/10.1080/23754931.2015.1084359
Lenaerts, B., Collard, B.C.Y., Demont, M., 2019. Review: Improving global food security
through accelerated plant breeding. Plant Science 287, 110207.
https://doi.org/10.1016/j.plantsci.2019.110207
Lin, X., Harrington, J., Ciampitti, I., Gowda, P., Brown, D., Kisekka, I., 2017. Kansas Trends

and Changes in Temperature, Precipitation, Drought, and Frost-Free Days from the 1890s
to 2015. Journal of Contemporary Water Research & Education 162, 18–30.
https://doi.org/10.1111/j.1936-704X.2017.03257.x
Long, N.V., Assefa, Y., Schwalbert, R., Ciampitti, I.A., 2017. Maize Yield and Planting Date
Relationship: A Synthesis-Analysis for US High-Yielding Contest-Winner and Field
Research Data. Front. Plant Sci. 8. https://doi.org/10.3389/fpls.2017.02106
Lopez, J.R., Erickson, J.E., Asseng, S., Bobeda, E.L., 2017. Modification of the CERES grain
sorghum model to simulate optimum sweet sorghum rooting depth for rainfed production
on coarse textured soils in a sub-tropical environment. Agricultural Water Management
181, 47–55. https://doi.org/10.1016/j.agwat.2016.11.023
Lu, H., Xue, J., Guo, D., 2017. Efficacy of planting date adjustment as a cultivation strategy to
cope with drought stress and increase rainfed maize yield and water-use efficiency.
Agricultural Water Management, Special Issue on Improving Agricultural Water
Productivity to Ensure Food Security under Changing Environments Overseen by: Brent
Clothier 179, 227–235. https://doi.org/10.1016/j.agwat.2016.09.001
Lyons, J.M., 1973. Chilling Injury in Plants. Annual Review of Plant Physiology 24, 445–466.
https://doi.org/10.1146/annurev.pp.24.060173.002305
Maiga, A., 2012. Effects of planting practices and nitrogen management on grain sorghum
production [WWW Document]. URL /paper/Effects-of-planting-practices-and-nitrogen-
on-grain-Maiga/2b32e6a422fe1df3c6cd402388eec91fddc44520 (accessed 5.28.20).
Marla, S.R., Burow, G., Chopra, R., Hayes, C., Olatoye, M.O., Felderhoff, T., Hu, Z.,
Raymundo, R., Perumal, R., Morris, G.P., 2019. Genetic Architecture of Chilling
Tolerance in Sorghum Dissected with a Nested Association Mapping Population. G3:
Genes, Genomes, Genetics 9, 4045–4057. https://doi.org/10.1534/g3.119.400353
Maulana, F., Tesso, T.T., 2013. Cold Temperature Episode at Seedling and Flowering Stages
Reduces Growth and Yield Components in Sorghum. Crop Science 53, 564–574.
https://doi.org/10.2135/cropsci2011.12.0649
McMaster, G.S., Edmunds, D.A., Jones, S.M., Johnson, J.J., Vigil, M.F., 2016. Simulating the
Probability of Grain Sorghum Maturity before the First Frost in Northeastern Colorado.
Agronomy 6, 44. https://doi.org/10.3390/agronomy6040044
Messina, C.D., Sinclair, T.R., Hammer, G.L., Curan, D., Thompson, J., Oler, Z., Gho, C.,
Cooper, M., 2015. Limited-Transpiration Trait May Increase Maize Drought Tolerance in
the US Corn Belt. Agronomy Journal 107, 1978–1986.
https://doi.org/10.2134/agronj15.0016
Mocoeur, A., Zhang, Y.-M., Liu, Z.-Q., Shen, X., Zhang, L.-M., Rasmussen, S.K., Jing, H.-C.,
2015. Stability and genetic control of morphological, biomass and biofuel traits under
temperate maritime and continental conditions in sweet sorghum (Sorghum bicolour).
Theor Appl Genet 128, 1685–1701. https://doi.org/10.1007/s00122-015-2538-5
Mourtzinis, S., Gaspar, A.P., Naeve, S.L., Conley, S.P., 2017. Planting Date, Maturity, and
Temperature Effects on Soybean Seed Yield and Composition. Agronomy Journal 109,
2040–2049. https://doi.org/10.2134/agronj2017.05.0247
Ostmeyer, T., Bheemanahalli, R., Srikanthan, D., Bean, S., Peiris, K.H.S., Madasamy, P.,
Perumal, R., Jagadish, S.V.K., 2020. Quantifying the agronomic performance of new
grain sorghum hybrids for enhanced early-stage chilling tolerance. Field Crops Research
258, 107955. https://doi.org/10.1016/j.fcr.2020.107955
Pachta, C.J., 2007. Improving irrigated cropping systems on the high plains using crop

simulation models (Thesis). Kansas State University.

Parra-Londono, S., Fiedler, K., Kavka, M., Samans, B., Wieckhorst, S., Zacharias, A., Uptmoor,
R., 2018. Genetic dissection of early-season cold tolerance in sorghum: genome-wide
association studies for seedling emergence and survival under field and controlled
environment conditions. Theor Appl Genet 131, 581–595.
https://doi.org/10.1007/s00122-017-3021-2

Pražnikar, J., 2017. Particulate matter time-series and Köppen-Geiger climate classes in North
America and Europe. Atmospheric Environment 150, 136–145.
https://doi.org/10.1016/j.atmosenv.2016.11.056

Priestley, C.H.B., Taylor, R.J., 1972. On the Assessment of Surface Heat Flux and Evaporation
Using Large-Scale Parameters. Monthly Weather Review 100, 81–92.
https://doi.org/10.1175/1520-0493(1972)100<0081:OTAOSH>2.3.CO;2

R Core Team, 2017. , R: A language and environment for statistical computing. Foundation for
Statistical Computing. Viena, Austria.

Ritchie, J.T., Singh, U., Godwin, D.C., Bowen, W.T., 1998. Cereal growth, development and
yield, in: Tsuji, G.Y., Hoogenboom, G., Thornton, P.K. (Eds.), Understanding Options
for Agricultural Production, Systems Approaches for Sustainable Agricultural
Development. Springer Netherlands, Dordrecht, pp. 79–98. https://doi.org/10.1007/978-
94-017-3624-4_5

Roozeboom, K., Fjell, D., 1998. Selection of grain sorghum hybrids, in: Grain Sorghum
Production Handbook. Kansas State University.

Schaffasz, A., Windpassinger, S., Friedt, W., Snowdon, R., Wittkop, B., 2019. Sorghum as a
Novel Crop for Central Europe: Using a Broad Diversity Set to Dissect Temperate-
Adaptation. Agronomy 9, 535. https://doi.org/10.3390/agronomy9090535

Schlegel, A.J., Assefa, Y., Haag, L.A., Thompson, C.R., Holman, J.D., Stone, L.R., 2017. Yield
and Soil Water in Three Dryland Wheat and Grain Sorghum Rotations. Agronomy
Journal 109, 227–238. https://doi.org/10.2134/agronj2016.07.0387

Schlegel, A.J., Havlin, J.L., 1997. Green Fallow for the Central Great Plains. Agronomy Journal
89, 762–767. https://doi.org/10.2134/agronj1997.00021962008900050009x

Seifert, C.A., Roberts, M.J., Lobell, D.B., 2017. Continuous Corn and Soybean Yield Penalties
across Hundreds of Thousands of Fields. Agronomy Journal 109, 541–548.
https://doi.org/10.2134/agronj2016.03.0134

Shavrukov, Y., Kurishbayev, A., Jatayev, S., Shvidchenko, V., Zotova, L., Koekemoer, F., de
Groot, S., Soole, K., Langridge, P., 2017. Early Flowering as a Drought Escape
Mechanism in Plants: How Can It Aid Wheat Production? Front. Plant Sci. 8.
https://doi.org/10.3389/fpls.2017.01950

Shroyer, James., Kok, H., Fjell, D., 1998. Seedbed preparation and planting practices, in: Grain
Sorghum Production Handbook. Kansas State University, pp. 5–9.

Singh, P., Nedumaran, S., Traore, P.C.S., Boote, K.J., Rattunde, H.F.W., Prasad, P.V.V., Singh,
N.P., Srinivas, K., Bantilan, M.C.S., 2014. Quantifying potential benefits of drought and
heat tolerance in rainy season sorghum for adapting to climate change. Agricultural and
Forest Meteorology 185, 37–48. https://doi.org/10.1016/j.agrformet.2013.10.012

Singh, S.P., 1985. Sources of cold tolerance in grain Sorghum. Can. J. Plant Sci. 65, 251–257.
https://doi.org/10.4141/cjps85-037

Singh, V., Nguyen, C.T., van Oosterom, E.J., Chapman, S.C., Jordan, D.R., Hammer, G.L.,
2015. Sorghum genotypes differ in high temperature responses for seed set. Field Crops

728   Research 171, 32–40. https://doi.org/10.1016/j.fcr.2014.11.003

729 Smith, C.W., Frederiksen, R.A., 2000. Sorghum: Origin, History, Technology, and Production.
730   John Wiley & Sons.

731 Soil Survey Staff, 2020. Web Soil Survey - Home [WWW Document]. URL
732   https://websoilsurvey.sc.egov.usda.gov/App/HomePage.htm (accessed 6.7.19).

733 Staggenborg, S.A., Fjell, D.L., Devlin, D.L., Gordon, W.B., Maddux, L.D., Marsh, B.H., 1999a.
734   Selecting Optimum Planting Dates and Plant Populations for Dryland Corn in Kansas.
735   Journal of Production Agriculture 12, 85–90. https://doi.org/10.2134/jpa1999.0085

736 Staggenborg, S.A., Vanderlip, R.L., 2005. Crop Simulation Models Can be Used as Dryland
737   Cropping Systems Research Tools. Agronomy Journal 97, 378–384.
738   https://doi.org/10.2134/agronj2005.0378

739 Staggenborg, S.A., Vanderlip, R.L., 1996. Sorghum Grain Yield Reductions Caused by Duration
740   and Timing of Freezing Temperatures. Agronomy Journal 88, 473–477.
741   https://doi.org/10.2134/agronj1996.00021962008800030019x

742 Staggenborg, S.A., Vanderlip, R.L., Roggenkamp, G.J., Kofoid, K.D., 1999b. Methods of
743   simulating freezing damage during sorghum grain fill. Agronomy journal.

744 Stickler, F.C., Pauli, A.W., Casady, A.J., 1962. Comparative Responses of Kaoliang and Other
745   Grain Sorghum Types to Temperature. Crop Science 2, 136–139.
746   https://doi.org/10.2135/cropsci1962.0011183X000200020015x

747 Suleiman, A.A., Ritchie, J.T., 2003. Modeling Soil Water Redistribution during Second-Stage
748   Evaporation. Soil Science Society of America Journal 67, 377–386.
749   https://doi.org/10.2136/sssaj2003.3770

750 Taylor, A.O., Rowley, J.A., 1971. Plants under Climatic Stress: I. Low Temperature, High Light
751   Effects on Photosynthesis. Plant Physiol 47, 713–718.
752   https://doi.org/10.1104/pp.47.5.713

753 Tilman, D., Cassman, K.G., Matson, P.A., Naylor, R., Polasky, S., 2002. Agricultural
754   sustainability and intensive production practices. Nature 418, 671–677.
755   https://doi.org/10.1038/nature01014

756 Tirfessa, A., McLean, G., Mace, E., Oosterom, E. van, Jordan, D., Hammer, G., 2020.
757   Differences in temperature response of phenological development among diverse
758   Ethiopian sorghum genotypes are linked to racial grouping and agroecological
759   adaptation. Crop Science 60, 977–990. https://doi.org/10.1002/csc2.20128

760 van Ittersum, M.K., Leffelaar, P.A., van Keulen, H., Kropff, M.J., Bastiaans, L., Goudriaan, J.,
761   2003. On approaches and applications of the Wageningen crop models. European Journal
762   of Agronomy, Modelling Cropping Systems: Science, Software and Applications 18,
763   201–234. https://doi.org/10.1016/S1161-0301(02)00106-5

764 Villanueva, R.A.M., Chen, Z.J., 2019. ggplot2: Elegant Graphics for Data Analysis, 2nd edition.
765   Meas.-Interdiscip. Res. Perspect. 17, 160–167.
766   https://doi.org/10.1080/15366367.2019.1565254

767 Wallach, D., Makowski, D., Jones, J.W., Brun, F., 2014. Chapter 9 - Model Evaluation, in:
768   Wallach, D., Makowski, D., Jones, J.W., Brun, F. (Eds.), Working with Dynamic Crop
769   Models (Second Edition). Academic Press, San Diego, pp. 345–406.
770   https://doi.org/10.1016/B978-0-12-397008-4.00009-5

771 White, J.W., Alagarswamy, G., Ottman, M.J., Porter, C.H., Singh, U., Hoogenboom, G., 2015.
772   An Overview of CERES–Sorghum as Implemented in the Cropping System Model
773   Version 4.5. Agronomy Journal 107, 1987. https://doi.org/10.2134/agronj15.0102

Wylie, P., 2008. Managing sorghum for high yields: a blueprint for doubling sorghum
production. Grains Research and Development Corp., Kingston, A.C.T.
Yakoub, A., Lloveras, J., Biau, A., Lindquist, J.L., Lizaso, J.I., 2017. Testing and improving the
maize models in DSSAT: Development, growth, yield, and N uptake. Field Crops
Research 212, 95–106. https://doi.org/10.1016/j.fcr.2017.07.002
Yang, X., Zheng, L., Yang, Q., Wang, Z., Cui, S., Shen, Y., 2018. Modelling the effects of
conservation tillage on crop water productivity, soil water dynamics and
evapotranspiration of a maize-winter wheat-soybean rotation system on the Loess Plateau
of China using APSIM. Agricultural Systems 166, 111–123.
https://doi.org/10.1016/j.agsy.2018.08.005
Zhang, H., Link to external site, this link will open in a new window, Li, Y., Jian-Kang, Z.,
2018. Developing naturally stress-resistant crops for a sustainable agriculture. Nature
Plants; London 4, 989–996. http://dx.doi.org.er.lib.k-state.edu/10.1038/s41477-018-0309-
4

Table 1. Cultivar parameters for CERES-Sorghum model for full- and short-season
hybrids*.

Code	Full season	Short season	Description
P1	393	250	Potential duration from emergence to end of juvenile phase (°C day).
P2	102	102	Potential duration from juvenile phase to panicle initiation (°C day).
P2O	15.0	15.0	Critical daylength above which development slows (short day response) (h).
P2R	40	40	Photoperiod sensitivity as the extent to which development is delayed for each hour of photoperiod above P2O (°C day).
PANTH	617	617	Duration from panicle initiation to anthesis (°C day).
P3	152	152	Duration from end of flag leaf expansion to anthesis (°C day).
P4	82	82	Duration from anthesis to onset of grain filling (°C day).
P5	640	640	Duration of grain-filling phase (onset of grain filling to physiological maturity) (°C day).
PHINT	49	49	Phyllochron interval (°C day/leaf).
G1	3	3	Scaler for relative leaf size (unitless).
G2	6.5	6.5	Scalar for partitioning of assimilates to the panicle (unitless).

790 * Reproduced from Araya et al. (2018).

791   **Table 2. Characteristics for the study locations in Kansas.**

Location	Soil family	Soil texture	Organic matter (%)	Average annual precipitation (mm)	Koppen Climate classification
Colby	Richfield	Silty clay loam	2	520	Cold semi-arid
Garden City	Park-Ulysses	Silt loam	2	480	Cold semi-arid
Hays	Harney	Silt loam	2	680	Hot-summer humid continental
Manhattan	Smolan	Silt loam	3	900	Hot-summer humid continental

792
793

a

b

c

Figure 1. Study system to evaluate the impact of early planting with CERES-Sorghum. (a) Annual precipitation gradients across Kansas and locations to simulate the growth and development for sorghum in Kansas. (b) Cumulative growing degree units (GDU) or thermal time between mid-April to mid-June. The map was generated using the daily average mean over five years of daily data (https://prism.oregonstate.edu/). (c) The rate of sorghum and plantings in Kansas, barplots represent the five-year mean (2010-2016), black lines represent the cumulative planting (https://www.nass.usda.gov/) and the pink triangles represent the planting dates under study: very early (April 15), early (May 15), and normal (June 15).

Figure 2. CERES-Sorghum model performance for days to seedling emergence. (a) CERES-Sorghum model performance for days to seedling emergence in different planting dates in Kansas for Tbase of 6 °C, 8 °C, and 10 °C, observed information corresponds to experiments conducted in Hays (HA), Manhattan (MA), and Ashland Bottom (AB; 10 km from MA) in 2016−2018 that were planted in April, May and June. (b) Variation in simulated days to emergence as a function of Tbase in very early (April 15; black points), early (May 15, dark gray points), or normal (June 15, light gray points) planting. Each point represents the mean over 30 years and error lines indicate the standard error.

Figure 3. Effect of precipitation on soil moisture at planting for a simulated full-season hybrid. (a) Variability of monthly precipitation in four representative locations in Kansas, red triangles represent the monthly mean over a 30 years period. (b) Daily soil extractable water (soil moisture) in different planting dates, vertical dashed lines indicate the planting date at each location. (c) Extractable water for very early (mid-April), early (mid-May), and normal (mid-June) dates planting. Each boxplot and violin plot represent the annual variability over 30 years (1986-2015) in Kansas. Letters represent significant differences ($\alpha < 0.05$) using the Tukey HSD test.

Figure 4. Crop phenology for a simulated full-season sorghum hybrid in two contrasting locations. (a) Days for emergence, (b) days for vegetative development and (c) days for reproductive development for very early (mid-April), early (mid-May), and normal (mid-June) planting dates. Environmental conditions during (d) days to emergence, (e) days for vegetative, and (f) reproductive stage in different planting dates. Each violin plot represents the annual variability of 30 years (1986-2015) at each location in Kansas. Letters indicate significant differences ($\alpha < 0.05$) using the Tukey HSD test.

Figure 5. Water budgets under different levels of precipitation for a simulated full-season sorghum hybrid. (a) Total soil evaporation across planting dates (upper panel), soil evaporation during fallow periods (upper bars, light colors) and growing periods (lower bars, dark colors) under different planting dates. (b) Plant transpiration or water capture during the growing period (upper panel), and during vegetative (upper bars, light colors) and reproductive stages (lower bars, dark colors) under different planting dates. (c) Water stress events during the growing period (upper panel), and during vegetative (upper bars, light colors) and reproductive stages (lower bars, dark colors). Each violin plot represents the annual variability of 30 years (1986-2015) in Kansas. Each bar plot represents the median of each variable on each phenological stage. Letters represent significant differences ($\alpha < 0.05$) using the Tukey HSD test. Annual precipitation for dry, moderate, and wet years was 161-428 mm, 428-580 mm and 580-1380 mm, respectively.

Figure 6. Grain yield under different levels of precipitation and planting dates for simulated full- and short-season hybrids. Each violin plot represents the annual variability of 30 years (1986-2015) in Kansas. Letters indicate significant differences ($\alpha < 0.05$) of all pairwise comparisons using the Tukey HSD test. Annual precipitation for dry, moderate, and wet years was 161-428 mm, 428-580 mm, and 580-1380 mm, respectively.

SUPPLEMENTARY DATA

Included in the review PDF file:

Supplementary Tables 1-6

Supplementary Figures 1-2

Uploaded separately:

Supplementary File 1 - Code

Supplementary File 2 - Data

Table S1. Soil physical characteristics of study sites in Kansas.

Location	Depth (cm)	LL (cm cm^{-1})	DUL (cm cm^{-1})	SAT (cm cm^{-1})	RGF (-)	BD (g cm^{-3})	OC (%)	Clay (%)	Silt (%)	pH
Colby	0-10	0.21	0.39	0.43	1.00	1.43	1.16	30	52	7.2
	10-27	0.25	0.43	0.43	1.00	1.43	0.87	39	54	7.5
	10-56	0.16	0.34	0.44	0.44	1.43	0.29	25	63	7.5
	56-194	0.16	0.32	0.49	0.08	1.28	0.29	25	52	8.5
Garden City	0-15	0.18	0.37	0.49	1.00	1.3	1.2	24	60	7.9
	10-32	0.17	0.35	0.48	0.63	1.3	0.6	26	59	7.9
	32-74	0.15	0.33	0.46	0.35	1.4	0.3	23	64	8.5
	74-194	0.15	0.33	0.46	0.07	1.4	0.3	23	64	8.5
Hays	0-15	0.18	0.39	0.43	1.00	1.44	1.16	24	67	6.7
	15-22	0.19	0.40	0.47	1.00	1.31	1.16	26	67	6.7
	22-32	0.22	0.40	0.45	0.58	1.38	0.58	35	60	7.3
	32-64	0.23	0.41	0.45	0.38	1.38	0.46	39	56	7.3
	64-88	0.18	0.37	0.44	0.22	1.42	0.29	30	64	7.9
	88-194	0.16	0.35	0.44	0.06	1.42	0.29	25	69	7.9
Manhattan	0-17	0.19	0.41	0.44	1.00	1.4	1.7	24	64	6.5
	17-29	0.21	0.40	0.45	0.63	1.4	1.2	30	58	6.5
	29-39	0.22	0.40	0.44	0.51	1.4	0.9	33	58	6.5
	39-69	0.25	0.43	0.44	0.34	1.4	0.6	42	51	6.7
	69-91	0.23	0.40	0.45	0.20	1.4	0.4	39	54	6.7
	91-130	0.21	0.38	0.44	0.11	1.4	0.2	36	54	7.2
	130-194	0.19	0.36	0.44	0.04	1.4	0.1	32	56	7.2

LL, DUL, SAT and RGF were estimated using a pedotransfer function embedded in the Sbuild tool of DSSAT suit.
LL: lower limit
DUL: drainage upper limit
SAT: saturation
RGF: root growth factor
BD: bulk density
OC: organic carbon

Table S2. Phenology development and environmental conditions during the growing period
and during vegetative and reproductive stage for a simulated full- and short-season
hybrids in three planting dates across representative sites in Kansas. Each value represents
the overall mean over four locations and 30 years (1986-2015).

Planting date	Phenology	Days		Tmax (°C)		Tmin (°C)		Prec (mm)	
		mean	sd	mean	sd	mean	sd	mean	sd
Full season hybrid									
Very early	Total growing season	139	7	28.7	1.2	14.2	1.3	347	157
	Emergence	16	5	20.6	2.9	5.2	2.3	26	27
	Vegetative	79	1	28.6	1.4	14.1	1.2	220	111
	Reproductive	44	2	32.0	2.3	17.6	2.3	101	73
Early	Total growing season	124	6	30.5	1.6	16.0	1.6	319	151
	Emergence	10	2	25.3	3.4	10.9	2.4	25	31
	Vegetative	68	2	31.1	1.6	16.7	1.3	194	104
	Reproductive	46	2	30.8	2.4	16.2	2.5	100	71
Normal	Total growing season	116	2	30.2	1.6	15.5	1.4	263	139
	Emergence	6	1	31.0	2.7	16.1	2.3	19	26
	Vegetative	63	2	32.5	1.7	18.1	1.5	162	104
	Reproductive	48	5	27.4	2.3	12.2	2.2	82	62
Short season hybrid									
V. early	Total growing season	121	4	28.4	1.2	13.8	1.4	310	147
	Emergence	16	5	20.6	2.9	5.2	2.3	26	27
	Vegetative	68	1	27.9	1.5	13.4	1.3	185	100
	Reproductive	37	2	32.7	2.1	18.4	1.7	99	72
Early	Total growing season	107	5	30.8	1.4	16.4	1.4	286	136
	Emergence	10	2	25.3	3.4	10.9	2.4	25	31
	Vegetative	57	1	30.8	1.6	16.3	1.3	165	95
	Reproductive	40	2	32.2	2.1	17.9	1.8	97	67
Normal	Total growing season	102	2	31.2	1.9	16.5	1.7	240	135

Emergence	6	1	31.0	2.7	16.2	2.3	19	26
Vegetative	51	2	32.7	1.8	18.2	1.5	136	95
Reproductive	45	1	29.7	2.6	14.9	2.5	85	64

Emergence: from planting to emergence.

Vegetative: from emergence to anthesis.

Reproductive: from anthesis to harvest.

Table S3. Transpiration (mm) during the growing period and during vegetative and
reproductive stage for a simulated full- and short-season hybrids in three planting dates
across representative sites in Kansas. Each value represents the great mean over four
locations and 30 years (1986-2015).

Hybrid	Planting date	Total (mm)		Vegetative (mm)		Reproductive (mm)	
		mean	sd	mean	sd	mean	sd
Full season							
	V. early	236	104	149	58	87	61
	Early	232	97	147	53	86	58
	Normal	211	84	141	44	70	53
Short season							
	V. early	200	99	119	56	80	56
	Early	202	81	120	42	81	52
	Normal	178	67	107	36	72	44

Source of variation	df	Total	Vegetative	Reproductive
Planting date (PD)	2	<0.001	<0.001	<0.001
Hybrid (V)	1	<0.001	<0.001	0.1
Precipitation levels (dnw)	2	<0.001	<0.001	<0.001
PD x dnw	4	<0.001	0.3	<0.001

883 **Table S4. Water stress events (days) during the growing period and during vegetative and**
884 **reproductive stage for a simulated full- and short-season hybrids in three planting dates**
885 **across representative sites in Kansas. Each value represents the great mean over four**
886 **locations and 30 years (1986-2015).**

Hybrid	Planting date	Total mean	sd	Vegetative mean	sd	Reproductive mean	sd
Full season							
	V. early	31	19	10	10	21	13
	Early	32	19	9	9	23	14
	Normal	28	20	7	9	21	16
Short season							
	V. early	23	17	7	8	16	12
	Early	25	17	6	8	18	12
	Normal	24	19	5	7	19	15

Source of variation	df	Total	Vegetative	Reproductive
Planting date (PD)	2	0.04	<0.001	0.15
Hybrid (V)	1	<0.001	<0.001	<0.001
Precipitation levels (dnw)	2	<0.001	<0.001	<0.001
PD x dnw	4	0.16	0.2	<0.001

888 **Table S5. Grain yield (Mg ha⁻¹) for a simulated full- and short-season hybrids in three**
889 **planting dates across representative sites in Kansas. Each value represents the great mean**
890 **over four locations and 30 years (1986-2015).**

Hybrid	Planting date	Grain (Mg ha⁻¹) mean	sd	group (PDxV)
Full season				
	V. early	4.4	2.9	b
	Early	4.4	2.9	b
	Normal	4.0	2.5	a
Short season				
	V. early	3.7	2.5	a
	Early	3.7	2.4	a
	Normal	3.5	1.9	a

Source of variation	df	P>F	
Planting date (PD)	2	0.005	**
Hybrid (V)	1	<0.001	***
Precipitation levels (dnw)	2	<0.001	***
PD x V	2	0.08	
PD x dnw	4	<0.001	***

893 **Table S6. Simulated grain yield (Mg ha^{-1}) in different planting dates for a full and short**
894 **season hybrid in four contrasting sites in Kansas (1986-2015). Each value represents the**
895 **great mean over four locations and 30 years (1986-2015)**

Location	Planting date	Full season		Short season	
		mean	sd	mean	sd
All	V. early	4.4	2.9	3.7	2.5
	Early	4.4	2.9	3.7	2.3
	Normal	3.9	2.5	3.5	1.9
Colby	V. early	3.7	2.8	3.0	2.7
	Early	3.6	2.9	3.3	2.5
	Normal	2.6	1.8	2.9	2.0
Garden City	V. early	2.8	2.4	2.5	2.3
	Early	2.9	2.3	2.9	2.4
	Normal	2.5	1.9	2.5	1.9
Hayes	V. early	4.8	2.7	4.0	2.3
	Early	5.1	2.8	4.2	2.3
	Normal	4.9	2.2	4.3	1.9
Manhattan	V. early	6.2	2.6	5.0	2.5
	Early	6.0	2.6	4.4	1.9
	Normal	5.9	2.3	4.2	1.5

Figure S1. Seasonal evapotranspiration in four contrasting locations and different planting dates for a simulated full-and short-season hybrids. Each violin plot represents the annual variability of 30 years (1986-2015) in Kansas.

Figure S2. Soil extractable water at (A) anthesis and (B) maturity in four contrasting
locations and different planting dates in Kansas for a simulated full-and short-season
hybrids. Each violin plot represents the annual variability of 30 years (1986-2015).