Comparative metabolomics of fruits and leaves in a hyperdiverse lineage suggests fruits are a key incubator of phytochemical diversification

Gerald F. Schneider[1,2*], Diego Salazar[3], Sherry B. Hildreth[1,4], Richard F. Helm[4], Susan R. Whitehead[1]

[1] Department of Biological Sciences, Virginia Polytechnic Institute and State University, Blacksburg, VA, USA

[2] Department of Biology, Utah State University, Logan, UT, USA

[3] Department of Biological Sciences, International Center for Tropical Botany, Florida International University, Miami, FL, USA

[4] Department of Biochemistry, Fralin Life Sciences Institute, Virginia Polytechnic Institute and State University, Blacksburg, VA, USA

* Correspondence:
Gerald F. Schneider
jerry.schneider@usu.edu

Keywords: chemical diversity$_1$, molecular networking$_2$, metabolomics$_3$, fruit$_4$, seed$_5$, leaf$_6$, *Piper*$_7$

Word count: 6,272
Figures: 6
Tables: 4

Abstract

Interactions between plants and leaf herbivores have long been implicated as the major driver of plant secondary metabolite diversity. However, other plant-animal interactions, such as those between fruits and frugivores, may also be involved in phytochemical diversification. Using 12 species of *Piper*, we conducted untargeted metabolomics and molecular networking with extracts of fruits and leaves. We evaluated organ-specific secondary metabolite composition and compared multiple dimensions of phytochemical diversity across organs, including richness, structural complexity, and variability across samples at multiple scales within and across species. Plant organ identity significantly influenced secondary metabolite composition, both independent of and in interaction with species identity. Leaves and fruit shared a majority of compounds, but fruits contained more unique compounds and had higher total estimated chemical richness. While organ-level chemical richness and structural complexity varied substantially across species, fruit diversity exceeded leaf diversity in more species than the reverse. Furthermore, the variance in chemical composition across samples was higher for fruits than leaves. By documenting a broad pattern of high phytochemical diversity in fruits relative to leaves, this study lays groundwork for incorporating fruit into a

comprehensive and integrative understanding of the ecological and evolutionary factors shaping
secondary metabolite composition at the whole-plant level.

Introduction

Phytochemistry plays a key role in mediating the ecological and evolutionary dynamics of plant
interactions (Kessler & Baldwin, 2002; Wittstock & Gershenzon, 2002; Hartmann, 2007). As
functional traits, secondary metabolites can significantly affect plant fitness by defending plants
against antagonists, directly affecting the competitive ability of neighboring plants, protecting plants
from harsh environmental conditions, and attracting and rewarding mutualists, both above and below
ground (Iason *et al*., 2012). However, research on secondary metabolites and their role in the ecology
and evolution of plants has been disproportionately focused on vegetative organs, specifically the leaf
(e.g. Kursar *et al*., 2009; Richards *et al*., 2015; Volf *et al*., 2018; Salazar *et al*., 2018). While
secondary metabolites have numerous demonstrated functions mediating plant-animal interactions
surrounding leaves, they also likely perform a crucial and complex set of functions in reproductive
organs.

Plant reproductive organs have been a nexus of plant-animal interactions since before the emergence
of angiosperms. However, the ecological role that secondary metabolites play in the biology of these
plant organs has not been deeply explored. Fruits, and the seeds they contain, provide a direct link to
plant fitness and are therefore likely to be under intense selection pressure to attract mutualists and
deter antagonists. These complex and contrasting selective pressures are distinct from those acting on
leaves, and may lead to the occurrence of secondary metabolites not found in other organs. Indeed,
given the complex and often contrasting nature of selective pressures to which fruits and seeds are
exposed, fruits and seeds are likely to serve as evolutionary incubators of novel secondary
metabolites, and disproportionately contribute to the diversity of phytochemical traits. This is
especially likely in systems involving animal-mediated seed dispersal (zoochory), in which plants
face the ecological and physiological challenge of attracting and offering a nutritional reward to
dispersal vectors while also repelling seed predators, pathogens, and non-target frugivores (Herrera,
1982; Tewksbury, 2002; Whitehead *et al*., 2016).

Secondary metabolites endemic to fruits, and with demonstrated functional significance in seed
dispersal and/or fruit defense, have been shown in several systems, including iridoid glycosides in
honeysuckles (Whitehead & Bowers, 2013a, 2013b), capsaicinoids in *Capsicum* (Suzuki & Iwai,
1984; Tewksbury & Nabhan, 2001; Tewksbury *et al.*, 2008), and amides and alkenylphenols in *Piper*
(Whitehead *et al*., 2013, 2016; Whitehead & Bowers, 2014; Maynard *et al*., 2020). Further,
capsaicinoids in *Capsicum* and alkenylphenols in *Piper* are synthesized only in the fruits of these
taxa (Suzuki & Iwai, 1984; Maynard *et al*., 2020). Overall, these studies suggest that unique and
potentially contrasting selective pressures on fruits may be an important factor shaping
phytochemical diversification in plants. However, our understanding of the relative importance of
interactions across plant organs in shaping phytochemical diversity is limited by a paucity of studies
that compare chemical composition and metabolomic diversity across plant organs in an ecological
context.

Comparative metabolomic studies across plant organs have the potential to greatly expand our
understanding of secondary metabolite function and evolution. Given that metabolites may be organ-
specific, the location in which they are expressed in the plant (and consequently, the ecological
interactions in which they are involved) can provide valuable insight into both the evolutionary
origins and ecological consequences of the vast diversity of undescribed plant secondary metabolites.

Despite the likelihood of distinct selective pressures promoting divergent evolution of secondary
metabolites across plant organs, it is likely that the phytochemical diversity in one organ may be
constrained by physiological or genetic linkages with the phytochemistry of other organs (Adler *et*
al., 2006, 2012; Kessler & Halitschke, 2009; Keith & Mitchell-Olds, 2019). Physiological
constraints may result when a majority of the steps in a secondary metabolite pathway are localized
to a particular part of the plant, yielding complete or nearly complete end products that are then
transported to the organs in which they are utilized, e.g. glucosinolates in the Brassicaceae (Keith &
Mitchell-Olds, 2019). Such a pathway has a limited capacity to generate organ-specific modifications
of its end products prior to transport, and the sink organs may lack the metabolic machinery required
for such modifications. Other secondary metabolites are locally synthesized, but in this case organ-
specific metabolites derived from a shared metabolic pathway may be limited by genetic linkage,
through co-localization of genes responsible for modifications within a metabolic pathway, e.g.
terpene synthase clusters (Falara *et al.*, 2011; Chen *et al.*, 2020; Xu *et al.*, 2020). Certainly,
evolutionary processes may overcome these constraints when there are conflicting selection pressures
among organs, as evidenced by the examples above of compounds occurring only in specific organs.
Furthermore, even when fruits and leaves do share compounds, these compounds may be
quantitatively uncorrelated (Cipollini *et al.*, 2004; Whitehead & Bowers, 2013; Berardi *et al*., 2016).
Thus, while all plant species are biochemically circumscribed to some extent by the biosynthetic
pathways acquired through their evolutionary history, broad evolutionary patterns of such constraints
across plant organs have yet to be elucidated. Comparative metabolomics provide us with the tools to
define and characterize these patterns of constraint in conjunction with patterns of phytochemical
innovation.

In this study, we use comparative untargeted metabolomics to explore whether and how differential
selective pressures and constraints across reproductive and vegetative organs have shaped the
diversity and distribution of secondary metabolites in *Piper*, a pantropical species-rich genus. *Piper*
are diverse and dominant members of neotropical lowland forest understories and are known to
contain a rich array of secondary metabolites (Kato & Furlan, 2007; Richards *et al.*, 2015). Their
well-studied chemical composition and a long history of ecological research have made them a model
system for understanding phytochemical diversification and its role in shaping plant interactions and
community structure (Dyer & Palmer, 2004; Richards *et al.*, 2015; Salazar *et al.*, 2016).

Our overall objective in this study is to test the hypothesis that fruits can act as incubators of
phytochemical diversification in plants. First, we describe the occurrence patterns of secondary
metabolites across leaves, fruit pulp, and seeds in 12 *Piper* species, providing baseline data for
understanding *Piper* secondary metabolite function. We use untargeted mass spectrometry-based
metabolomics, molecular networking, and in-silico fragmentation modeling to characterize
undescribed metabolites, followed by machine learning and distance-based methods to compare
composition across organs and species. Second, we use these data to test predictions of high relative
diversity in fruits derived from our hypothesis of fruit-driven phytochemical diversification. We
compare multiple dimensions of phytochemical diversity across leaves and fruit organs, including the
richness at multiple scales (alpha and gamma diversity), variability (beta diversity), and structural
complexity of secondary metabolites.

Materials and Methods

Study system

Encompassing over 1,000 species across the Neotropics (Quijano-Abril *et al.*, 2006), the genus *Piper*
is diverse and abundant in forest understories, clearings, and edges (Gentry, 1990; Dyer & Palmer,

2004). *Piper* growth forms range from herbs and vines to shrubs and small trees (Gentry, 1990; Dyer
& Palmer, 2004). Fruits of Neotropical *Piper* are borne on distinct spike-shaped infructescences that
are dispersed primarily by bats of the genus *Carollia* (Phyllostomidae). Fruit antagonists of *Piper*
include insect seed predators, which have been found to consume up to 87% of seeds (Greig, 1993),
and a largely uncharacterized suite of pathogens, which rapidly attack fruit upon ripening (Thies &
Kalko, 2004; Whitehead & Bowers, 2014; Maynard *et al*., 2020). Leaves of *Piper* are subject to
herbivory from a broad array of arthropods, including a genus of specialist geometrid moths, *Eois*,
estimated to include over 1,000 species in the Neotropics (Brehm *et al.*, 2016), as well as other
geometrid moths, coleopterans, and orthopterans (Dyer & Palmer, 2004).

Field collections

All field collections took place between 2009 and 2012 at La Selva Biological Station, Heredia
Province, Costa Rica. Samples were collected during a phenology census across 28 species of *Piper*
during 2009-10 and opportunistically from 2010-12 when ripe fruits were available. Ripe fruits were
distinguished by a distinct softening and swelling of the fruit along an infructescence combined with
a partial senescence of the infructescence from the branch (presumably to allow bats to easily remove
the entire infructescence in flight). In most *Piper* species included in this study, one or a few
infructescences ripen per day per plant during the fruiting period, and the vast majority of these are
removed on the same night of ripening by bats (Thies & Kalko, 2004; Maynard *et al.*, 2020). Those
that are not removed rapidly decompose; therefore, we always took care to collect freshly-ripened
infructescences. We chose 12 species for inclusion in this study for which we were able to obtain
collections from at least three individual plants. For each individual, we collected 1-2 ripe
infructescences and the unripe infructescences that were immediately distal to the ripe ones on the
same branch. Fruits on a *Piper* branch mature sequentially from the proximal to the distal end of the
branch; thus, these adjacent unripe infructescences were the next closest to maturity on that branch.
Leaves were collected from the same branch. We chose the youngest fully-expanded leaf that did not
have extensive herbivore damage. All samples were transported immediately to the laboratory
(within 2 hours) and frozen at -80^{o}C prior to analysis. Subsequent analyses involve four sample
types: complete leaves, pulp from unripe and ripe infructescences, and seeds from ripe
infructescences.

Chemical extractions

The frozen plant material was freeze-dried ($-20°$ C/ $-55 °$ C, shelf/condenser), then ground to a fine
powder using a FastPrep-24 homogenizer. Seeds and pericarp were separated prior to grinding by
gently rubbing the dried fruit over fine mesh; the lignified central rachis of the infructescence was
discarded. In unripe fruit, seeds that were not sufficiently developed to be separated from the pericarp
by this method were homogenized with the pericarp. For each sample, 50 mg of homogenized
powder was weighed into a 2 mL Eppendorf tube using a microbalance. To isolate the broadest
possible range of phytochemicals while excluding the broadest possible range of primary
metabolites, extracts were prepared using buffered acetonitrile and acetone in series. The acetonitrile
and acetone extraction solutions were prepared with an aqueous acetate buffer (44.3 mmol/L
ammonium acetate), both at 70:30 solvent: buffer, v/v. The solutions were prepared with Nanopure®
water, Fisher HPLC-grade acetic acid, and Fisher Optima®-grade ammonium acetate, acetonitrile,
and acetone. All containers and instruments coming into contact with the extracts were rinsed with
Fisher Optima®-grade methanol. Each 50 mg sample was extracted twice with 1.5 mL buffered
acetonitrile, then twice more with 1.5 mL buffered acetone (6.0 mL total extraction solution). During
each of these four extractions, the sample was mixed with the extraction solvent for 5 min in a

vortexer, and then centrifuged for 5 min at 15870 rcf, after which the supernatant was removed and
added to a 20 mL glass scintillation vial. The supernatant from each of the four extractions was
combined in the same 20 mL vial. The combined extract was dried at 30° C using a nitrogen
evaporator until no solvent was visible, then further dried in a lyophilizer for 12 h (-20° C/ -55 ° C,
shelf/condenser) before being transferred to storage at -80° C until analysis.

Untargeted metabolomics

LC-MS data were collected using an Acquity I-class UPLC coupled to a Waters Synapt G2-S
quadrupole time-of-flight mass spectrometer (Waters). For analysis, dried extracts were resuspended
at 10 mg/mL in 75:25 water: acetonitrile + 0.1 % formic acid, with 1.0 µg/mL *N*-oleoylglycine as an
internal standard. The extract was then sonicated for 10 min, after which a 20 µL aliquot was taken
and diluted 10-fold with 75:25 water: acetonitrile + 0.1 % formic acid. The diluted aliquot was then
vortexed and centrifuged (10 min,13,000 xg) and an aliquot (180 µL) was transferred to an LC-MS
vial for analysis. Solvent blanks and combined, quality-control samples were injected at regular
intervals during data collection. The autosampler temperature was 10°C and the injection volume was
1.5 µL. The column employed was a reverse-phase Acquity BEH C18 (2.1 mm ID x 150 mm, 1.7 um
particle size, Waters) maintained at 35 °C at a flow rate of 0.2 mL/min. Solvent A was water with
0.1% formic acid and solvent B was acetonitrile with 0.1% formic acid (LCMS grade, Fisher
Chemical). Solvent gradient: 0-0.5 min, 90% A; 0.5-1.0 min, 75% A; 1.0-8.0 min, 5% A; 8.0-10.0
min, held at 5% A; 10.0-11.0 min, 90% A; 11.0-15.0 min, held at 90% A. Mass spectra and
fragmentation spectra were collected simultaneously using Waters' MS^E in positive-ion mode, with
the following parameters: peak data recorded in centroid mode; 0.185 s MS scan time; 20-35 V
collision energy ramp; argon collision gas; 125° C source temperature; 3 V capillary voltage; 30 V
sample cone voltage; 350° C desolvation temperature; nitrogen desolvation at 500 L/hr; 10 µL/min
lockspray flow rate; 0.1 s lockspray scan time; 20 s lockspray scan frequency; 3 lockspray scans to
average; 0.5 Da lockspray mass window; 3 V lockspray capillary voltage. The lockspray solution was
1 ng/ µL leucine enkephalin, and sodium formate was used to calibrate the mass spectrometer.

Alignment, deconvolution, and annotation of molecular and adduct ions were conducted using the
XCMS and CAMERA packages in R statistical software (Smith *et al.*, 2006; Tautenhahn *et al.*, 2008;
Benton *et al.*, 2010; Kuhl *et al.*, 2012).

Molecular networking

Molecular networking was used to quantify and visualize the dimensions of the chemical structural
trait space occupied by the secondary metabolites in our study (Aron *et al.,* 2020). This technique
employs tandem mass spectrometry to generate fragmentation spectra for each putative compound.
These fragmentation spectra are diagnostic of molecular structure, and through pairwise comparison
they are used to generate a network linking putative compounds to one another based on structural
similarity.

In our study, fragmentation spectra data files were aligned, deconvoluted, and converted to .mgf
using MS-DIAL software (v4.10) and were then uploaded to the Global Natural Products Social
Molecular Networking (GNPS) online workflow for molecular networking and library-based
annotation. The following parameters were used for the GNPS workflow METABOLOMICS-
SNETS-V2 (v14): 0.02 Da precursor ion mass tolerance; 0.02 Da fragment ion mass tolerance;
minimum matched fragment peaks = 6; minimum cluster size = 3; minimum cosine score for network
pairs = 0.7; network TopK = 1000; maximum connected component size = 0. All mass spectral
libraries available through GNPS which contained data collected in positive ion mode were used for

annotation. Library search parameters were: minimum matched peaks = 6; cosine score threshold =
0.6; maximum analog mass difference = 100. Workflow options for advanced filtering, advanced
GNPS repository search, and advanced output were not used.

For further annotation via *in-silico* modeling, results of the METABOLOMICS-SNETS-V2
workflow were passed to a second GNPS workflow, Network Annotation Propagation (NAP_CCMS
v1.2.5). The parameters used for NAP_CCMS were as follows: all clusters selected; subselection
cosine value = 0.7; first candidates for consensus score = 10; fusion results used for consensus;
accuracy for exact mass candidate search = 15 ppm; acquisition mode = positive; adduct ion types =
$[M+H]^+$ and $[M+Na]^+$; all structure databases selected; no custom database or parameter file;
compound class not specified; parent mass selection enabled; maximum number of graphed
candidate structures = 10; standard workflow type.

Finally, the outputs from METABOLOMICS-SNETS-V2 and NAP_CCMS were combined and
exported for visualization using the GNPS workflow MolNetEnhancer (v15). Network visualization
and curation was conducted using Cytoscape software (v3.7.2). Parent masses of features in the
molecular network were curated based on the XCMS-CAMERA output described above, with
primary metabolites and artefactual or pseudoreplicated features removed from the network and
subsequent analyses. Features in the molecular network were annotated to the level of chemical class,
e.g. flavonoid or prenol lipid, based on ClassyFire chemical taxonomy as applied by
MolNetEnhancer. The list of annotated molecular features returned by XCMS-CAMERA processing
was used to compare overall phytochemical composition across organs and species.

Unfragmented ions collected during single-mass-spectrometry and subsequently aligned,
deconvoluted, and annotated, as described above, were used to compare overall phytochemical
composition across organs and species. Ion abundance data were transformed to presence/absence
data using the peak recognition parameters in XCMS (R code repository). Ion presence/absence was
used for analyses rather than relative ion abundance for two reasons: 1) our sample size affords
limited capacity to account for variation in abundance within a given organ of a given species, and 2)
the scale of variation in ion abundance is likely to differ widely across the structurally diverse
compounds in *Piper* due to variation in ionization efficiency (Cech & Enke, 2001).

Comparisons of phytochemical composition across organs and species

To compare metabolome-level patterns of phytochemical composition across organs and species, we
conducted two separate analyses of the multivariate sample composition, focused first on compound
occurrences (presence/absence data) and second on the structural composition of samples. First, to
visualize differences in patterns of compound occurrence across samples, we used non-metric
multidimensional scaling (NMDS) based on the Sørensen dissimilarity index (binary Bray-Curtis).
We then tested for effects of organ, species, and their interaction on compound composition using
PERMANOVA, implemented with the 'adonis2' function in the R package 'vegan'. The individual
plant identity was included in these analyses as a 'strata' (i.e. random effect), and we used 999
permutations (note that this means the minimum possible *P*-value is $P = 0.001$, indicating that the
observed differences in sample composition could not be replicated in any of the 999 permutations).
To further understand specific differences among the four organ types, we followed this analysis with
post-hoc pairwise PERMANOVAs for all possible combinations of organ types, correcting for
multiple comparisons using the 'pairwise.adonis2' function (Martinez Arbizu, 2020). In addition,
based on strongly supported interactions between organ and species (see results), we also divided the

data by species and tested for the effects of organ on compound composition for each species
individually. All analyses were conducted using the 'vegan' package in R (Oksanen *et al.*, 2019).

In addition to our analysis of compound occurrence, we also examined how the structural
composition of samples was affected by organ, species, and their interaction. To account for
structural features, we generated a multivariate structural dissimilarity index that was a modification
of Sedio *et al.*'s (2017) Chemical Structural and Compositional Similarity (CSCS) index, which
quantifies the pairwise similarity of samples by calculating the maximum cosine similarity of the
aligned MS-MS ion fragmentation spectra for each inter-sample pair of molecular features. We
modified this index by representing ion abundance as a binary term and expressing the index in terms
of dissimilarity (1-CSCS). The structural dissimilarity matrix was then used as the basis for NMDS
and PERMANOVAs as above that examined the effects of organ, species, and their interaction on
structural composition.

Machine learning

To identify molecular features that distinguished different organs, we used random forest analysis via
the "randomForest" and "Boruta" packages for R statistical software (Liaw & Wiener, 2002; Kursa &
Rudnicki, 2010). All molecular features distinguished in XCMS-CAMERA processing were used as
variables in these analyses. The random forest analysis used a decision tree model to assign samples
to our four organ groups (Breiman, 2001; 2002). In the process, the analysis ranked molecular feature
variables according to their importance in the model's group assignments. Boruta analysis
complemented the random forest analysis by applying a search for molecular features that were
important in informing group assignments. This is accomplished by comparing the features'
importance with importance achievable at random, using "shadow" variables which are generated by
permuting the original variables (Kursa & Rudnicki, 2010).

Comparisons of chemical diversity across organs

Phytochemical diversity is a multifarious concept that includes the number of compounds (richness),
their relative abundances (evenness), their structural complexity, and their variation in space and time
(Wetzel & Whitehead, 2020). Considering the challenges associated with estimating abundances in
untargeted LC-MS-MS data, we focus here on richness and structural complexity, both of which
were examined at multiple scales within and across species. For each organ type, we define gamma
diversity as the total diversity observed across all samples for that organ, alpha diversity as the
average diversity within a single sample from one organ from one *Piper* individual, and beta-
diversity as the variation (both intra- and inter-specific) across samples.

Gamma diversity. To compare the gamma diversity (total number of compounds detected across all
species) of different organs, we used a rarefaction analysis analogous to those commonly used to
assess species diversity (Gotelli & Colwell, 2011) with compounds as "species" as in Wetzel &
Whitehead (2020). This allowed us to: 1) explicitly visualize the relationship between chemical
diversity and sampling scale across different organs (i.e. alpha, beta, and gamma diversity), and 2)
estimate the total compound richness in each organ type. Because our individual samples were not
independent (we collected three samples per species for 12 species), we used a constrained
rarefaction that is similar conceptually to spatially-constrained rarefaction (Chiarucci *et al.,* 2009).
Briefly, samples were added to bootstrapped accumulation curves in a semi-random manner in which
samples from the same species were grouped. For each iteration, a random sample was chosen as a
starting point, then other samples from that species were added in random order prior to choosing

another sample at random, following with all other samples from that species, and so on until all
species were included. We estimated total species richness from these curves using the
'fitspecaccum' function in 'vegan' based on an asymptotic regression model. Accumulation curves
and fits were averaged across 5000 bootstrapped samples with random starting points.

Alpha diversity. To compare the average compound richness in a sample (i.e. alpha diversity) across
organs, we used a linear mixed model with organ, species, and their interaction as fixed effects and
plant identity as a random effect. For hypothesis testing, we compared the full model to simplified
versions with fixed effects terms deleted using likelihood ratio tests. Based on a strong interaction
between organ and species (see results), we further divided the data by species and examined
differences in richness among organs for each species separately.

Structural complexity. To compare structural complexity across organ types, we first calculated an
index of structural complexity for each sample that was modified from the CSCS index described in
Sedio *et al*. (2017) to include only presence/absence data. This within-sample CSCS represents the
mean pairwise similarity among all individual molecular features detected in a sample. We used the
inverse of this similarity index (1-CSCS) as a measure of overall structural complexity present in a
sample. To examine how structural complexity varied across organs and species, we used a linear
mixed model with species, organ, and their interaction as fixed effects and plant identity as a random
effect. Hypothesis testing was conducted as described above using likelihood ratio tests. Based on
strong interactions between organ and species (see results), we examined differences among organs
separately for each *Piper* species.

Beta diversity. We examined differences in beta-diversity (i.e. sample-to-sample variance in
composition) across organs in two ways, focusing first on variation in compound occurrences
(presence/absence) and second on structural features. These analyses were based on the same
distance matrices described above that we used to assess overall differences in composition across
samples, but instead focused on variance (i.e. dispersion) among samples. This was assessed using
the function 'betadisper' in the R package 'vegan' to compare the dispersion around the group
centroid across the four organ types. The 'betadisper' function calculated the distances from each
sample to the group centroid, and statistical support for differences in dispersion across organs was
assessed using a permutation test (N = 999 permutations) followed by a post-hoc Tukey HSD test to
assess pairwise differences among individual organs. Because this analysis focused on sample-to-
sample variance and our dataset included multi-level sampling (multiple species and multiple
individuals within species), significant differences in beta diversity across organs could be due to
both intraspecific and interspecific variance among samples. Thus, we followed this analysis with a
set of PERMANOVAs, conducted separately for each organ type, with *Piper* species as an
explanatory factor. This analysis allowed us to test if *Piper* species explained a significant portion of
the variation in composition within an organ, and partitioned sample-to-sample variance within an
organ type according to the percent of variance explained by species and the percent explained by
differences among individuals within species (i.e. the residual variance).

Results

Untargeted metabolomics and molecular networking reveal high chemical diversity and many
compounds unique to fruits

Alignment, deconvolution, and annotation of molecular and adduct ions via XCMS and CAMERA
yielded 1,311 unique molecular features across all species and organs. Most of these compounds
(1,126) occurred in all organ types, 92 of these were unique to fruits (unripe pulp, ripe pulp, and/or

seeds) and 4 were unique to leaves (Fig. 1a). It is important to note that, like all other metabolomic
approaches, our analytical approach is likely to overestimate the true number of individual chemical
compounds present in our samples. The combination of XCMS-CAMERA followed by manual
curation unfortunately cannot condense all features (m/z and retention time pairs) into individual
compounds. In-source fragmentation, ion clusters, centroid peak splitting of highly abundant ions,
and centroid merging of ions near the noise level can all contribute to expanding the dataset beyond
individual compounds. The 1,311 features described in this work thus overestimates the number of
individual molecular species, though to a lesser extent than in uncurated datasets. Nevertheless, this
overestimation is likely to represent a small fraction of the total chemical diversity captured in our
analysis. Furthermore, this overestimation is also likely to be of equal magnitude across all species
and organs and therefore, will not have a significant impact on the general conclusions of our study.
Regarding terminology, these 1,311 features meet or exceed the level of curation beyond which
features have, for clarity, been described as "compounds" in the chemical ecology literature (e.g.:
Sedio *et al.*, 2017; Christian *et al.*, 2020; Ricigliano *et al.*, 2020). Thus, for the sake of consistency
and clarity, we refer to our curated features as compounds.

Tandem mass spectrometry yielded fragmentation spectra for 706 of these compounds (Table 1, Fig.
2). Library- and *in silico*-based classification of fragmentation spectra and parent ions via GNPS
resulted in annotation at the level of "class" *sensu* ClassyFire chemical taxonomy for 527 compounds
in 23 classes and 179 were unknowns (Table 1, Fig. 2). Of the 706 compounds that yielded
fragmentation spectra, 62 were unique to fruits and 2 were unique to leaves (Table 1). Fruit-specific
richness was higher than leaf-specific richness across all chemical classes, these differences were
statistically supported for carboxylic acids and derivatives and prenol lipids (Table 1; Fig. 2).

Phytochemical composition differs across organs and species

The multivariate patterns of phytochemical occurrence were strongly affected by organ, species, and
their interaction (organ: $F_{3,95} = 24.19$, $P = 0.001$; species: $F_{11,95} = 27.65$, $P = 0.001$; organ x species:
$F_{33,95} = 1.99$, $P = 0.001$; Fig. **3a**). Pairwise comparisons among organs indicated strong differences
among all organs ($P = 0.001$ for all comparisons). Further examination of differences among organs
for each of the 12 *Piper* species individually also revealed strong effects of organ in all cases (Table
2). Similarly, when we assessed factors influencing the multivariate patterns of structural
composition across samples, we found a strong effect of organ, species, and their interaction (organ:
$F_{3,95} = 17.34$, $P = 0.001$; species: $F_{11,96} = 21.28$, $P = 0.001$; organ x species: $F_{33,96} = 2.31$, $P = 0.001$;
Fig. **3b**), significant differences among organs in all pairwise comparisons ($P = 0.001$ for all
comparisons), and differences among organs for each individual species (Table 2).

Machine learning, informed by numerous compounds from diverse chemical classes in each
organ, accurately distinguishes between reproductive and vegetative organs

The random forest decision tree model used 2000 trees with 36 variables at each split. Our analysis
showed an overall out-of-bag (OOB) mean error rate of 11.72% across the four organ groups. In
other words, using secondary metabolites alone, the algorithm was able to predict if a sample was
from a leaf, ripe fruit, unripe fruit, or seed approximately 9 times out of every 10 samples. Examining
the error rate of each organ group, it was apparent that correctly assigning pulp samples to the correct
ripeness stage was the main source of OOB error, with error rates of 27.78% and 18.92% for unripe
and ripe pulp respectively. Leaves and ripe seeds both exhibited zero OOB error. Boruta analysis,
designed to both identify important classification features and assess their relative contribution to the

final classification performance, identified 23 features exhibiting a significantly higher variable
importance score (VIS) than shadow variables. These 23 features are detailed in Table S1.

Comparisons of chemical diversity across organs

Gamma diversity

Of the 1,311 compounds detected across all organ types, 1,126 were shared across all organs, 92
were found only in fruit (unripe pulp, ripe pulp, and/or seeds), and four were found only in leaves.
Rarefaction analysis indicated that the observed richness of secondary metabolites was approaching
the asymptote for all organ types (Fig. 1b). Further, this analysis showed that the estimated total
gamma diversity (total number of compounds across all 12 species of *Piper*) was highest in unripe
and ripe fruit pulp, intermediate in seeds, and lowest in leaves, with only unripe and ripe pulp
exhibiting overlapping confidence intervals (Fig. 1b, Table 3).

Alpha diversity

In our analysis of average differences in compound richness across organs and species, we found a
strong interaction between organ and species ($X^2 = 128.99$, $P < 0.0001$) and further examined
differences among organs for each species separately. Organs often showed clear differences in
average richness, but the patterns were highly variable across species (Fig. 4). In three species (*P.
glabrescens, P. reticulatum,* and *P. slivivagum*), pulp and/or seeds had higher compound richness
than leaves. However, in two species (*P. multiplinervum* and *P. generalense*) leaves had higher
compound richness than all other fruit organs.

Structural complexity

In our analysis of average differences in structural complexity across organs and species, we found a
strong interaction between organ and species ($X^2 = 131.13$, $P < 0.0001$) and further examined
differences among organs for each species separately. For seven of twelve species, organs showed
differences in average structural complexity, but the patterns were variable across species (Fig. 5). In
two species (*P. glabrescens* and *P. slivivagum*), one or more fruit organs had higher complexity than
leaves. In another species (*P. generalense*), leaves had higher complexity than all other fruit organs.
Often, seeds had the lowest structural complexity, or at least lower structural complexity than unripe
or ripe fruit pulp (Fig. 5).

Beta diversity

We found that beta-diversity in chemical composition was higher for fruits than leaves when
considering only compound occurrences as well as structural composition. First, for compound
occurrences, there was strong support for overall differences in beta diversity across organ types
($F_{3,139} = 7.56$, $P = 0.001$), with higher average distances to the group centroid for seeds, unripe pulp,
and ripe pulp relative to leaves (Fig. **6a**). Next, for structural composition, there was also strong
support for overall differences in beta diversity across organ types ($F_{3,139} = 4.10$, $P = 0.009$). In this
case, leaves had lower beta diversity than seeds or ripe pulp, and unripe pulp was intermediate (Fig.
6b). Further analyses conducted separately for each organ type showed that the differences in beta-
diversity among organ types was due to variation both at the interspecific and intraspecific level
(Table 4). A large proportion of the sample-to-sample variation within organ types (67-86%) was
explained by differences among species relative to that explained by variation within species (14-
33%), and this was especially true for unripe and ripe fruit pulp (Table 4).

Discussion

Secondary metabolites occur in all plant parts, but theory in chemical ecology has largely focused on
interactions between leaves and leaf herbivores. In this study, we surveyed and compared the
compositional and structural diversity of secondary metabolites across vegetative and reproductive
organs in 12 species of the genus *Piper*. Most metabolites were shared across organs, but fruits
contained many more unique compounds than leaves. Furthermore, relative to leaves, fruits contained
a higher total number of compounds (gamma diversity) and a higher sample-to-sample variance in
composition (beta diversity). While *Piper* species varied in terms of whether fruits or leaves had a
higher average compound richness per sample (alpha diversity), richness was more often higher in
fruits than in leaves. Taken together, these patterns reveal that, in neotropical *Piper*, fruits are overall
more chemically diverse than leaves and point to a key role for mutualistic and antagonistic fruit-
frugivore interactions in shaping phytochemical evolution and diversity.

Our untargeted metabolomic survey of phytochemical occurrence patterns revealed that fruit organs
harbor fruit-specific metabolites from a variety of chemical classes (Table 1, Fig. 2), in each class
equal to or greater in number than those that were leaf-specific (Table 1). This included classes of
compounds that have previously been found to be more numerous and abundant in fruit organs (e.g.
amides; Whitehead *et al.*, 2013), as well as numerous chemical classes previously described in
studies of *Piper* spp. leaf chemistry (Parmar *et al.*, 1997; Baldoqui *et al.*, 1999; Kato & Furlan, 2007;
Richards *et al.*, 2015). The occurrence of numerous fruit-specific secondary metabolites from a
variety of unlinked biosynthetic pathways suggests a pattern of fruit-specific secondary metabolite
trait evolution, likely a result of fruit-specific selective pressures.

The evolution of organ-specific phytochemical traits across our target plant species is also made
evident by the results of our machine learning analysis. Here, our random forest model was very
successful at distinguishing among organ types based solely on their secondary metabolite
composition. Most notably, the exceptional performance of the classification algorithm to distinguish
between vegetative and reproductive organs can only be explained by the presence of strong
association between chemical composition and organ type. Despite the fact that our species set
included vines, understory shrubs, and pioneering taxa, all adapted to very different local habitats,
these associations are consistent across all 12 focal species.

The clustering patterns found by our NMDS analysis (Fig. 3) show clustering at two different levels.
First, the samples from different organs from the same species cluster together. This pattern strongly
suggests the presence of physiological or genetic linkage constraints in the organ specific evolution
of phytochemicals. The strong chemical similarity across organs within a species could point toward
the influence that changes in the expression or composition of secondary compounds in one plant
organ could have on the expression or composition of other organs. Although our data do not allow
us to disentangle the precise mechanisms that give rise to these patterns, it is clear that chemical
changes in one plant organ are likely to be mirrored, to some extent, by changes in the chemical
architecture of the whole plant. Second, as expected, and despite the strong chemical similarity
exhibited by organs within a *Piper* species, samples also show a clear pattern of clustering by organ
type (Fig. 3). This pattern reinforces the expectation that the distinctive regimes of selective pressures
imposed upon the different plant organs are sufficiently strong to create convergent organ-specific
patterns of the chemical composition, and that these selective regimes are likely to be consistent
across species and habitats. The Boruta variable importance model, a widely used machine learning
algorithm designed to identify statistically important classification variables from large datasets,
revealed specific compounds from at least six different chemical classes as key features that
distinguish vegetative and reproductive organs (Table S1).

While the majority of significant Boruta variables, like the overall majority of secondary metabolites
cataloged in our study, exhibited some overlap in occurrence across leaf and fruit organs when the 12
Piper species were evaluated as a group (Fig. 1a), there was substantially less overlap at the level of
individual species (Fig. S1). In many cases, these patterns of variance were the result of numerous
compounds occurring in only one organ type in a certain species or subset of species, but occurring
more widely in another species or subset of species.

The broad overlap across organs in compound occurrence at the genus level provides a degree of
insight into the extent of constraints on organ-specific chemical trait evolution at this taxonomic
scale. However, to a degree this overlap can also be attributed to the shared demand for defensive
compounds across vegetative and reproductive organ types. While phylogenetic data will be required
in order to infer the ancestral organ localizations of phytochemical traits of *Piper*, the widespread
variation in organ localizations that we observed across species suggests that genetic constraints have
not bound these traits to a certain organ type over the course of *Piper* speciation. Further, the
apparent mobility of secondary metabolite traits across organ types within the genus suggests a
bidirectional exchange of these traits, which, when vegetative and reproductive organs are each
threatened by separate assemblages of consumers, may allow more rapid defense trait adaptation than
can arise from novel mutations.

Our untargeted metabolomic survey has shown that fruit organs are at the very least a reservoir of
phytochemical richness. While the alpha diversity of organs at the species level was highly variable
(Fig. 4), rarefaction analysis of gamma diversity showed a small but clear trend towards higher
richness of secondary metabolites in reproductive organs (Fig. 1b; Table 3). Similarly, while
chemical structural complexity of organs at the species level was highly variable (Fig. 5), chemical
structural variance ($\square$-diversity) across species was significantly higher for reproductive organs than
for leaves (Fig. 6). In summary, these trends indicate not only that reproductive organs accumulate a
higher number of secondary metabolite traits than do leaves, but also that these traits are more
divergent from one another across species than those of leaves. These trends are consistent with
higher overall evolutionary diversification of phytochemical traits in reproductive organs, suggesting
that fruits may be an important, but underappreciated, force in shaping chemical trait evolution at the
whole plant level.

Acknowledgements

This research was supported by National Science Foundation (Grants No. DEB-1210884 and DEB-
1856776 to SRW) and start-up funds to SRW from the Virginia Tech Department of Biological
Sciences. The mass spectrometry resources used in this work were maintained with funds from the
Fralin Life Science Institute as well as the Virginia Agricultural Experiment Station Hatch Program
(VA-160085). We thank the Organization for Tropical Studies for logistical support at La Selva
Biological Station. Marisol Luna Martinez assisted with sample collection and Orlando Vargas
Ramírez assisted with plant identification. Natalie Rodeman and Katherine Berg assisted with sample
processing for chemical analyses.

Author contributions

GFS, SRW, and DS designed the research; SRW collected field samples; GFS, SBH, and RFH
conducted chemical analyses, GFS conducted molecular networking and data curation; GFS and
SRW conducted the statistical analysis with contributions from DS; GFS wrote the first draft of the

manuscript with contributions from SRW, and all authors contributed substantially to revisions and
approved the final version.

Data availability

520 UPLC-MS data and associated programming scripts will be made publicly available through Global
521 Natural Products Social Molecular Networking (https://gnps.ucsd.edu/) and all other data and
522 programming scripts that support the findings of this study will be made publicly available at Zenodo
523 https://www.zenodo.org/ upon publication of this manuscript.

References

Adler LS, Seifert MG, Wink M, Morse GE, Turlings T. **2012**. Reliance on pollinators predicts
defensive chemistry across tobacco species. *Ecology Letters* **15**: 1140–1148.

Adler LS, Wink M, Distl M, Lentz AJ. **2006**. Leaf herbivory and nutrients increase nectar
alkaloids. *Ecology Letters* **9**: 960–7.

**Aron AT, Gentry EC, McPhail KL, Nothias LF, Nothias-Esposito M, Bouslimani A, Petras
D, Gauglitz JM, Sikora N, Vargas F, *et al.* 2020**. Reproducible molecular networking of
untargeted mass spectrometry data using GNPS. *Nature Protocols* **15**: 1954–1991.

Benton HP, Want EJ, Ebbels TMD. 2010. Correction of mass calibration gaps in liquid
chromatography-mass spectrometry metabolomics data. *BIOINFORMATICS* **26**: 2488.

Baldoqui DC, Kato MJ, Cavalheiro AJ, Bolzani VDS, Young MCM, Furlan M. 1999. A
chromene and prenylated benzoic acid from *Piper aduncum*. *Phytochemistry* **51**: 899–902.

Berardi AE, Hildreth SB, Helm RF, Winkel BSJ, Smith SD. **2016**. Evolutionary correlations in
flavonoid production across flowers and leaves in the Iochrominae (Solanaceae). *Phytochemistry*
130: 119–127.

**Brehm G, Hebert PDN, Colwell RK Adams MO, Bodner F, Friedemann K, Möckel
L,Fiedler K. 2016.** Turning up the heat at a hotspot: DNA barcodes reveal 80% more species of
geometrid moths along an Andean elevational gradient. *PlosOne* **11:** e0150327. doi:
10.1371/journal.pone.0150327

Breiman L. 2001. Random forests. *Machine learning* **45**: 5-32.

Breiman L. 2002. Manual on setting up, using, and understanding random forests v3. 1. URL:
www.stat.berkeley.edu/~breiman/Using_random_forests_V3.1.pdf [accessed 20 January 2021].

Cech NB, Enke CG. 2001. Practical implications of some recent studies in electrospray ionization
fundamentals. *Mass Spectrometry Reviews* **20**: 362–387.

Chen H, Köllner TG, Li G, Wei G, Chen X, Zeng D, Qian Q, Chen F. 2020. Combinatorial
evolution of a terpene synthase gene cluster explains terpene variations in *Oryza*. *Plant Physiology*
182: 480–492.

Chiarucci A, Bacaro G, Rocchini D, Ricotta C, Palmer M, Scheiner S. 2009. Spatially
constrained rarefaction: Incorporating the autocorrelated structure of biological communities into
sample-based rarefaction. *Community Ecology* **10**: 209–214.

Christian N, Sedio BE, Florez-Buitrago X, Ramírez-Camejo LA, Rojas EI, Mejía LC, Palmedo S, Rose A, Schroeder JW, Herre EA. 2020. Host affinity of endophytic fungi and the potential for reciprocal interactions involving host secondary chemistry. *American Journal of Botany* **107**: 219–228.

Dyer LA, Palmer ADN. **2004**. *Piper: A model Genus for Studies of Phytochemistry, Ecology, and Evolution*. Kluwer Academic/ Plenum Publishers.

Dyer LA, Philbin CS, Ochsenrider KM, Richards LA, Massad TJ, Smilanich AM, Forister ML, Parchman TL, Galland LM, Hurtado PJ*, et al.* **2018**. Modern approaches to study plant - insect interactions in chemical ecology. *Nature Reviews Chemistry* **2**: 50–64.

Falara V, Akhtar TA, Nguyen TTH, Spyropoulou EA, Bleeker PM, Schauvinhold I, Matsuba Y, Bonini ME, Schilmiller AL, Last RL*, et al.* **2011.** The tomato terpene synthase gene family. *Plant Physiology* **157**: 770–789.

Gentry AH. 1990. *Four neotropical rainforests.* Yale University Press, New Haven.

Gotelli NJ, Colwell RK. 2011. Estimating species richness. In: Magurran AE, McGill BJ, eds. *Biological Diversity: Frontiers in Measurement and Assessment*. Oxford, UK: Oxford University Press, 39–54.

Greig N. 1993. Predispersal seed predation on five *Piper* species in tropical rain-forest. *Oecologia* **93**: 412–420

Hartmann T. **2007**. From waste products to ecochemicals: Fifty years research of plant secondary metabolism. *Phytochemistry* **68**: 2831–2846.

Herrera CM. 1982. Defense of ripe fruit from pests: Its significance in relation to plant-disperser interactions. *The American Naturalist* **120**: 218–241.

Iason GR, Dicke M, Hartley SE. 2012. *The ecology of plant secondary metabolites: From genes to global processes.* Cambridge, UK: Cambridge University Press.

Kato MJ, Furlan M. **2007**. Chemistry and evolution of the Piperaceae. *Pure and Applied Chemistry* **79**: 529–538.

Keith RA, Mitchell-Olds T. **2019**. Antagonistic selection and pleiotropy constrain the evolution of plant chemical defenses. *Evolution* **73**: 947–960.

Kessler A, Baldwin IT. **2002**. Plant responses to insect herbivory: The emerging molecular analysis. *Annual Review of Plant Biology* **53**: 299–328.

Kessler A, Halitschke R. **2009**. Testing the potential for conflicting selection on floral chemical traits by pollinators and herbivores: Predictions and case study. *Functional Ecology* **23**: 901–912.

Kuhl C, Tautenhahn R, Boettcher C, Larson TR, Neumann S. 2012. CAMERA: an integrated strategy for compound spectra extraction and annotation of liquid chromatography/mass spectrometry data sets. *Analytical Chemistry*, **84**, 283–289.

Kursa MB, Rudnicki WR. 2010. Feature selection with the Boruta package. *Journal of Statistical Software* **36**: 1-13.

Kursar TA, Dexter KG, Lokvam J, Pennington RT, Richardson JE, Weber MG, Murakami ET, Drake C, McGregor R, Coley PD. **2009**. The evolution of antiherbivore defenses and their contribution to species coexistence in the tropical tree genus *Inga*. *Proceedings of the National Academy of Sciences of the United States of America* **106**: 18073–18078.

Liaw A, Wiener M. 2002. Classification and regression by randomForest. *R News* **2**: 18-22.

Martinez Arbizu P. 2020. pairwiseAdonis: Pairwise multilevel comparison using adonis. R
package version 0.4. URL https://github.com/pmartinezarbizu/pairwiseAdonis . [accessed 9
January 2021].

Maynard LD, Slinn HL, Glassmire AE, Matarrita-Carranza B, Dodson CD, Nguyen TT,
Burroughs MJ, Dyer LA, Jeffrey CS, Whitehead SR. 2020. Secondary metabolites in a
neotropical shrub: spatiotemporal allocation and role in fruit defense and dispersal. *Ecology* **101**:
1–15.

Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR,
O'Hara RB, Simpson GL, Solymos P, *et al.* 2019. vegan: Community Ecology Package. R
package version 2.5-6. URL https://CRAN.R-project.org/package=vegan . [accessed 9 January
2021]

Parmar VS, Jain SC, Bisht KS, Jain R, Taneja P, Jha A, Tyagi OD, Prasad AK, Wengel J,
Olsen CE, et al. 1997. Phytochemistry of the genus *Piper*. *Phytochemistry* **46**: 597–673.

Richards LA, Dyer LA, Forister ML, Smilanich AM, Dodson CD, Leonard MD, Jeffrey CS.
2015. Phytochemical diversity drives plant–insect community diversity. *Proceedings of the*
National Academy of Sciences of the United States of America **112**: 10973–10978.

Ricigliano VA, Sica VP, Knowles SL, Diette N, Howarth DG, Oberlies NH. 2020. Bioactive
diterpenoid metabolism and cytotoxic activities of genetically transformed Euphorbia lathyris
roots. *Phytochemistry*. doi:10.1016/j.phytochem.2020.112504

Salazar D, Jaramillo MA, Marquis RJ. **2016**. Chemical similarity and local community
assembly in the species rich tropical genus *Piper*. *Ecology* **97**: 3176–3183.

Salazar D, Lokvam J, Mesones I, Pilco MV, Milagros J, Zuñiga A, Valpine P De, Fine PVA.
2018. Origin and maintenance of chemical diversity in a species-rich tropical tree lineage. *Nature*
Ecology & Evolution **2**: 983–990.

Schneider GF, Coley PD, Younkin GC, Forrister DL, Mills AG, Kursar TA. 2019. Phenolics
lie at the centre of functional versatility in the responses of two phytochemically diverse tropical
trees to canopy thinning. *Journal of Experimental Botany* **70**: 5853–5864.

Sedio BE, Echeverri JCR, Boya CA, Wright SJ. **2017**. Sources of variation in foliar secondary
chemistry in a tropical forest tree community. *Ecology* **98**: 616–623.

Smith CA, Want EJ, O'Maille G, Abagyan R, Siuzdak G. 2006. XCMS: Processing mass
spectrometry data for metabolite profiling using nonlinear peak alignment, matching and
identification. *Analytical Chemistry* **78**: 779–787.

Suzuki T, Iwai K. 1984. Constituents of red pepper species: Chemistry, biochemistry,
pharmacology, and food science of the pungent principle of Capsicum species. In: Brossi A, ed.
The Alkaloids, Volume 23. Academic Press, 227–299.

Tautenhahn R, Boettcher C, Neumann S. 2008. Highly sensitive feature detection for high
resolution LC/MS. *BMC Bioinformatics*. doi: 10.1186/1471-2105-9-504.

Tewksbury JJ. 2002. Fruits, frugivores and the evolutionary arms race. *New Phytologist* **156**:
137–139.

Tewksbury JJ, Nabhan GP. **2001**. Seed dispersal: Directed deterrence by capsaicin in chillies.
Nature **412**: 403–404.

Tewksbury JJ, Reagan KM, Machnicki NJ, Carlo TA, Haak DC, Peñaloza ALC, Levey DJ.
2008. Evolutionary ecology of pungency in wild chilies. *Proceedings of the National Academy of*
Sciences of the United States of America **105**: 11808–11811.

Thies W, Kalko EKV. 2004. Phenology of Neotropical pepper plants (Piperaceae) and their
association with their main dispersers, two short-tailed fruit bats, *Carollia perspicillata* and *C.*
castanea (Phyllostomidae). *Oikos* **104**: 362–376.

Volf M, Segar ST, Miller SE, Isua B, Sisol M, Aubona G, Šimek P, Moos M, Laitila J, Kim J,
***et al.* 2018**. Community structure of insect herbivores is driven by conservatism, escalation and
divergence of defensive traits in *Ficus*. *Ecology Letters* **21**: 83–92.

Wang M, Carver JJ, Phelan V V, Sanchez LM, Garg N, Peng Y, Nguyen DD, Watrous J,
Kapono CA, Luzzatto-Knaan T, *et al.* 2016. Sharing and community curation of mass
spectrometry data with Global Natural Products Social Molecular Networking. *Nature*
Biotechnology **34**: 828–837.

Wetzel WC, Whitehead SR. **2020**. The many dimensions of phytochemical diversity: linking
theory to practice. *Ecology Letters* **23**: 16–32.

Whitehead SR, Bowers MD. 2013a. Iridoid and secoiridoid glycosides in a hybrid complex of
bush honeysuckles (*Lonicera* spp., Caprifoliaceae): Implications for evolutionary ecology and
invasion biology. *Phytochemistry* **86**:57–63.

Whitehead SR, Bowers MD. 2013b. Evidence for the adaptive significance of secondary
compounds in vertebrate-dispersed fruits. *The American Naturalist* **182**: 563–577.

Whitehead SR, Bowers MD. 2014. Chemical ecology of fruit defence: synergistic and
antagonistic interactions among amides from *Piper*. *Functional Ecology* **28**: 1094–1106.

Whitehead SR, Jeffrey CS, Leonard MD, Dodson CD, Dyer LA, Bowers MD. 2013. Patterns
of secondary metabolite allocation to fruits and seeds in *Piper reticulatum*. *Journal of Chemical*
Ecology **39**: 1373–1384.

Whitehead SR, Quesada MFO, Bowers MD. **2016**. Chemical tradeoffs in seed dispersal:
defensive metabolites in fruits deter consumption by mutualist bats. *Oikos* **125**: 927–937.

Wittstock U, Gershenzon J. **2002**. Constitutive plant toxins and their role in defense against
herbivores and pathogens. *Current Opinion in Plant Biology* **5**: 300–307.

Xu S, Kreitzer C, McGale E, Lackus ND, Guo H, Köllner TG, Schuman MC, Baldwin IT,
Zhou W. 2020. Allelic differences of clustered terpene synthases contribute to correlated
intraspecific variation of floral and herbivory-induced volatiles in a wild tobacco. *New*
Phytologist. doi: 10.1111/nph.16739

Figure 1: Chemical gamma diversity parsed by organ type. A Venn diagram (**A**) shows the total

number of compounds detected across all samples that were unique and shared across organ type.

The rarefaction curve (**B**) shows how compound richness accumulates with sampling scale in each

organ type. Curves represent an average across 5000 bootstrapped accumulation curves with random

starting points. Because samples from the same species were not independent, the rarefaction was

constrained by species such that samples from the same species were always added in sequence.

Figure 2: Molecular network of 706 compounds from 12 *Piper* species color-coded by ClassyFire

chemical classification annotation (**A**) or by organ-level occurrence across the 12 species (**B**). Node

and edge arrangement and compound annotation are as described in "Molecular Networking"

methods. Enlarged, diamond-shaped nodes represent compounds identified by the Boruta analysis as

important for distinguishing among organs. In (**B**), compounds are coded as occurring in "fruit" if

they occur in one or more of the three sample types (unripe pulp, ripe pulp, or seeds).

Figure 3: NMDS plots showing the effects of organ and species on two aspects of multivariate

chemical composition across samples: (**A**) compound occurrences (presence/absence) and (**B**)

structural composition.

Figure 4: Average chemical richness differs across species and organ type (leaf, seed, unripe pulp,

and ripe pulp). Letters indicate results of pairwise Tukey post-hoc comparisons of organs within each

species, with non-shared letters indicating a significant difference at $P < 0.05$. Each species plot

includes $\square^2$ and P-values from species-level LMMs.

Figure 5: Average structural complexity differs across species and organ type (leaf, seed unripe pulp,

and ripe pulp). Letters above each box plot column indicate results of pairwise Tukey post-hoc

comparisons of organs within each species, with non-shared letters indicating a significant difference

at $P < 0.05$. Each species plot includes $\square^2$ and P-values from species-level LMMs.

Figure 6: Beta diversity in chemical composition is higher for reproductive organs than leaves when

considering variance in compound occurrences (**A**) or structural composition (**B**). Letters indicate

results of pairwise Tukey post-hoc comparisons of organs, with non-shared letters indicating a

significant difference at *P* < 0.05.

Table 1: Summary of GNPS molecular network annotations. Compound richness indicates the number of putative compounds, for which fragmentation spectra were obtained, that fall under the given category. Chemical classes are per ClassyFire chemical taxonomy. Chemical classes which represented ≤ 1% of all annotated compounds were categorized as "Other". A compound was labeled as fruit or leaf-specific if it was detected only in that organ within the 12 focal *Piper* species. Asterisks indicate statistical support (P < 0.05) for differences between fruit-specific and leaf-specific richness (binomial test with probability = 0.5).

Chemical Class	Examples of class known from *Piper* spp.	Total Compound Richness	Fruit-specific Compound Richness	Leaf-specific Compound Richness
Benzene and substituted derivatives	Cyanogenic benzoates, Non-prenylated benzoic acids	75	5	0
Carboxylic acids and derivatives	Amides, Chromenes, Kavalactones	122	25*	1
Flavonoids	Flavonoids	104	3	0
Organo-oxygen compounds	Oxygenated or glycosidic derivatives of other classes	65	5	0
Other	Amides, Chalcones, Chromenes, Imides	37	4	0
Prenol lipids	Chalcones, Prenylated benzoic	124	6*	0

acids, Neryl catechol diols, Terpenes		
Unknown	179	14*	1
Total	706	62*	2

Table 2: Results from PERMANOVAs, conducted separately for each species, testing the effects of organ type (leaves, seed, unripe pulp, or ripe pulp) on two aspects of phytochemical composition: compound occurrences and structural composition

Piper species	Compound Occurrence		Structural Composition	
	$F_{3,11}$	*P*	$F_{3,11}$	*P*
aduncum	3.07	0.001	3.07	0.001
auritum	3.51	0.002	3.51	0.002
biolleyi	2.19	0.009	2.19	0.009
colonense	4.39	0.001	4.39	0.001
generalense	4.11	0.001	4.11	0.002
glabrescens	4.18	0.002	4.18	0.001
multiplinervum	3.45	0.001	3.45	0.001
peltatum	4.50	0.001	4.50	0.001
reticulatum	5.38	0.001	5.38	0.002
sancti-felicis	3.67	0.001	3.67	0.001

silvivagum	5.19	0.001	5.19	0.001
umbricola	2.89	0.002	2.89	0.001

Table 3: Rarefaction results showing total estimated richness for each organ across all 12 Piper species sampled

Organ	Observed Richness	Estimated Richness	SE	95% CI high	95% CI low
leaf	1219	1226	1.5	1229.3	1223.5
seed	1222	1276	3.0	1282.0	1270.1
unripe pulp	1293	1312	2.0	1315.8	1307.9
ripe pulp	1285	1311	2.7	1316.5	1306.1

Table 4: Results from PERMANOVAs showing a large percentage of sample-to-sample variance in composition within organ types (i.e. beta diversity) is explained by species

	$F_{11,35}$[b]	P[b]	η^2 (Species)[c]	η^2 (Residual)[d]
Compound Occurrences [a]				
leaf	6.29	0.001	0.74	0.26
seed	5.17	0.001	0.70	0.30
unripe pulp	12.46	0.001	0.86	0.14
ripe pulp	13.18	0.001	0.86	0.14

Structural Composition [a]

leaf	4.51	0.001	0.67	0.33
seed	6.58	0.001	0.75	0.25
unripe pulp	7.90	0.001	0.79	0.21
ripe pulp	10.70	0.001	0.83	0.17

[a] Separate sets of PERMANOVAs were conducted for each aspect of compound composition: compound occurrences (presence/absence) and structural composition

[b] Statistical results from permutation tests showing strong support for an effect of Piper species on composition

[c] Proportion of sample-to-sample variance explained by species (i.e. interspecific variation)

[d] Proportion of sample-to-sample variance explained by individual and within-individual (residual) variance

(A)

Chemical class

- Benzene and substituted derivatives
- Carboxylic acids and derivatives
- Flavonoids
- Organooxygen compounds
- Prenol lipids
- Other
- Unknown

(B)

Compound presence by tissue type

- Fruit
- Leaf
- Fruit & Leaf