1 **Regulation of blood–brain barrier integrity and cognition by the microbiome-**
2 **associated methylamines trimethylamine-*N*-oxide and trimethylamine**
3

4 Lesley Hoyles[1]*, Matthew G. Pontifex[2], Ildefonso Rodriguez-Ramiro[2,3], M. Areeb Anis-
5 Alavi[4], Tom Snelling[5], Egle Solito[6,7], Sonia Fonseca[8], Ana L. Carvalho[8], Simon R.
6 Carding[2,8], Michael Müller[2], Robert C. Glen[5,9] David Vauzour[2] & Simon McArthur[4]*
7

[1]Department of Biosciences, School of Science and Technology, Nottingham Trent
University, Clifton, Nottingham, UK
[2]Norwich Medical School, University of East Anglia, Norwich, UK
[3]Metabolic Syndrome Group, Madrid Institute for Advanced Studies (IMDEA) in Food,
Madrid, E28049, Spain
[4]Institute of Dentistry, Barts & the London School of Medicine & Dentistry, Blizard
Institute, Queen Mary University of London, London, UK
[5]Faculty of Medicine, Department of Metabolism, Digestion and Reproduction,
Imperial College London, London, UK
[6]William Harvey Research Institute, Barts & the London School of Medicine &
Dentistry, Queen Mary, University of London, London, UK
[7]Dipartimento di Medicina molecolare e Biotecnologie mediche, Federico II University,
Naples, Italy
[8]The Gut Microbes and Health Research Programme, The Quadram Institute, Norwich
Research Park, Norwich, UK
[9]Centre for Molecular Informatics, Department of Chemistry, University of Cambridge,
Cambridge, UK
25

*Corresponding authors: Lesley Hoyles, lesley.hoyles@ntu.ac.uk; Simon McArthur,
s.mcarthur@qmul.ac.uk

Running title: Dietary methylamines affect the blood–brain barrier
Abbreviations: BBB, blood–brain barrier; DI, discrimination index; GO, gene
ontology; KEGG, Kyoto Encyclopedia of Genes and Genomes; LPS,
lipopolysaccharide; NOR, novel object recognition; SCFA, short-chain fatty acid; OFT,
open field test; SPIA, signalling pathway impact analysis; TEER, transendothelial
electrical resistance; TMA, trimethylamine; TMAO, trimethylamine *N*-oxide.

Supplementary materials associated with the article are available from figshare

(https://doi.org/10.6084/m9.figshare.13549334.v1).

ABSTRACT

Communication between the gut microbiota and the brain is primarily mediated *via* soluble microbe-derived metabolites, but the details of this pathway remain poorly defined. Methylamines produced by microbial metabolism of dietary choline and L-carnitine have received attention due to their proposed association with vascular disease, but their effects upon the cerebrovascular circulation have not hitherto been studied. Here we use an integrated *in vitro*/*in vivo* approach to show that physiologically relevant concentrations of the dietary methylamine trimethylamine *N*-oxide (TMAO) enhanced and protected blood-brain barrier (BBB) integrity, acting through the tight junction regulator annexin A1. In contrast, the TMAO precursor trimethylamine (TMA) impaired BBB function and disrupted tight junction integrity. Moreover, we show that long-term exposure to TMAO has beneficial effects upon cognition in mice, improving visual recognition memory. Our findings demonstrate a direct interaction of microbiome-associated metabolites with the mammalian BBB, with consequences for cerebrovascular and cognitive function.

INTRODUCTION

As the role of the gut microbiota in host physiology and disease is categorised, novel pathways through which these interactions are mediated continue to emerge. We and others recently identified the blood–brain barrier (BBB) as a target for gut microbe-derived short-chain fatty acid (SCFA) activity, with butyrate and propionate acting to promote BBB integrity and protect the cerebral vasculature from insult[1,2]. SCFAs represent just one of many classes of gut microbe-derived metabolites, with little known as to how these other classes may influence BBB function.

Dietary methylamines, such as choline, phosphatidylcholine, betaine and trimethylamine-*N*-oxide (TMAO), are a class of metabolites receiving considerable attention as modulators of vascular function[3,4], although the mechanism(s) by which they affect human physiology remain poorly understood. The aforementioned methylamines can be broken down by members of the gut microbiota into trimethylamine (TMA)[5], which is carried from the gut through the portal vasculature to the liver and rapidly converted into TMAO by flavin monooxygenases[6]. TMAO then enters the systemic circulation, reaching fasting plasma concentrations of between 2 and 40 μM in humans[7–9], prior to excretion through the urine[5]. Approximately ten-fold lower concentrations of TMA compared with TMAO are found in the circulation under normal physiological conditions.

Early observational work reported an apparent association between atherosclerosis and elevated levels of TMAO[10,11]. Similarly, pre-clinical studies demonstrate the damaging effects of supraphysiological TMAO doses in atherosclerosis-prone mice[12] and upon thrombus formation[13]. Despite this, the impact of TMAO upon the vasculature remains uncertain, with a number of detailed studies encompassing both human and murine systems failing to replicate these initial findings[14], instead suggesting that this negative relationship disappears upon correction for renal function[4,15–17]. This indicates that the raised TMAO levels may in fact reflect impaired excretion rather than being a causative factor of disease. Moreover, protective roles for TMAO have been reported in rodent models of hypertension[18], atherosclerosis[19] and non-alcoholic steatohepatitis[20] and we have previously shown TMAO to improve glucose homeostasis and insulin secretion in mice fed a high-fat diet[21]. Perhaps helping to clarify this apparent contradiction, recent studies have established that

89 intravenous treatment of rats with the TMAO precursor TMA, but not TMAO itself,
90 increases mean arterial blood pressure[22]. Notably, the majority of reports describing
91 associations of plasma TMAO with cardiovascular disease have not concurrently
92 monitored levels of TMA; TMA but not TMAO has been shown to associate with severe
93 aortic stenosis[22] and gestational diabetes risk[23].
94
Beyond vascular health, dietary methylamines have implications for cognition, with a
positive correlation observed between choline intake and cognitive function in both
humans[24,25] and mice[26,27]. In contrast, cerebrospinal fluid TMAO levels have been
indicated as predictive of cognitive decline in Alzheimer's disease[28], while suppression
of microbial TMA/TMAO production improves cognitive function in the murine
APP/PS1 model of Alzheimer's disease[29]. Given the disparities in the literature
regarding the effects of methylamines upon the vasculature, and our increasing
awareness of the BBB as a major actor in the pathology of multiple neurological
conditions, we investigated the effects of physiologically relevant concentrations of
TMAO and its precursor TMA upon BBB integrity and cognitive behaviour.
105

METHODS

Endothelial cell culture

The human cerebromicrovascular endothelial cell line hCMEC/D3 was maintained and treated as described previously[2,30]. Cells bearing shRNA sequences targeting annexin A1 (*ANXA1*) or non-specific scramble sequences were produced as described previously[31]; the degree of *ANXA1* knock-down was confirmed by flow cytometry analysis (Suppl. Fig. 1). For all lines, cells were cultured to confluency in complete EBM-2MV microvascular endothelial cell growth medium (Promocell GmbH, Heidelberg, Germany), whereupon medium was replaced by EBM-2MV without VEGF and cells were further cultured for a minimum of 4 days to enable intercellular tight junction formation prior to experimentation.

In vitro barrier function assessments

Paracellular permeability and transendothelial electrical resistance (TEER) were measured on 100 % confluent hCMEC/D3 cultures polarised by growth on 24-well plate polyethylene terephthalate (PET) transwell inserts (surface area: 0.33 cm^2, pore size: 0.4 µm; Greiner Bio-One GmbH, Kremsmünster, Austria) coated with calf-skin collagen and fibronectin (Sigma-Aldrich, UK). The permeability of hCMEC/D3 cell monolayers to 70 kDa FITC-dextran (2 mg/ml) was measured as described previously[31–33]. TEER measurements were performed using a Millicell ERS-2 Voltohmmeter (Millipore, Watford, UK) and were expressed as Ω.cm^2. In all cases, values obtained from cell-free inserts similarly coated with collagen and fibronectin were subtracted from the total values. In some cases, barrier integrity was tested by challenge with bacterial lipopolysaccharide (LPS). Confluent hCMEC/D3 monolayers were treated with TMAO or TMA for 12 h, whereupon LPS (*Escherichia coli* O111:B4; 50 ng/ml, comparable to circulating levels of LPS in human endotoxemia[34]) was added for a further 12 h, without wash-out. Barrier function characteristics were then interrogated as described above.

Cell adhesion assays

hCMEC/D3 cells were cultured to confluency on transwell inserts (0.4 µm pore size, 0.33 cm^2 diameter, Greiner Bio-One Gmbh, Austria) prior to 16 h treatment with 10 ng/ml TNFα. Monolayers were then incubated for 2 h with U937 monocytic cells pre-

labelled according to manufacturer's instructions with CMFDA cell tracker dye
(ThermoFisher Scientific, UK). Co-cultures were washed vigorously with ice-cold PBS
three times and fixed by incubation for 10 min in 1 % formaldehyde in 0.1 M PBS. Co-
cultures were mounted and examined using an Axiovert 200M inverted microscope
(Zeiss) equipped with a 20x objective lens. Images were captured with ZEN imaging
software (Carl Zeiss Ltd, UK) and analysed using ImageJ 1.53c (National Institutes of
Health, USA).

Microarrays
hCMEC/D3 cells were grown on 6-well plates coated with calf-skin collagen (Sigma-
Aldrich, Gillingham, UK), and collected in TRIzol (Thermo-Fisher Scientific, UK) as
described previously[2]. Total RNA was extracted using a TRIzol Plus RNA purification
kit (Thermo-Fisher Scientific, UK) and quantified using a CLARIOstar
spectrophotometer equipped with an LVis microplate (BMG Labtech GmbH,
Germany).

Hybridization experiments were performed by Macrogen Inc. (Seoul, Republic of
Korea) using Illumina HumanHT-12 v4.0 Expression BeadChips (Illumina Inc., San
Diego, CA) as described previously[2].

Processing and analyses of array data
Raw data supplied by Macrogen were quality-checked, $\log_2$-transformed and loess-
normalized (2 iterations) using affy[35]. Probe filtering and matching of probes not
classified as 'Bad' or 'No match' to Entrez identifiers were done as described
previously[2]. Average gene expression values were used for identification of
differentially expressed genes. Array data have been deposited in ArrayExpress under
accession number E-MTAB-6662. Normalized data are available (Supplementary
Table 1).

Enrichr[36,37] was used to perform Gene Ontology (GO) analysis. Signaling Pathway
Impact Analysis (SPIA) was used to determine whether Kyoto Encyclopedia of Genes
and Genomes (KEGG) pathways were activated or inhibited in hCMEC/D3 cells
exposed to TMAO or TMA[38]. Human KEGG pathways (KGML format) downloaded

from the KEGG PATHWAY database[39] were used for network (KEGGgraph, RBGL[40])

analysis.

Immunofluorescence microscopy

hCMEC/D3 cells were cultured on 24-well plate polyethylene terephthalate (PET)

transwell inserts (surface area: 0.33 cm^2, pore size: 0.4 µm; Greiner Bio-One GmbH,

Kremsmünster, Austria) coated with calf-skin collagen and fibronectin (Sigma-Aldrich,

UK), prior to immunostaining according to standard protocols[2,31] and using a primary

antibody directed against zonula occludens-1 (ZO-1; 1:100, ThermoFisher Scientific,

UK) or Alexafluor 488-conjugated phalloidin (1:140, ThermoFisher Scientific, UK).

Nuclei were counterstained with DAPI (Sigma-Aldrich, UK). Images were captured

using an LSM880 confocal laser scanning microscope (Carl Zeiss Ltd, Cambridge,

UK) fitted with 405 nm and 488 nm lasers, and a 63x oil immersion objective lens (NA,

1.4 mm, working distance, 0.17 mm). Images were captured with ZEN imaging

software (Carl Zeiss Ltd, UK) and analysed using ImageJ 1.53c (National Institutes of

Health, USA).

Flow cytometry analysis

Following experimental treatment, hCMEC/D3 cells were detached using 0.05%

trypsin and incubated with an unconjugated rabbit polyclonal antibody directed against

ANXA1 (1:1000, ThermoFisher Scientific, UK) on ice for 30 minutes, followed by

incubation with an AF488-conjugated goat anti-rabbit secondary antibody (1:500,

ThermoFisher Scientific, UK). Similarly detached hCMEC/D3 cells were incubated

with APC-conjugated mouse monoclonal anti-BCRP (1:100, BD Biosciences, Oxford,

UK), or PE-conjugated mouse monoclonal anti-MDR1A (1:100, BD Biosciences, UK)

antibodies on ice for 30 minutes, alongside fluorescence minus one controls.

Immunofluorescence was analysed for 20,000 events per treatment using a BD

FACSCanto II (BD Biosciences, UK) flow cytometer; data were analysed using FlowJo

8.0 software (Treestar Inc., CA, USA).

Efflux transporter assays

Activity of the major efflux transporters P-glycoprotein and BCRP was determined

through the use of commercially available assays (PREDEASY™ ATPase Assay Kits,

Solvo Biotechnology Inc., Budapest, Hungary), performed according to the

manufacturer's instructions. Stepwise dose–response curves centred around reported

physiological circulating concentrations of TMA (4.9 nM – 10.8 µM) and TMAO (0.5

µM – 1.08 mM) were constructed ($n = 4$) to investigate inhibitory effects of the

methylamines upon transporter activity.

ELISA

Culture medium ANXA1 content was assayed by specific ELISA as described

previously[41]. Serum TNFα and IL-1β concentrations were assayed using commercial

ELISA kits according to the manufacturer's instructions (ThermoFisher Scientific, UK).

Animal experiments

All animal experiments were performed according to the UK Animals (Scientific

Procedures) Act of 1986, under UK Home Office Project Licences PFA5C4F4F (short

term studies) and 70/8710 (long term studies), following ethical review by the Animal

Welfare and Ethical Review Boards of Queen Mary, University of London or the

University of East Anglia, respectively. Wild-type male C57Bl/6J mice (Charles River

Ltd., Harlow, UK) aged 8 weeks at the start of procedures were used throughout, with

a group size of n=6 for short term studies and n=8 for long-term/behavioural analyses.

Animals were housed in individually ventilated cages on a daily 12 h:12 h light/dark

cycle with, unless otherwise indicated, *ad libitum* access to standard mouse chow and

drinking water. Experimental procedures were started at 9 am to minimise variation

associated with circadian rhythms.

Assessment of acute effects of TMAO on BBB integrity

Mice (n=4-6 per group) were injected i.p. with 1.8 mg/kg body weight TMAO in 100 µl

saline vehicle, a dose calculated to approximate human circulating TMAO levels[42],

followed 2 h, 6 h or 24 h later by assessment of Evans blue extravasation as described

below. Alternatively, mice were injected i.p. with 3 mg/kg body weight LPS or 100 µl

0.9% saline vehicle, followed 2 h later by i.p. injection of either 1.8 mg/kg body weight

TMAO or 100 µl 0.9% saline vehicle for assessment of Evans blue extravasation 2 h

later. In both experiments, one hour before assessment animals were injected i.p. with

100 µl of a 2% (w/v) solution of Evans blue dye in 0.9 % saline (Sigma–Aldrich Ltd,

Poole, UK). Dye was permitted to circulate for 1 h before animals were transcardially

perfused with 0.9% saline at 4 °C to remove circulating dye. Brains were removed,

bisected and homogenized by maceration in 0.9% saline. Suspended macromolecules

were precipitated by incubation with 60% trichloroacetic acid, and dye content of

resulting supernatants was detected using a CLARIOstar spectrophotometer (BMG

Labtech GmbH, Germany) alongside a standard curve of defined concentrations of

Evans blue in the same buffer. Brain Evan's blue content was expressed as µg of dye

per mg of brain tissue, normalized to circulating plasma concentrations.

Long-term LPS and TMAO treatments

To assess the long-term impact of both LPS and TMAO on cognitive performance,

mice were divided into four groups (n=8 per group): 1) Water + PBS; 2) Water + TMAO;

3) LPS + PBS; 4) LPS + TMAO. C57Bl/6 mice were administered phosphate-buffered

saline (PBS) or LPS (*Escherichia coli* O55:B5, Sigma-Aldrich, UK) via intraperitoneal

(i.p.) injection (0.5 mg/kg/wk) for 8 weeks[43]. A final LPS treatment was administered

the day before sacrifice for nine total injections. Body weights were recorded prior to

each injection. TMAO was provided in the drinking water (500 mg/L) and water bottles

replaced every other day. Drinking volumes were recorded before bottle change.

Processing and analyses of RNAseq data

Mice were transcardially perfused with 0.9% saline at 4 °C to remove circulating blood,

and brains were removed and collected into RNAlater (Thermofisher Scientific Ltd.,

UK) prior to storage at -20 °C for later analysis. Whole brain total RNA was extracted

using a PureLink RNA Mini Kit (Thermofisher Scientific Ltd., UK) and quantified using

a CLARIOstar spectrophotometer equipped with an LVis microplate (BMG Labtech

GmbH, Germany). RNA samples (*n*=3 TMAO, *n*=3 control) were sent to Macrogen Inc.

(Republic of Korea) where they were subject to quality checks (RIN analysis); libraries

were prepared (TruSeq Stranded mRNA LT Sample Prep Kit) for paired-end (2x 100

nt) sequencing on an Illumina HiSeq 4000 apparatus. Raw RNAseq sequence data

(delivered in fastq format) were processed in house as follows. Reads were mapped

onto the mouse genome (mm10) using HISAT2 v2.1.0[44]. Number of reads in each

sample that mapped to genes in the BAM files returned by HISAT2 was determined

using featureCounts v1.6.4[45]. Entrez gene identifiers were converted to gene symbols

using *Mus musculus* annotations downloaded from NCBI on 26 November 2020; only

those genes with valid Entrez gene identifiers were retained in analyses. Raw RNAseq

data have been deposited with ArrayExpress under accession number E-MTAB-9869.

Significantly differentially expressed genes ($P<0.1$) were analysed by mouse KEGG

pathway over-representation analysis using Enrichr and manual curation.

Behavioural analyses

Behavioural tests were performed in the order they are introduced below. Apparatus

was cleaned using 70 % ethanol upon completion of each trial, eliminating any residual

odour.

Open field test (OFT) was conducted as previously described[46]. Briefly, mice were

placed in the centre of the OFT, a grey 50 x 50 x 50 cm apparatus illuminated with low

lux (100 lux) lighting. Total travel distance and time spent in the centre of the field was

determined at 5 min with a video tracking system (Smart 3.0 tracking software, Panlab,

Kent, UK).

The novel object recognition (NOR), a measure of recognition memory, was performed

as described previously[47,48], with slight modifications. Briefly, on day 1 mice were

habituated in a grey 50 x 50 x 50 cm apparatus illuminated with low lux (100 lux)

lighting, mice were placed into the empty maze and allowed to move freely for 10 min.

On day 2, mice were conditioned to a single object for a 10 min period. On day 3, mice

were placed into the same experimental area in the presence of two identical objects

for 15 min, after which they were returned to their respective cages and an inter-trial

interval of 1 h was observed. One familiar object was replaced with a novel object.

Mice were placed back within the testing area for a final 10 min. Videos were analysed

for a 5 min period, after which if an accumulative total of 15 s with both objects failed

to be reached, analysis continued for the full 10 min or until 15 s was achieved. Those

not achieving 15 s were excluded from the analysis[49]. A discrimination index (DI) was

calculated as follows: DI = (TN−TF)/(TN+TF), where TN is the time spent exploring

the novel object and TF is the time spent exploring the familiar object.

Y-maze spontaneous alternation test, a measure of spatial working memory, was

performed on the final day of behavioural testing as previously described[50]. Briefly, the

Y-maze apparatus comprised white Plexiglas (dimensions 38.5 × 8 × 13 cm, spaced

120° apart) and was illuminated with low lux (100 lux) lighting. Mice were placed in the

maze and allowed to explore freely for 7 min while tracking software recorded zone

transitioning and locomotor activity (Smart 3.0 tracking software, Panlab, Kent, UK).

Spontaneous alternation was calculated using the following formula: Spontaneous

Alternation = (Number of alternations/ Total Arm entries - 2) x 100.

Extravasation assay and sample processing following long-term treatment

Twenty-four hours after the final injection of LPS, mice were injected i.p. with 200 µl of

2% sodium fluorescein in sterile ddH$_2$O and anesthetized 30 min later with isoflurane

(1.5%) in a mixture of nitrous oxide (70%), and oxygen (30%). Once sedated, blood

was collected by cardiac puncture and centrifuged at 1,500 × g for 15 min at 4 °C to

collect the serum. The samples were analysed immediately for sodium fluorescein

extravasation or snap-frozen in liquid nitrogen and stored at -80 °C until further

analysis.

Mice were then transcardially perfused with saline containing 10 kU/ml heparin

(Sigma, Devon, UK). Dissected left hemi-brains were fixed in 4% PFA for 24 h and

embedded into paraffin before being processed for immunohistochemical analysis.

Right hemi-brains were stored at -80 °C until further analysis; cerebellums were

processed immediately for the sodium fluorescein extravasation assay. Cleared

volume of sodium fluorescein that passed from the plasma into the brain was

calculated as described previously[43].

Statistical analyses

Sample sizes were calculated to detect differences of 15 % or more with a power of

0.85 and α set at 5 %, calculations being informed by previously published data[2,31]. *In*

vitro experimental data (except those for *in vitro* microarray experiments) are

expressed as mean ± SEM, with a minimum of *n* = 3 independent experiments

performed in triplicate for all studies. In all cases, normality of distribution was

established using the Shapiro–Wilks test, followed by analysis with two-tailed

Student's *t*-tests to compare two groups or, for multiple comparison analysis, 1- or 2-

way ANOVA followed by Tukey's HSD *post hoc* test. Where data were not normally

distributed, non-parametric analysis was performed using the Wilcoxon signed rank

test. A *P* value of less than or equal to 5 % was considered significant. Differentially

expressed genes were identified in microarray data using LIMMA (Ritchie *et al*, 2015);

P values were corrected for multiple testing using the Benjamini–Hochberg procedure

(False Discovery Rate); a *P* value of less than or equal to 10 % was considered

significant in this case; *n* = 5 for control, TMAO and TMA groups. Significantly

differentially expressed genes (P_{FDR}<0.1) in RNAseq data (Supplementary Table 11)

were identified using DESeq2 v1.22.1[51].

RESULTS

To provide an initial assessment of the effects of the methylamines TMA and TMAO upon the BBB we used a well-established *in vitro* BBB model, hCMEC/D3 immortalised human cerebromicrovascular cell monolayers grown under polarising conditions on a Transwell filter, examining two key barrier properties: paracellular permeability to a protein-sized tracer and TEER. Exposure of hCMEC/D3 cells for 24 h to TMA (0-40 µM) caused a clear dose-dependent increase in paracellular permeability to 70 kDa FITC-dextran (Fig. 1A), with normal circulating levels (0.4 µM) of TMA and upwards significantly enhancing permeability. In contrast, exposure for 24 h to TMAO (0-4000 µM) caused a biphasic dose-dependent response (Fig. 1A), with normal circulating concentrations (40 µM) significantly reducing permeability to the tracer, an effect lost at 10-fold greater TMAO concentrations and reversed at 100-fold greater TMAO (4 mM), where a significant increase in paracellular permeability was apparent. In contrast, TMA had no effect upon TEER at any concentration studied, while TMAO enhanced TEER by approximately 65%, an effect that was notably dose-independent (Fig. 1B).

The physical barrier that the BBB provides is only one aspect by which it separates the brain parenchymal environment from the periphery, equally important is the immunological barrier that it represents. To model this, we employed a simple system in which adhesion of CMFDA-labelled U937 monocytic cells to TNFα-activated (10 ng/ml, 16 h) hCMEC/D3 monolayers was quantified in response to TMA or TMAO treatment. Treatment with a physiologically relevant concentration of TMA (0.4 µM[42], 24 h post-TNFα) had no effect on the density of adherent U937 cells, but exposure of hCMEC/D3 monolayers to physiological levels of TMAO (40 µM[42], 24 h post-TNFα) significantly reduced U937 cell adhesion by approximately 50% compared to cultures stimulated with TNFα alone (Fig. 1C).

The endothelial cells of the BBB express numerous efflux transporter proteins that serve to limit entry of endogenous and exogenous molecules into the parenchyma, with BCRP and P-glycoprotein being two of the most important. Consequently, we examined whether treatment with TMA or TMAO affected function or expression of either of these two transporters. Using commercially available *in vitro* assays, neither

381 methylamine affected BCRP or P-glycoprotein activity across a wide concentration
382 range (TMA: 4.9 nM to 10.8 µM; TMAO 0.5 µM to 1.08 mM) (Suppl. Fig. 2A-D).
383 Similarly, treatment of hCMEC/D3 cells for 24 h with physiologically relevant
384 concentrations of TMA (0.4 µM) or TMAO (40 µM) was without effect on cell surface
385 expression of either BCRP or P-glycoprotein (Suppl. Fig. 2E-F).
386
Methylamine-induced changes in gene expression
Having identified significant TMA-/TMAO-induced functional changes in endothelial
barrier characteristics *in vitro*, we undertook a microarray analysis of hCMEC/D3 cells
treated with either TMA (0.4 µM, 24 h) or TMAO (40 µM, 24 h) to investigate the
transcriptional changes underlying these effects. Treatment with TMA had a significant
(P_{FDR}<0.1) effect on 49 genes, with the expression of 39 upregulated and 10
downregulated (Fig 2A, Supplementary Table 2). In contrast, treatment with TMAO
had a significant (P_{FDR}<0.1) effect on 440 genes with 341 upregulated and 99
downregulated (Fig. 2B, Supplementary Table 3). *FMO3* gene expression was not
affected by TMA or TMAO at the physiological concentrations employed (Suppl. Fig.
3).
398
399 SPIA of the 440 TMAO-affected genes showed activation of the tight junction pathway
400 (P = 0.031), but significance was lost after correction for multiple testing
401 (Supplementary Table 4). No pathways were shown to be activated or inactivated by
402 the 49 TMA-affected genes (data not shown).
403
Gene ontology (GO) analysis was performed on TMA- and TMAO-regulated genes
using Enrichr[36,37]. TMA up-regulated and down-regulated genes were significantly
(P_{FDR}<0.2) associated with processes indicative of a degree of cellular stress (Fig 2C,
Supplementary Table 5, Supplementary Table 6). In contrast, genes up-regulated by
TMAO treatment were associated with regulation of the cytoskeleton and cell
morphology and with actin bundle formation (P_{FDR}<0.2), whereas pathways associated
with inflammatory signalling were down-regulated (Fig 2D, Supplementary Table 7,
Supplementary Table 8).
412
We then assessed the topology of a directional network of the 440 TMAO-associated
genes mapped onto all human KEGG pathways. In line with the GO analysis described

above, a number of genes of differing function were regulated by TMAO treatment,
with two principal groupings being particularly evident, namely those associated with
aspects of cellular metabolism and with regulation of actin cytoskeletal dynamics
(Figure 2E). Finally, we compared the 19,309 genes represented on the microarray
with a set of 203 genes[2] known to be associated with the BBB. While TMA treatment
had no significant effects on expression of these genes (Supplementary Table 9),
TMAO significantly ($P_{FDR}<0.1$) upregulated expression of four genes from this set
associated with transporter proteins and barrier integrity (Table 1, Supplementary
Table 10).

Given these transcriptional indications, and the fact that the restrictive properties of
the BBB are largely governed by inter-endothelial cell tight junctions linked *via* the
zonula occludens complex to the actin cytoskeleton[52], we hypothesised that TMA and
TMAO may affect barrier permeability through modification of the links between tight
junctions and the actin cytoskeleton. Confocal immunofluorescence microscopy of
hCMEC/D3 monolayers treated with a physiologically relevant concentration of TMA
(0.4 µM, 24 h) or TMAO (40 µM, 24 h) revealed clear changes to both ZO-1 and
fibrillar actin disposition within cells (Fig. 2F). Compared to untreated cells in which
both ZO-1 and F-actin fibres clearly defined the cellular perimeter, cells treated with
TMA exhibited a broken, patchy distribution of perimeter ZO-1 expression, and the
appearance of marked cytoplasmic F-actin stress fibres. In contrast, cells treated with
TMAO showed little change in ZO-1 distribution, but a marked enhancement of cortical
F-actin fibre thickness and intensity.

The actions of TMAO are mediated through annexin A1 signalling
Of the four BBB-associated genes identified as upregulated by TMAO, *ANXA1* is of
particular interest as we have previously shown this protein to regulate BBB tightness
in vitro and *in vivo* through modulation of the actin cytoskeleton[53]. Examination of
ANXA1 expression in hCMEC/D3 cells revealed that while total cellular levels of the
protein were not changed by either TMA (0.4 µM, 24 h) or TMAO (40 µM, 24 h)
treatment (Fig. 3A), TMA significantly suppressed and TMAO significantly augmented
medium ANXA1 content (Fig. 3B), a finding of interest given that autocrine/paracrine
effects are a major route of ANXA1 action[54].

To establish the importance of ANXA1 in mediating the effects of TMAO, we
investigated the effects of its depletion through use of hCMEC/D3 clones stably
transfected with shRNA sequences targeting *ANXA1* mRNA (Suppl. Fig. 1).
Suppression of *ANXA1* expression significantly inhibited the effects of TMAO (40 μM,
24 h) upon both paracellular permeability and TEER (Fig. 3C-D), an effect not seen in
cells bearing non-targeting scramble shRNA sequences. The actions of ANXA1 are
mediated to a large extent through the G protein-coupled receptor formyl peptide
receptor 2 (FPR2)[55]. Hence, we investigated how inclusion of a well-characterised
antagonist to this receptor, WRW$_4$ (10 μM, 10 min pre-treatment), would affect the
functional response to TMAO. Pre-treatment with WRW$_4$ was able to significantly
attenuate the effects of TMAO treatment on both TEER (Fig. 3E) and paracellular
permeability (Fig. 3F), further indicating the role of ANXA1 signalling as the principal
mediator of TMAO actions on hCMEC/D3 cells.

Acute beneficial effects of TMAO treatment in vivo
While hCMEC/D3 endothelial cells are a widely used and generally representative
model, they cannot reflect all aspects of the multicellular neurovascular unit that
underlies BBB function, hence we investigated whether the beneficial effects of TMAO
identified *in vitro* translate to an *in vivo* situation. Initial studies revealed that systemic
administration of TMAO to wild-type male mice (1.8 mg/kg, i.p.) induced a time-
dependent reduction in BBB permeability to the tracer Evans blue (2 % in saline, 100
μl, i.p.), with a significant reduction in dye extravasation to the brain parenchyma being
apparent 2 h following TMAO administration, an effect lost at longer time-points (Fig.
4A), presumably due to the relatively short plasma half-life of TMAO *in vivo*[56]. To
further investigate this effect of TMAO, we employed a simple model of enhanced BBB
permeability, namely acute peripheral administration of bacterial LPS[31]. Treatment
with LPS (*E. coli* O111:B4, 3 mg/kg, i.p.) significantly enhanced intraparenchymal
extravasation of Evans blue within 4 h, an effect significantly attenuated by subsequent
treatment with TMAO (1.8 mg/kg, i.p.) 2 h post-LPS (Fig. 4B), further confirming a
beneficial action of TMAO at physiological concentrations upon the BBB *in vivo*.

TMAO treatment rapidly alters brain transcriptional activity

To investigate the wider actions of TMAO upon the brain we performed whole brain RNAseq transcriptomic analysis of wild-type male mice 2 h following TMAO administration (1.8 mg/kg i.p.). We identified 76 significantly differentially expressed genes (P_{FDR}<0.1), with expression of 41 upregulated and 35 downregulated (Figure 5A; Supplementary Table 11). KEGG pathway analysis using Enrichr identified a number of significantly regulated murine pathways (Figure 5B), including oxidative phosphorylation, Parkinson's disease and Alzheimer's disease. Closer analysis of regulated genes identified a number of general groupings (Figure 5C), with downregulated genes associated with the mitochondrial respiratory chain (*COX1*, *COX3*, *ATP6*, *ND4L*, *CYTB*, *ND1*, *ND3*, *ND4*, *ND6*) and ribosomal function (*mt-Rnr2*, *mt-Rnr1*, *Rps23rg1*) and upregulated genes associated with cellular or axonal growth (*Nme7*, *B3gat2*, *Fuz*, *Nefm*, *Basp1*, *Mtg1*, *Vps37a*, *Smim1*, *Araf*). Of the 203 BBB-associated human genes previously identified[2], 197 had matches in our mouse brain data set. Here, two genes were identified as significantly differentially expressed at P_{FDR}<0.1: reduced *Cpe* (carboxypeptidase E) and increased *App* (amyloid precursor protein) expression (Figure 5D; Supplementary Table 12).

Chronic low-dose TMAO treatment prevents LPS-induced BBB disruption and memory impairment

The fundamental role of the BBB is to protect the brain, preserving its homeostatic environment; damage to BBB integrity is therefore detrimental, and is believed to directly contribute towards cognitive impairment[57–59]. Having shown TMAO to exert a protective effect upon BBB function/integrity in response to acute inflammatory insult, we next examined whether a similar effect held true for chronic conditions, and whether this protection extended to cognition. TMAO was administered to male C57Bl/6J mice through drinking water (0.5 mg/ml) over 2 months, in combination with chronic low-dose LPS administration (0.5 mg/kg/week, i.p.) to model a mild inflammatory stress known to impact cognitive behaviour[43]. There were no differences in volumes of water drunk or, where relevant, final consumption of TMAO between any groups (Table 2). The serum inflammatory cytokines TNFα and IL-1β were both nominally elevated in response to LPS treatment, although not reaching statistical significance, indicating a sub-clinical inflammatory response; TMAO had no effect on TNFα nor IL-1β levels (Suppl. Fig. 4). Notably, animals exposed to LPS exhibited a

significant reduction in body weight gain compared to their untreated counterparts, an effect reversed by TMAO treatment (Fig. 6A). Treatment with LPS increased cerebellar FITC extravasation, an effect that was prevented by TMAO treatment, although this did not reach statistical significance on *post hoc* analysis (Fig. 6B). To corroborate these findings, we investigated a second marker of impaired BBB integrity, confocal microscopic detection of brain perivascular IgG deposition. In comparison with sham-treated animals, exposure to LPS caused a significant accumulation of IgG in the perivascular compartment, an effect prevented by TMAO treatment (Fig. 6C).

The OFT confirmed neither LPS nor TMAO treatment affected motor function, with movement speed and distance travelled comparable across treatment groups (Fig. 6D-E). Similarly, no effect was apparent on the proportion of time animals spent in the centre of the field, suggesting limited effects upon anxiety (Fig. 6F). Working memory, however, determined via NOR indicated a significant reduction in performance in animals exposed to LPS, a behavioural deficit notably prevented in animals co-treated with TMAO (Fig. 6G). In contrast, no effect of either LPS or TMAO treatment was apparent in the Y-maze spontaneous alternation task (Fig. 6H) or in distance travelled during this task (Fig. 6I), indicating no differences in spatial memory.

DISCUSSION

The relationship between the BBB and cognitive behaviour is complex and far from being fully understood, but it is clear from both human and animal studies that deficits in barrier integrity can exert a profound and deleterious effect upon memory, language and executive function[60–63]. Indeed, BBB impairment is among the first events to occur in the course of Alzheimer's disease, and may aggravate the pathological processes that underlie the condition[64]. Strategies to promote BBB function may thus have significant value in helping to protect the brain from progressive neurological diseases such as dementia. In this study we identify novel and distinct roles for the microbiome-associated dietary methylamines TMA and TMAO in regulating BBB function *in vitro* and *in vivo* and provide evidence that the beneficial action of TMAO upon the BBB coincides with similar protective effects upon cognition. These data reinforce the position of the cerebral vasculature as a major target for the gut-brain axis, and extend our knowledge of its interactions with microbial metabolites beyond SCFAs[1,2] to another major class of molecules, the dietary methylamines.

Notably, our data show that while both TMA and TMAO have activity upon the endothelium, there is a marked distinction between their effects despite their close structural similarity. TMA, the direct product of microbial choline, L-carnitine and TMAO metabolism in the upper gut[5], had a deleterious effect upon the endothelium, disrupting cytoskeletal arrangement, inducing signs of metabolic stress and ultimately impairing endothelial barrier integrity. In contrast, TMAO, largely derived from hepatic FMO3-mediated oxidation of TMA taken up from the gut via the hepatic portal vein[6], was beneficial for cerebral vascular integrity *in vitro* and *in vivo*. These differences suggest that host conversion of TMA (a gas) to TMAO (a stable metabolite) may be an effective detoxification pathway, emphasising the importance of host metabolic pathways in modulating communication in the gut-brain axis, and underlining the importance of using a systems-level approach to understand the interactions between the host and its resident microbiota.

TMAO and cognitive function

Numerous groups have investigated the putative relationship between TMAO and cognition following reports of an association between cerebrospinal fluid TMAO

content and Alzheimer's disease[28], with negative correlations between plasma TMAO content and cognitive function having been identified in both clinical[65–67] and experimental[29,68,69] settings. Whether this relationship is truly deterministic remains unclear, however, as the role of the immediate precursor to TMAO, TMA, in cognition and vascular function has largely been overlooked. This omission may be important in light of studies reporting negative correlations between cognitive impairment and serum TMA[70–72] and our own data showing a potent detrimental effect of physiological levels of TMA upon the cerebrovascular endothelium *in vitro*. Given that TMA has been shown to be detrimental in contexts other than cognitive function[22,23], the contribution that this metabolite plays in disease is evidently in need of closer attention.

Interpreting associations between circulating TMAO and cognition is further complicated by studies indicating that consumption of the TMAO precursors choline and L-carnitine can improve cognitive function[24,25,73], evidence that patients with Parkinson's disease have lower circulating TMAO than healthy controls[74], and more recent Mendelian randomisation analysis indicating that serum TMAO and Alzheimer's disease are not causally related[75]. Given this background, our data indicating that physiologically relevant concentrations of TMAO have positive effects upon both BBB integrity and cognition *in vivo* thus serve as a useful counterweight to population-level correlation studies. Interestingly, a number of previous interventional studies have been performed in mice, suggesting that substantially higher doses of TMAO may have detrimental effects upon learning and memory[68,69,76], although as we and others[77] have identified dose-dependency in the effects of TMAO *in vitro*, it seems plausible that this may reflect a similar phenomenon *in vivo*. The importance of investigating the impact of TMAO under physiologically relevant conditions is further emphasised by a recent study showing TMAO treatment to impair novel object recognition in mice[65], ostensibly an opposite finding to our data achieved with a similar dosing regimen. Importantly, however, mice in this study were maintained on a reduced choline diet, a condition known to alter hepatic metabolism[78]; what impact such changes might have on handling of (TMA and) TMAO by the body is unknown. These discrepancies may be instructive in guarding against incautious extrapolation of TMAO effects from healthy to diseased populations.

Could TMAO contribute to the beneficial effects of a seafood-rich diet?

Consumption of a diet rich in fish and other seafood, known to provide significant quantities of TMAO[79], associates with a reduced risk of cognitive decline[80,81] and protection against cerebrovascular disease[82]. These effects have in large part been attributed to beneficial actions of the omega-3 polyunsaturated fatty acids[83], although there is little evidence that their direct supplementation improves cognitive function[84] or stroke risk[85]. Here we provide evidence that another component of a seafood-rich diet, TMAO, has protective effects on the cerebral vasculature and upon cognition, indicating that broadening the scope of nutritional analyses beyond the omega-3 fatty acids may be worthwhile. In particular, we demonstrate that TMAO acts to stimulate endothelial secretion of a major regulator of BBB permeability, ANXA1[53], leading *via* autocrine/paracrine activation of its receptor FPR2 to cytoskeletal actin rearrangement and enhanced BBB integrity.

This novel interaction between TMAO and ANXA1 may have implications beyond BBB integrity, given that ANXA1 is a major mediator of inflammatory resolution[86], and highlighted by our finding that TMAO treatment inhibited model immune cell adhesion to activated endothelia *in vitro*. As with the link between a seafood-rich diet and cerebrovascular health, fish consumption has been associated with reduced inflammatory disease, again attributed primarily to a role for omega-3 fatty acids[87]. While it is too early to definitively claim an anti-inflammatory role for dietary methylamines, particularly given the opposing actions of TMA and TMAO, our data do suggest that investigation of the bioactive properties of seafood-rich diets should progress beyond omega-3 fatty acids to focus on other potential contributors.

Conclusions

Interest in the role played by the gut microbiota in communication through the gut-brain axis has grown dramatically in the last few years, with much attention focused on the mediating actions of microbe-derived metabolites[88]. While a number of studies have shown patterns in microbial metabolite production that associate with different brain functions[89], detailed understanding of the role of individual molecules remains in its infancy, with defined roles characterised for only a subset of the many molecules known to be released by gut microbes. Here we show that the dietary methylamine TMAO can beneficially modulate both BBB integrity and cognitive function *in vivo*, providing direct mechanistic evidence for a positive role of this microbiome-associated

metabolite, and reinforce the position of the BBB as an interface in the gut-brain axis.

Notably, the positive effects of TMAO that we report stand in contrast to previous work

describing deleterious effects of TMAO exposure at high concentrations or under non-

physiological conditions[77], emphasising the importance of taking a holistic approach

to understanding gut microbiota-host interactions.

COMPETING INTERESTS

The authors declare that they have no competing interests.

ACKNOWLEDGEMENTS

This work was funded by Alzheimer's Research UK Pilot Grant No. ARUK-PPG2016B-
6. PREDEASY™ efflux transporter analysis kits were generously provided through the
SOLVO Biotechnology Research and Academic Collaborative Transporter Studies
(ReACTS) Program. This work used the computing resources of the UK MEDical
BIOinformatics partnership – aggregation, integration, visualisation and analysis of
large, complex data (UK MED-BIO), which was supported by the Medical Research
Council (grant number MR/L01632X/1). SF was supported by Fundación Alfonso
Martín Escudero. TS was supported by a bursary from the Imperial College London
Undergraduate Research Opportunities Programme. LH, DV and SM designed the
experiments. SM performed cellular assays and acute *in vivo* analyses. TS carried out
the initial permeability and TEER assays. MAA performed IgG extravasation studies.
ES produced and provided shRNA treated hCMEC/D3 clones. LH undertook all
processing and analyses of transcriptomic data. RCG provided valuable insight and
advice throughout the project. DV, MP, IR and MM performed the chronic *in vivo* LPS
challenge study and undertook all analyses of behavioural data. SRC, ALC and SF
contributed to preliminary animal work. LH, DV and SM wrote the manuscript. All
authors read and approved the final version of the manuscript.

REFERENCES

1. Braniste, V. *et al.* The gut microbiota influences blood-brain barrier permeability in mice. *Sci. Transl. Med.* **6**, 263ra158-263ra158 (2014).

2. Hoyles, L. *et al.* Microbiome–host systems interactions: protective effects of propionate upon the blood–brain barrier. *Microbiome 2018 61* **6**, 55–55 (2018).

3. Tang, W. H. W. & Hazen, S. L. Microbiome, trimethylamine N-oxide, and cardiometabolic disease. *Transl. Res. J. Lab. Clin. Med.* **179**, 108–115 (2017).

4. Dinicolantonio, J. J., McCarty, M. & Okeefe, J. Association of moderately elevated trimethylamine N-oxide with cardiovascular risk: Is TMAO serving as a marker for hepatic insulin resistance. *Open Heart* **6**, e000890–e000890 (2019).

5. Hoyles, L. *et al.* Metabolic retroconversion of trimethylamine N-oxide and the gut microbiota. *Microbiome* **6**, 73–73 (2018).

6. Zeisel, S. H., Wishnok, J. S. & Blusztajn, J. K. Formation of methylamines from ingested choline and lecithin. *J. Pharmacol. Exp. Ther.* **225**, 320–324 (1983).

7. Kühn, T. *et al.* Intra-individual variation of plasma trimethylamine-N-oxide (TMAO), betaine and choline over 1 year. *Clin. Chem. Lab. Med. CCLM* **55**, 261–268 (2017).

8. Duranton, F. *et al.* Normal and pathologic concentrations of uremic toxins. *J. Am. Soc. Nephrol. JASN* **23**, 1258–70 (2012).

9. Bain, M., Faull, R., Fornasini, G., Milne, R. & Evans, A. Accumulation of Trimethylamine and trimethylamine-N-oxide in End-Stage Renal Disease Patients Undergoing Haemodialysis. *Nephrol. Dial. Transplant. Off. Publ. Eur. Dial. Transpl. Assoc. - Eur. Ren. Assoc.* **21**, 1300–4 (2006).

10. Tang, W. H. W. *et al.* Intestinal microbial metabolism of phosphatidylcholine and cardiovascular risk. *N. Engl. J. Med.* **368**, 1575–1584 (2013).

11. Tang, W. H. W. *et al.* Prognostic value of elevated levels of intestinal microbe-generated metabolite trimethylamine-N-oxide in patients with heart failure: refining the gut hypothesis. *J. Am. Coll. Cardiol.* **64**, 1908–1914 (2014).

12. Wang, Z. *et al.* Non-lethal Inhibition of Gut Microbial Trimethylamine Production for the Treatment of Atherosclerosis. *Cell* **163**, 1585–1595 (2015).

13. Zhu, W. *et al.* Gut Microbial Metabolite TMAO Enhances Platelet Hyperreactivity and Thrombosis Risk. *Cell* **165**, 111–124 (2016).

14. Aldana-Hernández, P. *et al.* Dietary Choline or Trimethylamine N-oxide Supplementation Does Not Influence Atherosclerosis Development in Ldlr−/− and Apoe−/− Male Mice. *J. Nutr.* (2019) doi:10.1093/jn/nxz214.

15. Miller, C. A. *et al.* Effect of egg ingestion on trimethylamine-N-oxide production in
humans: a randomized, controlled, dose-response study. *Am. J. Clin. Nutr.* **100**,
778–786 (2014).
16. Jia, J. *et al.* Assessment of Causal Direction Between Gut Microbiota-Dependent
Metabolites and Cardiometabolic Health: Abi-Directional Mendelian
Randomisation Analysis. *Diabetes* (2019) doi:10.2337/db19-0153.
17. Winther, S. A. *et al.* Utility of Plasma Concentration of Trimethylamine N-Oxide in
Predicting Cardiovascular and Renal Complications in Individuals With Type 1
Diabetes. *Diabetes Care* **42**, 1512–1520 (2019).
18. Huc, T. *et al.* Chronic, low-dose TMAO treatment reduces diastolic dysfunction and
heart fibrosis in hypertensive rats. *Am. J. Physiol.-Heart Circ. Physiol.* **315**, H1805–
H1820 (2018).
19. Collins, H. L. *et al.* L-Carnitine intake and high trimethylamine N-oxide plasma
levels correlate with low aortic lesions in ApoE−/− transgenic mice expressing
CETP. *Atherosclerosis* **244**, 29–37 (2016).
20. Zhao, Z.-H. *et al.* Trimethylamine N-oxide attenuates high-fat high-cholesterol diet-
induced steatohepatitis by reducing hepatic cholesterol overload in rats. *World J.*
Gastroenterol. **25**, 2450–2462 (2019).
21. Dumas, M.-E. *et al.* Microbial-Host Co-metabolites Are Prodromal Markers
Predicting Phenotypic Heterogeneity in Behavior, Obesity, and Impaired Glucose
Tolerance. *Cell Rep.* **20**, 136–148 (2017).
22. Jaworska, K., Bielinska, K., Gawrys-Kopczynska, M. & Ufnal, M. TMA
(trimethylamine), but not its oxide TMAO (trimethylamine-oxide), exerts
haemodynamic effects: implications for interpretation of cardiovascular actions of
gut microbiome. *Cardiovasc. Res.* **115**, 1948–1949 (2019).
23. Huo, X. *et al.* Trimethylamine N-Oxide Metabolites in Early Pregnancy and Risk of
Gestational Diabetes: A Nested Case-Control Study. *J. Clin. Endocrinol. Metab.*
104, 5529–5539 (2019).
24. Poly, C. *et al.* The relation of dietary choline to cognitive performance and white-
matter hyperintensity in the Framingham Offspring Cohort. *Am. J. Clin. Nutr.* **94**,
1584–1591 (2011).
25. Nurk, E. *et al.* Plasma free choline, betaine and cognitive performance: the
Hordaland Health Study. *Br. J. Nutr.* **109**, 511–519 (2013).

26. Leathwood, P. D., Heck, E. & Mauron, J. Phosphatidyl choline and avoidance
performance in 17 month-old SEC/1ReJ mice. *Life Sci.* **30**, 1065–1071 (1982).

27. Bartus, R. T., Dean, R. L., Goas, J. A. & Lippa, A. S. Age-related changes in passive
avoidance retention: modulation with dietary choline. *Science* **209**, 301–3 (1980).

28. Vogt, N. M. *et al.* The gut microbiota-derived metabolite trimethylamine N-oxide is
elevated in Alzheimer's disease. *Alzheimers Res. Ther.* **10**, 124–124 (2018).

29. Gao, Q. *et al.* Decreased levels of circulating trimethylamine N-oxide alleviate
cognitive and pathological deterioration in transgenic mice: A potential therapeutic
approach for Alzheimer's disease. *Aging* **11**, 8642–8663 (2019).

30. Weksler, B. B. *et al.* Blood-brain barrier-specific properties of a human adult brain
endothelial cell line. *FASEB J. Off. Publ. Fed. Am. Soc. Exp. Biol.* **19**, 1872–4
(2005).

31. Maggioli, E. *et al.* Estrogen protects the blood-brain barrier from inflammation-
induced disruption and increased lymphocyte trafficking. *Brain. Behav. Immun.* **51**,
212–222 (2015).

32. Abbott, N. J., Hughes, C. C., Revest, P. A. & Greenwood, J. Development and
characterisation of a rat brain capillary endothelial culture: towards an in vitro
blood-brain barrier. *J. Cell Sci.* **103 (Pt 1**, 23–37 (1992).

33. Coisne, C. *et al.* Mouse syngenic in vitro blood-brain barrier model: a new tool to
examine inflammatory events in cerebral endothelium. *Lab. Investig. J. Tech.*
Methods Pathol. **85**, 734–46 (2005).

34. Pais de Barros, J.-P. *et al.* Quantitative lipopolysaccharide analysis using
HPLC/MS/MS and its combination with the limulus amebocyte lysate assay. *J.*
Lipid Res. **56**, 1363–9 (2015).

35. Gautier, L., Cope, L., Bolstad, B. M. & Irizarry, R. A. affy--analysis of Affymetrix
GeneChip data at the probe level. *Bioinforma. Oxf. Engl.* **20**, 307–15 (2004).

36. Chen, E. Y. *et al.* Enrichr: interactive and collaborative HTML5 gene list enrichment
analysis tool. *BMC Bioinformatics* **14**, 128–128 (2013).

37. Kuleshov, M. V. *et al.* Enrichr: a comprehensive gene set enrichment analysis web
server 2016 update. *Nucleic Acids Res.* **44**, W90-7 (2016).

38. Tarca, A. L. *et al.* A novel signaling pathway impact analysis. *Bioinforma. Oxf. Engl.*
25, 75–82 (2009).

39. Hoyles, L. *et al.* Molecular phenomics and metagenomics of hepatic steatosis in
non-diabetic obese women. *Nat. Med.* **24**, 1070–1080 (2018).

40. Zhang, J. D. & Wiemann, S. KEGGgraph: a graph approach to KEGG PATHWAY
in R and bioconductor. *Bioinforma. Oxf. Engl.* **25**, 1470–1471 (2009).
41. McArthur, S. *et al.* Annexin A1: a central player in the anti-inflammatory and
neuroprotective role of microglia. *J Immunol* **185**, 6317–6328 (2010).
42. Wishart, D. S. *et al.* HMDB 4.0: the human metabolome database for 2018. *Nucleic*
Acids Res. **46**, D608–D617 (2018).
43. Marottoli, F. M. *et al.* Peripheral Inflammation, Apolipoprotein E4, and Amyloid-β
Interact to Induce Cognitive and Cerebrovascular Dysfunction: *ASN Neuro* (2017)
doi:10.1177/1759091417719201.
44. Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner with low
memory requirements. *Nat. Methods* **12**, 357–360 (2015).
45. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose
program for assigning sequence reads to genomic features. *Bioinforma. Oxf. Engl.*
30, 923–930 (2014).
46. Hölter, S. M. *et al.* Tests for Anxiety-Related Behavior in Mice. *Curr. Protoc. Mouse*
Biol. **5**, 291–309 (2015).
47. Davis, K. E., Eacott, M. J., Easton, A. & Gigg, J. Episodic-like memory is sensitive
to both Alzheimer's-like pathological accumulation and normal ageing processes
in mice. *Behav. Brain Res.* **254**, 73–82 (2013).
48. Leger, M. *et al.* Object recognition test in mice. *Nat. Protoc.* **8**, 2531–2537 (2013).
49. Denninger, J. K., Smith, B. M. & Kirby, E. D. Novel Object Recognition and Object
Location Behavioral Testing in Mice on a Budget. *J. Vis. Exp. JoVE* (2018)
doi:10.3791/58593.
50. Thomas, R., Morris, A. W. J. & Tai, L. M. Epidermal growth factor prevents APOE4-
induced cognitive and cerebrovascular deficits in female mice. *Heliyon* **3**, e00319
(2017).
51. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).
52. Lochhead, J. J., Yang, J., Ronaldson, P. T. & Davis, T. P. Structure, Function, and
Regulation of the Blood-Brain Barrier Tight Junction in Central Nervous System
Disorders. *Front. Physiol.* **11**, 914 (2020).
53. Cristante, E. *et al.* Identification of an essential endogenous regulator of blood-
brain barrier integrity, and its pathological and therapeutic implications. *Proc. Natl.*
Acad. Sci. U. S. A. **110**, 832–841 (2013).

54. McArthur, S. *et al.* Annexin A1 regulates hormone exocytosis through a mechanism
involving actin reorganization. *FASEB J* **23**, 4000–4010 (2009).

55. Bena, S., Brancaleone, V., Wang, J. M., Perretti, M. & Flower, R. J. Annexin A1
interaction with the FPR2/ALX receptor: identification of distinct domains and
downstream associated signaling. *J. Biol. Chem.* **287**, 24690–7 (2012).

56. Cho, C. E. *et al.* Trimethylamine-N-oxide (TMAO) response to animal source foods
varies among healthy young men and is influenced by their gut microbiota
composition: A randomized controlled trial. *Mol. Nutr. Food Res.* **61**, 1600324
(2017).

57. Montagne, A. *et al.* Blood-Brain Barrier Breakdown in the Aging Human
Hippocampus. *Neuron* **85**, 296–302 (2015).

58. Taheri, S. *et al.* Blood-brain barrier permeability abnormalities in vascular cognitive
impairment. *Stroke J. Cereb. Circ.* **42**, 2158–63 (2011).

59. Bowman, G. L. *et al.* Blood-brain barrier impairment in Alzheimer disease: stability
and functional significance. *Neurology* **68**, 1809–14 (2007).

60. Merino, J. G. *et al.* Blood-brain barrier disruption after cardiac surgery. *AJNR Am.*
J. Neuroradiol. **34**, 518–523 (2013).

61. Hu, N. *et al.* Involvement of the blood-brain barrier opening in cognitive decline in
aged rats following orthopedic surgery and high concentration of sevoflurane
inhalation. *Brain Res.* **1551**, 13–24 (2014).

62. Abrahamov, D. *et al.* Blood-Brain Barrier Disruption After Cardiopulmonary
Bypass: Diagnosis and Correlation to Cognition. *Ann. Thorac. Surg.* **104**, 161–169
(2017).

63. Yang, S. *et al.* Anesthesia and Surgery Impair Blood-Brain Barrier and Cognitive
Function in Mice. *Front. Immunol.* **8**, 902–902 (2017).

64. Zenaro, E., Piacentino, G. & Constantin, G. The blood-brain barrier in Alzheimer's
disease. *Neurobiol. Dis.* (2016) doi:10.1016/j.nbd.2016.07.007.

65. Brunt, V. E. *et al.* The gut microbiome–derived metabolite trimethylamine N-oxide
modulates neuroinflammation and cognitive function with aging. *GeroScience*
(2020) doi:10.1007/s11357-020-00257-2.

66. He, W. *et al.* Trimethylamine N-Oxide, a Gut Microbiota-Dependent Metabolite, is
Associated with Frailty in Older Adults with Cardiovascular Disease. *Clin. Interv.*
Aging **15**, 1809–1820 (2020).

67. Zhu, C. *et al.* Association of plasma trimethylamine-N-oxide levels with post-stroke
cognitive impairment: a 1-year longitudinal study. *Neurol. Sci. Off. J. Ital. Neurol.*
Soc. Ital. Soc. Clin. Neurophysiol. **41**, 57–63 (2020).

68. Zhao, L. *et al.* Higher Circulating Trimethylamine N-oxide Sensitizes Sevoflurane-
Induced Cognitive Dysfunction in Aged Rats Probably by Downregulating
Hippocampal Methionine Sulfoxide Reductase A. *Neurochem. Res.* **44**, 2506–
2516 (2019).

69. Meng, F., Li, N., Li, D., Song, B. & Li, L. The presence of elevated circulating
trimethylamine N-oxide exaggerates postoperative cognitive dysfunction in aged
rats. *Behav. Brain Res.* **368**, 111902–111902 (2019).

70. Sanguinetti, E. *et al.* Microbiome-metabolome signatures in mice genetically prone
to develop dementia, fed a normal or fatty diet. *Sci. Rep.* **8**, (2018).

71. Liu, J. *et al.* Baicalin ameliorates neuropathology in repeated cerebral ischemia-
reperfusion injury model mice by remodeling the gut microbiota. *Aging* **12**, 3791–
3806 (2020).

72. Wang, Q.-J. *et al.* Concomitant memantine and Lactobacillus plantarum treatment
attenuates cognitive impairments in APP/PS1 mice. *Aging* **12**, 628–649 (2020).

73. Sawicka, A. K., Renzi, G. & Olek, R. A. The bright and the dark sides of L-carnitine
supplementation: a systematic review. *J. Int. Soc. Sports Nutr.* **17**, 49 (2020).

74. Chung, S. J. *et al.* Gut microbiota-derived metabolite trimethylamine N-oxide as a
biomarker in early Parkinson's disease. *Nutr. Burbank Los Angel. Cty. Calif* **83**,
111090 (2020).

75. Zhuang, Z. *et al.* Causal relationships between gut metabolites and Alzheimer's
disease: a bidirectional Mendelian randomization study. *Neurobiol. Aging* (2020)
doi:10.1016/j.neurobiolaging.2020.10.022.

76. Li, D. *et al.* Trimethylamine-N-oxide promotes brain aging and cognitive impairment
in mice. *Aging Cell* **17**, (2018).

77. Papandreou, C., Moré, M. & Bellamine, A. Trimethylamine N-Oxide in Relation to
Cardiometabolic Health-Cause or Effect? *Nutrients* **12**, (2020).

78. Zeisel, S. H. Dietary choline: biochemistry, physiology, and pharmacology. *Annu.*
Rev. Nutr. **1**, 95–121 (1981).

79. Lundstrom, R. C. & Racicot, L. D. Gas chromatographic determination of
dimethylamine and trimethylamine in seafoods. *J. - Assoc. Off. Anal. Chem.* **66**,
1158–1163 (1983).

80. Zeng, L.-F. *et al.* An exploration of the role of a fish-oriented diet in cognitive
decline: a systematic review of the literature. *Oncotarget* **8**, 39877–39895 (2017).
81. Keenan, T. D. *et al.* Adherence to a Mediterranean diet and cognitive function in
the Age-Related Eye Disease Studies 1 & 2. *Alzheimers Dement.* **16**, 831–842
(2020).
82. Zhao, W. *et al.* Fish Consumption and Stroke Risk: A Meta-Analysis of Prospective
Cohort Studies. *J. Stroke Cerebrovasc. Dis. Off. J. Natl. Stroke Assoc.* **28**, 604–
611 (2019).
83. Zhang, Y. *et al.* Intakes of fish and polyunsaturated fatty acids and mild-to-severe
cognitive impairment risks: a dose-response meta-analysis of 21 cohort studies.
Am. J. Clin. Nutr. **103**, 330–340 (2016).
84. Cooper, R. E., Tye, C., Kuntsi, J., Vassos, E. & Asherson, P. Omega-3
polyunsaturated fatty acid supplementation and cognition: A systematic review and
meta-analysis. *J. Psychopharmacol. Oxf. Engl.* **29**, 753–763 (2015).
85. Abdelhamid, A. S. *et al.* Omega-3 fatty acids for the primary and secondary
prevention of cardiovascular disease. *Cochrane Database Syst. Rev.* **11**,
CD003177 (2018).
86. Gobbetti, T. & Cooray, S. N. Annexin A1 and resolution of inflammation: tissue
repairing properties and signalling signature. *Biol. Chem.* **397**, 981–993 (2016).
87. Wall, R., Ross, R. P., Fitzgerald, G. F. & Stanton, C. Fatty acids from fish: the anti-
inflammatory potential of long-chain omega-3 fatty acids. *Nutr. Rev.* **68**, 280–289
(2010).
88. Nicholson, J. K. *et al.* Host-gut microbiota metabolic interactions. *Science* **336**,
1262–7 (2012).
89. Needham, B. D., Kaddurah-Daouk, R. & Mazmanian, S. K. Gut microbial
molecules in behavioural and neurodegenerative conditions. *Nat. Rev. Neurosci.*
21, 717–731 (2020).
897

898 **Table 1.** BBB-associated genes whose expression was upregulated upon exposure

899 of hCMEC/D3 cells to TMAO.

Gene	Entrez ID	Description	Log$_2$ fold change	Category	P_{FDR}
TFRC	7037	Transferrin receptor	0.23	Transporter proteins	0.054
ABCC4	10257	ATP binding cassette subfamily C member 4	0.20	Transporter proteins	0.088
ANXA1	301	Annexin A1	0.16	Cell Adhesion/Junctional proteins/Cytoskeletal factors	0.088
CDH2	1000	Cadherin 2	0.31	Cell Adhesion/Junctional proteins/Cytoskeletal factors	0.095

904 **Table 2.** Daily consumption of TMAO and water in mice chronically treated with TMAO
905 and/or LPS.

907 Data are mean ± standard deviation.

Variable	Sham-treated mice	LPS-treated mice	*P* value
TMAO intake (mg/day/mouse)	2.79 ± 0.38	2.82 ± 0.25	0.7
TMAO intake (mg/kg/mouse)	85.0 ± 11.4	88.9 ± 7.9	0.14
Water consumption (ml/day/mouse)	5.57 ± 0.8	5.64 ± 0.5	0.48

FIGURE LEGENDS

Fig. 1. Effects of TMAO and TMA on integrity of hCMEC/D3 cell monolayers. (A) Assessment of paracellular permeability of hCMEC/D3 monolayers to a 70 kDa FITC-dextran tracer following treatment for 24 h with varying doses of TMA (0.4 – 40 µM) or TMAO (40 – 4000 µM). Data are expressed as mean ± s.e.m., n=4 independent experiments. (B) Assessment of TEER of hCMEC/D3 monolayers to a 70kDa FITC-dextran tracer following treatment for 24 h with varying doses of TMA (0.4 – 40 µM) or TMAO (40 – 4000 µM). Data are expressed as mean ± s.e.m., n=4 independent experiments. (C) Adhesion of U937 monocytic cells to TNFα-stimulated hCMEC/D3 monolayers (10 ng/ml, 16 h) that had been treated or not for 24 h with 0.4 µM TMA or 40 µM TMAO. Data are expressed as mean ± s.e.m., n=3 independent experiments.

Fig. 2. Effects of TMA and TMAO on gene expression in hCMEC/D3 cells. (A) Heatmap showing expression of the 49 genes found to be significantly (P_{FDR}<0.1) differentially expressed upon exposure of hCMEC/D3 cells to 0.4 µM TMA (n=5 per group). (B) Heatmap showing expression of the 440 genes found to be significantly (P_{FDR}<0.1) differentially expressed upon exposure of hCMEC/D3 cells to 40 µM TMAO (n=5 per group). (C) Biological processes associated with genes found to be significantly upregulated (n=39) or downregulated (n=10) upon exposure of cells to TMA. (D) Biological processes of genes found to be significantly upregulated (n=341) or downregulated (n=99) upon exposure of cells to TMAO. Images in (C, D) shown based on Enrichr P value ranking from GO analysis. (E) Topological analysis of the KEGG networks associated with the 440 genes whose expression was significantly affected upon exposure of cells to TMAO (blue, significantly downregulated; red, significantly upregulated); genes of similar cellullar role are highlighted. (F) Confocal microscopic analysis of expression of fibrillar actin (F-actin) and the tight junction component zonula occludens-1 (ZO-1) in hCMEC/D3 cells following treatment for 24 h with 0.4 µM TMA or 40 µM TMAO. Images are representative of at least three independent experiments.

Fig. 3. Annexin A1 (ANXA1) signalling mediates effects of TMAO on hCMEC/D3 cells. (A) Total cellular expression of ANXA1 in hCMEC/D3 cells treated for 24 h with 0.4 µM TMA or 40 µM TMAO. Data are expressed as mean ± s.e.m., n=5-7 independent

experiments. (B) Medium ANXA1 content of hCMEC/D3 monolayers treated for 24 h with 0.4 µM TMA or 40 µM TMAO. Data are expressed as mean $\pm$ s.e.m., n=3 independent experiments. (C) Assessment of paracellular permeability of monolayers of wild-type hCMEC/D3 cells, or hCMEC/D3 cells stably transfected with either a scramble shRNA sequence or an shRNA sequence targeting ANXA1 to a 70kDa FITC-dextran tracer following treatment for 24 h with 40 µM TMAO. Data are expressed as mean $\pm$ s.e.m., n=4 independent experiments. (D) Assessment of TEER of monolayers of wild-type hCMEC/D3 cells, or hCMEC/D3 cells stably transfected with either a scramble shRNA sequence or an shRNA sequence targeting ANXA1 following treatment for 24 h with 40 µM TMAO. Data are expressed as mean $\pm$ s.e.m., n=4 independent experiments. (E) Assessment of paracellular permeability of hCMEC/D3 cells to a 70kDa FITC-dextran tracer following treatment for 24 h with 40 µM TMAO, with or without 10 min pre-treatment with the FPR2 antagonist WRW$_4$ (10 µM). Data are expressed as mean $\pm$ s.e.m., n=3 independent experiments. (F) Assessment of TEER of hCMEC/D3 cells following treatment for 24 h with 40 µM TMAO, with or without 10 min pre-treatment with the FPR2 antagonist WRW4 (10 µM). Data are expressed as mean $\pm$ s.e.m., n=3 independent experiments.

Fig. 4. Acute treatment with TMAO promotes BBB integrity *in vivo*. (A) Extravasation of Evans blue dye into brain parenchyma over a 1 h period in 2-month-old male C57Bl/6J mice following i.p. injection of 1.8 mg/kg TMAO for 2 h, 6 h or 24 h *vs.* a saline injected control. Data are normalised to plasma Evans blue content, and are expressed as mean $\pm$ s.e.m., n=5-6 mice. (B) Extravasation of Evans blue dye into brain parenchyma over a 1 h period in 2-month-old male C57Bl/6J mice following i.p. injection of saline or E. coli O111:B4 LPS (3 mg/kg) with or without subsequent i.p. injection of 1.8 mg/kg TMAO according to the schedule shown. Data are normalised to plasma Evans blue content, and are expressed as mean $\pm$ s.e.m., n=4-6 mice.

Fig. 5. Acute exposure of mice to TMAO significantly alters the whole brain transcriptome. (A) Heatmap showing expression of the 76 genes found to be significantly (P_{FDR}<0.1) differentially expressed in the mouse brain after 2 h exposure to 1.8 mg/kg TMAO (n=3 per group). Data were scaled by row. (B) Over-representation analysis (Enrichr) showing KEGG pathways associated with the 76 genes. (C)

Comparative analysis of significantly differentially expressed genes identified groupings associated with distinct biological functions. (D) Among the 197 BBB-specific genes identified in the data set, only *App* and *Cpe* were significantly (P_{FDR}<0.1) differentially expressed in the mouse brain after 2 h exposure to TMAO. Data are shown as mean ± s.d, *n*=3 per group. Individual data points are not shown due to the negligible values of the s.d.

Fig. 6. Effect of long-term TMAO exposure on BBB integrity and cognitive function of mice. (A) Body weight gain in mice treated with TMAO through their drinking water (0.5 mg/ml) over 2 months, combined with a chronic low dose administration of LPS (0.5 mg/kg/week, i.p.). Data are expressed as mean ± s.e.m., *n*=8 mice, columns with different letters are significantly different at *P*<0.05. (B) Cerebellar permeability index to sodium fluorescein 2h following administration in animals previously treated with TMAO through their drinking water (0.5 mg/ml) over 2 months, combined with a chronic low dose administration of LPS (0.5 mg/kg/week, i.p.). Data are expressed as mean ± s.e.m., *n*=8 mice, columns with different letters are significantly different at *P*<0.05. (C) Typical confocal microscopic images of perivascular IgG deposition in male C57Bl/6J mice treated with TMAO through their drinking water (0.5 mg/ml) over 2 months, combined with a chronic low dose administration of LPS (0.5 mg/kg/week, i.p.). *Griffonia simplicifolia* isolectin B$_4$ (red) defines endothelial cells, areas of IgG deposition (white) are highlighted by arrow heads. (D) Distance travelled, (E) movement speed and (F) percentage of time in the centre as measured in the OFT in animals previously treated with TMAO through their drinking water (0.5 mg/ml) over 2 months, combined with a chronic low dose administration of LPS (0.5 mg/kg/week, i.p.). Data are expressed as mean ± s.e.m., *n*=8 mice. (G) Novel object discrimination index, calculated as described in Methods, of animals previously treated with TMAO through their drinking water (0.5 mg/ml) over 2 months, combined with a chronic low dose administration of LPS (0.5 mg/kg/week, i.p.). Data are expressed as mean ± s.e.m., *n*=8 mice, columns with different letters are significantly different at *P*<0.05. (H) Percentage of spontaneous alternation and (I) total distance travelled in the Y-maze test for animals previously treated with TMAO through their drinking water (0.5 mg/ml) over 2 months, combined with a chronic low dose administration of LPS (0.5 mg/kg/week, i.p.). Data are expressed as mean ± s.e.m., *n*=8 mice.

Supplementary Fig. 1. Confirmation of ANXA1 targeting and knock-down in hCMEC/D3 cells stably transfected with appropriate shRNA sequences. (A) Typical flow cytometric profiles of wild-type cells, cells transfected with a scramble shRNA sequence and cells transfected with an ANXA1-targeting shRNA sequence, alongside unstained and second antibody controls. (B) Expression of ANXA1 in wild-type hCMEC/D3 cells, cells transfected with a scramble shRNA sequence and cells transfected with an ANXA1-targeting shRNA sequence. Data are expressed as mean $\pm$ s.e.m., n=4 independent experiments.

Supplementary Fig. 2. Neither TMAO nor TMA treatment affected the major endothelial efflux transporters BCRP or P-glycoprotein. (A-D) *In vitro* analysis revealed no significant effects of either TMAO or TMA upon BCRP (A, C) or P-glycoprotein (B, D) activity. Dashed lines represent Loess regression fits, with shading representing 95% confidence intervals. (E) hCMEC/D3 cell surface expression of BCRP was unaffected by 24 h exposure to TMA (0.4 µM) or TMAO (40 µM). Data are expressed as mean $\pm$ s.e.m., n=3 independent experiments. (F) hCMEC/D3 cell surface expression of P-glycoprotein was unaffected by 24 h exposure to TMA (0.4 µM) or TMAO (40 µM). Data are expressed as mean $\pm$ s.e.m., n=3 independent experiments.

Supplementary Fig. 3. *In vitro* (hCMEC/D3 cell) *FMO3* gene expression is unaffected by exposure to TMA (0.4 µM) or TMAO (40 µM). Dark blue lines represent the standard deviation (+/-). There was no statistically significant difference between TMA and the control, nor between TMAO and the control (Supplementary Tables 2 and 3). Individual data points are not shown due to the negligible values of the standard deviations.

Supplementary Fig. 4. Neither chronic low-dose LPS nor TMAO treatment significantly increases serum inflammatory cytokines. (A) Serum TNFα concentrations in male C57Bl/6J mice treated with TMAO through their drinking water (0.5 mg/ml) over 2 months, combined with a chronic low dose administration of LPS (0.5 mg/kg/week, i.p.). Data are expressed as mean $\pm$ s.e.m., n=7-8 mice. (B) Serum IL-1β concentrations in male C57Bl/6J mice treated with TMAO through their drinking water (0.5 mg/ml) over 2 months, combined with a chronic low dose administration of

1046 LPS (0.5 mg/kg/week, i.p.). Data are expressed as mean $\pm$ s.e.m., *n*=7-8 mice.

1047

1048 **Supplementary Table 1.** Normalized microarray data for hCMEC/D3 treated or not

1049 with TMA (0.4 µM) or TMAO (40 µM)

1050

1051 **Supplementary Table 2.** Differential gene expression for control compared with TMA

1052 treatment, as assessed using LIMMA

1053

1054 **Supplementary Table 3.** Differential gene expression for control compared with

1055 TMAO treatment, as assessed using LIMMA

1056

1057 **Supplementary Table 4.** KEGG-based SPIA of genes whose expression was

1058 significantly affected by TMAO

1059

1060 **Supplementary Table 5.** GO (biological process) analysis of TMA-up-regulated

1061 genes as assessed using Enrichr

1062

1063 **Supplementary Table 6.** GO (biological process) analysis of TMA-down-regulated

1064 genes as assessed using Enrichr

1065

1066 **Supplementary Table 7.** GO (biological process) analysis of TMAO-up-regulated

1067 genes as assessed using Enrichr

1068

1069 **Supplementary Table 8.** GO (biological process) analysis of TMAO-down-regulated

1070 genes as assessed using Enrichr

1071

1072 **Supplementary Table 9.** Differential gene expression of BBB-associated genes (*in*

1073 *vitro* TMA treatment)

1074

1075 **Supplementary Table 10.** Differential gene expression of BBB-associated genes (*in*

1076 *vitro* TMAO treatment)

1077

1078   **Supplementary Table 11.** Differential gene expression in mouse brain for control

1079   compared with TMAO treatment, as assessed using DESeq2

1080

1081   **Supplementary Table 12.** Differential expression of BBB-associated genes in mouse

1082   brain for control compared with TMAO treatment, as assessed using DESeq2

1083

Figure 1

A

B

C

Figure 2

Figure 3

Figure 4

A

B

Figure 5

Figure 6

Supplementary Figure 1

Supplementary Figure 4