Glucose inhibits neutrophil migration and haemostasis, and accelerates diet-induced

hyperlipidaemia in zebrafish embryos

Simone Morris[1], Pradeep Cholan[1]†, Warwick J Britton[1,2], Stefan H Oehlers[1,3] *

[1] Tuberculosis Research Program at the Centenary Institute, The University of Sydney,

Camperdown NSW 2050 Australia

[2] Department of Clinical Immunology, Royal Prince Alfred Hospital, Camperdown, NSW 2050,

Australia

[3] The University of Sydney, Discipline of Infectious Diseases & Immunology and Marie Bashir

Institute, Camperdown NSW 2050 Australia

† Co-corresponding author email: p.cholan@centenary.org.au

* Corresponding author email: stefan.oehlers@sydney.edu.au

ABSTRACT

Hyperglycaemia damages the microvasculature in part through the reduced recruitment of immune

cells and interference with platelet signalling, leading to poor wound healing and accelerated lipid

deposition in mammals. We investigated the utility of zebrafish embryos to model the effect of

glucose on leukocyte recruitment, thrombosis, and hyperlipidaemia by treating embryos with

exogenous glucose. Exogenous glucose supplementation reduced neutrophil recruitment, and

thrombocyte and fibrin plug formation following tail transection in zebrafish embryos, but did not

affect the recruitment of macrophages. We observed accelerated lipid accumulation in glucose-

supplemented embryos challenged with high fat chicken egg yolk feeding. Our study identifies

conserved and divergent effects of high glucose in the tractable zebrafish embryo model system

compared to mammals.

INTRODUCTION

Hyperglycaemic damage to the microvasculature is hypothesised to underpin much of the pathology associated with diabetes in mammals, including perturbations to leukocyte biology, haemostasis, and the accumulation of lipid laden macrophages in the vessel wall [1-4]. Previous mammalian research has demonstrated that hyperglycaemia damages the microvasculature resulting in reduced expression of endothelial adhesion molecules for immune cell recruitment [2, 4]. As a result, fewer neutrophils and macrophages are recruited to diabetic wounds [1, 2, 5]. Of the macrophages and neutrophils that eventually arrive, there is a skewing of differentiation towards an inflammatory phenotype owing to the inflammatory nature of the diabetic wound microenvironment [6, 7].

Mammals with hyperglycaemia demonstrate perturbed coagulation and platelet signalling, causing disruption of haemostasis [3, 8]. Reduced efficiency of haemostasis results in unstable and ineffective clots within diabetic foot ulcers [3, 9]. Treatments for diabetic foot ulcers can involve platelet and fibrin therapy, indicating an important role for inadequate fibrin clot production in the ulceration process [10, 11].

Hyperglycaemia-induced damage to the microvasculature also increases vascular lipid accumulation in conjunction with hyperlipidaemia [12]. Hyperlipidaemia and hyperglycaemia are compounding risk factors for the development of Type 2 diabetes, associated with the 'Western Diet' consisting of fat and sugar alongside limited exercise [13, 14]. There are multiple mechanisms by which hyperglycaemia damages the microvasculature: through the formation of advanced glycation products [3], the induction of oxidative stress [15], interfering with nitric oxide production [16], and inducing macrophages to form lipid-laden foam cells [17]. The degradation of the endothelial structural integrity increases the rate of lipid deposition by providing a physical niche for lipid infiltration [12]. This can ultimately lead to the development of atheroma, which are common in diabetic patients [17].

Our study investigates the effect of high glucose environments on inflammation, thrombosis and hyperlipidaemia using a zebrafish model. Zebrafish are established models for investigating each of these processes in isolation [18-21]. Pertinent to this study, zebrafish have similar clotting, metabolic, and immune systems to mammals, and these have been used to provide insight into the shared function of these systems across vertebrates [19, 20, 22]. Here we have combined these models to investigate the role of high glucose environments in disrupting immune cell recruitment to wounds, thrombosis, and lipid accumulation in zebrafish embryos.

METHODS

Zebrafish Husbandry

Zebrafish embryos were produced through natural spawning (Sydney Local Health District Animal Welfare Committee Approval 17-036). The strains used were $Tg(lyzC:DsRed)^{nz50}$ to visualise neutrophils, $Tg(mfap4:turquoise)^{xt27}$ to visualise macrophages, $Tg(itga2b:gfp)^{la2}$ to visualise thrombocytes, and $Tg(fabp10a:fgb-gfp)^{mi4001}$ to visualise fibrin deposition. From 1-5 days post fertilisation (dpf) embryos were kept in a dark incubator at 28°C.

Induction of High Glucose Concentrations in the Zebrafish

Injection method: Eggs were injected with approximately 15 nmol of glucose, or PBS as a control within four hours of fertilisation (Figure 1A). Immersion method: Dechorionated 2 dpf embryos were immersed in 5% mannitol (osmolarity control) or glucose dissolved in E3 media (Figure 1B), media was changed daily to reduce microbial growth.

Glucose Oxidase Assay

Embryos were snap frozen and stored at -20°C. Embryos were lysed in the buffer solution from the Amplex Red Oxidase kit (Sigma: A22188), lysates were sedimented by centrifugation, and the

Amplex Red Glucose oxidase kit was used to measure the glucose content of the supernatant in
accordance with the manufacturer's instructions.

Tail Transection Assays

Caudal fin amputations were performed on embryos at 5 dpf. Embryos were anesthetised with 2.5%

(v/v) ethyl-3-aminobenzoate methanesulfonate (tricaine) (Sigma, E10521), wounded posterior to

the notochord using a sterile scalpel and kept in a 28°C incubator to recover as previously described

[23]. Wounded embryos were imaged at 6 hours (neutrophil and macrophages) or 2.5 hours (fibrin

and thrombocytes).

Hyperlipidaemia Assay

Post 5 dpf, embryos were transferred to a 28°C incubator with a 14/10 hour light/dark cycle.

Embryos were placed in an E3 solution containing 0.05% of emulsified chicken egg yolk from 5

dpf. Each day, a random sample of embryos were removed, euthanised, and fixed in

paraformaldehyde. Embryos were stained with Oil Red O to quantitate lipid accumulation, as

previously described [21, 24, 25].

Imaging and Image Analysis

Embryos were imaged using a Leica M205FA fluorescent microscope. ImageJ software was used to

quantify fluorescent pixel count within 100 μm of the wound site for transgenic wound assays as

previously described [23, 26].

Oil Red O staining was quantified in ImageJ by adjusting the colour threshold to eliminate non-red

signal. The image was then converted to a binary mask, and the tail region posterior to the swim

bladder was selected to measure the number and area of particles [25].

Statistical analysis

Outliers were excluded using a 1% ROUT test. Statistical testing was carried out by ANOVA with

correction for multiple comparisons or Student's *t*-tests as appropriate using GraphPad Prism. Data

are expressed as mean ± SD. Every datapoint represents a single embryo unless otherwise noted.

RESULTS

Exogenous glucose exposure increases glucose in zebrafish embryos

Figure 1: Injection and immersion methods increase glucose levels in zebrafish embryos.

A. Relative concentration of glucose in 5 dpf embryos that had been injected with 15 nmol glucose as eggs. Statistical testing by *t*-test, each data point is representative of a group of n = 10-30 embryos.

B. Relative concentration of glucose in 5 dpf embryos immersed in 5% solutions of glucose or mannitol since 2 dpf. Statistical testing by ANOVA, each data point is representative of a group of n = 10-30 embryos.

To establish the efficacy of the injection and immersion techniques to increase glucose concentrations in 5 dpf zebrafish embryos, we conducted a glucose oxidase assay. Consistent with past literature, we observed an increase in the glucose concentration contained within the glucose-injected and -immersed embryos compared to controls (Figure 1) [27, 28]. The injection method was selected for our study because it decreased the increased risk of microbial overgrowth compared to the immersion method.

Glucose impedes neutrophil, but not macrophage, chemotaxis in zebrafish embryos.

Figure 2: High glucose reduced neutrophil, but not macrophage, recruitment to a tail wound.

A. Schematic of experiment to measure immune cell recruitment to a tail wound.

B. Quantification of macrophage recruitment following tail transection.

C. Quantification of neutrophil recruitment following tail transection.

Statistical testing by *t*-test. Data are from 2 biological replicates.

Altered innate immune cell recruitment to wounds is a conserved feature of hyperglycaemia in mammals [4, 5, 29-31]. To determine if this phenomenon was conserved in zebrafish embryos, we utilised the tail transection wound model which causes reproducible leukocyte recruitment [23] (Figure 2A). We first performed this assay using transgenic *Tg(mfap4:turquoise)^{xt27}* embryos, where macrophages are marked by turquoise fluorescent protein to quantify macrophage recruitment. Surprisingly, we observed no difference in the number of recruited macrophages between the glucose-injected and control embryos at 6 hours post wounding (hpw) (Figure 2B).

We then used *Tg(lyzC:DsRed)*[nz50] embryos, where neutrophils are marked by DsRed fluorescent

protein, to measure the recruitment of neutrophils. We observed a reduction in neutrophil

recruitment between the control and glucose-injected embryos at 6 hpw (Figure 1C).

Together, these results indicate that neutrophil, but not macrophage, recruitment is perturbed by

high glucose in zebrafish embryos.

Glucose impedes haemostasis in zebrafish embryos

Figure 3: High glucose reduced thrombocyte and fibrin accumulation to a tail wound.

A. Schematic of experiment to visualise haemostasis following tail transection.

B. Representative overlay images of thrombocytes (green) demonstrating recruitment to tail

wound in control and glucose *Tg(itga2b:gfp)*[la2] embryos. Arrowhead indicates location of

severed major blood vessels at tail transection site.

C. Quantification of thrombocyte plug size following tail transection.

D. Quantification of fibrin clot size following tail transection.

Statistical testing by *t*-test. Data are representative of 2 biological replicates.

Hyperglycaemia perturbs coagulation and platelet activation in mammals, resulting in ineffective

haemostasis [3, 8]. To visualise the effects of high glucose on haemostasis in zebrafish embryos, we

first used the *Tg(itga2b:gfp)* *la2* line where thrombocytes are tagged with GFP [32, 33]. Following

tail transection (Figure 3A), high glucose *Tg(itga2b:gfp)* *la2* embryos demonstrated reduced

thrombocyte accumulation (Figure 3B-C).

To determine if the reduction in thrombocyte recruitment was mirrored by perturbed clotting, we

conducted tail transections on *Tg(fabp10a:fgb-gfp)*[mi4001] embryos which express fluorescently

tagged fibrinogen and allow visualisation of clots [34]. Glucose-injected *Tg(fabp10a:fgb-gfp)*[mi4001]

embryos had reduced fibrin accumulation following tail transection compared to PBS-injected

embryos (Figure 3D).

Together, these results demonstrate that a high glucose environment inhibits haemostasis in

zebrafish embryos.

High glucose accelerated hyperlipidaemia in zebrafish embryos challenged with a high fat diet

Figure 4: Glucose-injected embryos have increased lipid accumulation following a high fat diet.

A. Schematic of the high fat feeding challenge assay to measure lipid accumulation in glucose-injected embryos.

B. Bright field images of 6 dpf Oil Red O-stained embryos, demonstrating darker vascular staining in glucose-injected embryos. Box indicates location of inset, arrowheads indicate stained intercostal vessels in inset, asterisk indicates intestinal lumen which was excluded from analysis.

C. Quantification of lipid accumulation in 5 dpf glucose-injected embryos.

D. Quantification of lipid accumulation in 6 dpf glucose-injected embryos challenged with a high fat diet from 5 dpf.

E. Quantification of lipid accumulation in 7 dpf glucose-injected embryos challenged with a high fat diet from 5 dpf.

Statistical testing by *t*-test. Data are representative of 2 biological replicates.

Hyperglycaemia and hyperlipidaemia are intimately associated in mammals [35]. To determine if this interaction is conserved in zebrafish embryos, we fed glucose-injected embryos a high fat diet

consisting of emulsified chicken egg yolk from 5-7 dpf (Figure 4A). Glucose- and PBS-injected

embryos had similar Oil Red O vascular staining prior to the initiation of chicken egg yolk feeding

at 5 dpf (Figure 4B). Quantification of Oil Red O staining revealed glucose-injected embryos had

increased vascular lipid content at one and two days post feeding (Figure 4D-E).

These results recapitulate the interaction between hyperglycaemia and hyperlipidaemia seen in

mammals [15], demonstrating that zebrafish embryos are a robust model to study this conserved

interaction.

DISCUSSION

In this study, we explored the effect of high glucose levels on inflammation, thrombosis, and lipid

accumulation in the zebrafish embryo model. We found that high glucose reduces neutrophil

recruitment and haemostasis around wound sites. In addition, we determined that high glucose

accelerates lipid accumulation when embryos are challenged with a high fat diet.

Hyperglycaemia in mammals causes vascular dysfunction that restricts the recruitment of a broad

range of immune cells to wounds [16]. It was therefore surprising that high glucose appeared to

have no effect on the recruitment of macrophages to the tail wound in this study. It is possible that

high glucose in zebrafish embryos does not affect abluminal crawling of macrophages along blood

vessels, since wound-responsive macrophages have been demonstrated to move predominantly

through interstitial tissue in zebrafish embryos [5, 36].

Our finding that high glucose environments reduce the recruitment of neutrophils to a tail wound is

consistent with studies of diabetic foot ulcers [2, 4]. In mammals, high glucose environments have

been shown to interfere with metabolites and enzymes related to neutrophil function [37].

Neutrophils may also be restricted by hyperglycaemia mediated vascular dysfunction as they are

more reliant on extravasation to reach distal sites in zebrafish embryos than macrophages [16, 36].

Although neutrophil dysfunction was conserved, macrophage function was unaffected by high

glucose environments suggesting that glucose injection of zebrafish embryos may not be a suitable

platform for studying the impact of high glucose on leukocyte recruitment.

The relationship between hyperglycaemia and hyperlipidaemia has been previously reported in

various mammalian species [15, 28, 38], and a recent study by Wang *et al.* has demonstrated a

similar the interaction of a high cholesterol diet with glucose immersion on vascular lipid

accumulation in zebrafish after 10 days of feeding [28]. Our study using chicken egg yolk as a high

fat diet challenge demonstrates a dramatically accelerated accumulation of lipid after just one day of

feeding. This chicken egg yolk feeding-based model, therefore provides a more rapid model to

investigate lipid accumulation.

In summary, we report glucose-supplemented zebrafish embryos as a tractable platform to

investigate the conserved interactions between glucose and neutrophil biology, haemostasis, and

diet-induced hyperlipidaemia.

ACKNOWLEDGEMENTS

We thank other members of the Oehlers lab at Centenary Institute for their input into this work and

maintenance of the zebrafish aquarium; Dr Angela Kurz and Sydney Cytometry for assistance with

imaging; and A/Prof Jordan Shavit for supplying the *Tg(fabp10a:fgb-gfp)*[mi4001] line.

FUNDING

This work was supported by a Centenary Institute Summer Scholarship to S.M.; University of

Sydney Fellowship [grant number G197581] and NSW Ministry of Health under the NSW Health

Early-Mid Career Fellowships Scheme [grant number H18/31086] to S.H.O.; and the NHMRC

Centre of Research Excellence in Tuberculosis Control (APP1153493) to W.J.B..

REFERENCES

1.   Salazar, J.J., W.J. Ennis, and T.J. Koh, *Diabetes medications: Impact on inflammation and wound*
healing. Journal of diabetes and its complications, 2016. **30**(4): p. 746-752.

2.   Galkowska, H., U. Wojewodzka, and W.L. Olszewski, *Low recruitment of immune cells with*
increased expression of endothelial adhesion molecules in margins of the chronic diabetic foot
ulcers. Wound Repair and Regeneration, 2005. **13**(3): p. 248-254.

3.   Vazzana, N., et al., *Diabetes mellitus and thrombosis.* Thrombosis Research, 2012. **129**(3): p. 371-
377.

4.   Sawaya, A.P., et al., *Deregulated immune cell recruitment orchestrated by FOXM1 impairs human*
diabetic wound healing. Nature communications, 2020. **11**(1): p. 1-14.

5.   Son, S.M., I.J. Kim, and Y.K. Kim, *High Glucose Induces Injury of Endothelial Cells by Activated*
Neutrophils: Implication for Inflammation in Vascular Complication, in *Diabetes.* 2000. p. A147.

6.   Lucas, T., et al., *Differential roles of macrophages in diverse phases of skin repair.* Journal of
immunology (Baltimore, Md. : 1950), 2010. **184**(7): p. 3964-3977.

7.   Miriam, A.M.L., et al., *Differences in Cellular Infiltrate and Extracellular Matrix of Chronic Diabetic*
and Venous Ulcers Versus Acute Wounds. Journal of Investigative Dermatology, 1998. **111**(5): p.
850.

8.   Rao, A.K., et al., *Alterations in insulin-signaling and coagulation pathways in platelets during*
hyperglycemia-hyperinsulinemia in healthy non-diabetic subject. Thrombosis Research, 2014.
134(3): p. 704-710.

9.   Tsimerman, G., et al., *Involvement of microparticles in diabetic vascular complications.* Thromb
Haemost, 2011. **106**(2): p. 310-21.

10.   Ding, Y., et al., *Platelet-Rich Fibrin Accelerates Skin Wound Healing in Diabetic Mice.* Annals of
Plastic Surgery, 2017. **79**(3): p. e15-e19.

11.   Babaei, V., et al., *Management of chronic diabetic foot ulcers using platelet-rich plasma.* Journal of
Wound Care, 2017. **26**(12): p. 784-787.

12.   Okon, E.B., et al., *Hyperglycemia and hyperlipidemia are associated with endothelial dysfunction*
during the development of type 2 diabetes. Canadian Journal of Physiology and Pharmacology,
2007. **85**(5): p. 562-567.

13.   Andreadou, I., et al., *Hyperlipidaemia and cardioprotection: Animal models for translational studies.*
British journal of pharmacology, 2019.

14.   Crawford-Faucher, A., *Preventing or Delaying Type 2 Diabetes Mellitus with Diet and Exercise.*
American family physician, 2018. **98**(11): p. 643-644.

15.   Pulakazhi Venu, V.K., et al., *Minimizing Hyperglycemia-Induced Vascular Endothelial Dysfunction by*
Inhibiting Endothelial Sodium-Glucose Cotransporter 2 and Attenuating Oxidative Stress:
Implications for Treating Individuals With Type 2 Diabetes. Canadian Journal of Diabetes, 2019.
43(7): p. 510-514.

16.   Williams, S.B., et al., *Acute Hyperglycemia Attenuates Endothelium-Dependent Vasodilation in*
Humans In Vivo. Circulation, 1998. **97**(17): p. 1695-1701.

17.   Lamharzi, N., et al., *Hyperlipidemia in Concert With Hyperglycemia Stimulates the Proliferation of*
Macrophages in Atherosclerotic Lesions. Diabetes, 2004. **53**(12): p. 3217.

18.   Oyelaja-Akinsipo, O.B., E.O. Dare, and D.P. Katare, *Protective role of diosgenin against*
hyperglycaemia-mediated cerebral ischemic brain injury in zebrafish model of type II diabetes
mellitus. Heliyon, 2020. **6**(1).

19.   Richardson, R., et al., *Adult Zebrafish as a Model System for Cutaneous Wound-Healing Research.*
Journal of Investigative Dermatology, 2013. **133**(6): p. 1655-1665.

20.   Martin, P. and Y. Feng, *Wound healing in zebrafish.* Nature, 2009. **459**(7249): p. 921-923.

295	21.	Fang, L., C. Liu, and Y.I. Miller, *Zebrafish models of dyslipidemia: relevance to atherosclerosis and*
296		*angiogenesis.* Translational Research, 2014. **163**(2): p. 99-108.
297	22.	Kinkel, M.D. and V.E. Prince, *On the diabetic menu: Zebrafish as a model for pancreas development*
298		*and function.* 2009, WILEY-VCH Verlag: Weinheim. p. 139-152.
299	23.	Cholan, P.M., et al., *Conserved anti-inflammatory effects and sensing of butyrate in zebrafish.* Gut
300		Microbes, 2020. **12**(1): p. 1-11.
301	24.	Passeri, M.J., et al., *Hepatic steatosis in response to acute alcohol exposure in zebrafish requires*
302		*sterol regulatory element binding protein activation.* Hepatology, 2009. **49**(2): p. 443-452.
303	25.	Johansen, M.D., et al., *Mycobacterium marinum infection drives foam cell differentiation in*
304		*zebrafish infection models.* Developmental & Comparative Immunology, 2018. **88**: p. 169-172.
305	26.	Matty, M.A., S.H. Oehlers, and D.M. Tobin, *Live Imaging of Host-Pathogen Interactions in Zebrafish*
306		*Larvae.* Methods Mol Biol, 2016. **1451**: p. 207-23.
307	27.	Rocha, F., et al., *Glucose overload in yolk has little effect on the long-term modulation of*
308		*carbohydrate metabolic genes in zebrafish (Danio rerio).* The Journal of Experimental
309		Biology, 2014. **217**(7): p. 1139-1149.
310	28.	Wang, Z., et al., *Impact of a Combined High Cholesterol Diet and High Glucose Environment on*
311		*Vasculature.* PLOS ONE, 2013. **8**(12): p. e81485.
312	29.	Delamaire, M., et al., *Impaired Leucocyte Functions in Diabetic Patients.* Diabetic Medicine, 1997.
313		**14**(1): p. 29-34.
314	30.	Wong, S.L., et al., *Diabetes primes neutrophils to undergo NETosis, which impairs wound healing.*
315		Nature medicine, 2015. **21**(7): p. 815-819.
316	31.	Zykova, S.N., et al., *Altered cytokine and nitric oxide secretion in vitro by macrophages from diabetic*
317		*type II-like db/db mice.* Diabetes, 2000. **49**(9): p. 1451.
318	32.	Bagamery, K., et al., *Flow cytometric analysis of CD41-labeled platelets isolated by the rapid,*
319		*one-step OptiPrep method from human blood.* Cytometry Part A, 2005. **65**(1): p. 84-87.
320	33.	Huarng, M.C. and J.A. Shavit, *Simple and rapid quantification of thrombocytes in zebrafish larvae.*
321		Zebrafish, 2015. **12**(3): p. 238-242.
322	34.	Vo, A.H., et al., *Loss of fibrinogen in zebrafish results in symptoms consistent with human*
323		*hypofibrinogenemia.* PloS one, 2013. **8**(9): p. e74682-e74682.
324	35.	Goldberg, I.J., *Diabetic Dyslipidemia: Causes and Consequences.* The Journal of Clinical
325		Endocrinology & Metabolism, 2001. **86**(3): p. 965-971.
326	36.	Barros-Becker, F., et al., *Live imaging reveals distinct modes of neutrophil and macrophage*
327		*migration within interstitial tissues.* Journal of Cell Science, 2017. **130**(22): p. 3801.
328	37.	Alba-Loureiro, T.C., et al., *Diabetes causes marked changes in function and metabolism of rat*
329		*neutrophils.* J Endocrinol, 2006. **188**(2): p. 295-303.
330	38.	Chait, A. and K.E. Bornfeldt, *Diabetes and atherosclerosis: is there a role for hyperglycemia?* Journal
331		of lipid research, 2009. **50 Suppl**(Suppl): p. S335-S339.

332