Glucose inhibits haemostasis and accelerates diet-induced hyperlipidaemia in zebrafish larvae

Simone Morris[1], Pradeep Manuneedhi Cholan[1]†, Warwick J Britton[1,2], Stefan H Oehlers[1,3] *

[1] Tuberculosis Research Program at the Centenary Institute, The University of Sydney,

Camperdown NSW 2050 Australia

[2] Department of Clinical Immunology, Royal Prince Alfred Hospital, Camperdown, NSW 2050,

Australia

[3] The University of Sydney, Discipline of Infectious Diseases & Immunology and Marie Bashir

Institute, Camperdown NSW 2050 Australia

† Co-corresponding author email: pradeep.manuneedhicholan@mq.edu.au

* Corresponding author email: s.oehlers@centenary.org.au

ABSTRACT

Hyperglycaemia damages the microvasculature in part through the reduced recruitment of immune

cells and interference with platelet signalling, leading to poor wound healing and accelerated lipid

deposition in mammals. We investigated the utility of zebrafish larvae to model the effect of

glucose on leukocyte recruitment, thrombosis, and hyperlipidaemia by treating larvae with

exogenous glucose. Exogenous glucose supplementation reduced thrombocyte and fibrin plug

formation following tail transection in zebrafish larvae but did not affect the recruitment of

neutrophils and macrophages. We observed accelerated lipid accumulation in glucose-supplemented

larvae challenged with high fat chicken egg yolk feeding. Our study identifies conserved and

divergent effects of high glucose on inflammation in the tractable zebrafish embryo model system

compared to mammals.

INTRODUCTION

Hyperglycaemic damage to the microvasculature is hypothesised to underpin much of the pathology

associated with diabetes in mammals, including perturbations to leukocyte biology, haemostasis,

and the accumulation of lipid laden macrophages in the vessel wall (1-4). Previous mammalian

research has demonstrated that hyperglycaemia damages the microvasculature resulting in reduced

expression of endothelial adhesion molecules for immune cell recruitment (2, 4). As a result, fewer

neutrophils and macrophages are recruited to diabetic wounds (1, 2, 5). Of the macrophages and

neutrophils that eventually arrive, there is a skewing of differentiation towards an inflammatory

phenotype owing to the inflammatory nature of the diabetic wound microenvironment (6, 7).

Mammals with hyperglycaemia demonstrate perturbed coagulation and platelet signalling, causing

disruption of haemostasis (3, 8). Reduced efficiency of haemostasis results in unstable and

ineffective clots within diabetic foot ulcers (3, 9). Treatments for diabetic foot ulcers can involve

platelet and fibrin therapy, indicating an important role for inadequate fibrin clot production in the

ulceration process (10, 11).

Hyperglycaemia-induced damage to the microvasculature also increases vascular lipid

accumulation in conjunction with hyperlipidaemia (12). Hyperlipidaemia and hyperglycaemia are

compounding risk factors for the development of Type 2 diabetes, associated with the 'Western

Diet' consisting of fat and sugar alongside limited exercise (13, 14). There are multiple mechanisms

by which hyperglycaemia damages the microvasculature: through the formation of advanced

glycation products (3), the induction of oxidative stress (15), interfering with nitric oxide

production (16), and inducing macrophages to form lipid-laden foam cells (17). The degradation of

the endothelial structural integrity increases the rate of lipid deposition by providing a physical

niche for lipid infiltration (12). This can ultimately lead to the development of atheroma, which are

common in diabetic patients (17).

These phenotypes have not been previously investigated in fish. Our study investigates the effect of

high glucose environments on inflammation, thrombosis and hyperlipidaemia using a zebrafish

model. Zebrafish are established models for investigating each of these processes in isolation (18-

21). Pertinent to this study, zebrafish have similar clotting, metabolic, and immune systems to

mammals, and these have been used to provide insight into the shared function of these systems

across vertebrates (19, 20, 22). Here we have combined these models to investigate the role of high

glucose environments in disrupting thrombosis and lipid accumulation, but not immune cell

recruitment to a wound, in zebrafish larvae.

METHODS

Zebrafish Husbandry

Zebrafish embryos were produced through natural spawning (Sydney Local Health District Animal

Welfare Committee Approval 17-036). The strains used were *Tg(lyzC:DsRed)*[nz50] to visualise

neutrophils (23), *Tg(mfap4:turquoise)*[xt27] to visualise macrophages (24), *Tg(itga2b:gfp)*[la2] to

visualise thrombocytes (25), and *Tg(fabp10a:fgb-gfp)*[mi4001] to visualise fibrin deposition (26). From

1-5 days post fertilisation (dpf) embryos were kept in a dark incubator at 28°C.

Induction of High Glucose Concentrations in the Zebrafish

Injection method: Eggs were injected with approximately 15 nmol of glucose, or an equal volume

of PBS as a control within four hours of fertilisation. Immersion method: Dechorionated 2 dpf

embryos were immersed in 5% mannitol (osmolarity control) or glucose dissolved in E3 media,

media was changed daily to reduce microbial growth.

Glucose Oxidase Assay

Larvae were snap frozen and stored at -20°C. Larvae were lysed in the buffer solution from the

Amplex Red Oxidase kit (Sigma: A22188), lysates were sedimented by centrifugation, and the

Amplex Red Glucose oxidase kit was used to measure the glucose content of the supernatant in

accordance with the manufacturer's instructions.

Tail Transection Assays

Caudal fin amputations were performed on larvae at 5 dpf. Larvae were anesthetised with 2.5%
(v/v) ethyl-3-aminobenzoate methanesulfonate (tricaine) (Sigma, E10521), wounded posterior to
the notochord using a sterile scalpel and kept in a 28°C incubator to recover as previously described
(27). Wounded larvae were imaged at 6 hours (neutrophil and macrophages) or 2.5 hours (fibrin and
thrombocytes).

Hyperlipidaemia Assay

Post 5 dpf, larvae were transferred to a 28°C incubator with a 14/10 hour light/dark cycle. Larvae
were placed in an E3 solution containing 0.05% of emulsified chicken egg yolk from 5 dpf. Each
day, a random sample of larvae were removed, euthanised, and fixed in paraformaldehyde. Larvae
were stained with Oil Red O to quantitate lipid accumulation, as previously described (21, 28, 29).

Imaging and Image Analysis

Larvae were imaged using a Leica M205FA fluorescent microscope. ImageJ software was used to
quantify fluorescent pixel count within 100 µm of the wound site for transgenic wound assays as
previously described (27, 30).

Oil Red O staining was quantified in ImageJ by adjusting the colour threshold to eliminate non-red
signal. The image was then converted to a binary mask, and the tail region posterior to the swim
bladder was selected to measure the number and area of particles (29).

Statistical analysis

Outliers were excluded using a 1% ROUT test. Statistical testing was carried out by ANOVA with

correction for multiple comparisons or Student's *t*-tests as appropriate using GraphPad Prism. Data

are expressed as mean ± SD. Every datapoint represents a single embryo unless otherwise noted.

RESULTS

Exogenous glucose exposure increases glucose in zebrafish larvae

To establish the efficacy of the injection and immersion techniques to increase glucose

concentrations in 5 dpf zebrafish larvae, we conducted a glucose oxidase assay. Consistent with past

literature, we observed an increase in the glucose concentration contained within the glucose-

injected and -immersed larvae compared to controls (Figure 1A and 1B) (31, 32).

Glucose immersion caused sporadic microbial overgrowth and we observed reduced growth of

larvae immersed in either glucose or mannitol as measured by total body area (Figure 1C) or eye

area (Figure 1D). These changes were not observed in glucose-injected larvae.

We next analysed the effect of glucose injection on the development of key blood cells. We

estimated the quantity of macrophages in transgenic *Tg(mfap4:turquoise)^{xt27}* larvae, where

macrophages are marked by turquoise fluorescent protein, and found similar numbers of

macrophages (Figure 1E). We estimated the quantity of neutrophils in *Tg(lyzC:DsRed)^{nz50}* larvae,

where neutrophils are marked by DsRed fluorescent protein, and found reduced numbers of

neutrophils (Figure 1F). Finally, we estimated the quantity of thrombocytes in *Tg(itga2b:gfp)^{la2}*

larvae, where thrombocytes are tagged with GFP, and found similar numbers of thrombocytes

(Figure 1G).

Glucose does not affect neutrophil and macrophage recruitment to wounds in zebrafish

larvae.

Altered innate immune cell recruitment to wounds is a conserved feature of hyperglycaemia in

mammals (4, 5, 33-35). To determine if this phenomenon was conserved in zebrafish larvae, we

utilised the tail transection wound model which causes reproducible leukocyte recruitment (27)

(Figure 2A). We first performed this assay using transgenic *Tg(mfap4:turquoise)*[xt27] larvae to

quantify macrophage recruitment to the tail wound (Figure 2B). Surprisingly, we observed no

difference in macrophage recruitment between the glucose-injected and control larvae at 6 hours

post wounding (hpw) (Figure 2C). We also observed no difference in macrophage recruitment in

the glucose immersion model (Figure 2D).

We then used *Tg(lyzC:DsRed)*[nz50] larvae to quantify the recruitment of neutrophils to the tail wound

(Figure 2E). We observed great variability between experiments with two out of four experiments

finding significantly reduced neutrophil recruitment between the control and glucose-injected larvae

at 6 hpw and two out of four experiments finding significantly increased neutrophil recruitment

(Figure 2F). Consistently, we did not observe any difference in neutrophil recruitment in the

glucose immersion model (Figure 2G).

Together, these results indicate that zebrafish neutrophil and macrophage recruitment is not affected

by exogenous glucose supplementation in zebrafish larvae.

Glucose impedes haemostasis in zebrafish larvae

Hyperglycaemia perturbs coagulation and platelet activation in mammals, resulting in ineffective

haemostasis (3, 8). To visualise the effects of high glucose on haemostasis in zebrafish larvae, we

first used the *Tg(itga2b:gfp)*[la2] line to visualise thrombocyte plug formation in the severed blood

vessel (25, 36). Following tail transection (Figure 3A), glucose-injected *Tg(itga2b:gfp)*[la2] larvae

demonstrated reduced thrombocyte accumulation (Figure 3B-C). This effect was replicated in

glucose-immersed embryos compared to control embryos (Figure 3D).

To determine if the reduction in thrombocyte recruitment was mirrored by perturbed clotting, we

conducted tail transections on *Tg(fabp10a:fgb-gfp)*^{mi4001} larvae (Figure 3E), which express

fluorescently tagged fibrinogen and allow visualisation of clots (26). Glucose-injected

Tg(fabp10a:fgb-gfp)^{mi4001} larvae had reduced fibrin accumulation following tail transection

compared to PBS-injected larvae (Figure 3F). We did not observe any effect of glucose immersion

on the deposition of fluorescently tagged fibrinogen (Figure 3G).

Together, these results demonstrate that a high glucose environment inhibits the thrombocyte

component of haemostasis in zebrafish larvae.

High glucose accelerated hyperlipidaemia in zebrafish larvae challenged with a high fat diet

Hyperglycaemia and hyperlipidaemia are intimately associated in mammals (37). To determine if

this interaction is conserved in zebrafish larvae, we fed glucose-injected larvae a high fat diet

consisting of emulsified chicken egg yolk from 5-7 dpf (Figure 4A). Glucose- and PBS-injected

larvae had similar Oil Red O vascular staining prior to the initiation of chicken egg yolk feeding at

5 dpf (Figure 4B-C). Quantification of Oil Red O staining revealed glucose-injected larvae had

increased vascular lipid content at one and two days post feeding (Figure 4D-E).

These results recapitulate the interaction between hyperglycaemia and hyperlipidaemia seen in

mammals (15), demonstrating that zebrafish larvae are a robust model to study this conserved

interaction.

DISCUSSION

In this study, we explored the effect of high glucose levels on inflammation, thrombosis, and lipid

accumulation in the zebrafish embryo model. We found that high glucose reduces haemostasis

around wound sites. In addition, we determined that high glucose accelerates lipid accumulation

when larvae are challenged with a high fat diet.

The reduced size of glucose and mannitol immersed larvae compared to control larvae is a major

caveat when interpreting the datasets generated with the immersion model. We thus preferred the

injection method for the chicken egg yolk challenge experiments and for the interpretation of the

effects of glucose on larval zebrafish immunity and haemostasis. Despite daily changes of media,

we experienced further difficulty rearing zebrafish larvae from 2 to 5 dpf in 5% glucose solutions

due to microbial overgrowth which may have slowed larval development by consuming oxygen or

damaging larval mucosal surfaces. Although the glucose injection model did not achieve as high a

fold change in glucose at 5 dpf, we did not observe developmental delays and haematopoiesis was

largely comparable to PBS-injected control larvae.

Hyperglycaemia in mammals causes vascular dysfunction that restricts the recruitment of a broad

range of immune cells to wounds (16). It was therefore surprising that high glucose appeared to

have no effect on the recruitment of leukocytes to the tail wound in this study. It is possible that

high glucose in zebrafish larvae does not affect abluminal crawling of leukocytes along blood

vessels, since wound-responsive leukocytes have been demonstrated to move predominantly

through interstitial tissue in zebrafish larvae (5, 38). Overall our findings suggest exogenous supply

of glucose to zebrafish larvae may not be a suitable platform for studying the impact of high

glucose on leukocyte biology.

The relationship between hyperglycaemia and hyperlipidaemia has been previously reported in

various mammalian species (15, 32, 39), and a recent study by Wang *et al.* has demonstrated a

similar the interaction of a high cholesterol diet with glucose immersion on vascular lipid

accumulation in zebrafish after 10 days of feeding (32). Our study using chicken egg yolk as a high

fat diet challenge demonstrates a dramatically accelerated accumulation of lipid after just one day of

feeding. This chicken egg yolk feeding-based model, therefore provides a more rapid model to

investigate lipid accumulation.

In summary, we report glucose-supplemented zebrafish larvae as a tractable platform to investigate

the conserved interactions between glucose and haemostasis, and glucose and diet-induced

hyperlipidaemia.

ACKNOWLEDGEMENTS

We thank other members of the Oehlers lab at the Centenary Institute for their input into this work

and maintenance of the zebrafish aquarium; Dr Angela Kurz and Sydney Cytometry for assistance

with imaging; and A/Prof Jordan Shavit for supplying the $Tg(fabp10a:fgb\text{-}gfp)^{mi4001}$ line.

FUNDING

This work was supported by a Centenary Institute Summer Scholarship to S.M.; University of

Sydney Fellowship [grant number G197581] and NSW Ministry of Health under the NSW Health

Early-Mid Career Fellowships Scheme [grant number H18/31086] to S.H.O.; and the NHMRC

Centre of Research Excellence in Tuberculosis Control (APP1153493) to W.J.B..

REFERENCES

1.   Salazar JJ, Ennis WJ, Koh TJ. Diabetes medications: Impact on inflammation and wound healing. J
Diabetes Complications. 2016;30(4):746-52.
2.   Galkowska H, Wojewodzka U, Olszewski WL. Low recruitment of immune cells with increased
expression of endothelial adhesion molecules in margins of the chronic diabetic foot ulcers. Wound Repair
and Regeneration. 2005;13(3):248-54.
3.   Vazzana N, Ranalli P, Cuccurullo C, Davì G. Diabetes mellitus and thrombosis. Thrombosis Research.
2012;129(3):371-7.
4.   Sawaya AP, Stone RC, Brooks SR, Pastar I, Jozic I, Hasneen K, et al. Deregulated immune cell
recruitment orchestrated by FOXM1 impairs human diabetic wound healing. Nature communications.
2020;11(1):1-14.
5.   Son SM, Kim IJ, Kim YK. High Glucose Induces Injury of Endothelial Cells by Activated Neutrophils:
Implication for Inflammation in Vascular Complication. Diabetes. 2000 2000/05//:A147.

6.   Lucas T, Waisman A, Ranjan R, Roes J, Krieg T, Müller W, et al. Differential roles of macrophages in
diverse phases of skin repair. Journal of immunology (Baltimore, Md : 1950). 2010;184(7):3964-77.
7.   Miriam AML, Evert NL, Jimmy Z, Jan RM, Jan DB, Esther M. Differences in Cellular Infiltrate and
Extracellular Matrix of Chronic Diabetic and Venous Ulcers Versus Acute Wounds. Journal of Investigative
Dermatology. 1998;111(5):850.
8.   Rao AK, Freishtat RJ, Jalagadugula G, Singh A, Mao G, Wiles A, et al. Alterations in insulin-signaling
and coagulation pathways in platelets during hyperglycemia-hyperinsulinemia in healthy non-diabetic
subject. Thrombosis Research. 2014;134(3):704-10.
9.   Tsimerman G, Roguin A, Bachar A, Melamed E, Brenner B, Aharon A. Involvement of microparticles
in diabetic vascular complications. Thromb Haemost. 2011;106(2):310-21.
10.   Ding Y, Cui L, Zhao Q, Zhang W, Sun H, Zheng L. Platelet-Rich Fibrin Accelerates Skin Wound Healing
in Diabetic Mice. Annals of Plastic Surgery. 2017;79(3):e15-e9.
11.   Babaei V, Afradi H, Gohardani HZ, Nasseri F, Azarafza M, Teimourian S. Management of chronic
diabetic foot ulcers using platelet-rich plasma. Journal of Wound Care. 2017;26(12):784-7.
12.   Okon EB, Chung AWY, Zhang H, Laher I, van Breemen C. Hyperglycemia and hyperlipidemia are
associated with endothelial dysfunction during the development of type 2 diabetes. Canadian Journal of
Physiology and Pharmacology. 2007;85(5):562-7.
13.   Andreadou I, Schulz R, Badimon L, Adameová A, Kleinbongard P, Lecour S, et al. Hyperlipidaemia
and cardioprotection: Animal models for translational studies. British journal of pharmacology. 2019.
14.   Crawford-Faucher A. Preventing or Delaying Type 2 Diabetes Mellitus with Diet and Exercise.
American family physician. 2018;98(11):643-4.
15.   Pulakazhi Venu VK, El-Daly M, Saifeddine M, Hirota SA, Ding H, Triggle CR, et al. Minimizing
Hyperglycemia-Induced Vascular Endothelial Dysfunction by Inhibiting Endothelial Sodium-Glucose
Cotransporter 2 and Attenuating Oxidative Stress: Implications for Treating Individuals With Type 2
Diabetes. Canadian Journal of Diabetes. 2019;43(7):510-4.
16.   Williams SB, Goldfine AB, Timimi FK, Ting HH, Roddy M-A, Simonson DC, et al. Acute Hyperglycemia
Attenuates Endothelium-Dependent Vasodilation in Humans In Vivo. Circulation. 1998;97(17):1695-701.
17.   Lamharzi N, Renard CB, Kramer F, Pennathur S, Heinecke JW, Chait A, et al. Hyperlipidemia in
Concert With Hyperglycemia Stimulates the Proliferation of Macrophages in Atherosclerotic Lesions.
Diabetes. 2004;53(12):3217.
18.   Oyelaja-Akinsipo OB, Dare EO, Katare DP. Protective role of diosgenin against hyperglycaemia-
mediated cerebral ischemic brain injury in zebrafish model of type II diabetes mellitus. Heliyon. 2020;6(1).
19.   Richardson R, Slanchev K, Kraus C, Knyphausen P, Eming S, Hammerschmidt M. Adult Zebrafish as a
Model System for Cutaneous Wound-Healing Research. Journal of Investigative Dermatology.
2013;133(6):1655-65.
20.   Martin P, Feng Y. Wound healing in zebrafish. Nature. 2009;459(7249):921-3.
21.   Fang L, Liu C, Miller YI. Zebrafish models of dyslipidemia: relevance to atherosclerosis and
angiogenesis. Translational Research. 2014;163(2):99-108.
22.   Kinkel MD, Prince VE. On the diabetic menu: Zebrafish as a model for pancreas development and
function. Weinheim: WILEY-VCH Verlag; 2009. p. 139-52.
23.   Hall C, Flores MV, Storm T, Crosier K, Crosier P. The zebrafish lysozyme C promoter drives myeloid-
specific expression in transgenic fish. BMC Dev Biol. 2007;7(1):42.
24.   Walton EM, Cronan MR, Beerman RW, Tobin DM. The Macrophage-Specific Promoter mfap4 Allows
Live, Long-Term Analysis of Macrophage Behavior during Mycobacterial Infection in Zebrafish. PLoS ONE.
2015;10(10):e0138949.
25.   Lin HF, Traver D, Zhu H, Dooley K, Paw BH, Zon LI, et al. Analysis of thrombocyte development in
CD41-GFP transgenic zebrafish. Blood. 2005;106(12):3803-10.
26.   Vo AH, Swaroop A, Liu Y, Norris ZG, Shavit JA. Loss of fibrinogen in zebrafish results in symptoms
consistent with human hypofibrinogenemia. PLoS ONE. 2013;8(9):e74682.
27.   Cholan PM, Han A, Woodie BR, Watchon M, Kurz AR, Laird AS, et al. Conserved anti-inflammatory
effects and sensing of butyrate in zebrafish. Gut Microbes. 2020;12(1):1-11.
28.   Passeri MJ, Cinaroglu A, Gao C, Sadler KC. Hepatic steatosis in response to acute alcohol exposure
in zebrafish requires sterol regulatory element binding protein activation. Hepatology. 2009;49(2):443-52.

29.   Johansen MD, Kasparian JA, Hortle E, Britton WJ, Purdie AC, Oehlers SH. Mycobacterium marinum
infection drives foam cell differentiation in zebrafish infection models. Developmental & Comparative
Immunology. 2018;88:169-72.
30.   Matty MA, Oehlers SH, Tobin DM. Live Imaging of Host-Pathogen Interactions in Zebrafish Larvae.
Methods Mol Biol. 2016;1451:207-23.
31.   Rocha F, Dias J, Engrola S, Gavaia P, Geurden I, Dinis MT, et al. Glucose overload in yolk has little
effect on the long-term modulation of carbohydrate metabolic genes in zebrafish (Danio rerio).
The Journal of Experimental Biology. 2014;217(7):1139-49.
32.   Wang Z, Mao Y, Cui T, Tang D, Wang XL. Impact of a Combined High Cholesterol Diet and High
Glucose Environment on Vasculature. PLoS ONE. 2013;8(12):e81485.
33.   Delamaire M, Maugendre D, Moreno M, Le Goff MC, Allannic H, Genetet B. Impaired Leucocyte
Functions in Diabetic Patients. Diabetic Medicine. 1997;14(1):29-34.
34.   Wong SL, Demers M, Martinod K, Gallant M, Wang Y, Goldfine AB, et al. Diabetes primes
neutrophils to undergo NETosis, which impairs wound healing. Nature medicine. 2015;21(7):815-9.
35.   Zykova SN, Jenssen TG, Berdal M, Olsen R, Myklebust R, Seljelid R. Altered cytokine and nitric oxide
secretion in vitro by macrophages from diabetic type II-like db/db mice. Diabetes. 2000;49(9):1451.
36.   Huarng MC, Shavit JA. Simple and rapid quantification of thrombocytes in zebrafish larvae.
Zebrafish. 2015;12(3):238-42.
37.   Goldberg IJ. Diabetic Dyslipidemia: Causes and Consequences. The Journal of Clinical Endocrinology
& Metabolism. 2001;86(3):965-71.
38.   Barros-Becker F, Lam P-Y, Fisher R, Huttenlocher A. Live imaging reveals distinct modes of
neutrophil and macrophage migration within interstitial tissues. Journal of Cell Science. 2017;130(22):3801.
39.   Chait A, Bornfeldt KE. Diabetes and atherosclerosis: is there a role for hyperglycemia? Journal of
lipid research. 2009;50 Suppl(Suppl):S335-S9.

Figure legends

Figure 1: Injection and immersion methods increase glucose levels in zebrafish larvae.

A. Relative concentration of glucose in 5 dpf larvae that had been injected with 15 nmol glucose as

eggs. Statistical testing by t-test, each data point is representative of a group of n = 10-30 larvae.

B. Relative concentration of glucose in 5 dpf larvae immersed in 5% solutions of glucose or

mannitol from 2 dpf. Statistical testing by ANOVA, each data point is representative of a group of n

= 10-30 larvae.

C. Total body area calculated from lateral images of 5 dpf larvae immersed in 5% solutions of

glucose or mannitol from 2 dpf. Statistical testing by ANOVA. Data are representative of 2

biological replicates.

D. Eye area calculated from lateral images of 5 dpf larvae immersed in 5% solutions of glucose or

mannitol from 2 dpf. Statistical testing by ANOVA. Data are representative of 2 biological

replicates.

E. Quantification of total macrophage number from lateral images of 5 dpf larvae. Statistical testing

by t-test. Data are representative of 2 biological replicates.

F. Quantification of total neutrophil number from lateral images of 5 dpf larvae. Statistical testing

by t-test. Data are representative of 2 biological replicates.

G. Quantification of total thrombocyte number from lateral images of 5 dpf larvae. Statistical

testing by t-test. Data are representative of 2 biological replicates.

Figure 2: High glucose does not affect neutrophil and macrophage recruitment to a tail wound.

A. Schematic of experiment to measure immune cell recruitment to a tail wound.

B. Representative images of macrophage (red) recruitment to a tail wound in glucose-injected

larvae.

C. Quantification of macrophage recruitment following tail transection in the glucose injection

model.

D. Quantification of macrophage recruitment following tail transection in the glucose immersion

model.

E. Representative images of neutrophil (red) recruitment to a tail wound in glucose-injected larvae.

F. Quantification of neutrophil recruitment following tail transection in the glucose injection model.

Each paired data point represents the average of an biological replicate with n>10 embryos per

condition.

G. Quantification of neutrophil recruitment following tail transection in the glucose immersion

model.

Scale bars represent 100 μm. Statistical testing by t-test. Data are representative of 3 biological

replicates.

Figure 3: High glucose reduced thrombocyte and fibrin accumulation at a tail wound.

A. Schematic of experiment to visualise haemostasis following tail transection.

B. Representative overlay of thrombocytes (red) at 2.5 hours after tail transection in glucose-

injected larvae.

C. Quantification of thrombocyte plug size following tail transection in the glucose injection model.

D. Quantification of thrombocyte plug size following tail transection in the glucose immersion

model.

E. Representative images of fibrinogen deposition (red) at 2.5 hours after tail transection in glucose-

injected larvae.

F. Quantification of fibrin clot size following tail transection in the glucose injection model.

G. Quantification of fibrin clot size following tail transection in the glucose immersion model.

Scale bars represent 100 μm. Statistical testing by t-test. Data are representative of 3 biological

replicates.

Figure 4: Glucose-injected larvae have increased lipid accumulation following a high fat diet.

A. Schematic of the high fat feeding challenge assay to measure lipid accumulation in glucose-

injected larvae.

B. Bright field images of 6 dpf Oil Red O-stained larvae, demonstrating darker vascular staining in

glucose-injected larvae. Box indicates location of inset, arrowheads indicate stained intersegmental

vessels in inset, asterisk indicates intestinal lumen which was excluded from analysis.

C. Quantification of lipid accumulation in 5 dpf glucose-injected larvae.

D. Quantification of lipid accumulation in 6 dpf glucose-injected larvae challenged with a high fat

diet from 5 dpf.

E. Quantification of lipid accumulation in 7 dpf glucose-injected larvae challenged with a high fat

diet from 5 dpf.

Statistical testing by t-test. Data are representative of 2 biological replicates.