Title: Methylation of histone H3 lysine 4 is required for maintenance of beta cell function in adult mice

Authors: Ben Vanderkruk,[1,2] Nina Maeshima,[2] Daniel J. Pasula,[1,2] Meilin An,[2] Cassandra L. McDonald,[2]

Francis C. Lynn, [1,2] Dan S. Luciani,[1,2] and Brad G. Hoffman[1,2]*

Affiliations: [1]Diabetes Research Group, BC Children's Hospital Research Institute, Vancouver, BC, Canada

[2]Department of Surgery, University of British Columbia, Vancouver, BC, Canada

*Correspondence. Email: brad.hoffman@ubc.ca

Keywords: H3K4me3; DPY30; COMPASS; Transcription; Beta cell; Type 2 diabetes

Abstract

Abstract

Pancreatic β-cells control glucose homeostasis via regulated production and secretion of insulin. This

function arises from a highly specialized gene expression program which is established during

development and then sustained, with limited flexibility, in terminally differentiated β-cells. Dysregulation

of this program is seen in type 2 diabetes (T2D) but mechanisms that preserve gene expression or underlie

its dysregulation in mature β-cells are not well resolved. Here we show that trithorax group-dependent

histone H3 lysine 4 trimethylation (H3K4me3) maintains expression of genes important for insulin

biosynthesis and glucose-responsiveness in β-cells. Transcriptional changes in H3K4me3-deficient β-cells

lead to severe hyperglycemia in adult mice. We show that H3K4me3 deficiency leads to a less active and

more repressed epigenome profile, which locally correlates with gene expression deficits but does not

globally reduce gene expression. Instead, developmentally regulated genes and genes in weakly active or

suppressed states particularly rely on H3K4 methylation. We then show that H3K4me3 is re-organized in

diabetic *Lepr^db/db* mouse islets in favour of weakly active and disallowed genes at the expense of terminal

β-cell markers with broad H3K4me3 peaks. Our results point to key roles of H3K4me3 in maintaining

mature β-cell function and establishing a dysfunctional transcriptome in diabetic islets.

Introduction

Pancreatic islets are endocrine micro-organs that regulate glucose homeostasis. In adults, islet β-cells are

the exclusive source of circulating insulin, the major blood glucose-lowering hormone. Insufficient insulin

release by β-cells is the primary etiology of type 2 diabetes (T2D). The rising global incidence of T2D [1] has

prompted intensive research into mechanisms underlying the progressive dysfunction of β-cells that leads

to T2D. Transcriptome remodeling, involving activation of developmentally silenced genes and decreased

expression of genes important for insulin biosynthesis, glucose-responsiveness, or those that enforce the

β-cell identity, emerges in β-cells exposed to chronic metabolic stress [2]. While β-cell-enriched

transcription factors have been the primary focus of studies defining the transcriptional program in

mature β-cells, there is mounting evidence that chromatin modifications play a central role in the

maintenance of mature β-cell identity and function, and that the chromatin landscape is reshaped in β-

cells exposed to chronic metabolic stress and T2D [3–10].

Tissue-specific transcription programs rely on precise activation and maintenance of specific

genes and stable repression of other genes. Expression patterns are established during cell differentiation

and maturation, and then maintained during the balance of that cell's life, which may span decades, cell

divisions, and environmental perturbations. Active and repressed genes have stereotypical patterns of

chromatin modification which may stabilize their transcription state and are useful in their systematic

classification [11]. For example, it has been shown that H3K4me3 is reliably enriched on nucleosomes at the

5' end of actively transcribed genes and is therefore the canonical marker of active promoters [12]. Despite

its utility as a marker of promoter activity, global (i.e., genome-wide) decreases in H3K4 methylation are

remarkably well tolerated in embryonic stem cells and during development in genetic knockout models

[13–16]. Evidence in multicellular animal models points to a requirement for H3K4 methyltransferase

activities during certain specialized developmental events, whereas evidence for continued requirement

in mature tissues is rare [16–21].

Mono-, di-, and trimethylation of H3K4 in mammalian cells is predominantly catalyzed by

COMPASS and COMPASS-related complexes. They are composed of the core subunits WDR5, RBBP5,

ASH2L, and DPY30, plus one of six catalytic subunits MLL1-4 or SET1A-B, and a variable retinue of
accessory factors [22] . We previously defined roles for DPY30 during embryonic pancreas development.
DPY30 fine-tunes cell fate decisions to decrease endocrine acinar cell specification in favour of exocrine
cell specification [21] and helps to activate terminal marker genes during endocrine cell maturation [23]. By
contrast, it is not known whether the continuous activity of H3K4 methyltransferases is necessary in
terminally differentiated islet cells, where lineage-specific transcription programs have already been
established. In islets, recruitment of H3K4 methyltransferase proteins by transcription factors—for
example, recruitment of SETD7 by PDX1 [7], and MLL3/4 by MAFA [24]—support expression of insulin and
other genes that are important for mature β-cell functions. Whether these functions are performed by
H3K4 methylation or by independent coactivator activities of these complexes is not clear [14,21,25]. We
therefore sought to define the role for H3K4 methylation in the regulation of gene expression in mature
β-cells.
67
Results
Reduction to H3K4 methylation in beta cells of adult mice leads to glucose intolerance and
hyperglycemia.
With the aim of reducing H3K4 methylation we targeted DPY30, a noncatalytic core subunit of all
COMPASS complexes which is not required for complex assembly or coactivator activity but boosts
methyltransferase activities [26,27]. To induce synchronized deletion of *Dpy30* in mature β-cells, mice
harbouring a Pdx1:*CreER^TM* transgene [28] and *LoxP* sites flanking the fourth exon of *Dpy30* [21] were
administered tamoxifen at eight weeks of age (herein called *Dpy30*-KO) (Fig 1A). As expected, knockout
of *Dpy30* did not impact assembly of the other core subunits of the COMPASS complex WDR5, RBBP5, and
ASH2L, or their association with chromatin (Fig 1B) but did lead to reductions of H3K4me3 and H3K4me1
in islets (Fig 1C).
79
We tracked the loss of H3K4 methylation in *Dpy30*-KO β-cells using immunofluorescence at 15-
day intervals after tamoxifen administration. As seen in Fig 1D-E, H3K4me3 was stable for at least 15 days
and then gradually lost during the subsequent 45 days. H3K4me1 showed a similar pattern, but its loss
was even more gradual, whereas DPY30 was undetectable by 15 days after tamoxifen administration (Fig
1D). We took advantage of the slow turnover of H3K4me3 and H3K4me1 to parse the relative contribution
of H3K4 methylation, versus DPY30 itself, in controlling gene expression in mature β-cells. We reasoned
that defects arising in *Dpy30*-KO β-cells by 15-days post-tamoxifen (when DPY30 is lost but H3K4
methylation is unchanged) highlight functions of DPY30 itself. Defects arising later, by 45-days post-
tamoxifen (when H3K4 methylation is reduced) highlight functional contributions of H3K4 methylation.
To measure the β-cell transcriptome, GFP-positive islet cells labeled with the mTmG reporter [29] were
enriched using fluorescence-activated cell sorting (FACS) from *Dpy30*-KO and -WT mice 15- and 45-days
post-tamoxifen and processed for calibrated bulk RNA-seq analysis [30]. Remarkably, loss of DPY30 had little
effect on gene expression in β-cells, with only three genes (*Dpy30*, *Edn3*, *C3*) identified as differentially
expressed 15-days post-tamoxifen (Fig 1F). In contrast, 828 genes were dysregulated 45-days post-
tamoxifen, the majority of which (634) showed lower expression in *Dpy30*-KO cells (Fig 1F). We conclude
that transcriptome remodeling in *Dpy30*-KO β-cells results from loss of H3K4 methylation.
96
To determine whether loss of DPY30 and/or H3K4 methylation in β-cells impairs glucose
homeostasis in adult mice, we measured glucose clearance and insulin levels after intraperitoneal

injection of glucose (2 g/kg body weight) in mice that had been fasted for 6 hrs. While glucose tolerance
was no different between *Dpy30*-WT and -KO mice 15- and 30-days after tamoxifen administration,
Dpy30-KO mice developed impaired blood glucose clearance and reduced serum insulin levels 45-days
post-tamoxifen (Fig 1G). Glucose intolerance quickly progressed to hyperglycemia (Fig 1H) prompting
euthanasia by 60-days.
104
Since the Pdx1:*CreER*[TM] transgene is known to also drive recombination in pancreatic islet δ-cells
[31] and hypothalamic neurons [32] we additionally used the *Ins1*[Cre] mouse model [33] to confirm that *Dpy30*
deletion specifically in β-cells is sufficient to drive the *in vivo* phenotype. Deletion of *Dpy30* from maturing
β-cells using *Ins1*[Cre] caused reduction of H3K4me3 and H3K4me1 in islets by 5-weeks of age (Fig S1A-B).
Unfasted blood glucose was dramatically elevated from 4- to 5-weeks of age, whereas body mass was
similar among *Dpy30*-KO, -WT, and -heterozygous *Ins1*[Cre] mice (Fig S1C-D). *Ins1*[Cre] *Dpy30*-KO mice showed
impaired glucose clearance and low serum insulin following intraperitoneal glucose injection (Fig S1E-F),
confirming that β-cell-intrinsic dysfunction is sufficient to drive metabolic defects in *Dpy30*-KO mice. In
summary, reduction of H3K4 methylation in β-cells leads to impaired glucose tolerance, reduced serum
insulin, and hyperglycemia.
115
H3K4me3 maintains expression of genes involved in insulin production and glucose-stimulated activity.
The reduction of serum insulin in *Dpy30*-KO mice prompted us to examine insulin production and
secretion. We first confirmed that H3K4me3 was lost from the *Ins1* and *Ins2* promoters in Pdx1:*CreER*[TM]
Dpy30-KO β-cells (Fig 2A) using calibrated chromatin immunoprecipitation sequencing (ChIP-seq) from
FACS-enriched GFP-positive β-cells 45 days after tamoxifen administration. H3K4me1 also showed
dramatic reduction (Fig 2A). Despite this, RNA-seq data showed a surprisingly modest reduction of *Ins1*
and *Ins2* transcripts (Fig 2B). Therefore, robust expression of *Ins1* and *Ins2* RNA can occur in the absence
of local H3K4me3. Notably, expression of several genes involved in insulin peptide maturation and
packaging was downregulated (Fig S2A). For example *Nnat* and *Spcs3*, subunits of the signal peptidase
complex which co-translationally converts preproinsulin to proinsulin [34], showed reduced expression in
Dpy30-KO β-cells (Fig S2A-B). Proinsulin-to-insulin convertases *Pcsk1*, *Pcsk2*, and *Cpe* were not
downregulated (Fig S2A). *Slc30a8*, encoding a zinc transporter required for maturation of insulin secretory
granules [35] was highly downregulated in *Dpy30*-KO cells (Fig S2A). Insulin peptide content was reduced
much more strongly than insulin RNA in *Dpy30*-KO islets (Fig 2B-C). Reduction of insulin content was
confirmed in individual β-cells, where significant reduction to insulin granule size and density was seen in
transmission electron micrographs of β-cells from *Dpy30*-KO mice 45 days after tamoxifen administration
(Fig 2D-E). We conclude that loss of H3K4 methylation from mature β-cells impairs insulin biosynthesis.
133
Gene ontology analysis of genes showing reduced expression suggested the additional
impairment of biological processes required for stimulated insulin secretion, namely vesicle fusion,
calcium-dependent exocytosis, and glucose homeostasis (Fig 2F). Energy production processes including
oxidation/reduction and the electron transport chain, meanwhile, were upregulated (Fig 2F). We
therefore examined *ex vivo* islet activity during glucose stimulation. First, we measured insulin secretion
as a fraction of insulin content to control for the 3.4-fold reduction on insulin in *Dpy30*-KO islets. Notably,
the reduced insulin content of *Dpy30*-KO islets did not fully account for the impairment of insulin
secretion. The persistent impairment was specific to high glucose stimulation, whereas secretion under
nonstimulatory conditions, or bypassing glucose metabolism by depolarizing cells directly using KCl,

showed no difference in insulin secretion between *Dpy30*-WT and -KO islets (after adjusting to total insulin
content) (Fig 2G). Measurement of cytosolic calcium dynamics further supported an impairment of high
glucose stimulated activity:. consistent with lower expression of calcium signaling genes (Fig 2F), high
glucose-stimulated calcium influx showed a delayed response and lower magnitude in *Dpy30*-KO islets
(Fig 2H). Cytosolic calcium concentration was not different in nonstimulatory conditions, and direct
depolarization stimulated comparable calcium influx between islets from *Dpy30*-KO and *Dpy30*-WT mice
(Fig 2H) further supporting a specific impairment in glucose responsiveness. We therefore wondered if
Dpy30-KO impaired glucose catabolism. Expression of *Slc2a2*, encoding the glucose transporter GLUT2,
was dramatically reduced in *Dpy30*-KO cells (Fig S2A-B), but expression of the rate-limiting enzyme of
glycolysis, *Gck* [36], was not significantly altered (Fig S2A-B) nor was the rate of mitochondrial oxygen
consumption in high glucose media altered (Fig 2I-J). In fact, mitochondrial respiratory capacity and area
were increased in *Dpy30*-KO islets (Fig 2I-J, Fig S2C) , consistent with elevated expression of oxidative
phosphorylation genes in *Dpy30*-KO β-cells (Fig 2F). Therefore, glucose metabolism in β-cells was not
appreciably impaired by reduction of H3K4 methylation. These results indicate that H3K4 methylation is
essential for the expression of genes that couple glucose metabolism to insulin secretion in mature β-
cells.
159
Loss of H3K4me3 leads to gene downregulation in mature beta cells.
It is notable that, while H3K4 methylation is well correlated with [37] and predictive of [38] transcription, our
RNA-seq analysis indicates that globally only 4.3% of the 14,677 expressed genes were downregulated by
depletion of H3K4 methylation in β-cells. This mirrors previous findings in yeast [39], *Drosophila* embryos [15],
and mouse embryonic stem cells [13,14,40] that H3K4 methylation is not required for expression of most
genes. We therefore wondered whether gene promoters that are sensitive to H3K4 methylation in β-cells
have generalizable features. We first confirmed that H3K4me3 and H3K4me1 were depleted at active
gene promoters in *Dpy30*-KO chromatin (Fig 3A). We further confirmed genome-wide reduction of
H3K4me3 and H3K4me1 by comparing enrichment between WT and *Dpy30*-KO chromatin in 10-kb bins
spanning the genome (Fig 3B-C), confirming that *Dpy30* is necessary for maintenance of β-cell H3K4
methylation. Despite this, the transcriptomes of *Dpy30*-KO and -WT cells were well correlated, and the
total RNA expression of *Dpy30*-KO cells was not decreased (Fig 3D). Therefore, global depletion of H3K4
methylation does not cause a global decrease in gene expression in mature β-cells.
173
To explore further, we segregated genes into three groups based on residual enrichment for H3K4
methylation, determined using MACS2 [41]. Group (i) retained at least some H3K4me3 enrichment at the
transcription start site (TSS) (10,045 genes), group (ii) lost H3K4me3 but retained at least some H3K4me1
enrichment (2,302 genes), and group (iii) lost H3K4me3 and H3K4me1 (111 genes) (Fig 3E). As a group,
genes that retained at least some enrichment for H3K4me3 at the TSS were not generally downregulated
in *Dpy30*-KO cells (Fig 3F). Loss of both H3K4me3 and H3K4me1 from the TSS was linked to significant
impairment of gene expression, and genes that lost H3K4me3 but retained H3K4me1 showed an
intermediate phenotype (Fig 3F). To visualize this relationship in a threshold-free manner, we plotted the
change in H3K4me3, H3K4me1, and RNA expression for all active TSSs (Fig 3G). The majority of genes
clustered in the bottom left quadrant, indicating some reduction of both H3K4me3 and H3K4me1 from
the TSS. Strikingly, genes that were highly downregulated (red) clustered at the extreme edge of the
distribution, indicating that genes experiencing the greatest relative loss of both H3K4me3 and H3K4me1
were highly downregulated (Fig 3G). Together, these data support a role for H3K4me3 in contributing to

active gene expression: while the genome-wide reduction of H3K4me3 did not influence total cellular
mRNA content, excess loss of H3K4me3 on a per-gene basis leads to downregulation of that gene.
189
H3K4me3 enforces expression of weakly active and repressed genes in mature beta cells.
We then compared the chromatin profile of downregulated and stably expressed genes by measuring
H3K27me3, a marker of developmentally repressed chromatin, and H3K27ac, a marker of active
chromatin, using ChIP-seq. The downregulated gene set had a weakly active profile even in WT cells:
enrichment for H3K4me3 and H3K27ac were low and H3K4me1 and H3K27me3 were high in comparison
to stably expressed genes (Fig 4A-B). Accordingly, the downregulated gene set showed lower absolute
expression in WT cells than stably expressed genes (Fig 4C). The DNA sequence of downregulated genes
also tended to be relatively GC rich (Fig 4E). Imprinted genes, comprising genomic loci for which one allele
is silenced by DNA methylation, tended to be downregulated in *Dpy30*-KO cells (Fig 4F), raising the
possibility that H3K4me3 impedes *de novo* DNA methylation at some genes in mature β-cells, as it does
in mouse embryonic stell cells [42]. To summarize, genes that lose an especially large fraction of H3K4
methylation tend to be downregulated in *Dpy30*-KO cells; they tend to already be in a weakly active state
in WT β-cells and may be more susceptible to silencing in the absence of H3K4 methylation.
203
H3K4me3 deficiency leads to a less active and more repressed epigenome in mature beta cells.
Our finding that H3K4 methylation is especially important for maintaining expression of weakly expressed
genes is intriguing in light of a recent report by Dobrinić *et al.* (ref [43]) that demonstrated Polycomb-
mediated gene repression especially constrains weakly expressed genes [43]. The trithorax group was
originally defined as antagonistic to Polycomb-mediated repressive activity [44]. During development,
repressive (H3K27me3) and active (H3K4me3) states are dynamically regulated and resolution of the
genome into active and repressive domains is fundamental to the establishment of terminal cell-type-
specific transcriptomes [45]. As such, it is notable that expression of genes that are dynamically regulated
during, and important for, β-cell maturation [46] tend to be lower in *Dpy30*-KO cells (Fig 4F). This led us to
ask whether H3K27me3 expands in H3K4me3-deficient β-cells. As shown in Fig 5A, the large-scale
distribution of H3K27me3-positive compartments is unchanged in *Dpy30*-KO chromatin. H3K7me3
enrichment in 10-kb windows across the genome remained similar between *Dpy30*-KO and -WT (Fig 5B),
together indicating that global reduction of H3K4me3 and H3K4me1 did not result in expansion of
Polycomb-repressed compartments. However, H3K27me3 enrichment specifically increased at promoters
of genes that were downregulated in *Dpy30*-KO cells (Fig 5C-E). These data indicate that genome-wide
H3K27me3 organization is largely insensitive to reduction of H3K4 methylation in this model; however,
maintenance of H3K4 methylation is necessary to prevent targeted Polycomb gene repression in mature
β-cells.
222
In contrast to the locus-specific accumulation observed of H3K27me3, H3K27ac, a marker of
active chromatin, was diminished throughout the genome in *Dpy30*-KO cells (Fig 5B). Like H3K4me3,
H3K27ac was reduced even at promoters of genes that were stably expressed or upregulated in *Dpy30*-
KO cells (Fig 5D), indicating that the generalized decrease in H3K27ac was not a consequence of gene
downregulation. Despite the global reduction, however, H3K27ac was still positively associated with gene
expression in *Dpy30*-KO chromatin: H3K27ac peaks gaining intensity were near upregulated genes, and
H3K27ac peaks losing intensity were near downregulated genes, more frequently than expected by

chance (Fig S3). These observations are consistent with a model wherein acetylation of H3K27 is partially
a consequence of H3K4 methylation [14] and of transcription [47].

Reduction to transcriptional consistency in *Dpy30*-KO beta cells.
H3K4me3 enrichment has also been linked with transcriptional consistency—that is, low variability, or
"noise", in the expression of a given gene between cells [48,49]. A related metric, transcriptional entropy, is
a measure of the "specialization" of a cell's transcriptome, where a low entropy describes a narrow
distribution of highly expressed genes in each cell, versus a high entropy where gene expression becomes
less predictable [50]. In islets, transcriptional entropy decreases during maturation [50] and increases in T2D [4].
As such, we asked whether reduction of H3K4 methylation impairs transcriptional consistency in mature
β-cells. To measure transcriptional consistency, we processed islets from a WT and *Dpy30*-KO mouse for
single cell RNA-seq 45-days after tamoxifen administration, generating 2,510 WT and 2,597 KO cell
transcriptomes that passed quality filtering and that were identified as β-cells based on unsupervised
clustering and high expression of *Ins1 and Ins2* (Fig S4A). A population of β-cells expressing *tdTomato*,
indicating that they were not Cre-recombined, formed a distinct cluster from recombined *GFP*-positive β-
cells from the *Dpy30*-KO mouse (Fig 6A). To model divergence of *GFP*-positive from *tdTomato*-positive β-
cells (hereafter, mG and mT cells, respectively), we performed pseudotime ordering using the mT cluster
as the ground state (Fig 6B). As seen in Fig 6C, *Ins1* and *Ins2* expression became notably more stochastic
over pseudotime. To quantify variability of global transcription, we generated transcriptomic entropy
scores for each β-cell. In *Dpy30*-KO β-cells, there is an overall increase in transcriptional entropy along
pseudotime (Fig 6D) and in mG cells compared to mT cells (Fig 6E). This supports a role for H3K4
methylation in maintaining high transcriptional consistency in mature β-cells. Since higher entropy is a
feature of immature β-cells [50], we examined expression of genes associated with immaturity and
dedifferentiation and again did not observe any induction of these genes in Dpy30-KO cells (*c.f.* Figs 4F &
S4B-C). Therefore, *Dpy30*-KO in mature β-cells led to a more stochastic gene expression profile but did
not alter cell fate or re-activate an immature cell state.

We then asked whether the loss of transcriptional consistency was linked to the initial breadth of
the H3K4me3 peak, since it has been suggested that H3K4me3 peaks that span an especially large stretch
of DNA (i.e. broad peaks) may enforce high transcriptional consistency of associated genes [48,49]. We
hypothesized that genes with broader H3K4me3 peaks are more susceptible to increasingly stochastic
expression when H3K4me3 levels are reduced, compared to other genes with narrower peaks. To test
this, we ranked H3K4me3 peaks in WT cells in order of increasing breadth, grouped them into 20 quantiles,
each representing 5% (622) of H3K4me3 peaks with an associated gene. We then compared the variance
in expression, normalized to absolute expression level, of genes in each quantile between mT and mG
cells. Overall transcriptional variability between cells was increased in mG cells (Fig 6F). There was a
positive relationship between H3K4me3 peak breadth and the change in transcriptional variability, where
greater H3K4me3 peak breadth was linked to greater gain of variability in *Dpy30*-KO cells (Fig 6G). This
supports a role for H3K4me3 in constraining transcriptional variability in β-cells, which is positively linked
to peak breadth. Intriguingly, a similar pattern emerges in β-cells from human donors with T2D. Analysis
of publicly available datasets for H3K4me3 ChIP-seq from healthy human β-cells (GSE50386, ref. [51]) and
scRNA-seq from human donors with or without T2D (GSE124742, ref. [52]) showed that variability of gene
expression was higher in β-cells from donors with T2D (Fig 5H) and that the gain in variability is positively
correlated with initial H3K4me3 peak breadth (Fig 5I).

H3K4me3 peak breadth stratifies genes dysregulated in type 2 diabetes.

We further explored the relationship between H3K4me3 peak breadth, gene expression, and diabetes. H3K4me3 forms 18,936 distinct peaks with a median breadth of 1.2 kb in WT mouse β-cells. Ranking H3K4me3 peaks by breadth revealed a class of exceptionally broad peaks spanning up to 21 kb, including those at essential β-cell-enriched transcription factor genes (Fig 7A), consistent with the idea that genes important for maintaining cell identity are marked with broad H3K4me3 peaks [48,53]. As expected for repressed genes, the majority (29 of 39, or 74%) of islet "disallowed genes" [54], meanwhile, are H3K4me3-negative in healthy β-cells. Gene expression was moderately correlated with promoter H3K4me3 peak breadth (Spearman rank correlation R = 0.4287); however, genes with exceptionally broad peaks did not have exceptionally high RNA expression (Fig S5A). For example, Fig 7B shows the H3K4me3 profiles of a housekeeping gene with a typical H3K4me3 peak (*Rplp0*), and an expression-matched β-cell-enriched transcription factor with a broad peak (*Nkx6-1*). Therefore, β-cell lineage-enriched genes are marked by broad H3K4me3 peaks.

To explore the relationship between H3K4me3 peak breadth and transcriptome remodeling associated with diabetes, we turned to a mouse model of T2D. Leptin receptor-deficient *Lepr^db/db^* mice exhibit overfeeding, obesity, fasting hyperglycemia, and glucose intolerance, along with β-cell dysfunction [55]. We retrieved a published bulk RNA-seq dataset from *Lepr^db/db^* and *Lepr^+/+^* islets [56] and compared H3K4me3 peak breadth with gene expression changes. Interestingly, genes that become downregulated in *Lepr^db/db^* islets were explicitly enriched in the group of genes with broad H3K4me3 peaks (Fig 7C, left). Furthermore, housekeeping genes (GSEA:M11197, ref [57]), genes involved in endocrine pancreas development (GSEA:M12875), and maturity onset diabetes of the young (MODY) genes (GSEA:M18312) were enriched in the broadest H3K4me3 peaks (Fig S5B). In contrast, genes that are upregulated in *Lepr^db/db^* islets were uniquely enriched in the narrowest H3K4me3 peaks (Fig 7C, right). As H3K4me3 peak breadth was positively correlated with expression level (Fig S5A), we tested whether these observations could be secondary to a relationship wherein highly expressed genes become downregulated, and weakly expressed genes become upregulated, in *Lepr^db/db^*. Ranking genes by RNA expression level instead of H3K4me3 peak breadth proved less effective at stratifying each gene set except housekeeping genes (Fig S5B), supporting a primary link with H3K4me3. Therefore, genes that are differentially expressed in *Lepr^db/db^* islets are diametrically stratified according to H3K4me3 breadth, with upregulated genes having narrow peaks and downregulated genes having broad peaks (in non-diabetic conditions). These observations are replicable in humans using publicly available H3K4me3 ChIP-seq data of β-cells from donors without diabetes (GSE50386, ref [51]) and RNA-seq data of islets from donors with and without T2D (GSE50398, ref [58]) (Fig S5C-E).

H3K4me3 peak breadth dynamics encode gene expression changes in a mouse model of type 2 diabetes.

The positive association between H3K4me3 peak breadth and differential gene expression in T2D prompted us to ask if H3K4me3 is altered in diabetic islets. Immunoblots showed that, overall, H3K4me3 levels were not altered in islets from *Lepr^db/db^* mice (Fig 7D). Calibrated ChIP-seq similarly showed that average enrichment for H3K4me3 across all active promoters is remarkably similar between *Lepr^+/+^* and *Lepr^db/db^* islets (Fig 7E). While we did not find evidence for a global shift in H3K4me3 levels, we observed contraction of H3K4me3 at downregulated genes, for example the transcription factor *Pdx1* and zinc transporter *Slc30a8*, and expansion of H3K4me3 at upregulated genes, for example the disallowed

Aldh1a3 and *Ldha* (Fig 7F, Fig S5F). In general, broad and narrow H3K4me3 peaks showed significant
tendency to shrink or expand, respectively (Fig 7G). Similarly, genes that were transcriptionally up- or
downregulated in *Lepr^db/db* islets showed corresponding expansion or contraction of HK4me3 levels,
respectively (Fig 7H). Therefore, a pattern emerges in *Lepr^db/db* islets wherein weakly active genes tend to
gain H3K4me3 and become upregulated, while genes with broad H3K4me3 peaks tend to lose some
H3K4me3 and become downregulated.
324
Finally, we asked whether genes that are dysregulated in *Lepr^db/db* islets are regulated by H3K4
methylation, i.e., are downregulated in the day-45 *Dpy30*-KO RNA-seq. To test the overlap between
differential gene expression in the *Dpy30*-KO and *Lepr^db/db* models, we used a threshold-free rank-rank
hypergeometric overlap test (RRHO) [59]. There was significant overlap between genes downregulated in
Dpy30-KO mice with genes that were differentially expressed (either up or down) in *Lepr^db/db* mice (Fig 7J).
Thus, H3K4 methylation helps maintain expression of genes that become dysregulated in the diabetic
Lepr^db/db model. Accordingly, genes downregulated in *Dpy30*-KO mice were enriched in both the narrow
and broad H3K4me3 peak groups (Fig 7I-J), thus mirroring the pattern for dysregulated genes in the
Lepr^db/db model (Fig 7C). This observation in *Dpy30*-KO cells is in agreement with observations presented
above: that weakly active genes are most downregulated in *Dpy30*-KO (Fig 4), and that β-cell transcription
factor genes have broad peaks (Fig 7A) and tend to be downregulated by *Dpy30*-KO (Fig 4F). Together,
this supports the idea that differential gene expression between *Lepr^+/+* and *Lepr^db/db* mice is partially
driven by changes in H3K4me3 peak breadth.
338
Discussion
Here we set out to address whether H3K4me3 is required for maintenance of gene expression in mature
β-cells. To this end we deleted a core component of the COMPASS H3K4 methyltransferase complexes,
Dpy30, leading to dramatic genome-wide reduction of H3K4me3 in mouse β-cells. Loss of H3K4
methylation led to a less active and more repressed epigenome profile which locally correlated with gene
downregulation. Gene expression became more stochastic, a feature also shared with T2D. H3K4me3-
dependent gene expression meaningfully contributed to maintenance of mature β-cell function and
maturity. H3K4me3-deficient cells showed loss of maturity including reduced glucose stimulated insulin
secretion, reduced expression of terminal markers, and higher transcriptome entropy. These analyses
support a role for H3K4me3 in maintaining β-cell gene expression and function.
349
In *Dpy30*-KO cells, changes to RNA expression were linked to local changes to histone methylation,
leading to significant dysregulation of around 5% of genes, but overall cellular RNA levels were not
reduced. H3K4 methylation may, therefore, be important for allocation of transcriptional machinery to
particular genes, rather than their absolute activity. In other words, promoter-associated H3K4me levels
modulate the expression of target genes but global H3K4me3 or H3K4me1 levels should not be taken as
a reflection of a cell's transcriptional output. We note, however, that total mature RNA content, which we
measured here, is not a direct measure of transcriptional activity, and measurement of nascent RNA may
have uncovered more widespread roles in transcription regulation.
358
359 Hinting at more general roles in transcription is the observation that *Dpy30*-KO β-cells displayed
360 a more stochastic gene expression profile. High gene expression variability between cells in a tissue is a
361 beneficial feature in some circumstances – it can increase adaptability to stress and can reflect a high

degree of cell subtype specialization. In other circumstances, high variability is associated with aging and disease [60–63], and we find that it is increased in β-cells from donors with T2D. Very broad H3K4me3 peaks are proposed to confer high transcriptional stability [48], and experimental perturbation of H3K4me3 using *Dpy30*-KO supports this idea. Further, transcription of genes with H3K4me3 peaks of any breadth tended to become more variable in *Dpy30*-KO, meaning that H3K4me3 also generally confers consistent mRNA transcription. Defining exactly how H3K4me3 influences transcriptional activity of select genes will require detailed consideration of the larger epigenetic environment and RNA polymerase activity.

Since H3K4 cannot be simultaneously mono- and trimethylated, enrichment of H3K4me3 causes H3K4me1 to be generally excluded near TSSs. The degree of exclusion relates to the transcriptional output of the TSS, where weakly expressed genes display closer infiltration of H3K4me1 [64]. In *Dpy30*-KO cells, replacement of H3K4me3 with H3K4me1 in the TSS was associated with transcriptional downregulation, consistent with a report that promoter-associated H3K4me1 is repressive [65]. Surprisingly, replacement of H3K4me3 with H3K4me1 buffered gene downregulation compared to complete demethylation, suggesting that promoters with H3K4me1 are not repressed but simply less active, and H3K4me1 can partially compensates for loss H3K4me3 in promoters. This mirrors a finding that enhancers can function with H3K4me3 instead of H3K4me1 [13]. These results argue against an immutable functional distinction between H3K4me1 and H3K4me3, at least in this model.

One of the most surprising findings is that promoters in a weakly active state, with little H3K4me3, show the greatest reliance on H3K4 methylation for the maintenance of gene expression. It. A possible explanation is that these promoters are particularly targeted for epigenetic silencing [66] since H3K4 methylation sterically hinders repressive H3K27 and DNA methyltransferase activity [42,67]. Genes that were downregulated in *Dpy30*-KO cells showed high initial enrichment of H3K27me3 and further accumulation after loss of H3K4me3. Weakly-active genes are also the most likely to be upregulated after inactivation of the Polycomb system [43] and in β-cells, inactivation of the Polycomb system leads to activation of previously silenced genes [4]. Therefore, weakly active and suppressed genes may be the most responsive targets in ongoing competition between Polycomb and trithorax systems in mature β-cells. Future studies may unravel the degree to which H3K4me3 is an activating versus anti-repressive signal *in vivo*.

Peak breadth has emerged as an informative dimension of H3K4me3 enrichment: H3K4me3 peak breadth predicts gene expression level [38]; breadth dynamics predict gene expression dynamics during methionine restriction *in vivo* and *in vitro* [68]; and, exceptionally broad H3K4me3 peaks in a given cell type are predictive of genes critical to the maintenance of that cell's lineage [48]. From our analyses of peak breadth, we confirm that terminal markers of β-cells, including *Ins1/2* and mature β-cell enriched transcription factor genes, are marked with exceptionally broad H3K4me3 peaks in mature β-cells. *Dpy30*-KO led to nonspecific reduction of H3K4me3 genome-wide but transcriptional downregulation was largely restricted to genes that originally had either very narrow, or very broad, H3K4me3 peaks, indicating that these two classes of promoters in particular rely on H3K4me3 for the maintenance of gene expression. Compared to the effect at narrow peaks described above, the transcriptional defects for genes with broad H3K4me3 enrichment was more modest. For example, *Ins1* and *Ins2* were downregulated by 15 to 25%, remaining the two most highly expressed genes in *Dpy30*-KO cells despite suffering complete loss of their broad H3K4me3 peaks. Since broad peaks colocalize with many other activating chromatin marks [48,53], the modest effect likely results from compensation from other regulatory inputs. In any case, our results show

406 that broad H3K4me3 increases expression of associated genes but is not strictly required for their
407 transcription.
408
The *Dpy30*-KO model highlighted the central importance of relative H3K4me3 levels on a per-gene
basis rather than absolute enrichment; thus, reduction of peak breadth may exert a stronger impairment
on gene expression in scenarios where H3K4me3 is not simultaneously reduced at all other genes. To that
point, we find that promoters with broad H3K4me3 peaks showed a unique tendency to shrink, and for
the associated gene to be downregulated, in *Lepr^{db/db}* islets. Meanwhile, promoters marked with narrow
H3K4me3 peaks showed a significant tendency to be upregulated and gain H3K4me3 in *Lepr^{db/db}* islets. It
is notable that these two classes of promoters are the same that are most sensitive to H3K4me3
enrichment. More work will be required to determine what drives H3K4me3 peak dynamics in *Lepr^{db/db}*
islets and to what extent those mechanisms are active in human T2D. In any case, these results favour a
model wherein redistribution of H3K4me3 away from promoters of genes with broad H3K4me3 peaks,
that enforce β-cell identity or functions, in favour of relatively inactive, weakly-expressed genes, including
developmentally repressed and disallowed genes, contributes to transcriptome remodeling in diabetes.
421

Methods
Animal strains and care
Mice harboring conditional alleles of *Dpy30* (derived from EUCOMM EM:09575 [21]) and *Rosa26^{mTmG}* (Jax
007576 [29]) were crossed with *Pdx1-CreER^{Tg}* (Jax 024968 [28]) or *Ins1^{Cre}* (Jax 026801 [33]) lines. Genotypes used
for *Pdx1-CreER* studies were: KO: *Pdx1-CreER^{Tg/0}*; *Dpy30^{flox/flox}*; *Rosa26^{mTmG/+}*; WT: *Pdx1-CreER^{Tg/0}*; *Dpy30^{+/+}*;
Rosa26^{mTmG/+} or *Pdx1-CreER^{0/0}*; *Dpy30^{flox/flox}*; *Rosa26^{mTmG/+}*. Genotypes used for *Ins1^{Cre}* studies were: KO:
Ins1^{Cre/+}; *Dpy30^{flox/flox}*; WT: *Ins1^{+/+}*; *Dpy30^{flox/flox}*; HET: *Ins1^{Cre/+}*; *Dpy30^{flox/+}*. Mice were maintained on a mixed
genetic background. At 8-weeks-old, *Pdx1-CreER* mice were administered 8 mg tamoxifen (Sigma)
dissolved in 100 μL corn oil by oral gavage three times, with ~48-hours between administrations. The first
day of injection is considered day 0. BKS *Lepr^{db/db}* mice and BKS *Dock7^{m/m}* controls (Jax 000642) were used
at 12-weeks-old. Procedures were restricted to male mice and approved by the University of British
Columbia Animal Care Committee under certificates A17-0045 and A18-0111. Up to five littermates were
housed per cage with temperature and humidity control under a 12-hour light/dark cycle and ad libitum
access to water and food (Teklad 2918).
436

Cell culture
438 Primary mouse islet cells were maintained in RPMI 1640 medium containing 11 mM glucose (Gibco), 10%
439 (v/v) FBS, 50 U/ml penicillin, 50μg/mL streptomycin (complete RPMI) in 37°C, 5% CO_2 atmosphere.
440 *Drosophila* S2 cells were maintained at ambient temperature and air in Schneider's Drosophila Medium
441 (Gibco) containing 10% (v/v) heat-inactivated FBS, 50 U/ml penicillin, 50μg/mL streptomycin.
442

Islet and β-cell isolation
Unless otherwise indicated, mouse islets were isolated 45-days after tamoxifen administration (i.e. at 14-
weeks old) and processed for analysis immediately after isolation. Pancreata were perfused with 700
U/mL collagenase type XI (Sigma-Aldrich) in Hank's Buffered Saline Solution (138 mM NaCl, 5.3 mM KCl,
0.44 mM KH_2PO_4, 0.34 mM Na_2HPO_4, 5.5 mM D-Glucose, pH 7.3) through the bile duct, excised, and held
for 15-minutes at 40°C before manual disruption and hand-picking of islets. To isolate β-cells, islets from
Pdx1-CreER^{Tg/0} Rosa26^{mTmG/-} transgenic mice were washed 3× in room temperature PBS + 2 mM EDTA and

then dispersed by constant gentle pipetting in 1 mL of 0.025% Trypsin-EDTA (Thermo Fisher Scientific) in
PBS at room temperature for 8-10 minutes. Trypsin was quenched by addition of 9 mL room temperature
complete RPMI, cells were centrifuged for 3-minutes at 600 × g at 4°C, and resuspended in 0.3-0.5 mL ice-
cold sorting buffer (PBS, 2 mM EDTA, 1% (w/v) BSA). After passing through a 70 µm cell strainer, singlet
eGFP+ tdTomato- cells were purified by fluorescence-activated cell sorting (FACS) using a FACS-Aria II with
a 100 µm nozzle.
456
Chromatin immunoprecipitation sequencing (ChIP-seq)
A native ChIP protocol [69] was used to generate ChIP-seq libraries in biological duplicate, with
modifications. For *Pdx1-CreER$^{Tg/0}$*; *Dpy30$^{+/+}$*; *Rosa26$^{mTmG/+}$* and *Pdx1-CreER$^{Tg/0}$*; *Dpy30$^{flox/flox}$*; *Rosa26$^{mTmG/+}$*
mice, 100,000 eGFP+ tdTomato- islet cells were pooled with 50,000 *Drosophila* S2 cells by FACS. For
Lepr$^{db/db}$ and *Lepr$^{+/+}$* mice, 100,000 dispersed islet cells were counted by hemacytometer and pooled with
50,000 *Drosophila* S2 cells without sorting. Pooled cells were centrifuged at 600 × g for 5 minutes at 4°C,
supernatant removed, and then cell pellets were flash-frozen and stored at -80°C for up to one month.
Upon thawing, cells were permeabilized in nuclear isolation buffer (0.1% Triton X-100, 0.1% sodium
deoxycholate) containing 1 mM phenylmethylsulfonyl fluoride (PMSF) and EDTA-free protease inhibitor
cocktail (Roche)) on ice. Chromatin was fragmented to mononucleosomes by micrococcal nuclease
(MNase) (NEB) in digestion buffer (50 mM Tris pH 7.9, 5 mM CaCl$_2$, 1.5 mM DTT) for 7.5-minutes at 37°C,
then quenched by addition of 0.1 volume of a solution containing 2% Triton X-100, 2% sodium
deoxycholate, and 100 mM EDTA. Soluble chromatin was pre-cleared with Protein A Dynabeads (Thermo
Fisher Scientific) pre-adsorbed with normal rabbit IgG (Millipore, 12-370) for 2-hours at 4°C rotating 9
rpm, and then rotated overnight at 4°C with 1 µg of H3K4me3 (Abcam, ab1012), 2 µg H3K4me1 (Abcam,
ab8895), 1 µg H3K27ac (Active Motif, 39034), or 2 µg H3K27me3 (Millipore, 07-449) antibodies pre-
adsorbed to 10 µL of Protein A Dynabeads (Thermo Fisher Scientific) in 150 µL of ChIP buffer (20 mM Tris
pH 8.0, 2 mM EDTA, 150 mM NaCl, 0.2% Triton X-100, 1 mM PMSF, and EDTA-free protease inhibitor
cocktail (Roche)). After washing the beads twice with ice-cold wash buffer (20 mM Tris pH 8, 150 mM
NaCl, 2 mM EDTA, 1% (v/v) Triton X-100, 0.1% (w/v) SDS) and twice with ice-cold high-salt wash buffer (20
mM Tris pH 8, 500 mM NaCl, 2 mM EDTA, 1% (v/v) Triton X-100, 0.1% (w/v) SDS), DNA was eluted in
elution buffer (100 mM NaHCO$_3$, 1% (w/v) SDS, 50 µg/mL RNase A) for 1-hour at 55°C, with addition of 1
µL 20 mg/mL proteinase K after the first 30-minutes and occasional manual inversion. DNA was purified
using standard Phenol:chloroform:isoamyl alcohol extraction and ethanol precipitation and used for
library preparation with the NEBNext Ultra II DNA Library Prep Kit for Illumina (NEB) with 9 (H3K4me1,
H3K27me3) or 13 (H3K4me3, H3K27ac) amplification cycles. Indexed libraries were analyzed for size
distribution using the Bioanalyzer High Sensitivity DNA chip (Agilent) and for concentration using the Qubit
dsDNA HS Assay (Thermo Fisher Scientific). *Dpy30*-KO samples were sequenced on the NextSeq 500
platform (Illumina) for 2 × 78 nucleotide paired end reads and *Lepr$^{db/db}$* samples on the NextSeq 2000
platform (illumina) for 2 × 61 nucleotide paired end reads. Sequence data is available in the Gene
Expression Omnibus (GEO) under accession number GSE181951.
488
ChIP-seq data analysis
ChIP-seq reads were aligned to concatenated mouse (GRCm38/mm10) and *Drosophila* (BDGP6.28)
genome assemblies using Bowtie2 v2.3.4.1 with options '--very-sensitive --no-unal --no-discordant' [70].
Multi-mapped reads, reads with mapping quality MAPQ < 20, and suspected PCR-duplicates were
discarded using Samtools [71]. To scale enrichment according to the *Drosophila* cell spike-ins, the fraction

of remaining reads that mapped to the *Drosophila* genome was determined, and a scaling factor was
calculated for each sample as *(minimum* Drosophila *read fraction per histone mark)/(sample* Drosophila
read fraction). After determining a scaling factor, each sample was downsampled to the lowest mapped
read count per each histone mark using Samtools, and then reads mapped to the *Drosophila* genome were
removed. For visualization, libraries were converted to bedgraph format using the calculated scaling
factor in the '-scale' argument of BEDtools v2.26.0 genomecov [72]. Genome browser views were generated
for merged replicates using Spark [73] and TSS profiles using Deeptools v3.4.2 [74]. Peak locations and breadth
were determined using MACS2 v2.2.6 with parameters '--broad --nolambda' [41]. Peaks detected in only
one biological replicate or that overlapped with high-background blacklisted regions defined by the
ENCODE project [75] were excluded. Chromosomes X, Y, and M were also excluded from the analyses.
DESeq2 [76] was used for differential peak width analysis using a significance cutoff of $P ≤ 0.05$ (Wald test
with Benjamini and Hochberg adjustment), and enrichment for differential breadth in peak quantiles was
determined by one-sided Fisher's exact test. We defined TSS locations as the 5' end of the most abundant
annotated transcript of each gene detected in our RNA-seq data. H3K4me3 peaks were assigned to genes
for which they overlap the region ±1 kb of the TSS.
509
RNA-seq
eGFP+ tdTomato- β-cells (70,000-226,000 per mouse, biological triplicate) were purified and counted by
FACS and supplemented with a 10% spike-in of *Drosophila* S2 cells. *Drosophila* cells were included to
detect global changes in cell RNA content [77]. The cells were sorted directly into Trizol LS Reagent (Thermo
Fisher Scientific) and total RNA was extracted according to manufacturer's instructions. Residual DNA was
digested by Turbo DNase (Thermo Fisher Scientific) for 30-minutes at 37°C. Purified total RNA content was
measured using the Qubit RNA HS assay (Thermo Fisher Scientific). Mature mRNA was enriched from 400
ng of total RNA per sample using the NEBNext Poly(A) mRNA Magnetic Isolation Module (NEB) and used
to prepare libraries with the NEBNext Ultra II Directional RNA Library Prep Kit for Illumina (NEB) with 11
amplification cycles. Uniquely indexed libraries were quantified, pooled, and sequenced for 2 × 38
nucleotide reads as described above.
521
RNA-seq data analysis
Sequenced reads were aligned to the concatenated mouse (GRCm38/mm10; Gencode vM24) and
Drosophila (BDGP6.28; Ensembl release 99) genomes and transcriptomes using STAR v2.7.3a [78] with
default settings. Transcript abundance was calculated using Salmon v1.4.0 [79] in alignment mode with
seqBias and gcBias options. Gene counts and differential expression were calculated using DESeq2 [76].
Genes with fewer than two counts in any sample were filtered out and genes showing ≥ 2-fold difference
in expression with $P ≤ 0.01$ (Wald test with Benjamini and Hochberg adjustment for multiple comparisons)
are considered differentially expressed. Genes showing <1.1 fold difference in expression are considered
stable. Except in Fig. 6F, Drosophila transcripts were removed after abundance calculation; for Fig. 6F,
Drosophila transcripts were used as control genes in the estimateSizeFactors function of DESeq2, and then
removed.
533
Single-cell RNA-seq
Islets from one WT and one KO mouse were dissociated to single-cell suspensions as described above
without FACS-enrichment. The cell suspensions were processed through the Chromium Single Cell 3'
protocol using the Chromium Controller (firmware v4.0) with Reagent Kit v3.1 (10X Genomics) and Dual

Index Kit TT Set A (10X Genomics) according to the manufacturer's instructions for a targeted 5,000-cell
recovery. Total cDNA was amplified for 11-cycles, and then 13-cycles during index-ligation. cDNA
concentrations were quantified by qPCR with the NEBNext Library Quant Kit for Illumina (NEB). Libraries
were pooled and sequenced for 28-90 paired-end nucleotide reads to an average depth of ~70,000
mapped fragments per cell.
543

Single cell RNA-seq data analysis
Cellranger (v5.0.0, 10X Genomics) was used to generate FASTQ files and demultiplex reads. Cell barcode
detection, read mapping, quality filtering, and transcript counting were performed using Alevin (Salmon
v1.4.0) [80] against the mouse protein-coding transcriptome (Gencode vM24), which we appended with the
Tdtomato and *Egfp* sequences, as well as the mouse genome (GRCm38/mm10) as mapping decoys for
selective alignment [81]. Additional cell quality filtering was performed using Seurat v4.0 [82] based on unique
features (> 1,000 genes) and mitochondrial RNA content (< 25%). Putative doublets were excluded using
DoubletFilter [83]. Genes detected in < 3 cells were excluded from analysis. After quality filtering, the dataset
comprised 3,703 WT and 3,641 KO cells. Filtered data were scaled and normalized using SCTransform [84]
with default parameters. PCA and UMAP dimensionality reduction using the first 60 principal components
was used to cluster and visualize cell populations. We then used Slingshot [85] to perform pseudotime
analysis on β-cell clusters. Entropy scores are a modified Shannon entropy calculated using a published R
function (ref [86]).
557

Immunohistochemistry
Mouse pancreata were isolated and fixed in 4% (w/v) paraformaldehyde (PFA) in PBS at 4°C overnight.
Following washing with PBS and dehydration in ethanol and xylenes, mouse tissues were embedded in
paraffin and sectioned with a thickness of 5 μm. Immunofluorescent analyses of tissue sections was
performed as previously described[23] in a blinded fashion from 4-6 sections per biological sample. Primary
antibodies used were: H3K4me3 (Cell Signaling Technologies, C42D8, 1:500), H3K4me1 (Abcam, ab8895,
1:1,000), H3 (Abcam, ab1791, 1:1,000), DPY30 (Atlas Antibodies, HPA043761, 1:500), Insulin (Agilent, IR-
002, 1:4), and Glucagon (Sigma Aldrich, G2654, 1:1,000).
566

Transmission electron microscopy
Islets were fixed in 2% glutaraldehyde (Sigma Aldrich) in PBS and submitted to the Facility for Electron
Microscopy Research of McGill University (Hamilton, ON, Canada). Islets were post-fixed with 1% osmium
tetroxide, dehydrated in ethanol, and embedded in Spurr's resin before sectioning with a Leica UCT
ultramicrotome. Sections were stained with uranyl acetate and lead citrate and imaged with a JEOL JEM
1200 EX TEMSCAN transmission electron microscope (JEOL, Peabody, MA, USA).
573

Immunoblotting
Islets were lysed in Laemmli sample buffer (2% (w/v) SDS, 10% (v/v) glycerol, 60 mM Tris pH 6.8, 1mM
NaF, 1 mM PMSF, protease inhibitor cocktail (Roche)) for 5-minutes at 95°C. Protein concentration was
measured by BCA assay (Thermo Fisher Scientific), then β-mercaptoethanol was added to 5% (v/v) and
lysates were reboiled for 5-minutes and stored at -80°C. 10 μg lysates were resolved in polyacrylamide
gels, transferred to PVDF membranes, blocked for 60-minutes in 5% (w/v) skim milk powder in TBST (20
mM Tris pH 7.6, 150 mM NaCl, 0.1% (v/v) Tween-20) at room temperature and probed with primary
antibodies overnight in 5% (w/v) BSA in TBST at 4°C. The next day, after washing membranes three times

with TBST, HRP-linked secondary antibodies were added for 1-hour at room temperature, membranes were washed three times with TBST, and detected using ECL system with film. When normalizing to total H3, membranes were stripped for 20-minutes in mild stripping buffer (200 mM glycine, 0.1% (w/v) SDS, 1% (v/v) Tween-20, pH 2.2) at room temperature, rinsed twice in PBS and twice in TBST for 10 minutes each at room temperature, then re-blocked and probed for histone H3. Primary antibodies used for immunoblots were: H3K4me3 (Cell Signaling Technologies, C42D8, 1:1,000), H3K4me1 (Abcam, ab8895, 1:100,000), H3K27ac (Cell Signaling Technologies, D5E4, 1:1,000), H3K27me3 (Millipore, 07-449, 1:100,000), H3 (Abcam, ab1791, 1:1,000,000), WDR5 (Bethyl, A302-430A, 1:2,000), RBBP5 (Bethyl, A300-109A, 1:10,000), and ASH2L (Bethyl, A300-489A, 1:10,000).

Protein co-immunoprecipitation

Immunoprecipitation of WDR5 was performed in triplicate in 750-900 dispersed islets pooled from 3 mice. First, nuclei were enriched from dispersed islet cells by incubation in nuclear isolation buffer for 10-minutes on ice and pelleting at 600 × g for 5-minutes at 4°C. Chromatin was solubilized in IP buffer (20 mM Tris pH 7.4, 150 mM NaCl, 1 mM EDTA, 5% (v/v) glycerol, 0.5% (v/v) Igepal ca-630, 5 mM CaCl$_2$, 1 mM PMSF, protease inhibitor cocktail (Roche)) with MNase (NEB) for 10-minutes at room temperature. EDTA was added to 5 mM to stop MNase and samples were centrifuged at 14,000 × g for 5-minutes at 4°C. The protein concentration in the supernatant was measured by BCA assay (Thermo Fisher Scientific), and an equal mass of WT and KO nuclear lysate (~30 μg protein) was immunoprecipitated by 1 μg WDR5 (Bethyl, A302-429A) or normal rabbit IgG (Millipore, 12-370) antibodies rotating overnight at 4°C. The next day, 10 μL protein A Dynabeads (Thermo Fisher Scientific) was added to the antibody:lysate mixture and rotated for 2-hours at 4°C. Finally, bead:antibody:antigen complexes were washed six times in ice-cold IP buffer and resuspended in Laemmli sample buffer with 5% (v/v) β-mercaptoethanol for immunoblotting.

Insulin secretion assays

For in vivo glucose tolerance tests, 2 g/kg glucose in a 20% (w/v) water solution was injected intraperitoneally following a 6-hour fast. Blood glucose was sampled immediately prior to and 15, 30, 60, and 120-minutes after glucose injection from a lateral saphenous vein. Serum was collected at the 0, 15, and 60-minute time points by centrifuging blood samples at 9,000 × g for 9 minutes at 4°C and stored at -80°C. Serum insulin was measured by ELISA (Alpco) according to the manufacturer's instructions. Ex vivo static insulin secretion assays were performed in technical duplicate in groups of 50 islets per mouse one day after isolation. Islets were preincubated in KRBH buffer (2.8 mM glucose, 20 mM HEPES, 114 mM NaCl, 4.7 mM KCl, 1.2 mM MgSO$_4$, 1.2 mM KH$_2$PO$_4$, 2.5 mM CaCl$_2$, 24 mM NaHCO$_3$, 0.1% (w/v) BSA, pH 7.3) for one hour at 37°C and then sequentially for 45-minutes in KRBH, either 16.7 mM glucose or 15 mM dimethyl α-ketoglutarate (Sigma Aldrich), and finally 30 mM KCl. Conditioned medium was collected after each incubation. At the completion of the assay islets were lysed in RIPA buffer (Pierce) containing 1 mM PMSF and EDTA-free protease inhibitor cocktail (Roche)) at 95°C for 5 minutes. Insulin concentration was measured in conditioned medium and islet lysate by ELISA (Alpco). DNA concentration of islet lysate was determined by Qubit dsDNA HS Assay (Thermo Fisher Scientific) to normalize islet insulin content.

Calcium imaging

Intact islets were cultured for two days to allow for attachment onto glass coverslips. Prior to imaging, coverslips were transferred to a 2 mL imaging chamber and stained with 5 μM Fura-2AM (Molecular Probes, F1221) for 30-minutes at 37°C in complete RPMI. The imaging chamber loaded with a coverslip

was then mounted on a Leica SP8 Laser Scanning Confocal Microscope with a 10× objective lens and
continuously washed with Ringer's solution (5.5 mM KCl, 2 mM CaCl$_2$, 1 mM MgCl$_2$, 20 mM HEPES, 144
mM NaCl, 2.8 mM glucose, pH 7.4) for 30-minutes. Throughout the experiment, islets were continuously
perifused at a flow rate of 2.5 mL/min with Ringer's solution. Glucose and KCl concentrations were
adjusted by iso-osmotic substitution of NaCl where indicated. Fura-2 was excited at 340 nm and 380 nm
and emitted fluorescence was detected at approximately 510 nm. Cytosolic Ca^{2+} levels are expressed as
the mean ratio of Fura-2 fluorescence emission intensity (F_{340}/F_{380}) of 3-9 islets per biological replicate.
633

Respirometry
Islets were dissociated to single cells 3-4 hours after isolation and plated at 100 islets/well in a Seahorse
24-well cell culture dish in technical duplicate. Cells were cultured for 2-days in complete RPMI to allow
for attachment to the culture dish and then for 1-hour in XF RPMI medium (Agilent) with 2.8 mM glucose,
1% FBS, 1 mM pyruvate, and 2 mM glutamine to reach a metabolic baseline. Oxygen consumption was
then measured using a Seahorse XF24 Extracellular Flux Analyser (Agilent). The injection ports were
loaded with the following compounds in XF Assay Media (Agilent): A, 167 mM glucose; B, 10 µM
oligomycin; C, 15 µM FCCP; D, 10 µM antimycin A and 1 µM rotenone. Measurements were normalized
to the DNA content of assayed cells, which were determined using Qubit HS DNA Assay (Thermo Fisher
Scientific) after lysing in RIPA buffer upon completion of the assay.
644

645 **Analysis of publicly available data**
646 Publicly available data were reprocessed for consistency. GEO: GSE50244, GSE50386, GSE107489,
647 GSE124742 were downloaded from the Sequence Read Archive (SRA) using the sratoolkit
648 (https://github.com/ncbi/sra-tools) and analyzed according to the experimental application, as described
649 above.
650

Statistics
Unless otherwise stated, plots show mean ± standard deviation (SD) with points representing each
biological replicate. In boxplots, the central horizontal line marks median, the upper and lower limits of
the box the first and third quartiles, and whiskers span 1.5× the interquartile range, in the style of Tukey.
Spearman's rank coefficient was used to estimate correlation. Regression lines were generated using a
generalized additive model. *P*-values were calculated using Student's t-tests with Welch's correction,
Wilcoxon rank sum test, Wald test, mixed-effect model, one- or two-way ANOVA, Fisher's exact test, or
permutation test, as indicated. Correction for multiple comparison using the Benjamini-Hochberg method
was applied where indicated. Statistical calculations were performed in R or Prism 8 or 9 (GraphPad
Software, La Jolla, CA, USA). *P* < 0.05 was deemed significant.
661

Author contributions
Experiments were conducted by B.V., N.M., D.J.P., M.A., and C.L.M, and conceptualized by B.V. and B.G.H.
D.J.P. and D.S.L. performed calcium imaging experiments and designed respirometry experiments.
Bioinformatics were performed by B.V. and B.G.H. Funding and supervision was provided by B.G.H. and
F.C.L; the manuscript was written by B.V. B.G.H. is the guarantor of this work and, as such, had full access
to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the
data analysis.
669

Declaration of interests

All authors declare they have no competing interests.

Acknowledgements

The authors thank the staff of the Animal Care Facility at British Columbia Children's Hospital Research Institute for daily maintenance of the mouse colonies, and W. Gibson (UBC) for providing Ins1[Cre] animals. We thank M. Reid at the Facility for Electron Microscopy Research of McGill University for help with transmission electron microscopy, L. Xu at the Flow Core of British Columbia Children's Hospital Research Institute for help with FACS. This work was supported by operating grants from the British Columbia Children's Hospital Research Institute, the Natural Sciences and Engineering Research Council of Canada (RGPIN-2016-04292) and the Canadian Institute for Health Research (RN310864 − 375894). Fellowship support for B.V. was provided by the British Columbia Children's Hospital Research Institute, University of British Columbia, and Natural Sciences and Engineering Research Council of Canada.

References

1. Saeedi, P. *et al.* Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045: Results from the International Diabetes Federation Diabetes Atlas, 9th edition. *Diabetes Res. Clin. Pract.* **157**, 1–10 (2019).

2. Salinno, C. *et al.* β-Cell Maturation and Identity in Health and Disease. *Int. J. Mol. Sci.* **20**, 1–20 (2019).

3. Kim, H. *et al.* PRMT1 is required for the maintenance of mature β-cell identity. *Diabetes* **69**, 355–368 (2020).

4. Lu, T. T. H. *et al.* The Polycomb-Dependent Epigenome Controls β Cell Dysfunction, Dedifferentiation, and Diabetes. *Cell Metab.* **27**, 1294-1308.e7 (2018).

5. Dhawan, S., Georgia, S., Tschen, S. ing, Fan, G. & Bhushan, A. Pancreatic β Cell Identity Is Maintained by DNA Methylation-Mediated Repression of Arx. *Dev. Cell* **20**, 419–429 (2011).

6. Dayeh, T. *et al.* Genome-Wide DNA Methylation Analysis of Human Pancreatic Islets from Type 2 Diabetic and Non-Diabetic Donors Identifies Candidate Genes That Influence Insulin Secretion. *PLoS Genet.* **10**, e1004160 (2014).

7. Deering, T. G., Ogihara, T., Trace, A. P., Maier, B. & Mirmira, R. G. Methyltransferase set7/9 maintains transcription and euchromatin structure at islet-enriched genes. *Diabetes* **58**, 185–193 (2009).

8. Kuo, T. *et al.* Identification of C2CD4A as a human diabetes susceptibility gene with a role in β cell insulin secretion. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 20033–20042 (2019).

9. De Jesus, D. F. *et al.* m6A mRNA methylation regulates human β-cell biology in physiological states and in type 2 diabetes. *Nat. Metab.* **1**, 765–774 (2019).

10. Miguel-Escalada, I. *et al.* Human pancreatic islet three-dimensional chromatin architecture provides insights into the genetics of type 2 diabetes. *Nat. Genet.* **51**, 1137–1148 (2019).

11. Jenuwein, T. & Allis, C. D. Translating the Histone Code. *Science (80-.).* **293**, 1074–1080 (2001).

12. Barski, A. *et al.* High-Resolution Profiling of Histone Methylations in the Human Genome. *Cell* **129**, 823–837 (2007).

13. Rickels, R. *et al.* Histone H3K4 monomethylation catalyzed by Trr and mammalian COMPASS-like proteins at enhancers is dispensable for development and viability. *Nat. Genet.* **49**, 1647–1653 (2017).

14. Dorighi, K. M. *et al.* Mll3 and Mll4 Facilitate Enhancer RNA Synthesis and Transcription from Promoters Independently of H3K4 Monomethylation. *Mol. Cell* **66**, 568-576.e4 (2017).

15. Hödl, M. & Basler, K. Transcription in the absence of histone H3.2 and H3K4 methylation. *Curr.*

717   *Biol.* **22**, 2253–2257 (2012).

718 16. Howe, F. S., Fischl, H., Murray, S. C. & Mellor, J. Is H3K4me3 instructive for transcription
719   activation? *BioEssays* **39**, 1–12 (2017).

720 17. Ernst, P. *et al.* Definitive hematopoiesis requires the mixed-lineage leukemia gene. *Dev. Cell* **6**,
721   437–443 (2004).

722 18. Glaser, S. *et al.* The histone 3 lysine 4 methyltransferase, Mll2, is only required briefly in
723   development and spermatogenesis. *Epigenetics and Chromatin* **2**, 1–16 (2009).

724 19. Lee, J. E. *et al.* H3K4 mono- And di-methyltransferase MLL4 is required for enhancer activation
725   during cell differentiation. *Elife* **2013**, 1–25 (2013).

726 20. Bledau, A. S. *et al.* The H3K4 methyltransferase Setd1a is first required at the epiblast stage,
727   whereas Setd1b becomes essential after gastrulation. *Dev.* **141**, 1022–1035 (2014).

728 21. Campbell, S. A., McDonald, C. L., Krentz, N. A. J., Lynn, F. C. & Hoffman, B. G. TrxG Complex
729   Catalytic and Non-catalytic Activity Play Distinct Roles in Pancreas Progenitor Specification and
730   Differentiation. *Cell Rep.* **28**, 1830-1844.e6 (2019).

731 22. Piunti, A. & Shilatifard, A. Epigenetic balance of gene expression by polycomb and compass
732   families. *Science (80-.).* **352**, (2016).

733 23. Campbell, S. A. *et al.* H3K4 trimethylation is required for postnatal pancreatic endocrine cell
734   functional maturation. *bioRxiv* (2020). doi:https://doi.org/10.1101/2020.11.29.402990

735 24. Scoville, D. W. *et al.* MLL3 and MLL4 methyltransferases bind to the MafA and MAFB
736   transcription factors to regulate islet β-cell function. *Diabetes* **64**, 3772–3783 (2015).

737 25. Sze, C. C. *et al.* Histone H3K4 methylation-dependent and -independent functions of
738   set1A/COMPASS in embryonic stem cell self-renewal and differentiation. *Genes Dev.* **31**, 1732–
739   1737 (2017).

740 26. Jiang, H. *et al.* Role for Dpy-30 in ES Cell-Fate Specification by Regulation of H3K4 Methylation
741   within Bivalent Domains. *Cell* **144**, 513–525 (2011).

742 27. Haddad, J. F. *et al.* Structural Analysis of the Ash2L/Dpy-30 Complex Reveals a Heterogeneity in
743   H3K4 Methylation. *Structure* **26**, 1–10 (2018).

744 28. Gu, G., Dubauskaite, J. & Melton, D. A. Direct evidence for the pancreatic lineage: NGN3+ cells
745   are islet progenitors and are distinct from duct progenitors. *Development* **129**, 2447–2457
746   (2002).

747 29. Muzumdar, M. D., Tasic, B., Miyamichi, K., Li, L. & Luo, L. A Global Double-Fluorescent Cre
748   Reporter Mouse. *Genesis* **45**, 593–605 (2007).

749 30. Taruttis, F. *et al.* External calibration with Drosophila whole-cell spike-ins delivers absolute mRNA
750   fold changes from human RNA-Seq and qPCR data. *Biotechniques* **62**, 53–61 (2017).

751 31. Zhang, J. *et al.* The diabetes gene Hhex maintains δ -cell differentiation and islet function. *Genes*
752   *Dev.* **28**, 829–834 (2014).

753 32. Sabatini, P. V. *et al.* Neuronal PAS Domain Protein 4 Suppression of Oxygen Sensing Optimizes
754   Metabolism during Excitation of Neuroendocrine Cells. *Cell Rep.* **22**, 163–174 (2018).

755 33. Thorens, B. *et al.* Ins1 Cre knock-in mice for beta cell-specific gene recombination. *Diabetologia*
756   **58**, 558–565 (2015).

757 34. Millership, S. J. *et al.* Neuronatin regulates pancreatic β cell insulin content and secretion. *J. Clin.*
758   *Invest.* **128**, 3369–3381 (2018).

759 35. Wijesekara, N. *et al.* Beta cell-specific Znt8 deletion in mice causes marked defects in insulin
760   processing, crystallisation and secretion. *Diabetologia* **53**, 1656–1668 (2010).

761 36. Matschinsky, F. M. & Wilson, D. F. The central role of glucokinase in glucose homeostasis: A
762   perspective 50 years after demonstrating the presence of the enzyme in islets of Langerhans.
763   *Front. Physiol.* **10**, (2019).

764 37. Ernst, J. *et al.* Mapping and analysis of chromatin state dynamics in nine human cell types. *Nature*

473, 43–49 (2011).

38. Karlić, R., Chung, H. R., Lasserre, J., Vlahoviček, K. & Vingron, M. Histone modification levels are
predictive for gene expression. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 2926–2931 (2010).

39. Choudhury, R., Singh, S., Arumugam, S., Roguev, A. & Stewart, A. F. The Set1 complex is dimeric
and acts with Jhd2 demethylation to convey symmetrical H3K4 trimethylation. *Genes Dev.* **33**, 1–
15 (2019).

40. Clouaire, T. *et al.* Cfp1 integrates both CpG content and gene activity for accurate H3K4me3
deposition in embryonic stem cells. *Genes Dev.* **26**, 1714–1728 (2012).

41. Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, (2008).

42. Ooi, S. K. T. *et al.* DNMT3L connects unmethylated lysine 4 of histone H3 to de novo methylation
of DNA. *Nature* **448**, 714–717 (2007).

43. Dobrinić, P., Szczurek, A. T. & Klose, R. J. PRC1 drives Polycomb-mediated gene repression by
controlling transcription initiation and burst frequency. *Nat. Struct. Mol. Biol.* **28**, 811–824
(2021).

44. Kennison, J. A. The Polycomb and trithorax group proteins of Drosophila: Trans-Regulators of
Homeotic Gene Function. *Annu. Rev. Genet.* **29**, 289–303 (1995).

45. Schuettengruber, B., Bourbon, H.-M., Di Croce, L. & Cavalli, G. Genome Regulation by Polycomb
and Trithorax: 70 Years and Counting. *Cell* **171**, 34–57 (2017).

46. Blum, B. *et al.* Functional beta-cell maturation is marked by an increased glucose threshold and
by expression of urocortin 3. *Nat. Biotechnol.* **30**, 261–264 (2012).

47. Martin, B. J. E. *et al.* Transcription shapes genome-wide histone acetylation patterns. *Nat.*
Commun. **12**, 1–9 (2021).

48. Benayoun, B. A. *et al.* H3K4me3 breadth is linked to cell identity and transcriptional consistency.
Cell **158**, 673–688 (2014).

49. Collins, B. E., Sweatt, J. D. & Greer, C. B. Broad domains of histone 3 lysine 4 trimethylation are
associated with transcriptional activation in CA1 neurons of the hippocampus during memory
formation. *Neurobiol. Learn. Mem.* **161**, 149–157 (2019).

50. Kannan, S., Farid, M., Lin, B. L., Miyamoto, M. & Kwon, C. Transcriptomic entropy benchmarks
stem cell-derived cardiomyocyte maturation against endogenous tissue at single cell level. *PLoS*
Comput. Biol. **17**, 1–21 (2021).

51. Bramswig, N. C. *et al.* Epigenomic plasticity enables human pancreatic α to β cell reprogramming.
J. Clin. Invest. **123**, 1275–1284 (2013).

52. Camunas-Soler, J. *et al.* Patch-Seq Links Single-Cell Transcriptomes to Human Islet Dysfunction in
Diabetes. *Cell Metab.* **31**, 1017-1031.e4 (2020).

53. Chen, K. *et al.* Broad H3K4me3 is associated with increased transcription elongation and
enhancer activity at tumor-suppressor genes. *Nat. Genet.* **47**, 1149–1157 (2015).

54. Pullen, T. J. *et al.* Identification of genes selectively disallowed in the pancreatic islet. *Islets* **2**, 89–
95 (2010).

55. Talchai, C., Xuan, S., Lin, H. V., Sussel, L. & Accili, D. Pancreatic β Cell Dedifferentiation as a
Mechanism of Diabetic β Cell Failure. *Cell* **150**, 1223–1234 (2012).

56. Neelankal John, A., Ram, R. & Jiang, F.-X. RNA-Seq Analysis of Islets to Characterise the
Dedifferentiation in Type 2 Diabetes Model Mice db/db. *Endocr. Pathol.* **29**, 207–221 (2018).

57. Hsiao, L. L. *et al.* A compendium of gene expression in normal human tissues. *Physiol. Genomics*
2002, 97–104 (2002).

58. Fadista, J. *et al.* Global genomic and transcriptomic analysis of human pancreatic islets reveals
novel genes influencing glucose metabolism. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 13924–13929
(2014).

59. Cahill, K. M., Huo, Z., Tseng, G. C., Logan, R. W. & Seney, M. L. Improved identification of

concordant and discordant gene expression signatures using an updated rank-rank
hypergeometric overlap approach. *Sci. Rep.* **8**, 1–11 (2018).

60. Li, J., Liu, Y., Kim, T. H., Min, R. & Zhang, Z. Gene expression variability within and between
human populations and implications toward disease susceptibility. *PLoS Comput. Biol.* **6**, (2010).

61. Ecker, S., Pancaldi, V., Valencia, A., Beck, S. & Paul, D. S. Epigenetic and Transcriptional Variability
Shape Phenotypic Plasticity. *BioEssays* **40**, 1–11 (2018).

62. Rangaraju, S. *et al.* Suppression of transcriptional drift extends C. elegans lifespan by postponing
the onset of mortality. *Elife* **4**, 1–39 (2015).

63. Benninger, R. K. P. & Kravets, V. The physiological role of β-cell heterogeneity in pancreatic islet
function. *Nat. Rev. Endocrinol.* **0123456789**, (2021).

64. Soares, L. M. *et al.* Determinants of Histone H3K4 Methylation Patterns. *Mol. Cell* **68**, 773-785.e6
(2017).

65. Cheng, J. *et al.* A role for H3K4 monomethylation in gene repression and partitioning of
chromatin readers. *Mol. Cell* **53**, 979–992 (2014).

66. Kumar, D., Cinghu, S., Oldfield, A. J., Yang, P. & Jothi, R. Decoding the function of bivalent
chromatin in development and cancer. *Genome Res.* **31**, 2170–2184 (2021).

67. Schmitges, F. W. *et al.* Histone Methylation by PRC2 Is Inhibited by Active Chromatin Marks. *Mol.*
Cell **42**, 330–341 (2011).

68. Dai, Z., Mentch, S. J., Gao, X., Nichenametla, S. N. & Locasale, J. W. Methionine metabolism
influences genomic architecture and gene expression through H3K4me3 peak width. *Nat.*
Commun. **9**, 1–12 (2018).

69. Brind'Amour, J. *et al.* An ultra-low-input native ChIP-seq protocol for genome-wide profiling of
rare cell populations. *Nat. Commun.* **6**, 1–8 (2015).

70. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**, 357–
359 (2012).

71. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079
(2009).

72. Quinlan, A. R. & Hall, I. M. BEDTools: A flexible suite of utilities for comparing genomic features.
Bioinformatics **26**, 841–842 (2010).

73. Kurtenbach, S. & William Harbour, J. SparK: A Publication-quality NGS Visualization Tool. *bioRxiv*
1–4 (2019). doi:10.1101/845529

74. Ramírez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data analysis.
Nucleic Acids Res. **44**, W160–W165 (2016).

75. Amemiya, H. M., Kundaje, A. & Boyle, A. P. The ENCODE Blacklist: Identification of Problematic
Regions of the Genome. *Sci. Rep.* **9**, 1–5 (2019).

76. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biol.* **15**, 1–21 (2014).

77. Taruttis, F. *et al.* External calibration with Drosophila whole cell spike ins delivers absolute mRNA
fold changes from human RNA seq and qPCR data. *Biotechniques* **62**, 53–61 (2017).

78. Dobin, A. *et al.* STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

79. Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast and bias-
aware quantification of transcript expression. *Nat. Methods* **14**, 417–419 (2017).

80. Srivastava, A., Malik, L., Smith, T., Sudbery, I. & Patro, R. Alevin efficiently estimates accurate
gene abundances from dscRNA-seq data. *Genome Biol.* **20**, (2019).

81. Sarkar, H., Zakeri, M., Malik, L. & Patro, R. Towards selective-alignment: Bridging the accuracy
gap between alignment-based and alignment-free transcript quantification. *bioRxiv* 1–7 (2017).
doi:10.1101/138800

82. Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902.e21 (2019).

83. McGinnis, C. S., Murrow, L. M. & Gartner, Z. J. DoubletFinder: Doublet Detection in Single-Cell
RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst.* **8**, 329-337.e4 (2019).
84. Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-seq data
using regularized negative binomial regression. *bioRxiv* 1–15 (2019). doi:10.1101/576827
85. Street, K. *et al.* Slingshot: cell lineage and pseudotime inference for single-cell transcriptomics.
BMC Genomics **19**, 1–16 (2018).
86. Kannan, S., Farid, M., Lin, B., Miyamoto, M. & Kwon, C. Transcriptomic entropy benchmarks stem
cell-derived cardiomyocyte maturation against endogenous tissue at single cell level. *bioRxiv*
(2020). doi:10.1101/2020.04.02.022632
87. Huang, D. W., Sherman, B. T. & Lempicki, R. A. Systematic and integrative analysis of large gene
lists using DAVID bioinformatics resources. *Nat. Protoc.* **4**, 44–57 (2009).
88. Brideau, C. M., Eilertson, K. E., Hagarman, J. A., Bustamante, C. D. & Soloway, P. D. Successful
Computational Prediction of Novel Imprinted Genes from Epigenomic Features. *Mol. Cell. Biol.*
30, 3357–3370 (2010).

Figure legends
Figure 1. Reduction to H3K4 methylation in beta cells leads to glucose intolerance and hyperglycemia
in adult mice.
A, Top: schematic showing the core subunits of COMPASS complexes and a nucleosome methylated on
H3K4. Bottom: genome-aligned RNA-seq reads at the *Dpy30* gene locus in *Dpy30*-WT and -KO cells. **B,**
Immunoblots showing COMPASS core subunits RBBP5, ASH2L, and WDR5, and nucleosome protein
histone H3, coimmunoprecipitated with WDR5 or an IgG control from *Dpy30*-WT or -KO islet cell nuclei 45
days after tamoxifen administration. Representative immunoblots of three independent coIPs are shown.
IP: immunoprecipitation. **C,** Immunoblots showing H3K4me3, H3K4me1, H3K27ac, H3K27me3, and total
histone H3 in islets from *Dpy30*-WT and -KO mice 45 days after tamoxifen administration. Numbers
beneath each band indicate the band intensity normalized to the left-most sample. **D,** Mean
immunofluorescent intensity of DPY30, H3K4me3, and H3K4me1 in *Dpy30*-WT and -KO β-cell nuclei at the
indicated days after tamoxifen administration. Data are normalized to fluorescence intensity of α-cell
nuclei; n = 3 mice of each genotype and time-point. **E,** Example IHC images of *Dpy30*-KO islets used for
measurements in panel D showing H3K4me3 (cyan), insulin (magenta), and glucagon (yellow). Scale bars:
100 μm. **F,** Scatterplots showing $\log_2$(fold-change) and $-\log_{10}$(*P*-value) of gene expression in *Dpy30*-KO
versus -WT cells 15- and 45-days after tamoxifen administration. Genes showing a ≥2-fold increase or
decrease in expression with *P* ≤ 0.01 (Wald test with Benjamini-Hochberg correction) are coloured red or
green, respectively, and enumerated above. The full-length transcript of *Dpy30* is outlined and labeled in
blue. **G,** Blood glucose (top) and serum insulin (bottom) concentration during IPGTT in *Dpy30*-WT and -KO
mice 15-, 30-, and 45-days after tamoxifen administration. Data presented as mean ± SD (n = 8 mice for
glucose measurements, 8-15 mice for insulin measurements). *P*-values were calculated from area under
the curve (AUC) using multiple two-tailed t-tests with Welch's and Benjamini-Hochberg corrections; only
values ≤ 0.05 are shown. **H,** Unfasted blood glucose (top) and body mass (bottom) of *Dpy30*-WT and -KO
mice during the 60-days after tamoxifen administration. Data are presented as individual measurements
with mean (n = 8, however tracking was stopped after a blood glucose reading ≥ 20 mM).
902

Figure 2. Genes involved in insulin production and glucose-induced activity are regulated by H3K4me3.
A, Genome browser views of H3K4me3 and H3K4me1 enrichment at the *Ins1* and *Ins2* gene loci in *Dpy30*-
WT and -KO β-cells 45-days after tamoxifen administration. **B,** *Ins1* and *Ins2* RNA levels in *Dpy30*-WT and

-KO cells. Data presented as mean ± SD with values from individual mice (n = 3). Expression and *P*-values
were calculated from RNA-Seq data using DESeq2 (Wald test with Benjamini-Hochberg correction). **C,**
Insulin content in *Dpy30*-WT and -KO islets. Data presented as mean ± SD with values from individual mice
(n = 8 WT, 6 KO). *P*-value calculated by two-tailed t-test with Welch's correction. **D,** Representative
transmission electron micrographs showing insulin granules in β-cells from a *Dpy30*-WT and a -KO mouse.
Scale bars: 2 μm. **E,** Quantification of median insulin core granule size and density. Data presented as
mean ± SD with values from individual mice (n = 3). *P*-values calculated by two-tailed t-test with Welch's
correction. **F,** Overrepresentation analysis of Gene Ontology: biological process terms in differentially
expressed genes 45-days after tamoxifen administration. *P*-values represent EASE scores, a modified
Fisher's exact *P*-value, calculated using the Database for Annotation, Visualization, and Integrated
Discovery (DAVID) v.6.8 [87]. **G,** Insulin secretion from *Dpy30*-WT and -KO islets during static *in vitro*
stimulation with glucose and KCl solutions, normalized to islet insulin content. *P*-values calculated by
multiple two-tailed t-tests with Welch's and Benjamini-Hochberg corrections; only values ≤ 0.05 are
shown. **H,** Cytosolic Ca^{2+} concentration in islets from *Dpy30*-WT and -KO mice during *in vitro* perifusion of
glucose and KCl solutions. *P*-values calculated for AUC in each time block by one-way ANOVA between
genotypes (n = 4 WT, 3 KO); only values ≤ 0.05 are shown. **I,** Oxygen consumption rate in *Dpy30*-WT and -
KO dispersed islet cells during treatment with the indicated compounds. n = 3 WT, 4 KO. **J,** Mitochondrial
respiration in *Dpy30*-WT and -KO islet cells in 16.7 mM glucose and their maximal respiration capacity,
inferred from the data shown in panel I. *P*-values calculated by two-tailed t-test with Welch's correction
(n = 3 WT, 4 KO).
926
Figure 3. TSS-associated H3K4 methylation maintains gene expression in mature beta cells.
A, Average enrichment profiles of H3K4me3 and H3K4me1 at the TSS of all expressed genes in *Dpy30*-WT
and -KO chromatin. **B-C,** Scatterplot of H3K4me3 (B) and H3K4me1 (C) enrichment in 10-kb bins spanning
the genome in *Dpy30*-KO versus -WT cells. **D,** Scatterplot showing spike-in normalized gene expression in
Dpy30-KO versus -WT cells. **E,** Genome browser views show examples of genes that retain H3K4me3 (i),
lose H3K4me3 but retain H3K4me1 (ii), or lose H3K4me3 and H3K4me1 (iii) from the TSS in *Dpy30*-KO
chromatin. **F,** Box and whisker plot showing $\log_2$(fold-change) of RNA expression for gene groups
described in panel E. *P* indicates the Wilcoxon signed-rank test result with Benjamini-Hochberg correction.
G, Contour scatterplot showing $\log_2$(fold-change) of H3K4me1 enrichment (x-axis) plotted against
$\log_2$(fold-change) of H3K4me3 enrichment (y-axis) in the TSS ±1 kb of all expressed genes. The $\log_2$(fold-
change) of RNA expression of the associated genes is shown as a colour gradient.
938
Figure 4. Weakly active and developmentally regulated genes are susceptible to downregulation in
H3K4me3-deficient beta cells.
A, Average enrichment profiles of H3K4me3, H3K4me1, H3K27ac, and H3K27me3 in *Dpy30*-WT cells at
TSSs of genes that are stably expressed or downregulated by *Dpy30*-KO. **B,** Box and whiskers plots showing
quantification of data shown in panel A. **C,** Box and whiskers plot showing mRNA expression in *Dpy30*-WT
cells for genes stably expressed or downregulated by *Dpy30*-KO. **D,** Average %GC content profile in the
gene body ±5 kb of genes that are stably expressed or downregulated by *Dpy30*-KO. **E,** Box and whiskers
plot showing quantification of data shown in panel D. **F,** Running enrichment plot for imprinted genes (ref.
[88]) and genes that are up- or downregulated during mouse β-cell maturation (adults versus postnatal day
10, ref. [46]), and mature β-cell transcription factor genes. NES: normalized enrichment score. *P* values

calculated using Wilcoxon signed-rank test (panels B,C,E) or permutation test (F) with Benjamini-Hochberg
correction.
951

Figure 5. H3K4me3 deficiency leads to a less active and more repressed epigenome in mature beta cells.
A, Genome browser view of H3K4me3, H3K4me1, H3K27ac, and H3K27me3 in *Dpy30*-WT and -KO cells
showing preservation of mega base-scale organization of H3K27me3-positive PcG compartments. **B,**
Scatterplots of H3K27ac (top) and H3K27me3 (bottom) enrichment in 10-kb bins spanning the genome in
Dpy30-KO versus -WT cells. **C,** Average enrichment profiles of H3K27ac (top) and H3K27me3 (bottom) at
the TSS of all expressed genes in *Dpy30*-WT and -KO cells. **D,** Box and whisker plots showing the $\log_2$(fold-
change) of H3K27ac (top) and H3K27me3 (bottom) enrichment in the TSS of transcriptionally
downregulated, stable, or upregulated genes in *Dpy30*-KO compared to -WT cells. *P*-values calculated with
Wilcoxon rank sum test with Benjamini-Hochberg correction. **E,** Genome browser view of a
downregulated gene *Rab34* showing loss of H3K4me3, H3K4me1, H3K27ac, and RNA, and accumulation
of H3K27me3, in *Dpy30*-KO cells.
963

Figure 6. Reduction of transcriptional consistency in *Dpy30*-KO and diabetic beta cells.
A, UMAP visualization of β-cell transcriptomes from a *Dpy30*-KO mouse. Expression of *Tdtomato*, *Egfp*,
and their overlap are shown as a colour gradient. **B,** Pseudotime representation of A. **C,** Scatterplot
showing *Ins1* and *Ins2* expression in β-cells plotted against pseudotime. **D,** Scatterplot showing gene
expression entropy scores of β-cells plotted against pseudotime. **E,** Box and whisker plot showing the
transcriptional entropy of β-cells in *Tdtomato+* and *Egfp+* clusters (n = 274 and 2,323 cells, respectively).
F, Box and whisker plot showing the variance/mean ratio of gene expression across β-cells in *Tdtomato+*
and *Egfp+* clusters. **G,** Variance/mean ratio of β-cell gene expression in *Egfp+* versus *Tdtomato+* clusters
(y-axis) plotted as a function of H3K4me3 breadth in quantiles (x-axis) (622 genes/quantile). Data
presented as mean ± SD. **H-I,** same as F-G in β-cells from human donors with or without T2D, using
H3K4me3 ChIP-seq data from GSE50386 (ref. [51]) and scRNA-seq data from GSE124742 ref. [52]).
975

Figure 7. H3K4me3 peak breadth encodes gene expression dynamics in a mouse model of type 2
diabetes.
A, H3K4me3 peaks in *Dpy30*-WT β-cells ranked from narrow to broad. Peaks associated with β-cell
transcription factor genes are labeled. **B,** Genome browser views showing H3K4me3 enrichment in WT β-
cells at a housekeeping gene (*Rplp0*), and an expression-matched β-cell transcription factor gene (*Nkx6-*
1). **C,** Enrichment *P*-values (y-axis) of genes down- or upregulated in islets from *Lepr^db/db* mice, where genes
have been ranked and grouped by H3K4me3 peak breadth from narrow to broad (x-axis). Each group
contains 622 genes. *P*-values were calculated using one-sided Fisher's exact test. **D,** Immunoblots of
H3K4me3 and total histone H3 in islet lysate from *Lepr^+/+* and *Lepr^db/db* mice. H3-normalized H3K4me3
band densities are listed. **E,** Average TSS enrichment profile of H3K4me3 in *Lepr^+/+* and *Lepr^db/db* islets. **F,**
Genome browser views of H3K4me3 in *Lepr^+/+* or *Lepr^db/db* islets at notable genes that are downregulated
(*Pdx1*, *Slc30a8*) or induced (*Aldh1a3*, *Ldha*) in *Lepr^db/db* islets. **G,** Enrichment *P*-values of H3K4me3 peaks
showing significant reduction or expansion of peak breadth in *Lepr^db/db* versus *Lepr^+/+* islets (y-axis) in the
different gene groups ranked from narrow to broad (x-axis). Enrichment *P*-values were calculated using
one-sided Fisher's exact test. **H,** Box and whisker plots showing the $\log_2$(fold-change) of H3K4me3 peak
breadth for genes that are downregulated, stably expressed, or upregulated in *Lepr^db/db* islets. *P*-values
calculated with Wilcoxon rank sum test with Benjamini-Hochberg correction. **I,** Stratified rank/rank

hypergeometric overlap plot comparing gene expression changes caused by *Dpy30*-KO (x-axis) and by
Lepr^{db/db} (y-axis) mutations. Colourscale shows the enrichment *P*-value calculated using the RRHO2 R
package [59]. **J,** Same as panel C for genes that are differentially expressed in *Dpy30*-KO cells.
996
Figure S1. Deletion of *Dpy30* in mouse β-cells using *Ins1^{Cre}* results in reduction of H3K4 methylation,
hyperglycemia, hypoinsulinemia, and impaired glucose tolerance.
A, Immunoblots of H3K4me3, H3K4me1, and total histone H3 in islet lysate from 5-week-old *Dpy30*-WT,
-KO, and -HET mice. **B,** Bar graphs showing band intensities of data in panel A. *P*-values were calculated
using one-way ANOVA with Tukey's multiple comparisons correction; n = 4. **C-D,** Unfasted blood glucose
concentration (C) and body mass (D) of *Dpy30*-WT, -KO, and -HET mice during 4-10 weeks of age. Values
from each mouse and time point are shown (n = 10 WT, 15 KO, 7 HET, however tracking was stopped after
1-2 blood glucose reading ≥ 20 mM). **E-F,** Blood glucose (E) and serum insulin (F) concentration during
intraperitoneal glucose tolerance tests in 5-week-old *Dpy30*-WT, -KO, and -HET mice. *P*-values were
calculated by comparison of *Dpy30*-KO versus -HET (top, teal) or -KO versus -WT (bottom, grey) AUC's
using one-way ANOVA with Tukey's multiple comparisons correction; only values ≤ 0.05 are shown. WT:
Ins1^{+/+}; *Dpy30^{flox/flox}*. KO: *Ins1^{Cre/+}*; *Dpy30^{flox/flox}*. HET: *Ins1^{Cre/+}*; *Dpy30^{flox/+}*.
1009
Figure S2. Genes involved in insulin production and glucose-induced activity are regulated by H3K4me3.
A, Heatmap showing expression Z-scores in RNA-seq data from *Dpy30*-WT and -KO 45-days post-
tamoxifen for selected genes important for glucose metabolism, insulin production or secretion. *P*-values
calculated by Wald test with Benjamini-Hochberg correction; n = 3. **B,** Genome browser views of
H3K4me3, H3K4me1, and RNA in *Dpy30*-WT and -KO chromatin at selected genes. **C,** Bar graph showing
the fraction of cytoplasm area occupied by mitochondria in transmission electron micrographs of *Dpy30*-
WT and -KO β-cells. *P*-value calculated using two-tailed t-test with Welch's correction. **D,** Representative
immunofluorescence images showing EdU incorporation in *Dpy30*-WT and -KO mouse islets that were
dispersed and cultured in 5.5 mM or 16.7 mM glucose for 48-hours, as indicated, showing EdU (white)
and DAPI (white) staining and eGFP (green) and tdTomato (red) endogenous fluorescence. Scale bars: 100
μm. **E,** Bar graph showing quantification of EdU+ eGFP+ double positive cells as a fraction of total eGFP+
cells (n = 3 per treatment and genotype). *P*-values calculated using two-way ANOVA with Tukey's
correction. ns: not significant (*P* > 0.05).
1023
Figure S3. H3K27ac peak dynamics are linked to gene expression dynamics in *Dpy30*-KO cells.
A, Pie chart showing the number of H3K27ac peaks that are increased >1.5-fold, decreased >1.5-fold, or
<1.5-fold changed for enrichment of H3K27ac in *Dpy30*-KO versus -WT chromatin. **B,** Association
frequency of H3K27ac peaks classified in panel A with genes that are upregulated or downregulated in
Dpy30-KO cells. Association was tested between H3K27ac peaks with the nearest active TSS and compared
with association frequencies after randomizing peak locations 100 times. ** *P* < 0.01 by permutation test.
1030
Figure S4. *Dpy30*-KO β-cells do not activate genes associated with dedifferentiated β-cells.
A, UMAP visualization of islet cell transcriptomes from a *Dpy30*-WT and -KO mouse. Markers of β-cells
(*Ins1*), α-cells (*Gcg*), and δ-cells (*Sst*) are shown as a colour gradient (n = 3,703 WT cells, 3,641 KO cells).
B, Violin plots showing expression *Ins1*, *Ins2*, *Iapp*, *Glp1r*, *Slc2a2*, and *G6pc2* in *Dpy30*-WT and -KO β-cells
(n = 2,510 WT, 2,597 KO β-cells). **C,** Lowess curves of gene expression of the indicated maturity and
immaturity genes in *Dpy30*-KO β-cells along pseudotime defined in Fig 6B.

1037

Figure S5. H3K4me3 peak breadth stratifies β-cell gene sets more effectively than gene expression level.

A, H3K4me3 peaks ranked by increasing breadth (x-axis) plotted against RNA expression of the associated

gene (y-axis). R indicates Spearman's rank correlation coefficient. **B,** Overrepresentation analysis of

expressed genes that are ranked and grouped by H3K4me3 peak breadth from narrow to broad (blue) or

ranked and grouped by gene expression level from low to high expression (green). The number in each

square indicates the number of genes of each geneset matching to each quantile. Enrichment *P*-values

calculated using two-sided Fisher's exact tests. **C,** H3K4me3 peaks in human β-cells [51] ranked by peak

breadth. Peaks associated with β-cell transcription factor genes are labeled. **D,** H3K4me3 peaks ranked by

breadth (x-axis) plotted against RNA expression [58] of the associated gene (y-axis) in non-diabetic human

islets. R indicates Spearman's rank correlation coefficient. **E,** Overrepresentation analysis of human islet

genes that are ranked and grouped by H3K4me3 peak breadth from narrow to broad (blue) or ranked and

grouped by gene expression level from low to high expression (green). The number in each square

indicates the number of genes of each gene set matching to each quantile. Each quantile includes 783

genes. Enrichment *P*-values calculated using one-sided Fisher's exact test. **F,** Running enrichment plot of

genes disallowed in islets [54] and β-cell-enriched transcription factor genes in *Lepr$^{db/db}$* RNA-seq data [56] .

NES: normalized enrichment score. *P*-values calculated by permutation test.

Figure 1

Figure 2

Figure 3

Figure 4

A

Gene expression (*Dpy30*-KO/WT): —— Stable —— Downregulated

H3K4me3 H3K4me1 H3K27ac H3K27me3

Signal in *Dpy30*-WT

B

☐ Stable ☐ Downregulated

H3K4me3 H3K4me1 H3K27ac H3K27me3

Signal in TSS ±3kb in *Dpy30*-WT

$P = 1.445 \times 10^{-83}$ $P = 8.025 \times 10^{-55}$ $P = 7.853 \times 10^{-48}$ $P = 9.064 \times 10^{-46}$

C

RNA

Gene expression in *Dpy30*-WT

$P = 1.978 \times 10^{-26}$

D

GC Content

Mean %GC

−5kb TSS TES +5kb

E

GC Content

%GC in gene body ±5kb

$P = 2.686 \times 10^{-61}$

F

Enrichment score

Gene rank

Up-regulated <----> Down-regulated

—— Imprinted
NES = −1.731; $q = 7.82 \times 10^{-4}$

—— Up in β-cell maturation
NES = −1.799; $q = 5.850 \times 10^{-4}$

—— Down in β-cell maturation
NES = −2.045; $q = 2.567 \times 10^{-5}$

—— β-cell transcription factors
NES = −1.708; $q = 1.252 \times 10^{-3}$

Figure 5

Figure 6

Figure 7

Figure S1

Figure S2

A

B

C

Figure S3

A

H3K27ac peak intensity (KO/WT)

B

H3K27ac peak intensity (KO/WT)

Figure S4

A

B

C

Figure S5

A

R = 0.4287

log₁₀ (Gene expression) vs H3K4me3 peak rank

B

🔵 Genes ranked & grouped by promoter H3K4me3 peak breadth
🟢 Genes ranked & grouped by RNA expression level

Enrichment −log₁₀ (*P*-value)

Number of geneset genes in quantile

Housekeeping

🔵	3	5	5	9	7	5	11	5	16	14	18	24	19	19	16	16	17	30	20	40
🟢	2	2	0	1	6	1	5	4	4	1	5	12	7	6	10	16	24	29	52	112

0 — 80

Endocrine pancreas development

🔵	2	0	0	0	0	0	1	0	0	0	0	1	4	1	3	4	0	1	0	17
🟢	2	1	0	2	1	1	1	1	0	1	1	2	1	1	1	2	3	2	7	4

0 — 15

Maturity onset diabetes of the young

🔵	1	0	1	0	0	0	0	0	0	0	1	0	2	0	0	0	3	2	1	12
🟢	1	0	1	1	0	1	1	1	1	1	0	1	1	0	1	0	0	2	1	8

0 — 10

Downregulated in *Lepr*^db/db islets

🔵	13	12	16	16	15	20	18	16	13	19	18	15	26	20	27	26	28	29	40	67
🟢	2	5	16	32	33	23	20	22	18	22	25	25	22	24	19	15	18	19	37	57

0 — 15

Upregulated in *Lepr*^db/db islets

🔵	72	55	45	21	28	19	23	21	26	12	14	25	23	16	9	13	15	21	13	13
🟢	39	58	67	48	49	22	16	25	28	9	15	11	7	11	13	9	6	12	19	20

0 — 20

Quantile: 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20

Increasing breadth or expression ⟶

C

H3K4me3 peak breadth (kb) vs H3K4me3 peak rank

MEIS2, NKX2-2, ISL1, PDX1, PAX6, ST18, MAFB, NKX6-1, INSM1, MAFA, FOXO1, NEUROD1, MYT1L, HNF1B, FOXA2

D

R = 0.4274

log₁₀ (Gene expression) vs H3K4me3 breadth quantile

E

🔵 Genes ranked & grouped by promoter H3K4me3 peak breadth
🟢 Genes ranked & grouped by RNA expression level

Enrichment −log₁₀ (*P*-value)

Number of geneset genes in quantile

Housekeeping

🔵	3	4	6	6	12	7	19	20	18	14	16	12	16	18	24	23	36	27	28	42
🟢	0	0	0	1	0	2	1	2	2	3	7	3	4	9	11	15	20	36	43	192

0 — 200

Endocrine pancreas development

🔵	0	1	1	1	1	2	1	1	1	0	2	1	1	0	1	5	0	2	2	17
🟢	0	1	0	0	0	2	3	1	0	2	5	2	1	1	3	5	3	7	4	0

0 — 15

Maturity onset diabetes of the young

🔵	0	0	1	0	0	1	1	1	0	0	1	3	0	0	0	0	2	1	1	10
🟢	0	0	0	1	0	0	2	0	0	3	4	1	2	1	2	0	1	1	2	2

0 — 8

Downregulated in T2D islets

🔵	1	5	5	12	9	13	16	10	12	14	13	26	20	22	15	21	32	27	29	59
🟢	0	0	4	4	12	13	16	16	17	19	13	26	27	24	22	24	38	26	37	23

0 — 15

Upregulated in T2D islets

🔵	18	26	24	25	19	9	23	16	21	12	14	9	16	13	23	12	12	8	12	13
🟢	0	0	14	16	17	27	23	17	17	28	14	19	12	17	10	11	14	20	20	29

0 — 3

Quantile: 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20

Increasing breadth or expression ⟶

F

Lepr^db/db vs *Lepr*^+/+

Enrichment score vs Gene rank

NES = 2.201
P = 2.035 × 10⁻⁶

NES = −2.102
P = 2.125 × 10⁻⁵

— disallowed in islets
— β-cell-enriched transcription factors