INFERRING LARGE-SCALE STRUCTURAL CONNECTIVITY FROM LOCAL ACTIVITY IN A MODEL OF THE MOUSE LARGE-SCALE CORTICAL NETWORK

Ronaldo V. Nunes[1], Marcelo Bussotti Reyes[1], Jorge F. Mejias[2], and Raphael Y. de Camargo[1]

[1]Center for Mathematics, Computing and Cognition, Universidade Federal do ABC, São Bernardo do Campo, Brazil

[2]Swammerdam Institute for Life Sciences, University of Amsterdam, Amsterdam, The Netherlands

March 11, 2021

ABSTRACT

Inferring the structural connectivity from electrophysiological measurements is a fundamental challenge in systems neuroscience. Directed functional connectivity measures, such as the Generalized Partial Directed Coherence (GPDC), provide estimates of the causal influence between areas. However, the relation between causality estimates and structural connectivity is still not clear. We analyzed this problem by evaluating the effectiveness of GPDC to estimate the connectivity of a ground-truth, data-constrained computational model of a large-scale network model of the mouse cortex. The model contains 19 cortical areas comprised of spiking neurons, with areas connected by long-range projections with weights obtained from a tract-tracing cortical connectome. We show that GPDC values provide a reasonable estimate of structural connectivity, with a Pearson correlation of 0.73. Moreover,

even in a typical electrophysiological recording scenario containing five areas, the

mean correlation was above 0.6. These results indicate that GPDC estimates provide

statistical information on structural connections, but individual connection estimates

are not reliable.

1   Introduction

The communication between brain regions is often analyzed through structural and functional

connectivity [1]. The former refers to anatomical connections between brain regions generally

quantified using tracer injections or diffusion magnetic resonance imaging [2]. The map of these

connections is called "connectome" [3], whose analysis includes methods to evaluate the networks

defined by nodes (brain regions) and edges (synapses) [4, 5]. Functional connectivity evaluates

brain communication from statistical relations between recorded brain signals [6, 1]. Particularly,

directed functional connectivity methods use the concept of causality to infer both the intensity and

the direction of the connections between brain regions [7]. Even though there is some association

between structural and functional connectivity, the relationship between them is not straightfor-

ward [1]. While the former is practically static and compose the map of possible pathways for

information flow between brain regions, the latter changes continuously and depends, for example,

on the dynamical states of brain regions, noise, and strength of structural connections [8].

During electrophysiological procedures, researchers typically record brain signals using electrodes

positioned in different depths of brain regions. Even with the improvement in technologies for

recording signals, it is usually possible to record signals only from a few areas compared to the

number of sources of activity in the brain [9, 10, 11]. Thus, the functional connectivity analysis

presents a problem because many unrecorded regions may indirectly influence other regions as

common inputs [7, 12, 6]. Therefore, the comparison between structural and functional connectivity

becomes more complicated since spurious inferred causality relations can lead to misinterpretations

of electrophysiological data.

Some simulation studies evaluated the relation between directed functional connectivity and structural connections [8, 13, 14, 15]. However, most of these studies used either autoregressive models for the dynamics of each node [16] or rate-based models [13]. These studies provided important steps towards evaluating the reliability of causality measures, but they are still distant from laboratory experimental conditions.

In this work, we investigate the relationship between directed functional connectivity and structural connectivity in a large-scale network model of the cortex, derived from a cortical connectome of the mouse obtained using tracer injections [17]. Causality measures obtained using GPDC explained most of the variance of structural connection strengths. The mean correlation remained high ($r > 0.6$) even when only a few cortical areas were considered in the GPDC calculation, indicating that this causality measure provides reliable results in typical experimental conditions in which only recordings from a subset of areas are available.

2 Methods

2.1 Neuron model

We modeled the neurons using a single-compartment Hodgkin–Huxley-type model, where the membrane potential of the i-th neuron described by,

$$C_m \frac{dV_i}{dt} = -g_{Na} m_i^3 h_i (V - E_{Na}) - g_K n_i^4 (V - E_k) - g_L (V - E_L) + I_{\text{ext},i} + I_{\text{syn},i}, \quad (1)$$

the membrane capacitance C_m is 0.50 nF (0.125 nF) for excitatory (inhibitory) neurons. The maximal conductances values were $g_{Na} = 12.5$ μS, $g_K = 4.74$ μS and $g_L = 0.025$ μS. The reversal potentials $E_{Na} = 40$ mV, $E_K = -80$ mV, and $E_L = -65$ mV correspond to the sodium, potassium and leakage channel, respectively [18]. The dynamics of the voltage-gated ion channels are described by activation and inactivation variables m, n, and h, where m and n accounts for the

dynamics of Na channels and h for K channels. The probability that an ion channel is open evolves according to a set of ordinary differential equations [19],

$$\frac{dm}{dt} = \alpha_m(V)(1-m) - \beta_m(V)m, \tag{2}$$

$$\frac{dh}{dt} = \alpha_h(V)(1-h) - \beta_h(V)h, \tag{3}$$

$$\frac{dn}{dt} = \alpha_n(V)(1-m) - \beta_n(V)n, \tag{4}$$

where,

$$\alpha_m(V) = 0.1\frac{(V+16)}{1-\exp(-(V+16)/10)}, \tag{5}$$

$$\beta_m(V) = 4\exp(-(V+41)/18), \tag{6}$$

$$\alpha_h(V) = 0.07\exp(-(V+30)/20), \tag{7}$$

$$\beta_h(V) = [1+\exp(-V/10)]^{-1}, \tag{8}$$

$$\alpha_n(V) = 0.01\frac{(V+20)}{1-\exp(-(V+20)/10)}, \tag{9}$$

$$\beta_h(V) = 0.125\exp(-(V+30)/80), \tag{10}$$

The parameters used in this neuron model was previously reported and applied in some studies that modeled cortical neuronal populations. [20, 19, 21]

2.2 Spiking neuronal population model

Each spiking neuronal population was composed of 2000 neurons, 1600 excitatory and 400 inhibitory. Neurons within each spiking neuronal population were randomly connected with probability $p_{\text{intra}} = 10\%$. The synaptic current I_{syn} that arrives to postsynaptic neuron i is modeled by,

$$I_{\text{syn},i}(t) = \sum_{j \in \text{presyn}} g_{\text{syn},i,j}(t) \left[E_{\text{syn}} - V_i(t) \right], \tag{11}$$

68 where the index j represent a presynaptic neuron connected to neuron i and the sum over j accounts

69 for all the synapses that impinge on neuron i. E_{syn} is the synaptic reversal potential which is 0 mV

70 for excitatory and -70 mV for inhibitory synapses. The dynamics of synaptic conductance $g_{\text{syn},i,j}$

71 is described by an exponential function as follows [22],

$$\frac{dg_{\text{syn},i,j}}{dt} = -\frac{g_{\text{syn},i,j}}{\tau}. \tag{12}$$

When a presynaptic neuron j fires a spike, a synaptic weight w is added on the synaptic conductance

$g_{\text{syn},i,j}$ after the axonal delay, which was set as 1 ms [19]. The value of w depends on the excita-

tory/inhibitory nature of the presynaptic and postsynaptic neurons. The synaptic time constant τ is

2 ms and 8 ms for excitatory and inhibitory synapses, respectively. Furthermore, all neurons receive

a background input given by a heterogeneous Poisson-process spiking activity with a rate of 7.3

kHz [19]. The effect of the background input in each neuron is an excitatory synaptic current. To

add heterogeneity in our model, all synaptic weights w for recurrent connections and background

input were taken from a Gaussian distribution (Table 1).

Table 1: **Synaptic weights for intra-areal connections**. Mean synaptic weight $\overline{w}$ and standard deviation σ_w for all possible synapses. E, I, and Input represent excitatory neurons, inhibitory neurons, and external input, respectively. The arrow indicates the direction of the connection.

Synapses	$\overline{w}$ (nS)	σ_w (nS)
$E \rightarrow E$	2.5	1.0
$E \rightarrow I$	2.5	1.0
$I \rightarrow E$	240	10
$I \rightarrow I$	240	10
$Input \rightarrow E$	3.2	1.0
$Input \rightarrow I$	3.2	1.0

2.3 Mouse large-scale cortical network

The mouse cortex's large-scale network model comprises 19 cortical areas where a spiking neuronal population models each area with long-range and recurrent synapses. Parameters related to recurrent synapses were described in the previous session. Neurons from different areas are randomly connected with probability $p_{\text{inter}} = 5\%$. The synaptic weights between cortical areas are based on the recently published anatomical connectivity dataset for the mouse cortex [17] obtained by retrograde tracer injections [23]. This technique consists in injecting a tracer that flows from the target area to the cell bodies, allowing to identify neurons projecting to the target area. The Fraction of Labeled Neurons (FLN) was measured as the ratio of the number of labeled neurons in a source area to the total quantity of labeled neurons in all source areas [24, 25]. The synaptic weights for directed long-range connections are the FLNs scaled by the global scaling parameters $\mu_{\text{E}} = 50$ and $\mu_{\text{I}} = 25$,

$$w_{\text{lr,E}}^i = \mu_{\text{E}} \sum_j^N \text{FLN}_{i,j} \tag{13}$$

$$w_{\text{lr,I}}^i = \mu_{\text{I}} \sum_j^N \text{FLN}_{i,j}. \tag{14}$$

Long-range connections as excitatory, targeting either excitatory or inhibitory neurons with synaptic weight, $w_{\text{lr,E}}^i$, and $w_{\text{lr,I}}^i$, respectively. The index j represents the source area, i the target area, and N is the total number of simulated cortical areas. The axonal delay for long-range connections is given by the ratio between the anatomical distance estimates between cortical areas and the conduction speed set as 3.5 m/s [26].

2.4 LFP signal

We computed the local field potential (LFP) signal as a sum of the currents' absolute values acting upon excitatory neurons in a spiking neuronal population [27, 28]. Thus, for a cortical area in our

100 model, the LFP signal will be given by,

$$\text{LFP} = R\frac{\sum_i^{N_\text{E}}(|I_{\text{E},i}| + |I_{\text{I},i}| + |I_{\text{bkg},i}|)}{N_\text{E}}, \tag{15}$$

$I_{\text{E},i}$ accounts for both the local (within population) and global (inter-areal projections) excitatory

synaptic currents, while $I_{\text{I},i}$ corresponds to the local inhibitory current. $I_{\text{bkg},i}$ is the synaptic current

related to the background Poisson input. R represents the resistance of a typical electrode used for

extracellular measurements, here chosen to be $1\ \text{M}\Omega$ [19]. N_E is the number of excitatory neurons

in each neuronal population.

The mean was subtracted from the simulated LFP signal. The resultant signal was filtered using a

low-pass filter in 1 kHz to avoid aliasing and downsampled to 1 kHz.

2.5 Generalized partial directed coherence

109 Generalized partial directed coherence (GPDC) is a frequency-domain method of directed functional

110 connectivity established on multivariate vector autoregressive (MVAR) model [29]. Considering

111 that LFP signals from all areas in the model are represented by a set $\mathbf{x}(t) = [x_1(t)\cdots x_N(t)]^T$ of

112 simultaneously observed time series. The MVAR model for $\mathbf{x}(t)$ is defined as:

$$\mathbf{x}(t) = \sum_{k=1}^{p} \mathbf{A}_k\mathbf{x}(t-k) + \epsilon(t) \tag{16}$$

113 where p is the MVAR model order. $\mathbf{A}_k$ are coefficient matrices in which the element $a_{ij}^{(k)}$ define the

114 effect of $x_j(t-k)$ on $x_i(t)$. The term $\epsilon(t)$ is a vector of N white noises with covariance matrix $\mathbf{\Sigma}$.

115 The GPDC from the time series x_j to the time series x_i at frequency λ is defined as,

$$\text{GPDC}_{ij}(\lambda) = \left|\frac{\frac{1}{\sigma_i}\overline{A}_{ij}(\lambda)}{\sqrt{\sum_{k=1}^{N}\frac{1}{\sigma_k^2}\overline{A}_{kj}(\lambda)\overline{A}_{kj}^*(\lambda)}}\right|^2, \tag{17}$$

116 where

$$\overline{A}_{ij}(\lambda) = \begin{cases} 1 - \sum_{k=1}^{p} A_{ij,k} e^{-2\pi\sqrt{-1}\lambda k}, & \text{if } i = j \\ -\sum_{k=1}^{p} A_{ij,k} e^{-2\pi\sqrt{-1}\lambda k}, & \text{if } i \neq j, \end{cases} \tag{18}$$

117 and σ_i^2 refers to the variance of white noise $\epsilon_i(t)$ [29]. λ is a normalized frequency where $\lambda = 0.5$

118 means one half of the sampling rate fs [30]. The MVAR model was estimated by the method of

119 ordinary least squares (OLS) [31]. Akaike's information criterion gave the model order indicated

120 that the best model order p that was lesser than or equal to 50 [30]. For all analysis it was evaluated

121 the peak of GPDC.

2.6   Weighted FLN and weighted GPDC

123 In order to investigate the propagation of neuronal activity through the links defined by structural

124 and directed functional connectivity, we defined the weighted FLN (wFLN) and the weighted GPDC

125 (wGPDC) [32],

$$\text{wFLN}_{ij} = \frac{1}{n} \sum_{j=1}^{n} \left(\frac{\text{FLN}_{ij} r_j}{\sum_{j \in S} r_j} \right) \tag{19}$$

$$\text{wGPDC}_{ij} = \frac{1}{n} \sum_{j=1}^{n} \left(\frac{\text{GPDC}_{ij} r_j}{\sum_{j \in S} r_j} \right) \tag{20}$$

126 where FLN_{ij} is the FLN from area j to area i, r_j is the firing rate for area j, GPDC_{ij} is the peak of

127 GPDC from area j to area i, S is the set of source areas for target area i and finally and n is the

128 cardinatility of set S.

2.7   Numerical simulations

130 All simulations were performed using the simulator Brian2 [33] applying the exponential Euler

131 method [34] to integrate the differential equations with an integration step of 0.1 ms. Each simulating

132 was 30 s long, generating sufficient data points to apply GPDC on the simulated LFP signals [35].

3 Results

The large-scale network model of the mouse cortex contains 19 spiking neural populations with recurrent connections and excitatory long-range connections between populations, constrained by the directed and weighted structural connectome (Figure 1A and Figure 1B). The dynamical behavior of each simulated cortical area is predominantly asynchronous with transient spike synchronization [36, 37] (Figure 1C), with the typical power spectral density (PSD) of LFP signals displaying a peak in the gamma band (Figure 1D and 1E) [38]. The firing rate of inhibitory neurons is 4.74 ± 0.11, higher than excitatory neurons rate of 3.64 ± 0.42 (Figure 1F). Differences in population behavior are mostly due to inputs from other areas since we sample their parameters from the same distributions (Section 2.2).

Figure 1: **Large-scale cortical network.** (A) Local neuronal population where E and I are populations of spiking neurons. (B) Map of structural connectivity given by the FLNs. These values define the strength of long-range projections in the large-scale network model. (C) Raster plot of $50ms$ of activity for each cortical area. (D) Simulated LFP signal for an area in the large-scale network model. (E) Power spectral density for simulated LFP signals from one area. Continuous black line corresponds to the average over ten simulations, and the gray shaded area delimits its standard deviation. (F) Firing-rate for excitatory (blue) and inhibitory (red) populations computed using a sliding window of $100ms$. The continuous line corresponds to an average firing-rate over ten simulations, and the shaded area is the standard deviation. To exemplify, we used data from area MOp in (D), (E), and (F).

We first compared the FLN values to the average GPDC over ten simulations of the model. Most medium to strong connections from the structural connectome were also captured by the directed

functional connectivity (Figure 2A and Figure 2B). We used the GPDC largest value (peak), but other approaches such as the average of GPDC over frequencies and area under the GPDC curve (Supplementary Figure S1) produced similar results.

Although the graph density of the structural connectome is 97% [17], most structural connections are weak, which leads to a prevalence of weak average GPDC values. Weak structural connections are characteristics shared by connectomes from different mammals, with FLNs varying by several orders of magnitude, log-normally distributed [23, 39, 17, 40]. To evaluate the relation between structural and directed functional connectivity, we plotted GPDC values from ten simulations against FLNs and fitted a linear model, obtaining the Pearson correlation r (Figure 2C). The scatter plot presents most points close to the origin due to the predominance of small values for the GPDC and FLN. Interestingly, for some FLNs, there is a high variation of GPDC values over different simulations. The Pearson correlation between FLN and GPDC is 0.73, and GPDC explains approximately 54% of the variance in FLN values. This explainability level similar to those obtained by other works that analyzed different structural connectomes using functional connectivity applied to empirical data [41, 42] or firing-rate models [43].

Figure 2: **Relation between structural and directed functional connectivity.** (A) Map of structural connectivity given by the FLNs. (B) Map of directed functional connectivity given by the average of GPDC peaks over ten simulations. (C) Scatter plot of FLNs *versus* GPDC peak. The red line corresponds to the linear fit. The Pearson correlation between FLNs and GPDC is 0.73.

We also investigated the propagation of activity between cortical areas through the pathways defined by structural connectivity. The propagation of activity in the cortex is constrained by direct anatomical connections between areas and indirect paths [44], with the propagation of activity

occurring mainly through the strongest long-range projections [24]. The weighted FLNs are strongly correlated to the target areas' firing rate (Figure 3A), while the correlation of weighted GPDCs and firing rates was 0.54 (Figure 3B). This indicates that neural activity propagation is directly dependent on the strength of structural connections and that GPDC can capture part of this activity propagation.

Figure 3: **Propagation of activity through structural and directed functional pathways**. (A) Weighted FLN *versus* firing-rate for target areas ($r = 0.94$). (B) Weighted GPDC *versus* firing-rate for target areas ($r = 0.54$). Red lines are linear fits.

We analyzed the behavior of GPDC estimates when considering a reduced number of areas, reproducing typical experimental setups. We considered a visual and a frontoparietal cluster, each containing 7 cortical areas [17]. We evaluated the GPDC estimates between all areas of each cluster (Figures 4C and 4F), conditioned on the whole connectome (Figures 4B and 4E), conditioned on the areas in each cluster, and using only pairwise (bivariate) estimates (Figures 4D and 4G). This analysis simulates the situations where an electrophysiologist only has information from a single cluster of cortical areas or a pair of areas. The highest correlations between the GPDC and FLN occurred when we conditioned GPDC to the whole connectome, followed by GPDC conditioned to the cluster area, and pairwise GPDC. Also, the correlation for the frontoparietal cluster was higher in all scenarios.

We extended the analysis to evaluate the effect of cluster size on GPDC correlation to FLN. We used cluster sizes ranging from 2 to 15 areas. We created 150 random clusters sampled from all areas in the connectome for each cluster size and computed the Pearson correlation for the GPDC

Figure 4: **Correlation between FLN and GPDC for the visual and frontoparietal clusters.** (A) Graph representing the mouse cortical connectome. Nodes represents cortical areas and edges, directed long-range projections between them. Green nodes are cortical areas belonging to the frontoparietal cluster. Red nodes are cortical areas belonging to the visual cluster. Each cluster contains 7 cortical areas. (B-D) FLN *versus* GPDC for the visual cluster, when (B) conditioned to the whole connectome, (C) conditioned to the areas in the cluster, and (D) using bivariate data. (E-G) FLN *versus* GPDC for the frontoparietal cluster when (E) conditioned to the whole connectome, (F) conditioned to the areas in the cluster, and (G) using bivariate data. Red lines are linear fits and r is the Pearson correlation.

(A) conditioned on the whole connectome, (B) conditioned on the cluster areas, and (C) evaluated using pairwise data. For cases (A) and (B), the Pearson correlation increases, and the standard variation decreases as we increase the cluster size (Figure 5), showing that it is advantageous to include more areas in the GPDC calculation. Surprisingly, the correlation between structural and directed functional connectivity when using signals measured in a few cortex areas (blue dots) is similar to using signals from the whole cortex (black dots), with few points showing statistically different results. The correlation between whole connectome GPDC and cluster GPDC estimates increases for large clusters, as expected (Supplementary Figure S2). The bivariate GPDC (Figure 5) had a statistically significant lower average Pearson correlation for all cluster sizes with four or more areas, indicating that these measures are affected by interference from ignored signals.

Figure 5: **Pearson correlations between FLN and GPDC for the different cluster sizes with randomly chosen areas.**. The graph shows the average (dots) and error bars of the GPDC conditioned to the whole connectome (in black), to areas in the cluster (in blue), and to pairwise signals (in red). Stars represent statistically different averages when compared to the whole connectome (Holm-Bonferroni corrected Welch t-test).

4   Discussion

Our results shed light on the relationship between structural and directed functional connectivity in circumstances similar to those faced by electrophysiologists. They indicate that the reliability of directed functional connectivity estimates and their relationship with structural connectivity depends on the number of areas considered. Nevertheless, the GPDC conditioned on few cortical areas had similar results to the GPDC conditioned on all areas, providing evidence that it is possible to obtain a reasonable relationship between structural and directed functional connectivity in electrophysiological experiments even with signals from few areas.

Previous studies evaluated the relationship between structural and functional network connectivity strength on electrophysiological data [45], with some using undirected functional connectivity measures [46, 47]. But in electrophysiology studies researchers do not have access to signals from unrecorded areas and have only estimates of structural strengths from tracers. Using large-scale

network models solves this problem, as the researcher has access to all variables in the system,

allowing a better understanding of the obtained functional connectivity results.

The relationships between structural and functional connectivity has been largely unexplored through

large-scale network models [48], and the existing models use neural mass descriptions (rate models)

to describe each area's activity [49, 50]. However, information propagated between brain regions

can be characterized not only by the rate code but also by the temporal code [51, 52, 53, 54, 55],

and hypotheses are pointing to spike-timing and spike coherence as essential components of cortical

communication [56, 36, 57]. Spiking neuronal populations have richer dynamical behaviors than

rate models and better resemble cortical activity, including the effects of neuronal synchronization

and its effects on information propagation. Moreover, our obtained correlations are in the same range

of the studies using more complex electrophysiological data [45], indicating that our simulations

did not produce artificially high correlations due to model simplifications.

The propagation of activity between simulated cortical areas is strongly correlated with weighted

FLNs and moderately correlated with weighted GPDC estimates, when using the concept of

activity flow proposed by Cole *et al.* [32]. This result is interesting since it indicates that structural

connectivity is more relevant to activity propagation than direct functional connectivity extracted

from GPDC peak values. In other words, GPDC do not directly correspond to activity flow between

areas and should be used with caution in this case.

The relationship between structural and directed functional connectivity is the largest when GPDC

is conditioned to all areas in the connectome and and decreases as we reduce the number of areas.

Gămănut et al. identified 6 clusters in the mouse connectome (prefrontal, frontal, parietal, cingulate,

temporal and visual) [17] based on the same approach used to investigate the macaque cortex [58].

We evaluated the relationship between GPDC and FLNs in the visual cluster and in a combination

of the prefrontal, frontal and parietal clusters, which we called frontoparietal. We did not use the

other clusters, which had a small a small number of regions. The obtained correlation was in the

range of random clusters, with $r = 0.74$ for the frontoparietal and $r = 0.54$ for the visual cortex.

This indicates that within anatomical clusters the relationship between GPDC and FLNs does not change in relation to randomly selected areas. These results also show that GPDC estimates provide statistical information on structural connections even when considering a few areas, but individual connection estimates are not reliable.

It is well known that indirect connections and common input from areas not included in GPDC influence the estimates [7, 14, 6, 49], but the magnitude of this influence had not been quantified. We verified that the correlation between whole connectome GPDC and cluster GPDC increases with the number of areas considered in the measurement (Supplementary Figure S2A). For seven areas, the Pearson correlation between the whole connectome GPDC and cluster GPDC was 0.65, resulting in a explained variance of 42%. This shows that areas not considered in the GPDC calculations can have an important impact in the GPDC estimates. We also showed that the difference between the whole connectome GPDC and cluster GPDC values decreases with larger cluster sizes (Supplementary Figure S2B) as expected.

Our large-scale network model has some limitations. First, modeled neuronal population parameters are drawn from the same distributions with activity in the gamma band range (Figure 1). The activity of cortical areas in mice occurs in multiple frequency ranges [59, 60] and the relationship between structural and functional connectivity depends on the frequency [61]. A second limitation is that we do not model changes in network states, which are known to influence functional connectivity [62]. Some studies in computational neuroscience have already explored multistability and temporal patterns of functional connectivity [63, 64, 65]. Finally, we considered only cortical areas in our large-scale network model, excluding subcortical areas, which have a more complex dynamic [66, 67]. Future studies can overcome these limitations by creating richer spiking network models, with different operating frequencies and evolving neuronal dynamics. These models are difficult to create, but would allow one to compare functional connectivity values to structural connection strength in more dynamical settings.

References

[1] Andrea Avena-Koenigsberger, Bratislav Misic, and Olaf Sporns. Communication dynamics in complex brain networks. *Nature Reviews Neuroscience*, 19(1):17, 2018.

[2] Karen S Ambrosen, Simon F Eskildsen, Max Hinne, Kristine Krug, Henrik Lundell, Mikkel N Schmidt, Marcel AJ van Gerven, Morten Mørup, and Tim B Dyrby. Validation of structural brain connectivity networks: The impact of scanning parameters. *NeuroImage*, 204:116207, 2020.

[3] Olaf Sporns, Giulio Tononi, and Rolf Kötter. The human connectome: a structural description of the human brain. *PLoS Comput Biol*, 1(4):e42, 2005.

[4] Danielle S Bassett, Perry Zurn, and Joshua I Gold. On the nature and use of models in network neuroscience. *Nature Reviews Neuroscience*, 19(9):566–578, 2018.

[5] Danielle S Bassett and Olaf Sporns. Network neuroscience. *Nature neuroscience*, 20(3):353–364, 2017.

[6] Andrew T Reid, Drew B Headley, Ravi D Mill, Ruben Sanchez-Romero, Lucina Q Uddin, Daniele Marinazzo, Daniel J Lurie, Pedro A Valdés-Sosa, Stephen José Hanson, Bharat B Biswal, et al. Advancing functional connectivity research from association to causation. *Nature neuroscience*, 22(11):1751–1760, 2019.

[7] André M Bastos and Jan-Mathijs Schoffelen. A tutorial review of functional connectivity analysis methods and their interpretational pitfalls. *Frontiers in systems neuroscience*, 9:175, 2016.

[8] Ronaldo V Nunes, Marcelo B Reyes, and Raphael Y De Camargo. Evaluation of connectivity estimates using spiking neuronal network models. *Biological cybernetics*, 113(3):309–320, 2019.

[9] Guosong Hong and Charles M Lieber. Novel electrode technologies for neural recordings. *Nature Reviews Neuroscience*, 20(6):330–345, 2019.

[10] Kenneth D Harris, Rodrigo Quian Quiroga, Jeremy Freeman, and Spencer L Smith. Improving data quality in neuronal population recordings. *Nature neuroscience*, 19(9):1165–1174, 2016.

[11] Marieke L Schölvinck, David A Leopold, Matthew J Brookes, and Patrick H Khader. The contribution of electrophysiology to functional connectivity mapping. *Neuroimage*, 80:297–306, 2013.

[12] Ruben Sanchez-Romero and Michael W Cole. Combining multiple functional connectivity methods to improve causal inferences. *Journal of Cognitive Neuroscience*, pages 1–15, 2019.

[13] Xichao Mi, Ning Cheng, and Tao Zhang. Performance comparison between gpdc and pcmi for measuring directionality of neural information flow. *Journal of Neuroscience Methods*, 227:57–64, 2014.

[14] Lionel Barnett and Anil K Seth. The mvgc multivariate granger causality toolbox: a new approach to granger-causal inference. *Journal of neuroscience methods*, 223:50–68, 2014.

[15] Luiz A Baccalá and Koichi Sameshima. Partial directed coherence: a new concept in neural structure determination. *Biological cybernetics*, 84(6):463–474, 2001.

[16] Leonardo Novelli, Patricia Wollstadt, Pedro Mediano, Michael Wibral, and Joseph T Lizier. Large-scale directed network inference with multivariate transfer entropy and hierarchical statistical testing. *Network Neuroscience*, 3(3):827–847, 2019.

[17] Răzvan Gămănuţ, Henry Kennedy, Zoltán Toroczkai, Mária Ercsey-Ravasz, David C Van Essen, Kenneth Knoblauch, and Andreas Burkhalter. The mouse cortical connectome, characterized by an ultra-dense cortical graph, maintains specificity by distinct connectivity profiles. *Neuron*, 97(3):698–715, 2018.

[18] Yoram Gutfreund, I Segev, et al. Subthreshold oscillations and resonant frequency in guinea-pig cortical neurons: physiology and modelling. *The Journal of physiology*, 483(3):621–640, 1995.

[19] Belén Sancristóbal, Raul Vicente, and Jordi Garcia-Ojalvo. Role of frequency mismatch in neuronal communication through coherence. *Journal of computational neuroscience*, 37(2):193–208, 2014.

[20] Alessandro Barardi, Belen Sancristóbal, and Jordi Garcia-Ojalvo. Phase-coherence transitions and communication in the gamma range between delay-coupled neuronal populations. *PLoS computational biology*, 10(7):e1003723, 2014.

[21] Belén De Sancristóbal, Raul Vicente, Jose Maria Sancho, and Jordi Garcia-Ojalvo. Emergent bimodal firing patterns implement different encoding strategies during gamma-band oscillations. *Frontiers in computational neuroscience*, 7:18, 2013.

[22] Petar Tomov, Rodrigo FO Pena, Michael A Zaks, and Antonio C Roque. Sustained oscillations, irregular firing, and chaotic dynamics in hierarchical modular networks with mixtures of electrophysiological cell types. *Frontiers in computational neuroscience*, 8:103, 2014.

[23] Nikola T Markov, MM Ercsey-Ravasz, AR Ribeiro Gomes, Camille Lamy, Loic Magrou, Julien Vezoli, P Misery, A Falchier, R Quilodran, MA Gariel, et al. A weighted and directed interareal connectivity matrix for macaque cerebral cortex. *Cerebral cortex*, 24(1):17–36, 2014.

[24] Madhura R Joglekar, Jorge F Mejias, Guangyu Robert Yang, and Xiao-Jing Wang. Inter-areal balanced amplification enhances signal propagation in a large-scale circuit model of the primate cortex. *Neuron*, 98(1):222–234, 2018.

[25] Jorge F Mejias, John D Murray, Henry Kennedy, and Xiao-Jing Wang. Feedforward and feedback frequency-dependent interactions in a large-scale laminar network of the primate cortex. *Science advances*, 2(11):e1601335, 2016.

[26] Hannah Choi and Stefan Mihalas. Synchronization dependent on spatial structures of a mesoscopic whole-brain network. *PLoS computational biology*, 15(4):e1006978, 2019.

[27] Alberto Mazzoni, Henrik Linden, Hermann Cuntz, Anders Lansner, Stefano Panzeri, and Gaute T Einevoll. Computing the local field potential (lfp) from integrate-and-fire network models. *PLoS computational biology*, 11(12), 2015.

[28] Alberto Mazzoni, Stefano Panzeri, Nikos K Logothetis, and Nicolas Brunel. Encoding of naturalistic stimuli by local field potential spectra in networks of excitatory and inhibitory neurons. *PLoS Comput Biol*, 4(12):e1000239, 2008.

[29] Luiz A Baccala, K Sameshima, and DY Takahashi. Generalized partial directed coherence. In *2007 15th International conference on digital signal processing*, pages 163–166. IEEE, 2007.

[30] Koichi Sameshima and Luiz Antonio Baccala. *Methods in brain connectivity inference through multivariate time series analysis*. CRC press, 2014.

[31] JD Hamilton. Times series analysis, one of princeton university press's notable centenary titles. *Hard over*, 1994.

[32] Michael W Cole, Takuya Ito, Danielle S Bassett, and Douglas H Schultz. Activity flow over resting-state networks shapes cognitive task activations. *Nature neuroscience*, 19(12):1718–1726, 2016.

[33] Marcel Stimberg, Romain Brette, and Dan FM Goodman. Brian 2, an intuitive and efficient neural simulator. *Elife*, 8:e47314, 2019.

[34] James M Bower and David Beeman. *The book of GENESIS: exploring realistic neural models with the GEneral NEural SImulation System*. Springer Science & Business Media, 2012.

[35] Sara Sommariva, Alberto Sorrentino, Michele Piana, Vittorio Pizzella, and Laura Marzetti. A comparative study of the robustness of frequency-domain connectivity measures to finite data length. *Brain topography*, 32(4):675–695, 2019.

[36] Agostina Palmigiano, Theo Geisel, Fred Wolf, and Demian Battaglia. Flexible information routing by transient synchrony. *Nature neuroscience*, 20(7):1014, 2017.

[37] Peter Uhlhaas, Gordon Pipa, Bruss Lima, Lucia Melloni, Sergio Neuenschwander, Danko Nikolić, and Wolf Singer. Neural synchrony in cortical networks: history, concept and current status. *Frontiers in integrative neuroscience*, 3:17, 2009.

[38] György Buzsáki and Xiao-Jing Wang. Mechanisms of gamma oscillations. *Annual review of neuroscience*, 35:203–225, 2012.

[39] Panagiota Theodoni, Piotr Majka, David H Reser, Daniel K Wójcik, Marcello GP Rosa, and Xiao-Jing Wang. Structural attributes and principles of the neocortical connectome in the marmoset monkey. *bioRxiv*, 2020.

[40] György Buzsáki and Kenji Mizuseki. The log-dynamic brain: how skewed distributions affect network operations. *Nature Reviews Neuroscience*, 15(4):264–278, 2014.

[41] Patric Hagmann, Leila Cammoun, Xavier Gigandet, Reto Meuli, Christopher J Honey, Van J Wedeen, and Olaf Sporns. Mapping the structural core of human cerebral cortex. *PLoS Biol*, 6(7):e159, 2008.

[42] Maxwell H Turner, Kevin Mann, and Thomas R Clandinin. The connectome predicts resting state functional connectivity across the drosophila brain. *bioRxiv*, 2020.

[43] Rishidev Chaudhuri, Kenneth Knoblauch, Marie-Alice Gariel, Henry Kennedy, and Xiao-Jing Wang. A large-scale circuit mechanism for hierarchical dynamical processing in the primate cortex. *Neuron*, 88(2):419–431, 2015.

[44] Bertha Vézquez-Rodríguez, Zhen-Qi Liu, Patric Hagmann, and Bratislav Misic. Signal propagation via cortical hierarchies. *Network Neuroscience*, 4(4):1072–1090, 2020.

[45] Milou Straathof, Michel RT Sinke, Rick M Dijkhuizen, and Willem M Otte. A systematic review on the quantitative relationship between structural and functional network connectivity strength in mammalian brains. *Journal of Cerebral Blood Flow & Metabolism*, 39(2):189–209, 2019.

[46] James M Stafford, Benjamin R Jarrett, Oscar Miranda-Dominguez, Brian D Mills, Nicholas Cain, Stefan Mihalas, Garet P Lahvis, K Matthew Lattal, Suzanne H Mitchell, Stephen V David, et al. Large-scale topology and the default mode network in the mouse connectome. *Proceedings of the National Academy of Sciences*, 111(52):18745–18750, 2014.

[47] Joanes Grandjean, Valerio Zerbi, Joshua Henk Balsters, Nicole Wenderoth, and Markus Rudin. Structural basis of large-scale functional connectivity in the mouse. *Journal of Neuroscience*, 37(34):8092–8101, 2017.

[48] Kanika Bansal, Johan Nakuci, and Sarah Feldt Muldoon. Personalized brain network models for assessing structure–function relationships. *Current Opinion in Neurobiology*, 52:42–47, 2018.

[49] Leonardo Novelli and Joseph T Lizier. Inferring network properties from time series using transfer entropy and mutual information: validation of multivariate versus bivariate approaches. *Network Neuroscience*, (Just Accepted):1–52, 2020.

[50] Christopher J Honey, Olaf Sporns, Leila Cammoun, Xavier Gigandet, Jean-Philippe Thiran, Reto Meuli, and Patric Hagmann. Predicting human resting-state functional connectivity from structural connectivity. *Proceedings of the National Academy of Sciences*, 106(6):2035–2040, 2009.

[51] Malte Bieler, Kay Sieben, Nicole Cichon, Sandra Schildt, Brigitte Röder, and Ileana L Hanganu-Opatz. Rate and temporal coding convey multisensory information in primary sensory cortices. *eneuro*, 4(2), 2017.

[52] Anil K Seth. Neural coding: rate and time codes work together. *Current Biology*, 25(3):R110–R113, 2015.

[53] Gerald Hahn, Adrian Ponce-Alvarez, Gustavo Deco, Ad Aertsen, and Arvind Kumar. Portraits of communication in neuronal networks. *Nature Reviews Neuroscience*, 20(2):117–127, 2019.

[54] Arvind Kumar, Stefan Rotter, and Ad Aertsen. Spiking activity propagation in neuronal networks: reconciling different perspectives on neural coding. *Nature reviews neuroscience*, 11(9):615–627, 2010.

[55] Artur Luczak, Bruce L McNaughton, and Kenneth D Harris. Packet-based communication in the cortex. *Nature Reviews Neuroscience*, 16(12):745–755, 2015.

[56] Gerald Hahn, Alejandro F Bujan, Yves Frégnac, Ad Aertsen, and Arvind Kumar. Communication through resonance in spiking neuronal networks. *PLoS computational biology*, 10(8):e1003811, 2014.

[57] Paul HE Tiesinga and Terrence J Sejnowski. Mechanisms for phase shifting in cortical networks and their role in communication through coherence. *Frontiers in human neuroscience*, 4:196, 2010.

[58] Mária Ercsey-Ravasz, Nikola T Markov, Camille Lamy, David C Van Essen, Kenneth Knoblauch, Zoltán Toroczkai, and Henry Kennedy. A predictive network model of cerebral cortical connectivity based on a distance rule. *Neuron*, 80(1):184–197, 2013.

[59] Tomokazu Tsurugizawa, Boucif Djemai, and Andrew Zalesky. The impact of fasting on resting state brain networks in mice. *Scientific reports*, 9(1):1–12, 2019.

[60] Maxwell A Sherman, Shane Lee, Robert Law, Saskia Haegens, Catherine A Thorn, Matti S Hämäläinen, Christopher I Moore, and Stephanie R Jones. Neural mechanisms of transient neocortical beta rhythms: Converging evidence from humans, computational modeling, monkeys, and mice. *Proceedings of the National Academy of Sciences*, 113(33):E4885–E4894, 2016.

[61] Julien Vezoli, Martin Vinck, Conrado Arturo Bosman, André Moraes Bastos, Christopher Murphy Lewis, Henry Kennedy, and Pascal Fries. The role of anatomical connection strength for interareal communication in macaque cortex. *Available at SSRN 3751057*, 2020.

[62] Iain Stitt, Karl J Hollensteiner, Edgar Galindo-Leon, Florian Pieper, Eva Fiedler, Thomas Stieglitz, Gerhard Engler, Guido Nolte, and Andreas K Engel. Dynamic reconfiguration of cortical functional connectivity across brain states. *Scientific reports*, 7(1):1–14, 2017.

[63] Mathieu Golos, Viktor Jirsa, and Emmanuel Daucé. Multistability in large scale models of brain activity. *PLoS computational biology*, 11(12):e1004644, 2015.

[64] Patricio Orio, Marilyn Gatica, Rubén Herzog, Jean Paul Maidana, Samy Castro, and Kesheng Xu. Chaos versus noise as drivers of multistability in neural networks. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 28(10):106321, 2018.

[65] Gustavo Deco and Viktor K Jirsa. Ongoing cortical activity at rest: criticality, multistability, and ghost attractors. *Journal of Neuroscience*, 32(10):3366–3375, 2012.

[66] Csaba Erö, Marc-Oliver Gewaltig, Daniel Keller, and Henry Markram. A cell atlas for the mouse brain. *Frontiers in neuroinformatics*, 12:84, 2018.

[67] Joseph E Knox, Kameron Decker Harris, Nile Graddis, Jennifer D Whitesell, Hongkui Zeng, Julie A Harris, Eric Shea-Brown, and Stefan Mihalas. High-resolution data-driven model of the mouse connectome. *Network Neuroscience*, 3(1):217–236, 2018.

Supplementary Material

Figure S1: **Area Under Curve (AUC) and mean of GPDC estimates** . (A) GPDC AUC *versus* FLN. (B) GPDC mean *versus* FLN. In both cases, the Pearson correlation between GPDC mean or AUC and FLN is 0.66. Red lines are linear fit.

Figure S2: **Correlation and difference between GPDC etimates conditioned to the whole con-
nectome and GPDC estimates conditioned to the cluster areas**. A)Pearson correlation increases
with the cluster size. Each bullet represents the average pearson correlation over 150 clusters
composed of randomly chosen areas. Comparing all clusters size, the highest value for standard
deviation of pearson correlation 0.07. B) Average difference between GPDC estimates conditoned
to the whole connnectome and GPDC estimates conditioned to the cluster areas. Average and
standard deviation decrease with the cluster size.

Table S1: **Name of areas in mouse cortical connectome.** Adapted from Gămănut et. al., 2018.

Abbreviations	Areas
ACAd	Anterior cingulate area dorsal part
AL	Anterolateral area
AM	Anteromedial area
AUDpo	Auditory cortex posterior area
DP	Dorsal posterior area
GU	Gustatory area
LM	Lateromedial area
MM	Mediomedial area
MOp	Motor cortex primary
P	Posterior area
PL	Prelimbic area
PM	Posteromedial area
RL	Rostrolateral area
RSPd	Rostroplenial area dorsal part
SSp-bfd	Somatosensory cortex primary barrel field
SSp-un	Somatosensory cortex primary unassigned
SSs	Somatosensory cortex secondary
V1	Primary visual area
VISC	Visceral area