# Adaptations and responses of the common dandelion to low

# atmospheric pressure in high altitude environments

Carla C. M. Arce[1]* (carla.marques@unine.ch), Zoe Bont[2]* (zoe.bont@ips.unibe.ch), Ricardo

A. R. Machado[3] (ricardo.machado@unine.ch), Paulo F. Cristaldo[4] (pfellipec@gmail.com),

Matthias Erb[2] (matthias.erb@ips.unibe.ch)

*[1]Laboratory of Fundamental and Applied Research in Chemical Ecology, University of*

*Neuchâtel, rue Emile-Argand 11, 2000 Neuchâtel, Switzerland*

*[2]Institute of Plant Sciences, University of Bern, Altenbergrain 21, 3013 Bern, Switzerland*

*[3]Experimental Biology Research Group, Institute of Biology, University of Neuchâtel, rue*

*Emile-Argand 11, 2000 Neuchâtel, Switzerland*

*[4]Agronomy Department, Rural Federal University of Pernambuco, 52171900 – Recife, Brazil*

* shared first authorship

**Correspondence:** Matthias Erb (matthias.erb@ips.unibe.ch), University of Bern, Institute of

Plant Sciences, Altenbergrain 21, 3013 Bern, Switzerland, tel: +41 31 631 86 68

**Abstract**

Atmospheric pressure is an important, yet understudied factor that may shape plant ecology and evolution. We evaluated the impact of ecologically realistic atmospheric pressures between 660 and 950 hPa on the growth and secondary metabolism of dandelion populations originating from different altitudes. We find that growth under low atmospheric pressure suppresses root growth and the production of defensive sesquiterpene lactones. The suppression of defence at low atmospheric pressure is most pronounced for populations originating from lower altitudes. Populations from higher altitudes constitutively produce less latex and sesquiterpene lactones, and do not suffer from suppression under low atmospheric pressure. We conclude that atmospheric pressure modulates plant growth and defence traits, and that evolutionary history shapes plant phenotypic plasticity in response to atmospheric pressure. Our findings have important implications for our understanding of altitudinal gradients and the future use of plants as a source of food and bioactive metabolites in extraterrestrial habitats.

**Introduction**

Plants successfully colonize a wide range of terrestrial habitats. Their capacity to survive under variable and challenging environmental conditions is a prerequisite for their contribution to biodiversity and sustainable agriculture in the face of climate change, and their potential use as a source of food and medicine in future extraterrestrial habitats (Jump & Peñuelas 2005; Wheeler 2017). Atmospheric pressure plays a potentially important role in terrestrial and extraterrestrial environments, albeit for different reasons. On earth, climate change research often relies on altitudinal gradients to understand how plants and terrestrial ecosystems will be affected by future climates. Such studies typically assume that differences in climate at different altitudes account for changes in plant performance, and that effects of atmospheric pressure are negligible. In contrast, space plant biology has long been interested in understanding effects of atmospheric pressure on plants, as it would be more practical to maintain low pressure greenhouses in future moon or mars colonies (Corey *et al.* 1997; Iwabuchi & Kurata 2003; Richards *et al.* 2006). Thus, understanding the impact of atmospheric pressure on plants and the capacity of plants to withstand and adapt to different atmospheric pressures is of substantial interest for the present and future of humanity. Yet, to date, our understanding of how atmospheric pressure influences plants is not well developed.

How are plants affected by low atmospheric pressure? One consequence of low atmospheric pressure at high altitudes is the decrease of partial pressures of $O_2$ and $CO_2$, which are essential substrates for respiration and photosynthesis (Zabalza *et al.* 2009; Xu *et al.* 2015). Changes in $O_2$ and $CO_2$ partial pressures have been linked to changes in plant physiology and growth (Paul *et al.* 2004; He *et al.* 2007; Kammer *et al.* 2015; Zhou *et al.* 2017). Effects are complex, since the decrease of the partial pressure of the atmospheric gases is accompanied by an increase in diffusion rates, which may compensate for the low ambient concentration of the essential gases (Terashima *et al.* 1995). Further, as the diffusion coefficient for water vapor is increased, transpiration increases, (Smith & Geller 1979), which can impose water stress on plants growing under reduced atmospheric pressure (Iwabuchi & Kurata 2003; Paul *et al.* 2004; Richards *et al.* 2006).

Can plants exhibit phenotypic plasticity to cope with low atmospheric pressure? Several studies suggest that plants respond dynamically to reduced atmospheric pressure

(Spanarkel & Drew 2002; He *et al.* 2003; Iwabuchi & Kurata 2003; Richards *et al.* 2006). In the

mountain plant *Arabis alpina*, for instance, stomatal density increases and stomata aperture

narrows at low atmospheric pressure (Kammer *et al.* 2015). This induced response likely

benefits the plant, because at low partial pressures of atmospheric gases, a higher stomatal

density may ensure optimal supply of $CO_2$ for photosynthesis (Woodward & Bazzaz 1988; Xu

*et al.* 2016), while a narrow aperture of the stomata can restrict water loss to counteract

increased transpiration rate (Buckley 2005). At a molecular level, it is assumed that low

atmospheric pressure represents an environmental stress to which plants must respond with

changes in their metabolic pathways in order to survive successfully (Ferl *et al.* 2002; Paul &

Ferl 2006). Recent research has documented extensive changes in gene expression patterns

in *Arabidopsis thaliana* when exposed to a low atmospheric pressure environment, including

genes associated with hypoxia and water loss (Paul *et al.* 2004; Zhou *et al.* 2017).

Can plants adapt to low atmospheric pressure over evolutionary time? Vascular plants

colonize habitats between 0 and 6150 m above sea corresponding to atmospheric pressures

between 101 and 46 kPa (Angel *et al.* 2016). Since atmospheric pressure influences plant

performance, and many plant species occur along altitudinal gradients, local adaptation to

atmospheric pressure can be expected (Kammer *et al.* 2015; Ward & Strain 1997; Ward *et al.*

2000). Using growth chambers simulating high and low altitude pressure conditions, Kammer

*et al*. (2015) found that *Arabis alpina* adjusts stomatal density in response to low atmospheric

pressure, while the low altitude plant *Arabidopsis thaliana* does not. Evidence for genetic

differences in the response of stomatal density to pressure conditions was also found within

species. Woodward & Bazzaz (1988) showed that in the grass *Nardus stricta*, plants from

higher altitudes developed greater declines in stomatal density at experimentally increased $CO_2$

partial pressure than plants from lower altitudes. Apart from stomatal development, very little is

known about evolutionary adaptations of plants to low atmospheric pressure.

Plant secondary or specialized metabolites play important roles in plant responses and

adaptations to diverse environments and stress factors (Hartmann 2007; Moore *et al.* 2014),

including herbivores and pathogens (Ehrlich & Raven 1964; Kessler & Baldwin 2001; Bednarek

& Osbourn 2009; Moles *et al.* 2013), abiotic stress (Ramakrishna & Ravishankar 2011; Arbona

*et al.* 2013; Nakabayashi & Saito 2015), mutualists (Pichersky & Gershenzon 2002; Schliemann

*et al.* 2008; Stevenson *et al.* 2017) and other plants (Baldwin *et al.* 2006; Semchenko *et al.*

2014). The production, transport and storage of specialized metabolites is assumed to be costly

(Neilson *et al.* 2013), and plants therefore constantly fine-tune their chemical arsenal to the

demands of their environment and internal condition through phenotypic plasticity (Metlen *et al.*

2009). Over evolutionary times, environmental conditions may act as selective forces on plant

genotype selection and can shape genetically determined chemical profiles of plants

(Cunningham *et al.* 1999; Agrawal *et al.* 2012; Züst *et al.* 2012; Kessler & Kalske 2018). With

increasing altitude, mountain habitats impose different environmental demands on plants,

including harsher abiotic conditions and a lower intensity of biotic interactions (Rasmann *et al.*

2014; Buckley *et al.* 2019; Midolo & Wellstein 2020). Several studies found evidence that the

genetic variation of plant secondary metabolites is shaped by these environmental gradients

(Bernal *et al.* 2013; Moreira *et al.* 2018; Bakhtiari *et al.* 2019; Buckley *et al.* 2019; Bont *et al.*

2020). However, the role of decreasing atmospheric pressure at increasing altitudes in shaping

the evolution and expression of plant secondary metabolites is poorly understood. Levine *et al.*

(2008) found that the glucosinolate content in the roots of radish was decreased under

hypobaric, normoxic conditions. In lettuce, the concentration of phenolics, anthocyanins and

carotenoids in the leaves was increased under hypobaric and hypoxic conditions (He *et al.*

2013). However, all these experiments were conducted in low pressure chambers under

conditions far beyond the natural range of plants. How a plant's evolutionary history shapes its

response to ecologically realistic low-pressure environments is unknown.

Along environmental gradients, closely related asexual and sexual plant taxa often

have different distribution patterns – a phenomenon called geographic parthenogenesis

(Glesener & Tilman 1978). It is generally predicted that apomictic plants with asexual

reproduction tend to have larger distribution ranges including higher latitudes and higher

altitudes, and thus lower atmospheric pressure, than their sexual relatives (Bierzychudek 1985;

Kearney 2005; Cosendai *et al.* 2013). Hypotheses to explain these environmental distribution

patterns include different colonization abilities of sexual and asexual organisms as well as

different capacities to co-evolve with other organisms (e.g. reviewed in Hörandl 2006 and in

Tilquin & Kokko 2016). Although many studies empirically support the predicted geographical

parthenogenesis patterns, some taxa with asexual and sexual organisms show opposite or

mixed trends in geographic distribution. For the common dandelion *Taraxacum officinale* agg.

(Asteraceae), a species complex that largely consists of diploid sexuals and triploid asexuals

(Verduijn *et al.* 2004), larger latitudinal distribution of triploids has been observed, with triploids

colonizing more extreme environments in the north of Europe (Menken *et al.* 1995; Van Dijk *et*

*al.* 2003; Verhoeven & Biere 2013). Surprisingly however, triploids are less frequent at higher

altitudes than diploids, at least along certain transects (Calame & Felber 2000; Bont *et al.* 2020).

The reason for this pattern is currently unclear. One possibility is that high altitude environments

impose selection pressures that are different from high latitude environments. Therefore, one

hypothesis to explain the lower success of triploids at high altitude is that they are constrained

by low resistance to low atmospheric pressure.

*Taraxacum officinale* possesses a reservoir of secondary metabolites stored in

specialized cells, so called laticifers, throughout almost all organs. The latex of *T. officinale* is

most abundant in the taproot. The latex is characterized by three major classes of secondary

metabolites: hydroxyphenylacetate inositol esters with either two or three side chains (di-PIEs

and tri-PIEs), the sesquiterpene lactone taraxinic acid β-d-glucopyranosyl ester (TA-G), and

triterpene acetates (TritAc) (Huber *et al.* 2015). Latex is mainly connotated with defensive

functions against herbivores and pathogens (Konno 2011), and our previous work on the

bioactivity and ecological role of the latex metabolites of *T. officinale* confirms this hypothesis

(Huber *et al.* 2016b, a; Bont *et al.* 2017). TA-G in particular reduces the attractiveness of *T.*

*officinale* to white grubs and thereby increases plant performance (Huber *et al.* 2016b).

Interestingly, in our recent study on the latex metabolites of 63 natural *T. officinale* populations

across Switzerland we found a strong association of the latex metabolites TA-G and di-PIEs

with the climatic history of the natural populations, which may be suggestive of a role of latex

secondary metabolites in abiotic stress tolerance (Bont *et al.* 2020). Inside the lacticifers, latex

is maintained at positive pressure, which allows it to be expelled after tissue disruption. Latex

likely responds to turgor pressure (Agrawal & Konno 2009), which in turn is determined by the

water balance of the plant. At low atmospheric pressure, evaporation is increased, which could

therefore have a direct or indirect effect on latex quality or quantity.

Here, we took advantage of experimental stations in Switzerland between 526 and

3450 m with standardized abiotic conditions to study the effects of atmospheric pressure on the

growth and latex composition of *T. officinale*. We included offspring from nine natural

populations from Switzerland from different altitudes, including populations containing both

diploid and triploid cytotypes. This setup allowed us to test (i) whether low atmospheric pressure

affects plant growth and latex composition through environmental plasticity; (ii) whether there

are signatures of local adaptation to atmospheric pressure, with populations from higher

altitudes performing better at low atmospheric pressure; and (iii) whether low atmospheric

pressure has a stronger impact on triploid than diploid plants, thus potentially restricting the

expansion of triploids towards higher altitudes.

**Material and methods**

*Study species and seed collection*

**Figure 1** Geographical positions of the origins of the *T. officinale* populations (blue squares 1-9) and of the experimental stations (orange circles). (a) Spatial distribution of the sites across Switzerland. © Photographs: Bern, CC0 public domain; Adlemsried, Zoe Bont; Jungfraujoch, Julius Silver, CC BY-SA 4.0; Kleine Scheidegg, Grindel1, CC BY-SA 3.0. (b) Altitude and atmospheric pressure of the sites and representation of the linear relationship between these two parameters. (c) Visualization of plant growth conditions at experimental stations, including controlled abiotic parameters (grey bars) and varying atmospheric pressure (orange bars).

The common dandelion is a latex-producing species complex with a worldwide,

cosmopolitan distribution (Stewart-Wade *et al.* 2002). The wind-dispersed perennial herb can

tolerate a broad range of environmental conditions and can be found from sea level to altitudes

of up to 4000 m.a.s.l. (Molina-Montenegro *et al.* 2013; Sandoya *et al.* 2017). In Switzerland, the

plant occurs at altitudes of up to 2000 m.a.s.l. (Calame & Felber 2000). In this study, plants

from nine natural populations of Switzerland were included. The nine populations are a subset

of 63 populations that were characterized in our previous work (Bont *et al.* 2020). They cover

an altitudinal range from 302 to 1607 m.a.s.l (Fig. 1ab, Table S1 in Supporting Information).

Each population was located within a maximal distance of 1 km from a meteorological

monitoring station of MeteoSwiss, the Swiss Federal Office for Meteorology and Climatology,

which enabled us to obtain long-term data on climatic conditions. According to Bont *et al.*

(2020), 10 variables were selected from the MeteoSwiss database, representing average air

pressure, temperature, precipitation and light conditions of the populations for the years 1996

– 2015. For most variables, data was available for all 9 populations (Fig. S1c). Seeds from the

natural populations were collected and F2 plants were generated for each population as

described in Bont *et al.* 2020. The cytotype distribution of each population was determined by

analysing the ploidy level of the plants with flow cytometry (Bont *et al.* 2020). The nine F2

populations consisted of populations with only diploid plants (5 populations) and populations

with both triploid and diploid plants (4 populations) (Table S1).

*Experimental design*

In order to investigate the effect of atmospheric pressure on the growth and chemical

defence of *T. officinale*, we cultivated the F2 plants from the nine natural populations in four

experimental stations at 526 m.a.s.l. (Bern), 1046 m.a.s.l. (Adlemsried), 2061 m.a.s.l. (Kleine

Scheidegg) and 3450 m.a.s.l. (Jungfraujoch) in Switzerland (Fig. 1). Although *T. officinale* does

not occur naturally in Switzerland at 3450 m.a.s.l. due to the cold climate, the plant is frequently

reported at this altitude in the South American Andes (Sandoya *et al.* 2017) and is therefore

not restricted in its occurrence by the associated low atmospheric pressure. In the experimental

stations, the plants were grown inside under controlled light supply, temperature and relative

humidity and ambient indoor air quality. This allowed us to specifically test the influence of the

atmospheric pressure gradient, ranging from 950 hPa to 660 hPa (Fig. 1c).

For experiments, seeds from 6-8 mother plants (F1) per population were germinated

on moist seedling substrate and transplanted into individual 1l pots filled with potting soil (5

parts field soil, 4 parts peat, 1 part sand) after 18 ± 2 days. All plants were grown under

controlled conditions (25 ± 2 °C, 60 ± 5% RH, 16:8 light:dark cycle) in Bern. 24 h after

transplantation, the plants were distributed among the four experimental stations, so that 6-8

plants from each of the nine populations were cultivated in each station. The temperature and

relative humidity were measured hourly at all stations with RHT10-Data Logger (Extech

Instruments, China). To standardize light supply, we grew the plants in rooms with reduced

natural light and supplied them with LED lights (400 LED beads, LED type SMD 5730, 400 ±

20 µmol*m$^2$*s$^1$) placed one meter above the plants. The plants were watered three times a

week and fertilized once a week.

*Assessment of plant traits and chemical analysis*

After 45 days of growth in the experimental stations, we transported the plants back to

Bern to analyse growth and defence traits. To assess vegetative growth, we quantified the plant

biomass by measuring the dry weight of roots and shoots separately and then calculated

root:shoot ratios. To study defence traits, we measured the amount of taproot latex and

quantified the concentration of the latex secondary metabolites TA-G, di-PIEs and tri-PIEs.

To quantify latex traits, plants were cut 0.5 cm below the tiller and the taproot latex that

was released over 20 seconds was collected and weighted. Two µl of latex was immediately

transferred into 200 µl methanol for chemical analysis. The roots were carefully washed with

tap water and roots and shoots were placed in a drying oven at 50°C until constant dry mass

was reached. The chemical analysis of the latex metabolites was carried out as described in

Bont *et al.*, 2017. Briefly, the samples were vortexed for 10 min, ultrasonicated for 10 min,

centrifuged at 4 °C and 14000 rpm for 20 min and supernatants were used for further analysis.

Relative concentrations of TA-G, di-PIEs and tri-PIEs were determined by injecting the latex

extracts into an Acquity UPLC-PDA-MS (Waters, Milford MA, USA) with electrospray ionization

in positive mode, consisting of an ultra-performance liquid chromatograph (UPLC) coupled to

a photodiode array detector (PDA) and a single quadrupole mass detector (QDa). For

quantification, peak areas were integrated at 245 nm for TA-G and at 275 nm for di- and tri-

PIES, while concurrently recorded characteristic mass features were used to confirm

compound identities (Bont *et al.* 2017). For absolute quantification of TA-G, we established an

external standard curve with loganin (CAS: 18524-94-2, Sigma-Aldrich Chemie GmbH, Buchs,

Switzerland) and calculated the corresponding response factor to pure TA-G. (Bont et al. 2017).

For di-and tri-PIEs, relative concentrations were calculated separately.

*Statistical analysis*

All statistical analyses were performed in R 4.0.2 (R Core Team 2017). To represent

the climatic conditions associated with the different altitudes of the populations we first

conducted a principal component analysis (PCA), as some of the meteorological variables of

the population origins were highly correlated (Fig. S1c). This approach is widely used to analyze

the impact of climate on the evolution of plant traits (Keller *et al.* 2009; Kooyers *et al.* 2015;

Villaverde *et al.* 2017). We applied the function 'prcomp' on scaled variables to reduce

dimensionality of the data and selected the axis that explained most of the cumulative variance

and represented the climatic conditions associated with the different altitudes at which the

populations evolved (climPCA1).

For data exploration, we calculated a correlation matrix of all parameters included in

the experiment. Then, we analysed all data with linear mixed-effects models (LMEMs) (Bates

2020). The models were fit using the function 'lmer' from the package 'lme4' (Bates *et al.* 2015)

with restricted maximum likelihood estimation (REML). Variables representing fixed effects

were scaled and centred prior to computation to reduce nonessential multicollinearity

(Iacobucci *et al.* 2016). If necessary, log-transformation was applied to the response variable

to improve distribution of variance. Model assumptions were validated using 'plotresid' from the

package 'RVAideMemoire' (Hervé 2018). The significances of the fixed effects were estimated

using the package 'lmerTest' (Kuznetsova *et al.* 2017) by calculating type II analysis of variance

tables with Kenward-Rodger's approximation to degrees of freedom (Halekoh & Højsgaard

2014).

To test the overall effect of the varying atmospheric pressure of the experimental

stations on plant growth and defence, we performed LMEMs separately for each plant trait (root

dry weight, shoot dry weight, root:shoot ratio, taproot latex, TA-G, di-PIEs, tri-PIEs). We used

the mean value per trait per population per station as response variable, tested the fixed effect

of the atmospheric pressure of the experimental station ('$P_{Station}$') and included '(1|Population)'

as random effect and grouping factor to allow for varying intercepts between populations. We

performed a similar analysis to assess the overall effect of the climatic conditions associated

with the different altitudes of the population origins on plant growth and defence by testing the

fixed effect of 'climPCA1' and including '(1|Station)' as random effect and grouping factor to

allow for varying intercepts between experimental stations.

In order to further investigate interacting effects of the climatic conditions associated

with different altitudes of the population origins (genetic effects) and the atmospheric pressure

of the experimental stations (environmental effects) on the plants, we next conducted a full

model analysis for each plant trait, including the abiotic environment of the stations as covariate.

Since temperature and relative humidity were highly correlated (Pearson's $r$ = -0.95, Fig. S3),

we used only temperature as covariate in the model. Trait values of individual plants were used

as response variables, with '(1|Population)' as random effect to allow the calculation of separate

intercepts for each population. Concurrently, the impact of ploidy was tested by adding the

ploidy level (diploid or triploid) of each plant as fixed effect. The full model syntax was the

following: plant trait ~ climPCA1 x $P_{Station}$ + $P_{Station}$ x temperature + $P_{Station}$ x Ploidy +

(1|Population). As expected, temperature influenced all experimental parameters (LMEMs:

Temp, $P$ < 0.05, Table 1), but did not interact with atmospheric pressure (LMEMs: $P_{Station}$ x

Temp, $P$ < 0.05, Table 1). A confounding effect of residual variation in temperature on the

detected effects of atmospheric pressure is, therefore, unlikely. To visualize the  significant

interactive effect of 'climPCA1 x $P_{Station}$' on TA-G and on di-PIEs concentration, we then used

the package 'effects' (Fox $et~al.$ 2019) for model prediction with unscaled and uncentered fixed

effects, excluding the non-significant interaction of '$P_{Station}$ x temperature' from the model.

Likewise, we predicted and visualized the nonsignificant effects of $P_{Station}$ x Ploidy on all

measured plant traits. All results were visualized using 'ggplot2' (Wickham 2016) and 'effects'

(Fox $et~al.$ 2019).

**Results**

*Validation of experimental setup*

We first tested whether our efforts to standardize environmental parameters across experimental sites were successful. We found that indoor temperature and relative humidity varied slightly between sites (Fig. 1c), but neither variable was correlated to atmospheric pressure (Fig. S3). As plants were also grown using identical potting soil, watering, lighting and comparable indoor environments with respect to air pollution, we estimate that major abiotic parameters were successfully standardized to allow for an assessment of the role of atmospheric pressure on *T. officinale*. By growing *T. officinale* plants from seeds of diploid and triploid populations originating from different altitudes, we tested (i) the effect of atmospheric pressure on growth and defence; (ii) the impact of population origin on plant responses to atmospheric pressure (as an indicator for local adaptation to atmospheric pressure); and (iii) the importance of a plant's cytotype (diploid vs. triploid) on its capacity to grow under different atmospheric pressure, as a test of the hypothesis that low atmospheric pressure restricts the establishment of triploids at high altitudes. To this end, we first describe environmental (Fig. 2) and heritable contributions (Fig. 3) to variation in defence and growth independently. Using a full model, we then address adaptive phenotypic plasticity (Table 1, Fig. 4) and differences between cytotypes (Table 1, Fig. 5).

*Atmospheric pressure modulates root growth and defences*

Across all populations and irrespective of population origin and cytotype, *T. officinale* plants showed distinct phenotypes as a function of atmospheric pressure at the different experimental sites (Fig. 2). Both root growth and root:shoot ratio were overall lower at higher altitudes with lower atmospheric pressure (LMEM, $P$ = 0.022, Fig. 2 resp. LMEM, $P$ < 0.001, Fig. 2). By contrast, no significant effects on shoot growth and on the amount of exuded taproot latex were observed (LMEMs, $P$ > 0.05, Fig. 2). The concentration of the defence metabolite TA-G in the taproot latex was reduced in plants growing under lower atmospheric pressure (LMEM, $P$ = 0.001, Fig. 2). The concentrations of di-PIEs and tri-PIEs did not vary significantly with atmospheric pressure (LMEMs, $P$ > 0.05, Fig. 2). Thus, growth under low atmospheric pressure is specifically associated with lower root growth and lower accumulation of a defensive sesquiterpene lactone in *T. officinale*.

**Figure 2** Overall effect of atmospheric pressure of experimental station on mean plant performance parameters (root dry weight, shoot dry weight, root:shoot ratio, taproot latex) and chemical composition of latex (TA-G, di-PIEs, tri-PIEs). Blue dots represent average values across all populations per experimental station, while black dots represent average values per population per experimental station ($N$ = 5-8 per population and station). The significance of the effect was tested with linear mixed-effects models (LMEM) and corresponding $P$-values are displayed. Blue lines indicate statistically significant effects ($P$ < 0.05). TA-G: taraxinic acid ß-D-glucopyranosyl ester; di-PIEs: di-4-hydroxyphenylacetate inositol esters; tri-PIEs: tri-4-hydroxyphenylacetate inositol esters.

*Climatic history associated with altitude shapes heritable variation in root defences*

To characterize the climatic conditions under which the *T. officinale* populations evolved, we conducted a principal component analysis (PCA) and then scaled variables to reduce the dimensionality of the data. The first two axes (climPCA1 and climPCA2) together

explained 90% of the cumulative variance in climate parameters (Fig. S1a). 54.3% of the

variance was explained by the first axis (climPCA1) which mainly represents variation in

temperature-related parameters and atmospheric pressure (Fig. S1a, Fig. S1c). ClimPCA1, but

not climPCA2 was highly correlated with the altitude of the population origin (Pearson's $r$ =

0.941, $P$ < 0.001, Fig. S1b). We therefore used climPCA1 to represent the climatic conditions

associated with the different altitudes of the populations.

**Figure 3** Overall effect of climatic conditions associated with different altitudes of population
origins (climPCA1) on mean plant performance parameters (root dry weight, shoot dry weight,
root:shoot ratio, taproot latex) and chemical composition of latex (TA-G, di-PIEs, tri-PIEs).
climPCA1 represents a climatic gradient ranging from high altitude environment to low altitude
environment, which is illustrated by icons. Blue dots represent average values per population
across all experimental stations, while black dots represent average values per population per
experimental station ($N$ = 5-8 per population and station). The significance of the effect was
tested with linear mixed-effects models (LMEM) and corresponding $P$-values are displayed.

Across experimental stations, we found no significant association between climPCA1 and *T. officinale* growth (LMEMs, $P > 0.05$, Fig. 3). However, climPCA1 was correlated to the amount of taproot latex and the concentration of TA-G, with populations originating from lower altitudes producing more latex and more TA-G than populations collected at high altitudes (LMEMs, $P < 0.001$, Fig. 3). No clear effects were observed for di- and tri-PIEs (LMEMs, $P > 0.05$, Fig. 3). Thus, climatic conditions that are correlated with altitude are associated with heritable differences in the production and chemical composition of root latex, with plant originating from higher altitudes producing less latex and defensive sesquiterpene lactones.

*Root defences show patterns of adaptation to high atmospheric pressure*

**Table 1:** Effects of experimental atmospheric pressure ($P_{Station}$), of climatic conditions associated with different altitudes of population origins (climPCA1), of the plant's ploidy level (Ploidy) and of selected interactions on performance and latex profile of *T. officinale* are shown. Temperature of experimental station (Temp) is included as control variable. Results of full mixed-effects model analyses are displayed separately for each performance parameter (root dry weight, shoot dry weight, root:shoot ratio, amount of latex) and for each class of latex secondary metabolites. Significances of fixed effects were assessed by *F*-tests. Estimated *F*-values$_{(NumDF, DenDF)}$ are shown. Levels of statistical significance are indicated with asterisks (***$P$ < 0.001; **$P$ < 0.01; *$P$ < 0.05; (*) $P$ < 0.1). TA-G: taraxinic acid ß-D-glucopyranosyl ester; di-PIEs: di-4-hydroxyphenylacetate inositol esters; tri-PIEs: tri-4-hydroxyphenylacetate inositol esters.

| | Root | Shoot | Root:shoot | Latex | TA-G | di-PIEs | tri-PIEs[1] |
|---|---|---|---|---|---|---|---|
| climPCA1 | $0.83_{(1,6)}$ | $2.08_{(1,6)}$ | $0.00_{(1,6)}$ | $5.20_{(1,6)}$ (*) | $11.00_{(1,6)}$ * | $2.05_{(1,6)}$ | $0.35_{(1,6)}$ |
| $P_{station}$ | $10.75_{(1,209)}$ ** | $1.29_{(1,209)}$ | $18.22_{(1,209)}$ *** | $0.01_{(1,209)}$ | $14.12_{(1,211)}$ *** | $0.01_{(1,211)}$ | $5.60_{(1,211)}$ * |
| Temp | $6.26_{(1,210)}$ * | $4.16_{(1,210)}$ * | $1.62_{(1,209)}$ | $11.41_{(1,210)}$ *** | $4.17_{(1,211)}$ * | $8.03_{(1,211)}$ ** | $8.21_{(1,211)}$ ** |
| Ploidy | $0.06_{(1,108)}$ | $0.92_{(1,107)}$ | $0.38_{(1,143)}$ | $0.06_{(1,114)}$ | $0.23_{(1,101)}$ | $1.36_{(1,95)}$ | $0.38_{(1,209)}$ |
| climPCA1 x $P_{Station}$ | $0.72_{(1,209)}$ | $2.47_{(1,209)}$ | $0.40_{(1,209)}$ | $0.08_{(1,209)}$ | $7.26_{(1,211)}$ ** | $6.49_{(1,211)}$ * | $2.06_{(1,211)}$ |
| $P_{Station}$ x Temp | $0.49_{(1,210)}$ | $0.13_{(1,210)}$ | $1.33_{(1,210)}$ | $3.32_{(1,210)}$ (*) | $0.27_{1,212}$ | $0.00_{(1,212)}$ | $0.09_{(1,211)}$ |
| $P_{Station}$ x Ploidy | $0.67_{(1,209)}$ | $1.75_{(1,209)}$ | $1.08_{(1,209)}$ | $0.50_{(1,209)}$ | $0.91_{(1,211)}$ | $0.48_{(1,211)}$ | $3.30_{(1,211)}$ |

[1] log-transformed

**Figure 4** Visualization of the significant interaction effect of the climatic conditions associated with different altitudes of the population origins (climPCA1) on TA-G concentration (a) and on concentration of di-PIEs (b) in the latex depending on the atmospheric pressure of the experimental station. Blue lines indicate predicted slopes from the mixed-effects model, with 95% confidence interval shaded. TA-G: taraxinic acid ß-D-glucopyranosyl ester; di-PIEs: di-4-hydroxyphenylacetate inositol ester.

In a next step, we constructed a full model to detect interactions between the climatic history of the populations (climPCA1) and the atmospheric pressure of the experimental sites at which the plants were grown ($P_{Station}$). The full model confirmed that the concentration of TA-G in the latex is influenced by climPCA1 (LMEM: climPCA1, $P = 0.017$, Table 1, Fig. 3), and that atmospheric pressure at the experimental sites is associated with changes in root growth (LMEM: $P_{Station}$, $P = 0.001$, Table 1, Fig. 2), root:shoot ratio (LMEM: $P_{Station}$, $P < 0.001$, Table 1, Fig. 2) and TA-G concentration in root latex (LMEM: $P_{Station}$, $P < 0.001$, Table 1, Fig. 2). In addition, we detected a negative effect of atmospheric pressure on the concentration of tri-PIEs

(LMEM: $P_{Station}$, $P$ = 0.019, Table 1, Fig. 2). We detected no significant interactions between the

climatic history of the populations (climPCA1) and atmospheric pressure at the experimental

sites for root or shoot growth (LMEM: climPCA1 x $P_{Station}$, $P$ = 0.05, Table 1). However, a

significant interaction was observed for TA-G concentration (LMEM: climPCA1 x $P_{Station}$, $P$ =

0.008, Table 1), indicating natural selection for phenotypic variability of TA-G production under

different atmospheric pressure. Closer inspection of the data revealed that TA-G concentrations

in populations originating from low altitude environments were more strongly influenced by

atmospheric pressure than in populations from high altitude environments (Fig. 4a). Under high

atmospheric pressure, populations from low altitude environments produced significantly more

TA-G than populations from high altitude environments (Fig. 4a). With decreasing atmospheric

pressure, this difference disappeared, leading to similar TA-G concentrations at 660 hPa. A

similar pattern, albeit with more variability, was observed for di-PIEs (LMEM: climPCA1 x

$P_{Station}$, $P$ = 0.012, Table 1, Fig. 4b). These results are suggestive of adaptation of root defence

expression of low altitude populations to high atmospheric pressure.

*The performance of triploid plants is not constrained by resistance to atmospheric pressure*

To test whether plant cytotype influences the capacity of *T. officinale* to grow at different

atmospheric pressures, we tested for interactions between cytotype and atmospheric pressure

at the different experimental sites. Across experimental sites, we did not detect any significant

differences between diploid and triploid plants for any of the measured parameters (LMEMs:

Ploidy, $P$ > 0.05, Table 1, Fig. 5). We also found no significant interactions between cytotype

and atmospheric pressure (LMEMs: $P_{Station}$ x Ploidy, $P$ > 0.05, Table 1, Fig. 5). Thus, the

hypothesis that low atmospheric pressure restricts the establishment of triploids at high

altitudes is not supported by our data.

**Figure 5** Visualization of influence of ploidy on plant parameters depending on atmospheric
pressure of experimental station. Predicted mean values with standard errors from mixed-
effects models are shown. None of the plant parameters differ statistically significantly between
diploid and triploid plants (n.s.: *P*>0.05). 2n: diploid; 3n: triploid; TA-G: taraxinic acid ß-D-
glucopyranosyl ester; di-PIEs: di-4-hydroxyphenylacetate inositol esters; tri-PIEs: tri-4-
hydroxyphenylacetate inositol esters.
424

**Discussion**

The sum of environmental factors varying with changing altitude can simultaneously impose multiple selection pressure on plant traits, making it difficult to disentangle single effects. By growing several *T. officinale* populations in controlled environments, our study identifies atmospheric pressure as factor that influences plant root growth and chemical defence irrespective of the cytotype level of the plants and shows that the natural habitats of the populations shape the potential for phenotypic variability in response to varying atmospheric pressure. Here, we discuss our findings in an eco-evolutionary and plant-physiological context.

Although many plants seem to be capable to grow vegetatively at pressures even below 25 kPa (Richards *et al.* 2006), they suffer from stress associated with hypoxia and desiccation and in turn develop responses and adaptations, which are often organ-specific (Zhou *et al.* 2017). In environments with reduced $O_2$ and $CO_2$ availability, roots may react more strongly to oxygen deficiency because, as heterotrophic organs, they are highly dependent on oxygen for mitochondrial energy production, while shoots, as autotrophic plant organs, may be restricted in photosynthesis due to the reduced $CO_2$ diffusion rate (Mustroph *et al.* 2014). Our results show that root but not shoot growth of *T. officinale* is reduced at low atmospheric pressure, resulting in a decrease of the root:shoot ratio. In our experimental stations, the reduction in atmospheric pressure is associated with naturally decreased levels of atmospheric gases including a reduced partial pressure of $O_2$, and we assume that the resulting mild hypoxia restricts root growth of *T. officinale*. This finding is in line with previous studies showing that roots are particularly sensitive to reduced $O_2$ partial pressure (He *et al.* 2007; Tang *et al.* 2015), whereas several studies with plants cultivated at reduced atmospheric pressure, but with a partial pressure of $O_2$ experimentally maintained at the same level as at ambient pressure, found no reduction in root biomass (Iwabuchi *et al.* 1996; Spanarkel & Drew 2002; Levine *et al.* 2008).

Exposure to low atmospheric pressure also affects plant chemistry (He *et al.* 2013; Levine *et al.* 2008; Zhou *et al.* 2017). In our study we detected a decline in the concentration of the secondary metabolite TA-G in the latex of *T. officinale* with decreasing atmospheric pressure. TA-G is constitutively produced by the plant, acts repellent against root feeders and therefore defends the plant against herbivores (Huber *et al.* 2016b; Bont *et al.* 2017). To reduce

the costs associated with constitutively produced chemical defenses (Neilson *et al.* 2013),

plants may use abiotic conditions as external stimuli to adjust the level of defensive secondary

metabolites to the expected herbivore pressure. Temperature, for instance, is assumed to be

a good indicator for herbivore attack in the field, because herbivore appearance and activity is

often modulated by temperature (Bale *et al.* 2002). In our previous work, we found evidence

that *T. officinale* can use seasonal temperature variation to synchronize deployment of

chemical defenses with expected herbivore attack intensity in the field, indicating an important

role of abiotic conditions in fine-tuning the level of constitutively produced defensive metabolites

(Huang *et al.* 2019). With increasing altitude, it is often assumed that herbivore pressure

decreases (Moreira *et al.* 2018). *T. officinale* might thus use low atmospheric pressure as an

indicator of high-altitude growth conditions and associated expected lower herbivore pressure,

and the observed decrease in TA-G production might be a fine-tuning of defense deployment

to reduce costs and maximize plant fitness. However, these are highly speculative conclusions

and require further investigations.

Our previous work has not only shown that TA-G is involved in herbivore defense in *T.*

*officinale*, but it has also shown an association of TA-G with the climatic history of the plants'

natural habitats, suggesting an additional role of TA-G in abiotic stress management (Bont *et*

*al.* 2020). The climatic parameters associated with TA-G production included sun and rain

intensity and were related to the latitude but not to the altitude of the natural populations –

plants from the rainier North of Switzerland produced more TA-G than plants from the sun-

intense regions in the South of Switzerland (Bont *et al.* 2020), leading to the hypothesis that

TA-G may passively or actively be involved in moisture regulation. At low atmospheric pressure,

water evaporation is increased, and evidence exists that when plants are grown under such

conditions, their perceptual mechanisms of water movement are altered even though the plants

are fully hydrated and do not experience actual desiccation (Paul *et al.* 2004; Zhou *et al.* 2017).

The decline in TA-G concentration with decreasing atmospheric pressure observed in our study

could therefore be a direct or indirect consequence of increased evapotranspiration and would

then support an involvement of TA-G in moisture regulation.

The capacity of phenotypic plasticity allows a plant with a given genotype to adjust its

phenotype to the demands of contrasting environments (Nicotra *et al.* 2010; Sultan 2000). In

our experiment, we exposed the same group of *T. officinale* genotypes to different atmospheric pressures while other abiotic parameters were kept constant, and subsequently detected plastic responses in growth and chemistry. Exposure to the atmospheric pressure gradient was sufficient to stimulate phenotypic plasticity, which again emphasizes the importance of atmospheric pressure for plant growth and development (Paul & Ferl 2006). For the latex metabolites TA-G and di-PIEs, plastic responses differed among populations, indicating heritable within-species variation for plasticity in these traits, likely shaped by the climatic histories of the populations. Under high atmospheric pressure, plants from low altitude environments produced more of the metabolites than plants from high altitude environments, whereas under low atmospheric pressure, the difference between the populations vanished. The capacity of low altitude plants to increase production of TA-G and di-PIEs when grown under low altitude atmospheric pressure could benefit these plants and may be an adaptive trait, as latex metabolites have defensive functions against root feeders (Huber *et al.* 2016a, b; Bont *et al.* 2017) and herbivore pressure is often expected to increase with decreasing altitude (Rasmann *et al.* 2014; Moreira *et al.* 2018). These findingsg are consistent with previous studies showing that within species, constitutive defenses often decrease with increasing altitude (Bakhtiari *et al.* 2019; Buckley *et al*. 2019; Meyer & Carlson 2001; Pellissier *et al.* 2014). Our results further suggest that climatic conditions characterizing low altitude environments select for genotypes with high plasticity in latex metabolite production, although whether the observed plasticity is a passive or active plastic response (van Kleunen & Fischer 2005) and whether its expression increases plant fitness remain to be elucidated.

Along altitudinal transects, diploid sexual *T. officinale* predominate over triploid asexual individuals at higher altitudes (Calame & Felber 2000), while along latitudinal gradients, triploids predominate over diploids at higher latitudes (Van Dijk *et al.* 2003). In our study, we tested the hypothesis that the establishment of triploids at high altitudes is restricted by low atmospheric pressure. However, we found no evidence for a disadvantage of triploids at low atmospheric pressure, as diploids and triploids did not differ in any of the measured plant traits at different atmospheric pressures. We conclude that atmospheric pressure is unlikely to influence geographic parthenogenesis of *T. officinale* and that other factors, for example recolonization

patters after glaciation (Calame & Felber 2000; Kearney 2005), might have influenced the

spatial distribution of diploids and triploids along certain transects.

Studies comparing different species have shown that plants growing in high altitude

environments have developed adaptations to cope with the challenges of these environments

(Halbritter *et al.* 2018) and that some of these adaptations, for example changes in stomatal

development, are directly related to atmospheric pressure (Kammer *et al.* 2015). Our study

confirms that atmospheric pressure is an important abiotic factor that influences plastic

responses in plant growth and development and shows that across *T. officinale* populations,

exposure to varying atmospheric pressures evokes heritable responses in growth and defense

traits - responses that are shaped by the climatic conditions of the natural habitats of the

populations. Given the non-negligible impact of atmospheric pressure on the expression and

likely also on the evolution of plant traits, we suggest that atmospheric pressure should be

included by default as an abiotic factor when studying plant variation along altitudinal gradients

to prevent results from being obscured by possibly unrecognized effects of atmospheric

pressure variation. This may be particularly important when altitudinal gradients are used to

surrogate climate change (Carlyle *et al.* 2014; Frei *et al.* 2014; Michalet *et al.* 2014), since

atmospheric pressure, unlike temperature or precipitation, is not expected to change under

global warming.

Our results emphasize that within species, plants from different populations respond

differently to varying atmospheric pressure, especially in the production of secondary

metabolites. For successful cultivation of plants in extraterrestrial habitats as food source, it

therefore may be worthwhile to screen various populations of the species of interest under low

atmospheric pressure to detect resistant populations with secondary metabolite profiles

suitable for human nutrition.

**Acknowledgements**

We thank Marc Pfander for his support with statistical analysis. We are grateful to Andrea Bonini for his help with field and lab work and to the gardeners of the University of Bern for taking care of the plants. We thank Hildegard Hinni, Annerös Erb, André Hofer, Christoph Steiner, Joan Fischer and Martin Fischer for watering the plants during the experiments at the research stations in Adlemsried, Kleine Scheidegg and Jungfraujoch. We thank the Kleine Scheidegg station of the Swiss Federal Railways to provide the room to perform the experiment. We thank the International Foundation High Altitude Research Station Jungfraujoch and Gornergrat (HFSJG) for the opportunity to conduct experiments. Meteorological data was provided by MeteoSwiss, the Swiss Federal Office for Meteorology and Climatology. This study was supported by the Swiss National Science Foundation (Grant No. 153517) and the Seventh Framework Programme for Research and Technological Development of the European Union (FP7 MC-CIG 629134).

**References**

Agrawal, A.A. & Konno, K. (2009). Latex: a model for understanding mechanisms, ecology, and evolution of plant defense against herbivory. *Annu. Rev. Ecol. Evol. Syst.*, 40, 311–331.

Agrawal, A.A., Hastings, A.P., Johnson, M.T.J., Maron, J.L. & Salminen, J.-P. (2012). Insect herbivores drive real-time ecological and evolutionary change in plant populations. *Science,* 338, 113–116.

Angel, R., Conrad, R., Dvorsky, M., Kopecky, M., Kotilínek, M., Hiiesalu, I., *et al.* (2016). The root-associated microbial community of the world's highest growing vascular plants. *Microb. Ecol.*, 72, 394–406.

Arbona, V., Manzi, M., de Ollas, C. & Gómez-Cadenas, A. (2013). Metabolomics as a tool to investigate abiotic stress tolerance in plants. *Int. J. Mol. Sci.*, 14, 4885–4911.

Bakhtiari, M., Formenti, L., Caggìa, V., Glauser, G. & Rasmann, S. (2019). Variable effects on growth and defense traits for plant ecotypic differentiation and phenotypic plasticity along elevation gradients. *Ecol. Evol.*, 9, 3740–3755.

Baldwin, I.T., Halitschke, R., Paschold, A., Von Dahl, C.C. & Preston, C.A. (2006). Volatile signaling in plant-plant interactions: "Talking trees" in the genomics era. *Science*, 311, 812–815.

Bale, J.S., Masters, G.J., Hodkinson, I.D., Awmack, C., Bezemer, T.M., Brown, V.K., *et al.* (2002). Herbivory in global climate change research: direct effects of rising temperature on insect herbivores. *Glob. Chang. Biol.*, 8, 1–16.

Bates, D., Maechler, M., Bolker, B.M. & Walker, S. (2015). Fitting linear mixed-effects models using {lme4}. *J. Stat. Softw.*, 67, 1–48.

Bates, D.M. (2020). lme4 : Mixed-effects modeling with R.

Bednarek, P. & Osbourn, A. (2009). Plant-microbe interactions: chemical diversity in plant defense. *Science*, 324, 746–748.

Bernal, M., Llorens, L., Julkunen-Tiitto, R., Badosa, J. & Verdaguer, D. (2013). Altitudinal and seasonal changes of phenolic compounds in *Buxus sempervirens* leaves and cuticles. *Plant Physiol. Biochem.*, 70, 471–482.

Bierzychudek, P. (1985). Patterns in plant parthenogenesis. *Experientia*, 41, 1255–1264.

Bont, Z., Arce, C., Huber, M., Huang, W., Mestrot, A., Sturrock, C.J., *et al.* (2017). A herbivore

tag-and-trace system reveals contact- and density-dependent repellence of a root toxin.

*J. Chem. Ecol.*, 43, 295–306.

Bont, Z., Züst, T., Arce, C.C.M., Huber, M. & Erb, M. (2020). Heritable variation in root

secondary metabolites is associated with recent climate. *J. Ecol.*, 1–14.

Buckley, J., Widmer, A., Mescher, M.C. & De Moraes, C.M. (2019). Variation in growth and

defence traits among plant populations at different elevations: implications for

adaptation to climate change. *J. Ecol.*, 107, 2478–2492.

Buckley, T.N. (2005). The control of stomata by water balance. *New Phytol.*, 168, 275–292.

Calame, F. & Felber, F. (2000). Distribution of diploid sexual and triploid apomictic dandelions

(*Taraxacum* sect. *Ruderalia*) along two altitudinal gradients in Switzerland. *Bot. Helv.*,

110, 109–114.

Carlyle, C.N., Fraser, L.H. & Turkington, R. (2014). Response of grassland biomass

production to simulated climate change and clipping along an elevation gradient.

*Oecologia*, 174, 1065–1073.

Corey, K.A., Barta, D.J. & Henninger, D.L. (1997). Photosynthesis and respiration of a wheat

stand at reduced atmospheric pressure and reduced oxygen. *Adv. Sp. Res.*, 20, 1869–

1877.

Cosendai, A.C., Wagner, J., Ladinig, U., Rosche, C. & Hörandl, E. (2013). Geographical

parthenogenesis and population genetic structure in the alpine species *Ranunculus*

*kuepferi* (Ranunculaceae). *Heredity*, 110, 560–569.

Cunningham, S.A., Summerhayes, B. & Westoby, M. (1999). Evolutionary divergences in leaf

structure and chemistry, comparing rainfall and soil nutrient gradients. *Ecology*, 69,

569–588.

Van Dijk, P.J., Barrett, S.C.H. & Vinkenoog, R. (2003). Ecological and evolutionary

opportunities of apomixis: insights from *Taraxacum* and *Chondrilla*. *Philos. Trans. R.*

*Soc. B Biol. Sci.*, 358, 1113–1121.

Ehrlich, P.R. & Raven, P.H. (1964). Butterflies and plants: a study in coevolution. *Evolution*,

18, 586–608.

Ferl, R.J., Schuerger, A.C., Paul, A.-L., Gurley, W.B., Corey, K. & Bucklin, R. (2002). Plant

adaptation to low atmospheric pressures: potential molecular responses. *Life Support*

*Biosph. Sci.*, 8, 93–101.

Fox, J., Weisberg, S., Price, B., Friendly, M., Firth, D. & Taylor, S. (2019). Package ' effects .'

Frei, E.R., Ghazoul, J., Matter, P., Heggli, M. & Pluess, A.R. (2014). Plant population

differentiation and climate change: responses of grassland species along an elevational

gradient. *Glob. Chang. Biol.*, 20, 441–455.

Glesener, R.R. & Tilman, D. (1978). Sexuality and the components of environmental

uncertainty: clues from geographic parthenogenesis. *Am. Nat.*, 112, 659–673.

Halbritter, A.H., Fior, S., Keller, I., Billeter, R., Edwards, P.J., Holderegger, R., *et al.* (2018).

Trait differentiation and adaptation of plants along elevation gradients. *J. Evol. Biol.*, 31,

784–800.

Halekoh, U. & Højsgaard, S. (2014). A Kenward-Roger approximation and parametric

bootstrap methods for tests in linear mixed models - the R package pbkrtest. *J. Stat.*

*Softw.*, 59, 1–32.

Hartmann, T. (2007). From waste products to ecochemicals: fifty years research of plant

secondary metabolism. *Phytochemistry*, 68, 2831–2846.

He, C., Davies, F.T. & Lacey, R.E. (2007). Separating the effects of hypobaria and hypoxia on

lettuce: growth and gas exchange. *Physiol. Plant.*, 131, 226–240.

He, C., Davies F.T., Lacey, R.E., Drew, M.C. & Brown, D.L. (2003). Effect of hypobaric

conditions on ethylene evolution and growth of lettuce and wheat. *J. Plant Physiol.*, 160,

1341–1350.

He, C., Jacobo-Velázquez, D.A., Cisneros-Zevallos, L. & Davies, F.T. (2013). Hypobaria and

hypoxia affects phytochemical production, gas exchange, and growth of lettuce.

*Photosynthetica*, 51, 465–473.

Hervé, M. (2018). Package "RVAideMemoire."

Hörandl, E. (2006). The complex causality of geographical parthenogenesis. *New Phytol.*,

171, 525–538.

Huang, W., Bont, Z., Hervé, M.R., Robert, C.A.M. & Erb, M. (2019). Impact of seasonal and

temperature-dependent variation in root defense metabolites on herbivore preference in

*Taraxacum officinale. J. Chem. Ecol.*

Huber, M., Bont, Z., Fricke, J., Brillatz, T., Aziz, Z., Gershenzon, J., *et al.* (2016a). A below-

ground herbivore shapes root defensive chemistry in natural plant populations. *Proc.*

*Biol. Sci.*, 283, 1–10.

Huber, M., Epping, J., Schulze Gronover, C., Fricke, J., Aziz, Z., Brillatz, T., *et al.* (2016b). A

latex metabolite benefits plant fitness under root herbivore attack. *PLoS Biol.*, 14, 1–27.

Huber, M., Triebwasser-Freese, D., Reichelt, M., Heiling, S., Paetz, C., Chandran, J.N., *et al.*

(2015). Identification, quantification, spatiotemporal distribution and genetic variation of

major latex secondary metabolites in the common dandelion (*Taraxacum officinale*

agg.). *Phytochemistry*, 115, 89–98.

Iacobucci, D., Schneider, M.J., Popovich, D.L. & Bakamitsos, G.A. (2016). Mean centering

helps alleviate "micro" but not "macro" multicollinearity. *Behav. Res. Methods*, 48, 1308–

1317.

Iwabuchi, K., Goto, E. & Takakura, T. (1996). Germination and growth of spinach under

hypobaric conditions. *Environ. Control Biol.*, 34, 169–178.

Iwabuchi, K. & Kurata, K. (2003). Short-term and long-term effects of low total pressure on

gas exchange rates of spinach. *Adv. Sp. Res.*, 31, 241–244.

Jump, A.S. & Peñuelas, J. (2005). Running to stand still: adaptation and the response of

plants to rapid climate change. *Ecol. Lett.*, 8, 1010–1020.

Kammer, P.M., Steiner, J.S. & Schöb, C. (2015). *Arabis alpina* and *Arabidopsis thaliana* have

different stomatal development strategies in response to high altitude pressure

conditions. *Alp. Bot.*, 125, 101–112.

Kearney, M. (2005). Hybridization, glaciation and geographical parthenogenesis. *Trends Ecol.*

*Evol.*, 20, 495–502.

Keller, S.R., Sowell, D.R., Neiman, M., Wolfe, L.M. & Taylor, D.R. (2009). Adaptation and

colonization history affect the evolution of clines in two introduced species. *New Phytol.*,

183, 678–690.

Kessler, A. & Baldwin, I.T. (2001). Defensive function of herbivore-induced plant volatile

emissions in nature. *Science*, 291, 2141–2144.

Kessler, A. & Kalske, A. (2018). Plant secondary metabolite diversity and species

interactions. *Annu. Rev. Ecol. Evol. Syst.*, 49, 115–138.

van Kleunen, M. & Fischer, M. (2005). Constraints on the evolution of adaptive phenotypic

plasticity in plants. *New Phytol.,* 166, 49–60.

Konno, K. (2011). Plant latex and other exudates as plant defense systems: roles of various

defense chemicals and proteins contained therein. *Phytochemistry*, 72, 1510–1530.

Kooyers, N.J., Greenlee, A.B., Colicchio, J.M., Oh, M. & Blackman, B.K. (2015). Replicate

altitudinal clines reveal that evolutionary flexibility underlies adaptation to drought stress

in annual *Mimulus guttatus. New Phytol.*, 206, 152–165.

Kuznetsova, A., Brockhoff, P.B. & Christensen, R.H.B. (2017). lmerTest package: tests in

linear mixed effects models. *J. Stat. Softw.*, 82.

Levine, L.H., Bisbee, P.A., Richards, J.T., Birmele, M.N., Prior, R.L., Perchonok, M., *et al.*

(2008). Quality characteristics of the radish grown under reduced atmospheric pressure.

*Adv. Sp. Res.*, 41, 754–762.

Menken, S.B.J., Smit, E. & Den Nijas, H.C.M. (1995). Genetic population structure in plants:

gene flow between diploid sexual and triploid asexual dandelions (*Taraxacum* section

*ruderalia*). *Evolution*, 49, 1108–1118.

Metlen, K.L., Aschehoug, E.T. & Callaway, R.M. (2009). Plant behavioural ecology: dynamic

plasticity in secondary metabolites. *Plant, Cell Environ.*, 32, 641–653.

Meyer, S.E. & Carlson, S.L. (2001). Achene mass variation in *Ericameria nauseosus*

(Asteraceae) in relation to dispersal ability and seedling fitness. *Funct. Ecol.*, 15, 274–

281.

Michalet, R., Schöb, C., Lortie, C.J., Brooker, R.W. & Callaway, R.M. (2014). Partitioning net

interactions among plants along altitudinal gradients to study community responses to

climate change. *Funct. Ecol.*, 28, 75–86.

Midolo, G. & Wellstein, C. (2020). Plant performance and survival across transplant

experiments depend upon temperature and precipitation change along elevation. *J.*

*Ecol.*, 1–14.

Moles, A.T., Peco, B., Wallis, I.R., Foley, W.J., Alistair, G.B., Seabloom, E.W., *et al.* (2013).

Correlations between physical and chemical defences in plants: tradeoffs, syndromes,

or just many different ways to skin a herbivorous cat? *New Phytol.*, 198, 252–263.

Molina-Montenegro, M., Palma-Rojas, C., Alcayaga-Olivares, Y., Oses, R., Corcuera, L.J.,

Cavieres, L. a., *et al.* (2013). Ecophysiological plasticity and local differentiation help

explain the invasion success of *Taraxacum officinale* (dandelion) in South America.

*Ecography*, 36, 718–730.

Moore, B.D., Andrew, R.L., Külheim, C. & Foley, W.J. (2014). Explaining intraspecific diversity

in plant secondary metabolites in an ecological context. *New Phytol.*, 201, 733–50.

Moreira, X., Petry, W.K., Mooney, K.A., Rasmann, S. & Abdala-Roberts, L. (2018).

Elevational gradients in plant defences and insect herbivory: recent advances in the field

and prospects for future research. *Ecography*, 41, 1485–1496.

Mustroph, A., Barding, G.A., Kaiser, K.A., Larive, C.K. & Bailey-Serres, J. (2014).

Characterization of distinct root and shoot responses to low-oxygen stress in

*Arabidopsis* with a focus on primary C- and N-metabolism. *Plant, Cell Environ.*, 37,

2366–2380.

Nakabayashi, R. & Saito, K. (2015). Integrated metabolomics for abiotic stress responses in

plants. *Curr. Opin. Plant Biol.*, 24, 10–16.

Neilson, E.H., Goodger, J.Q.D., Woodrow, I.E. & Møller, B.L. (2013). Plant chemical defense:

at what cost? *Trends Plant Sci.*, 18, 250–258.

Nicotra, A.B., Atkin, O.K., Bonser, S.P., Davidson, A.M., Finnegan, E.J., Mathesius, U., *et al.*

(2010). Plant phenotypic plasticity in a changing climate. *Trends Plant Sci.*, 15, 684–

692.

Paul, A.-L. & Ferl, R.J. (2006). The biology of low atmospheric pressure – implications for

exploration mission design and advanced life support. *Gravitational Sp. Biol.*, 19, 3–18.

Paul, A.L., Schuerger, A.C., Popp, M.P., Richards, J.T., Manak, M.S. & Ferl, R.J. (2004).

Hypobaric biology: *Arabidopsis* gene expression at low atmospheric pressure. *Plant*

*Physiol.*, 134, 215–223.

Pellissier, L., Roger, A., Bilat, J. & Rasmann, S. (2014). High elevation *Plantago lanceolata*

plants are less resistant to herbivory than their low elevation conspecifics: is it just

temperature? *Ecography.*, 37, 950–959.

Pichersky, E. & Gershenzon, J. (2002). The formation and function of plant volatiles:

perfumes for pollinator attraction and defense. *Curr. Opin. Plant Biol.*, 5, 237–243.

R Core Team. (2017). *R: a language and environment for statistical computing. R Found.*

*Stat. Comput. Vienna, Austria.* Available at: http://www.r-project.org/.

Ramakrishna, A. & Ravishankar, G.A. (2011). Influence of abiotic stress signals on secondary

metabolites in plants. *Plant Signal. Behav.*, 6, 1720–1731.

Rasmann, S., Buri, A., Gallot-Lavallée, M., Joaquim, J., Purcell, J. & Pellissier, L. (2014).

Differential allocation and deployment of direct and indirect defences by *Vicia sepium*

along elevation gradients. *J. Ecol.*, 102, 930–938.

Richards, J.T., Corey, K.A., Paul, A.L., Ferl, R.J., Wheeler, R.M. & Schuerger, A.C. (2006).

Exposure of *Arabidopsis thaliana* to hypobaric environments: implications for low-

pressure bioregenerative life support systems for human exploration missions and

terraforming on Mars. *Astrobiology*, 6, 851–866.

Sandoya, V., Pauchard, A. & Cavieres, L.A. (2017). Natives and non-natives plants show

different responses to elevation and disturbance on the tropical high Andes of Ecuador.

*Ecol. Evol.*, 7, 7909–7919.

Schliemann, W., Ammer, C. & Strack, D. (2008). Metabolite profiling of mycorrhizal roots of

*Medicago truncatula*. *Phytochemistry*, 69, 112–146.

Semchenko, M., Saar, S. & Lepik, A. (2014). Plant root exudates mediate neighbour

recognition and trigger complex behavioural changes. *New Phytol.*, 204, 631–637.

Smith, W.K. & Geller, G.N. (1979). Plant transpiration at high elevations: theory, field

measurements, and comparisons with desert plants. *Oecologia*, 41, 109–122.

Spanarkel, R. & Drew, M.C. (2002). Germination and growth of lettuce (*Lactuca sativa*) at low

atmospheric pressure. *Physiol. Plant.*, 116, 468–477.

Stevenson, P.C., Nicolson, S.W. & Wright, G.A. (2017). Plant secondary metabolites in

nectar: impacts on pollinators and ecological functions. *Funct. Ecol.*, 31, 65–75.

Stewart-Wade, S.M., Neumann, S., Collins, L.L. & Boland, G.J. (2002). The biology of

Canadian weeds. 117. *Taraxacum officinale* G. H. Weber ex Wiggers. *Can. J. Plant Sci.*,

825–853.

Sultan E. Sonia. (2000). Phenotypic plasticity for plant development, function and life history.

*Trends Plant Sci.*, 5, 537–542.

Tang, Y., Gao, F., Guo, S. & Li, F. (2015). The morphology, physiology and nutritional quality

of lettuce grown under hypobaria and hypoxia. *Acta Astronaut.*, 112, 29–36.

Terashima, I., Masuzawa, T., Ohba, H. & Yokoi, Y. (1995). Is photosynthesis suppressed at

higher elevations due to low CO2 pressure ? *Ecology*, 76, 2663–2668.

Tilquin, A. & Kokko, H. (2016). What does the geography of parthenogenesis teach us about

sex? *Philos. Trans. R. Soc. Lond. B. Biol. Sci.*, 371, 20150516–20150538.

Verduijn, M.H., Van Dijk, P.J. & Van Damme, J.M.M. (2004). The role of tetraploids in the

sexual-asexual cycle in dandelions (*Taraxacum*). *Heredity*, 93, 390–398.

Verhoeven, K.J.F. & Biere, A. (2013). Geographic parthenogenesis and plant-enemy

interactions in the common dandelion. *BMC Evol. Biol.*, 13.

Villaverde, T., González-Moreno, P., Rodríguez-Sánchez, F. & Escudero, M. (2017). Niche

shifts after long-distance dispersal events in bipolar sedges (*Carex*, Cyperaceae). *Am.*

*J. Bot.*, 104, 1765–1774.

Ward, J.K., Antonovics, J., Thomas, R.B. & Strain, B.R. (2000). Is atmospheric CO2 a

selective agent on model C3 annuals? *Oecologia*, 123, 330–341.

Ward, J.K. & Strain, B.R. (1997). Effects of low and elevated CO2 partial pressure on growth

and reproduction of *Arabidopsis thaliana* from different elevations. *Plant, Cell Environ.*,

20, 254–260.

Wheeler, R.M. (2017). Agriculture for space: people and places paving the way. *Open Agric.*,

2, 14–32.

Wickham, H. (2016). *ggplot2: elegant graphics for data analysis*. Springer-Verlag New York.

Woodward, F.I. & Bazzaz, F.A. (1988). The responses of stomatal density to CO2 partial

pressure. *J. Exp. Bot.*, 39, 1771–1781.

Xu, Z., Jiang, Y., Jia, B. & Zhou, G. (2016). Elevated-CO2 response of stomata and its

dependence on environmental factors. *Front. Plant Sci.*, 7, 1–15.

Xu, Z., Jiang, Y. & Zhou, G. (2015). Response and adaptation of photosynthesis, respiration,

and antioxidant systems to elevated CO2 with environmental stress in plants. *Front.*

*Plant Sci.*, 6, 1–17.

Zabalza, A., Van Dongen, J.T., Froehlich, A., Oliver, S.N., Faix, B., Gupta, K.J., *et al.* (2009).

Regulation of respiration and fermentation to control the plant internal oxygen

concentration. *Plant Physiol.*, 149, 1087–1098.

Zhou, M., Callaham, J.B., Reyes, M., Stasiak, M., Riva, A., Zupanska, A.K., *et al.* (2017).

Dissecting low atmospheric pressure stress: transcriptome responses to the

components of hypobaria in *Arabidopsis. Front. Plant Sci.*, 8, 1–13.

Züst, T., Heichinger, C., Grossniklaus, U., Harrington, R., Kliebenstein, D.J. & Turnbull, L.A.

(2012). Natural enemies drive geographic variation in plant defenses. *Science*, 338,

116–120.