Heterologous vaccination regimens with self-amplifying RNA and Adenoviral

COVID vaccines induce superior immune responses than single dose vaccine

regimens in mice

Alexandra J Spencer[1*], Paul F McKay[2], Sandra Belij-Rammerstorfer[1], Marta Ulaszewska[1],

Cameron D Bissett[1], Kai Hu[2], Karnyart Samnuan[2], Daniel Wright[1], Hannah R Sharpe[1], Ciaran

Gilbride[1], Adam Truby[1], Elizabeth R Allen[1], Sarah C Gilbert[1], Robin J Shattock[2], Teresa

Lambe[1]

Affiliations:

[1]The Jenner Institute, Nuffield Department of Medicine, University of Oxford,

United Kingdom

[2]Department of Infectious Disease, Imperial College London, United Kingdom

Correspondence to: Alexandra J Spencer, The Jenner Institute, ORCRB, Roosevelt Drive,

Oxford OX3 7DQ. Email: alex.spencer@ndm.ox.ac.uk.

Short title: Superior immunity with heterologous vaccination regimens

Abstract

Several vaccines have demonstrated efficacy against SARS-CoV-2 mediated disease, yet

there is limited data on the immune response induced by heterologous vaccination

regimens using alternate vaccine modalities. Here, we present a detailed description of the

immune response, in mice, following vaccination with a self-amplifying RNA (saRNA) vaccine

and an adenoviral vectored vaccine (ChAdOx1 nCoV-19/AZD1222) against SARS-CoV-2. We

demonstrate that antibody responses are higher in two dose heterologous vaccination

regimens than single dose regimens, with high titre neutralising antibodies induced.

Importantly, the cellular immune response after a heterologous regimen is dominated by

cytotoxic T cells and Th1$^+$ CD4 T cells which is superior to the response induced in

homologous vaccination regimens in mice.

Introduction

A number of vaccines against SARS-CoV-2 have reached late-stage clinical trials with

encouraging efficacy readouts and mass vaccination schemes already initiated across

several countries. Multiple vaccine technologies are being advanced [1], but there is limited

information on how these vaccine modalities may work in combination. With ongoing

clinical trial assessment of numerous SARS-CoV-2 vaccines and new mass vaccination

incentives, there is a recognised real-world scenario where individuals may be vaccinated

with different vaccine modalities. However, the utility of vaccination regimens combining

different types of vaccine approaches remains to be determined.

Self-amplifying RNA (saRNA) typically encodes the alphaviral replicase and a target antigen.

Upon entry into the cytoplasm, the RNA is amplified with subsequent translation of the

target antigen [2]. The SARS-CoV-2 saRNA vaccine has been demonstrated to be highly

immunogenic in preclinical animal models [3] and is progressing through clinical trial

assessment with promising results. Adenoviruses are a frequently used viral vector vaccine

technology, they can be rapidly made to GMP at large scale, and a single vaccination can be

sufficient to provide rapid immunity in individuals [4]. In particular, chimpanzee derived

adenoviruses (ChAd) have good safety profiles, whilst inducing strong cellular and humoral

immune response against multiple target disease antigens [5-7]. Previous studies have

demonstrated efficacy after vaccination with ChAdOx1 nCoV-19/AZD1222 against SARS-

CoV-2 mediated disease [8], with concomitant high titre humoral immune responses and a

measurable Th1-dominated cellular immune response [9,10]. Importantly, an immune profile

consistently observed across different animal species [11,12].

In this study, the immunogenicity of saRNA and ChAdOx1 vaccines expressing full-length

SARS-CoV-2 spike protein were assessed in mice following vaccination with different vaccine

modalities combination. We demonstrate robust antibody responses following

heterologous vaccination regimens, with high titre neutralising antibodies. The cellular

immune response is dominated by cytotoxic T cells secreting IFNγ and TNFα and antigen

specific CD4+ T cells of a Th1 phenotype, with significantly higher antigen specific responses

observed following heterologous vaccination than those responses induced in single vaccine

regimens.

Results

Heterologous vaccination potentiates SARS-CoV-2 spike-specific antibody response

It was previously shown that homologous prime-boost immunization with saRNA [3] or ChAd

[11] induced a SARS-CoV-2 spike-specific IgG response with neutralisation capacity. In this

series of experiments, we compared the immune response induced by heterologous,

homologous and single dose vaccination with ChAd and saRNA vaccine modalities.

Heterologous vaccination with either ChAd or saRNA prime and alternative boost (i.e. ChAd-

saRNA or saRNA-ChAd) or two doses of saRNA induced the highest IgG responses, compared

to mice vaccinated with a single dose of either ChAd or saRNA (Fig. 1A and Fig. S1A), with

strong correlation between the two independent IgG ELISA methods (Fig. S1B). This IgG

response showed mixed profile, with mainly IgG2 (IgG2a, IgG2b in both mouse strains and

IgG2c in CD1 mice) and IgG1 subclasses (in both mouse strains) (Fig. 2A and B), which was

similar to previously reported results after vaccination with ChAd alone (Silva-Cayetano in

press) or saRNA alone [3].

Comparable amounts of SARS-CoV-2 spike-specific IgM were detected in all vaccinated

groups (Fig. 1B), while heterologous vaccination with saRNA-ChAd induced higher serum

SARS-CoV-2 spike-specific IgA levels compared to single vaccination with either vaccine, in

both CD1 and BALB/c mice (Fig. 1B).

Two doses of vaccine were also shown to increase the avidity of the IgG towards SARS-CoV-

2 spike (Fig. 1C) compared to single administration of either vaccine, while antibody-

mediated neutralisation was measured across all groups of vaccinated mice, with

heterologous and two dose saRNA inducing significantly higher levels of neutralisation

compared to single dose vaccination regimens (Fig. 1D). Not surprisingly, a correlation was

measured between neutralisation and levels of IgG /IgA SARS-CoV-2 spike-specific

antibodies (Fig. S1C).

Flow cytometry staining enabled identification of SARS-CoV-2 spike-specific B cells in the

spleen of mice. The data demonstrated that all vaccination regimens induced a similar

number of antigen specific B cells (Fig. S2B), with similar numbers of germinal centre (GC)

and isotype class switched B cells observed in all groups of vaccinated mice.

101   Heterologous vaccination induces strong Th1 type response

Cell mediated immunity was also investigated following heterologous or homologous

ChAdOx1 and saRNA vaccination regimens. The highest IFNγ response detected by ELISpot

was observed in mice that received a heterologous combination of vaccines, this increase

was only statistically significant when compared to single administration of saRNA in both

strains of mice (Fig. 3A and B), or saRNA-ChAd compared to ChAd in BALB/c mice (Fig. 3B). In

agreement with earlier reports, ELISpot responses were primarily directed towards the S1

portion of the spike protein (Fig S3A), with a consistent breadth of response measured with

all vaccine combinations and in both strains of mice (Fig. 3A and B).

Phenotype and functional capacity of the T cell response was measured by intracellular

cytokine staining, together with memory T cell marker staining. Consistent with previously

published data, saRNA and ChAd vectors induced antigen specific CD4$^+$ T cells of a Th1 bias,

with minimal IL4 and IL10 production measured and a response dominated by production of

IFNγ and TNFα, regardless of the vaccination regimen (Fig. 4A). Antigen specific CD4$^+$ T cells

displayed mixed Teff and Tem phenotype, with no statistically significant differences

observed in the total number of cells (Fig. 4A) and cell subsets (Fig. S3C). As before, in mice

the cell-mediated response following vaccination was dominated by CD8$^+$ T cells, with much

higher levels of IFNγ and TNFα (in addition to upregulation of CD107a) seen in all vaccine

groups (Fig. 4B) compared to CD4$^+$ T cell responses. In addition, the highest frequency of

antigen specific cells, in both outbred and inbred strains of mice, was measured following

heterologous vaccination, regardless of vaccine order (Fig. 4B). Antigen specific CD8$^+$ T cells

displayed a predominantly Teff phenotype, with the highest number of total antigen specific

cells (Fig. 4B) observed in heterologous vaccinated animals.

Discussion

The current pandemic and extraordinary efforts to develop effective vaccines with subsequent mass vaccination roll-out has highlighted the 'real-world' practicalities of global vaccination campaigns. There are around 20 vaccines in Phase 3 clinical trial assessment, and several vaccines have already reported efficacy with subsequent emergency licensure granted in some countries. While each individual vaccine candidate has been thoroughly tested for safety and efficacy, there have been no studies reported to date that have examined the safety, efficacy or any added benefit of mixed modality vaccinations. Given the real-world vaccination initiatives that are being progressed, there may be a scenario wherein an individual receives a vaccine prime and a boost dose from different manufacturers or of different vaccine types. This current pre-clinical study examined the cellular and humoral immune responses in mice following vaccination with either the ChAdOx adenoviral vector or the saRNA LNP in homologous or heterologous prime-boost combinations and compared immunogenicity to single moiety vaccinations.

All prime-boost regimens elicited high levels of SARS-CoV-2 spike-specific antibodies with neutralisation capacity and high avidity, levels that were greater than single vaccines alone. While heterologous vaccination regimens induced the highest antibody responses post-vaccination, homologous saRNA and ChAd induced higher antibody responses than single dose regimens, consistent with previous data in mice, pigs and NHPs [11,12]. Importantly in human clinical trials, strong enhancement of the antibody response was observed following a booster dose of ChAdOx1[13] , with this regimen shown to provide efficacy against SARS-CoV-2 disease in late-stage clinical trials[8]. The cell-mediated response was dominated by cytotoxic T cells, with heterologous regimens inducing higher frequencies and total numbers

of antigen specific CD8[+] T cells. Although CD4[+] T cell responses were overall lower in

frequency and number than CD8[+] T cell responses, all vaccination regimens elicited CD4[+] T

cell responses of a Th1 type response, even in the Th2 biased BALB/c background, a

response indicating that the potential for antibody dependent enhancement (ADE) and

subsequent enhancement of respiratory disease (ERD) caused by Th2-type lung

immunopathology is reduced [14-16].

While there are no defined correlates of protection, Rhesus Macaques studies have

demonstrated a clear role for neutralising antibodies and also CD8[+] T cells [17]. In agreement,

human studies have demonstrated neutralising antibodies and T cells play an important role

in preventing severe disease and augmenting recovery from COVID-19[18]. Both vaccine

modalities elicited high numbers of antigen specific T cells, which were further increased in

the heterologous regimens. The majority of the IFNγ ELISpot response was directed against

the S1 spike protein, particularly the first half (317 AA) which does not include the RBD. Not

surprisingly two doses regimens induced higher antibody responses, with combination

regimens leading to simultaneous high levels of neutralising antibodies and T cell responses.

Vaccination regimens that induce a broad immune response (humoral and cell-mediated)

will likely be the best option for long-term protection against severe disease and will be best

placed if booster vaccinations are considered in the future. The logistical challenges of

administering vaccines in a rapidly evolving landscape of mass vaccination schemes, coupled

with limited global supply, underpins the need to generate data on mixing vaccine

modalities. The data described herein suggests clinical trials should progress to assess the

safety, immunogenicity and efficacy of heterologous vaccination regimens to help progress

global deployment and vaccine uptake against SARS-CoV-2.

Methods

Ethics Statement; Mice were used in accordance with the UK Animals (Scientific

Procedures) Act under project license number P9804B4F1 granted by the UK Home Office.

Age matched animals were purchased from commercial suppliers as a batch for each

experiment and randomly split into groups on arrival at our facility. Animals were group

housed in IVCs under SPF conditions, with constant temperature and humidity with lighting

on a 13:11 light-dark cycle (7am to 8pm). For induction of short-term anaesthesia, animals

were anaesthetised using vaporised IsoFlo®. All animals were humanely sacrificed at the end

of each experiment by an approved Schedule 1 method.

Animals and Immunizations; Outbred CD1Hsd:ICR (CD-1) (Envigo) (n=8 per group), Crl:CD1

(ICR) (Charles River) (n=8 per group) and inbred BALB/cOlaHsd (BALB/c) (Envigo) (n=6 per

group) mice of 7 weeks of age, were immunized intramuscularly (i.m.) in the musculus

tibialis with 10^8 infectious units (iu) of ChAdOx1 nCoV-19 [12] or 1μg saRNA [3]. Mice were

boosted i.m. with the relevant vaccine candidate 4 weeks later. All mice were sacrificed 3

weeks after the final vaccination with serum and spleens collected for analysis of humoral

and cell-mediated immunity.

Pseudotype virus neutralisation assay; A HIV-pseudotyped luciferase-reporter based

system was used to assess the neutralization ability of sera from vaccinated mice. In brief,

CoV S-pseudotyped viruses were produced by co- transfection of 293T/17 cells with a HIV-1

gag-pol plasmid (pCMV-Δ8.91, a kind gift from Prof. Julian Ma, St George's University of

London), a firefly luciferase reporter plasmid (pCSFLW, a kind gift from Prof. Julian Ma, St

George's University of London) and a plasmid encoding the S protein of interest (pSARS-

CoV2-S) at a ratio of 1:1.5:1. Virus- containing medium was clarified by centrifugation and

filtered through a 0.45 µm membrane 72 h after transfection, and subsequently aliquoted

and stored at −80 °C. For the neutralization assay, heat-inactivated sera were first serially

diluted and incubated with virus for 1 h, and then the serum-virus mixture was transferred

into wells pre-seeded Caco2 cells. After 48 h, cells were lysed, and luciferase activity was

measured using Bright-Glo Luciferase Assay System (Promega). The IC50 neutralization was

then calculated using GraphPad Prism (version 8.4). Statistical analyses were performed on

log transformed data.

Antigen-specific IgG ELISA; The antigen-specific IgG titres in mouse sera were assessed by a

semi-quantative ELISA. MaxiSorp high binding ELISA plates (Nunc) were coated with 100 µL

per well of 1 µg mL−1 recombinant SARS-CoV-2 protein with the pre-fusion stabilized

conformation in PBS. After overnight incubation at 4 °C, the plates were washed 4 times

with PBS-Tween 20 0.05% (v/v) and blocked for 1 h at 37 °C with 200 µL per well blocking

buffer (1% BSA (w/v) in PBS-Tween-20 0.05%(v/ v)). The plates were then washed and the

diluted samples or a 5-fold dilution series of the standard IgG added using 50 µL per well

volume. Plates were incubated for 1 h at 37 °C, then washed and secondary antibody added

at 1:2000 dilution in blocking buffer (100 µL per well) and incubated for 1 hr at 37 °C. After

incubation and washes, plates were developed using 50 µL per well SureBlue TMB (3,3', 5,5'-

tetramethylbenzidine) substrate and the reaction stopped after 5 min with 50 µL per well

stop solution (Insight Biotechnologies). The absorbance was read on a Versamax

Spectrophotometer at 450 nm (BioTek Industries). Statistical analyses were performed on

log-transformed data.

Antigen specific Isotype ELISA; MaxiSorp plates (Nunc) were coated with 50 μl of 2 μg/mL

or 50ul of 5 μg/mL ng/well SARS-CoV-2 FL-S protein overnight at 4 °C for detection of IgG

(250ng/well) or IgM and IgA (500ng/well), respectively, prior to washing in PBS/Tween

(0.05% v/v) and blocking with Blocker Casein in PBS (Thermo Fisher Scientific) for 1 h at

room temperature (RT). Standard positive serum (pool of mouse serum with high endpoint

titre against FL-S protein), individual mouse serum samples, negative and an internal control

(diluted in casein) were incubated for 2 hrs at room temperature for detection of specific

IgG or 1h at 37 °C for detection of specific IgM or IgA. Following washing, bound antibodies

were detected by addition of alkaline phosphatase (AP)-conjugated goat anti-mouse IgG

(Sigma-Aldrich) or anti-mouse IgM (Abcam) or anti-mouse IgA (SouthernBiotech) for 1h at

room temperature and addition of p-Nitrophenyl Phosphate, Disodium Salt substrate

(Sigma-Aldrich). An arbitrary number of ELISA units (EU) were assigned to the reference

pool and optical density values of each dilution were fitted to a 4-parameter logistic curve

using SOFTmax PRO software. ELISA units were calculated for each sample using the optical

density values of the sample and the parameters of the standard curve. IgM limit of

detection was defined a 2 ELISA Units, IgA limit of detection set as 6 ELISA Units. All data

was log-transformed for statistical analyses.

Antigen Specific IgG Subclass ELISAs; MaxiSorp plates (Nunc) were coated with 50 μL of 2

μg/mL per well of SARS-CoV-2 FL-S protein overnight at 4 °C prior to washing in PBS/Tween

(0.05% v/v) and blocking with Blocker Casein in PBS (Thermo Fisher Scientific) for 1 h at

room temperature (RT). For detection of IgG subclasses all serum samples were diluted to 1

total IgG EU and incubated at 37 °C for 1 h prior to detection with Alkaline Phosphatase

conjugated anti-mouse IgG subclass-specific secondary antibodies (Southern Biotech or

Abcam) incubated for 1 h at 37 °C . The results of the IgG subclass ELISA are presented using

optical density values.

Avidity ELISA; Anti-SARS-CoV-2 spike-specific total IgG antibody avidity was assessed by

sodium thiocyanate (NaSCN)-displacement ELISA. Nunc MaxiSorp ELISA plates (Thermo

Fisher Scientific) coated overnight at 4 °C with 2μg/well SARS-CoV-2 trimeric spike protein

diluted in PBS were washed with PBS/Tween (0.05% v/v) and blocked for 1 h with 100 μl per

well of Blocker Casein in PBS (Thermo Fisher Scientific) at 20 °C. Test samples and a positive

control serum pool were diluted in blocking buffer to 1 total IgG EU and incubated for 2 h at

20 °C. After washing, increasing concentrations of NaSCN (Sigma-Aldrich) diluted in PBS

were added and incubated for 15 min at 20 °C. Following another wash, bound antibodies

were detected by addition of AP-conjugated goat anti-mouse IgG (Sigma-Aldrich) for 1 h at

room temperature and addition of p-Nitrophenyl Phosphate, Disodium Salt substrate

(Sigma-Aldrich). For each sample, concentration of NaSCN required to reduce the OD405 to

50% of that without NaSCN (IC50) was interpolated from this function and reported as a

measure of avidity.

Antigen specific B cell staining; Spike, RBD and a decoy NANP$_9$C (repeat region from *P.*

falciparum CSP protein) tetramers were prepared in house by mixing biotinylated proteins

with streptavidin conjugated flurochromes (A488, A647 or r-PE) in a 4:1 molar ratio and

incubating on ice for 30 minutes. Splenocytes were stained with Spike-PE and RBD-A647 at a

final concentration of 0.04mM, whilst decoy tetramer, NANP$_9$C-A488 was used at a final

concentration of 0.4mM. Splenocytes were stained with Live-Dead Aqua and Fc block (anti-

CD16/32 mAb, Clone 93) prior to staining with the antibody cocktail containing NANP$_9$C-

Alexa488, GL7-PerCPCy5.5 (Clone GL7), CD138 BV421 (Clone 281-2), CD95-BV605 (Clone

SA367H8), CD4-BV650 (Clone GK1.5), CD279-BV711 (Clone 29F.1A12), CD19-BV780 (6D5),

RBD-A647, IgD-A700 (Clone 11-26c.2a), IgM-APCCy7 (Clone 11/41), Spike-PE, CD38-PECY5

(Clone 90), CD69 PeCy7 (Clone H1.2F3), CD45R-BUV395 (Clone RA3-6B2) and CD3-BUV496

(Clone 145-2C11), antibodies purchased from BioLegend, BD or Invitrogen. Antigen specific

B cells were identified by gating on LIVE/DEAD negative, doublet negative (FSC-H vs FSC-A),

size (FSC-H vs SSC), CD45RA$^+$, CD19$^+$ and NANP-A488$^-$, RBD-A647$^+$ and Spike-PE$^+$.

ELISpot and ICS staining; Spleen single cell suspension were prepared by passing cells

through 70μM cell strainers and ACK lysis prior to resuspension in complete media. For

analysis of IFNγ production by ELISpot, splenocytes were stimulated with two pools of S1

(pools 1 and 2) and two pools S2 (pools 3 and 4) peptides (final concentration of 2μg/mL) on

IPVH-membrane plates (Millipore) coated with 5μg/mL anti-mouse IFNγ (AN18). After 18-20

hours of stimulation at 37 °C, IFNγ spot forming cells (SFC) were detected by staining

membranes with anti-mouse IFNγ biotin (1mg/mL) (R46A2) followed by streptavidin-

Alkaline Phosphatase (1mg/mL) and development with AP conjugate substrate kit (BioRad,

UK).

For analysis of intracellular cytokine production, cells were stimulated at 37 °C for 6 hours

with 2μg/mL a pool of S1 (ELISpot pools 1 and 2) or S2 (ELISpot pools 3 and 4) peptides,

media or positive control cell stimulation cocktail (containing PMA-Ionomycin, BioLegend),

together with 1μg/mL Golgi-plug (BD) and 2μl/mL CD107a-Alexa647 (Clone 1D4B). Following

surface staining with CD3-A700 (Clone 17A2), CD4-BUV496 (Clone GK1.5), CD8-BUV496

(Clone 53-6.7), CD44-BV780 (Clone IM7), CD62L-BV711 (Clone MEL-14), CD69-PECy7 (Clone

H1.2F3) and CD127-BV650 (Clone A7R34) or CD127-APCCy7 (Clone A7R34), cells were fixed

with 4% paraformaldehyde and stained intracellularly with TNFa-A488 (Clone MP6-XT22),

IL2-PerCPCy5.5 (Clone JES6-5H4), IL4-BV605 (Clone 11B11), IL10-PE (Clone JES5-16E3) and

IFNγ-e450 (Clone XMG1.2) diluted in Perm-Wash buffer (BD). Sample acquisition was

performed on a Fortessa (BD) and data analyzed in FlowJo V10 (TreeStar). An acquisition

threshold was set at a minimum of 5000 events in the live CD3$^+$ gate. Antigen specific T cells

were identified by gating on LIVE/DEAD negative, doublet negative (FSC-H vs FSC-A), size

(FSC-H vs SSC), CD3$^+$, CD4$^+$ or CD8$^+$ cells and cytokine positive. Cytokine positive responses

are presented after subtraction of the background response detected in the corresponding

media stimulated control sample for each mouse and summing together the response

detected to each pool of peptides. T effector (Teff) cells were defined as CD62L$^-$ CD127$^-$, T

central memory cells defined as CD62L$^-$ CD127$^+$ and T central memory (Tcm) cells defined as

CD62L^{hi} CD127^{hi} (Figure S3B). All graphs and statistical analysis were performed using Prism

v9 (Graphpad), total cell counts were log transformed prior to statistical analysis.

References

1   Tregoning, J. S. *et al.* Vaccines for COVID-19. *Clin Exp Immunol* **202**, 162-192,
314   doi:10.1111/cei.13517 (2020).
315   2   Blakney, A. K. *et al.* Big Is Beautiful: Enhanced saRNA Delivery and Immunogenicity
316   by a Higher Molecular Weight, Bioreducible, Cationic Polymer. *ACS Nano* **14**, 5711-
317   5727, doi:10.1021/acsnano.0c00326 (2020).
318   3   McKay, P. F. *et al.* Self-amplifying RNA SARS-CoV-2 lipid nanoparticle vaccine
319   candidate induces high neutralizing antibody titers in mice. *Nature communications*
320   **11**, 3523, doi:10.1038/s41467-020-17409-9 (2020).
321   4   Ewer, K. *et al.* Chimpanzee adenoviral vectors as vaccines for outbreak pathogens.
322   *Hum Vaccin Immunother* **13**, 3020-3032, doi:10.1080/21645515.2017.1383575
323   (2017).
324   5   Antrobus, R. D. *et al.* Clinical assessment of a novel recombinant simian adenovirus
325   ChAdOx1 as a vectored vaccine expressing conserved Influenza A antigens. *Mol Ther*
326   **22**, 668-674, doi:10.1038/mt.2013.284 (2014).
327   6   Venkatraman, N. *et al.* Safety and Immunogenicity of a Heterologous Prime-Boost
328   Ebola Virus Vaccine Regimen in Healthy Adults in the United Kingdom and Senegal. *J
329   Infect Dis* **219**, 1187-1197, doi:10.1093/infdis/jiy639 (2019).

7   Folegatti, P. M. *et al.* Safety and immunogenicity of a candidate Middle East
respiratory syndrome coronavirus viral-vectored vaccine: a dose-escalation, open-
label, non-randomised, uncontrolled, phase 1 trial. *Lancet Infect Dis* **20**, 816-826,
doi:10.1016/S1473-3099(20)30160-2 (2020).
8   Voysey, M. *et al.* Safety and efficacy of the ChAdOx1 nCoV-19 vaccine (AZD1222)
against SARS-CoV-2: an interim analysis of four randomised controlled trials in Brazil,
South Africa, and the UK. *Lancet* **397**, 99-111, doi:10.1016/S0140-6736(20)32661-1
(2021).
9   Ewer, K. J. *et al.* T cell and antibody responses induced by a single dose of ChAdOx1
nCoV-19 (AZD1222) vaccine in a phase 1/2 clinical trial. *Nat Med*,
doi:10.1038/s41591-020-01194-5 (2020).
10 Barrett, J. R. *et al.* Phase 1/2 trial of SARS-CoV-2 vaccine ChAdOx1 nCoV-19 with a
booster dose induces multifunctional antibody responses. *Nat Med*,
doi:10.1038/s41591-020-01179-4 (2020).
11 Graham, S. P. *et al.* Evaluation of the immunogenicity of prime-boost vaccination
with the replication-deficient viral vectored COVID-19 vaccine candidate ChAdOx1
nCoV-19. *NPJ Vaccines* **5**, 69, doi:10.1038/s41541-020-00221-3 (2020).
12 van Doremalen, N. *et al.* ChAdOx1 nCoV-19 vaccine prevents SARS-CoV-2 pneumonia
in rhesus macaques. *Nature* **586**, 578-582, doi:10.1038/s41586-020-2608-y (2020).
13 Ramasamy, M. N. *et al.* Safety and immunogenicity of ChAdOx1 nCoV-19 vaccine
administered in a prime-boost regimen in young and old adults (COV002): a single-
blind, randomised, controlled, phase 2/3 trial. *Lancet* **396**, 1979-1993,
doi:10.1016/S0140-6736(20)32466-1 (2021).
14 Wang, S. F. *et al.* Antibody-dependent SARS coronavirus infection is mediated by
antibodies against spike proteins. *Biochem Biophys Res Commun* **451**, 208-214,
doi:10.1016/j.bbrc.2014.07.090 (2014).
15 Tseng, C. T. *et al.* Immunization with SARS coronavirus vaccines leads to pulmonary
immunopathology on challenge with the SARS virus. *PLoS One* **7**, e35421,
doi:10.1371/journal.pone.0035421 (2012).
16 Lee, W. S., Wheatley, A. K., Kent, S. J. & DeKosky, B. J. Antibody-dependent
enhancement and SARS-CoV-2 vaccines and therapies. *Nat Microbiol* **5**, 1185-1191,
doi:10.1038/s41564-020-00789-5 (2020).
17 Mercado, N. B. *et al.* Single-shot Ad26 vaccine protects against SARS-CoV-2 in rhesus
macaques. *Nature* **586**, 583-588, doi:10.1038/s41586-020-2607-z (2020).
18 Rydyznski Moderbacher, C. *et al.* Antigen-Specific Adaptive Immunity to SARS-CoV-2
in Acute COVID-19 and Associations with Age and Disease Severity. *Cell* **183**, 996-
1012 e1019, doi:10.1016/j.cell.2020.09.038 (2020).
367

Acknowledgments: The authors would like to thank D. Pulido for provision of spike and RBD

proteins, BMS staff for animal husbandry and management and A. Worth, J.Furze, M.

Mykhaylyk and R. Evans for facilities support.

Funding: This report is independent research funded by the National Institute for Health

Research (UKRI Grant Ref: MC_PC_19055, NIHR Ref: COV19 OxfordVacc-01). The views

expressed in this publication are those of the author(s) and not necessarily those of the

National Institute for Health Research or the Department of Health and Social Care.

Author Contributions: AJS, MU, SB-J, CB, KS, KM & PMK performed experiments. AJS, MU,

HS, CG, EA & AT performed animal procedures and/or sample processing. AJS, SBJ, KH &

PMK, analyzed data. AJS, TL, PMK, RS & SG designed the study. AJS, SBJ, TB wrote the

manuscript. All authors reviewed the final version of the manuscript.

Competing interests: SCG is co-founder and board member of Vaccitech (collaborators in

the early development of this vaccine candidate) and named as an inventor on a patent

covering use of ChAdOx1-vectored vaccines and a patent application covering this SARS-

CoV-2 vaccine. TL is named as an inventor on a patent application covering this SARS-CoV-2

vaccine and was consultant to Vaccitech. PMK and RJS are co-founders and RJS is a board

member of VaxEquity and VacEquity and are named inventors on a patent application

covering the SARS-CoV-2 saRNA vaccine candidate.

Figure legends

Fig 1: Antibody responses following ChAd and saRNA vaccination

Antibody responses were measured in the serum of mice collected 3 weeks after the final

immunisation. Graphs show SARS CoV-2 spike-specific IgG (**A.**), IgM and IgA (**B.**) and IgG

avidity (**C.**) measured by ELISA, and SARS-CoV-2 pseudotyped virus neutralisation (IC50) (**D.**).

Individual mice are represented by a single data point, bars represent the median response

in each group (CD1 n=8; BALB/c n=6). Data in each graph was analysed with a Kruskal-Wallis

and post-hoc positive test to compare differences between vaccination groups, p values

indicate significant differences between groups.

Fig 2: SARS-CoV-2 spike-specific IgG subclasses following ChAd and saRNA vaccination

For detection of IgG subclasses, each sample was diluted to 1 IgG ELISA Unit. Graphs show optical density measured against each IgG subclass where individual data points were expressed as an OD and shown here as scatter dot plots with bars showing the median (**A.**), followed by the heatmap summary representation with median response in each group to each IgG subclass (**B.**). Individual mice are represented by a single data point, bars represent the median response in each group (CD1 n=8; BALB/c n=6). Data in each graph was analysed with a Kruskal-Wallis and post-hoc positive test to compare differences between vaccination groups, p-values indicate significant differences between groups.

Fig 3: Breadth of T cell response measured by ELISpot

Graphs represent the total spike specific IFNγ response (sum of peptide pools) measured in outbred CD1 (**A.**) or inbred BALB/c (**B.**). Pie charts represent the response to each peptide pool as a proportion of total response. Data points represent individual mice and bars represent the median response in each group. Data in each graph was analysed with a Friedman test and post-hoc Dunn's multiple comparison to compare between vaccination regimens, p values are indicated on the graph.

Fig 4: Phenotype of the T cells response following vaccination

Splenocytes were stimulated for 6 hours with pools of SARS-CoV-2 peptides prior to staining

for effector and memory T cells markers and intracellular cytokines. **A.)** Graphs show the

frequency of the spike-specific CD4+ T cells responses (left) in CD1 (top panel) and BALB/c

mice (bottom panel), heatmaps (middle) show the proportion of the response producing

each cytokine and total number of antigen specific cells of a T effector (Teff), T effector

memory (Tem) or T central memory (Tcm) phenotype (right). **B.**) Graphs show the

frequency of the spike-specific CD8$^+$ T cells responses (left) in CD1 (top panel; n=8) and

BALB/c mice (bottom panel; n=6), heatmaps (middle) show the proportion of the response

producing each cytokine and total number of antigen specific cells of a T effector (Teff), T

effector memory (Tem) or T central memory (Tcm) phenotype (right). Data points indicate

individual mice, bars represent the median response in each group. Total numbers of each

population are displayed in Fig S3C. Data in each graph was analysed with a two-way anova

and post-hoc Tukey's multiple comparison test to compare between vaccination regimens, p

values are indicated on the graph.

443
444

Fig S1: Comparison of assays measuring antibody responses after vaccination

SARS-CoV-2 spike-specific IgG responses were measured in a standardised ELISA (**A.**) and

compared to IgG responses measured as a concentration (**B.**)

C.) Graphs show relationship between IgG, IgM and IgA responses to SARS-CoV-2

pseudoneutralisation IC50 values presented by linear

450   regression on log-transformed values. Data points represent individual animals from all

451   groups together, r^2 and p values are indicated on each graph.

Fig S2: Antigen specific B cell responses

SARS-CoV-2 spike-specific B cells responses were measured in the spleen of BALB/c mice 3

weeks after the final immunisation. **A.**) Antigen specific cells were identified by staining with

spike-PE and RBD-A647 tetramers. Germinal center (GC) B cells were identified as GL7+,

CD95+, switched B cells identified as IgD- and IgM- antigen-specific B cells **B.**) Graphs show

the total number of antigen-specific B cells, antigen-specific GC B cells and antigen-specific

461   switched B cells, data points are representative of individual mice, median responses per

462   group indicated by bars.

Fig S3: T cell responses measured by ELISpot and ICS

A.) Graphs show the IFNγ SFC detected to each pool of peptides by ELISpot. Each data point

represents an individual mouse, bars represent the median response per group.

B.) Plots show the gating strategy used to identify antigen specific T cell responses.

C.) Graphs show the total number of Teff, Tem and Tcm responses detected in the spleens

of mice. Each point indicates an individual mouse, bars represent the median response per

group.