Resequencing 250 soybean accessions: new insights into genes associated with agronomic traits and genetic networks

Chunming Yang[1,a#], Jun Yan[2,b#], Shuqin Jiang[2,c], Xia Li[3,d], Haowei Min[4,e*], Xiangfeng Wang[2,f*], Dongyun Hao[1,g*]

[1]*Key Laboratory for Agricultural Biotechnology of Jilin Provincial, Institute of Agricultural Biotechnology, Jilin Academy of Agricultural Sciences (JAAS), Jilin 130033, China*

[2]*Department of Crop Genomics and Bioinformatics, College of Agronomy and Biotechnology, China Agricultural University, Beijing 100094, China*

[3]*Key Laboratory of Molecular Cytogenetics and Genetic Breeding of Heilongjiang Province, College of Life Science and Technology, Harbin Normal University, Harbin 150025, Heilongjiang, China*

[4]*BioTrust Technology. Inc., Beijing 100094, China*

\# Equal contribution

* Corresponding authors

E-mail:

dyhao@cjaas.com (Hao D);

xwang@cau.edu.cn (Wang X);

biotrust_st@163.com (Min H);

a ORCID: 0000-0002-3806-6457

b ORCID: 0000-0000-0000-0000

c ORCID: 0000-0000-0000-0000

d ORCID: 0000-0000-0000-0000

e ORCID: 0000-0000-0000-0000

30   f ORCID: 0000-0002-6406-5597

31   f ORCID: 0000-0002-6406-5597

40   **Running title:** Chunming Yang *et al.* / Soybean resequence

Abstract

Limited knowledge on genomic diversity and the functional genes associated with soybean variety traits has resulted in slow breeding progress. We sequenced the genome of 250 soybean landraces and cultivars from China, America and Europe, and investigated their population structure, genetic diversity and architecture and selective sweep regions of accessions. We identified five novel agronomically important genes and studied the effects of functional mutations in respective genes. We found candidate genes *GSTT1*, *GL3* and *GSTL3* associated with isoflavone content, *CKX3* associated with yield traits, and *CYP85A2* associated with both architecture and yield traits. Our phenotype-gene network analysis revealed that hub nodes play a role in complex phenotypic associations. In this work, we describe novel agronomic trait associated genes and a complex genetic network, providing a valuable resource for future soybean molecular breeding.

Introduction

Soybean *Glycine max* [L.] Merr. is one of the most important crops worldwide of vegetable oil and proteins source for human and livestock feed etc. Soybean originated in China and its wild species (*G. soja* Sieb. & Zucc.) were domesticated in approximately 3,000 B.C. before introduced to Korea and Japan about 3,000 years later. It was brought to Europe and North America in the 18th century, and extensively cultivated on a global scale since the 19th century[1].

With the rapid development of modern molecular biology and the high-throughput sequencing technologies, whole-genome resequencing and genome wide association studies (GWAS) have become common methods used to study population genetic diversity and locating phenotypic related quantitative trait loci (QTL) or genes. This has improved our knowledge extensively in crop genomes and selective breeding. In recent years, for example, there are increasing number of reports on the domestication and improvement of soybean at the genome-wide level.

This includes genes and genetic networks related to soybean agronomic traits and

functions[2-4]. However, due to the diversity of soybean varieties and their complex

genetic background, our knowledge of the soybean genome and functional genes is

still limited in comparison to rice and maize. A greater number of soybean varieties

need further exploration at the genomic level, particularly in relation to molecular

traits associated with edible quality, soybean "ideotype" and the underlying genetic

network of high-yielding varieties.

In this study, we collected 250 soybean varieties from the core Northeast China

soybean germplasm pool, which consisted of 134 accessions of landrace and cultivar

from Northeast and Northwest China, as well as 116 accessions from European and

North American cultivars. The genomes of the most accessions are not sequenced

previously. We performed the high-depth whole-genome resequencing and

comprehensive analyses of this 250 soybean population. The generated dataset

revealed valuable new information on soybean genome structure, novel genes

associated with important agronomic traits and the genetic networks. These genetic

resources provide unique references into molecular breeding and evolution study in

soybean.

Results

Genome resequencing and variation calling

High-depth whole genome resequencing was performed on 250 soybean accessions,

including 51 landraces and 83 cultivars originating from provinces in Northeast China

(i.e. Heilongjiang, Jilin, Liaoning and Northeast Inner Mongolia) and Northwest

China (i.e. Xinjiang, Ningxia and Gansu), as well as 116 cultivars originating from

Europe and North America **(Figure 1a, Supplementary Table 3).** In total, we

obtained approximately 10G of pair-end reads and 3T bases. The maximum

sequencing depth of a single accession was 22.5x, with the average depth at 11x. After

filtering out the raw sequencing data (see methods), the remaining high-quality

cleaned data were compared with the soybean reference genome *G. max v2.0[5].* The

effective mapping rates ranged from 74.8% to 87.6%, while the genome coverage

ranged from 94.8% to 97.0% (**Supplementary Table 1**). The high mapping rates and

coverage guarantee that the sequenced data is reliable and of high quality.

Through standard variation detection, genotype filtering and imputation steps

(see methods), we detected in total 6,333,721 single nucleotide polymorphisms (SNPs)

and 2,565,797 insertion & deletions (Indels). This includes 244,360 SNPs and 62,714

Indels located in the exon regions. The ratio of non-synonymous SNP to synonymous

SNP substitution was 1.37. There are 4,311,814 SNPs with a minor allele frequency

(MAF) larger than 0.05 (**Supplementary Table 2 & Supplementary Figure 1**). In

summary, we achieved over 6M high-density and high-quality genotype data from

250 soybean accessions with a density of one SNP per 15 bases.

Population structure of soybean landraces and cultivars

Using the 6M SNP genotype dataset, we constructed a phylogenetic tree using the

neighbour joining (NJ) method. This resulted in the classification of the 250 soybean

accessions into four groups (**Figure 1b**). Among them, *Group 1* included 65 Chinese

varieties, four European and six American cultivars, whereas *Group 2* contained 56

Chinese varieties, one European and 13 American varieties. In *Group 3*, there were 21

European, two Chinese and six American cultivars, while 65 North American

cultivars and 11 Chinese varieties were clustered within *Group 4* (**Figure 1c**).

Principal Component Analysis (PCA) results were consistent with the phylogenetic

tree results. Three groups, *Group 1*, *3*, and *4* radiated away from *Group 2* within the

rectangular coordinate system projected using eigenvector 1 and eigenvector 2 data on

X and Y axes, respectively. Concurrently, the distribution of varieties in the four

groups had continuity, indicating varieties located in different groups also have

genetic similarities (**Figure 1e**). A Bayesian clustering algorithm based on a mixed

model was used to estimate the proportion of ancestors in each accession. That is,

when $K = 2$, the main ancestor component (yellow) of *Group 4* was split, indicating

that *Group 4* has the highest level of selection. When $K = 3$, the main ancestor

component (blue) of *Group 3* was split, indicating that *Group 3* has the second level

of selection. However, when $K = 4$ and $K = 5$, *Group 1 and Group 2* exhibit complex

differentiated mixed ancestor components, indicating a higher genetic diversity and

lower selection level in *Groups 1* and *2* (**Supplementary Table 3, Figure 1d**).

These results indicate that group classification of the 250 soybean accessions is

closely related to their geographical distribution. That is, varieties with similar

geographical distribution have similar genetic backgrounds. Generally speaking, the

group classification was also related to the level of domestication. Landraces have a

lower level of domestication, while cultivars have higher levels of domestication.

Varieties with similar domestication levels tend to have a higher similarity in genetic

backgrounds. However, there are still differences in geographical distribution and

domestication level among breeds with similar genetic backgrounds, indicating that

gene exchange may have occurred between accessions of different groups. This

observation reflected the complexity of soybean domestication history.

Genetic diversity and selective sweep analysis

Linkage disequilibrium (LD) analysis showed that the overall LD decay distance was

more than 100 kb, and the LD decay distance of the landraces was smaller than that of

the cultivars (**Figure 2A**). Further LD decay analysis of the four groups showed that

the LD decay distance of *Group 1* was the smallest, followed by *Groups 2* and *3*,

while *Group 4* had the largest LD decay distance (**Figure 2B**). In addition, the LD

levels varied for different chromosomes or different regions across one chromosome.

Identical by state (IBS) analysis can reflect the degree of relatedness among

individuals by calculating the consistency of all genetic markers. The IBS values of

all comparisons in each group were calculated, and it was found that the average IBS

values of landraces were less than that of cultivars (**Figure 2C**). The IBS values of

Groups 1-4 followed the same trends as that of LD decay distance. In particular, the

IBS values of *Group 1* were the lowest and the IBS values of *Group 4* were the

highest of all groups (**Figure 2D**). $\theta\pi$ values can reflect the genetic diversity within a

population by calculating the number of different sites between any two sequences or

individuals within a population. *Fst* is a calculation used to measure the

differentiation and genetic distance between two populations. $\theta\pi$ values were calculated for landraces, cultivars, all accessions, and *Groups 1-4*. *Fst* values were calculated between landraces and cultivars, and between each comparison of the four Groups. Results show that a population with a higher level of LD decay distance or higher IBS values correlate with a smaller $\theta\pi$ (**Figure 2E, F**). This pattern is opposite to that of the LD decay and IBS values. The lowest *Fst* value was for *Group 1* versus *Group 2*, while the highest value was for *Group 3* versus *Group 4*. We also observed that the *Fst* value of *Group 2* versus *Group 3* was higher than that of *Group 1* versus *Group 3* (**Figure 2G**). The *Fst* value of *Group 2* versus *Group 4* was smaller than that of *Group 1* versus *Group 4*. In addition, the results of allele frequency distribution (AFD) analysis, as an alternative population similarity measurement, were consistent with the *Fst* results (**Supplementary Figure 2**). The population diversity analysis results, when combined with population structure and geographical distribution information, infer that the European and American soybean varieties may have originated from different Chinese ancestors before undergoing independent selection. Results indicate that European cultivars and the Chinese landrace group (*Group 1*) in our study have a more recent common ancestor, while North American cultivars and the Chinese cultivar group (*Group 2*) have a more recent common ancestor.

Tajima' D (based on a neutral test), $\theta\pi$ (based on genetic diversity within a population), and *Fst* (based on genetic diversity between two populations), have provided us with highly effective tools that screen selective sweep signals across a genome[6]. We combined methods in pairs for mining potential selective sweep regions in the soybean genome that may have underwent artificial selection. One pair was *Tajima' D* combined with $\theta\pi$ for the whole population. Another pair was *Fst* combined with $\theta\pi$ ratios between two subpopulations, landrace and cultivar. We used a sliding window method to calculate the values of *Tajima' D, $\theta\pi$, and Fst* in each window across the whole genome, and selected the top 5% most significant windows as potential selective sweep regions (**Supplementary Figure 3A, B**). A total of 148 and 222 potential selective sweep regions were screened by the two methods, and they covered 36.09 Mb and 88.15 Mb genome regions, respectively (**Supplementary**

Table 4). These potential selective sweep regions covered 9,128 genes, accounting for approximately one sixth of all soybean genes. A total of 1,876 genes were screened by both methods (**Supplementary Figure 3C**). A runs of homozygosity (ROH) region is a continuous homozygous chromosome region in a genome, which may relate to domestication or artificial selection[7]. Through ROH analysis, 71 ROH regions larger than 300 kb were obtained from all 250 accessions, with a total length of 27.84 Mb. The longest ROH regions up to 911 kb were located at the beginning of chromosome 10 (**Supplementary Table 5**). There were 3,397 genes located in these ROH regions, 924 of which were also located in the potential selected sweep region (**Supplementary Figure 3D**).

Phenotype-related loci and genes identified using GWAS analysis

We measured 50 agronomic traits in 250 soybean accessions from three geographic locations for three years, and then integrated them using best linear unbiased prediction (BLUP). The 50 traits included traits related to architecture (15), colour (5), isoflavone (1), oil (4), protein (18) and yield (7) and were classified into six categories (**Supplementary Table 6**). We calculated Pearson correlation coefficients for traits so as to compare within and between categories, and found that traits in the same categories were more strongly correlated than traits in different categories. For example, there were strong positive or negative correlations between almost all protein-related traits, oil-related traits and yield-related traits. Linoleic acid content was positively correlated with linolenic acid content, but negatively correlated with oleic acid content. Stem intension was negatively correlated with lodging (**Supplementary Figure 4**). Some traits were evenly distributed, while others were ranked (**Supplementary Figure 5-54**).

Using 4,311,814 SNPs with a MAF > 0.05 as an input, we performed GWAS analysis using the mixed linear model (MLM) method for 50 agronomic traits. For each trait, we used a clump based method[8] and defined a significant associated loci (SAL) at a chromosome region with a substantial amount of SNPs associated with the trait. A total of 203 SALs were detected in 43 traits (**Supplementary Figure 5-54,**

Supplementary Table 7). Since each SAL may contain dozens of genes, we used a

functional mutation based haplotype test method for further mining of the most

reliable candidate trait associated genes[9]. In particular, we considered only the

non-synonymous SNPs, frameshift Indels, mutations within a gene that happened on a

start or stop codon, splice site or transcription start sites as effective functional

mutations. We used these mutations to classify a gene into different haplotypes, and

subsequently tested the phenotypic differences of the accessions belonging to each

haplotype. A gene with significant phenotypic differences was defined as significant

associated gene (SAG), and 3,165 SAGs were screened in 43 traits. These SAGs

include some QTL or genes that have been previously identified, such as: the flower

colour   related   *chr13:16551728-19506795*;   pubescence   colour   related

chr6:16930159-19168772;   seed   coat   lustre   related   *chr15:8910798-10281804*;

palmitic acid content related *chr5:879095-1682551[4]*; isoflavone content related

chr5:38880530-39142565[10]; plant height related *Dt1[4]*; and oil content related

FAD2 and *SAT1[11]*, among others. They also contain genes that we have identified

for the first time in soybean, such as: the isoflavone content related *GL3* and *GSTL3*;

the yield traits related *CKX3*; and the architecture and yield traits related *CYP85A2*.

Novel genes related to isoflavone content

Isoflavone content is an important quality-related trait in soybean, but its molecular

mechanism is still unclear. Here we identified four SALs related to isoflavone content,

namely   *chr3:38590023-38728718*,   *chr5:3888053-39142565*,

chr13:18342836-18541809,   and   *chr5:24726091-24852447*.   Only   one   SAL

chr5:24726091-24852447 overlaps with a previously reported QTL that contains a

GST (Glutathione S-transferase) gene *GSTT1*[10]. All other SALs are newly identified.

There are 48 genes located within these SALs (**Supplementary Table 8**), and three

genes (*GSTT1, GL3* and *GSTL3)* may be related to isoflavone content (**Figure 3A, B**).

There are two functional mutation sites at *c5s38936266* and *c5s38940717*, forming

two haplotypes for *GSTT1a* and *GSTT1b*, respectively. For each *GSTT1* gene, soybean

accessions with a different haplotype have significantly different isoflavone contents.

Since *GSTT1a* and *GSTT1b* are approximately only 1 kb apart from each other in the

same genome region, we considered the two genes to be one in further analysis. Three

haplotypes were formed by two functional mutation sites when the two *GSTT1* genes

were analysed as one. *Haplotype 1* versus *Haplotype 2*, as well as *Haplotype 2* versus

Haplotype 3 showed significant differences in isoflavone content, while *Haplotype 1*

versus *Haplotype 3* showed no significant differences (analysed using Tukey's test).

This suggests that *GSTT1a* is associated with isoflavone content due to its linkage

with *GSTT1b*. However, *c5s38936266* did not contribute to the isoflavone content

difference. Thus, only *c5s38940717* on *GSTT1b* was associated with isoflavone

content (**Figure 3C**).

We found that two functional mutations, *c5s39035509* and *c5s39036346*

producing two haplotypes in *GL3*, were associated with different isoflavone contents

in soybean accessions (**Figure 3D**). We also identified another *GST* gene, *GSTL3,*

which was located in chromosome 13. Two functional mutations produced three

haplotypes, and significant associations between the different haplotypes and

isoflavone content was detected for each comparison (**Figure 3E**). Based on the

above results, we drew a schematic diagram of the roles of candidate genes according

to their biological functions where we indicate that *GL3* regulates isoflavone synthesis,

whilst *GSTT1* and *GSTL3* participate in isoflavone transport (**Figure 3F**).

Yield related traits and the artificial selection of *CKX3*

Four yield related traits (pod number per plant, seed number per plant, one hundred

seed weight and seed size) have a common SAL located at the ~4.0 Mb to ~4.2 Mb

region of chromosome 17 (**Figure 4A**). Further analysis revealed that this SAL

contains two tandem repeat *CKX* (cytokinin oxidase/dehydrogenase) genes named

CKX3 and *CKX4*, approximately 15 kb apart from each other (**Figure 4F**).

We further analysed the relationship between functional mutations in *CKX3* and

CKX4. There were three non-synonymous SNPs on *CKX3* and two non-synonymous

SNPs on *CKX4*. As these two genes are only approximately 3 kb apart from each

other in the same genome region, we analysed the two genes separately as well as

combined as one in relation to their association with haplotypes and traits (**Figure**

4B-E). Results showed that the functional mutations can both form three haplotypes

for *CKX3* and *CKX4 separately*, and four haplotypes for *CKX3+4* combined. For all

comparisons in all traits, *Haplotype 1* always showed significant differences

compared with the other haplotypes. The relationship between different haplotypes in

terms of pod or seed number per plant showed a consistent trend, while that of one

hundred seed weight or seed size showed a consistent but opposite trend. There was a

phenotypic correlation between pod number per plant and seed number per plant

(0.92), and between one hundred seed weight and seed size (0.67) (**Figure 4F**).

Furthermore, we observed that *CKX3* and *CKX4* were located in different strands of

the same chromosome, suggesting they are more likely to have independent functions.

However, we did not detect expression of *CKX4* in the subsequent qPCR validation.

Thus, only *CKX3* is regarded as a real candidate gene, while the role of *CKX4* need

further study.

When we compared the soybean accession information of each haplotype for the

four yield-related traits, we observed that most accessions with the *Haplotype 1*

genotype had dominant traits (lower pod or seed numbers and larger seeds and seed

weights), and were more associated with cultivars. The other haplotypes were mainly

landrace-specific haplotypes, and these accessions all belong to *Group 1*. We found

that *CKX3* was also located on a strong selective sweep region. This indicates that the

functional mutation sites in *CKX3* experienced strong directed artificial selection,

resulting in genotype differences and affecting yield related traits. Furthermore, we

compared all SAGs and selective sweep regions for all traits, and found that

approximately 12% of genes are located in the selected sweep regions, which have

experienced artificial selection (**Supplementary Table 9**). It is interesting to note that

of all the SAGs located in selective sweep regions, about 55% are related to yield

traits, 36% related to protein traits, and less than 10% are related to other traits

(**Figure 4G, H**).

CYP85A2 is associated with architecture and yield traits

There is one SAL on chromosome 18 which is associated with six traits including

plant height, main stem number, stem strength, lodging, podding habit, and seed

weight per plant. Interestingly, these traits include both architecture and yield related

traits. A cytochrome P450 family gene named *CYP85A2* is located within a 4.37 kb

region of this SAL (**Figure 5A, B**). The association of the *CPY85A2* gene with

architecture and yield traits in soybean is a novel finding. We also observed that a

non-synonymous mutation site *c18s55526062* is involved in producing two

haplotypes. The haplotype with a *CC* genotype has a dwarf plant height, a low main

stem node number, a high stem strength and a low lodging rate. When plants produced

mostly limited or semi-limited pods, their seed weight per plant was also found to

increase (**Figure 5C**). Phenotypically, plant height was positively correlated with

main stem node number (0.95), while stem strength and lodging were negatively

correlated (-0.81), showing a trend consistent with the genotype (**Figure 5D**). The *CC*

genotype of *c18s55526062* is a dominant genotype, which is useful when designing

an ideal plant type and increasing soybean yield.

Different phenotypes coupled through hub gene modules form a complex

phenotype-gene network

Based on in-depth exploration of the GWAS results, we observed that one trait is

associated with multiple genes and vice versa. At the same time, due to widespread

protein-level interactions between genes, a complex network was also found between

various phenotypes and genes. In order to explore this further, we used a functional

mutation-based haplotype test to screen SAGs in all SALs for all traits. We then

constructed a phenotype-gene network which included 34 traits and 853 SAGs

(**Figure 6**). At the trait level, they were divided into six categories, namely

architecture, colour, oil, isoflavone, protein, and yield. At the gene level, besides the

six categories, there emerged a mixed category with which genes associated more

than with any one trait category. We found that traits in the same category were

closely linked within the entire network. However, some trait categories were also

linked with each other, such as yield, oil, protein and colour, and they were all closely

linked to architecture through common SAGs. This suggests that there are subtle

relationships between architecture and other trait categories. In this genetic network,

six trait categories were linked through 15 hub nodes containing a total of 367 genes

(**Supplementary Table 10**). The largest hub was *HA1* (short form for *Hub*

Architecture 1). The genes in this hub were only associated with two or more

architecture-related traits. Unlike *HA1*, the genes in the *HA2* node were associated

with two or more architecture-related traits, but also had protein interactions with

other genes. Multiple yield traits were associated with hub *HY1*, containing *CKX3*,

while hub *HM4*, which contains *CYP85A2*, was connected with architecture and yield

traits.

Discussion

In this study, we deeply sequenced 250 representative landrace and cultivar soybean

accessions. Through population genetics and GWAS analyses, the genetic structure of

European soybean varieties was analysed for the first time. Novel candidate genes

related to seed isoflavone content, yield and architecture traits were identified.

Moreover, we constructed a soybean phenotype-gene interaction network, and found

evidence of the improvement of soybean yield related traits at molecular level.

A total of ~3T bases and 6M SNPs were obtained, the maximum sequencing

depth of a single accession was 22.5x, with the average depth at 11x (higher than

previous soybean resequencing studies[2-4]. Eighty-four percent of the accessions

with their genome were sequenced for the first time, which provides new data for

soybean genome research. Previous soybean research mainly focused on varieties

from Asia and North America, but not Europe[3, 4]. This study completed the

resequencing of 26 European accessions and, for the first time, outlines a breeding

history of European soybean. It was found that European soybean cultivars had higher

genetic diversities and lower breeding levels compared to North American cultivars.

Both European and American soybean cultivars may have been introduced from

different ancestors in China. This theory is based on the following findings: there is a

small population difference between European varieties and Chinese landraces, and

between American varieties and Chinese cultivars, whilst there is a large population

difference between American and European varieties (**Figure 2G, Supplementary**

Figure 2). Our findings are consistent with the current hypothesis that soybean

originated in China, and they show that ancestral components from the area of origin

are the most complex. This study showed that the heterozygosity rates of most

accessions are less than 0.2, except four accessions with a higher heterozygosity

which may be caused by their complex ancestral compositions (**Supplementary**

Table S3). Further combination analysis of the selective sweep and GWAS revealed

that artificial selection of soybean at the phenotypic level is consistent with the

genome level. Genomic regions associated with yield and quality traits are more

likely to experience artificial selection. This may be a reflection of yield- and

quality-directed artificial selection of soybean breeding at the genetic level.

Furthermore, evidence of functional mutations under artificial selection for a

candidate gene *CKX3* related to multiple yield traits were identified. The results of

this study provide valuable information for marker-assisted selection, which is vital in

the improvement of soybean breeding.

Isoflavone is a secondary metabolite produced via phenylpropane metabolic

pathways in higher plants. Isoflavone is associated with plant stress resistance,

defence against microbial and insect infection, promotion of rhizobium chemotaxis,

and the development of rhizome and nitrogen fixation in plants. It also provides health

benefits to human, such as in reducing the incidence of cancer and cardiovascular

diseases, and regulating the immune response[12]. Therefore, increasing the seed

isoflavone content of soybean can improve its nutritional and health benefits.

However, few genome-wide studies have investigated the molecular mechanism of

soybean's isoflavone content. Isoflavone is synthesized in the cytoplasm, but due to

cell cytotoxicity, it cannot accumulate in the cytoplasm and must be continuously

transported to vacuoles for storage. Therefore, isoflavone content mainly depends on

two factors: synthesis efficiency and transport efficiency[13]. The transcript factor

GL3 is a *bHLH* gene family member which can form the MYB-bHLH-WD40 (MBW)

complex with two other transcription factors (*MYB* and *WD40*) to jointly regulate the synthesis of flavonoids and anthocyanin in plants[14]. *GST* can bind with glutathione (GSH) to form an ABC transporter to transport and catalyse the entry of flavonoids into vacuoles for accumulation[13]. In this study, we identified four novel genes that may be associated with isoflavone content. These genes include transcription factors *GL3*, which participate in the regulation of multi-enzyme systems from phenylpropanoid to isoflavone biosynthesis pathways, and two *GST* genes, *GSTT1* and *GSTL3*, which facilitate the transporting of isoflavone from the cytoplasm to vacuoles (**Figure 3F**). In addition, we observed many other genes in the SALs, such as cation/H+ exchanger (*CHX20*), pyrophosphorylase 4 (*PPa4*), an actin depolymerizing factor 7 (*ADF7*), a mitochondrial substrate carrier family protein, a myosin heavy chain-related protein, an *ATP* synthase alpha/beta family protein, and a protein kinase superfamily protein, among others (**Supplementary Table 8**) are all related to isoflavone transport. This over-representation of transport-related genes further suggests that the accumulation of soybean isoflavone is related to its transport to the vacuole. In conclusion, soybean isoflavone content is not merely determined by one or several genes or loci, but by a multiple gene system involved in synthesis, regulation, transport, and storage.

We observed that other novel candidate genes, such as *CKX3*, is associated with multiple yield traits. We also observed, for the first time, that *CYP85A2* is associated with multiple architecture and yield traits in soybean. It is well-known that cytokinin promotes cell division and plant growth, and *CKX* is one of the key enzymes in cytokinin metabolism. A functional variation in the *CKX* gene may affect the cytokinin metabolism, thus affecting grain yield and related traits. A number of studies on *Arabidopsis thaliana*, rice and other crops have shown that mutation or reduced expression levels of *CKX* family genes are related to a decrease in seed setting rate and an increase in seed weight[15, 16]. *CYP85A2* is involved in the brassinosteroid biosynthesis pathway in *Arabidopsis thaliana* and it converts 6-deoxocastasterone to castasterone, which is followed by the conversion of castasterone to brassinolide[17]. Brassinosteroids (BRs) are broad-spectrum plant

growth regulators, playing an important role in plant growth and development, as well as in biological and abiotic stress responses[18]. Mutations in the *CYP85A2* gene have led to an increased production of the dwarf phenotype[19], and an overexpression of the *CYP85A* family gene resulting in increased BR content, biomass, plant height, plant fresh weight and fruit yield[20]. These results showed that, *CKX3* and *CYP85A2* may affect soybean yield and architecture related traits through different molecular mechanisms. The potential effect of functional mutations in these genes on the phenotypes was further confirmed by our haplotype tests. However, to verify whether these candidate genes and functional mutations are the true cause of the phenotypic differences, further functional verification of these genes is necessary. Multiple methods such as the construction of isolated populations, transgene, gene knockout, gene editing, and expression verification could be used for this purpose. In this study, we performed expression verification in seedlings with different haplotypes/phenotypes for six genes *GL3*, *GSTL3*, *GSTT1b*, *CKX3*, *CKX4* and *CYP85A2*. The results show that, except for no expression were detected for *CKX4*, all the other five genes were expressed differently for different haplotypes/phenotypes in seedlings; the expression levels of *GL3*, *GSTL3* and *GSTT1b* related to isoflavone content in the strains with high isoflavone content values was significantly higher than that in the strains with low isoflavone content values (T-test, P<0.05); the expression level of *CKX3* in the strains with high yield phenotype values was significantly higher than that in the strains with low yield phenotype values (T-test, P<0.05) (**Supplementary Figure 55**).

The highest goal of plant breeding is to aggregate many desired traits into a single genome. Breeders need to simultaneously select and improve multiple related traits. However, because multiple traits are interrelated, it is possible that when screening for a favourable trait one also selects an unfavourable one. Understanding the genetic network behind different traits can help breeders increase breeding efficiency. Although soybean genetic networks for multiple agronomic traits have been established at the loci level[4], we built a new phenotype-gene network which includes 34 traits and 853 genes. This network reflects the relationship between

phenotypes and genes more directly than the previous phenotype-SAL network, and is

more conducive to the discovery of important candidate genes. For example, the *Hub*

Mixed 1 (*HM1*) node was associated with two or more trait types (architecture, colour

or protein), while the *HBT* gene in the *HM1* node was associated with six architectural

traits (branch number, main stem number, plant height, stem strength, lodging, and

podding habit) and four protein-related traits (phenylalanine content, isoleucine

content, tyrosine content, and glycine content). It is known that the *HBT* gene belongs

to the *CDC27b* gene family and is involved in cell cycle regulation, which is related

to cell development and division[21]. Therefore, soybean architecture is likely

affected by *HBT*, despite its relationship with amino acid content is unclear. There are

also many other interesting examples of the above. Leaf shape is known to affect

photosynthesis efficiency, followed by carbohydrate accumulation, and, as a

consequence, oil accumulation, while Hub *HM2*, containing *FAD2*, connects oil

content and leaf shape[22]. It has also been reported that oil traits and seed coat lustre

traits experienced parallel selection during bean domestication[23]. The hub *HM3*

node connects oil-related traits and seed coat lustre. Anthocyanin synthesis and

isoflavone synthesis share part of their metabolic pathways and hub *HM5* connects

colour traits and isoflavone content. Our phenotype-gene network may surpass the

phenotype-SAL network in terms of candidate gene selection, which is also beneficial

to polymerization breeding programs. For example, breeders can achieve

polymerization breeding by directly selecting a favourable gene (such as *CYP85A2*) in

Hub *HM4*, which is related to both yield and architecture traits, and eliminate the

confusion of other adverse genes located in the same SAL. Furthermore, it is worth

noting that the architecture related traits, which centrally connect various other trait

categories, have the most extensive connectivity. In other words, there are numerous

relationships between architecture related traits and other trait categories in the

phenotype-gene network (**Figure 6**), suggesting that some candidate genes related to

architecture traits may also be related to other trait types. This may provide theoretical

support and practical guidance for parallel selection breeding and promote "ideotype"

breeding in soybean. The next step is to conduct more in-depth functional

investigations on genes with a potential application value, such as *CKX3* and *CYP85A2*. This would help promote the design and breeding process of soybean varieties with a higher yield and quality. Overall, our work is conducive to promoting soybean genome functional research and genomic breeding.

Materials and Methods

Plant materials and phenotyping

A total of 250 soybean varieties were analysed in this study, which were provided by the National Crop Germplasm Resources Platform, Institute of Crop Genetics, Institute of Crop Sciences, Chinese Academy of Agricultural Sciences. All materials were planted and phenotyped at three locations: the Gongzhuling experimental site in the Jilin Academy of Agricultural Sciences (latitude 43.51°, east longitude 124.80°), the Harbin experimental site in the Heilongjiang Academy of Agricultural Sciences (45.68° north latitude, 126.61° east longitude), and the Chifeng experimental site un the Agricultural Science Institute in Inner Mongolia (42.27° north latitude, 118.90° east longitude) in late April of 2008, 2009 and 2010, respectively. Grain protein content was measured using the Kjeldahl method from National Food Safety Standard GB5009.5-2010 China[24], while the grain fatty acid contents were determined using the Soxhlet extraction method from National Food Safety Standard GB/T5512-2008, China[25]. Amino acid content was determined using high performance liquid chromatography (S433D, Seckam, Germany) following a previous amino acid determination method from National Food Safety Standard GB/T 18246-2000, China[26]. Grain isoflavone content was determined using high performance liquid chromatography from National Food Safety Standard GB/T23788-2009, China[27]. Finally, the phenotypic data were integrated using the BLUP (Best Linear Unbiased Prediction) method using the R package lme4[28] in order to remove environmental effects and obtain stable genetic phenotypes. Seeds were planted in (CLC-BIV-M/CLC404-TV, MMM, Germany) at 20°C (with 12-h day/12-h night) and a relative humidity of 60–80% till six leaves stage (about two-week-old).

Two-week-old seedlings (24°C, 12-h day/12-h night cycle) were used in this qPCR

validation.

DNA preparation and sequencing

The genomic DNA for all soybean accessions were extracted from soybean leaves

after three weeks of growth. DNA extraction was performed using the

cetyltrimethylammonium bromide (CTAB) method[29]. The library for each

accession was constructed with an insert size of approximately 500 base pairs,

following manufacturer's instructions (Illumina Inc., San Diego, CA, USA). All

soybean accessions were sequenced and paired-end 150 bp reads were produced using

an Illumina NovaSeq 6000 sequencer at the BerryGenomics Company

(http://www.berrygenomics.com/ Beijing, China).

Total RNA extraction, cDNA synthesis and qRT-PCR analysis

Total RNA was isolated for each sample using TRIzol Reagent (Invitrogen,

Nottingham, UK) according to the manufacturer's instructions. The purified RNA was

stored at -80°C until subsequent analyses. According to the manufacturer's

instructions (Takara, Shiga, Japan), first-strand cDNA synthesis was performed using

M-MLV reverse transcriptase. Quantitative real-time PCR (qRT-PCR) was performed

using a SYBR Premix Ex Taq Kit (Takara) and a real-time PCR machine (CFX96;

Bio-Rad, Hercules, CA, USA), following the manufacturer's instructions. The

procedure used for qRT-PCR was 95°C at 10 minutes, followed by 38 cycles of 15 s at

95°C and 60 s at 61-62°C. β-actin was used as a reference gene for analysis of relative

expression patterns of mRNA. The reactions were carried out with three biological

replicates, with at least two technical replicates for each sample. The data were

analyzed using the method according to the previous study[30], and the means ±

standard errors (SE) of three biological replicates are presented.

Mapping, variant calling and annotation

Raw paired-end resequencing reads were first cleaned by removing reads with

adaptors, reads of low quality and reads with "N"s. The high-quality clean reads were
then mapped to the soybean reference genome (Williams 82 assembly v2.1) with
BWA[31]. Statistical analyses of mapping rate and genomic coverage of clean reads
were performed using in-house scripts. The Speedseq pipeline[32] was used for SNP
and Indel calling, and vcftools[33] was used for genotype filtering. Missing genotypes
were imputed and phased through a localized haplotype clustering algorithm
implemented using Beagle v3.0[34]. Variant annotation was performed using
ANNOVAR[35] against the soybean gene model set v2.1.42. After annotation, SNPs
and Indels were categorized into exonic, intronic, intergenic, splicing, 5ʹUTRs,
3ʹUTRs, upstream, and downstream. Exonic SNPs were further categorized into
synonymous, nonsynonymous, stop gain, and stop loss. Exonic Indels were further
categorized into frameshift, non-frameshift, stop gain, and stop loss.

Population structure analysis
Approximately 6M SNPs from the 250 soybean accessions were concatenated for the
construction of a phylogenetic tree. Using a neighbour joining algorithm with a
pairwise gap deletion method for 100 bootstrap replications, a phylogenetic tree was
constructed with MegaCC[36]. The output was displayed using the iTOL[37] web
tool. With the whole genome genotype as the input, a principal component analysis
(PCA) was done using flashPCA [38] and the first two eigenvectors were plotted. A
population admixture analysis with $k = 2$ to $k = 5$ parameters were set to infer the
admixture of ancestors using fastSTRUCTURE.

Genetic diversity analysis
Linkage disequilibrium analyses for each subpopulation were performed using
PLINK[39] by calculating the correlation coefficient (r^2) of any two SNP pairs in one
chromosome. An LD decay plot was drawn using the average r^2 value for the distance
from 0 to 1,000 kb. Pairwise IBS calculations were also performed using PLINK and
a distance matrix was generated for each subpopulation. Population genetic diversities
were measured using VCFtools[33] by calculating $\theta\pi$ and *Fst*. $\theta\pi$ was used to measure

the genetic diversity of each subpopulation, while *Fst*, plus the allele frequency

distribution (AFD) plot (which was generated by in-house scripts), were used to

measure genetic diversity between subpopulations. In addition, sliding window

calculations of r^2, $\theta\pi$, *Fst* and *Tajima' D* values were also performed for genome-wide

displays of soybean genetic diversities with a 100 kb window and a 10 kb step.

Selective sweep analysis

We used two methods to detect selective sweep regions across the soybean genome:

Tajima' D combined $\theta\pi$ and *Fst* combined $\theta\pi$ ratios. Firstly, a genome-wide sliding

window calculation of $\theta\pi$, *Fst*, and *Tajima' D* values (with a 100 kb window and a 10

kb step) were performed on landraces, cultivars, and the whole population,

respectively. Secondly, the top 5% of the *Tajima' D* and $\theta\pi$ windows for the whole

population were selected. In addition, the top 5% of the *Fst* and $\theta\pi$ ratio windows for

the landraces versus cultivars were also selected. Thirdly, the selected windows from

these two methods were merged together to become the final selective sweep regions.

ROH analyses for each accession were performed using PLINK[39] with the

parameters of a minimum ROH length set to 300 kb.

GWAS and significantly associated loci

Association analysis for each trait on each SNP with an MAF larger than 0.05 was

performed using a single-locus mixed linear model (MLM) implemented in

GEMMA[40] (which corrects confounding by population structure and the

relatedness matrix). The GWAS results were displayed using a Manhattan plot and a

QQ-plot created with the R package CMplot[41] . A clump based method

implemented in PLINK[39] was used to reduce a false peak and to detect real SALs.

The P-value cut-off was set to 10^{-5} so as to, firstly, uncover significant associated

SNPs. Following this, for each significantly associated SNP, if there were more than

10 SNPs within a 100 kb distance that had P-values smaller than 10^{-4}, then the region

was regarded as a potential SAL. Finally, all overlapping SALs were merged to

generate final SAL sets and the SNP with the smallest P-value in a SAL was defined

as a peak.

Detection of significantly associated genes

There are usually tens of genes in a SAL, and it is difficult to determine which genes

are truly associated with traits and which are irrelevant. We improved a functional

mutation-based haplotype test method for SAG discovery in SAL. As most variants

within a gene are non-functional, the gene's amino acid sequence and its function will

not change. Only a few variants have the potential to change a gene's amino acid

sequence, such as nonsynonymous SNPs, frameshift Indels, variants in splicing sites,

promoter regions, start codons, and stop codons. These combined functional

mutations can only produce two or three different gene haplotypes. It is possible to

test the relationship between gene haplotypes and traits. If they are significantly

associated, then the gene is also most likely associated with the trait, which is how

SAG is defined. In this study, Welch's test was used for a two-group haplotype test

and a Tukey's test was used for a multiple group haplotype test to detect SAGs.

Functional annotation of SAGs was directly retrieved from SoyBase[42].

Network construction

Of all the genes located in the SALs, the most significant SAGs with a P-value

smaller than 10^{-5}, and their corresponding traits, were retained to build the

phenotype-gene network for soybean. Protein-protein interaction information for

soybean was retrieved form the String database[43] and mapped to the soybean genes

using BLAST[44]. Construction, visualization and exploration of the network was

performed using Cytoscape[45].

Data availability

651   The raw sequence data reported in this paper have been deposited in the Genome

652   Sequence Archive[46] in BIG Data Center[47], Beijing Institute of Genomics (BIG),

653   Chinese Academy of Sciences, under accession numbers CRA002552 that are

publicly accessible at https://bigd.big.ac.cn/gsa. The variation data reported in this paper have been deposited in the Genome Variation Map [48] under accession number GVM000076 that can be publicly accessible at http://bigd.big.ac.cn/gvm/getProjectDetail?project=GVM000076. The bioinformatics analysis scripts used in this paper can be download through https://github.com/yjthu/GPB_250SoyReseq.

CRediT statement

Chunming Yang: Resources, Investigation, Validation. **Jun Yan:** Methodology, Formal analysis, Writing, Revision. **Shuqin Jiang:** Formal analysis. **Xia Li:** Investigation, Validation. **Haowei Min:** Conceptualization, Supervision, Formal analysis, Writing, Revision. **Xiangfeng Wang:** Conceptualization, Supervision, Writing, Revision. **Dongyun Hao:** Conceptualization, Supervision, Writing, Revision. All authors read and approved the final manuscript.

Competing interests

The authors declare no competing financial interests.

Acknowledgements

This research was supported by grants from the Agricultural Science and Technology Innovation Project, Jilin Province [CXGC2017ZY027], and from Program of Accurate Identification and Display of Soybean Germplasm, China [NB08-2130315-(25-31)-06, NB07-2130315-(25-30)-06, NB06-070401-(22-27)-05, NB2010-2130315-25-05]. We are grateful to Prof. Lijuan Qiu for her agreement of using the 250 soybean varieties from her laboratory at China Academy of Agricultural Sciences. We appreciate Dr. Zhangxiong Liu of China Academy of Agricultural Sciences for the technical guidance in soybean phenotypic characterization. Acknowledgement also goes to Yunshan Wei(Inner Mongolia Academy of Agriculture

& Animal Husbandry Sciences), Shuhong Wei and Qiang Wang(Heilongjiang

Academy of Agricultural Sciences), for partially phenotypic characterization of the

soybean population used in this work.

References

[1] Sedivy, E.J., F. Wu, and Y. Hanzawa. Soybean domestication: the origin, genetic
architecture and molecular bases. New Phytol 2017;214:539-553.
[2] Lam, H.M., X. Xu, X. Liu, W. Chen, G. Yang, F.L. Wong, et al. Resequencing of
31 wild and cultivated soybean genomes identifies patterns of genetic diversity
and selection. Nat Genet 2010;42:1053-9.
[3] Zhou, Z., Y. Jiang, Z. Wang, Z. Gou, J. Lyu, W. Li, et al. Resequencing 302 wild
and cultivated accessions identifies genes related to domestication and
improvement in soybean. Nat Biotechnol 2015;33:408-14.
[4] Fang, C., Y. Ma, S. Wu, Z. Liu, Z. Wang, R. Yang, et al. Genome-wide association
studies dissect the genetic networks underlying agronomical traits in soybean.
Genome Biol 2017;18:161.
[5] Schmutz, J., S.B. Cannon, J. Schlueter, J. Ma, T. Mitros, W. Nelson, et al. Genome
sequence of the palaeopolyploid soybean. Nature 2010;463:178-83.
[6] Nielsen, R. Molecular signatures of natural selection. Annu Rev Genet
2005;39:197-218.
[7] Purfield, D.C., S. McParland, E. Wall, and D.P. Berry. The distribution of runs of
homozygosity and selection signatures in six commercial meat sheep breeds.
PLoS One 2017;12:e0176780.
[8] Crowell, S., P. Korniliev, A. Falcao, A. Ismail, G. Gregorio, J. Mezey, et al.
Genome-wide association and high-resolution phenotyping link Oryza sativa
panicle traits to numerous trait-specific QTL clusters. Nat Commun
2016;7:10527.
[9] Yano, K., E. Yamamoto, K. Aya, H. Takeuchi, P.C. Lo, L. Hu, et al. Genome-wide
association study using whole-genome sequencing rapidly identifies new
genes influencing agronomic traits in rice. Nat Genet 2016;48:927-34.
[10] Meng, S., J. He, T. Zhao, G. Xing, Y. Li, S. Yang, et al. Detecting the QTL-allele
system of seed isoflavone content in Chinese soybean landrace population for
optimal cross design and gene system exploration. Theor Appl Genet
2016;129:1557-76.
[11] Zhang, J., X. Wang, Y. Lu, S.J. Bhusal, Q. Song, P.B. Cregan, et al.
Genome-wide Scan for Seed Composition Provides Insights into Soybean
Quality Improvement and the Impacts of Domestication and Breeding. Mol
Plant 2018;11:460-472.
[12] Messina, M. A brief historical overview of the past two decades of soy and
isoflavone research. J Nutr 2010;140:1350S-4S.
[13] Braidot, E., M. Zancani, E. Petrussa, C. Peresson, A. Bertolini, S. Patui, et al.
Transport and accumulation of flavonoids in grapevine (Vitis vinifera L.).

Plant Signal Behav 2008;3:626-32.

[14] Xu, W., C. Dubos, and L. Lepiniec. Transcriptional control of flavonoid
biosynthesis by MYB-bHLH-WDR complexes. Trends Plant Sci
2015;20:176-85.

[15] Li, J., X. Nie, J.L. Tan, and F. Berger. Integration of epigenetic and genetic
controls of seed size by cytokinin in Arabidopsis. Proc Natl Acad Sci U S A
2013;110:15479-84.

[16] Ashikari, M., H. Sakakibara, S. Lin, T. Yamamoto, T. Takashi, A. Nishimura, et al.
Cytokinin oxidase regulates rice grain production. Science 2005;309:741-5.

[17] Kim, T.W., J.Y. Hwang, Y.S. Kim, S.H. Joo, S.C. Chang, J.S. Lee, et al.
Arabidopsis CYP85A2, a cytochrome P450, mediates the Baeyer-Villiger
oxidation of castasterone to brassinolide in brassinosteroid biosynthesis. Plant
Cell 2005;17:2397-412.

[18] De Bruyne, L., M. Hofte, and D. De Vleesschauwer. Connecting growth and
defense: the emerging roles of brassinosteroids and gibberellins in plant innate
immunity. Mol Plant 2014;7:943-959.

[19] Kwon, M.F., Shozo & Ji, H.J. & Kim, Ho Bang & Takatsuto, Suguru & Yoshida,
Shigeo & An, Chung & Choe, Sunghwa. A double mutant for the CYP85A1
and CYP85A2 genes of Arabidopsis exhibits a brassinosteroid dwarf
phenotype. Journal of Plant Biology 2005;48:237-244.

[20] Northey, J.G., S. Liang, M. Jamshed, S. Deb, E. Foo, J.B. Reid, et al.
Farnesylation mediates brassinosteroid biosynthesis to regulate abscisic acid
responses. Nat Plants 2016;2:16114.

[21] Perez-Perez, J.M., O. Serralbo, M. Vanstraelen, C. Gonzalez, M.C. Criqui, P.
Genschik, et al. Specialization of CDC27 function in the Arabidopsis thaliana
anaphase-promoting complex (APC/C). Plant J 2008;53:78-89.

[22] Dar, A.A., A.R. Choudhury, P.K. Kancharla, and N. Arumugam. The FAD2 Gene
in Plants: Occurrence, Regulation, and Role. Front Plant Sci 2017;8:1789.

[23] Zhang, D., L. Sun, S. Li, W. Wang, Y. Ding, S.A. Swarm, et al. Elevation of
soybean seed oil content through selection for seed coat shininess. Nat Plants
2018;4:30-35.

[24] GB/T 5009.5-2010 National food safety standard Determination of protein in
foods.

[25] GB/T 5512-2008 Inspect of grain and oilseeds—Determination of crude fat
content in grain.

[26] GB/T 18246-2000 Determination of amino acids in feeds.

[27] GB/T 23788-2009 Determination of soybean isoflavone in health-care
food-High-performance liquid chromatography.

[28] https://github.com/lme4/lme4.

[29] Murray, M.G. and W.F. Thompson. Rapid isolation of high molecular weight
plant DNA. Nucleic Acids Res 1980;8:4321-5.

[30] Livak, K.J. and T.D. Schmittgen. Analysis of relative gene expression data using
real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods
2001;25:402-8.

[31] Li, H. and R. Durbin. Fast and accurate long-read alignment with
Burrows-Wheeler transform. Bioinformatics 2010;26:589-95.
[32] Chiang, C., R.M. Layer, G.G. Faust, M.R. Lindberg, D.B. Rose, E.P. Garrison, et
al. SpeedSeq: ultra-fast personal genome analysis and interpretation. Nat
Methods 2015;12:966-8.
[33] Danecek, P., A. Auton, G. Abecasis, C.A. Albers, E. Banks, M.A. DePristo, et al.
The variant call format and VCFtools. Bioinformatics 2011;27:2156-8.
[34] Browning, S.R. and B.L. Browning. Rapid and accurate haplotype phasing and
missing-data inference for whole-genome association studies by use of
localized haplotype clustering. Am J Hum Genet 2007;81:1084-97.
[35] Wang, K., M. Li, and H. Hakonarson. ANNOVAR: functional annotation of
genetic variants from high-throughput sequencing data. Nucleic Acids Res
2010;38:e164.
[36] Kumar, S., G. Stecher, D. Peterson, and K. Tamura. MEGA-CC: computing core
of molecular evolutionary genetics analysis program for automated and
iterative data analysis. Bioinformatics 2012;28:2685-6.
[37] Letunic, I. and P. Bork. Interactive Tree Of Life (iTOL) v4: recent updates and
new developments. Nucleic Acids Res 2019;47:W256-W259.
[38] Abraham, G. and M. Inouye. Fast principal component analysis of large-scale
genome-wide data. PLoS One 2014;9:e93766.
[39] Chang, C.C., C.C. Chow, L.C. Tellier, S. Vattikuti, S.M. Purcell, and J.J. Lee.
Second-generation PLINK: rising to the challenge of larger and richer datasets.
Gigascience 2015;4:7.
[40] Zhou, X. and M. Stephens. Genome-wide efficient mixed-model analysis for
association studies. Nat Genet 2012;44:821-4.
[41] https://github.com/YinLiLin/R-CMplot.
[42] Grant, D., R.T. Nelson, S.B. Cannon, and R.C. Shoemaker. SoyBase, the
USDA-ARS soybean genetics and genomics database. Nucleic Acids Res
2010;38:D843-6.
[43] Szklarczyk, D., J.H. Morris, H. Cook, M. Kuhn, S. Wyder, M. Simonovic, et al.
The STRING database in 2017: quality-controlled protein-protein association
networks, made broadly accessible. Nucleic Acids Res 2017;45:D362-D368.
[44] Madden, T. The BLAST sequence analysis tool. The NCBI Handbook [Internet].
2nd edition 2013;
[45] Shannon, P., A. Markiel, O. Ozier, N.S. Baliga, J.T. Wang, D. Ramage, et al.
Cytoscape: a software environment for integrated models of biomolecular
interaction networks. Genome Res 2003;13:2498-504.
[46] Wang, Y., F. Song, J. Zhu, S. Zhang, Y. Yang, T. Chen, et al. GSA: Genome
Sequence Archive<sup/>. Genomics Proteomics Bioinformatics
2017;15:14-18.
[47] National Genomics Data Center, M. and Partners. Database Resources of the
National Genomics Data Center in 2020. Nucleic Acids Res
2020;48:D24-D33.
[48] Song, S., D. Tian, C. Li, B. Tang, L. Dong, J. Xiao, et al. Genome Variation Map:

a data repository of genome variations in BIG Data Center. Nucleic Acids Res

2018;46:D944-D949.

Tables

819   **Table1.** Functional variants of representative significant associated genes

Variant ID	Chrom	Positon	Ref	Alt	Variant type	Gene ID	Gene symbol
c5s38936266	5	38936266	C	T	nonsynonymous SNV	GLYMA_05G206900	*GSTT1a*
c5s38940717	5	38940717	C	T	nonsynonymous SNV	GLYMA_05G207000	*GSTT1b*
c5s39035509	5	39035509	G	C	nonsynonymous SNV	GLYMA_05G208300	*GL3*
c5s39036346	5	39036346	T	C	nonsynonymous SNV	GLYMA_05G208300	*GL3*
c13s24804891	13	24804891	C	T	nonsynonymous SNV	GLYMA_13G135600	*GSTL3*
c13s24805363	13	24805363	A	T	splicing SNV	GLYMA_13G135600	*GSTL3*
c17s4143663	17	4143663	C	T	nonsynonymous SNV	GLYMA_17G054500	*CKX3*
c17s4143832	17	4143832	T	C	nonsynonymous SNV	GLYMA_17G054500	*CKX3*
c17s4146922	17	4146922	G	T	nonsynonymous SNV	GLYMA_17G054500	*CKX3*
c17s4151713	17	4151713	C	A	nonsynonymous SNV	GLYMA_17G054600	*CKX4*
c17s4151752	17	4151752	T	C	nonsynonymous SNV	GLYMA_17G054600	*CKX4*
c18s55526062	18	55526062	C	T	nonsynonymous SNV	GLYMA_18G272300	*CYP85A2*

Figures

Figure 1

Figure 1 Population structure of 250 soybean accessions. **A.** Geographic distribution of 250 soybean landraces and cultivars. Landraces are shown with green color and cultivars are shown with yellow color. **B.** Phylogenetic tree constructed for all soybean accessions. Group1-4 are shown with different colors, Landraces are labeled with green triangles and cultivars are labeled with yellow triangles. **C.** Statistics of the geographic origin for each subpopulation. **D.** Mixed ancestors analysis for soybean subpopulations. Each color represents an ancestral component. K from 2 to 5 are set to trace different ancestral components.   **E.** PCA plot of the first two eigenvectors for all soybean accessions. Landraces and cultivars are shown will different shape, while groups are shown with different colors.

Figure 2

Figure 2 Genetic diversity of soybean subpopulations. **A.** LD decay plots for landrace
(green), cultivars (yellow) and all soybean accessions (grey). **B.** LD decay plots for
soybean subpopulations. **C.** IBS values distribution for landrace (green), cultivars
(yellow) and all soybean accessions (grey). **D.** IBS values distribution for soybean
subpopulations.**E.** Comparison of $\theta\pi$ values for landrace (green), cultivars (yellow)
and all soybean accessions (grey). **F.** Comparison of $\theta\pi$ values for soybean
subpopulations. **G.** Comparison of *Fst* values between subpopulations. **H.** Landscape
of soybean genetic diversity across the whole genome. (a) Chromosomes. (b) Density
of genes (c) Density of SNPs (red) and Indels (blue). (d) LD values distribution for
landraces(green), cultivars(yellow) and all accessions(grey). (e) *Fst* values
distribution of landraces versus cultivars (f) $\theta\pi$ values distribution for
landraces(green), cultivars(yellow) and all accessions(grey). (g) *Tajima'D* values
distribution of all accessions. (h) Putative selective sweep regions detected by
Tajima'D combine $\theta\pi$. (i) Putative selective sweep regions detected by *Fst* combine
$\theta\pi$ ratios. (j) ROH region larger than 300 Kb.

Figure 3

Figure 3 GWAS of soybean isoflavone content. **A.** Manhantan plot and four candidate
genes for soybean isoflavone content**. B.** Chromosome location and transcripts
structure of the candidate genes. **C.** Soybean isoflavone content distribution for the
haplotypes of gene *GSTT1*. **D.** Soybean isoflavone content distribution for the
haplotypes of gene *GL3*. **E.** Soybean isoflavone content distribution for the
haplotypes of gene *GSTL3*. **F.** Diagram of soybean isoflavone synthesis and transport,
and the roles of candidate genes detected by GWAS. (*$P < 0.05$; **$P < 0.01$; n.s., not
significant)

Figure 4

Figure 4 Association of *CKX* and yield related traits in soybean. **A.** Manhantan plot of four yield related traits pod number per plant, seed number per plant, one hundred seed weight and seed size, and the candidate *CKX* genes. **B.** Pod number per plant distribution for the haplotypes of *CKX* genes.**C.** Seed number per plant distribution for the haplotypes of *CKX* genes. **D.** One hundred seed weight distribution for the haplotypes of *CKX* genes. **E.** Seed size distribution for the haplotypes of *CKX* genes. **F.** Chromosome location and transcripts structure of *CKX3* and *CKX4*. **G.** Phenotype correlation of four traits. **H.** Statistics of SAGs located in selective sweep regions and their percentage for trait categories. (*$P < 0.05$; **$P < 0.01$)

Figure 5

Figure 5 Association of *CYP85A2* and architecture or yield related traits in soybean.

A. Manhantan plot of plant height, main stem number, stem strength, lodging,

podding habit, seed weight per plant, and the candidate gene *CYP85A2*. **B.** Traits

distribution for the haplotypes of gene *CYP85A2*. **C.** Chromosome location and

transcripts structure of *CYP85A2*. **D.** Phenotype correlation of the six traits. (*$P <$

0.05; **$P <$ 0.01)

Figure 6

Figure 6 Phenotype-gene genetic network in soybean. Traits are solid rhombuses,

genes are solid circles, and hubs are hollow ellipses. Six trait categories, their

associated genes and links between them are colored accordingly, genes associated

with more than one trait categories are colored grey. Genes with protein-protein

interaction are linked with gray lines.

Supplementary material

Supplementary Tables 1-10; Supplementary Figures 1-55.

Supplementary Table 1 Summary of mapping and coverage

Supplementary Table 2 Summary of SNPs and Indels

Supplementary Table 3 The ancestry proportion estimates for each accession

Supplementary Table 4 Putative regions experiencing selective sweeps

Figure S6 GWAS of effective branch number using MLM. A. Density distribution of

effective branch number. B. Manhattan plots for effective branch number. Negative

log10 P-values from a genome-wide scan are plotted against SNP positions of 20

chromosomes. C. Quantile-quantile plot for effective branch number. The horizontal

red line indicates the significant threshold (10-5). Trait-associated SNPs above the

significant threshold are colored in red.

Figure S7 GWAS of pubescence density using MLM. **A.** Density distribution of

pubescence density. **B.** Manhattan plots for pubescence density. Negative $\log_{10}$

P-values from a genome-wide scan are plotted against SNP positions of 20

chromosomes. **C.** Quantile-quantile plot for pubescence density. The horizontal red

line indicates the significant threshold (10^{-5}). Trait-associated SNPs above the

significant threshold are colored in red.

Figure S8 GWAS of defollation using MLM. **A.** Density distribution of defollation. **B.**

Manhattan plots for defollation. Negative $\log_{10}$ P-values from a genome-wide scan are

plotted against SNP positions of 20 chromosomes. **C.** Quantile-quantile plot for

defollation. The horizontal red line indicates the significant threshold (10^{-5}).

Trait-associated SNPs above the significant threshold are colored in red.

Figure S9 GWAS of inflorenscence length using MLM. **A.** Density distribution of

inflorenscence length. **B.** Manhattan plots for inflorenscence length. Negative $\log_{10}$

P-values from a genome-wide scan are plotted against SNP positions of 20

chromosomes. **C.** Quantile-quantile plot for inflorenscence length. The horizontal red

line indicates the significant threshold (10^{-5}). Trait-associated SNPs above the

significant threshold are colored in red.

Figure S10 GWAS of leaf shape using MLM. **A.** Density distribution of leaf shape. **B.**

Manhattan plots for leaf shape. Negative $\log_{10}$ P-values from a genome-wide scan are

plotted against SNP positions of 20 chromosomes. **C.** Quantile-quantile plot for leaf

shape. The horizontal red line indicates the significant threshold (10^{-5}).

Trait-associated SNPs above the significant threshold are colored in red.

Figure S11 GWAS of leaflet size using MLM. **A.** Density distribution of leaflet size.

B. Manhattan plots for leaflet size. Negative $\log_{10}$ P-values from a genome-wide scan

are plotted against SNP positions of 20 chromosomes. **C.** Quantile-quantile plot for

leaflet size. The horizontal red line indicates the significant threshold (10^{-5}).

Trait-associated SNPs above the significant threshold are colored in red.

Figure S12 GWAS of lodging using MLM. **A.** Density distribution of lodging. **B.**

Manhattan plots for lodging. Negative $\log_{10}$ P-values from a genome-wide scan are

plotted against SNP positions of 20 chromosomes. **C.** Quantile-quantile plot for

lodging. The horizontal red line indicates the significant threshold (10^{-5}).

Trait-associated SNPs above the significant threshold are colored in red.

Figure S13 GWAS of number of nodes on main stem using MLM. **A.** Density

distribution of number of nodes on main stem. **B.** Manhattan plots for number of

nodes on main stem. Negative $\log_{10}$ P-values from a genome-wide scan are plotted

against SNP positions of 20 chromosomes. **C.** Quantile-quantile plot for number of

nodes on main stem. The horizontal red line indicates the significant threshold (10^{-5}).

Trait-associated SNPs above the significant threshold are colored in red.

Figure S14 GWAS of plant height using MLM. **A.** Density distribution of plant

height. **B.** Manhattan plots for plant height. Negative $\log_{10}$ P-values from a

genome-wide scan are plotted against SNP positions of 20 chromosomes. **C.**

Quantile-quantile plot for plant height. The horizontal red line indicates the significant

threshold (10^{-5}). Trait-associated SNPs above the significant threshold are colored in

red.

Figure S15 GWAS of plant type using MLM. **A.** Density distribution of plant type. **B.**

Manhattan plots for plant type. Negative $\log_{10}$ P-values from a genome-wide scan are

plotted against SNP positions of 20 chromosomes. **C.** Quantile-quantile plot for plant

type. The horizontal red line indicates the significant threshold (10^{-5}). Trait-associated

SNPs above the significant threshold are colored in red.

Figure S16 GWAS of stem termination using MLM. **A.** Density distribution of

podding habit. **B.** Manhattan plots for stem termination. Negative $\log_{10}$ P-values from

a genome-wide scan are plotted against SNP positions of 20 chromosomes. **C.**

Quantile-quantile plot for stem termination. The horizontal red line indicates the

significant threshold (10^{-5}). Trait-associated SNPs above the significant threshold are

colored in red.

Figure S17 GWAS of seed crack using MLM. **A.** Density distribution of seed crack.

B. Manhattan plots for seed crack. Negative $\log_{10}$ P-values from a genome-wide scan

are plotted against SNP positions of 20 chromosomes. **C.** Quantile-quantile plot for

seed crack. The horizontal red line indicates the significant threshold (10^{-5}).

Trait-associated SNPs above the significant threshold are colored in red.

Figure S18 GWAS of stem diameter using MLM. **A.** Density distribution of stem

diameter. **B.** Manhattan plots for stem diameter. Negative $\log_{10}$ P-values from a

genome-wide scan are plotted against SNP positions of 20 chromosomes. **C.**

Quantile-quantile plot for stem diameter. The horizontal red line indicates the

significant threshold (10^{-5}). Trait-associated SNPs above the significant threshold are

colored in red.

Figure S19 GWAS of stem intension using MLM. **A.** Density distribution of stem

intension. **B.** Manhattan plots for stem intension. Negative $\log_{10}$ P-values from a

genome-wide scan are plotted against SNP positions of 20 chromosomes. **C.**

Quantile-quantile plot for stem intension. The horizontal red line indicates the

significant threshold (10^{-5}). Trait-associated SNPs above the significant threshold are

colored in red.

Figure S20 GWAS of pubescence color using MLM. **A.** Density distribution of

pubescence color. **B.** Manhattan plots for pubescence color. Negative $\log_{10}$ P-values

from a genome-wide scan are plotted against SNP positions of 20 chromosomes. **C.**

Quantile-quantile plot for pubescence color. The horizontal red line indicates the

significant threshold (10^{-5}). Trait-associated SNPs above the significant threshold are

colored in red.

Figure S21 GWAS of flower color using MLM. **A.** Density distribution of flower

color. **B.** Manhattan plots for flower color. Negative $\log_{10}$ P-values from a

genome-wide scan are plotted against SNP positions of 20 chromosomes. **C.**

Quantile-quantile plot for flower color. The horizontal red line indicates the

significant threshold (10^{-5}). Trait-associated SNPs above the significant threshold are

colored in red.

Figure S22 GWAS of leaf color using MLM. **A.** Density distribution of leaf color. **B.**

Manhattan plots for leaf color. Negative $\log_{10}$ P-values from a genome-wide scan are

plotted against SNP positions of 20 chromosomes. **C.** Quantile-quantile plot for leaf

color. The horizontal red line indicates the significant threshold (10^{-5}).

Trait-associated SNPs above the significant threshold are colored in red.

Figure S23 GWAS of mature pod color using MLM. **A.** Density distribution of

mature pod color. **B.** Manhattan plots for mature pod color. Negative $\log_{10}$ P-values

from a genome-wide scan are plotted against SNP positions of 20 chromosomes. **C.**

Quantile-quantile plot for mature pod color. The horizontal red line indicates the

significant threshold (10^{-5}). Trait-associated SNPs above the significant threshold are

colored in red.

Figure S24 GWAS of seed coat luster using MLM. **A.** Density distribution of seed

coat luster. **B.** Manhattan plots for seed coat luster. Negative $\log_{10}$ P-values from a

genome-wide scan are plotted against SNP positions of 20 chromosomes. **C.**

Quantile-quantile plot for seed coat luster. The horizontal red line indicates the

significant threshold (10^{-5}). Trait-associated SNPs above the significant threshold are

colored in red.

Figure S25 GWAS of isoflavone content using MLM. **A.** Density distribution of

isoflavone content. **B.** Manhattan plots for isoflavone content. Negative $\log_{10}$ P-values

from a genome-wide scan are plotted against SNP positions of 20 chromosomes. **C.**

Quantile-quantile plot for isoflavone content. The horizontal red line indicates the

significant threshold (10^{-5}). Trait-associated SNPs above the significant threshold are

colored in red.

Figure S26 GWAS of linoleic acid content using MLM. **A.** Density distribution of

linoleic acid content. **B.** Manhattan plots for linoleic acid content. Negative $\log_{10}$

P-values from a genome-wide scan are plotted against SNP positions of 20

chromosomes. **C.** Quantile-quantile plot for linoleic acid content. The horizontal red

line indicates the significant threshold (10^{-5}). Trait-associated SNPs above the

significant threshold are colored in red.

Figure S27 GWAS of linolenic acid content using MLM. **A.** Density distribution of

linolenic acid content. **B.** Manhattan plots for linolenic acid content. Negative $\log_{10}$

P-values from a genome-wide scan are plotted against SNP positions of 20

chromosomes. **C.** Quantile-quantile plot for linolenic acid content. The horizontal red

line indicates the significant threshold (10^{-5}). Trait-associated SNPs above the

significant threshold are colored in red.

Figure S28 GWAS of oleic acid content using MLM. **A.** Density distribution of oleic

acid content. **B.** Manhattan plots for oleic acid content. Negative $\log_{10}$ P-values from a

genome-wide scan are plotted against SNP positions of 20 chromosomes. **C.**

Quantile-quantile plot for oleic acid content. The horizontal red line indicates the

significant threshold (10^{-5}). Trait-associated SNPs above the significant threshold are

colored in red.

Figure S29 GWAS of palmitic acid content using MLM. **A.** Density distribution of

palmitic acid content. **B.** Manhattan plots for palmitic acid content. Negative $\log_{10}$

P-values from a genome-wide scan are plotted against SNP positions of 20

chromosomes. **C.** Quantile-quantile plot for palmitic acid content. The horizontal red

line indicates the significant threshold (10^{-5}). Trait-associated SNPs above the

significant threshold are colored in red.

Figure S30 GWAS of crude protein content using MLM. **A.** Density distribution of

crude protein content. **B.** Manhattan plots for crude protein content. Negative $\log_{10}$

P-values from a genome-wide scan are plotted against SNP positions of 20

chromosomes. **C.** Quantile-quantile plot for crude protein content. The horizontal red

line indicates the significant threshold (10^{-5}). Trait-associated SNPs above the

significant threshold are colored in red.

Figure S31 GWAS of alanine content using MLM. **A.** Density distribution of alanine

content. **B.** Manhattan plots for alanine content. Negative $\log_{10}$ P-values from a

genome-wide scan are plotted against SNP positions of 20 chromosomes. **C.**

Quantile-quantile plot for alanine content. The horizontal red line indicates the

significant threshold (10^{-5}). Trait-associated SNPs above the significant threshold are

colored in red.

Figure S32 GWAS of arginine content using MLM. **A.** Density distribution of

arginine content. **B.** Manhattan plots for arginine content. Negative $\log_{10}$ P-values

from a genome-wide scan are plotted against SNP positions of 20 chromosomes. **C.**

Quantile-quantile plot for arginine content. The horizontal red line indicates the

significant threshold (10^{-5}). Trait-associated SNPs above the significant threshold are

colored in red.

Figure S33 GWAS of aspartic acid content using MLM. **A.** Density distribution of

aspartic acid content. **B.** Manhattan plots for aspartic acid content. Negative $\log_{10}$

P-values from a genome-wide scan are plotted against SNP positions of 20

chromosomes. **C.** Quantile-quantile plot for aspartic acid content. The horizontal red

line indicates the significant threshold (10^{-5}). Trait-associated SNPs above the

significant threshold are colored in red.

Figure S34 GWAS of glutamate content using MLM. **A.** Density distribution of

glutamate content. **B.** Manhattan plots for glutamate content. Negative $\log_{10}$ P-values

from a genome-wide scan are plotted against SNP positions of 20 chromosomes. **C.**

Quantile-quantile plot for glutamate content. The horizontal red line indicates the

significant threshold (10^{-5}). Trait-associated SNPs above the significant threshold are

colored in red.

Figure S35 GWAS of glycine content using MLM. **A.** Density distribution of glycine

content. **B.** Manhattan plots for glycine content. Negative $\log_{10}$ P-values from a

genome-wide scan are plotted against SNP positions of 20 chromosomes. **C.**

Quantile-quantile plot for glycine content. The horizontal red line indicates the

significant threshold (10^{-5}). Trait-associated SNPs above the significant threshold are

colored in red.

Figure S36 GWAS of histidine content using MLM. **A.** Density distribution of

histidine content. **B.** Manhattan plots for histidine content. Negative $\log_{10}$ P-values

from a genome-wide scan are plotted against SNP positions of 20 chromosomes. **C.**

Quantile-quantile plot for histidine content. The horizontal red line indicates the

significant threshold (10^{-5}). Trait-associated SNPs above the significant threshold are

colored in red.

Figure S37 GWAS of isoleucine content using MLM. **A.** Density distribution of

isoleucine content. **B.** Manhattan plots for isoleucine content. Negative $\log_{10}$ P-values

from a genome-wide scan are plotted against SNP positions of 20 chromosomes. **C.**

Quantile-quantile plot for isoleucine content. The horizontal red line indicates the

significant threshold (10^{-5}). Trait-associated SNPs above the significant threshold are

colored in red.

Figure S38 GWAS of leucine content using MLM. **A.** Density distribution of leucine

content. **B.** Manhattan plots for leucine content. Negative $\log_{10}$ P-values from a

genome-wide scan are plotted against SNP positions of 20 chromosomes. **C.**

Quantile-quantile plot for leucine content. The horizontal red line indicates the

significant threshold (10^{-5}). Trait-associated SNPs above the significant threshold are

colored in red.

Figure S39 GWAS of lysine content using MLM. **A.** Density distribution of lysine

content. **B.** Manhattan plots for lysine content. Negative $\log_{10}$ P-values from a

genome-wide scan are plotted against SNP positions of 20 chromosomes. **C.**

Quantile-quantile plot for lysine content. The horizontal red line indicates the

significant threshold (10^{-5}). Trait-associated SNPs above the significant threshold are

colored in red.

Figure S40 GWAS of methionine content using MLM. **A.** Density distribution of

methionine content. **B.** Manhattan plots for methionine content. Negative $\log_{10}$

P-values from a genome-wide scan are plotted against SNP positions of 20

chromosomes. **C.** Quantile-quantile plot for methionine content. The horizontal red

line indicates the significant threshold (10^{-5}). Trait-associated SNPs above the

significant threshold are colored in red.

Figure S41 GWAS of phenylalanine content using MLM. **A.** Density distribution of

phenylalanine content. **B.** Manhattan plots for phenylalanine content. Negative $\log_{10}$

P-values from a genome-wide scan are plotted against SNP positions of 20

chromosomes. **C.** Quantile-quantile plot for phenylalanine content. The horizontal red

line indicates the significant threshold (10^{-5}). Trait-associated SNPs above the

significant threshold are colored in red.

Figure S42 GWAS of proline content using MLM. **A.** Density distribution of proline

content. **B.** Manhattan plots for proline content. Negative $\log_{10}$ P-values from a

genome-wide scan are plotted against SNP positions of 20 chromosomes. **C.**

Quantile-quantile plot for proline content. The horizontal red line indicates the

significant threshold (10^{-5}). Trait-associated SNPs above the significant threshold are

colored in red.

Figure S43 GWAS of serine content using MLM. **A.** Density distribution of serine

content. **B.** Manhattan plots for serine content. Negative $\log_{10}$ P-values from a

genome-wide scan are plotted against SNP positions of 20 chromosomes. **C.**

Quantile-quantile plot for serine content. The horizontal red line indicates the

significant threshold (10^{-5}). Trait-associated SNPs above the significant threshold are

colored in red.

Figure S44 GWAS of threonine content using MLM. **A.** Density distribution of

threonine content. **B.** Manhattan plots for threonine content. Negative $\log_{10}$ P-values

from a genome-wide scan are plotted against SNP positions of 20 chromosomes. **C.**

Quantile-quantile plot for threonine content. The horizontal red line indicates the

significant threshold (10^{-5}). Trait-associated SNPs above the significant threshold are

colored in red.

Figure S45 GWAS of tyrosine content using MLM. **A.** Density distribution of

tyrosine content. **B.** Manhattan plots for tyrosine content. Negative $\log_{10}$ P-values

from a genome-wide scan are plotted against SNP positions of 20 chromosomes. **C.**

Quantile-quantile plot for tyrosine content. The horizontal red line indicates the

significant threshold (10^{-5}). Trait-associated SNPs above the significant threshold are

colored in red.

Figure S46 GWAS of valine content using MLM. **A.** Density distribution of valine

content. **B.** Manhattan plots for valine content. Negative $\log_{10}$ P-values from a

genome-wide scan are plotted against SNP positions of 20 chromosomes. **C.**

Quantile-quantile plot for valine content. The horizontal red line indicates the

significant threshold (10^{-5}). Trait-associated SNPs above the significant threshold are

colored in red.

Figure S47 GWAS of total amino acids content using MLM. **A.** Density distribution

of total amino acids content. **B.** Manhattan plots for total amino acids content.

Negative $\log_{10}$ P-values from a genome-wide scan are plotted against SNP positions

of 20 chromosomes. **C.** Quantile-quantile plot for total amino acids content. The

horizontal red line indicates the significant threshold (10^{-5}). Trait-associated SNPs

above the significant threshold are colored in red.

Figure S48 GWAS of hundred grain weight using MLM. **A.** Density distribution of

one hundred seed weight. **B.** Manhattan plots for hundred grain weight. Negative

$\log_{10}$ P-values from a genome-wide scan are plotted against SNP positions of 20

chromosomes. **C.** Quantile-quantile plot for hundred grain weight. The horizontal red

line indicates the significant threshold (10^{-5}). Trait-associated SNPs above the

significant threshold are colored in red.

Figure S49 GWAS of pod number per plant using MLM. **A.** Density distribution of

pod number per plant. **B.** Manhattan plots for pod number per plant. Negative $\log_{10}$

P-values from a genome-wide scan are plotted against SNP positions of 20

chromosomes. **C.** Quantile-quantile plot for pod number per plant. The horizontal red

line indicates the significant threshold (10^{-5}). Trait-associated SNPs above the

significant threshold are colored in red.

Figure S50 GWAS of pod size using MLM. **A.** Density distribution of pod size. **B.**

Manhattan plots for pod size. Negative $\log_{10}$ P-values from a genome-wide scan are

plotted against SNP positions of 20 chromosomes. **C.** Quantile-quantile plot for pod

size. The horizontal red line indicates the significant threshold (10^{-5}). Trait-associated

SNPs above the significant threshold are colored in red.

Figure S51 GWAS of seed number per plant using MLM. **A.** Density distribution of

seed number per plant. **B.** Manhattan plots for seed number per plant. Negative $\log_{10}$

P-values from a genome-wide scan are plotted against SNP positions of 20

chromosomes. **C.** Quantile-quantile plot for seed number per plant. The horizontal red

line indicates the significant threshold (10^{-5}). Trait-associated SNPs above the

significant threshold are colored in red.

Figure S52 GWAS of seed number per pod using MLM. **A.** Density distribution of

seed number per pod. **B.** Manhattan plots for seed number per pod. Negative $\log_{10}$

P-values from a genome-wide scan are plotted against SNP positions of 20

chromosomes. **C.** Quantile-quantile plot for seed number per pod. The horizontal red

line indicates the significant threshold (10^{-5}). Trait-associated SNPs above the

significant threshold are colored in red.

Figure S53 GWAS of seed size using MLM. **A.** Density distribution of seed size. **B.**

Manhattan plots for seed size. Negative $\log_{10}$ P-values from a genome-wide scan are

plotted against SNP positions of 20 chromosomes. **C.** Quantile-quantile plot for seed

size. The horizontal red line indicates the significant threshold (10^{-5}). Trait-associated

SNPs above the significant threshold are colored in red.

Figure S54 GWAS of seed weight per plant using MLM. **A.** Density distribution of

seed weight per plant. **B.** Manhattan plots for seed weight per plant. Negative $\log_{10}$

P-values from a genome-wide scan are plotted against SNP positions of 20

chromosomes. **C.** Quantile-quantile plot for seed weight per plant. The horizontal red

line indicates the significant threshold (10^{-5}). Trait-associated SNPs above the

significant threshold are colored in red.

Figure S55 Gene expression validation of different haplotypes/phenotypes for five

candidate genes. **A.** Phenotype distribution (left) and expression level (right) of *GL3*

for different haplotypes. **B.** Phenotype distribution (left) and expression level (right)

of *GSTL3* for different haplotypes. **C.** Phenotype distribution (left) and expression

level (right) of *GSTT1b* for different haplotypes. **D.** Phenotype distribution (left) and

expression level (right) of *CKX3* for different haplotypes. **E.** Phenotype distribution

(left) and expression level (right) of *CYP85A2* for different haplotypes.

1258

1259