RSC primes the quiescent genome for hypertranscription upon cell cycle re-entry

Christine E. Cucinotta, Rachel H. Dell, Keean C.A. Braceros, and Toshio Tsukiyama*

Division of Basic Sciences, Fred Hutchinson Cancer Research Center, Seattle WA

*To whom correspondence should be addressed.

Tel: 206-667-4996

 Email: ttsukiya@fredhutch.org

Abstract

Quiescence is a reversible G_0 state essential for differentiation, regeneration, stem cell

renewal, and immune cell activation. Necessary for long-term survival, quiescent

chromatin is compact, hypoacetylated, and transcriptionally inactive. How transcription

activates upon cell-cycle re-entry is undefined. Here we report robust, widespread

transcription within the first minutes of quiescence exit. During quiescence, the

chromatin-remodeling enzyme RSC was already bound to the genes induced upon

quiescence exit. RSC depletion caused severe quiescence exit defects: a global

decrease in RNA polymerase II (Pol II) loading, Pol II accumulation at transcription start

sites, initiation from ectopic upstream loci, and aberrant antisense transcription. These

phenomena were due to a combination of highly robust Pol II transcription and severe

chromatin defects in the promoter regions and gene bodies. Together, these results

uncovered multiple mechanisms by which RSC facilitates initiation and maintenance of

large-scale, rapid gene expression despite a globally repressive chromatin state.

Introduction

For decades scientists have used budding yeast to uncover mechanisms of chromatin

regulation of gene expression; and the vast majority of these studies were performed in

exponentially growing (hereafter log) cultures [1]. Log phase, however, is not a common

growth stage in unicellular organism lifecycles. Furthermore, many cell populations in

multicellular organisms, such as in humans, are not actively dividing [2–4]. Indeed, the

majority of "healthy" cells on Earth are not sustained in a persistently dividing state [3].

Non-proliferating cells reside in a G_0 state, which generally means these cells are either

terminally differentiated, senescent, or quiescent. The quiescent state provides

advantages to organisms: quiescence allows cells to remain dormant for long periods of

time to survive harsh conditions or to prevent over-proliferation [3–5,2]. Notwithstanding

this so-called "dormant state", quiescent cells can exit quiescence and re-enter the

mitotic cell-cycle in response to growth cues or environmental stimuli, which

distinguishes quiescence from other G_0 states. A major hallmark of quiescence is the

chromatin landscape—vast histone de-acetylation and chromatin compaction occur

during quiescence entry [6–8]. These events happen alongside a global narrowing of

nucleosome depleted regions (NDR) and increased resistance to micrococcal nuclease

(MNase) digestion, indicating a repressive chromatin environment [6]. Together, these

features of quiescent cells point to a critical role for chromatin regulation of the

quiescent state. However, the role of chromatin regulation upon exit from quiescence is

unknown.

Reversibility is a conserved hallmark of quiescent cells and is required for proper

stem-cell niche maintenance, T-cell activation, and wound healing in metazoans [4,9].

Therefore, we sought to elucidate molecular mechanisms by which cells can overcome

this repressive chromatin environment to re-enter the mitotic cell cycle. Given its genetic

tractability, the ease by which quiescent cells can be purified, and high level of

conservation among chromatin and transcription machinery, we turned to the budding

yeast *Saccharomyces cerevisiae* [10]. We can easily isolate quiescent yeast cells after

seven days of growth and density-gradient centrifugation. In this context, we can study

pure populations of quiescent yeast, a cell fate that is distinct from other cell types

present in a saturated culture [11].

Since DNA is wrapped around an octamer of histone proteins in increments of

~147bp to form nucleosomes [12], enzymes must move nucleosomes to give access to

transcription initiation factors [13]. One such enzyme is the SWI/SNF-family member,

RSC, which is a 17-subunit chromatin remodeling enzyme complex [14]. RSC contains

an ATP-dependent translocase, Sth1 [15–18], multiple subunits with bromodomains

(more than half of all bromodomains in the yeast genome are in RSC) and two zinc-

finger DNA-binding domains, which allow RSC to target and remodel chromatin [19,20].

Many components of the RSC complex are essential for viability in budding yeast and

the complex is conserved in humans, where it is named PBAF. In humans, mutations in

PBAF genes are associated with 40% of kidney cancers [21]; and 20% of all human

cancers contain mutations within SWI/SNF family genes [22], underscoring the

importance of such complexes in human health.

The best-described role for RSC in regulating chromatin architecture and

transcriptions is its ability to generate NDRs, by sliding or evicting nucleosomes [23–25].

Moving the +1 nucleosome allows for TATA binding protein (TBP) promoter binding and

transcription initiation [26]. To this end, RSC mostly localizes to the -1, +1, and +2

nucleosomes in log cells [27–29]. However, RSC has also been implicated in the

transcription elongation step where it tethers to RNA polymerase and can localize to

gene bodies [30–32]. Additionally, RSC binds nucleosomes within the so-called "wide

NDRs", where there are MNase-sensitive nucleosome-sized fragments, known as

"fragile" nucleosomes [33–36]. These RSC-bound nucleosomes are likely partially

unwrapped to aid in rapid gene induction [36–39].

In this study, we investigated how genes are transcribed during the first minutes

of quiescence exit. We were particularly interested in uncovering mechanisms to

overcome highly repressive chromatin found in quiescent cells. Unexpectedly, ~60% of

the yeast genome was transcribed by RNA polymerase II (Pol II) by the first 10-minutes

of exit, despite the highly repressive chromatin architecture present in quiescence. We

found that this hypertranscription [40] event is RSC dependent and that RSC binds

across the genome to ~80% of NDRs in quiescent cells. Upon RSC depletion, we

observed canonical abrogation of transcription initiation, defects in Pol II clearance past

the +1 nucleosome, and gross Pol II mislocalization, resulting in abnormal upstream

initiation and aberrant non-coding antisense transcripts. We further showed that RSC

alters chromatin structure to facilitate these processes. Taken together, we propose a

model in which RSC is bound to NDRs in quiescent cells to facilitate robust and

accurate burst of transcription upon quiescent exit through multiple mechanisms.

Results

Hypertranscription occurs within minutes of nutrient repletion post-quiescence

To determine the earliest time at which transcription reactivates during

quiescence exit, we fed purified quiescent cells YPD medium and took time points to

determine the kinetics of Pol II C-terminal domain (CTD) phosphorylation by western

blot analysis (Fig. 1A). Unexpectedly, Pol II CTD phosphorylation occurred within three

minutes (Fig. 1A, compare lanes 1 and 2), which was our physical limit of isolating cells

during this time course. To determine which transcripts were generated during these

early quiescence exit events, we performed nascent RNA-seq using 4-thio-uracil (4tU)

to metabolically label new transcripts [41,42]. In agreement with the western-blot

analysis, we observed a high level of transcriptional activation within a few minutes of

nutrient repletion (Fig. 1B). Based on our western-blot result, the highest Pol II CTD

phosphorylation is observed ~ten minutes after refeeding. Consistent with this result, we

observed the highest level of nascent transcripts at the ten-minute time point, where

~1000 mRNAs (20% of the genome) were statistically significantly increased compared

to the zero-minute time point (Fig.1B, Fig.1—supplement 1A). Given how quickly Pol II

was phosphorylated and transcripts were generated, we sought to determine if high

levels of Pol II were already bound to the early exit genes in the quiescent state, as was

observed previously in a heterogenous population of stationary phase cells [43]. To this

end, we performed spike-in-normalized ChIP-seq analysis of Pol II in quiescent cells

and at several time points following refeeding (Fig. 1C, Fig. 1—supplement 1B). Low

Pol II occupancy levels (compare heatmaps 1 and 5) were detected in quiescent cells,

which agrees with our western blot and RNA-seq analyses and previously published

literature [6–8]. This implied that Pol II is not paused (Fig. 1C, compare heatmaps 1 and

2) in quiescent cells, and suggested that Pol II needs to be recruited *de novo* for rapid

initiation and elongation. In support of this conclusion, we detected only low levels of the

pre-initiation complex subunit TFIIB bound to genes in quiescent cells, which increased

~3-fold by five minutes of exit (Fig. 1—supplement 1C).

Highlighting the high level of transcription occurring in the first ten minutes of

quiescence exit, we observed a drop-off in Pol II occupancy levels around the first G2/M

phase (240 minutes) (Fig. 1C-D, Fig. 1—supplement 1D). Indeed, when the data were

sorted into k-means clusters across the time course, we noticed that many of the genes

expressed in the 240-minute time point were similar, but still not identical, to those

expressed in log cells, suggesting a recovery to log-like gene expression profile takes

hours post refeeding (Fig. 1C, compare columns 4 and 5, Fig. 1D). There was a ~1.7-

fold increase in overall Pol II occupancy in the 10-minute time point relative to that of log

cells (Fig. 1D, Fig.1—supplement 1B). Together, these results demonstrate transcription

activates extremely rapidly and robustly in response to nutrient repletion.

Chromatin bears hallmarks of repression during early quiescent exit time points

Given the exceptionally high transcriptional response during the first ten minutes of

quiescence exit, we wondered whether chromatin changes reflected hypertranscription.

To this end, we performed ChIP-seq analysis of H3 to measure nucleosome occupancy

levels genome wide over time. Global H3 patterns during the early exit time points,

especially at the 5-minute time point, were more similar to that of the quiescent state

than to the 240-minute time point (Fig. 2A, compare columns 1-3), despite higher

transcription levels. The most striking changes in histone occupancy during the early

time-points were within NDRs, where the pattern at the 10-minute timepoint resembles

the 240-minute time point (Fig. 2A, B). However, the H3 profiles outside of NDRs (Fig.

2A, compare column 1-3 and 4 to the right of NDR, and Fig.2B) remain similar to that of

quiescent state during the early stage of quiescent exit. In addition to nucleosome

occupancy, we tested nucleosome positioning using MNase-seq analysis where

nucleosomes with 80% of the digested chromatin is represented by mononucleosomes.

Globally, nucleosome positions were stable across the early exit time points (Fig. 2C).

We next tested if a burst of histone acetylation occurred during these early exit

time points to help overcome the repressive quiescent chromatin environment. To test

this, we performed ChIP-seq analysis of H4ac using an antibody that recognizes penta-

acetylated H4. Similar to nucleosome occupancy and positions, a modest increase in

histone H4 acetylation occurred, but the levels did not reflect that of log cells (Fig. 2D,

E). This suggests that, while there was a strong transcriptional response during

refeeding, histone acetylation was delayed. This is consistent with a previous study of a

mixed population of saturated cultures where histone acetylation was found to occur

later in exit[44]. Together, our results are in agreement with a recent study

demonstrating that histone acetylation takes place mostly as a consequence of

transcription [45].

To assess a biological readout of the repressive chromatin environment, we

turned to phenotypic analysis of TFIIS disruption. TFIIS is a general elongation factor

that rescues stalled Pol II; and nucleosomal barriers have been shown to increase

stalled Pol II [46]. Given that Pol II stalling is common across the genome [47], it is

paradoxical that the gene encoding TFIIS is not essential for viability in actively dividing

cells, and its deletion does not cause strong growth defects [48]. Since Pol II must

achieve a high level of transcription in the repressive chromatin environment during

early quiescence exit, we hypothesized that TFIIS may play more critical roles during

this period than during log culture. Indeed, in the absence of TFIIS (*dst1Δ*), quiescent

yeast cells exhibited defects in cell cycle re-entry, where cells lacking TFIIS stall at the

first G1 during exit, which is not the case during the mitotic cell cycle (Fig. 2F, Fig. 2—

supplement 1B). These results collectively revealed that the chromatin environment

remains repressive during early quiescence exit.

In quiescence, RSC re-localizes to NDRs of genes expressed in exit

Given the modest changes in chromatin at most genes during the early stage of

quiescence exit (Fig. 2), we wondered whether MNase-sensitive or "fragile"

nucleosomes were present at the promoters of rapidly induced genes in quiescence and

were removed in early exit. Thus, we performed a weaker (low) MNase digestion (10%

mononucleosomes) (Fig. 3A) and compared it to the stronger (high) MNase digestion

(80% mononucleosomes) (Fig. 3B). Supporting our hypothesis, comparing the weaker

MNase digest to the stronger MNase digest revealed that ~1000 genes have fragile

nucleosomes in quiescent cells, which are reduced during exit (Fig. 3A). Additionally,

we noticed that the MNase-sensitivity of the +1 and +2 nucleosomes increased at the

10-minute time point, likely coinciding with transcription.

It has been recently suggested that that the ATP-dependent chromatin remodeler

RSC can remove fragile nucleosomes from promoters to activate transcription [26].

Additionally, it was proposed that RSC-bound nucleosomes are remodeling

intermediates that render such nucleosomes more MNase-sensitive [38]. Thus, RSC

was a strong candidate for regulating rapid transcription activation during quiescence

exit. We performed ChIP-seq analysis of the RSC catalytic subunit Sth1 in quiescent

cells. In quiescence, Sth1 exhibited a striking difference in binding pattern compared to

log cells (Fig. 3C, D). Sth1 bound to ~80% of NDRs at gene promoters in quiescent

cells (Fig. 3E, Figure 3—supplement 1A). This result was distinct from log cells, where

RSC was reported to occupy the widest NDRs but otherwise bind the -1, +1, and +2

nucleosomes for most highly expressed genes (Fig. 3C) [28,26,38]. The RSC binding

pattern in quiescent cells instead mirrored a recently described binding pattern in heat

shock, where RSC and other transcription regulators *transiently* relocate to the NDRs

[49]. In contrast to the heat shock response, however, we observed a stable, strong

binding pattern of RSC in NDRs regardless of NDR width (Fig. 3E). Another obvious

distinction of RSC binding patterns between log and quiescence was observed at tRNA

genes (Fig. 3F). RSC's role at tRNA expression has been well-studied in log cells [50–

52]. In quiescence, RSC was occluded from tRNAs genes. Whereas upon exit, RSC

rapidly targeted tRNAs, mimicking the log pattern. Together these data suggest that

RSC adopts a quiescence-specific binding profile, one in which RSC is bound to NDRs

more broadly across the genome.

We next sought to gain insight into how quiescent RSC occupancy patterns

might predict Pol II occupancy during exit. To this end, we compared localization of

RSC and Pol II in quiescence and exit. We first found that the presence of RSC at

NDRs in quiescent cells and strong transcription in exiting cells co-localized (Fig. 3—

supplement 1A). Next, we examined RSC occupancy changes during quiescence exit at

Pol II-transcribed genes. During quiescence exit, RSC began to move out of NDRs and

into gene bodies as transcription increased (Fig. 3G). These results suggested that

RSC facilitates transcriptional activation upon exit and raised the possibility that RSC

binding in NDRs may be a mechanism for cells to prepare for quiescence exit.

RSC depletion causes quiescent exit defects and global Pol II occupancy

reduction during quiescence exit

To test the requirement of RSC in quiescence exit, we simultaneously depleted two

essential subunits of the RSC complex, Sth1 and Sfh1, using the auxin degron system

[53], during quiescence entry (see methods; Figure 3—supplement 1B). Depletion of

these subunits throughout the exit process (hereafter "-RSC") caused a dramatic defect

in cell cycle progression upon quiescence exit, where the cells exhibited strong delays

in exiting the first G1 stage (Figure 4A). This result contrasted with that in cycling cells,

where *rsc* mutants or conditional alleles cause G2/M arrest [54].

To determine the impact of RSC depletion on hypertranscription during

quiescence exit, we performed Pol II ChIP-seq analysis on cells exiting quiescence. In

the presence of RSC, Pol II levels peaked at 10 minutes and substantially decreased at

30 minutes after the exit (Fig. 4B, compare columns 3 and 4). As is the case in log

cultures [50,55,56], Pol II occupancy decreased in the absence of an intact RSC

complex in Q-cells and upon nutrient repletion thereafter (Fig. 4B). Pol II occupancy did

eventually increase over time in the RSC-depleted samples. However, even after 30-

minutes, Pol II did not reach the peak level of occupancy seen at the 10-minute mark in

the +RSC condition (Fig. 4B, compare heatmaps 3 and 8, and 4C). This suggests that

the defect in Pol II occupancy during quiescence exit was not solely due to slower

kinetics.

As shown earlier in Figure 3G, we observed RSC leaving the NDRs and moving

into gene bodies during quiescence exit. Therefore, we examined the impact of RSC

depletion on nucleosome occupancy and positioning. H3 ChIP-seq showed that RSC is

required for removal of histones within NDRs (Fig. 4D), which is consistent with RSC's

role as the "NDR creator" [24]. Together, these data provide mechanistic explanations

for how RSC facilitates Pol II loading during early stages of quiescence exit.

RSC is required for Pol II passage through gene bodies

Given that RSC moves from NDRs into gene bodies during quiescence exit (Fig. 3G),

we next tested whether RSC could aid transcription after initiation. To this end, we

selected ~2000 genes where RSC moved toward gene bodies and examined RSC

localization at the 10-minute time point of quiescent exit. This analysis showed uniform

movement of RSC from NDR into gene bodies (Fig. 5A). We next tested whether this

RSC movement is dependent on Pol II transcription. To this end, we performed Sth1

ChIP-seq analyses during quiescence exit in the presence of a transcription inhibitor

1,10-phenanthroline (Fig. 5B, Pol II control in Fig. 5—supplement 1A). This experiment

demonstrated that the movement of RSC from NDRs into gene bodies was strongly

inhibited by 1,10-phenanthroline, establishing that RSC re-localization during quiescent

exit is dependent on Pol II transcription.

Co-transcriptional movement of RSC into gene bodies suggested a possibility

that RSC may help Pol II passage through gene bodies. To test this, we determined the

effects of RSC depletion on Pol II localization during early time points of quiescent exit.

Fig. 5C and D show that RSC depletion affects Pol II localization in at least two ways

during early quiescent exit. First, consistent with Fig 4B, the robust increase in the

amount of Pol II over genes is strongly decreased upon RSC depletion. In addition,

upon RSC depletion, Pol II sharply accumulates at TSSs at the 5-minute mark, which

continued to the 10-minute mark. In sharp contrast, Pol II accumulates at slightly more

downstream at the 5-minute mark and moves mostly to downstream regions at the 10-

minute time point in the presence of RSC. We also noticed a pile-up of Pol II at the 3'-

end of genes at the 5-minute timepoint upon RSC depletion (Fig. 5C). This is in

agreement with the possibility that RSC may be involved in proper transcription

termination [57]. At these loci, NDRs are relatively shallow in quiescence but histone

density rapidly decreases upon quiescence exit in the presence of RSC (Fig. 5E, Fig.

5—supplement 1B). In the absence of RSC at these sites, however, histone density is

unexpectedly lower at NDR in quiescence but does not change during quiescent exit

(Fig. 5F, Fig. 5—supplement 1B), suggesting defective chromatin structure at and

downstream of the NDR. Together, these results are consistent with the notion that co-

transcriptional movement of RSC facilitates passage of Pol II through nucleosomes

immediately downstream of TSSs through chromatin regulation.

RSC suppresses abnormal upstream transcription initiation

The fact that Pol II accumulated upstream of TSSs at the 5-minute mark upon RSC

depletion (Fig. 5C) suggested possible defects in transcription start site selection. To

test this possibility, we examined the 4tU-seq profiles of a subset of RSC targets (1426

genes) in which there appeared to be an enrichment of RNA signal directly upstream

and downstream of TSSs. We took the $\log_2$ ratio of RNA signal in the depleted condition

versus the non-depleted condition at the ten-minute time point (Fig. 6A). We sorted the

genes using k-means clusters and took into account RSC binding when determining

gene sets to analyze.

This analysis revealed that upon RSC depletion a large number of genes (~1400)

exhibited increased nascent sense-strand RNA signals starting upstream of their normal

TSSs, demonstrating wide-spread defects in TSS selection. Examination of individual

loci revealed that, in addition to filling of an NDR at the normal TSSs, an NDR is created

upstream, which overlaps with ectopic transcription observed at an upstream TSS (see

Fig. 6B for an example). These results suggest that RSC facilitates selection of accurate

transcription initiation sites through proper NDR formation upstream of protein coding

genes during the burst of transcription during quiescence exit. This is likely a

quiescence-specific function of RSC, or a result of the robust hypertranscription event

during exit, as depletion of Sth1 in cycling cells mostly repressed transcription initiation

with relatively few new upstream transcription start sites [55,56].

RSC is required for suppression of anti-sense transcripts during quiescence exit

Given the robust transcriptional response during the early minutes of quiescence exit

(Fig. 1), we examined whether aberrant transcripts might also arise during quiescence

exit when RSC was depleted. Indeed, in many cases we found antisense transcripts

arising in the absence of RSC. We found ~900 RSC targets that had generally reduced

sense transcript levels with strongly upregulated cognate antisense transcripts (Class I),

and ~600 genes (Class II) with only modest changes in both sense and anti-sense

transcript levels upon RSC depletion (Fig. 7A). Chromatin analyses of individual Class I

loci revealed that RSC depletion caused narrower NDRs upstream of the sense TSS

(see Fig. 7B for an example). In contrast, NDRs for sense transcripts remained largely

open at Class II genes (Fig. 7B). Both classes have RSC bound at the promoters of the

sense genes in quiescence, with slightly higher RSC binding in the class I genes (Fig.

7C). Strikingly, nucleosome positioning was heavily impacted in the Class I set of genes

upon RSC depletion in the sense direction, where NDRs became more resistant to

MNase and nucleosomes in gene bodies were shifted toward the 5'-ends of genes. This

was in contrast to that of Class II where NDRs were largely open (Fig. 7D). Consistent

with the MNase-mapping data, nucleosome occupancy at NDRs and in gene bodies are

much more strongly affected by RSC depletion at Class I genes than Class II genes

(Fig. 7E). It is likely that Class II genes overcome the absence of RSC by having more

"fragile" nucleosomes that can be readily removed by general regulatory factors [26].

These results collectively showed that chromatin structure at the Class I genes is

especially dependent on RSC. In both classes of genes, RSC signals and RSC-

dependent chromatin changes are not apparent around the start sites of anti-sense

transcripts. Therefore, suppression of anti-sense transcripts is unlikely to be a direct role

for RSC. Instead, it is likely that both Class I and II genes, especially the former, have

an intrinsic property to allow anti-sense transcription to occur when not properly

regulated, and RSC is targeted to them to ensure sense transcription takes place

through formation of proper NDRs.

Discussion

In this report we have shown that there is a rapid and robust transcriptional response

during the very early minutes of quiescence exit (Fig. 8A). This response is greatly

dependent on the chromatin remodeling enzyme RSC. We found that RSC promotes

transcription at the right place and time in four different ways: 1) RSC promotes

transcription initiation by creating NDRs in quiescence and maintaining them during exit

(Fig. 8B). 2) RSC moves into gene bodies and helps Pol II transcribe past the +1

nucleosome (Fig. 8C). 3) RSC maintains proper NDR locations to allow for accurate

transcription start site selection (Fig. 8D). 4) RSC suppresses cryptic antisense

transcription via generating NDRs at the cognate sense genes (Fig. 8E). Together, our

results suggest that the massive transcriptional response requires highly accurate

nucleosome positioning to allow for cells to exit from the quiescent state.

Quiescent yeast must downregulate their transcriptional program and generate a

repressive chromatin environment in order to survive harsh conditions for extended time

periods [10,6,58,59]. How, then, do cells rapidly escape the quiescent state when

conditions are favorable? In this study, we show that there is an incredibly strong

transcriptional response to nutrient repletion after quiescence, notwithstanding a

relatively repressive chromatin environment that persists until the first G2/M phase after

quiescence. Indeed, we identified a previously unidentified phenotype for the deletion of

the gene encoding yeast TFIIS, *dst1Δ*. High numbers of stalled Pol II are present in

cycling cells [47] despite the little impact of deleting *DST1* on cycling cell growth. We

speculate cells exiting quiescence may rely more heavily on TFIIS to transcribe through

repressive chromatin [60,61].

During quiescence, RSC relocates to NDRs upstream of Pol II transcribed genes

that are transcribed in exit. Although RSC binds and regulates chromatin around Pol III

genes [27,50], RSC is depleted at tRNA genes in quiescence and only returns during

quiescence exit, further supporting the notion that RSC is globally re-targeted in

quiescence. This is distinct from the transient NDR-relocalization observed in heat

shock [49], as what we observed in quiescence was a sustained and rather stable

localization. How RSC binds to these new locations in quiescence is unknown. Given

the distinct structure of quiescent chromatin there are several, non-mutually exclusive,

explanations for RSC's binding pattern in quiescence. 1) The genome is hypoacetylated

and thus RSC can no longer bind to acetylated nucleosomes in quiescence via its

bromodomains [19]. However, given the highly robust response to refeeding, RSC

activity must be poised to be active in this state. An intriguing possibility could be that

histone acetylation inhibits RSC activity to some extent as was recently reported *in vitro*

[62]. This would be consistent with the rapid changes in nucleosome positioning at

many genes during quiescence exit in the absence of high levels of histone acetylation.

2) Recent structural studies have shown that the nucleosome acidic patch is in direct

contact with subunits of the RSC complex [63–66]. If the acidic patch is occluded by

hypoacetylated H4 tails in quiescence for example [12,67–70], it is possible that RSC

can no longer interact with this region of the nucleosome, rendering its binding abilities

different in quiescence. Finally, 3) a lack of Pol II activity in quiescent cells could prevent

RSC from moving out of NDRs and into gene bodies. Indeed, transcription appears to

play a prominent role in RSC localization: RSC moves into gene bodies during

transcription activation and this movement is blocked when transcription is inhibited, as

we have reported above. It is likely that a combination of transcription and histone

acetylation helps pull RSC into gene bodies, given recent work showing that acetylation

is a consequence of transcription [45]. In a separate study, we recently found that the

SWI/SNF remodeling enzyme promotes transcription of a subset of hypoacetylated

genes during quiescence entry, implying a specialized transcription regulation program

for essential genes in the wake of widespread transcriptional shutdown [58]. In cycling

cells, it was recently shown that RSC and SWI/SNF cooperate at a subset of genes

[71]. Our results suggested that cooperation between the two SWI/SNF class

remodeling factors may also occur during quiescence entry.

Consistent with co-transcriptional re-localization, our data suggest RSC plays an

active role in helping Pol II transcribe past the +1 nucleosome in addition to initiating

transcription. Supporting this idea was our observation of a subset of genes where RSC

depletion caused a Pol II enrichment around the +1 nucleosome. Previous reports

showed that RSC can bind gene bodies and impact elongating and terminating Pol II

[31,57]; and one study showed interactions between the Rsc4 subunit and all three RNA

polymerases [30]. An intriguing possibility could be that RSC directly interacts with Pol II

to facilitate transcription past the first few nucleosomes.

The transcriptional response during quiescent exit was dampened by depleting

the essential chromatin remodeler, RSC, but it did not diminish completely. Pol II

occupancy was globally decreased ~2-fold at the 10-minute time point in RSC-depleted

cells. However, at ~900 genes we found that while sense transcription was reduced,

antisense transcripts were generated. This was largely due to a nearby NDR

susceptible to transcription initiation that could be co-opted for antisense transcription.

The mechanism that allows for this cryptic transcription is still unknown. Chromatin

remodeling enzymes are vastly important for repressing antisense lncRNAs [72].

Different chromatin remodeling enzymes function to repress lncRNA transcripts in

cycling cells, including RSC [73–75]. We speculate RSC is particularly suitable to

regulate global transcriptome during quiescence exit due to its high abundance, which

allows it to function through multiple mechanisms. The mouse embryonic stem cell-

specific BAF complex was also recently shown to globally repress lncRNA expression

[76]. This raises the possibility that some of our observations in yeast quiescent cells

could be conserved in mammalian quiescent cells. Given the robust transcriptional

response that occurs during quiescence exit, it is likely that chromatin structure is

crucial for maintaining the quality of the transcriptome. Indeed, we noted cases where

transcription occurred upstream of the canonical TSS when an NDR was not generated,

highlighting the defects in Pol II initiation and start site selection due to chromatin

defects in the absence of RSC. Hypertranscription events similar to the one observed

during quiescence exit occur throughout all organisms, particularly during development

[40]. Therefore, it is quite possible that we will see similar, multifaceted roles for RSC

homologues or other abundant chromatin remodeling factors in facilitating proper

hypertranscription in many other systems.

Materials and Methods

Yeast strains, yeast growth media, quiescent cell purification, and exit time

courses

The *S. cerevisiae* strains used in this study are listed in Supplementary Table S1 and

are isogenic to the strain W303-1a with a correction for the mutant *rad5* allele in the

original W303-1a [77]. Yeast transformations were performed as previously described

[78]. All cells were grown in YPD medium (2% Bacto Peptone, 1% yeast extract, 2%

glucose). We note that quiescent (Q) yeast need to be grown in YPD using "fresh"

(within ~three months) yeast extract as a source. To purify Q cells, liquid YPD cultures

were inoculated with a single colony into liquid cultures (colonies were no older than

one week). Yeast cells were grown in Erlenmeyer flasks ten times the liquid volume for

seven days at 30°C and shaking at 180 RPM. Q cells were purified by percoll gradient

centrifugation as previously described [11]. Briefly, percoll was diluted 9:1 with 1.5 M

NaCl into 25-mL Kimble tubes and centrifuged at 10,000 RPM for 15-minutes at 4°C.

Seven-day cultures were pelleted, washed with ddH2O, resuspended in 1 mL of

ddH2O, and gently pipetted over a pre-mixed percoll gradient. 400 OD_{660} were pipetted

onto a 25-mL gradient. Gradients with loaded cells were centrifuged for one hour at

1000 RPM, 4°C. The upper, non-quiescent cell population and the middle, ~8 mL

fraction, were carefully discarded via pipetting. The remaining volume was washed

twice with ddH2O in a 50 mL conical tube at 3,000 RPM, 10 minutes each.

422   Q exit experiments were performed as follows: Q cells were harvested and

added to YPD to 1 OD_{660}/mL. Cells were grown at 25°C to slow the kinetics for

feasibility. For ChIP-seq and MNase-seq experiments, cells were grown to the

appropriate time and then crosslinked for 20 minutes (described in more detail in the

sections below).

Depletion of RSC subunits, Sth1 and Sfh1

The yeast strains YTT 7222 and 7224 were grown in 5-mL overnight YPD cultures, back

diluted for four doublings, and inoculated to 0.002 OD_{660} into the appropriate YPD

volume for a given experiment. Cells were grown for 16 hours and monitored for

glucose exhaustion using glucose strips. Six hours after glucose exhaustion, 1mg/mL of

Indole-3-acetic acid (Sigma, I3750-5G-A) was added, in powder form, to the culture. Q

cells were purified as described above and depletion efficiency was determined by

western blot analysis (Supplementary Figure 1A).

Western Blot Analysis

Yeast cells were lysed by bead beating in trichloroacetic acid (TCA), as previously

described [79]. Proteins were resolved on 8% polyacrylamide gels and transferred to

nitrocellulose membranes. Membranes were incubated with primary antibodies: anti-

Rpb3 (Biolegend, 665003 1:1000 dilution), anti-Ser5p (Active Motif, 61085 1:1000

dilution), anti-Ser2p (Active Motif, 61083, 1:1000 dilution), and anti-HSV (Sigma, 1:500).

Following primary incubation, membranes were incubated with either anti-mouse or

anti-rabbit secondary antibodies (Licor, 1:10000). Protein signals were visualized by the

Odyssey CLx scanner.

ChIP-seq

100 OD_{660} U of cells were crosslinked and sonicated in biological duplicate using the

protocol described in [80]. Proteins were immunoprecipitated from 1 µg chromatin and 1

µL of anti-H3 (Abcam, 1791) conjugated to 20 µl protein G magnetic beads (Invitrogen,

10004D) per reaction. For Pol II ChIPs, we used an antibody against the Rpb3 subunit

(2 µl per reaction, Biolegend 665004) conjugated to 20 µl protein G magnetic beads

(Invitrogen, 10004D). For Sth1 ChIP experiments we used an antibody against the Flag-

epitope tag, FLAG M2 mouse monoclonal (Sigma Aldrich, F1804) and conjugated to 20

µl protein G beads (Invitrogen, 10004D) Libraries were generated using the Ovation

Ultralow v2 kit (NuGEN/Tecan, 0344) and subjected to 50-bp single-end sequencing on

an Illumina HiSeq 2500 at the Fred Hutchinson Cancer Research Center genomics

facility. We used bowtie2 to align raw reads to the sacCer3 reference genome [81].

Reads were then filtered using SAMtools [82]. Bigwig files of input-normalized ChIP-seq

data were generated from the filtered bam files using deepTools2 [83] and dividing the

IP data by the input data. Matrices for metaplots were generated in deepTools2 using

the annotation file from [84].

MNase-seq

Cell growth and crosslinking was done in the same fashion as in ChIP-seq experiments.

Generally, we followed the protocol in [80], with changes described here. Cells were

spheroplasted using 10 mg zymolyase (100T, AMSBIO, 120493-1) per 100 OD_{660} cells.

For Q cells, zymolyase treatment could take up to two hours. We monitored the cells via

microscopy and stopped the spheroplasting step when ~80% of the cells were

spheroplasted. MNase digestion was performed as described in [80]. High digests (80%

mononucleosomes) required 50U of micrococcal nuclease (Worthington, LS004798)

and for the low digests, chromatin was treated with 10 U of MNase. From this step,

chromatin was reverse crosslinked as described in [80]. Following reverse crosslinking,

RNase, and proteinase-K digestion, DNA was phenochloroform-extracted. Any large,

uncut genomic DNA species was separated out using Ampure beads (Beckman).

Sequencing libraries were generated from the purified DNA using the Ovation Ultralow

v2 kit (NuGEN, 0344). Libraries were subjected to 50-bp paired-end sequencing on an

Illumina HiSeq 2500 at the Fred Hutchinson Cancer Research Center genomics facility.

We used bowtie2 to align raw reads to the sacCer3 genome and filtered reads using

SAMtools as described above for ChIP-seq analysis. Bigwig files of input-normalized

ChIP-seq data were similarly generated from the filtered bam files using deepTools2

and the MNase option to center the reads around nucleosome dyads. Data represented

in the paper were filtered to mononucleosome sizes using deepTools2.

Nascent RNA-seq

Generally, nascent RNA-seq experiments were performed as described in

[85,42]. For the 0-minute and 5-minute samples, we added 100 and 50 OD_{660} of Q cells,

respectively, to YPD containing 5 mM 4-thiouracil (Sigma, 440736-1G). Cells were

incubated with 4tU for 5 minutes before pelleting (one minute, 3500 RPM) and flash

frozen in liquid nitrogen. For the 10-minute time points, 50 OD units of quiescent cells

were released into YPD for 5 minutes before an additional 5-minute incubation with 4tU

at a final concentration of 5 mM. All time points were labeled with 4tU for a total of 5

minutes before pelleting and freezing. Total RNA was isolated using Ambion's RiboPure

Yeast Kit (Thermo, AM1926). *S. cerevisiae cells* were lysed in the presence of

Kluvomyces lactis (*K. lactis*) cells in a 100:1 mixture. RNA was treated with DNAseI

according to the TURBO DNase kit (Thermo, AM2238). 40 ug RNA was then

biotinylated with MTSEA biotin-XX (diluted in 20% DMF) at a final concentration of 16.4

uM in 20mM HEPES pH 7.4 and 1 mM EDTA at room temperature for 30 minutes.

Unreacted MTS-biotin was removed from samples by PCI extraction and resuspended

in 100 uL nuclease-free water. Strepavidin beads (Invitrogen 65001) were washed with

high-salt wash buffer (100 mM Tris, 10 mM EDTA, 1 M NaCl, 0.05% Tween-20) and

blocked for one hour in high-salt wash buffer containing 40 ng/uL glycogen. 40 uL of

streptavidin beads were added to the RNA samples and incubated for 15 minutes at

room temperature. Beads were washed three times in 1 mL high salt wash buffer and

eluted for 15 minutes at room temperature in 50 uL streptavidin elution buffer (100 mM

DTT, 20 mM HEPES, 2.7, 1 mM EDTA, 100 mM NaCl, 0.05% Tween-20). The resulting

RNA was then purified and concentrated using the Qiagen miRNeasy kit (#217084).

Libraries were prepared from 5 ng of RNA using the Ovation SoLo kit (NuGEN/Tecan,

custom AnyDeplete; contact Tecan for ordering this kit for yeast). Libraries were

subjected to 50-bp paired-end sequencing on an Illumina HiSeq 2500 at the Fred

Hutchinson Cancer Research Center genomics facility. We used bowtie2 to align raw

reads to the sacCer3 and *K. lactis* (Ensembl ASM251v1) genomes and filtered reads

using SAMtools as described above for ChIP-seq analysis. Differential expression

analysis was performed using DESeq2 [86]

Data Availability

517 All sequencing data are uploading on the NCBI Gene Expression Omnibus under the

518 accession number GSE166789.

Acknowledgements

We are grateful to members of the Tsukiyama lab and Pravrutha Raman for helpful

comments and critical reading of this manuscript. We thank Sarah Hainer and Felix

Mueller-Planitz for advice on MNase-seq experiments. We thank Benjamin Martin, Rafal

Donczew and Sandipan Brahma for advice and feedback. We thank Mitchell Ellison and

Alex Francette for advice about analyzing nascent RNA-seq data. TT was supported by

the National Institutes of Health (R01 GM111428 and R35GM139429). CEC was

supported by the National Cancer Institute (T32CA009657) and National Institutes of

Health (F32GM131554).

FIGURE 1

Figure 1. Rapid hypertranscription occurs upon nutrient repletion of quiescent cells

(A) Western blots were probed with antibodies to detect Ser5p and Ser2p of the CTD of Rpb1 subunit of Pol II. An antibody against the Rpb3 subunit of Pol II was used as a loading control. **(B)** Strand-specific 4tU-seq analysis. "+" indicates Watson strand and "-" indicates Click strand. **(C)** Pol II ChIP-seq analysis. Heatmaps show k-means clusters of 6030 genes. Genes are linked across the heatmaps. **(D)** Metaplots of ChIP-seq data shown in (C) without k-means clustering.

Figure 1—supplement 1

(A) Volcano plot of nascent transcripts comparing significant changes in expression using a 2-fold cut off. **(B)** Boxplots illustrating the difference in Pol II ChIP-seq signals across genes. Log_2 ratio values were subtracted (ex: Q log_2 values were subtracted from 10 min. log_2 values). **(C)** TFIIB ChIP-seq analysis in Q cells and exit time points. Genes are linked across the time points and are aligned to TSS. **(D)** DNA content FACS analysis indicating cell cycle progress during Q exit.

FIGURE 2

Figure 2. Repressive chromatin persists during early quiescence exit
(A, B) ChIP-seq of total H3 in quiescent cells and exit time points sorted into quartiles based on NDR width. **(C)** MNase-seq analysis of 6030 genes in Q (pink line), Log (black line), and Q-exit time points 5 minutes (light grey line) and 10 minutes (dark grey line). **(D, E)** ChIP-seq analysis of penta-acetylated H4 (H4ac) in Q and Log cells and exit time points. Genes are separated as in (B). **(F)** DNA content FACS analysis following Q exit in WT and a TFIIS-absent strain (*dst1Δ*).

FIGURE 3

Figure 3. MNase sensitivity and quiescence-specific RSC relocalization indicate remodeling activity required for early exit
(A) MNase-digested chromatin to 10% mononucleosomes (low digestion). **(B)** Metaplot of MNase-digested chromatin to 80% mononucleosomes (high digestion) in Q and 10-minute time points. **(C,D)** ChIP-seq of the catalytic RSC subunit in quiescent and log cells at Pol II-transcribed genes. **(E)** ChIP-seq analysis of RSC shown across quartiles based on MNase-seq determined NDR width. **(F)** ChIP-seq of RSC at tRNA genes. **(G)** ChIP-seq of RSC and Pol II comparing RSC movement with Pol II into gene bodies.

Figure 3—supplement 1
(A) ChIP-seq analysis of RSC and Pol II using antibodies against Flag-tagged Sth1 and Rpb3, respectively. Genes are sorted into k-means clustered and are linked across the different ChIPs. **(B)** Western blot analysis of RSC depletion. Both Sth1 and Sfh1 contain C-terminal HSV and AID tags for detection and depletion using IAA. Western blot was probed with an antibody recognizing the HSV epitope tag and Rpb3 (Pol II subunit) as a loading control. The addition of IAA is indicated by – or +.

FIGURE 4

Figure 4. RSC is required for normal quiescence exit and hypertranscription upon nutrient repletion

(A) DNA content FACS analysis indicating cell cycle progression during Q exit in the presence (+) or absence (-) of RSC. **(B)** ChIP-seq analysis of Pol II across time in the presence or absence of RSC. Genes are sorted the same in all heatmaps. **(C)** Example tracks of data shown in **(B)** with RSC ChIP-seq in Q cells added. **(D)** H3 ChIP-seq sorted by NDR width (as determined by MNase-seq experiments).

Figure 5. RSC depletion causes severe Pol II mislocalization defects during quiescence exit.
C) ChIP-seq of RSC in Q and 10-minute time points. Genes are linked. **(B)**. ChIP-seq of RSC at 10-minutes of exit in the presence and absence of the transcription inhibitor 1,10-phenanthroline. **(C, D)** ChIP-seq of RSC and Pol II during exit. **(E-F)** H3 ChIP-seq in quiescence and during exit in the presence and absence of RSC.

Figure 5—supplement 1
(A) ChIP-seq analysis of Pol II in the absence and presence of the transcription inhibitor 1,10-phenanthroline. **(B)** MNase-seq analysis assessing differences in MNase sensitivity in Q and ten-minutes for cells with and without RSC. The +2 nucleosome MNase-digestion differences are highlighted by the pink arrows.

FIGURE 6

Figure 6. RSC depletion causes upstream transcription relative to canonical TSS
(A) Heatmap showing the Log_2 ratio of nascent sense transcripts in RSC-depleted versus non-depleted cells. Shown are 1426 genes that have upregulated transcripts at the 5'-ends of genes in the sense direction and have RSC ChIP signals. **(B)** Example gene of aberrant upstream transcript. Arrows direct to defects: blue arrow points to loss of NDR, yellow arrow points to gain of NDR, and pink arrow points to upstream RNA signal.

Figure 7. Aberrant antisense transcription arises when chromatin around sense transcripts is abrogated in the absence of RSC (A) Heatmaps of the Log$_2$ ratio of nascent RNAs that are RSC targets and give rise to antisense transcripts (cluster I, ~890 genes) when RSC is depleted and those where antisense transcripts are not made when the sense transcript NDR is unchanged (cluster II, ~600 genes). **(B)** Browser tracks of 4tU-seq and MNase-seq data. Boxes highlight defects (orange box) in NDRs and where antisense transcription arises from co-opted, intact NDRs (grey box). **(C)** ChIP-seq of RSC in quiescent cells and during exit at both cluster I and II. **(D)** MNase-seq at the 10-minute time point. (**E**.) 10-minute time point during exit of H3 ChIP-seq separated into the two clusters, I and II.

Figure 8. RSC safeguards the quiescent genome from aberrant transcription
In quiescent cells, RSC binds to NDRs upstream of Pol II transcribed genes. Upon quiescence exit, RSC shifts the +1 nucleosome to allow for Pol II occupancy and traverses into gene bodies **(A)**. In the absence of RSC NDRs are globally narrower and transcription initiation is blocked **(B).** At a subset of genes, RSC is required for efficient Pol II passage past the +1 nucleosome **(C)** and prevent upstream TSS selection **(D).** NDRs that are open despite RSC depletion become cryptic promoters and are utilized by transcription machinery to generate aberrant lncRNAs and antisense transcripts **(E)**.

References

1. Rando OJ, Winston F. Chromatin and Transcription in Yeast. Genetics. 2012;190: 351–387. doi:10.1534/genetics.111.132266

2. Sagot I, Laporte D. The cell biology of quiescent yeast – a diversity of individual scenarios. J Cell Sci. 2019;132: jcs213025. doi:10.1242/jcs.213025

3. Rittershaus ESC, Baek S-H, Sassetti CM. The Normalcy of Dormancy: Common Themes in Microbial Quiescence. Cell Host & Microbe. 2013;13: 643–651. doi:10.1016/j.chom.2013.05.012

4. Cheung TH, Rando TA. Molecular regulation of stem cell quiescence. Nat Rev Mol Cell Biol. 2013;14: 329–340. doi:10.1038/nrm3591

5. Tümpel S, Rudolph KL. Quiescence: Good and Bad of Stem Cell Aging. Trends in Cell Biology. 2019;29: 672–685. doi:10.1016/j.tcb.2019.05.002

6. McKnight JN, Boerma JW, Breeden LL, Tsukiyama T. Global Promoter Targeting of a Conserved Lysine Deacetylase for Transcriptional Shutoff during Quiescence Entry. Molecular Cell. 2015;59: 732–743. doi:10.1016/j.molcel.2015.07.014

7. Young CP, Hillyer C, Hokamp K, Fitzpatrick DJ, Konstantinov NK, Welty JS, et al. Distinct histone methylation and transcription profiles are established during the development of cellular quiescence in yeast. BMC Genomics. 2017;18: 107. doi:10.1186/s12864-017-3509-9

8. Swygert SG, Kim S, Wu X, Fu T, Hsieh T-H, Rando OJ, et al. Condensin-Dependent Chromatin Compaction Represses Transcription Globally during Quiescence. Mol Cell. 2019;73: 533-546.e4. doi:10.1016/j.molcel.2018.11.020

9. Chapman NM, Boothby MR, Chi H. Metabolic coordination of T cell quiescence and activation. Nat Rev Immunol. 2020;20: 55–70. doi:10.1038/s41577-019-0203-y

10. Gray JV, Petsko GA, Johnston GC, Ringe D, Singer RA, Werner-Washburne M. "Sleeping Beauty": Quiescence in Saccharomyces cerevisiae. MMBR. 2004;68: 187–206. doi:10.1128/MMBR.68.2.187-206.2004

11. Allen C, Büttner S, Aragon AD, Thomas JA, Meirelles O, Jaetao JE, et al. Isolation of quiescent and nonquiescent cells from yeast stationary-phase cultures. J Cell Biol. 2006;174: 89–100. doi:10.1083/jcb.200604072

12. Luger K, Mäder AW, Richmond RK, Sargent DF, Richmond TJ. Crystal structure of the nucleosome core particle at 2.8 Å resolution. Nature. 1997;389: 251–260. doi:10.1038/38444

13.   Lorch Y, LaPointe JW, Kornberg RD. Nucleosomes inhibit the initiation of transcription but allow chain elongation with the displacement of histones. Cell. 1987;49: 203–210. doi:10.1016/0092-8674(87)90561-7

14.   Hainer SJ, Kaplan CD. Specialized RSC: Substrate Specificities for a Conserved Chromatin Remodeler. BioEssays. 2020;42: 2000002. doi:10.1002/bies.202000002

15.   Cairns BR, Lorch Y, Li Y, Zhang M, Lacomis L, Erdjument-Bromage H, et al. RSC, an Essential, Abundant Chromatin-Remodeling Complex. Cell. 1996;87: 1249–1260. doi:10.1016/S0092-8674(00)81820-6

16.   Du J, Nasir I, Benton BK, Kladde MP, Laurent BC. Sth1p, a Saccharomyces cerevisiae Snf2p/Swi2p homolog, is an essential ATPase in RSC and differs from Snf/Swi in its interactions with histones and chromatin-associated proteins. Genetics. 1998;150: 987–1005.

17.   Saha A. Chromatin remodeling by RSC involves ATP-dependent DNA translocation. Genes & Development. 2002;16: 2120–2134. doi:10.1101/gad.995002

18.   Zofall M, Persinger J, Kassabov SR, Bartholomew B. Chromatin remodeling by ISW2 and SWI/SNF requires DNA translocation inside the nucleosome. Nat Struct Mol Biol. 2006;13: 339–346. doi:10.1038/nsmb1071

19.   Kasten M, Szerlong H, Erdjument-Bromage H, Tempst P, Werner M, Cairns BR. Tandem bromodomains in the chromatin remodeler RSC recognize acetylated histone H3 Lys14. EMBO J. 2004;23: 1348–1359. doi:10.1038/sj.emboj.7600143

20.   Angus-Hill ML, Schlichter A, Roberts D, Erdjument-Bromage H, Tempst P, Cairns BR. A Rsc3/Rsc30 Zinc Cluster Dimer Reveals Novel Roles for the Chromatin Remodeler RSC in Gene Expression and Cell Cycle Control. Molecular Cell. 2001;7: 741–751. doi:10.1016/S1097-2765(01)00219-2

21.   Varela I, Tarpey P, Raine K, Huang D, Ong CK, Stephens P, et al. Exome sequencing identifies frequent mutation of the SWI/SNF complex gene PBRM1 in renal carcinoma. Nature. 2011;469: 539–542. doi:10.1038/nature09639

22.   Kadoch C, Hargreaves DC, Hodges C, Elias L, Ho L, Ranish J, et al. Proteomic and bioinformatic analysis of mammalian SWI/SNF complexes identifies extensive roles in human malignancy. Nat Genet. 2013;45: 592–601. doi:10.1038/ng.2628

23.   Badis G, Chan ET, van Bakel H, Pena-Castillo L, Tillo D, Tsui K, et al. A Library of Yeast Transcription Factor Motifs Reveals a Widespread Function for Rsc3 in Targeting Nucleosome Exclusion at Promoters. Molecular Cell. 2008;32: 878–887. doi:10.1016/j.molcel.2008.11.020

24.   Hartley PD, Madhani HD. Mechanisms that Specify Promoter Nucleosome Location and Identity. Cell. 2009;137: 445–458. doi:10.1016/j.cell.2009.02.043

25.   Prajapati HK, Ocampo J, Clark DJ. Interplay among ATP-Dependent Chromatin Remodelers Determines Chromatin Organisation in Yeast. Biology. 2020;9: 190. doi:10.3390/biology9080190

26.   Kubik S, O'Duibhir E, de Jonge WJ, Mattarocci S, Albert B, Falcone J-L, et al. Sequence-Directed Action of RSC Remodeler and General Regulatory Factors Modulates +1 Nucleosome Position to Facilitate Transcription. Molecular Cell. 2018;71: 89-102.e5. doi:10.1016/j.molcel.2018.05.030

27.   Ng HH. Genome-wide location and regulated recruitment of the RSC nucleosome-remodeling complex. Genes & Development. 2002;16: 806–819. doi:10.1101/gad.978902

28.   Yen K, Vinayachandran V, Batta K, Koerber RT, Pugh BF. Genome-wide Nucleosome Specificity and Directionality of Chromatin Remodelers. Cell. 2012;149: 1461–1473. doi:10.1016/j.cell.2012.04.036

29.   Ramachandran S, Zentner GE, Henikoff S. Asymmetric nucleosomes flank promoters in the budding yeast genome. Genome Res. 2015;25: 381–390. doi:10.1101/gr.182618.114

30.   Soutourina J, Bordas-Le Floch V, Gendrel G, Flores A, Ducrot C, Dumay-Odelot H, et al. Rsc4 Connects the Chromatin Remodeler RSC to RNA Polymerases. MCB. 2006;26: 4920–4933. doi:10.1128/MCB.00415-06

31.   Spain MM, Ansari SA, Pathak R, Palumbo MJ, Morse RH, Govind CK. The RSC Complex Localizes to Coding Sequences to Regulate Pol II and Histone Occupancy. Molecular Cell. 2014;56: 653–666. doi:10.1016/j.molcel.2014.10.002

32.   Biernat E, Kinney J, Dunlap K, Rizza C, Govind CK. The RSC complex remodels nucleosomes in transcribed coding sequences and promotes transcription in *Saccharomyces cerevisiae*. Genomics; 2020 Mar. doi:10.1101/2020.03.11.987974

33.   Xi Y, Yao J, Chen R, Li W, He X. Nucleosome fragility reveals novel functional states of chromatin and poises genes for activation. Genome Research. 2011;21: 718–724. doi:10.1101/gr.117101.110

34.   Knight B, Kubik S, Ghosh B, Bruzzone MJ, Geertz M, Martin V, et al. Two distinct promoter architectures centered on dynamic nucleosomes control ribosomal protein gene transcription. Genes Dev. 2014;28: 1695–1709. doi:10.1101/gad.244434.114

35.   Teif VB, Beshnova DA, Vainshtein Y, Marth C, Mallm J-P, Höfer T, et al. Nucleosome repositioning links DNA (de)methylation and differential CTCF binding during stem cell development. Genome Res. 2014;24: 1285–1295. doi:10.1101/gr.164418.113

36.   Kubik S, Bruzzone MJ, Jacquet P, Falcone J-L, Rougemont J, Shore D. Nucleosome Stability Distinguishes Two Different Promoter Types at All Protein-

Coding Genes in Yeast. Molecular Cell. 2015;60: 422–434. doi:10.1016/j.molcel.2015.10.002

37.   Floer M, Wang X, Prabhu V, Berrozpe G, Narayan S, Spagna D, et al. A RSC/Nucleosome Complex Determines Chromatin Architecture and Facilitates Activator Binding. Cell. 2010;141: 407–418. doi:10.1016/j.cell.2010.03.048

38.   Brahma S, Henikoff S. RSC-Associated Subnucleosomes Define MNase-Sensitive Promoters in Yeast. Molecular Cell. 2019;73: 238-249.e3. doi:10.1016/j.molcel.2018.10.046

39.   Schlichter A, Kasten MM, Parnell TJ, Cairns BR. Specialization of the chromatin remodeler RSC to mobilize partially-unwrapped nucleosomes. eLife. 2020;9: e58130. doi:10.7554/eLife.58130

40.   Percharde M, Bulut-Karslioglu A, Ramalho-Santos M. Hypertranscription in Development, Stem Cells, and Regeneration. Developmental Cell. 2017;40: 9–21. doi:10.1016/j.devcel.2016.11.010

41.   Miller C, Schwalb B, Maier K, Schulz D, Dümcke S, Zacher B, et al. Dynamic transcriptome analysis measures rates of mRNA synthesis and decay in yeast. Mol Syst Biol. 2011;7: 458. doi:10.1038/msb.2010.112

42.   Duffy EE, Rutenberg-Schoenberg M, Stark CD, Kitchen RR, Gerstein MB, Simon MD. Tracking Distinct RNA Populations Using Efficient and Reversible Covalent Chemistry. Molecular Cell. 2015;59: 858–866. doi:10.1016/j.molcel.2015.07.023

43.   Radonjic M, Andrau J-C, Lijnzaad P, Kemmeren P, Kockelkorn TTJP, van Leenen D, et al. Genome-Wide Analyses Reveal RNA Polymerase II Located Upstream of Genes Poised for Rapid Response upon S. cerevisiae Stationary Phase Exit. Molecular Cell. 2005;18: 171–183. doi:10.1016/j.molcel.2005.03.010

44.   Mews P, Zee BM, Liu S, Donahue G, Garcia BA, Berger SL. Histone Methylation Has Dynamics Distinct from Those of Histone Acetylation in Cell Cycle Reentry from Quiescence. Molecular and Cellular Biology. 2014;34: 3968–3980. doi:10.1128/MCB.00763-14

45.   Martin BJE, Brind'Amour J, Kuzmin A, Jensen KN, Liu ZC, Lorincz M, et al. Transcription shapes genome-wide histone acetylation patterns. Nat Commun. 2021;12: 210. doi:10.1038/s41467-020-20543-z

46.   Noe Gonzalez M, Blears D, Svejstrup JQ. Causes and consequences of RNA polymerase II stalling during transcript elongation. Nat Rev Mol Cell Biol. 2021;22: 3–21. doi:10.1038/s41580-020-00308-8

47.   Churchman LS, Weissman JS. Nascent transcript sequencing visualizes transcription at nucleotide resolution. Nature. 2011;469: 368–373. doi:10.1038/nature09652

48.	Hubert JC, Guyonvarch A, Kammerer B, Exinger F, Liljelund P, Lacroute F. Complete sequence of a eukaryotic regulatory gene. The EMBO Journal. 1983;2: 2071–2073. doi:10.1002/j.1460-2075.1983.tb01702.x

49.	Vinayachandran V, Reja R, Rossi MJ, Park B, Rieber L, Mittal C, et al. Widespread and precise reprogramming of yeast protein–genome interactions in response to heat shock. Genome Res. 2018;28: 357–366. doi:10.1101/gr.226761.117

50.	Parnell TJ, Huff JT, Cairns BR. RSC regulates nucleosome positioning at Pol II genes and density at Pol III genes. EMBO J. 2008;27: 100–110. doi:10.1038/sj.emboj.7601946

51.	Mahapatra S, Dewari PS, Bhardwaj A, Bhargava P. Yeast H2A.Z, FACT complex and RSC regulate transcription of tRNA gene through differential dynamics of flanking nucleosomes. Nucleic Acids Research. 2011;39: 4023–4034. doi:10.1093/nar/gkq1286

52.	Kumar Y, Bhargava P. A unique nucleosome arrangement, maintained actively by chromatin remodelers facilitates transcription of yeast tRNA genes. BMC Genomics. 2013;14: 402. doi:10.1186/1471-2164-14-402

53.	Nishimura K, Kanemaki MT. Rapid Depletion of Budding Yeast Proteins via the Fusion of an Auxin-Inducible Degron (AID). Current Protocols in Cell Biology. 2014;64. doi:10.1002/0471143030.cb2009s64

54.	Tsuchiya E, Uno M, Kiguchi A, Masuoka K, Kanemori Y, Okabe S, et al. The Saccharomyces cerevisiae NPS1 gene, a novel CDC gene which encodes a 160 kDa nuclear protein involved in G2 phase control. EMBO J. 1992;11: 4017–4026.

55.	Kubik S, Bruzzone MJ, Challal D, Dreos R, Mattarocci S, Bucher P, et al. Opposing chromatin remodelers control transcription initiation frequency and start site selection. Nat Struct Mol Biol. 2019;26: 744–754. doi:10.1038/s41594-019-0273-3

56.	Klein-Brill A, Joseph-Strauss D, Appleboim A, Friedman N. Dynamics of Chromatin and Transcription during Transient Depletion of the RSC Chromatin Remodeling Complex. Cell Reports. 2019;26: 279-292.e5. doi:10.1016/j.celrep.2018.12.020

57.	Ocampo J, Chereji RV, Eriksson PR, Clark DJ. Contrasting roles of the RSC and ISW1/CHD1 chromatin remodelers in RNA polymerase II elongation and termination. Genome Res. 2019;29: 407–417. doi:10.1101/gr.242032.118

58.	Spain MM, Braceros KCA, Tsukiyama T, Fred Hutchinson Cancer Research Center, Division of Basic Sciences. SWI/SNF coordinates transcriptional activation through Rpd3-mediated histone hypoacetylation during quiescence entry. Molecular Biology; 2018 Sep. doi:10.1101/426288

59.   Poramba-Liyanage DW, Korthout T, Cucinotta CE, van Kruijsbergen I, van Welsem T, El Atmioui D, et al. Inhibition of transcription leads to rewiring of locus-specific chromatin proteomes. Genome Res. 2020;30: 635–646. doi:10.1101/gr.256255.119

60.   Kireeva ML, Hancock B, Cremona GH, Walter W, Studitsky VM, Kashlev M. Nature of the Nucleosomal Barrier to RNA Polymerase II. Molecular Cell. 2005;18: 97–108. doi:10.1016/j.molcel.2005.02.027

61.   Bondarenko VA, Steele LM, Újvári A, Gaykalova DA, Kulaeva OI, Polikanov YS, et al. Nucleosomes Can Form a Polar Barrier to Transcript Elongation by RNA Polymerase II. Molecular Cell. 2006;24: 469–479. doi:10.1016/j.molcel.2006.09.009

62.   Lorch Y, Maier-Davis B, Kornberg RD. Histone Acetylation Inhibits RSC and Stabilizes the +1 Nucleosome. Molecular Cell. 2018;72: 594-600.e2. doi:10.1016/j.molcel.2018.09.030

63.   Ye Y, Wu H, Chen K, Clapier CR, Verma N, Zhang W, et al. Structure of the RSC complex bound to the nucleosome. Science. 2019;366: 838–843. doi:10.1126/science.aay0033

64.   Patel AB, Moore CM, Greber BJ, Luo J, Zukin SA, Ranish J, et al. Architecture of the chromatin remodeler RSC and insights into its nucleosome engagement. eLife. 2019;8: e54449. doi:10.7554/eLife.54449

65.   Wagner FR, Dienemann C, Wang H, Stützer A, Tegunov D, Urlaub H, et al. Structure of SWI/SNF chromatin remodeller RSC bound to a nucleosome. Nature. 2020;579: 448–451. doi:10.1038/s41586-020-2088-0

66.   Baker RW, Reimer JM, Carman PJ, Turegun B, Arakawa T, Dominguez R, et al. Structural insights into assembly and function of the RSC chromatin remodeling complex. Nat Struct Mol Biol. 2021;28: 71–80. doi:10.1038/s41594-020-00528-8

67.   Shogren-Knaak M. Histone H4-K16 Acetylation Controls Chromatin Structure and Protein Interactions. Science. 2006;311: 844–847. doi:10.1126/science.1124000

68.   Robinson PJJ, An W, Routh A, Martino F, Chapman L, Roeder RG, et al. 30 nm Chromatin Fibre Decompaction Requires both H4-K16 Acetylation and Linker Histone Eviction. Journal of Molecular Biology. 2008;381: 816–825. doi:10.1016/j.jmb.2008.04.050

69.   Allahverdi A, Yang R, Korolev N, Fan Y, Davey CA, Liu C-F, et al. The effects of histone H4 tail acetylations on cation-induced chromatin folding and self-association. Nucleic Acids Research. 2011;39: 1680–1691. doi:10.1093/nar/gkq900

70.   Swygert SG, Lin D, Portillo-Ledesma S, Lin P-Y, Hunt DR, Kao C-F, et al. Chromatin Fiber Folding Represses Transcription and Loop Extrusion in Quiescent Cells. Molecular Biology; 2020 Nov. doi:10.1101/2020.11.24.396713

71.   Rawal Y, Chereji RV, Qiu H, Ananthakrishnan S, Govind CK, Clark DJ, et al. SWI/SNF and RSC cooperate to reposition and evict promoter nucleosomes at highly expressed genes in yeast. Genes Dev. 2018;32: 695–710. doi:10.1101/gad.312850.118

72.   Han P, Chang C-P. Long non-coding RNA and chromatin remodeling. RNA Biology. 2015;12: 1094–1098. doi:10.1080/15476286.2015.1063770

73.   Alcid EA, Tsukiyama T. ATP-dependent chromatin remodeling shapes the long noncoding RNA landscape. Genes Dev. 2014;28: 2348–2360. doi:10.1101/gad.250902.114

74.   Marquardt S, Escalante-Chong R, Pho N, Wang J, Churchman LS, Springer M, et al. A Chromatin-Based Mechanism for Limiting Divergent Noncoding Transcription. Cell. 2014;157: 1712–1723. doi:10.1016/j.cell.2014.04.036

75.   Gill JK, Maffioletti A, García-Molinero V, Stutz F, Soudet J. Fine Chromatin-Driven Mechanism of Transcription Interference by Antisense Noncoding Transcription. Cell Reports. 2020;31: 107612. doi:10.1016/j.celrep.2020.107612

76.   Hainer SJ, Gu W, Carone BR, Landry BD, Rando OJ, Mello CC, et al. Suppression of pervasive noncoding transcription in embryonic stem cells by esBAF. Genes Dev. 2015;29: 362–378. doi:10.1101/gad.253534.114

77.   Thomas BJ, Rothstein R. The genetic control of direct-repeat recombination in Saccharomyces: the effect of rad52 and rad1 on mitotic recombination at GAL10, a transcriptionally regulated gene. Genetics. 1989;123: 725–738.

78.   Ausubel FM, B.R. Kingston RE, Moore DD, Seidman JG, Smith JA, Struhl K. Current Protocols in Molecular Biology. 1988; NYWiley-Interscience.

79.   Cox JS, Chapman RE, Walter P. The unfolded protein response coordinates the production of endoplasmic reticulum protein and endoplasmic reticulum membrane. MBoC. 1997;8: 1805–1814. doi:10.1091/mbc.8.9.1805

80.   Rodriguez J, McKnight JN, Tsukiyama T. Genome-Wide Analysis of Nucleosome Positions, Occupancy, and Accessibility in Yeast: Nucleosome Mapping, High-Resolution Histone ChIP, and NCAM. Current Protocols in Molecular Biology. 2014;108. doi:10.1002/0471142727.mb2128s108

81.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012;9: 357–359. doi:10.1038/nmeth.1923

82.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map format and SAMtools. Bioinformatics. 2009;25: 2078–2079. doi:10.1093/bioinformatics/btp352

83.   Ramírez F, Dündar F, Diehl S, Grüning BA, Manke T. deepTools: a flexible platform for exploring deep-sequencing data. Nucleic Acids Research. 2014;42: W187–W191. doi:10.1093/nar/gku365

84.   Xu Y, Bernecky C, Lee C-T, Maier KC, Schwalb B, Tegunov D, et al. Architecture of the RNA polymerase II-Paf1C-TFIIS transcription elongation complex. Nat Commun. 2017;8: 15741. doi:10.1038/ncomms15741

85.   Bonnet J, Wang C-Y, Baptista T, Vincent SD, Hsiao W-C, Stierle M, et al. The SAGA coactivator complex acts on the whole transcribed genome and is required for RNA polymerase II transcription. Genes Dev. 2014;28: 1999–2012. doi:10.1101/gad.250225.114

86.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. 2014;15: 550. doi:10.1186/s13059-014-0550-8