Cell lineage as a predictor of immune response in neuroblastoma

Satyaki Sengupta[1,2,*], Sanjukta Das[1,2,*], Angela Crespo[2,3], Brian Miller[4,5,6,7], Bandana Sharma[1,2],

Shupei Zhang[8], Ruben Dries[1,2], Hao Huang[1,2], Malgorzata Krajewska[1,2], David N. Debruyne[1,2],

Luigi Soriano[1], Malkiel A. Cohen[8], Rogier Versteeg[9], Rudolf Jaenisch[8,10], Stefani Spranger[10,11],

Judy Lieberman[2,3], Rani E. George[1,2,12]

[1]Department of Pediatric Oncology, Dana-Farber Cancer Institute, Boston, MA 02215, USA

[2]Department of Pediatrics, Harvard Medical School, Boston, MA 02215, USA

[3]Program in Cellular and Molecular Medicine, Boston Children's Hospital, Boston, MA, USA.

[4]Broad Institute of MIT and Harvard, Cambridge, Massachusetts 02142, USA.

[5]Department of Medical Oncology, Dana-Farber Cancer Institute and Harvard Medical School,

Boston, Massachusetts 02215 USA.

[6]Department of Immunology, Blavatnik Institute, Harvard Medical School, Boston, MA, USA

[7]Evergrande Center for Immunological Diseases, Harvard Medical School and Brigham and

Women's Hospital, Boston, MA, USA.

[8]Whitehead Institute for Biomedical Research, 455 Main Street, Cambridge, MA 02142, USA

[9]Department of Oncogenomics, Amsterdam UMC University of Amsterdam, Meibergdreef 9,

1105 AZ Amsterdam, The Netherlands

[10]Department of Biology, Massachusetts Institute of Technology, Cambridge, MA 02142,

USA

[11]Koch Institute for Integrative Cancer Research, Cambridge, MA 02142, USA

*These authors contributed equally

[12]Corresponding author: rani_george@dfci.harvard.edu

SUMMARY

Immunotherapy for patients with neuroblastoma has met with limited success, partly due to an incomplete understanding of the mechanisms underlying immune responsiveness in this clinically and genetically heterogenic tumor. Here, we undertook an unbiased analysis using dimension reduction and UMAP visualization of transcriptional signatures derived from 498 primary neuroblastoma tumors. Four distinct clusters based on differentially expressed genes emerged, of which one, representing about 30% and comprising mainly of *MYCN*-nonamplified tumors, was notable for the high expression of genes associated with both immune response activation and suppression. This capacity to elicit a productive immune response resided exclusively in tumors with dominant populations of undifferentiated, neural crest-like or mesenchymal cells; by contrast, tumors comprising primarily of committed, adrenergic neuron-like cells were less immunogenic. Mesenchymal neuroblastoma cells were enriched for innate and adaptive immune gene signatures, demonstrated engagement with cytotoxic T and natural killer cells, and induced immune cell infiltration in an immunocompetent mouse model. Transcriptional or targeted therapy-induced reprogramming of adrenergic cells to the mesenchymal state led to reactivation of tumor cell-intrinsic immune genes. Key immune response genes in adrenergic tumor cells were found to be epigenetically silenced by the PRC2 complex, and such repression could be relieved by either mesenchymal cell state reprogramming or EZH2 inhibition, leading to increased activation of natural killer cells by the tumor cells. These data identify cell lineage as a major determinant of the immunogenic potential in neuroblastoma that could be used to stratify patients who are most likely to benefit from immunotherapy.

INTRODUCTION

The anti-disialoganglioside GD2 monoclonal antibody dinutuximab has significantly improved

event free survival rates in neuroblastoma[1]. Derived from the developing neural crest, this

common solid tumor of childhood manifests as an extracranial mass arising in the adrenal medulla

or sympathetic ganglia. Approximately half of all patients have high-risk features associated with

a poor outcome - age >18 months, distant metastases and unfavorable histologic and genetic

factors including amplification of the *MYCN* oncogene[2]. The success of anti-GD2 therapy that

relies on immune cell-mediated cytotoxicity suggests that patients with neuroblastoma would

benefit from other forms of immunotherapy; however, treatment results with use of cytotoxic CD8+

T lymphocytes directed against neuroblastoma antigens[3], adoptive transfer of chimeric antigen

receptor (CAR)-modified T cells[4-6] or checkpoint inhibition[7,8] have been suboptimal.

Major impediments to the effectiveness of immunotherapy in neuroblastoma are the immune

evasion tactics deployed by the tumor cells as well as the tumor microenvironment (TME)[9]. These

include downregulation of major histocompatibility complex (MHC) class I molecules and defects

in antigen-processing machinery (APM) that render neuroblastoma cells resistant to T-cell-

mediated cytotoxicity[10-12], downregulation of cell-surface ligands required for natural killer (NK)

cell receptor activation[13], upregulation of checkpoint proteins that exert a protective role from NK

cell-mediated lysis[14], inefficient homing of cytotoxic T-cells to the tumor site[15] or tumor cell

overexpression of the leukocyte surface antigen CD47, which enables avoidance of macrophage-

mediated phagocytosis[16]. Moreover, infiltration of suppressive immune cells such as T regulatory

cells[17], tumor-associated macrophages (TAMs)[18,19], myeloid-derived suppressor cells[20] and

secreted immunosuppressive factors such as TGF-β, contribute to the generation of a TME that

hinders an effective immune response and further dampens the effects of adoptive cell

therapies[21].

Amplification of the MYCN oncogene poses another distinct challenge to immunotherapy in

neuroblastoma. This transcription factor is amplified in approximately 50% of high-risk cases and

is associated with aggressive disease and a poor clinical outcome[22,23]. *MYCN*-amplified tumors

consistently evade immune destruction by downregulating MHC class I molecules[10] and are

associated with poor infiltration of cytotoxic CD8[+] T cells[24,25] and reduced expression of NK cell

ligands[26]. Interestingly, approximately half of high-risk neuroblastomas do not express amplified

MYCN, and their capacity to induce a productive immune response remains unclear. In a recent

study that analyzed the immune gene expression programs associated with *MYCN*-nonamplified

tumors from high-risk patients, tumors with low as well as high functional tumor *MYCN* signatures

were observed to have significantly higher levels of NK and CD8+ T-cell infiltrates compared to

MYCN-amplified tumors; although, somewhat counterintuitively, these findings translated into a

better outcome only in patients with high *MYCN* tumor signatures[25,27].

Thus, although many of the mechanisms of immune evasion in neuroblastoma are known,

further understanding of the tumor-host interaction will be crucial to enhancing the ability of

immunotherapy to target and eliminate tumor-initiating and propagating cell populations.

Especially challenging is the genetic and biologic heterogeneity of this tumor which makes it

difficult to identify factors that consistently indicate the likelihood of an effective immune response

and hence identify patients who are most likely to benefit from this form of therapy. Thus, we

undertook an unbiased analysis of gene expression signatures across diverse clinical subtypes

of primary tumors and identify tumor cell state as an important predictor of immune

responsiveness in neuroblastoma.

RESULTS

A subset of primary neuroblastomas express markers of a productive immune response

To determine whether neuroblastomas are capable of eliciting a productive immune response,

we first examined bulk RNA-sequencing data from 498 well-annotated primary human tumors

representing diverse clinical and genetic subtypes (SEQC-498; GSE49711; **Supplementary Fig.**

1a) to quantify tumor-to-tumor gene expression variability and cluster tumor types based on gene

expression profiles (see also Methods). In this unbiased analysis, all tumors within one cluster

would share similar gene expression profiles, while being dissimilar to those of tumors within other

clusters. Specifically, we first identified the top 5000 highly variably expressed genes within this

dataset based on the premise that these would be most likely to contribute to distinct molecular

subtypes[28,29] (**Supplementary Fig. 1b; Supplementary Table 1)**. The data were dimensionally

reduced using principal component analysis (PCA) and the top 20 leading principal components

selected for clustering analysis **(Supplementary Fig. 1c)**. Four distinct clusters were identified

and visualized using 2D-Uniform Manifold Approximation and Projection (UMAP), a non-linear

dimension-reduction tool[30,31] (**Fig. 1a**). To explore the transcriptional differences between the

clusters, we identified the differentially expressed genes (DEGs) in each cluster and noted that

tumors in cluster 1 (C1; n = 103), termed *Hi-MYCN,* were enriched for MYCN target genes

involved in cell proliferation and biosynthesis, and comprised 20% of the tumor set (**Fig. 1a-c;**

Supplementary Fig. 1d; Supplementary Table 2). Not surprisingly, this cluster segregated with

MYCN-amplified tumors in patients aged ≥ 18 months with stage 4 disease [according to the

international neuroblastoma staging system (INSS)][32] and annotated "high risk" status (based on

the Children's Oncology Group risk classification)[33] (**Fig. 1d; Supplementary Fig. 1e)**. The

remaining clusters consisted of *MYCN*-nonamplified tumors (**Fig. 1a**) of which, cluster 2 (C2, n =

241), or *neuronal*, made up the largest proportion of tumors, 48%, and comprised tumors that

were enriched for DEGs with roles in nervous system

Fig. 1.

Fig. 1. A subset of neuroblastomas exhibits a productive immune response. (a) Two-

dimensional UMAP representations of the gene expression profiles in 498 neuroblastoma (NB)

tumors. Each dot represents a tumor. The top 5000 highly variable genes were selected based

on the variance-stabilizing method[34] and the 20 significant principal components (PCs) selected

and processed in UMAP to generate four clusters representing four NB subtypes. The DEGs were

identified for each cluster using the receiver operating characteristics (ROC) curve to compare

one cluster with other three (log$_2$ FC > 0.25). **(b)** Gene ontology (GO) analysis of top DEGs in the

four clusters. **(c)** Heat map of expression values of 10 representative DEGs within each cluster.

Rows are z-score scaled average expression levels for each gene in all four clusters. **(d)** UMAP

visualization of the distribution of the indicated prognostic features in NB among the four different

clusters. **(e, f)** Heat map of z-score transformed log$_2$ normalized expression values of immune

activation **(e)** and evasion **(f)** genes in *MYCN*-nonamplified NBs (n=401). Tumors were ranked

based on increasing immune activation or evasion scores**.** Cluster annotations of the tumors are

indicated on the top horizontal bar.

Supplementary Fig. 1.

a

	MYCN (%)		Age (%)		Risk (%)		INSS stage (%)				
	Nonamp	Amp	< 18 mo.	≥ 18 mo.	Low	High	1	2	3	4	4S
Tumor (498)	401 (80.5)	92 (18.4)	300 (60.3)	198 (39.7)	322 (64.6)	176 (35.4)	121 (24.3)	78 (15.7)	63 (12.6)	183 (36.7)	53 (10.6)

Supplementary Fig. 1. Tumors within the immunogenic cluster express both immune

activation and evasion markers. **(a)** Distribution of the 498 primary NB tumors in the data set

(SEQC-498; GSE49711) within the indicated prognostic categories. **(b)** Scatter plot of the

standardized variance in expression of all protein coding genes within 498 tumors. Red dots

indicate the top 5000 variably expressed genes. **(c)** Elbow plot representing the percentage

variance for the top 20 principal components, PCs. **(d)** Violin plots showing the expression of

representative marker genes across the four clusters. **(e)** Stacked bar plots showing the

distribution of tumors within the defined prognostic features within each cluster. Amp, amplified;

Nonamp, nonamplified. **(f)** Two-dimensional UMAP representations of the gene expression

profiles in 394 NB tumors (GSE120572). Each dot represents a tumor. The top 3000 highly

variable genes were selected based on the variance-stabilizing method [34] and the 20 significant

principal components (PCs) selected and processed in UMAP to generate three clusters

representing three NB subtypes. The DEGs were identified for each cluster using the receiver

operating characteristics (ROC) curve to compare one cluster with other two ($\log_2$ FC > 0.25). **(g)**

Heat map of expression values of 10 representative DEGs within each cluster. Rows are z-score

scaled average expression levels for each gene in all three clusters.

Supplementary Fig. 2.

Supplementary Fig. 2. The immunogenic tumors are associated with markers of both

immune activation and evasion. (a, b) Box plots comparing immune activation **(a)** and evasion

(b) scores within the four clusters from the SEQC-498 tumor data set. All box plots are defined

by center lines (medians), box limits (25th and 75th percentiles), whiskers (minima and maxima;

the smallest and largest data range). Significance was determined by the Wilcoxon rank-sum test.

(c) UMAP visualization of the distribution of IA and IE scores among the three tumor clusters

derived from the 394 NBs in the GSE120572 dataset. Color bar represents normalized z-scores.

Values <2.5 and >2.5 were set to -2.5 and +2.5 respectively, to reduce the effects of extreme

outliers.

development **Fig. 1a-c; Supplementary Fig. 1d).** Cluster 3 (C3, n = 140), accounting for 28%

of the tumors, was enriched for tumors whose DEGs were involved in immune function, such as

interferon-gamma (IFN-γ) response and T cell inflammation and activation, and hence were

designated *immunogenic* **(Fig. 1a-c; Supplementary Fig. 1d)** . Cluster 4 (C4; n=14; 3%) was

clearly distinct from the other three clusters, and largely consisted of the spontaneously

regressing stage 4S tumors that were predominantly enriched for genes involved in fatty acid and

cholesterol homeostasis and hence were termed *metabolic* **(Fig. 1a-c; Supplementary Fig. 1d)**.

The neuronal and metabolic tumors arose predominantly in children <18 months of age, were of

stages 1-3 and 4S, while the tumors within the immunogenic cluster were associated with patient

age ≥18 months and metastatic disease (n= 66; 47%) **(Fig. 1d; Supplementary Fig. 1e)**. Thus,

our DEG-based analysis of almost 500 tumors categorized neuroblastoma into four largely distinct

groups that included a distinct subset, accounting for approximately one-third of the entire cohort,

whose gene expression profiles were closely linked to immune responsiveness. To ensure that

these results were not confined to one data set, we analyzed an independent data set of 394

tumors (GSE120572) using similar clustering methods. This cohort also segregated into *Hi-*

MYCN, neuronal and *immunogenic* clusters, again denoting the presence of immune response

gene expression in a subset of primary neuroblastomas, the majority of which lack MYCN

amplification **(Supplementary Fig. 1f, g)**.

To pursue the immune genes that were differentially enriched in the immunogenic cluster, we

generated an immune activation (IA) score based on the relative expression of a curated set of

41 genes known to have major roles in tumor cell-intrinsic immune functions, such as regulation

of MHC expression, antigen processing and presentation, NK cell recognition and T and NK cell

infiltration **(Supplementary Table 3)**. After assigning an IA score to each of the *MYCN-*

nonamplified tumors (n = 401) in the SEQC-498 data set and arranging them in ascending order

(Supplementary Table 4), we observed that a significant number with the highest IA scores

predominantly fell within the immunogenic and metabolic clusters **(Fig. 1e; Supplementary Fig.**

2a), while those with intermediate or lower scores were associated with the neuronal and Hi-

MYCN clusters, respectively **(Fig. 1e; Supplementary Fig. 2a)**. Because a cytotoxic immune

response is generally accompanied by immune suppression or evasion[35,36], we determined

whether immune suppression was also represented in the *MYCN*-nonamplified tumors by ranking

them in ascending order of an immune evasion (IE) score based on the relative expression of 19

genes, most of which were markers of T-cell dysfunction **(Supplementary Table 3)**. Again, the

immunogenic tumor cluster had significantly higher IE scores compared with the neuronal and

metabolic clusters **(Fig. 1f; Supplementary Fig. 2b).** Moreover, we observed enrichment for IA

and IE scores in the immunogenic cluster in the additional data set (GSE120572)

(Supplementary Fig. 2c), thus strengthening our premise that these tumors maybe capable of

eliciting an immune response.

Consistent with the known poor immunogenicity of *MYCN*-amplified tumors[24,25], we also

observed that tumors within the Hi-MYCN cluster had, on the whole, the lowest IA and IE scores

(Supplementary Fig. 2a, b). Surprisingly, however, a small subset within this cluster had scores

that were comparable to the highly immunogenic tumors within the immunogenic cluster [13 of

103 (12.6%) above the median for immunogenic tumors] **(Supplementary Fig. 2a, b)**. Thus, while

the majority of neuroblastomas do not possess an immune response gene signature, a subset

has significantly increased expression of both immune activation and evasion markers, pointing

to their ability to induce an anti-tumor immune response.

The mesenchymal lineage is preferentially associated with immune response signatures

in neuroblastoma

Having identified subsets of neuroblastomas with the potential for immunogenicity, we next sought

a biomarker that might consolidate the complex interactions between the immune system and the

tumor. To this end, we performed a modular gene co-expression analysis of the 140

transcriptomes within the immunogenic cluster in the SEQC-498 data set to identify biologically

relevant pathways based on similar gene expression patterns. Using the CEMiTool (co-

expression modules identification tool) package (Russo et al., 2018), we identified five gene co-

expression modules (M1-M5) within the immunological cluster **(Supplementary Fig. 3a).** Among

these modules, M1, with the highest number of co-expressed genes, contained gene sets

enriched for epithelial to mesenchymal transition (EMT), inflammatory response, and interferon

signaling, suggesting an association between EMT and the preponderant representation of

immune marker genes within the immunogenic cluster **(Fig. 2a; Supplementary Fig. 3b**).

Furthermore, integration of the co-expression data in module M1 with protein–protein interaction

data from the STRING 11.0 database identified mesenchymal lineage and immune markers as

top regulatory hubs **(Fig. 2b)**, leading us to hypothesize that in neuroblastoma, tumor

immunogenicity could be determined by cell state.

Two independent groups [37,38] recently described two distinct cell states in neuroblastoma: a

differentiated sympathetic neuron-like adrenergic (ADR) phenotype, defined by lineage markers

including *PHOX2B, DBH,* and *TH*, and a mesenchymal (MES) phenotype, characterized as

"neural crest cell-like" (NCC), and expressing genes such as *PRRX1, FOSL1,* and *FOSL2*. To

test our prediction, we first quantified the adrenergic and mesenchymal identities of each tumor

in our cohort based on the expression levels of the lineage-specific genes in each cell state as

established by Groningen et al[38] (see Methods). We ensured that there was no overlap between

the 369-gene adrenergic signature and the genes that made up the IA data set and removed the

6 IA genes that were also present in the 485-gene mesenchymal signature. Next, we assigned

either an adrenergic (A-score) or a mesenchymal (M-score) score to each tumor within our four

previously identified clusters. This analysis revealed significant enrichment of the mesenchymal

cell state within the immunogenic and metabolic clusters **(Fig. 2c; Supplementary Fig. 3c)**.

Fig. 2.

Fig. 2. The mesenchymal cell state is associated with an immunogenic signature in NB. (a)

GO analysis of co-expressed genes associated with module M1 using the KEGG (Kyoto

encyclopedia of genes and genomes) database. The vertical dashed line indicates the adjusted

P-value of 0.05. **(b)** Gene network representing all possible interactions in module M1. The

topmost connected genes (hubs) are indicated. Hubs derived from module M1 are colored blue

(co-expression) and those from the STRING database are indicated in red (interaction). The size

of each node corresponds to the degree of interaction. **(c)** UMAP visualization of the distribution

of adrenergic (top) and mesenchymal scores (bottom) among the four tumor clusters. Color bar

represents normalized z-scores. Values <2.5 and >2.5 were set to -2.5 and +2.5 respectively, to

reduce the effects of extreme outliers. **(d)** Dot plots showing the distribution of *MYCN*-

nonamplified tumors (n = 400) within each of the clusters based on ranked M-A scores. *Left*,

Tumors from the upper (high M-A) and lower (low M-A) M-A score quartiles are shown (n = 200;

P < 0.01 for C3). *Right*, Representations based on the median M-A scores of the entire tumor

cohort (n = 400; *P* = 0.05 for C3). Fisher's exact test was used for both calculations. **(e)** *Left*,

Heatmap representation of the expression of tumor cell-intrinsic immune activation genes in

MYCN-amplified tumors (n=92). Samples are ranked by increasing M-A score. Log$_2$ gene

expression values were z-score transformed for heatmap visualization. *Right*, Violin plots of the

distribution of immune activation scores in the tumors on the left, classified either as adrenergic

or mesenchymal, based on the median M-A score. The box plots within the violin plots are defined

by center lines (medians), box limits (25th and 75th percentiles), whiskers (minima and maxima;

1.5X the interquartile range). Significance was determined by the two-sided Kolmogorov-Smirnov

(KS) test. APM, antigen processing machinery. **(f)** Violin plots comparing the quantitative scores

of the indicated immune cell signatures in 100 tumors from the upper (mesenchymal) and lower

(adrenergic) quartiles of the tumor M-A scores using the two-sided KS test. The box plots within

the violin plots are defined as in (D). **(g)** *Left*, Bar diagram comparing regression coefficient (*β*)

values derived from multivariate multiple regression model analysis of *MYCN*-nonamplified

tumors. *β*-coefficient values were compared between three predictors: MYC score, M-A score and

MYCN signature. IA, IE, T cell, cytotoxic cell and NK cell scores were used as response variables

to generate the model[39]. *Right,* Heat map of the *P*-values associated with the three predictors. **(h)**

Bar diagram comparing the CIBERSORT-estimated fractional content of the indicated tumor-

infiltrating leukocytes between *MYCN*-nonamplified adrenergic and mesenchymal tumors.

Adrenergic and mesenchymal tumors were assigned as in **(f)**. Data represent the means, n = 100

tumors, **P* < 0.05, ***P* < 0.01, ****P* < 0.001 two-tailed Welch's t-test.

Supplementary Fig. 3.

c

Lineage (no of genes)	Hi-MYCN (2859) (*P*-value)	Neuronal (1416) (*P*-value)	Immunogenic (2213) (*P*-value)	Metabolic (2043) (*P*-value)
Adrenergic (369)	121 (<0.001)	78 (<0.001)	16 (n.s.)	13 (n.s.)
Mesenchymal (479)	46 (n.s.)	31 (n.s.)	129 (<0.001)	156 (<0.001)

Supplementary Fig. 3. Cell lineage markers are significantly associated with immune gene

signatures in NB. (a) Bar plots representing the numbers of co-expressed genes within each

module. **(b)** Profile plots depicting the expression levels (y-axis) of individual genes (colored lines)

and their mean expression (black line) in 140 tumors (x-axis) from modules M2, M3, M4 and M5.

GO analysis of co-expressed genes associated with each module using the KEGG database is

appended below the respective profile plot. The vertical dashed line indicates the adjusted *P*-

value of 0.05. **(c)** Summary of the overlap between the DEGs associated with the four tumor

clusters and the adrenergic or mesenchymal signature genes as per Groningen et al., 2017.

Significance was determined by Fisher's exact test. **(d)** Scatter plot of the 498 primary NB tumors

ranked based on increasing M-A score. **(e)** Violin plots of the distribution of normalized expression

levels of MYC in the four tumor clusters. **(f)** Pearson correlation matrix showing pairwise

correlation values among the indicated parameters. The colors and sizes of the circles indicate

the correlation coefficient values, with the least (smaller, orange circles) to the most (larger, blue

circles) degree of association between the parameters shown. **(g)** Scatter plot of the correlation

between MYC expression and immune activation (blue dots) or evasion (yellow dots) scores in

MYCN-nonamplified tumors.

Supplementary Fig. 4.

Supplementary Fig. 4. The relative mesenchymal score (M-A score) is positively correlated

with an immunogenic signature. (a) UMAP visualization of the distribution of adrenergic,

mesenchymal, and M-A scores among the three tumor clusters derived from 394 NBs in the

GSE120572 dataset. Color bar represents normalized z-scores. Values <2.5 and >2.5 were set

to -2.5 and +2.5 respectively, to reduce the effects of extreme outliers. **(b)** Heat maps of the

indicated immune cell signatures in *MYCN*-nonamplified tumors, ranked by increasing M-A

scores. Log_2 gene expression values were z-score transformed for heatmap visualization. **(c, d)**

Heatmaps depicting the immune activation **(c)** and evasion **(d)** signatures in *MYCN*-nonamplified

tumors, ranked by increasing M-A score. Log_2 gene expression values were z-score transformed

for visualization. Violin plots comparing the distribution of immune activation **(c)** and evasion **(d)**

signatures in 100 tumors from upper (mesenchymal) and lower (adrenergic) quartiles of the M-A

score are shown next to the heatmaps. Significance determined by the two-sided KS test. Box

plots within the violin plots are defined by center lines (medians), box limits (25^{th} and 75^{th}

percentiles), whiskers (minima and maxima; 1.5X the interquartile range). **(e)** Bar graph

comparing the CIBERSORT-estimated fractional content of the indicated tumor-infiltrating

leukocytes in six tumors with the uppermost (mesenchymal) and lowermost (adrenergic) M-A

scores.

By contrast, the Hi-MYCN and neuronal clusters were enriched for the adrenergic cell state **(Fig.**

2c; Supplementary Fig. 3c).

Next, to identify predictors of immunogenicity among the *MYCN*-nonamplified tumors, we

calculated the relative mesenchymal score for each tumor by subtracting the adrenergic from the

mesenchymal score (M-A score). This resulted in a continuum of low to high M-A scores,

corresponding to a less mesenchymal to a more mesenchymal tumor state **(Supplementary Fig.**

3d). To determine whether these cell states had any effect on immune response in the *MYCN*-

nonamplified tumors, we determined whether the mesenchymal, adrenergic, or M-A scores

correlated with our previously defined immune activation and evasion scores **(Fig. 1e, f)**. We also

tested the effect of *MYC*, one of the top differentially expressed genes in the immunogenic cluster

(Supplementary Fig. 3e). *MYC* is overexpressed in approximately 10% of *MYCN*-nonamplified

neuroblastomas[40] and regulates the expression of cell-intrinsic immune evasion markers in

lymphoma[41]. Of these variables, the M-A score showed the strongest correlation with

immunogenicity, not only in terms of the immune activation score ($R = 0.71$), but also the immune

evasion score ($R = 0.51$) **(Supplementary Fig. 3f)**. MYC expression was only modestly correlated

with immune evasion and activation scores ($R = 0.43$; $R = 0.38$, respectively) **(Supplementary**

Fig. 3g). With no overlap between the lineage marker and immune response gene sets, these

results suggest that the relative abundance of a mesenchymal signature (M-A score) is a better

predictor of immune response than individual adrenergic or mesenchymal signatures.   In

agreement, tumors with high M-A scores were represented at a significantly higher proportion

within the immunogeni*c* cluster compared to tumors with low M-A scores **(Fig. 2d),** a result that

was recapitulated in our second data set, (GSE120572) **(Supplementary Fig. 4a)**. Finally, our

finding of the subset of tumors marked by the relatively high expression of immune activation and

evasion genes within the Hi-MYCN cluster (**Supplementary Fig. 2a, b)** prompted us to further

evaluate the cell states of these tumors. Ranking these tumors based on increasing M-A scores

revealed a positive relationship between cell-intrinsic immunogenicity and the mesenchymal state

(Fig. 2e), suggesting that similar to our results in *MYCN*-nonamplified tumors, the presence of

immune gene expression in *MYCN*-amplified tumors is significantly correlated with the

mesenchymal phenotype.

The preferential overexpression in the mesenchymal phenotype of tumor cell-intrinsic genes

that induce a positive immune response (IA score), as well as its correlation with transcripts that

suppress the immune response (IE score), suggested that these tumors may support increased

immune cell infiltration**.** To test this prediction, we used two orthogonal approaches. First, we

assessed whether established signatures of immune cell infiltration [39,42] were present in the

tumors arranged according to increasing M-A scores, and observed enrichment for signatures of

infiltrating immune cells in tumors with mesenchymal phenotypes **(Supplementary Fig. 4b)**.

Intriguingly, we also noted that adrenergic tumors were enriched for CD276 (B7-H3) expression,

an immune checkpoint marker that protects neuroblastoma cells from NK cell-mediated

cytotoxicity, which may partly account for the decreased immune response signatures in these

cells[14] **(Supplementary Fig. 4b).** We next quantified the immune response signatures in a subset

of tumors (n = 100) from both the upper and lower quartiles of the M-A score that had significant

differences in their activation and evasion scores **(Supplementary Fig. 4c, d)**. High M-A scoring

(mesenchymal) tumors had significantly higher expression levels of cytotoxic T and NK cell

signatures compared with those of low M-A scoring (adrenergic) tumors (**Fig. 2f**). Considering

that the M-A score and, to a lesser extent, higher *MYC* expression were positively associated with

IA and IE scores in pairwise testing **(Supplementary Fig. 3f)**, we next assessed their relative

contributions as independent predictors of an immune response in a multivariate multiple

regression model consisting of immune activation and evasion scores and T and NK cell

signatures **(Fig. 2g)**. Because *MYCN* amplification is linked to immune suppression, we also

included the 157-gene *MYCN* signature generated by Valentijn et al[27] and subsequently used by

Wei et al to identify immune predictors in *MYCN*-nonamplified neuroblastoma[25]. In this analysis

also, the M-A score was a better predictor of tumor immunogenicity than *MYC* expression or

MYCN signature **(Fig. 2g)**.

Second, we used CIBERSORT (Cell type Identification By Estimating Relative Subsets Of

known RNA Transcripts)[43], as a deconvolution approach to estimate the fraction of immune cell

infiltration associated with the adrenergic and mesenchymal lineage tumors. We first established

the fraction of tumor-infiltrating leukocytes (TILs) in the most mesenchymal versus the most

adrenergic tumors (n = 6 each), observing significantly higher fractions in mesenchymal tumors

(Supplementary Fig. 4e). Next, we quantified the average immune cell content in 100 tumors

from the upper and lower quartiles of the M-A score **(Fig. 2h)**. Consistent with our previous results

(Fig. 2f), CIBERSORT analysis also showed significantly increased enrichment for cytotoxic CD8$^+$

T cells in mesenchymal tumors, which was also associated with a concomitant increase in

regulatory T cells, including those expressing markers of T-cell exhaustion **(Fig. 2h;**

Supplementary Fig. 4e). Interestingly, we also observed that mesenchymal tumors comprised a

higher fraction of naïve B cells, which were recently shown to take part in antitumor immunity [44]

(Fig. 2h; Supplementary Fig. 4e). Together, our findings indicate that the mesenchymal cell

state is a strong predictor of neuroblastoma immunogenicity.

Tumor cell-intrinsic upregulation of immune pathways in mesenchymal neuroblastoma

We next sought to understand the extent to which the presence or absence of an immunogenic

signature in the bulk RNA-sequencing data was intrinsic to tumor cells or was conferred by the

tumor microenvironment. Analysis of the lineage identities of a panel of 24 human neuroblastoma

cell lines (15 *MYCN*-amplified; 9 *MYCN*-nonamplified) from RNA-sequencing data (GSE28019)

revealed a gradient of adrenergic-to-mesenchymal scores **(Supplementary Fig. 5a)**. Consistent

Fig. 3.

Fig. 3. Tumor cell-intrinsic immune marker genes are upregulated in mesenchymal NBs.

(a) Scatter plot of the immune activation (IA) scores of human neuroblastoma cell lines (RNA-seq

data; GSE28019). Cell lines are arranged based on increasing IA scores and designated as

adrenergic or mesenchymal based on lineage-specific gene expression. **(b)** Western blot (WB)

analysis of adrenergic (PHOX2B) and mesenchymal (FN1, AXL) cell lineage markers and antigen

processing genes (TAP1/2, LMP2/7) in *MYCN*-nonamplified NB cell lines. GAPDH was used as

the loading control. Adrn, adrenergic; Mes, mesenchymal. **(c)** Waterfall plot of the fold-change in

RNA expression levels of up- and downregulated genes in SH-EP compared to SH-SY5Y NB

cells; selected immune genes are highlighted in green. **(d, e)** Heat maps of lineage marker (*blue*,

adrenergic; *red*, mesenchymal) and MHC and antigen processing machinery gene (*black*)

expression in the indicated *MYCN*-nonamplified **(d)** and *MYCN*-amplified (CHP-212, Kelly) and

overexpressing (NBL-S) **(e)** adrenergic and mesenchymal cells ($n = 2$ biological replicates). Rows

are z-scores calculated for each transcript in each cell type. **(f)** Fluorescence activated cell sorting

(FACS) analysis of cell surface HLA expression in the cells depicted in **e**. Isotype controls are

depicted in gray. The X-axis denotes fluorescence intensity of indicated proteins using

phycoerythrin (PE-A) tagged antibodies. Results representative of 2 independent experiments.

(g) RT-qPCR analysis of antigen processing and presentation genes in *MYCN*-amplified Kelly NB

cells engineered to express shMYCN or shGFP (control) with or without IFN-γ induction (100

ng/mL for 24 hr.). Data are normalized to GAPDH and represent means ± SD, $n = 2$ biological

replicates. Inset, WB analysis of MYCN in control and shMYCN cells. Actin was used as a loading

control.

Supplementary Fig. 5.

Supplementary Fig. 5. Mesenchymal lineage-specific marker and cell-intrinsic immune

gene expression are significantly correlated in NB. **(a)** Scatter plot of adrenergic and

mesenchymal scores in NB cell lines (GSE28019). **(b)** Volcano plot showing the gene expression

changes between mesenchymal SH-EP and adrenergic SH-SY5Y cells. The top ten lineage

marker genes are highlighted. The fold changes are represented in log_2 scale (X-axis) and the -

log_{10} of the *P*-values depicted on Y-axis (FDR < 0.1 and log_2FC > 1). **(c)** Venn diagram of the

overlap between DEGs in SH-EP cells (compared to SH-SY5Y cells) and the mesenchymal or

adrenergic signatures derived from Groningen et al[38]. Statistical significance was determined

using Fisher's exact test. **(d)** GO analysis of differentially upregulated genes in mesenchymal SH-

EP compared to adrenergic SH-SY5Y cells. **(e)** WB analysis of cell lineage marker and antigen

processing gene expression in the indicated NB cells. Dotted line indicates the margin where gel

images have been cut. **(f)** *Upper*, WB analysis of MYCN levels in the indicated numbers of Kelly

and NBL-S NB cells titrated against known amounts of purified GST-MYCN protein. *Lower,* WB

analysis comparing MYCN levels in whole cell extracts (WCE) from Kelly cells to titrated levels

from WCE in NBL-S cells. SH-SY5Y cells that do not express MYCN serve as a negative control.

Actin was used as a loading control in **(e)** and **(f)**.

with our observations in primary tumors, cell lines with higher mesenchymal gene signatures

grouped together and had significantly higher expression of tumor cell-intrinsic immune genes,

compared with the remainder, which had higher adrenergic scores and were mostly associated

with reduced immune marker gene expression (**Fig. 3a, b**). To further understand the association

of tumor cell-intrinsic immune pathways with lineage state, we focused on two neuroblastoma cell

lines –SH-SY5Y and SH-EP – subclones of the *MYCN*-nonamplified SK-N-SH cell line separated

on the basis of neuroblastic versus substrate-adherent morphology[45] and determined to be

adrenergic and mesenchymal[38], respectively. RNA sequencing showed that the differentially up-

and down-regulated genes in SH-EP compared with SH-SY5Y cells significantly overlapped with

established signatures of mesenchymal and adrenergic states, thus confirming their respective

phenotypes **(Supplementary Fig. 5b, c).** We noted that genes with roles in eliciting an immune

response were among the top differentially upregulated genes in mesenchymal SH-EP cells,

especially those involved in antigen processing and presentation and positive regulation of MHC

expression **(Fig. 3c, d)**. Moreover, gene ontology (GO) analysis of the upregulated transcripts

revealed enrichment for innate and adaptive immune responses including type-I interferon

signaling and ligands for the NK cell receptor, NKG2D (NK cell lectin-like receptor, KLRK1)

(Supplementary Fig. 5d). By contrast, adrenergic SH-SY5Y cells that showed upregulation of

neuronal lineage markers did not show significant enrichment of immune function genes **(Fig.**

3d), providing further evidence that cell-intrinsic immunogenicity is associated with the

mesenchymal phenotype.

The absence of a productive immune response has often been described in *MYCN-*

amplified neuroblastoma tumors[24,25]; indeed, the vast majority of such tumors in our cohort

exhibited similar findings **(Supplementary Fig. 2a, b)**. Nonetheless, based on our intriguing

finding of upregulation of immune response genes in a small number of *MYCN*-amplified tumors

(Supplementary Fig. 2a, b) that possessed mesenchymal cell signatures **(Fig. 2e)**, we sought

to understand the role of MYCN in mediating this immune response. We used Kelly and CHP-212

human neuroblastoma cells that expressed amplified *MYCN* but were of adrenergic and

mesenchymal phenotypes, respectively[37,38], and NBL-S cells that lacked *MYCN* amplification but

expressed moderate levels of MYCN RNA and protein[46] **(Supplementary Fig. 5e)** and were

classified as adrenergic (van Groningen et al., 2017 and this study). RNA-sequencing and flow

cytometry analysis suggested that tumor cell-intrinsic immune genes involved in antigen

processing and MHC regulation were highly expressed in mesenchymal CHP-212 compared to

adrenergic Kelly and NBL-S cells **(Fig. 3e, f; supplementary fig. 5e)**. Importantly, although

MYCN expression in NBL-S cells was lower than in Kelly cells **(Supplementary Fig. 5f)**, these

immune transcripts were expressed at lower levels in both cell lines, consistent with their

adrenergic status **(Fig. 3e; Supplementary Fig. 5f).** To further verify that cell state dictate tumor

cell-intrinsic immunogenicity, we depleted MYCN expression in Kelly cells and observed no

significant change to the IFN-γ-induced expression of HLA and antigen processing genes

compared to control cells **(Fig. 3g)**. Thus, our findings suggest that the lineage state of

neuroblastoma cells specifies the expression of tumor cell-intrinsic immune marker genes.

Cellular reprogramming to the mesenchymal state leads to increased immunogenicity

We next questioned whether acquisition of the mesenchymal phenotype would be sufficient to

render adrenergic neuroblastoma cells immunogenic. One of the top overexpressed genes in SH-

EP mesenchymal cells, *PRRX*1, encodes a core lineage-specific homeobox transcription factor

(TF), whose overexpression induces the mesenchymal state in neuroblastoma cells[37,38]

(Supplementary Fig. 5b). We therefore overexpressed doxycycline-inducible *PRRX1* in

adrenergic SH-SY5Y cells and observed a gradual loss of the adrenergic lineage marker

PHOX2B, together with increased expression of the mesenchymal markers, fibronectin, vimentin

and AXL **(Fig. 4a; Supplementary Fig. 6a)**. By contrast, overexpression of DNA-binding mutants

Fig. 4.

Fig. 4. Reprogramming of adrenergic NB cells to the mesenchymal cell state leads to

increased expression of immune response genes. (a) *Left*, WB analysis of PHOX2B and

antigen processing gene expression in adrenergic SH-SY5Y cells engineered to express

doxycycline (dox)-inducible PRRX1 in the presence or absence of dox (200 ng/mL) at the

indicated time points. *Right*, WB analysis of the indicated proteins in SH-SY5Y cells expressing

dox-inducible wild-type (WT) or DNA-binding mutants of PRRX1 at 10 days post dox-induction.

The DNA-binding (DB) mutants harbor individual deletions of the three α-helices (ΔH1, ΔH2 and

ΔH3) within the PRRX1 homeodomain. **(b)** RT-qPCR analysis of the indicated immune response

genes in the same cells as in **(a).** Data represent the means ± SD, *n* = 2 biological replicates. **(c)**

FACS analysis of cell surface HLA expression following dox-inducible expression of PRRX1 in

SH-SY5Y cells at the indicated time points. Data are representative of 2-3 independent

experiments. **(d)** FACS analysis of cell surface MICA/MICB expression after PRRX1 induction for

24 days in the same cells as in **(c).** A logscale expression value of 10^3 was used as a threshold

(vertical line) to gate MICA/MICB negative ($<10^3$) and positive ($\geq10^3$) populations. Numbers on

the right indicate the percentage of MICA/MICB-positive cells. Plots are representative of 2

independent experiments. **(e)** Heat map representation of adrenergic and mesenchymal gene

signatures in parental (5Y-par) and LDK-resistant (5Y-LDK-res) SH-SY5Y cells (*n* = 2 biological

replicates). Rows represent z-scores of $\log_2$ expression values for each gene in both cell types.

(f) Heat map depicting the expression of cell lineage markers (blue, adrenergic; red,

mesenchymal), antigen processing machinery genes and NKG2D ligands (black) in parental and

LDK-resistant SH-SY5Y cells (*n* = 2 biological replicates). Rows represent z-scores of $\log_2$

expression values. **(g)** WB analysis of lineage marker and antigen processing gene expression in

the indicated cells. Actin was used as a loading control in all immunoblots. **(h, i)** FACS analysis

of cell surface HLA **(h)** and NKG2D ligand **(i)** expression in the indicated cells.

Supplementary Fig. 6.

Supplementary Fig. 6. Acquired resistance to ceritinib (LDK387) in adrenergic SH-SY5Y

cells is associated with reprogramming to the mesenchymal lineage and increased

expression of immune response genes. **(a)** RT-qPCR analysis of PRRX1 expression in

adrenergic SH-SY5Y cells engineered to express doxycycline (dox)-inducible PRRX1 in the

presence or absence of dox (200 ng/mL) for 10 days. Data represent the means ± SD, *n* = 2

biological replicates. **(b)** Dose–response curves of ceritinib (LDK378)-sensitive (5Y-par.) and -

resistant (5Y-LDK-res.) SH-SY5Y cells treated with increasing concentrations of LDK378 for 72

h. Data represent means ± SD, *n* = 2 biological replicates. **(c)** Venn diagrams depicting the overlap

between the DEGs in LDK-resistant SH-SY5Y cells (compared to parental SH-SY5Y cells) and

the mesenchymal or adrenergic signatures derived from Groningen et al[38]. *P*-values were

determined by Fisher's exact test. **(d)** Waterfall plot of the fold-change in RNA expression levels

of up- and down-regulated genes in LDK-resistant SH-SY5Y cells compared to parental SH-SY5Y

cells; selected immune genes are highlighted in green. **(e)** GO analysis of differentially

upregulated genes in LDK-resistant SH-SY5Y compared to parental SH-SY5Y cells.

of *PRRX1* that contained homeodomain deletions had no effect on mesenchymal marker

expression, indicating that the lineage switch was a direct consequence of PRRX1-mediated

transcriptional control **(Fig. 4a)**. Next, to determine whether the phenotypic switch had any effect

on tumor cell-intrinsic pro-inflammatory pathways, we analyzed the expression of genes involved

in antigen processing (*TAP1, TAP2, LMP2, LMP7*) as well as *IFI16* and *STING* (*TMEM173*),

innate immune regulators that were differentially upregulated in mesenchymal NB cells **(Fig. 3c)**.

Induction of wild-type (WT) *PRRX1* led to increased RNA expression of these genes **(Fig. 4b)**.

Moreover, WT PRRX1 but not its DNA-binding mutants led to increased TAP1 and LMP7 protein

expression, which was accompanied by a sustained increase in cell surface MHC expression

(Fig. 4a, c). Additionally, PRRX1 induction led to increased cell surface expression of MICA and

MICB, ligands for the activating NK cell receptor NKG2D, in a minor population of cells **(Fig. 4d)**,

in agreement with elevated expression of these proteins in mesenchymal neuroblastoma cells

(Fig. 3c). These results suggest that conversion from the adrenergic to mesenchymal cell state

may be adequate to reprogram immune-insensitive cells toward immunocompetency.

Transition from the adrenergic to the mesenchymal state in neuroblastoma is

accompanied by resistance to chemotherapy [38]. Whether this transition in the face of treatment

pressure might include the acquisition of a pro-inflammatory signature is unclear, leading us to

compare adrenergic neuroblastoma cells that had gained mesenchymal features during the

development of treatment resistance with their sensitive, adrenergic counterparts. For this

purpose, we used an isogenic pair of cell lines comprising adrenergic SH-SY5Y neuroblastoma

cells that express the *ALK^{F1174L}* mutation and are sensitive to the small molecule inhibitor ceritinib

(LDK378) (parental SH-SY5Y, IC_{50} = 150 nM), and their ceritinib-resistant derivatives (LDK-

resistant SH-SY5Y, IC_{50} = 1101 nM) **(Supplementary Fig. 6b)**[47]. Comparison of the gene

expression signatures of these cell lines revealed significant downregulation of adrenergic

transcripts in LDK-resistant SH-SY5Y cells with concomitant upregulation of the mesenchymal

signature **(Fig. 4e)**. Moreover, a significant overlap was noted between the differentially up- or

downregulated transcripts in the LDK-resistant SH-SY5Y cells and established signatures of

mesenchymal and adrenergic states, respectively **(Supplementary Fig. 6c)**, suggesting that

these cells had acquired features of the mesenchymal phenotype with resistance. Consistent with

the key role of PRRX1 in triggering the conversion from an adrenergic to mesenchymal cell state,

we observed that this TF was among the top upregulated genes in LDK-resistant SH-SY5Y cells

(Supplementary Fig. 6d). The mesenchymal state of the LDK-resistant SH-SY5Y cells was

further supported by the loss of the pivotal adrenergic marker, PHOX2B, and increased

expression of additional mesenchymal markers AXL, YAP, TAZ, and IRF1, although these

changes were not as pronounced as those in SH-EP mesenchymal cells that served as a positive

control **(Fig. 4f, g)**. Further evidence supporting the conversion to the mesenchymal state came

from the differential upregulation in LDK-resistant SH-SY5Y cells of cell-intrinsic immune markers

engaged in antigen processing and presentation and NK cell activating receptor ligands (PSMB9,

MICA, MICB); in fact, these were among the top differentially upregulated genes in LDK-resistant

SH-SY5Y cells **(Supplementary Fig. 6d, e; Fig. 4f, g)**. These changes in immune genes

coincided with increases in cell surface expression of MHC receptors to levels comparable to

those in mesenchymal SH-EP cells **(Fig. 4h)**, as well as the increased expression of ligands for

the NK cell-activating receptor NKG2D **(Fig. 4i)**. Thus, the genetic reprogramming from the

adrenergic to the mesenchymal state that occurred with therapy resistance also led to the

upregulation of tumor cell-intrinsic pro-inflammatory pathway genes suggesting that such

conversion could render the tumor cells susceptible to recognition by T and NK cells.

Immune response gene expression during cell state transition is epigenetically regulated

As lineage plasticity in neuroblastoma is epigenetically driven[37,38,48,49], we next questioned

whether the altered expression of immune response genes observed in the individual cell states

could be the result of changes in chromatin organization. To this end, we analyzed the chromatin

occupancies of active and repressive histone marks at immune genes that were upregulated in

adrenergic SH-SY5Y cells upon induction of PRRX1 **(Fig. 4b)**. Indeed, PRRX1 induction resulted

in increased binding of the active H3K4me3 mark as well as loss of repressive H3K27me3 binding

at several candidate immune genes, including the APM genes *TAP1* and *PSMB9* **(Fig. 5a;**

Supplementary Fig. 7a). To understand epigenetic modifications that occur during the

spontaneous transition between the two lineage states (as compared with forced expression of

PRRXI) on a genome-wide basis, we compared histone occupancies between adrenergic

(parental SH-SY5Y) cells and those that had acquired mesenchymal characteristics with drug

resistance (LDK-resistant SH-SY5Y) **(Fig. 4e),** using SH-EP cells as a typical example of the

mesenchymal state. ChIP-seq analysis of active H3K27ac binding identified that the super-

enhancers (SEs) in LDK-resistant SH-SY5Y cells were associated with genes that conferred

mesenchymal identity while parental SH-SY5Y cells retained SEs at genes that conferred

adrenergic identity **(Supplementary Fig. 7b),** consistent with evidence that lineage plasticity is

driven by cell type-specific SEs[37,38]. To determine whether the SE-mediated regulation of lineage

genes also extended to genes associated with immune responsiveness, we analyzed the genes

in our 41-gene immune activation signature **(Supplementary Table 3;Fig. 1e)** as well as those

associated with an IFN-response signature (n = 91) in primary tumors and cell lines **(Fig. 1b;**

Supplementary Fig. 5d, see Methods**)**. Despite the higher expression of these genes in

mesenchymal cells (LDK-resistant SH-SY5Y and SH-EP), none was associated with an SE,

prompting us to focus on the promoter regions. We observed significantly higher enrichment of

H3K27ac and H3K4me3 binding at regions spanning the transcription start sites (TSS ± 2 kb) at

cell-intrinsic immune genes in LDK-resistant SH-SY5Y and SHEP compared to parental SH-SY5Y

Fig. 5

Fig. 5. Activation of **immune gene expression associated with cell state transition is**

epigenetically regulated. (a) *Upper,* Linear representation of TAP1 and PSMB9 gene loci

showing the locations of the bidirectional promoter and IRF1 and NF-κB binding sites. The

amplicons (1-8) analyzed for histone mark occupancy are shown in red. *Lower,* ChIP-qPCR

analysis of H3K4me3 and H3K27me3 enrichment at the indicated amplicons along the

TAP1/PSMB9 locus in adrenergic SH-SY5Y cells expressing dox-inducible PRRX1 in the

presence or absence of dox (200 ng/mL) for 10 days. Data represent the means ± SD, n = 2

biological replicates. **(b)** Metagene representations of average ChIP-seq occupancies of the

indicated histone marks at the promoters of tumor cell-intrinsic immune response genes (TSS ±

2 kb; n = 134) in parental (5Y-par.), LDK-resistant (5Y-LDK-res.) SH-SY5Y and SH-EP NB cells.

(c) Heat map representation of histone enrichment at the same immune gene promoters as in

(b), ranked in decreasing order of occupancy in the indicated cells. Each row represents the

normalized densities of histone marks within a ± 2 kb window centered on the TSS. **(d, e)**

Representation of pairwise comparisons between parental SH-SY5Y and LDK-resistant SH-SY5Y

(d), and parental SH-SY5Y and SH-EP cells **(e).** The changes (+, gained; -, lost) in occupancies

of the active (H3K4me3, H3K27ac) and repressive (H3K27me3) histone marks ($\log_2$ FC ≥ 0.75,

TSS ± 2 kb), together with the corresponding changes in RNA expression (+, overexpressed; $\log_2$

FC ≥ 1) of each of the 134 tumor cell-intrinsic immune genes analyzed in **(b)** are shown.

Representative genes showing either a switch from repressive to active chromatin (*MICB, ULBP3*)

or associated only with a gain of active chromatin (*IFIT3, STING*) are shown. **(f)** ChIP-seq tracks

depicting the gain of active histone binding together with the loss of repressive histone binding

(left) or gain of active marks without changes in repressive mark occupancy *(right)* at the indicated

immune gene loci. Signal intensity is given at the top left corner for each track. **(g)** Loess

regression analysis of the correlation between the ratios of active to repressive histone binding at

the promoters (TSS ± 2kb) of immune response genes and their RNA expression (*Upper,*

H3K4me3:H3K27me3; *Lower*, H3K27ac:H3K27me3). Genes are ranked based on increasing

expression. Shaded regions represent 95% confidence intervals.

Supplementary Fig. 7.

Supplementary Fig. 7. Immune gene activation associated with the mesenchymal cell state

is epigenetically regulated. (a) ChIP-qPCR analysis of H3K4me3 and H3K27me3 enrichment

at the promoters of the indicated immune genes in SH-SY5Y cells expressing doxycycline-

inducible PRRX1 in the presence or absence of dox (200 ng/mL) for 10 days**.** Enrichments at

TAP1 and PSMB9 loci correspond to amplicons 4 and 6 respectively, as described in **Fig. 5a**.

Data represent the means ± SD, $n = 2$ biological replicates. **(b)** Identification of enhancer regions

in parental SH-SY5Y, LDK-resistant SH-SY5Y and SH-EP cells. H3K27ac bound regions

identified as significant peaks were stitched together if they were within 12.5 kb of each other and

termed typical enhancers (plotted in grey). Super enhancers (SEs) were defined as stitched

enhancers surpassing the threshold signal based on the inclination point in all cell types (plotted

in red). In parental SH-SY5Y, LDK-resistant and SH-EP cells, 2.94% (915/31116), 6.56%

(1880/28635) and 4.18% (1215/29057) of the enhancers were classified as SEs respectively. The

top five SE-associated lineage-specific genes are highlighted. **(c)** ChIP-qPCR analysis of EZH2

and SUZ12 enrichment at the indicated genes in adrenergic 5Y-par. cells. Data represent the

means ± SD, $n = 2$ biological replicates, $^*P < 0.05$; $^{**}P < 0.01$ two-tailed Student's t-test. *P*-values

were calculated in comparison to enrichment observed at the Lin28B TSS (negative control

locus). **(d, e)** Scatter plots representing the differential binding of the indicated histone marks at

the promoter regions (TSS ± 2 kb) of all protein coding genes between parental SH-SY5Y (5Y-

par.) and LDK-resistant SH-SY5Y (5Y-LDK res.) **(d)**, and parental SH-SY5Y and SH-EP cells **(e)**.

rpm/bp, reads per million mapped reads per base pair. A ≥ 0.75 log$_2$FC threshold was used to

identify unique peaks for each individual histone mark. Unique and shared peaks are shown in

different colors. **(f)** Bar plots representing the numbers of immune genes with increased

deposition of H3K4me3 and H3K27ac (log$_2$ FC ≥ 0.75, TSS ± 2 kb) and loss of H3K27me3 (log$_2$

FC ≥ 0.75, TSS ± 2 kb) histone marks, together with increased RNA expression (log$_2$ FC ≥ 1) in

mesenchymal LDK-resistant SH-SY5Y (*left*) or SH-EP (*right*) as compared to adrenergic parental

SH-SY5Y cells. **(g)** Violin plots of the ratios of active to repressive histone marks (*left*,

H3K4me3:H3K27me3; *right*, H3K27ac:H3K27me3) surrounding immune gene promoters (TSS ±

2kb) in parental SH-SY5Y, LDK-resistant SH-SY5Y and SH-EP cells. Significance was

determined by the two-sided Wilcoxon rank-sum test.

cells **(Fig. 5b, c)**. On the other hand, the adrenergic parental SH-SY5Y cells showed significantly

higher occupancies of the H3K27me3 repressor mark at these immune gene promoters **(Fig. 5b,**

c). Analysis of the polycomb repressive complex 2 (PRC2) that promotes H3K27me3 deposition

at repressed chromatin[50] revealed that immune response genes enriched for H3K27me3 binding,

such as those encoding the NKG2D ligands *MICA/B*, *ULBP2/3* and *RAET1G* had significantly

higher occupancies for PRC2 subunits, EZH2 and SUZ12 in adrenergic parental SH-SY5Y cells

compared to negative control regions that lacked H3K27me3 binding (*PHOX2B* and *LIN28B*)

(Supplementary Fig. 7c), suggesting active immune gene repression in these cells.

We next sought to understand whether the activation of immune response genes observed

during the cell state transition from sensitivity to resistance represented a *switch* from repressive

to active chromatin or a *gain* of active chromatin marks. To this end, we quantified the changes

in histone binding occupancies between adrenergic parental SH-SY5Y and mesenchymal LDK-

resistant SH-SY5Y or SH-EP cells using pair-wise comparisons **(Supplementary Fig. 7d, e)**.

Compared to parental SH-SY5Y, LDK-resistant SH-SY5Y and SH-EP cells gained significant

H3K4me3 binding at the promoters of 60% and 62% immune genes (68 and 71 of 114)

respectively, which corresponded with their increased expression **(Fig. 5d, e; Supplementary**

Fig. 7f). A similar significant enrichment of the H3K27ac histone mark was observed at the

promoters of these immune genes [LDK-resistant SH-SY5Y, 58% (66/114); SH-EP, 67%

(76/114)]. Interestingly, gain of these active marks was accompanied by a concomitant loss of

H3K27me3 repressive histone binding at the promoters of 25% and 35% of (29 and 40 of 114)

immune genes in LDK-resistant SH-SY5Y and SH-EP cells respectively, as represented by the

NKG2D ligands, *MICB* and *ULBP3* **(Fig. 5d-f; Supplementary Fig. 7f).** On the other hand, in

LDK-resistant SH-SY5Y and SH-EP mesenchymal cells, 48% and 41% (55 and 47 of 114)

immune genes, such as the IFN-regulated factors *IFIT3 and STING,* gained either one or both

active marks without changes in occupancy of the repressive mark **(Fig. 5d-f; Supplementary**

Fig. 7f). Furthermore, the mesenchymal cells showed a significantly higher ratio of active to

repressor histone binding at the TSSs of immune-related genes (H3K4me3 or

H3K27ac/H3K27me3) **(Supplementary Fig. 7g)**, which importantly, also correlated with the

increased expression of these genes in this cell state **(Fig. 5g)**. Therefore, our results suggest

that the immune gene activation observed with the transition from the adrenergic to the

mesenchymal cell state represents either a switch from repressive to active chromatin or a gain

of active chromatin at promoter regions.

Mesenchymal neuroblastoma cells functionally engage cytotoxic T cells

To assess the functional consequences of the increased immunogenicity associated with a

mesenchymal phenotype, we utilized the murine neuroblastoma cell line NB-9464, which was

derived from tumors arising in the Th-MYCN genetically engineered mouse model (GEMM). This

model was generated in immunocompetent C57BL/6 mice and the tumors recapitulate the genetic

and immunological features of human neuroblastoma[51,52]. We observed that NB-9464 cells

consisted of distinct populations that could be sorted on the basis of surface MHC class I H-2Kb

expression into high (H-2Kb^{hi})- or low (H-2Kb^{lo})-expressing populations **(Supplementary Fig. 8a;**

Fig. 6a), both of which expressed transgenic human MYCN **(Fig. 6b)**. Consistent with our

hypothesis, H-2Kb^{hi} cells were enriched for *bona fide* mesenchymal markers - Prrx1, Sox9,

Notch1, and Snai2 (Slug) **(Fig. 6b)**, and showed enhanced migration and invasion, as might be

expected from their neural crest cell-like state **(Supplementary Fig. 8b, c)**. Importantly, the

mesenchymal H-2Kb^{hi} NB-9464 cells, but not their adrenergic H-2Kb^{lo} counterparts showed

augmented expression of cell surface class I MHC H-2Kb in response to IFN-γ, suggesting that

the mesenchymal cells had the potential for inducing a T cell-driven antitumor immune response

(Fig. 6c).

690   Hence, to determine whether the increased MHC class I expression of mesenchymal versus

691  adrenergic cells translated into T cell activation, we first asked whether mesenchymal H-2Kb[hi]

692  NB-9464 cells were capable of exogenous antigen presentation. Using the well-characterized

693  chicken ovalbumin-derived peptide (OVA$_{257-264}$ or SIINFEKL) antigen that binds to H-2Kb and can

694  be recognized by specific T cell receptors (TCRs) on CD8$^+$ T cells[53], we found that in comparison

695  to H-2Kb[lo] cells, H-2Kb[hi] cells expressed significantly higher levels of the H-2Kb-SIINFEKL

696  complex **(Fig. 6d)**. Next, we determined whether antigen presentation through H-2Kb enables

697  mesenchymal tumor cells to be recognized by antigen-specific T cell receptors (TCRs) on CD8$^+$

698  T cells, the first step towards a cytotoxic response. For this purpose, we used OT-I CD8$^+$ T cells

699  from C57BL/6 mice expressing a transgenic TCR that specifically recognizes the H-2Kb-

700  SIINFEKL complex[53]. H-2Kb[hi] or H-2Kb[lo] cells loaded with the SIINFEKL peptide were cocultured

701  with naïve OT-I cells, after which OT-I activation was measured through cell surface CD69

702  expression, an early marker of T-cell activation[54]. Mesenchymal H-2Kb[hi] cells led to significantly

703  higher OT-I activation in comparison with adrenergic H-2Kb[lo] cells, indicating specific recognition

704  of the H-2Kb-SIINFEKL complex by the TCR on OT-I cells **(Fig. 6e)**. By contrast, co-cultures of

705  OT-I cells and either H-2Kb[lo] or H-2Kb[hi] NB-9464 neuroblastoma cells without the SIINFEKL

706  peptide did not lead to T-cell recognition, confirming the specificity of the TCR-antigen interaction

707  **(Fig. 6e)**. Finally, we investigated whether the differential MHC-I expression between adrenergic

708  and mesenchymal neuroblastoma cells influenced tumor growth *in vivo* through subcutaneous

709  injection of H-2Kb[hi] or H-2Kb[lo] cells into syngeneic C57BL/6 (H-2Kb haplotype) mice

710  **(Supplementary Fig. 8d).** We noted an earlier onset of tumor formation with H-2Kb[lo] cells

711  compared to H-2Kb[hi] cells **(Supplementary Fig. 8d)**. However, once consistent tumor growth

712  was established, growth or survival rates did not change substantially between the two groups

713  **(Supplementary Fig. 8e)**, despite the persistence of higher H-2Kb and Prrx1 expression in the

714  H-2Kb[hi] tumors compared with the H-2Kb[lo] tumors **(Supplementary Fig. 8f,g).** While both types

715  of tumor cells had MYCN expression, histologically, in keeping with their adrenergic phenotype

Fig. 6.

Fig. 6. Mesenchymal NB cells functionally engage cytotoxic T cells. (a) FACS analysis of

H-2Kb expression in unsorted NB-9464 cells. Vertical dashed line denotes the logscale

expression value used as a threshold to gate H-2Kb^{lo} and H-2Kb^{hi} cell populations. **(b)** WB

analysis of the indicated lineage markers in H-2Kb^{lo} and H-2Kb^{hi} cell populations. Notch1-FL, full

length; -TM, transmembrane; -IC, intracellular. GAPDH is used as a loading control. **(c)** FACS

analyses of basal and IFN-γ-induced (100 ng/mL for 24 hr.) surface H-2Kb expression (darker

colored histograms) in H-2Kb^{lo} and H-2Kb^{hi} cells compared to isotype controls (lighter colored

histograms). Plots representative of 2 independent experiments. **(d)** *Upper,* Schematic of OVA

binding assay. *Lower*, FACS analysis of surface H-2Kb-bound SIINFEKL OVA peptide in H-2Kb^{hi}

and H-2Kb^{lo} cells under basal or IFN-γ-induced conditions as in **(c)** and in the absence or

presence of the OVA peptide. Plots representative of 2 independent experiments. **(e)** *Upper,*

Schematic of NB-9464-OT-I co-culture assay. *Lower*, Contour plots showing the percentage of

naïve OT-I cells that were activated (CD8$^+$ CD69$^+$) following co-culture with H-2Kb^{lo} and H-2Kb^{hi}

cells for 24 hr. with or without the OVA peptide. OT-I activation was measured by FACS analysis

of cell surface CD69. **(f)** *Left,* Immunohistochemical (IHC) staining for MYCN and CD8 expression

in representative murine NB xenograft tumors derived from NB-9464 H-2Kb^{lo} (adrenergic) and H-

2Kb^{hi} (mesenchymal) cells in immunocompetent syngeneic (C57BL/6) mice. Scale bars, 100 μm.

Right, Bar graphs showing the percentage of area occupied by CD8$^+$ T cells in H-2Kb^{lo} (0.9% ±

0.2%) vs. H-2Kb^{hi} (2.3% ± 0.5%) tumors; *P < 0.05, two-tailed Welch's t-test. Each dot represents

one of three independent measurements for each tumor. Data represent mean ± SEM.

Supplementary Fig. 8.

Supplementary Fig. 8. Tumors that arise from NB-9464 H-2Kb^{hi} (mesenchymal) cells show

cytotoxic T cell infiltration. **(a)** FACS scatter plot showing gating conditions used for sorting

NB-9464 cells into H-2Kb^{hi} and H-2Kb^{lo} populations. X axis represents side scatter (SSC-A); Y

axis denotes fluorescence intensity of surface H-2Kb detected using phycoerythrin (PE)-

conjugated antibody against H-2Kb. A logscale expression value of 10^3 was used as the threshold

(horizontal dashed line) to gate H-2Kb^{hi} (≥ 10^3) and H-2Kb^{lo} (< 10^3) populations. **(b)** *Left*, Bright

field images of crystal violet-stained H-2Kb^{hi} and H-2Kb^{lo} cells in transwell migration assays. Scale

bars, 100 μm. *Right,* Quantification of migrating cells per high-power field (HPF). Data represent

the means ± SD, *n* = 2 biological replicates, $^{***}P$ < 0.001; two-tailed Student's t-test. **(c)**

Quantification of the relative invasiveness of H-2Kb^{hi} and H-2Kb^{lo} cells. Data represent the means

± SD, *n* = 2 biological replicates, $^{***}P$ < 0.001; two-tailed Student's t-test. **(d)** Tumor volumes in

immunocompetent C57BL/6 mice injected subcutaneously with 1 x 10^6 H-2Kb^{lo} or H-2Kb^{hi} NB-

9464 cells. Measurements were started on day 3 after injection and continued three times weekly

for up to 50 days or until euthanized due to tumor growth. Graphs represent changes in tumor

volume until day 31 (means ± SD; n = 7 per group at all time points) to highlight an earlier onset

of tumor formation with H-2Kb^{lo} cells compared to H-2Kb^{hi} cells. Tumors were considered to be

established upon reaching a volume of ~250 mm^3 (observed between days 31-34 for both H-2Kb^{lo}

and H-2Kb^{hi} tumors), following which both tumor types displayed equal increases in tumor growth

(data not shown). Tumor onset was defined as the day following which tumor volumes showed a

consistent increase (24.3 ± 2.2 days for H-2Kb^{lo} and 27.4 ± 2.1 days for H-2Kb^{hi} cells, *P* < 0.05).

Closed arrows refer to the indicated days (13, 22, 27 and 31) on which there were significant

differences in tumor volumes (day 13, *P* = 0.02; day 22, *P* = 0.01; day 27, *P* = 0.02 and day 31,

P = 0.02; *n* = 7 per group at all time points). All *P*-values calculated using the two-tailed Student's

t-test. **(e)** Kaplan-Meier survival analysis of immunocompetent C57BL/6 mice bearing NB tumor

xenografts derived from H-2Kb^{lo} (adrenergic) and H-2Kb^{hi} (mesenchymal) cells (50.8 ± 3.7 vs.

49.8 ± 4.8 days; *P* = 0.7; n = 7 per group; log-rank test. **(f)** RT-qPCR analysis of H-2Kb expression

in H-2Kb^{lo} and H-2Kb^{hi} NB tumor xenografts and the cell lines used to generate the xenografts.

Data represent the means ± SD, *n* = 3 biological replicates. **(g)** Immunofluorescence images of

MYCN (green) and Prrx1 (red) expression in representative murine NB xenograft tumors derived

from NB-9464 H-2Kb^{hi} (mesenchymal) and H-2Kb^{lo} (adrenergic) cells in immunocompetent

syngeneic (C57BL/6) mice. Nuclei are counterstained with DAPI (blue). Insets depict cells with

nuclear co-staining of MYCN and Prrx1 (arrowheads) and are exclusively present in the H-2Kb^{hi}

mesenchymal tumor. Scale bars, 100 µm, insets 33.3 µm.

H-2Kb^{lo} tumors were composed of densely arranged small round blue cells, whereas H-2Kb^{hi}

tumors predominantly comprised elongated, spindle-like cells interspersed with clusters of small

round blue cells **(Fig. 6f)**. We next analyzed the immune status of these tumors, reasoning that

tumors arising from immunogenic H-2Kb^{hi} cells would be infiltrated by T cells. Indeed, H-2Kb^{hi}

tumors showed significantly higher CD8$^+$ T cell infiltration compared with H-2Kb^{lo} tumors **(Fig. 6f)**.

Taken together, these results suggest that the immunogenic traits of mesenchymal

neuroblastoma cells translate into the recruitment of cytotoxic T cells into the tumor

microenvironment.

Mesenchymal neuroblastoma cells induce NK cell degranulation

We next analyzed the functional relevance of the increased expression of the NKG2D NK cell

receptor ligands (ULBP1-3, MICA, MICB) seen in mesenchymal neuroblastoma tumors

(Supplementary Fig. 4c; Fig. 4i). Since the interaction of the receptor with its cognate ligands

on target cells is the first step towards a cytotoxic response **(Fig. 7a)**, we compared the ability of

adrenergic (parental SH-SY5Y) and mesenchymal (LDK-resistant SH-SY5Y and SH-EP) cells to

bind to the recombinant NKG2D receptor fusion protein in an *in vitro* binding assay. Both

mesenchymal cell types showed increased binding to the NKG2D receptor, in keeping with their

increased cell surface expression of the ULBP2/3/5/6, MICA and MICB ligands **(Fig. 7b)**. NK cell

cytotoxicity is mediated by granzyme proteases and the pore-forming protein perforin, which in

resting cells are stored in secretory lysosomes or lytic granules marked by lysosome-associated

membrane protein-1 (LAMP-1 or CD107a)[55]. Upon target recognition, NK cells undergo

degranulation, or exocytosis of the lytic granules, which is associated with relocation of the

CD107a antigen to the cell membrane. Using cell-surface CD107a as a specific marker of

degranulation, we measured degranulation of peripheral blood NK cells harvested from healthy

human donors in the presence of parental SH-SY5Y, LDK-resistant SH-SY5Y and SH-EP targets.

Fig. 7.

Fig. 7. Mesenchymal NB cells induce NK cell degranulation. (a) Schematic representation of

the interaction between NKG2D receptors on NK cells and cognate ligands on tumor cells, leading

to NK cell degranulation and receptor endocytosis. **(b)** FACS analysis of purified human NKG2D-

Fc protein binding (darker histograms) to adrenergic parental SH-SY5Y, and mesenchymal LDK-

resistant SH-SY5Y and SH-EP cells. Comp Alexa-647, compensated fluorescence intensity of

NKG2D-Fc protein detected using Alexa 647-conjugated anti-human IgG. Lighter histograms

indicate staining with Alexa 647-conjugated anti-human IgG only. Numbers indicate median

fluorescence intensity (MFI). Plots representative of 2 independent experiments. **(c)** X-Y plot

showing the percentage of degranulating NK cells following co-culture with the same cells as in

(b) for 4 hr. at an effector: target (E: T) cell ratio of 1:2. Degranulation was measured by FACS

analysis of cell-surface CD107a. Data represent the means ± SD, $n = 4$ biological replicates. **(d)**

X-Y plot showing the effect of a control IgG1 or an NKG2D blocking antibody on NK cell

degranulation following co-culture with the indicated cells for 4 hr. NK cell degranulation was

measured as in **(c).** Data represent the means ± SD, $n = 4$ biological replicates. **(e)** X-Y plot

depicting the MFI of NKG2D expression measured by FACS on naïve NK cells (NK alone) or

following co-culture with parental SH-SY5Y, LDK-resistant SH-SY5Y and SH-EP cells for 4 h.

Data represent the means ± SD, $n = 4$ biological replicates. **(f)** FACS analysis of surface

ULBP2/5/6, ULBP3, and MICA/MICB in adrenergic parental SH-SY5Y cells treated with DMSO

(vehicle control) or EED226 (5 µM for 8 days). **(g)** FACS analysis of purified human NKG2D-Fc

protein binding to parental SH-SY5Y cells treated with either DMSO or EED226 as in **(f)**. Light

gray histograms indicate human IgG1 isotype control. Numbers on the right represent MFI values

for FACS plots. Plots in **(f)** and **(g)** are representative of 2 independent experiments. **(h, i)** X-Y

plots showing NK cell degranulation **(h)** and MFI of surface-NKG2D **(i)** following co-culture of

naïve NK cells with parental SH-SY5Y cells treated with DMSO or EED226 as in **(f)**. NK cell

granulation and NKG2D MFI were measured as in **(c)** and **(e)**. Data represent means ± SD, $n =$

4 **(h)** and 5 **(i)** biological replicates. Significance for all results was calculated using the paired

two-tailed Student's t-test (*P < 0.05; **P < 0.01); *ns*, not significant

Supplementary Fig. 9.

Supplementary Fig. 9. LDK-resistant SH-SY5Y cells are more susceptible to NK-induced cell death.

(a) Bargraphs showing NK-induced cell death in parental SH-SY5Y and LDK-resistant SH-SY5Y cells assessed by SYTOX green uptake following 1 hour of co-culture at indicated effector-to-target (E:T) ratios. Experiments were performed in two biological replicates using NK cells harvested from two independent donors. Data represent means ± SD, n = 3 technical replicates. Significance for all results was calculated using the paired two-tailed Student's t-test (*P < 0.05; **P < 0.01).

Co-culture of NK cells with LDK-resistant SH-SY5Y and SH-EP cells, both of which express

ligands for the NKG2D receptor, resulted in increased NK cell degranulation compared to parental

SH-SY5Y cells that did not express these ligands (**Fig. 7c**). To confirm that the increased NK cell

degranulation in LDK-resistant SH-SY5Y and SH-EP cells was specific to the NKG2D receptor,

we blocked its function with an anti-NKG2D monoclonal antibody. Compared to the isotype

control, blockade of NKG2D receptor activity completely abrogated the increased NK cell

degranulation in LDK-resistant SH-SY5Y and SH-EP cells, but had no effects on parental SH-

SY5Y cells, signifying that the modest but robust increase in degranulation in the presence of

mesenchymal cells was specific to the NKG2D receptor on NK cells **(Fig. 7d).** Upon interaction

with their cognate ligands on target cells, NKG2D receptors are internalized via ubiquitin-

dependent endocytosis leading to their lysosomal degradation, rendering the loss of surface

NKG2D receptor expression a robust readout for ligand-receptor engagement[56] **(Fig. 7a)**. In line

with the presence of functional NKG2D ligands on mesenchymal cells, co-cultures with LDK-

resistant SH-SY5Y and SH-EP cells led to significant downregulation of NK cell surface-

associated NKG2D expression, whereas co-culture with the adrenergic parental SH-SY5Y cells

did not alter the abundance of surface NKG2D expression **(Fig. 7e)**. Moreover, consistent with

the increased levels of NK degranulation upon co-culture with mesenchymal cells, LDK-resistant

SH-SY5Y were more susceptible to NK-induced cell death compared to parental SH-SY5Y cells

(Supplementary Fig. 9a)

The observation that genes encoding NKG2D ligands are repressed by the PRC2 complex in

adrenergic parental SH-SY5Y cells **(Supplementary Fig. 7c)** prompted us to examine whether

PRC2 inhibitors could induce the expression of these transcripts and influence NK cell function.

Indeed, treatment of parental SH-SY5Y cells with EED226, an allosteric inhibitor of the PRC2

complex[57], led to increased expression and surface localization of ULBP2/3 and MICA/B NKG2D

ligands **(Fig. 7f)**. Moreover, such increased ligand expression led to their increased binding to the

NKG2D receptor fusion protein in EED226-treated cells compared to cells treated with DMSO

alone **(Fig. 7g)**. Consequently, PRC2 inhibition resulted in an ~20% increase in NK cell

degranulation **(Fig. 7h)**. Finally, co-culture of primary NK cells with adrenergic parental SH-SY5Y

cells treated with EED226 led to a significant loss of surface NKG2D receptor expression **(Fig.**

7i), suggesting that the increased degranulation resulting from PRC2 inhibition was driven by the

NKG2D receptor. Overall, these results suggest that the lineage-specific expression of NK cell

ligands in mesenchymal neuroblastoma cells has a functional impact on NK cell activity, and that

pretreatment of adrenergic neuroblastoma cells could potentially render these cells susceptible

to NK-cell mediated immunotherapy by upregulating ligand expression.

DISCUSSION

Despite the relatively poor track record of immunotherapy for neuroblastoma, growing evidence

suggests that subsets of these tumors have the potential to induce a productive immune

response[9]. Here, we demonstrate that the mesenchymal cell state, characterized by neural crest

cell (NCC)-like phenotypes, is a strong predictor of an antitumor immune response in

neuroblastoma. Induction of this state was accompanied by the expression of tumor cell-intrinsic

immune response-inducing genes that were epigenetically repressed in the more differentiated

(adrenergic) tumor cells. Importantly, inhibition of the PRC2 complex relieved such repression of

ligands for the activating NK cell receptor NKG2D, and led to NK cell degranulation, suggesting

that this strategy could be explored as a potential measure to improve the response of patients

with adrenergic neuroblastomas to NK cell-mediated therapy.

We sought to identify neuroblastomas capable of eliciting an immune response as those

characterized by the differential expression of immune gene signatures while remaining agnostic

to any of the established parameters that predict disease aggressiveness. Using UMAP

dimension reduction to analyze gene expression data from 498 primary neuroblastoma tumors,

we identified four clusters that were separated on the basis of differential activation of gene

networks that regulate the antitumor immune response, neuronal differentiation, MYCN-driven

processes and lipid metabolism. Intriguingly, the immunogenic cluster comprised almost equal

proportions of high- and low-risk tumors, raising the possibility that the molecular mechanisms

underlying immunogenicity in neuroblastoma are independent of disease aggressiveness. Thus,

our analysis using a cluster-based approach enabled the identification of antitumor immune

signatures as shared transcriptional programs between high- and low-risk neuroblastoma tumors,

a bridging feature that would have been missed in studies based solely on differential gene

expression between prognostically distinct groups of tumors.

A major finding of our study is the intimate link between tumor cell lineage and the propensity

of eliciting an immune response. Neuroblastoma tumors show lineage plasticity, underscored by

a phenotypic switch between undifferentiated NCC-like and more differentiated cells, two

divergent cell states driven by distinct transcriptional programs [37,38]. Our analysis revealed that

neuroblastomas enriched in NCC-derived signatures showed significantly higher antitumor

immunity featuring T and NK cells compared to tumors enriched for signatures of the adrenergic

lineage. Our finding is substantiated by studies showing that diverse cellular states such as

stemness, senescence and metastasis strongly influence the engagement of innate and adaptive

immune pathways [58-60]. Importantly, while the NCC-like/mesenchymal state promoted immune

response mechanisms such as upregulation of MHC class I and infiltration of cytotoxic

lymphocytes, these tumors were also characterized by the activation of immune checkpoints such

as regulatory T cells and exhaustion markers linked to immune suppression, similar to those

observed in chronic virally infected states [61]. The presence of such seemingly contradictory gene

signatures has important therapeutic implications for selecting patients who are likely to benefit

from T cell-based immunotherapies, as agents that target negative regulatory immune

checkpoints are likely to be most effective in those with a pre-existing but dampened antitumor

immune response [35]. Moreover, inhibition of cancer cell-intrinsic transcriptional programs that

promote T cell exclusion have been shown to effect changes in cell state and could potentially

employed to sensitize tumors to immunotherapy (Jerby-Arnon et al., 2018; Koyama et al., 2016;

Spranger et al., 2015).

In contrast to other *MYC*-driven cancers where high *MYC* levels restrain inflammatory

signaling and anti-tumor immune pathways[41,62], tumors within our immunogenic cluster, enriched

largely for mesenchymal/NCC signatures, had relatively high MYC expression that positively

correlated with immune cell infiltration. These diametrically opposite roles of MYC in regulating

immune response suggest that its transcriptional functions are likely to be very different in cancers

in which aberrant MYC expression is the main oncogenic driver compared to *MYCN*-nonamplified

neuroblastomas that are not dependent on MYC overexpression. This notion is supported by the

observation that inhibiting endogenous MYC function in non MYC-driven pancreatic cancer

models leads to decreased recruitment and retention of inflammatory cells[63]. As MYC is an

essential TF in NCCs[64,65], it is plausible that MYC promotes an immunogenic state by driving a

NCC-specific transcriptional program.

We also extended the analysis between cell state and immunogenicity to *MYCN*-amplified

tumors and identified a subset that incorporates both mesenchymal and immunogenic features.

This finding was substantiated by our data showing that expression of functional MHC class I is

retained in some *MYCN*-amplified mesenchymal neuroblastoma cells and is not perturbed by

changes in MYCN levels. Indeed, a robust antitumor immune response is observed during the

early stages of tumor development in Th-MYCN mice and in mouse-human chimeric tumors

derived from human NCCs expressing *MYCN* and oncogenic *ALK*[17]. Considering that these

mouse models are driven by gain of *MYCN* (~4-8 copies of the *MYCN* transgene)[17,52] rather than

the amplification seen in human tumors, and that absolute levels of MYCN protein dictate

transcriptional output[66], the relatively low MYCN dosage in these tumors may account for their

immunogenicity. These findings, if confirmed in additional data sets, should encourage us to

reconsider the notion that all *MYCN*-amplified tumors are intrinsically immune tolerant and that a

subset may in fact, be capable of inducing an immune response by virtue of their cell state.

The mutual exclusivity of the neuronal and immunogenic clusters suggests that

neuroblastomas with characteristics of neuronal differentiation (i.e. adrenergic phenotype) are

weakly immunogenic and thus incapable of inducing an effective immune response. Moreover,

the presence of NCC-derived gene expression signatures among immunogenic tumors indicates

that phenotypic reversal of sympathetic neuronal cells to an NCC-like state could contribute

significantly to the antitumor immune response. Such cell state-dependent immunogenic

switching could be mediated by the lineage-specific core regulatory circuitry (CRC) that drives

distinct transcriptomic states in neuroblastoma. Indeed, several TFs that constitute the NCC-like

(mesenchymal) CRC, including interferon regulatory factors 1-3 and IFI16, function as major

drivers of tumor cell-intrinsic innate and adaptive immune responses[67,68]. Furthermore, our data

identify PRRX1, another component of the mesenchymal CRC, as a regulator of MHC class I and

antigen-processing gene expression in neuroblastoma. Interestingly, PRRX1 was not identified

as a candidate regulator of MHC-I expression in genome-wide CRISPR knock-out screens to

identify NF-κB-dependent MHC-I suppressors in neuroblastoma[69], suggesting that the

mechanisms employed by PRRX1 could be independent of NF-kB activation and may involve

direct transcriptional activation of these genes. In addition, the upregulation of DNA damage

sensor proteins such as IFI16 and STING in mesenchymal neuroblastomas could also contribute

to the increased tumor-infiltrating lymphocyte abundance in these tumors, as suggested by results

in small cell lung cancer[70].

We have also established that the changes in immune gene expression accompanying the

adrenergic to mesenchymal transition are epigenetically regulated. Unlike lineage identity genes

that are regulated by super-enhancers, tumor cell-intrinsic immune genes involved in diverse

immune functions such as the inflammatory response, IFN-γ signaling and NK cell recognition,

are governed through changes in promoter structure, achieved by either *de novo* acquisition of

permissive chromatin or an epigenetic switch from PRC2-mediated repression to a permissive

chromatin landscape. Our findings support a role for the PRC2 complex in repressing genes that

encode ligands for the activating NK cell receptor NKG2D in adrenergic neuroblastoma cells and

the use of PRC2 inhibitors to augment the NK cell response against these cells. This approach is

justified on several grounds: our results add to the growing body of evidence for tumor-cell

autonomous function of PRC2 as a barrier to anti-tumor immunity, achieved through inhibition of

processes such as MHC expression[11,71], antigen processing and presentation and inflammatory

cytokine production[72,73]. Moreover, our observation that NKG2D ligands are enriched in

mesenchymal neuroblastomas coupled with the demonstration of the critical effector role of NK

cells in the antitumor immune response against this tumor [51] and the promising responses of

patients with other solid tumors to NKG2D-directed CAR NK cell therapy [74] strengthen this

premise. Considered together, the results of our analysis identify cell lineage as an important

determinant of the immune responsiveness of neuroblastoma and suggest rationales for the use

of immune-based therapies, either alone or in combination with epigenetic inhibitors, against the

two divergent phenotypes that define the lineage state of this pediatric tumor.

Acknowledgements

We thank Matthew Harlow from the George lab, Sumit Sen Santara, Ying Zhang, and Zhibin Zhang from the Lieberman lab for helpful discussions. We thank Mark Zimmerman, A. Thomas Look and Kimberly Stegmaier for sharing cell lines. We are thankful to the following members of the former Haining lab at DFCI for sharing resources and experimental advice: Ulrike Gerdemann, Dawn Comstock, Kathleen Yates, Anna Word, and Adrienne Long. The results shown here are in part based on data curated by the R2: Genomics Analysis and Visualization Platform: http://r2.amc.nl/. This work was supported by a St. Baldrick's Foundation Childhood Cancer Research Grant (R.E.G. and R.J.) DOD CA191000 (R.E.G. and R.J.) and NIH grants R01-CA197336; R01-CA148688 (R.E.G). St. S. is a recipient of the Pew Stewart Scholarship. Sa. S. and M.K. were supported by the Rally Foundation for Childhood Cancer Research and Infinite Love for Kids Fighting Cancer, B.C.M. by the National Center for Advancing Translational Sciences/NIH Award KL2 TR002542 and D.N.D. by an Alex's Lemonade Stand Foundation Young Investigator Fellowship. This manuscript is dedicated to the memory of John R. Gilbert, Scientific Editor.

Author contributions

Sa.S. and R.E.G. conceived the study. Sa.S., A.C., B.C.M., J.L., and R.E.G. designed the experiments. Sa.S. performed the molecular, cellular and genomic studies. S.D. conceived and performed the genomic and computational analysis with inputs from R.D., Sa.S. and R.E.G. Sa.S., A.C., and B.C.M. performed the T and NK cell studies. B.S. performed the animal and cloning experiments. S.Z. performed quantitative analysis of IHC images. H.H. performed the cell migration assays. M.K., D.N.D., and L.S. contributed to FACS analysis, generation of LDK378-resistant SH-SY5Y cells, and compound testing, respectively. M.C., R.V., R.J., and St.S. contributed ideas towards regulation of immune function and cell lineage state. Sa.S., S.D., and R.E.G. wrote the manuscript with input from all authors.

Competing Interests

R.J. is a cofounder of Fate Therapeutics, Fulcrum Therapeutics, and Omega Therapeutics and an advisor to Dewpoint Therapeutics.

References

1. Yu, A.L. *et al.* Anti-GD2 antibody with GM-CSF, interleukin-2, and isotretinoin for neuroblastoma. *N Engl J Med* **363**, 1324-34 (2010).

2. Cheung, N.K. & Dyer, M.A. Neuroblastoma: developmental biology, cancer genomics and immunotherapy. *Nat Rev Cancer* **13**, 397-411 (2013).

3. Singh, N. *et al.* T cells targeting NY-ESO-1 demonstrate efficacy against disseminated neuroblastoma. *Oncoimmunology* **5**, e1040216 (2016).

4. Park, J.R. *et al.* Adoptive transfer of chimeric antigen receptor re-directed cytolytic T lymphocyte clones in patients with neuroblastoma. *Mol Ther* **15**, 825-33 (2007).

5. Pule, M.A. *et al.* Virus-specific T cells engineered to coexpress tumor-specific receptors: persistence and antitumor activity in individuals with neuroblastoma. *Nat Med* **14**, 1264-70 (2008).

6. Louis, C.U. *et al.* Antitumor activity and long-term fate of chimeric antigen receptor-positive T cells in patients with neuroblastoma. *Blood* **118**, 6050-6 (2011).

7. Merchant, M.S. *et al.* Phase I Clinical Trial of Ipilimumab in Pediatric Patients with Advanced Solid Tumors. *Clin Cancer Res* **22**, 1364-70 (2016).

8. Davis, K.L. *et al.* Nivolumab in children and young adults with relapsed or refractory solid tumours or lymphoma (ADVL1412): a multicentre, open-label, single-arm, phase 1-2 trial. *Lancet Oncol* **21**, 541-550 (2020).

9. Richards, R.M., Sotillo, E. & Majzner, R.G. CAR T Cell Therapy for Neuroblastoma. *Front Immunol* **9**, 2380 (2018).

10. Bernards, R., Dessain, S.K. & Weinberg, R.A. N-myc amplification causes down-modulation of MHC class I antigen expression in neuroblastoma. *Cell* **47**, 667-74 (1986).

11. Burr, M.L. *et al.* An Evolutionarily Conserved Function of Polycomb Silences the MHC Class I Antigen Presentation Pathway and Enables Immune Evasion in Cancer. *Cancer Cell* **36**, 385-401 e8 (2019).

12. Raffaghello, L. *et al.* Multiple defects of the antigen-processing machinery components in human neuroblastoma: immunotherapeutic implications. *Oncogene* **24**, 4634-44 (2005).

13. Raffaghello, L. *et al.* Downregulation and/or release of NKG2D ligands as immune evasion strategy of human neuroblastoma. *Neoplasia* **6**, 558-68 (2004).

14. Castriconi, R. *et al.* Identification of 4Ig-B7-H3 as a neuroblastoma-associated molecule that exerts a protective role from an NK cell-mediated lysis. *Proc Natl Acad Sci U S A* **101**, 12640-5 (2004).

15. Coughlin, C.M. *et al.* Immunosurveillance and survivin-specific T-cell immunity in children with high-risk neuroblastoma. *J Clin Oncol* **24**, 5725-34 (2006).

16. Betancur, P.A. *et al.* A CD47-associated super-enhancer links pro-inflammatory signalling to CD47 upregulation in breast cancer. *Nat Commun* **8**, 14802 (2017).

17. Cohen, M.A. *et al.* Formation of Human Neuroblastoma in Mouse-Human Neural Crest Chimeras. *Cell Stem Cell* **26**, 579-592 e6 (2020).

18. Asgharzadeh, S. *et al.* Clinical significance of tumor-associated inflammatory cells in metastatic neuroblastoma. *J Clin Oncol* **30**, 3525-32 (2012).

19. Song, L. *et al.* Valpha24-invariant NKT cells mediate antitumor activity via killing of tumor-associated macrophages. *J Clin Invest* **119**, 1524-36 (2009).

20.   Mao, Y. *et al.* Targeting Suppressive Myeloid Cells Potentiates Checkpoint Inhibitors to
Control Spontaneous Neuroblastoma. *Clin Cancer Res* **22**, 3849-59 (2016).

21.   Tran, H.C. *et al.* TGFbetaR1 Blockade with Galunisertib (LY2157299) Enhances Anti-
Neuroblastoma Activity of the Anti-GD2 Antibody Dinutuximab (ch14.18) with Natural
Killer Cells. *Clin Cancer Res* **23**, 804-813 (2017).

22.   Brodeur, G.M., Seeger, R.C., Schwab, M., Varmus, H.E. & Bishop, J.M. Amplification of N-
myc in untreated human neuroblastomas correlates with advanced disease stage. *Science*
224, 1121-4 (1984).

23.   Seeger, R.C. *et al.* Association of multiple copies of the N-myc oncogene with rapid
progression of neuroblastomas. *N Engl J Med* **313**, 1111-6 (1985).

24.   Layer, J.P. *et al.* Amplification of N-Myc is associated with a T-cell-poor microenvironment
in metastatic neuroblastoma restraining interferon pathway activity and chemokine
expression. *Oncoimmunology* **6**, e1320626 (2017).

25.   Wei, J.S. *et al.* Clinically Relevant Cytotoxic Immune Cell Signatures and Clonal Expansion
of T-Cell Receptors in High-Risk MYCN-Not-Amplified Human Neuroblastoma. *Clin Cancer*
Res **24**, 5673-5684 (2018).

26.   Brandetti, E. *et al.* MYCN is an immunosuppressive oncogene dampening the expression
of ligands for NK-cell-activating receptors in human high-risk neuroblastoma.
Oncoimmunology **6**, e1316439 (2017).

27.   Valentijn, L.J. *et al.* Functional MYCN signature predicts outcome of neuroblastoma
irrespective of MYCN amplification. *Proc Natl Acad Sci U S A* **109**, 19190-5 (2012).

28.   Durbin, B.P., Hardin, J.S., Hawkins, D.M. & Rocke, D.M. A variance-stabilizing
transformation for gene-expression microarray data. *Bioinformatics* **18 Suppl 1**, S105-10
(2002).

29.   Lin, S.M., Du, P., Huber, W. & Kibbe, W.A. Model-based variance-stabilizing
transformation for Illumina microarray data. *Nucleic Acids Res* **36**, e11 (2008).

30.   McInnes L, H.J. UMAP: Uniform manifold approximation and projection for dimension
reduction. *arXiv preprint arXiv:1802.03426.* (2018).

31.   Becht, E. *et al.* Dimensionality reduction for visualizing single-cell data using UMAP. *Nat*
Biotechnol (2018).

32.   Brodeur, G.M. *et al.* Revisions of the international criteria for neuroblastoma diagnosis,
staging, and response to treatment. *Journal of Clinical Oncology* **11**, 1466-1477 (1993).

33.   Cohn, S.L. *et al.* The International Neuroblastoma Risk Group (INRG) classification system:
an INRG Task Force report. *J Clin Oncol* **27**, 289-97 (2009).

34.   Satija, R., Farrell, J.A., Gennert, D., Schier, A.F. & Regev, A. Spatial reconstruction of single-
cell gene expression data. *Nat Biotechnol* **33**, 495-502 (2015).

35.   Spranger, S. *et al.* Up-regulation of PD-L1, IDO, and T(regs) in the melanoma tumor
microenvironment is driven by CD8(+) T cells. *Sci Transl Med* **5**, 200ra116 (2013).

36.   Wherry, E.J. & Kurachi, M. Molecular and cellular insights into T cell exhaustion. *Nat Rev*
Immunol **15**, 486-99 (2015).

37.   Boeva, V. *et al.* Heterogeneity of neuroblastoma cell identity defined by transcriptional
circuitries. *Nat Genet* **49**, 1408-1413 (2017).

38.   van Groningen, T. *et al.* Neuroblastoma is composed of two super-enhancer-associated
differentiation states. *Nat Genet* **49**, 1261-1266 (2017).

39.   Bindea, G. *et al.* Spatiotemporal dynamics of intratumoral immune cells reveal the
1094   immune landscape in human cancer. *Immunity* **39**, 782-95 (2013).

1095   40.   Wang, L.L. *et al.* Augmented expression of MYC and/or MYCN protein defines highly
1096   aggressive MYC-driven neuroblastoma: a Children's Oncology Group study. *Br J Cancer*
1097   **113**, 57-63 (2015).

1098   41.   Casey, S.C. *et al.* MYC regulates the antitumor immune response through CD47 and PD-
1099   L1. *Science* **352**, 227-31 (2016).

1100   42.   Cursons, J. *et al.* A Gene Signature Predicting Natural Killer Cell Infiltration and Improved
1101   Survival in Melanoma Patients. *Cancer Immunol Res* **7**, 1162-1174 (2019).

1102   43.   Newman, A.M. *et al.* Robust enumeration of cell subsets from tissue expression profiles.
1103   *Nat Methods* **12**, 453-7 (2015).

1104   44.   Hu, X. *et al.* Landscape of B cell immunity and related immune evasion in human cancers.
1105   *Nat Genet* **51**, 560-567 (2019).

1106   45.   Ross, R.A., Spengler, B.A. & Biedler, J.L. Coordinate morphological and biochemical
1107   interconversion of human neuroblastoma cells. *J Natl Cancer Inst* **71**, 741-7 (1983).

1108   46.   Cohn, S.L. *et al.* Prolonged N-myc protein half-life in a neuroblastoma cell line lacking N-
1109   myc amplification. *Oncogene* **5**, 1821-7 (1990).

1110   47.   Debruyne, D.N. *et al.* ALK inhibitor resistance in ALK(F1174L)-driven neuroblastoma is
1111   associated with AXL activation and induction of EMT. *Oncogene* **35**, 3681-91 (2016).

1112   48.   Chipumuro, E. *et al.* CDK7 inhibition suppresses super-enhancer-linked oncogenic
1113   transcription in MYCN-driven cancer. *Cell* **159**, 1126-1139 (2014).

1114   49.   Durbin, A.D. *et al.* Selective gene dependencies in MYCN-amplified neuroblastoma
1115   include the core transcriptional regulatory circuitry. *Nat Genet* **50**, 1240-1246 (2018).

1116   50.   Margueron, R. & Reinberg, D. The Polycomb complex PRC2 and its mark in life. *Nature*
1117   **469**, 343-9 (2011).

1118   51.   Kroesen, M. *et al.* A transplantable TH-MYCN transgenic tumor model in C57Bl/6 mice for
1119   preclinical immunological studies in neuroblastoma. *Int J Cancer* **134**, 1335-45 (2014).

1120   52.   Weiss, W.A., Aldape, K., Mohapatra, G., Feuerstein, B.G. & Bishop, J.M. Targeted
1121   expression of MYCN causes neuroblastoma in transgenic mice. *EMBO J* **16**, 2985-95
1122   (1997).

1123   53.   Clarke, S.R. *et al.* Characterization of the ovalbumin-specific TCR transgenic line OT-I: MHC
1124   elements for positive and negative selection. *Immunol Cell Biol* **78**, 110-7 (2000).

1125   54.   Cibrian, D. & Sanchez-Madrid, F. CD69: from activation marker to metabolic gatekeeper.
1126   *Eur J Immunol* **47**, 946-953 (2017).

1127   55.   Uhrberg, M. The CD107 mobilization assay: viable isolation and immunotherapeutic
1128   potential of tumor-cytolytic NK cells. *Leukemia* **19**, 707-9 (2005).

1129   56.   Molfetta, R. *et al.* Regulation of NKG2D Expression and Signaling by Endocytosis. *Trends*
1130   *Immunol* **37**, 790-802 (2016).

1131   57.   Qi, W. *et al.* An allosteric PRC2 inhibitor targeting the H3K27me3 binding pocket of EED.
1132   *Nat Chem Biol* **13**, 381-388 (2017).

1133   58.   Dongre, A. & Weinberg, R.A. New insights into the mechanisms of epithelial-mesenchymal
1134   transition and implications for cancer. *Nat Rev Mol Cell Biol* **20**, 69-84 (2019).

1135   59.   Malladi, S. *et al.* Metastatic Latency and Immune Evasion through Autocrine Inhibition of
1136   WNT. *Cell* **165**, 45-60 (2016).

60. Pereira, B.I. *et al.* Senescent cells evade immune clearance via HLA-E-mediated NK and
CD8(+) T cell inhibition. *Nat Commun* **10**, 2387 (2019).

61. Shin, H. & Wherry, E.J. CD8 T cell dysfunction during chronic viral infection. *Curr Opin*
Immunol **19**, 408-15 (2007).

62. Kortlever, R.M. *et al.* Myc Cooperates with Ras by Programming Inflammation and
Immune Suppression. *Cell* **171**, 1301-1315 e14 (2017).

63. Sodir, N.M. *et al.* Endogenous Myc maintains the tumor microenvironment. *Genes Dev*
25, 907-16 (2011).

64. Bellmeyer, A., Krase, J., Lindgren, J. & LaBonne, C. The protooncogene c-myc is an
essential regulator of neural crest formation in xenopus. *Dev Cell* **4**, 827-39 (2003).

65. Rada-Iglesias, A. *et al.* Epigenomic annotation of enhancers predicts transcriptional
regulators of human neural crest. *Cell Stem Cell* **11**, 633-48 (2012).

66. Zeid, R. *et al.* Enhancer invasion shapes MYCN-dependent transcriptional amplification in
neuroblastoma. *Nat Genet* **50**, 515-523 (2018).

67. Chang, C.H., Hammer, J., Loh, J.E., Fodor, W.L. & Flavell, R.A. The activation of major
histocompatibility complex class I genes by interferon regulatory factor-1 (IRF-1).
Immunogenetics **35**, 378-84 (1992).

68. Unterholzner, L. *et al.* IFI16 is an innate immune sensor for intracellular DNA. *Nat*
Immunol **11**, 997-1004 (2010).

69. Spel, L. *et al.* Nedd4-Binding Protein 1 and TNFAIP3-Interacting Protein 1 Control MHC-1
Display in Neuroblastoma. *Cancer Res* **78**, 6621-6631 (2018).

70. Sen, T. *et al.* Targeting DNA Damage Response Promotes Antitumor Immunity through
STING-Mediated T-cell Activation in Small Cell Lung Cancer. *Cancer Discov* **9**, 646-661
(2019).

71. Ennishi, D. *et al.* Molecular and Genetic Characterization of MHC Deficiency Identifies
EZH2 as Therapeutic Target for Enhancing Immune Recognition. *Cancer Discov* **9**, 546-563
(2019).

72. Zingg, D. *et al.* The Histone Methyltransferase Ezh2 Controls Mechanisms of Adaptive
Resistance to Tumor Immunotherapy. *Cell Rep* **20**, 854-867 (2017).

73. Peng, D. *et al.* Epigenetic silencing of TH1-type chemokines shapes tumour immunity and
immunotherapy. *Nature* **527**, 249-53 (2015).

74. Xiao, L. *et al.* Adoptive Transfer of NKG2D CAR mRNA-Engineered Natural Killer Cells in
Colorectal Cancer Patients. *Mol Ther* **27**, 1114-1125 (2019).

75. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat*
Methods **9**, 676-82 (2012).

76. Kamentsky, L. *et al.* Improved structure, function and compatibility for CellProfiler:
modular high-throughput image analysis software. *Bioinformatics* **27**, 1179-80 (2011).

77. Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

78. Zhu, X. *et al.* Single-Cell Clustering Based on Shared Nearest Neighbor and Graph
Partitioning. *Interdiscip Sci* **12**, 117-130 (2020).

79. Russo, P.S.T. *et al.* CEMiTool: a Bioconductor package for performing comprehensive
modular co-expression analyses. *BMC Bioinformatics* **19**, 56 (2018).

80.   Yu, G., Wang, L.G., Han, Y. & He, Q.Y. clusterProfiler: an R package for comparing biological
1181   themes among gene clusters. *OMICS* **16**, 284-7 (2012).

1182   81.   Alexopoulos, E.C. Introduction to multivariate regression analysis. *Hippokratia* **14**, 23-8
1183   (2010).

1184   82.   Das, S. & Bansal, M. Variation of gene expression in plants is influenced by gene
1185   architecture and structural properties of promoters. *PLoS One* **14**, e0212678 (2019).

1186   83.   Debruyne, D.N. *et al.* BORIS promotes chromatin regulatory interactions in treatment-
1187   resistant cancer cells. *Nature* **572**, 676-680 (2019).

1188

1189

Methods

Cell culture

Human neuroblastoma (NB) cell lines (Kelly, NBL-S, CHP-212, SH-SY5Y, SH-EP, CHLA-20,

NB69, SK-N-FI) were obtained from the Children's Oncology Group cell line bank. ACN, GI-ME-

N, NB-EbC1 were kind gifts from A. Thomas Look and Kimberly Stegmaier at Dana Farber Cancer

Institute (DFCI). NB-9464 cells were provided by To-Ha Thai at Beth Israel Deaconess Medical

Center, Boston, MA. The cell lines were authenticated through STR analyses at the DFCI Core

facility and were routinely tested for mycoplasma. All NB cells were grown in RPMI-1640 medium

(Invitrogen) supplemented with 10% fetal bovine serum (FBS) (Invitrogen) and 1%

penicillin/streptomycin (Life Technologies). HEK293T cells obtained from the American Type

Culture Collection (ATCC) were grown in DMEM (Invitrogen) supplemented with 10% FBS and

1% penicillin/streptomycin (Life Technologies). SH-SY5Y cells resistant to the ALK inhibitor

ceritinib (LDK378) were described previously (Debruyne et al., 2016) and were grown in complete

RPMI-1640 in the presence of 1.5 µM ceritinib.

1204

Generation of PRRX1-inducible cell lines

Lentiviral vectors containing wild type and DNA-binding mutants of PRRX1 were generated by

cloning cDNAs encoding full length or homeodomain deletions of the human PRRX1A sequence

into the pInducer20 lentiviral plasmid (gift from Stephen Elledge, Addgene plasmid #44012). The

DNA-binding mutants harbor individual deletions of the three α-helices (ΔH1, ΔH2 and ΔH3) within

the PRRX1 homeodomain (amino acids (aa) 94-153). Amino acid boundaries of the deleted

regions are as follows: ΔH1 (aa 103-116); ΔH2 (aa 121-131); ΔH3 (aa 135-151). The lentivirus

was packaged by co-transfection of pInducer20 plasmid with the helper plasmids, pCMV-

deltaR8.91 and pMD2.G-VSV-G into HEK293T cells using the TransIT-LT1 Transfection Reagent

(Mirus Bio LLC). Virus-containing supernatants were collected 48 hr after transfection. SH-SY5Y

cells were transduced with the viral supernatant in the presence of 8 µg/ml polybrene (Sigma-

Aldrich) and 24 hr later selected using neomycin (G418) (5 µg/ml). Induction of gene expression

was achieved by treating cells every 2–3 days with doxycycline (dox; 200 ng/ml) in RPMI-1640

medium supplemented with 10% tetracycline-negative fetal bovine serum (tet-free FBS) (Gemini

Bio-Products) and 1% penicillin/streptomycin.

1220

MYCN shRNA knockdown and IFN-y treatment

The pLKO.1 shRNA construct targeting MYCN (TRCN0000020694) was purchased from Sigma-

Aldrich and the pLKO.1 GFP shRNA was a gift from D. Sabatini (Addgene plasmid #30323).

Lentiviral packaging was performed in HEK293T cells as decribed above and viral supernatant

was collected on days 2 and 3 after transfection. Kelly human NB cells were transduced with the

viral supernatant in the presence of 8 µg/ml polybrene (Sigma-Aldrich) and 24 h later selected

using puromycin (1 µg/ml) for 2 days. Puromycin-resistant Kelly cells were cultured for an

additional 6 days. Cells were then treated with 100 ng/ml of recombinant human IFNγ (Biolegend)

for 24h, following which they were harvested for analysis of RNA and protein expression.

1230

FACS analysis for cell surface protein staining

For each staining reaction 1 x 10^6 live cells were placed in a 12 x 75 mm polystyrene round bottom

tube (Falcon), resuspended in 100 µl 1x PBS and stained with the Zombie near-infrared (Zombie

NIR) viability dye (BioLegend) at a 1:1,000 dilution for 15 minutes at RT. Cells were then washed

once in FACS buffer (0.5% BSA in 1x PBS), resuspended in 100 µl of FACS buffer and incubated

in 5 µl of Human TruStain FcX™ (Fc receptor blocking solution, BioLegend) for 10 minutes at RT.

Next, appropriate volumes of conjugated fluorescent primary antibodies at predetermined

optimum concentrations were added and incubated on ice for 20 minutes in the dark. Cells were

then washed once in 2 ml of FACS buffer by centrifugation at 1500 rpm for 5 minutes. All FACS

samples were analyzed on a FACSCalibur flow cytometer (Becton Dickinson) using Cell Quest

software (Becton Dickinson). A minimum of 50,000 events was counted per sample and used for

further analysis. Data were analyzed using FlowJo v10 software (Becton Dickinson). The following

primary antibodies were used: PE-HLA (Biolegend; clone W6/32), PE-MICA/B (Biolegend clone

6D4), PE-ULBP2 (R&D Systems; clone 165903), PE-ULBP3 (R&D Systems; clone 166510), PE-

H-2Kb (Biolegend AF6-88.5), PE-H-2Kb SIINFEKL (Biolegend; clone 25-D1.16), PE-mouse

IgG2a k isotype control (Biolegend MOPC-173), PE- mouse IgG1 k isotype control (MOPC-21).

1247

Cell viability assay

SH-SY5Y parental and SH-SY5Y LDK-resistant cells were seeded in 96-well plates at a density

of 2×10^3 cells/well. After 24 h, cells were treated with increasing concentrations of LDK378

(ranging from 1 nM to 10 µM) dissolved in Dimethyl Sulfoxide (DMSO). DMSO solvent without the

drug served as a negative control. Following 72 h incubation, cells were analyzed for viability

using the CellTiter-Glo Luminescent Cell Viability Assay (Promega) according to the

manufacturer's instructions. Drug concentrations that inhibited cell growth by 50% (IC_{50}) were

determined using a non-linear regression curve fit with GraphPad Prism 8 software.

1256

Cell migration and invasion assays

Cell migration was measured using transwell chambers (Falcon). NB9464-H-2Kb^{lo} or

NB9464-H-2Kb^{hi} cells in serum-free medium (0.5×10^6 cells/ml) were added to the upper

chamber and inserts (8 µm pore size) were placed in the lower chamber containing medium

with 10% FBS. Following incubation at 37° C for 8 h, cells that migrated to the lower chamber

were fixed with methanol and stained with crystal violet (Sigma-Aldrich). The stained cells

were photographed with a light microscope at 100X magnification and migration was

quantified as the number of cells per high power field. Cell invasion was measured using the

fluorometric QCM^{TM} ECMatrix^{TM} Cell Invasion Assay (Millipore). NB9464-H-2Kb^{lo} or NB9464-

H-2Kb^{hi} cells in serum-free medium (0.5×10^6 cells/ml) were added to the upper chamber

and inserts (8 µm pore size) placed in the lower chamber containing medium with 10% FBS.

Following incubation at 37° C for 24 h, cell invasion was measured according to

manufacturer's instructions.

1270

1271   **EED226 treatment**

1272   5 x10^5 SH-SY5Y cells were seeded into 10 cm plates and treated with either 5 µM EED226

1273   (Selleck Chemicals) or DMSO (vehicle control) for 6-8 days, following which samples were

1274   harvested for downstream analyses. Cells were replenished with fresh media containing DMSO

1275   or EED226 every 2 days.

1276

1277   **Compounds**

1278   Ceritinib (LDK378) and EED226 were purchased from Selleck Chemicals. Doxycycline and

1279   dimethyl Sulfoxide (DMSO) was purchased from Sigma-Aldrich.

1280

RNA extraction and q-PCR

Total RNA was isolated using the RNAeasy Mini kit (Qiagen). Purified RNA was reverse

transcribed to cDNA using Superscript IV VILO master mix (Thermo Fisher Scientific) following

the manufacturer's protocol. Quantitative PCR was performed using 1 µl cDNA, 1x PowerTrack

SYBR Green PCR master mix (Thermo Fisher Scientific) and PCR primers (200 nM) in a total

volume of 25 µl and analyzed on a ViiA 7 Real-Time PCR system (Thermo Fisher Scientific). Each

individual biological sample was amplified in technical duplicate and normalized to GAPDH as an

internal control. Relative expression was calculated according to the $2^{-\Delta\Delta CT}$ quantification method

(Livak and Schmittgen, 2001). PCR primer sequences are shown in Table S5.

1290

1291

1292

Synthetic RNA spike-in and RNA-sequencing

1293

RNA-sequencing was performed on the following human NB cell lines: Kelly, NBL-S, CHP-212,

SH-SY5Y, SH-SY5Y LDK-resistant and SH-EP. Biological duplicates (5 x 10^6 cells per replicate)

were homogenized in 1 ml of TRIzol Reagent (Invitrogen) and purified using the mirVANA miRNA

isolation kit (Ambion) following the manufacturer's instructions. Total RNA was treated with DNA-

free™ DNase I (Ambion), spiked-in with ERCC RNA Spike-In Mix (Ambion) and analyzed on an

Agilent 2100 Bioanalyzer (Agilent Technologies) for integrity. Sequencing libraries were prepared

using LP-KAPA mRNA Hyper Prep and sequenced using Illumina HiSeq for 40 bases.

1301

Western blotting

1302

Cells were homogenized in NP40 lysis buffer (Life Technologies) containing 1× cOmplete EDTA-

free protease inhibitor cocktail and 1x PhosSTOP (Roche). Protein concentration was measured

using the DC Protein Assay (Bio-Rad). 100 µg total protein was denatured in LDS sample buffer

(Invitrogen), separated on pre-cast 4-12% Bis-Tris gels (Invitrogen) and transferred to

nitrocellulose membranes (Bio-Rad). Membranes were blocked using 5% dry milk (Sigma-Aldrich)

in Tris-buffered saline (TBS) supplemented with 0.2% Tween-20 (TBS-T) for 1 hr, and incubated

overnight with primary antibody in blocking buffer at 4 °C. Chemiluminescent detection was

performed with appropriate HRP-conjugated secondary antibodies and enhanced

chemiluminescence reagents (Thermo Scientific). Images were developed using Genemate Blue

ultra-autoradiography film (VWR).

1313

Antibodies

1314

1315 The following primary antibodies were used: MYCN (Cat #51705), MYC (13987), GATA3 (5852),

1316 TAP1 (12341), TAP2 (12259), LMP7 (13635), NOTCH1 (3608), cleaved NOTCH1 (4147), SOX9

1317 (82630), AXL (8661), GAPDH (2118), β-actin (3700), IRF1 (8478), VIM (5741), YAP1 (4912),

1318   TAZ1 (4883) (Cell Signalling Technologies (CST)); PHOX2B (Abcam;183741), FN1 (RnD

1319   systems; AF1918), LMP2 (Santa Cruz; 271354) and PRRX1 (Santa Cruz; 293386).

1320

Chromatin immunoprecipitation-quantitative PCR (ChIP-qPCR)

Soluble chromatin was prepared as above from SH-SY5Y cells without or with dox-inducible

PRRX1 expression (200 ng/ml dox for 10 days). ChIP was performed as described in the

preceding section using the following antibodies: H3K4me3 (Abcam 8580), H3K27me3 (Millipore

07-729), EZH2 (CST 5246), SUZ12 (CST 3737), EED (Millipore 17-10034), rabbit IgG (CST

2729). Purified ChIP DNA was dissolved in 60 µl of 1x TE. Quantitative PCR was performed on a

ViiA 7 Real-Time PCR system (Thermo Fisher Scientific) with 1 µl purified DNA, 1x PowerTrack

SYBR Green PCR master mix (Thermo Fisher Scientific) and PCR primers (200 nM) against the

genomic regions of interest. Each individual biological sample was amplified in technical

duplicate. Relative enrichment was quantified using the percent input method. PCR primer

sequences are shown in Table S5.

1332

Chromatin immunoprecipitation-sequencing (ChIP-seq)

Approximately 10-12 x 10^7 cells were crosslinked with 1% formaldehyde (Thermo Scientific) for

10 min at room temperature (RT) followed by quenching with 0.125 M glycine for 5 min. The cells

were then washed twice in ice-cold 1x Phosphate Buffered Saline (PBS), and the cell pellet

equivalent of 4 x 10^7 cells were flash frozen and stored at −80°C. Crosslinked cells were lysed in

lysis buffer 1 (50 mM HEPES-KOH pH7.5, 140 mM NaCl, 1 mM EDTA, 10% glycerol, 0.5% NP40,

0.25% Triton X-100). The resultant nuclear pellet was washed once in lysis buffer 2 (10 mM Tris-

HCl pH 8, 200 mM NaCl, 1 mM EDTA, 0.5 mM EGTA) and then resuspended in sonication buffer

(50 mM HEPES-KOH pH 7.5, 140 mM NaCl, 1mM EDTA, 1mM EGTA, 1% Triton X-100, 0.1%

sodium deoxycholate, 0.2% SDS). Chromatin was sheared using a Misonix 3000 sonicator

(Misonix) and at the following settings: 10 cycles, each for 30 s on, followed by 1 min off, at a

power of approximately 20 W. The lysates were then centrifuged for 15 min at 4 °C, supernatants

collected and diluted with an equal amount of sonication buffer without SDS to reach a final

concentration of 0.1% SDS. For each ChIP, the chromatin equivalent of 1×10^7 cells was used.

50 µl of Protein G Dynabeads per sample (Invitrogen) were blocked with 0.5% BSA (w/v) in 1x

PBS. Magnetic beads were loaded with the following antibodies: 10 µg of H3K27me3 (Millipore

07-729); 3 µg of H3K27ac (Abcam 4729), and 3 µg of H3K4me3 (Abcam 8580) and incubated

overnight at 4°C. The sonicated lysates were then incubated overnight at 4°C with the antibody-

bound magnetic beads, washed with low-salt buffer (50 mM HEPES-KOH (pH 7.5), 0.1% SDS,

1% Triton X-100, 0.1% sodium deoxycholate, 1 mM EGTA, 1 mM EDTA, 140 mM NaCl and 1×

complete protease inhibitor), high-salt buffer (50 mM HEPES-KOH (pH 7.5), 0.1% SDS, 1% Triton

X-100, 0.1% sodium deoxycholate, 1 mM EGTA, 1 mM EDTA, 500 mM NaCl and 1× complete

protease inhibitor), LiCl buffer (20 mM Tris-HCl (pH 8), 0.5% NP-40, 0.5% sodium deoxycholate,

1 mM EDTA, 250 mM LiCl and 1× complete protease inhibitor) and Tris-EDTA (TE) buffer. DNA

was then eluted in elution buffer (50 mM Tris-HCl (pH 8.0), 10 mM EDTA, 1% SDS), and high-

speed centrifugation performed to pellet the magnetic beads and collect the supernatants. The

crosslinking was reversed overnight at 65° C in the presence of 300 mM NaCl. RNA and protein

were digested using RNase A and proteinase K, respectively, and DNA was purified with phenol-

chloroform extraction and ethanol precipitation. Purified ChIP DNA was used to prepare Illumina

multiplexed sequencing libraries using the NEBNext Ultra II DNA Library Prep kit and the

NEBNext Multiplex Oligos for Illumina (New England Biolabs) according to the manufacturer's

protocol. Libraries with distinct indices were multiplexed and run together on the Illumina NextSeq

500 (SY-415-1001, Illumina) for 75 base pairs.

1366

IFN-gamma induction and antigen presentation in NB9464 cells

Approximately 1×10^6 cells were seeded onto 10 cm plates. 24 hr later, adherent cells were

treated with recombinant mouse IFN-γ (Biolegend) for 24h and harvested for H-2Kb analysis

using FACS as described above. For antigen presentation assays, cells treated with IFN-γ were

pulsed with SIINFEKL (OVA peptide) at 37°C. Cells were subsequently washed with 1x PBS to

remove unbound peptide and processed for analysis using FACS.

1373

T cell activation assays

OT-I T cell receptor (TCR) transgenic mice were purchased from Jackson Laboratories (Bar

Harbor, ME). Splenocytes were harvested and T cells were subsequently isolated from the

mononuclear layer using Ficoll separation and directly used in co-culture assays. Successful

enrichment of CD8$^+$ T cells was confirmed by FACS analysis using the FITC-CD8 antibody. Cells

were pulsed with SIINFEKL (OVA peptide) at 37 °C, to bind to cell surface H-2Kb. Cells were

subsequently washed with 1x PBS to remove unbound peptide and then co-cultured with

unstimulated OT-1 T cells for 24 h. OT-I cells were then harvested, and sequentially stained with

the Zombie NIR viability dye and FITC-CD8, PE-CD69 antibodies, followed by fixation with 1%

paraformaldehyde (Polysciences, Inc). OT-1 cells were analyzed by flow cytometry using a

FACSCanto II cell analyzer (Becton Dickinson) and FlowJo V10 software (Becton Dickinson).

1385

In vitro assays for NKG2D binding

SH-SY5Y parental, SH-SY5Y LDK-resistant, and SH-EP cells were incubated with recombinant

human NKG2D-Fc chimeric protein or an equivalent concentration of human IgG , following which

cells were washed, and sequentially stained with the Zombie NIR viability dye as described above

followed by incubation with an Alexa 647-conjugated anti-human IgG antibody for 30 minutes.

Cells were washed in FACS buffer and analyzed by flow cytometry using a FACSCanto II cell

analyzer (Becton Dickinson) and FlowJo V10 software (Becton Dickinson).

1393

1394

1395

NK cell degranulation assays

Human peripheral blood NK cells were isolated from blood collars using a RosetteSep™ human

NK cell enrichment cocktail (STEMCELL Technologies). NK cells were then co-cultured for 4 h

with confluent monolayers of SH-SY5Y parental, SH-SY5Y LDK-resistant, and SH-EP cells at in

the presence of CD107a antibody (Biolegend). Additionally, co-cultures of NK cells and 721.221

B cells were included as positive controls for degranulation. At the endpoint, NK cells were

harvested, stained with Zombie Yellow (Biolegend) and CD56 FITC (or NKp46 AlexaFluor 647^{TM})

and NKG2D PE (or mouse IgG1 PE) antibodies (Biolegend), followed by fixation with 1%

paraformaldehyde (Polysciences, Inc). For the NKG2D blocking assay, NK cells were incubated

with purified anti-NKG2D antibody or mouse IgG1 isotype control at 37 ºC, following which the

degranulation assay was performed as detailed above. NK cells were analyzed by flow cytometry

using a FACS Canto II (Becton Dickinson) and FlowJo V10 software (Becton Dickinson).

1408

Data sets

Publicly available RNA-seq data (GEO accession number GSE49711/GSE62564) from a cohort

of 498 primary human neuroblastoma tumors, microarray expression data from 394

neuroblastoma tumors (GSE120572) and 24 human neuroblastoma cell lines (GSE28019) were

accessed through the R2 genomics analysis and visualization platform

(https://hgserver1.amc.nl/cgi-bin/r2/). Clinical annotations for tumors were obtained from

GSE49711/GSE62564 regarding MYCN status (MYCN-nonamplified vs. MYCN-amplified, INSS

stage [high (stage 4) vs. low (1, 2, 3 and 4s], risk status (high vs. low) and age (< 18 months vs.

≥ 18 months).

1418

1419

1420

1421

Analysis of RNA-sequencing data

RNA-seq data processing and identification of differentially expressed genes

Single-end RNA-seq samples with 40 base pair (bp) read lengths were mapped to the human

genome (GRCh38) and ERCC spike-in sequences. Reads were mapped to the genome using

Bowtie2 (version 2.3.4.3) and default parameters. Reads that overlapped with the genomic

location for exonic regions were used to calculate gene counts with the FeatureCounts (Subread

package of version 1.6.3) package. Further, spike-in read counts were used for each sample to

normalize the library sizes. These read counts were used to calculate the sample-specific size

factor by using the function estimateSizeFactors (DESeq2) available in R. Normalized sample

coverage profiles were then created from previously determined size factors by using

bamCoverage (DeepTools v3.0.2) and parameters "--scaleFactor --skipNonCoveredRegions". To

check the reproducibility of biological replicates for each condition, principal component analysis

(PCA) and correlation (Spearman's rank coefficient) were assessed from the sample coverage

profiles at genome-wide scale and visualized using scatterplots and heatmaps. Because these

analyses showed a high correlation of sample coverage profiles between replicates, replicates

were merged using samtools merge and processed again as described for the individual

replicates. Next, differential gene expression analysis was performed using the DESeq2 in R. To

detect differentially expressed genes (DEGs) in each sample, raw read counts from RNA-seq

data were imported to the DESeq2 and the size factors calculated using the estimateSizeFactors

function. A transcript with an absolute log2 fold-change ≥ 1.5 and an adjusted *P*-value ≤ 0.01 was

considered significant.

1443

Enrichment analysis

Gene ontology enrichment for selected gene sets was performed by the Enrichr program

(https://amp.pharm.mssm.edu/Enrichr/). All GO terms were ranked based on the Enrichr

combined score, calculated by multiplying the adjusted *P*-value with the z-score using the Fisher's

exact test. The Fisher's exact test was used to determine significant overlaps between the queried

gene sets and other publicly available datasets. Enrichment of gene sets was considered

significant for an adjusted P-value ≤ 0.01, unless stated otherwise.

1451

Estimation of immune cell content in neuroblastoma tumors

Cell type identification by estimating relative subsets (CIBERSORT) [43], a deconvolution method

was used to evaluate immune cell fractions from gene expression data using the R package

'immunedeconv'. RNA transcript estimations were generated for all 498 neuroblastoma tumors

using the LM22 signature matrix available for 22 immune cell types. CIBERSORT was run in

"Absolute mode" with disabled quantile normalization as recommended for tumor RNA-seq data

and the overall immune content produced by the algorithm compared among tumors.

1459

ChIP-seq analysis

Data processing- All ChIP-seq raw datasets were processed as previously described [83]. The raw

read quality of the samples was accessed using the Fastqc tool (v0.11.7) to identify possible

sequencing errors and biases. Reads were aligned to the human genome (build hg19,

GRCh37.75) using the mapper Bowtie (v2.3.4.3) with default parameters. Unique and non-

duplicate reads that mapped to the reference genome were further processed using Samtools

(v1.9) and the MarkDuplicates (v2.1.1) command of Picard tools. Next, antibody enrichment in

each replicate as compared to input samples was verified using the PlotFingerprint command of

deepTools (v3.1.1). Peak caller MACS2 (2.1.1) was used to identity narrow peaks (H3K4me3 and

H3K27ac) with the parameters "--q 0.01--call-summits" and broad peaks (H3K27ac and

H3K27me3) with the parameters "--broad-cutoff 0.01". Peaks that overlapped with black-listed

regions (http://mitra.stanford.edu/kundaje/akundaje/release/blacklists/) of the reference genome

(mostly comprised of major satellite repeats of telomeric and pericentromeric regions) were

filtered out. Command "bamCompare" from the deepTools was used with the parameters "--

scaleFactorsMethod readCount --binSize 40 --operation subtract --smoothLength 80 --

extendReads 200" to create the input normalized bedgraph tracks for each replicate and

afterwards negative values were set to zero and counts were scaled to reads per million/base pair

(rpm/bp) to account for differences in the library size. Bigwig files were created for visualization

with bedGraphToBigWig. Subsequently, correlations among the ChIP-seq replicates were

accessed using bigwigs with the command "multiBigwigSummary" from deepTools and highly

correlated replicates merged at the BAM level. Peak identifications were then repeated in the

same manner for these merged BAM files.

1482

Identification of super-enhancer regions- Super-enhancers (SEs) were identified using the

ROSE algorithm (https://bitbucket.org/young_computation/rose/src/master/). Briefly, H3K27ac

binding regions identified by MACS2 as significant peaks, termed typical enhancers, were stitched

together if they were within 12.5 kb of each other. These stitched enhancers were ranked by

comparing the H3K27ac signal (density * length) with the input signal. The ROSE algorithm was

used to determine the inclination point for all stitched H3K27ac signals and to segregate regular

enhancers from SEs. To compare SEs in 5Y-parental, 5Y-LDK-resistant and SH-EP cells, the

same maximum threshold was used between the conditions.

1491

Analysis of histone binding changes between lineage states- To analyze the changes in

occupancies of active (H3K27ac and H3K4me3) and repressive (H3K27me3) histone marks

during the transition from adrenergic (5Y-par.) to mesenchymal (5Y-LDK-res., SHEP) states, we

compared the peaks of histone marks identified by MACS2 at the promoter regions. For this

purpose, we first extracted the promoter regions ± 2 kb with respect to the TSS (-2kb upstream

to +2kb downstream) of all annotated protein coding genes and subsequently, retrieved the peaks

of H3K27ac, H3K4me3 and H3K27me3 from 5Y-par., 5Y-LDK-res. and SH-EP cells. Now, to

determine the differential binding of each histone mark between 5Y-par. and 5Y-LDK-res. cells,

we first combined all significant peaks called by MACS2 at the promoter regions and merged the

peak regions that overlapped by at least 50%. This 50% threshold was used to avoid merging

peaks that had clear and distinct summits. Next, the normalized active or repressive histone

marks read densities were calculated for each region and a ratio of [log2 (5Y-LDK-res./ 5Y-par.)]

was calculated. Shared peaks had similar enrichment of either active or repressive histone marks

in both the cell types. Similar comparisons were made for active or repressive histone marks

between 5Y-par. and SH-EP cells. To further compare changes in all histone marks at the

promoters of immune genes, the gain of significant H3K27ac, H3K4me3 binding and loss of

H3K27me3 signals were listed in mesenchymal (5Y-LDK-res., SHEP) cells as compared to

adrenergic (5Y-par.) cells.

1510

Integrated analysis of histone binding and gene expression- Cell-type specific differential

enrichment of H3K27me3 and H3K4me3 binding in 5Y-par., 5Y-LDK-res. and SH-EP cells was

determined by calculating the log2 (H3K4me3+1/H3K27me3+1) ratios in the promoter regions

(TSS ± 2 kb) of immune genes. Next, to examine the association between gene expression and

differential enrichment of H3K27me3 and H3K4me3 binding in immune genes in 5Y-par., 5Y-LDK-

res. and SH-EP cells, genes were ranked based on their expression values and plotted against

the calculated ratios between the histone marks.