1   **Title Page**

3   **TAK1 blockade as a therapy for retinal neovascularization**

Fan-Li Lin[1,2†], Jiang-Hui Wang[3†], Jinying Chen[2,4], Linxin Zhu[2,4], Yu-Fan Chuang[1,2], Leilei Tu[4],

Chenkai Ma[5], Suraj Lama[2], Damien Ling[3], Raymond Ching-Bong Wong[3,6], Alex W. Hewitt[2,3], Ching-

Li Tseng[7], Bang V. Bui[8], Peter van Wijngaarden[3,6], Gregory J. Dusting[3,6], Peng-Yuan Wang[1], Guei-

Sheung Liu[2,6,9*]

[1]Shenzhen Key Laboratory of Biomimetic Materials and Cellular Immunomodulation, Shenzhen

Institute of Advanced Technology, Chinese Academy of Sciences, Shenzhen, China

[2]Menzies Institute for Medical Research, University of Tasmania, Hobart, Australia

[3]Centre for Eye Research Australia, Royal Victorian Eye and Ear Hospital, East Melbourne, Australia

[4]Department of Ophthalmology, the First Affiliated Hospital of Jinan University, Guangzhou, China

[5]Molecular Diagnostics Solutions, CSIRO Health and Biosecurity, North Ryde, New South Wales,

Australia

[6]Ophthalmology, Department of Surgery, University of Melbourne, East Melbourne, Victoria,

Australia

[7]Graduate Institute of Biomedical Materials and Tissue Engineering, College of Biomedical

Engineering, Taipei Medical University, Taipei, Taiwan

[8]Department of Optometry and Vision Sciences, University of Melbourne, Parkville, Victoria,

Australia

[9]Aier Eye Institute, Changsha, Hunan, China

†FLL and JHW contributed equally to this work.

*Correspondence and requests for materials should be addressed to Dr Guei-Sheung Liu

(rickliu0817@gmail.com). Menzies Institute for Medical Research, University of Tasmania. Address:

17 Liverpool Street, Hobart, TAS 7000, Australia. Tel: +61362264250.

Running title: Targeting TAK1 alleviates retinal neovascularization

ABSTRACT

Retinal neovascularization, or pathological angiogenesis in the retina, is a leading cause of blindness

in developed countries. Transforming growth factor-β-activated kinase 1 (TAK1) is a mitogen-

activated protein kinase kinase kinase (MAPKKK) activated by TGF-β1 and other pro-inflammatory

cytokines. TAK1 is also a key mediator of inflammation, innate immune responses, apoptosis and

tissue homeostasis and plays an important role in physiological angiogenesis. Its role in pathological

angiogenesis, particularly in retinal neovascularization, remains unclear. We investigated the

regulatory role of TAK1 in pathological angiogenesis in the retina. Transcriptome analysis of human

retina featuring retinal neovascularization revealed enrichment of known TAK1-mediated signaling

pathways. Selective inhibition of TAK1 activation by 5Z-7-oxozeaenol attenuated aberrant retinal

angiogenesis in rats following oxygen-induced retinopathy. Transcriptome profiling revealed that

TAK1 activation in human microvascular endothelial cells under TNFα stimulation led to increase the

gene expression related to cytokines and leukocyte-endothelial interaction, mainly through nuclear

factor kappa B (NFκB) signaling pathways. These results reveal that inhibition of TAK1 signaling

may have therapeutic value for the treatment of pathological angiogenesis in the retina.

INTRODUCTION

Retinal neovascularization is a severe complication of several ocular diseases, often resulting in permanent damage to the neurosensory retina and irreversible vision loss (1). Retinal neovascularization shares similar characteristics to pathological angiogenesis elsewhere in the body, exhibiting features including endothelial proliferation and migration, elevated vascular permeability and inflammation; all processes in which vascular endothelial growth factor (VEGF) plays a crucial role (2). Recently, the advent of ocular anti-VEGF therapies has been a significant advance in mitigating the risk of vison loss associated with retinal neovascularization. Despite this success, there is increasing evidence that some patients do not respond well to anti-VEGF treatments (3). Therefore, it is necessary to seek other therapeutic targets.

The common pathways shared between inflammation and angiogenesis play critical roles in retinal neovascularization (4). Several studies have reported increased expression of various pro-inflammatory cytokines, chemokines, pro-angiogenic factors and adhesion molecules in patients with retinal neovascularization, such as tumor necrosis factor α (TNFα), interleukin-1 (IL-1), chemokine (C-X-C motif) ligand 8 (CXCL8; known as IL-8), vascular endothelial growth factor (VEGF), as well as intercellular adhesion molecule 1 (ICAM-1 and VCAM-1) (5-9). TNFα was found to suppress the expression of tight junction proteins and is required for VEGF-mediated hyperpermeability in endothelial cells and the breakdown of the blood-retinal barrier in patients with diabetic retinopathy (DR) (10, 11). Interleukins are also primary mediators of inflammation that can promote angiogenesis via induction of proangiogenic factors such as VEGF (4). Moreover, TNFα and IL-1 stimulates expression of ICAM-1, CXCL8 and VEGF in endothelial cells and microglia cells, both of which promote progression of retinal neovascularization (12, 13). A key role for inflammatory mediators is highlighted by their growing use as prognostic markers of the severity of proliferative diabetic retinopathy (PDR) (4).

Transforming growth factor-β-activated kinase 1 (TAK1), a member of the mitogen-activated protein kinase (MAPK) family, is a critical serine/threonine kinase in several cellular signaling pathways. Such pathways can be activated by diverse pro-inflammatory stimuli including TNFα, IL-1, transforming growth factor beta (TGF-β), or toll-like receptor (TLR) ligands. The engagement of TAK1 in turn activates downstream signaling pathways, including nuclear factor kappa B (NFκB) and MAPK p38, extracellular signal-regulated kinase (ERK), and c-Jun kinase (JNK) signaling, modulating inflammatory responses and cell survival (14-16). TAK1 inhibition has been shown to reduce TNFα and CXCL8 secretion in leukocytes (17) and to suppress prostaglandin-endoperoxide synthase 2 (PTGS2; known as cyclooxygenase-2, COX-2), ICAM-1, and VEGF expression in cancer cells through NFκB signaling pathway (18). Expression of TAK1 provides protection for endothelial

cells against TNFα-induced apoptosis under inflammatory conditions (19). Indeed, TAK1 deletion

causes embryonic lethality as the capacity to prevent TNFα-induced endothelial cell death and vessel

regression is lost. Moreover, TAK1 is also important for supporting TNFα-independent vascular

formation and endothelial migration. These broad roles for TAK1 suggests that endothelial TAK1

inhibition may be a useful alternate anti-angiogenesis target in retinal disease (19, 20).

We hypothesized that TAK1 plays a crucial role in pathological angiogenesis and TAK1 inhibition

can suppress retinal neovascularization. Our findings indicate that TAK1 is an important regulator of

inflammatory and angiogenesis-related pathways that contribute to retinal neovascularization.

Pharmacological inhibition of TAK1 alleviated aberrant retinal angiogenesis by modulating cytokine

signaling and leukocyte-endothelium interactions in a rat model of ischemia-induced retinopathy.

RESULTS

TAK1 is expressed in human retina and is positively associated with retinal neovascularization

To understand TAK1 expression profiles in the human retina, we first analyzed published single nuclei

RNAseq (snRNAseq, GSE135133) data to examine TAK1 expression in different retinal cells.

Analysis of 121,799 individual cells from human retina revealed that both *TAK1* (*MAP3K7*) and its

binding protein, *TAB1-3,* are expressed moderately in most neural and non-neural retinal cells (**Fig.**

1A and S1). Notably, among the retinal cells examined, TAK1 is highly enriched in endothelial cells.

Immunostaining suggests that TAK1 is expressed in various layers of the human retina in both neural

and non-neural cells, such as mural cells and smooth muscle cells (**Fig. 1B**) in larger vessels (diameter

of artery or vein ranging from 0.6 - 16 mm) (21). We also found that co-localization of TAK1 and

endothelial cell marker CD31 was more likely to be found in the microvasculature, including arterioles

or venules (20 - 25 μm) and capillaries (9 μm).

To identify the key molecular signaling or biological pathways relevant to the angiogenic process

in patients with retinal neovascularization, a transcriptome dataset from humans with proliferative

diabetic retinopathy (PDR) was interrogated by taking advantage of Gene Set Enrichment Analysis

(GSEA), a knowledge-based approach for interpreting genome-wide expression profiles (22).

Compared with controls, those with retinal neovascularization showed enrichment of known TAK1

upstream signaling pathways, including IL1, TNF, TGF-β, TLR, T-cell receptor (TCR) and B-cell

receptor (BCR) (**Fig. 1C and S2**). Pathways related directly to TAK1 were enriched, such as "TAK1

activation of NFκB by phosphorylation and activation of inhibitor of κB (IκB) kinase (IKK) complex",

"activated TAK1 mediated p38 MAPK activation", and "JNK kinases phosphorylation and activation

mediated by activated TAK1" (**Fig. 1C and 1D**). In general, genes in association with TAK1 activation mediated by NFκB and MAPKs phosphorylation were up-regulated in patients with retinal neovascularization compared with controls (**Fig. S3**). There was negative enrichment of genes associated with several biological pathways in relation to visual phototransduction and neuronal systems (**Fig 1C**).

TAK1 is up-regulated in neovascular tufts and positively associated with retinal neovascularization in ischemia-induced retinopathy in rat

To investigate the role of TAK1 in pathological neovascularization in the retina, we subsequently used a rat model of oxygen-induced retinopathy (OIR) which yields ischemic avascular zones and preretinal neovascularization similar to that observed clinically in PDR and retinopathy of prematurity. Neonatal rats were subjected to daily cycles of 80% oxygen for 21 hours and ambient air for 3 hours from postnatal day 0 (P0) to P14 to induce vaso-obliteration. At P14 rats were returned to room air, where maximal preretinal neovascularization occurred by P18, which was followed by a phase of vascular regression (**Fig. 2A**). A significant increase in TAK1 expression at both RNA and protein levels was observed in OIR retinae at P18 compared with controls (**Fig. 2B and 2C**). Examination of the retinal distribution of TAK1 expression revealed that TAK1 expression was uniquely expressed in areas of active neovascularization (**Fig. 2D**). To gain further insight into the role of TAK1 activation at the peak of retinal neovascularization (P18), we then performed transcriptome analyses using bulk-RNA sequencing. GSEA-based analysis revealed a positive enrichment of several TAK1 upstream pathways, including IL1, TNF, TCR, BCR and TLR (**Fig. 2E and S4**), which is consistent with our findings in patients with retinal neovascularization due to PDR. Similar to findings in humans, there was a negative enrichment of genes associated with phototransduction and neuronal systems at P18 in the rats. Of note, certain canonical pathways and biological processes related to angiogenesis were positively enriched in the OIR model, such as VEGF-A/VEGFR2, PDGF pathway, the cellular response to hypoxia, angiogenesis and inflammation (**Fig. 2E and S4**). Overall, our data implicates active involvement of TAK1 in retinal neovascularization.

Inhibition of TAK1 activation attenuates retinal neovascularization in rat OIR model

Given that the ablation of TAK1 in endothelial cells impairs angiogenesis and causes lethality (19, 20), we therefore used a selective inhibitor of TAK1, 5Z-7-oxozeaenol, a resorcylic acid lactone derived from a fungus (23), to assess whether TAK1 signaling blockade can alleviate retinal neovascularization. 5Z-7-oxozeaenol was administered intravitreally to the OIR rats at P14 (**Fig. 3A**). Four days after drug

administration, retinal neovascularization was significantly suppressed by 51.3% and 57.1% in OIR rats that had received low (18 ng) and high doses (90 ng) of 5Z-7-oxozeaenol, respectively, compared to OIR rats that had vehicle only (**Fig. 3B and S5**). No significant difference in retinal vaso-obliteration was found among these groups (**Fig. 3C**). We also examined the development of deep capillary beds to assess the effects of TAK1 inhibition in physiological vessel growth (24). Normal control rats developed a complete deep layer vasculature, while such development was severely disrupted in OIR rats (**Fig. 3D**). We found that 5Z-7-oxozeaenol treatment did not improve or damage the formation of deep capillary beds in OIR retina, which is likely attributed to suppression of proliferative vessels by TAK1 inhibition. Moreover, the rats receiving low or high dose of 5Z-7-oxozeaenol did not show any abnormalities in the retinal vasculature under normoxic conditions (**Fig. S6**).

TAK1 deletion impairs the expression of multiple cytokines and cell adhesion molecules in endothelial cells responding to inflammatory insults

Given that TAK1 blockade prevented retinal neovascularization and previous studies have indicated that TAK1 is activated in response to various pro-inflammatory stimuli such as TNFα, IL-1β, TGF-β, or TLR ligand, we evaluated what signaling regulates TAK1 in the endothelial response to inflammation. We first generated TAK1-knockout (KO) human telomerase-immortalized microvascular endothelial (TIME) cells (TAK1-KO) using CRISPR/Cas9 (**Fig. S7**). To explore the effect of TAK1 deletion in human microvascular endothelial cells, we generated a transcriptome profile through bulk-RNA sequencing of wild-ype TIME cells (WT) and TAK1-KO TIME cells (TAK1-KO). We found that 2178 genes (743 upregulation and 1435 downregulation) were differentially expressed [false discovery rate (FDR), <0.05] following TAK1-KO (**Fig. S8A and S8B**). GSEA results suggested that TAK1 knockout negatively affected pathways related to tissue immune responses, cell adhesion, and cell-extracellular matrix interaction (**Fig. S8C**).

Since TNFα is strongly correlated with progression of retinal neovascularization and is a major activator of TAK1 signaling (10), we performed bulk-RNA sequencing to obtain transcriptomes from wild-type (WT) TIME cells and TAK1-KO TIME cells that were in the presence or absence of TNFα (**Fig. 4A**). We found that 462 and 3609 genes were differentially expressed (FDR<0.05 and Log2FoldChange >1 or <-1) in wild-type cells upon TNFα treatment (WT TNF *vs*. WT; hereafter referred to as the WT TNF set) and TAK1-KO cells upon TNFα treatment (TAK1-KO TNF *vs*. WT TNF; hereafter referred to as the TAK1-KO TNF set), respectively (**Fig. 4B**). To further explore the biological significance of our findings in differential gene expression, we undertook GSEA of WT TNF and TAK1-KO TNF sets. GSEA results showed the top 20 KEGG (Kyoto Encyclopedia of Genes and Genomes) gene sets were positively enriched due to TNFα stimulation in WT cells (**Fig. 4C**).

However, in TAK1-KO cells exposed to TNFα, 7 of the top 10 gene sets were negatively enriched,
suggesting that these biological functions (highlighted in red in **Fig.4C**) induced by TNFα were
adversely affected in cells due to TAK1 knockout. The gene sets related to cytokines ("Cytokine-
cytokine receptor interaction") and cell adhesion molecules ("Cell adhesion molecules-CAM") were
positively enriched in the wild-type cells exposed to TNFα yet not in TAK1-KO cells exposed to TNFα
(**Fig. 4C**), indicating the expression of cytokines and cell adhesion molecules induced by TNFα were
negatively affected in human microvascular endothelial cells due to TAK1 knockout. We further
analyzed the target genes of TAK1 in these two gene sets by a Venn diagram. A total of 74 cytokine-
related genes and 29 cell adhesion molecule-related genes were positively enriched in TNFα-treated
WT cells (**Fig. 4D and 4E**). Among of these genes, 59 cytokine-related genes and 23 cell adhesion
molecule-related genes were negatively enriched in TNFα-treated TAK1-KO cells (**Fig. 4D and 4E**).
Many of these genes are related to inflammation and angiogenesis, suggesting that TAK1 may be
involved in the regulation of the production of proinflammatory/proangiogenic cytokines (such as
CXCL8 and *VEGFA*), as well as leukocyte-endothelial interactions (leukostasis; such as *ICAM-1* and
VCAM-1) in retinal neovascularization (**Fig. 4F**).

TAK1 is required for activation of NFκB signaling in mediating the expression of multiple
cytokines and cell adhesion molecules in endothelial cells responding to inflammatory insults
NFκB and MAPKs signaling are the primary known downstream pathways mediated by TAK1
activation, both of which have been shown as key mediators involved in several pathological
conditions associated to angiogenesis, such as arthritis and cancer (25, 26) (**Fig. 5A**). We therefore
investigated whether the identified 59 cytokine-related genes and 23 cell adhesion molecule-related
genes from TNFα-treated TAK1-KO cells are regulated by NFκB and MAPKs signaling pathway. We
found that about 75% (38/59) of selected genes from the "cytokine gene set" and 35% (8/23) of
selected genes from the "adhesion molecule gene set" are known to be regulated by NFκB and activator
protein-1 (AP-1, a transcription factor regulates by MAPK signaling cascade), both of which were
identified from a NFκB target gene database (27) and ENCODE Transcription Factor Targets Database,
respectively (**Fig 5B and Table 1**).

To confirm the impairment of the activation of NFκB and MAPKs signaling caused by TAK1
knockout, we characterized the activation of NFκB and MAPKs in TNFα-treated TAK1-KO cells.
TNFα treatment significantly increased TAK1 phosphorylation in WT cells, an effect not observed in
TAK1-KO cells (**Fig. 5C and 5D**). IκB degradation and NFκB p65 phosphorylation are key events in
NFκB signaling (28). TNFα treatment significantly induced NFκB p65 phosphorylation and IκB

degradation in WT cells, while no such changes occurred in TAK1-KO cells (**Fig. 5C and 5E**). Likewise, phosphorylation of JNK, p38, and ERK in MAPK signaling was not increased in TNFα-treated TAK1-KO cells compared to TNFα-treated wild-type cells (**Fig. 5C and 5F**). Collectively, these data indicate that TAK1 is required for phosphorylation of the key mediators of NFκB and MAPK signaling in human microvascular endothelial cells.

Expression of the most significant genes (those also shown to be involved in retinal neovascularization) from the gene sets related to cytokines ("Cytokine-cytokine receptor interaction") and cell adhesion molecules ("Cell adhesion molecules-CAM") was also validated using qRT-PCR. Upon TNFα stimulation, the expression level of *CXCL8*, *VEGF-A, ICAM-1*, and *VCAM-1*, were substantially elevated in WT cells, while no TNFα induced change was observed in TAK1-KO cells (**Fig. 5G**). Similar results were found in IL1β-treated WT cells but not in IL1β-treated TAK1-KO cells (**Fig. S9**). Likewise, TAK1 knockdown using TAK1 siRNA (siTAK1) or TAK1 inhibitor (5Z-7-oxozeaenol) resulted in a similar effect in primary human retinal microvascular endothelial cells (HRMECs) (**Fig. S10 and S11**). These data further confirmed that TAK1 is required for activation of NFκB/AMPKs signaling to regulate expression of multiple cytokines and cell adhesion molecules in human microvascular endothelial cells responding to inflammation.

Pharmacological inhibition of TAK1 suppresses the expression of inflammatory cytokines and reduces leukocyte-endothelial interactions (leukostasis) in the rat OIR model, while has no impact on retinal structure and function

To further confirm the effect of TAK1 inhibition in suppressing the expression of inflammatory cytokines and reduces leukostasis in the development of retinal neovascularization, OIR rats were treated with 5Z-7-oxozeaenol at P14 and the retinae were harvested for qPCR and histological analysis at the time points indicated in the figure (**Fig. 6A**). Two days after drug administration, TAK1 inhibition attenuated the expression of inflammatory and angiogenic genes, including *TNFα*, *VEGFA* and *ICAM-1*, in the retina of OIR rats that had received low (18 ng) and high doses (90 ng) of 5Z-7-oxozeaenol, respectively, compared to OIR rats that had vehicle only, suggesting that 5Z-7-oxozeaenol can mitigate the production of inflammatory and angiogenic cytokines in the progression of retinal neovascularization (**Fig. 6B**). Since TAK1 was found to modulate the expression of cell adhesion molecules in human endothelial cells responding to proinflammatory cytokine, we therefore further evaluated the event of microglial adhesion to the vascular surface (leukostasis). Co-immunostaining of isolectin GS-IB4 (blood vessel) and Iba1 (a marker of microglia) illustrated that 5Z-7-oxozeaenol treatment at both low and high doses significantly attenuated the microglial adhesion to the vascular

surface in OIR rats (**Fig. 6C and 6D**). Together, our data suggest that active involvement of TAK1 in

the regulation of proinflammatory and proangiogenic cytokine expression as well as modulation of

leukostasis in the retinal vasculature. Thus, 5Z-7-oxozeaenol is therapeutically effective to suppress

retinal neovascularization in the rat model of retinopathy (**Fig. 6E**).

In addition to examining the therapeutic efficacy of 5Z-7-oxozeaenol in retinal neovascularization,

we further performed safety evaluations to test whether 5Z-7-oxozeaenol adversely affects rodent

retinae. Electroretinography (ERG) and optical coherence tomography (OCT) were performed to look

for any retinal functional or structural changes in rats 28 days post intravitreal injection of high dose

of 5Z-7-oxozeaenol. No difference in function was observed for photoreceptors, bipolar cells or

ganglion cells between vehicle- or 5Z-7-oxozeaenol-treated eyes (**Fig. S13A-S13F**). No significant

difference was observed in total retinal thickness or thickness of various retinal layers including the

nerve fiber layer (NFL) and outer retina layer (ORL) between 5Z-7-oxozeaenol- and vehicle-treated

eyes (**Fig. S13G and S13H**). Together, these results suggested that intravitreal injection of 5Z-7-

oxozeaenol does not result in adverse or toxic effects on retinal function and structure up to 28 days

post-injection.

DISCUSSION

TAK1 is a key modulator involved in a range of cellular functions including the immune response (14),

cell survival and death (16), angiogenesis (20), fibrosis (29) and tumor metastasis (30). TAK1 is

activated in response to various stimuli such as pro-inflammatory cytokines, hypoxia, oxidative stress,

metabolism, DNA damage, Wnt and osmotic shock (14, 31-33). Whether TAK1 is involved in

angiogenesis or inflammatory signaling in the retina, much less the molecular mechanisms by which

this occurs, has not been studied. Here, we show that TAK1 plays a central role in regulating

inflammatory and angiogenic signaling in the rat retina, and that the same signaling appears to be

involved in human diabetic retinopathy. Pharmacological antagonism of TAK1 signaling by 5Z-7-

oxozeaenol was shown to be a potential therapeutic option for retinal neovascularization.

NFκB and MAPK are critical regulators of stress responses, immunity, inflammation and

angiogenesis. A variety of proinflammatory cytokines such as TNFα and IL1β activate these signaling

pathways through a common upstream kinase TAK1 (34). For instance, TNFα binds to its receptor 1

(TNFR1) upon inflammatory stimuli, triggering TAK1 that in complex with TAB1-3 phosphorylate

IKK at Ser^{177} and Ser^{181}, which in turn leads to rapid proteasome-mediated degradation of the IκB

subunit. Subsequently, NFκB (p50/p65) is activated and translocated to the nucleus to promote gene

transcription (17). In addition, the TAK1 complex phosphorylates MAPKs to activate the JNK

pathway and the p38 MAPK pathways (both mediated by MKK4 and MKK7) as well as ERK by an

IKK2/tumor progression locus 2 (TPL2)-dependent pathway (35). Phosphorylation of p38 affects F-

actin polymerization in vascular endothelial cells and endothelial migration (36). Activation of JNK

and ERK subsequent to p21-activated kinase 1 stimulation promotes endothelial migration through

enhancement of cell motility (37). In particular, ERK is involved in VEGF-induced endothelial

proliferation, migration (via actin polymerization, lamellipodia and filopodia formation and myosin

contraction), morphogenesis (tube formation) and extracellular matrix degradation during

angiogenesis (37). In fact, deletion of TAK1 leads to the inactivation of both JNK and NFκB signaling,

as well as downregulation of expression of pro-survival genes in mice (38). Hence, deletion of TAK1

causes bone marrow and liver failure in mice due to the massive apoptotic death of hematopoietic cells

and hepatocytes, ultimately resulting in death within a few days. Moreover, endothelial-specific TAK1

knockout mice demonstrated intestinal and liver hemorrhage secondary to apoptosis of endothelial

cells and damaged vessels, resulting in rapid death (19). These data suggest that TAK1 protects

endothelial cells and maintains vascular integrity and homeostasis under inflammatory conditions.

Here, we provide the first evidence that both genetic and pharmacological inhibition of TAK1

activity in human microvascular endothelial cells prevents inflammatory stimuli from inducing

sequential phosphorylation of upstream kinases in both NFκB (p65) and MAPKs (JNK, p38 and ERK)

pathways and suppressed expression of downstream genes that drive inflammation and angiogenesis.

GSEA results confirmed that TAK1 knockout prevented TNFα induced cytokines and cell adhesion

molecules expression predominantly through the NFκB pathway, resulting in attenuated inflammation

and angiogenesis. In addition, TAK1 inhibition by 5Z-7-oxozeaenol in OIR rat significantly reduced

cytokines expression and adhesion of microglia to retinal endothelial cells. These data together suggest

a significant role for TAK1 in mediating crosstalk between inflammatory and angiogenic pathways,

both of which are crucial in pathological angiogenesis.

TAK1 is thus a central regulator of cell survival and death (16). 5Z-7-oxozeaenol, a resorcylic

acid lactone derived from a fungus, acts as a potent and selective inhibitor of TAK1, which displays

more than 33- and 62-fold selectivity for TAK1 over binding of MEKK1 and MEKK4, respectively

(23). It forms a covalent complex with TAK1 and impedes both the kinase and ATPase activity of

TAK1 following bi-phase kinetics (39). We demonstrated that intravitreal injection of 5Z-7-

oxozeaenol suppressed retinal neovascularization, and modulated elevated TNFα expression in the

retina of OIR rats, without causing degeneration of the normal vasculature. These data suggest that

TAK1 inhibition by 5Z-7-oxozeaenol may more specifically attenuate active angiogenesis with less

risk of damage to normal endothelial cells. However, further studies are required to understand the

effects of repeated 5Z-7-oxozeaenol treatment. Moreover, TAK1 Inhibition by 5Z-7-oxozeaenol is
known to block pro-inflammatory signaling by selectively inhibiting TAK1 MAPKKK (23). In an
animal model of early brain injury, 5Z-7-oxozeaenol was shown to inhibit subarachnoid hemorrhage-
induced phosphorylation of p38 and JNK, the nuclear translocation of NFκB p65 and degradation of
IκB, ultimately reducing neuronal apoptosis and early brain injury (40). Similarly, another study
demonstrated that 5Z-7-oxozeaenol alleviated experimental autoimmune encephalomyelitis by
reducing the levels of pro-inflammatory cytokines in splenocytes and the central nervous system, thus
reducing the number of activated microglia and blocking p38 MAPK, JNK, and ERK signaling
pathways (41). Overactive microglia can damage surrounding cells, via a range of detrimental effects
including release of pro-inflammatory cytokines, reactive nitrogen species, reactive oxygen, and
proteolytic enzymes (42). We also observed diminished adhesion of microglia to endothelial cells in
OIR rats receiving 5Z-7-oxozeaenol treatment, suggesting that 5Z-7-oxozeaenol alleviates
inflammation in the retina, in part by reducing activation of microglia. Altogether, our results indicate
that TAK1 inhibition by 5Z-7-oxozeaenol suppresses ocular inflammation and neovascularization by
inhibiting cellular proliferation and microglia activation through NFκB pathway.

5Z-7-oxozeaenol has been previously examined in a number of animal studies focusing on brain
injury (43-45). One study demonstrated that 5Z-7-oxozeaenol has neuroprotective function in a mouse
model of stroke (43). However, others suggest long-term TAK1 inhibition may not be protective in
neurodegenerative disorders (46). Chronic TAK1 inhibition resulted in apoptosis of a number of
normal cells, including keratinocytes, hepatocytes, and hematopoietic cells, raising safety concerns
(38, 47, 48). We demonstrated that TAK1 inhibition was capable of simultaneously reducing
proinflammatory cytokines and proangiogenic cytokines in the retina of OIR rats. In our studies, no
adverse effects of intravitreal injection of the TAK1 inhibitor on retinal function or structure were
apparent up to 28 days post-injection. Specifically, TAK1 inhibition by intravitreal injection of 5Z-7-
oxozeaenol did not affect the scotopic threshold response, a highly sensitive electroretinography
measure of retinal ganglion cell activity (49). Another study has shown that systemic TAK1 inhibition
by 5Z-7-oxozeaenol preserved β cell function, thus improving overall health condition in the animal
with autoimmune diabetes (50). It is worth noting that 5Z-7-oxozeaenol may have off-target effects,
potentially stemming from cross-reactivity with other human kinases that possess an analogous
cysteine (51). Whilst encouraging, further studies with chronic dosing of 5Z-7-oxozeaenol are clearly
needed to confirm safety.

In summary, the present study identifies TAK1 as a mediator of inflammatory and angiogenic
signaling pathways, with a central role in retinal neovascularization. We demonstrated elevated TAK1
signaling in both human and rat retinae associated with neovascularization. We showed that reducing

TAK1 suppressed the expression of multiple cytokines and leukostasis in endothelial cells responding

to inflammatory insults *in vitro* through NFκB signaling pathway. Using 5Z-7-oxozeaenol, a well-

studied selective TAK1 inhibitor, we demonstrated that pharmacological inhibition of TAK1 can

effectively suppress retinal neovascularization in *in vivo* models in rats. These findings provide new

insights into the molecular and cellular mechanisms by which TAK1 modulates pathological

angiogenesis, thus highlighting a potential alternative approach to anti-VEGF treatment to prevent

retinal neovascularization.

MATERIALS AND METHODS

Data accession

To assess TAK1 expression in human retinal cells, we utilized data single cell transcriptomes from human retina, from the Gene Expression Omnibus (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE135133) (accessed on September 20, 2020).

To access gene expression in retinas from patients with retinal neovascularization due to proliferative diabetic retinopathy, we obtained the RNA-seq raw counts and TPM matrices from the Gene Expression Omnibus (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE102485) (accessed on September 27, 2020). We specifically compared samples from patients with retinal neovascular proliferative membranes due to type II diabetes (GSM2739349, GSM2739350 and GSM2739351) and those from normal controls (GSM2739364, GSM2739365 and GSM2739366).

scRNA-seq data analysis

scRNA-seq data were retrieved from a public dataset (GSE135133) and analyzed in R (v3.6.3) using Seurat (v3.2) (52, 53) with customized scripts. In brief, cell identity was assigned according to the assignment labels kindly provided by the authors after we normalized and scaled all human samples. We assigned ganglion cells, horizontal cells, bipolar cells, amacrine cells, cones and rods as the neural cells while immune cells, basal cells, red cells, pericytes, endothelial cells, astrocytes and Müller cells were regarded as non-neural cells. Dot plots were utilized for visualization of gene expression levels and percentage difference for various cell types.

RNA isolation

Total cells or tissue RNA was isolated using Zymo Quick-RNA MiniPrep kit (R1055; Zymo Research, Irvine, CA) according to the manufacturer's instructions. The quantity of RNA was evaluated with a NanoDrop ND-1000 spectrophotometer (NanoDrop Technologies, Wilmington, DE).

RNA-seq analysis

Analysis was undertaken in triplicate on total RNAs (1 μg) from normoxic and OIR rats at P14, prepared as per the manufacturers' instructions. cDNA libraries were sequenced by using an Illumina Hiseq-2500 RNA-seq platform as 50 bp single end chemistry at the Australian Genome Research Facility (AGRF, Melbourne, VIC, Australia). The raw fastq files of single-end reads were removed the adapter sequences and the low-quality reads were dropped by Trimgalore 0.4.4 (Babraham

Bioinformatics). Filtering parameter were set as follow: -q 25 --length 50 -e 0.1 --stringency 5. The

trimmed reads were subjected to alignment with default setting using STAR 2.5.3. The reference

genomes for human and rat were hg38 and rn6, respectively. Aligned RNA-seq data was counted over

gene exon using feature Counts (subread 1.6.4). Genes were annotated as per the Gencode Version 33

annotation file.

Gene set enrichment analysis (GSEA)

GSEA was run on the human dataset of protein-coding genes in PDR versus controls and on the rodent

dataset of protein-coding genes in OIR versus controls, against canonical pathway genesets collection

(c2.cp.v7.2, Broad Institute) by using 1000 genesets permutations. Only genesets with FDR < 0.05

were considered. Readcount values were loaded into R (v3.6.3) for statistical analysis and the heatmap

function was used to perform hierarchical clustering analysis and for data representation.

Quantitative polymerase chain reaction (qPCR)

Total RNAs (100 ng) was reverse transcribed to cDNA using a High-Capacity cDNA Reverse

Transcription kit (4368814; Life Technologies, Carlsbad, CA). Quantitative PCR was performed using

TaqMan fast advanced master mix (4444553; Applied Biosystems, Foster City, CA) and gene TaqMan

probe sets to detect gene expression (**Table S2**). The expression levels of target genes were normalized

to the levels of rat beta-actin (Rn00821065_g1) or human GAPDH (Hs99999905_m1). Subsequently,

the ΔΔCt method was used to evaluate relative expression level (fold change) in each condition versus

the corresponding control condition.

Cell culture

Telomerase-immortalized human microvascular endothelial cell (TIME; CRL-4025™) and primary

human retinal microvascular endothelial cell (HRMEC; ACBRI 181V) were purchased from American

Type Culture Collection (ATCC; Manassas, VA) and Cell Systems (Kirkland, WA). Endothelial cells

were cultured in EBM™ Plus Basal Medium supplied with EGM™ Plus SingleQuots™ supplements

(CC-5035; Lonza, Basel, Switzerland). All cell lines were maintained in a humidified incubator at

37°C and 5% CO_2. Cell lines were mycoplasma free tested by MycoAlert™ Mycoplasma Detection

Kit (LT07; Lonza).

Western blotting

A standard immunoblotting protocol was used. Briefly, protein was extracted from cells or retinas with

Pierce RIPA buffer (89900; Life Technologies) supplied with protease inhibitor cocktail

(11697498001; Roche Diagnostics, Basel, Switzerland). Equal amounts of cell lysates from different
groups were subjected to Western blotting analysis with specific antibodies against IκB, phospho-
NFκB p65, NFκB p65, phospho-p44/42 MAPK (Erk1/2), p44/42 MAPK (Erk1/2), phospho-TAK1,
TAK1, phospho-SAPK/JNK, SAPK/JNK, phospho-p38 MAPK, p38 MAPK (4814, 3033, 4764, 9101,
4695, 9339, 4505, 9255, 9252, 9211, 8690, respectively; Cell Signaling), and β-actin (MAB1501;
Merck Millipore, Burlington, MA). The quantitative densitometry was analyzed using Image Pro Plus
software (Media Cybernetics, Rockville, MA).
435

Animals
Pregnant Sprague-Dawley rats (female, 12-14 weeks old) and Brown Norway rats (male, 12 weeks
old) were supplied by the Animal Resources Centre (Murdoch, WA, Australia) and Cambridge Farm
Facility of the University of Tasmania (Cambridge, TAS, Australia). Animals were housed in standard
cages with a temperature/humidity-controlled and a 12-hour light (50 lux illumination)/12-hour dark
(<10 lux illumination) cycle environment. Food and water were available *ad libitum*. All animal
experiments were conducted in accordance with guidelines of the ARVO Statement for the Use of
Animals in Ophthalmic and Vision Research and were approved by the Animal Ethics Committees of
St Vincent's Hospital (004/16), the University of Melbourne (13-044UM), and the University of
Tasmania (A0017311). Rats were randomly allocated to treatment groups such that littermates were
distributed equally between groups.
447

Rat model of oxygen-induced retinopathy and intravitreal injection of 5Z-7-oxozeaenol
The oxygen-induced retinopathy model was induced in rats as previously described (54). Briefly,
Sprague-Dawley litters (within 12 hours of birth; P0) and their nursing mothers were exposed to daily
cycles of 80% O_2 for 21 hours and room air for 3 hours in a custom-built and humidity-controlled (<
80%) chamber until P14. At P14 animals were returned to room air until P16 or P18, when they were
sacrificed to harvest retinae. An ProOx 110 oxygen controller (BioSpherix; Parish, NY) was used to
regulate and monitor the oxygen level. At P14, intravitreal injections were performed to administer
5Z-7-oxozeaenol visualizing with a microsurgical microscope. Under surface anesthesia, a puncture
in the superior temporal quadrant of the limbus was made by a 30-gauge needle. A hand-pulled glass
micropipette (Bio-Strategy, Australia) connected to a 10 μl Hamilton syringe (Hamilton Company,
Reno, NV) was inserted into the vitreous cavity, and one microliter of balanced salt solution containing
1% DMSO (as vehicle), low (18 ng) or high dose (90 ng) of 5Z-7-oxozeaenol was intravitreally
injected into the eye. For safety evaluation, Brown Norway rats were intravitreally injected with 1 μl
high dose (90 ng) of 5Z-7-oxozeaenol or vehicle. Eyes were excluded from the study if large backflow

after injection or severe hemorrhage were occurred. Electroretinography and optical coherence
tomography imaging evaluation was undertaken 28 days post-intravitreal injection in anesthetized
animals as previously described (55).

Immunohistochemistry and vessel quantification
Collection of donor samples was approved by the Human Research Ethics Committee of the Royal
Victorian Eye and Ear Hospital (HREC13/1151H) and carried out in accordance with the approved
guidelines. Post-mortem eye globes were obtained via the Lions Eye Donation Service (Royal
Victorian Eye and Ear Hospital, East Melbourne, VIC, Australia) and processed for paraffin
embedding and sectioning following standard histological procedures. Briefly, donated eyes were
isolated, fixed, and permeabilized. The cross sections were stained with anti-TAK1 (1:100; NBP1-
87819, Novus Biologicals, Littleton, CO) and anti-CD31 (1:25; M0823, DAKO, Jena, Germany)
primary antibody and then with the appropriate secondary antibodies. Images were digitized using a
confocal laser scanning microscope (FV3000; Olympus, Tokyo, Japan).

The rat retinal flat-mount were collected at P18 followed by retina dissection, staining overnight
with isolectin GS-IB$_4$ Alexa Fluor™ 488 conjugate (5 μg/mL; I21411, Life Technologies), anti-TAK1
(1:100; NBP1-87819, Novus Biologicals), or anti-Iba1 (1:300; 019-19741, FUJIFILM Wako
Chemicals, Richmond, VA), and then incubated with the appropriate secondary antibodies. The images
were captured by FV3000 confocal laser scanning microscope. The avascular (vaso-obliteration) and
neovascularized areas were quantified by two blinded assessors using Adobe Photoshop (San Jose,
CA) (56). Deep vessel area was analyzed by AngioTool software (National Institutes of Health,
Bethesda, MA) (57).

Statistical analysis
All data are expressed as mean ± SEM. Statistical analyses were performed with Graphpad Prism 7
(GraphPad, San Diego, CA) using student t-test for unpaired data, one-way or two-way ANOVA
wherever appropriate followed by Tukey's or Bonferroni's multiple comparisons test. P values < 0.05
were considered statistically significant.

Data and resource availability statement
The main datasets generated or analyzed during this study are included here in the article. Retinal RNA
profiles of OIR and control rats have been deposited in Gene Expression Omnibus and are accessible
through GEO Series accession number GSE104588
(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc= GSE104588). The original Western Blot

images and the details of statistical analysis for this study are available in the supplementary

information. Other original datasets will also be available to the research community upon reasonable

request.

ACKNOWLEDGEMENTS

The authors thank UTAS CFF animal technicians, Karen Shiels, Keri Smith, Heather Howard, Lisa Harding and Danielle Eastley for their assistance with rat chamber operation. We also thank Dr Jacqueline Y.K. Leung for assistance with intravitreal injection, and Zheng He for assistance with retinal imaging. This work was supported by grants from the National Health and Medical Research Council of Australia (1061912, 1185600 and 1123329), the Ophthalmic Research Institute of Australia, the National Natural Science Foundation of China (82000902) and the Shenzhen Key Laboratory of Biomimetic Materials and Cellular Immunomodulation (ZDSYS20190902093409851). GJD was a Principal Research Fellow of NHMRC. The Centre for Eye Research Australia receives Operational Infrastructure Support from the Victorian Government.

COMPETING INTERESTS

The authors have declared that no competing interest exists.

AUTHOR CONTRIBUTIONS

Conceptualization- F-L.L., J-H.W., P.vW., G.J.D., G-S.L. Methodology- F-L.L., C.M., J-H.W., B.V.B., P.vW., G-S.L. Formal Analysis- F-L.L., J-H.W., J.C., R.C.B.W., B.V.B., G-S.L. Investigation- F-L.L., J-H.W., J.C., L.Z., Y-F.C., L.T., C.M., S.L., D.L., C-L.T., B.V.B. Resources- R.C.B.W., A.W.H., C-L.T., G.J.D., P-Y.W., G-S.L. Data Curation- F-L.L., J-H.W., J.C., G-S.L. Writing (Original Draft)- F-L.L., J-H.W., G-S.L. Writing (Review & Editing)- J.C., C.M., R.C.B.W., B.V.B., P.vW., G.J.D., P-Y.W. Project Administration- F-L.L., J-H.W., G-S.L. Funding Acquisition- L.T., G.J.D., P-Y.W., G-S.L. Project Supervision- G.S.L.

REFERENCES

1.   Neely KA, and Gardner TW. Ocular neovascularization: clarifying complex interactions. *Am J Pathol.* 1998;153(3):665-70.

2.   Campochiaro PA. Retinal and choroidal neovascularization. *Journal of cellular physiology.* 2000;184(3):301-10.

3.   Singer MA, Kermany DS, Waters J, Jansen ME, and Tyler L. Diabetic macular edema: it is more than just VEGF. *F1000Res.* 2016;5.

4.   Qazi Y, Maddula S, and Ambati BK. Mediators of ocular angiogenesis. *J Genet.* 2009;88(4):495-515.

5.   Goldberg RB. Cytokine and cytokine-like inflammation markers, endothelial dysfunction, and imbalanced coagulation in development of diabetes and its complications. *J Clin Endocrinol Metab.* 2009;94(9):3171-82.

6.   Demircan N, Safran BG, Soylu M, Ozcan AA, and Sizmaz S. Determination of vitreous interleukin-1 (IL-1) and tumour necrosis factor (TNF) levels in proliferative diabetic retinopathy. *Eye (Lond).* 2006;20(12):1366-9.

7.   Capitao M, and Soares R. Angiogenesis and Inflammation Crosstalk in Diabetic Retinopathy. *J Cell Biochem.* 2016;117(11):2443-53.

8.   Murugeswari P, Shukla D, Rajendran A, Kim R, Namperumalsamy P, and Muthukkaruppan V. Proinflammatory cytokines and angiogenic and anti-angiogenic factors in vitreous of patients with proliferative diabetic retinopathy and eales' disease. *Retina.* 2008;28(6):817-24.

9.   Stevenson W, Cheng SF, Dastjerdi MH, Ferrari G, and Dana R. Corneal neovascularization and the utility of topical VEGF inhibition: ranibizumab (Lucentis) vs bevacizumab (Avastin). *Ocul Surf.* 2012;10(2):67-83.

10.   Aveleira CA, Lin CM, Abcouwer SF, Ambrosio AF, and Antonetti DA. TNF-alpha signals through PKCzeta/NF-kappaB to alter the tight junction complex and increase retinal endothelial cell permeability. *Diabetes.* 2010;59(11):2872-82.

11.   Clauss M, Sunderkotter C, Sveinbjornsson B, Hippenstiel S, Willuweit A, Marino M, et al. A permissive role for tumor necrosis factor in vascular endothelial growth factor-induced vascular permeability. *Blood.* 2001;97(5):1321-9.

12.   Yoshida S, Yoshida A, and Ishibashi T. Induction of IL-8, MCP-1, and bFGF by TNF-alpha in retinal glial cells: implications for retinal neovascularization during post-ischemic inflammation. *Graefes Arch Clin Exp Ophthalmol.* 2004;242(5):409-13.

13.   Joussen AM, Poulaki V, Mitsiades N, Kirchhof B, Koizumi K, Dohmen S, et al. Nonsteroidal anti-inflammatory drugs prevent early diabetic retinopathy via TNF-alpha suppression. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology.* 2002;16(3):438-40.

14.   Sato S, Sanjo H, Takeda K, Ninomiya-Tsuji J, Yamamoto M, Kawai T, et al. Essential function for the kinase TAK1 in innate and adaptive immune responses. *Nat Immunol.* 2005;6(11):1087-

95.

15.   Shim JH, Xiao C, Paschal AE, Bailey ST, Rao P, Hayden MS, et al. TAK1, but not TAB1 or

TAB2, plays an essential role in multiple signaling pathways in vivo. *Genes Dev.*

2005;19(22):2668-81.

16.   Mihaly SR, Ninomiya-Tsuji J, and Morioka S. TAK1 control of cell death. *Cell death and*

differentiation. 2014;21(11):1667-76.

17.   Scarneo SA, Mansourati A, Eibschutz LS, Totzke J, Roques JR, Loiselle D, et al. Genetic and

pharmacological validation of TAK1 inhibition in macrophages as a therapeutic strategy to

effectively inhibit TNF secretion. *Sci Rep.* 2018;8(1):17058.

18.   Harikumar KB, Sung B, Tharakan ST, Pandey MK, Joy B, Guha S, et al. Sesamin manifests

chemopreventive effects through the suppression of NF-kappa B-regulated cell survival,

proliferation, invasion, and angiogenic gene products. *Mol Cancer Res.* 2010;8(5):751-61.

19.   Naito H, Iba T, Wakabayashi T, Tai-Nagara I, Suehiro JI, Jia W, et al. TAK1 Prevents

Endothelial Apoptosis and Maintains Vascular Integrity. *Dev Cell.* 2019;48(2):151-66 e7.

20.   Morioka S, Inagaki M, Komatsu Y, Mishina Y, Matsumoto K, and Ninomiya-Tsuji J. TAK1

kinase signaling regulates embryonic angiogenesis by modulating endothelial cell survival and

migration. *Blood.* 2012;120(18):3846-57.

21.   Schoneberg J, De Lorenzi F, Theek B, Blaeser A, Rommel D, Kuehne AJC, et al. Engineering

biofunctional in vitro vessel models using a multilayer bioprinting technique. *Sci Rep.*

2018;8(1):10430.

22.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set

enrichment analysis: a knowledge-based approach for interpreting genome-wide expression

profiles. *Proc Natl Acad Sci U S A.* 2005;102(43):15545-50.

23.   Ninomiya-Tsuji J, Kajino T, Ono K, Ohtomo T, Matsumoto M, Shiina M, et al. A resorcylic

acid lactone, 5Z-7-oxozeaenol, prevents inflammation by inhibiting the catalytic activity of

TAK1 MAPK kinase kinase. *J Biol Chem.* 2003;278(20):18485-90.

24.   Hackett SF, Wiegand S, Yancopoulos G, and Campochiaro PA. Angiopoietin-2 plays an

important role in retinal angiogenesis. *J Cell Physiol.* 2002;192(2):182-7.

25.   Fechtner S, Fox DA, and Ahmed S. Transforming growth factor beta activated kinase 1: a

potential therapeutic target for rheumatic diseases. *Rheumatology (Oxford).* 2017;56(7):1060-

8.

26.   Mukhopadhyay H, and Lee NY. Multifaceted roles of TAK1 signaling in cancer. *Oncogene.*

2020;39(7):1402-13.

27.   Yang Y, Wu, J, and Wang, J.K. A database and functional annotation of NF-κB target genes. *Int*

J Clin Exp Med. 2016;9(5):7986-95.

28.   Ghosh S, and Karin M. Missing pieces in the NF-kappaB puzzle. *Cell.* 2002;109 Suppl:S81-

96.

29.   Li J, Liang C, Zhang ZK, Pan X, Peng S, Lee WS, et al. TAK1 inhibition attenuates both

inflammation and fibrosis in experimental pneumoconiosis. *Cell Discov.* 2017;3:17023.

30.   Yang L, Joseph S, Sun T, Hoffmann J, Thevissen S, Offermanns S, et al. TAK1 regulates
endothelial cell necroptosis and tumor metastasis. *Cell death and differentiation.*
2019;26(10):1987-97.

31.   Dai L, Aye Thu C, Liu XY, Xi J, and Cheung PC. TAK1, more than just innate immunity.
IUBMB Life. 2012;64(10):825-34.

32.   Onodera Y, Teramura T, Takehara T, Shigi K, and Fukuda K. Reactive oxygen species induce
Cox-2 expression via TAK1 activation in synovial fibroblast cells. *FEBS Open Bio.*
2015;5:492-501.

33.   Xie M, Zhang D, Dyck JR, Li Y, Zhang H, Morishima M, et al. A pivotal role for endogenous
TGF-beta-activated kinase-1 in the LKB1/AMP-activated protein kinase energy-sensor
pathway. *Proc Natl Acad Sci U S A.* 2006;103(46):17378-83.

34.   Sakurai H. Targeting of TAK1 in inflammatory disorders and cancer. *Trends in*
pharmacological sciences. 2012;33(10):522-30.

35.   Sabio G, and Davis RJ. TNF and MAP kinase signalling pathways. *Semin Immunol.*
2014;26(3):237-45.

36.   Landry J, and Huot J. Regulation of actin dynamics by stress-activated protein kinase 2
(SAPK2)-dependent phosphorylation of heat-shock protein of 27 kDa (Hsp27). *Biochem Soc*
Symp. 1999;64:79-89.

37.   Munoz-Chapuli R, Quesada AR, and Angel Medina M. Angiogenesis and signal transduction
in endothelial cells. *Cell Mol Life Sci.* 2004;61(17):2224-43.

38.   Tang M, Wei X, Guo Y, Breslin P, Zhang S, Zhang S, et al. TAK1 is required for the survival
of hematopoietic cells and hepatocytes in mice. *The Journal of experimental medicine.*
2008;205(7):1611-9.

39.   Wu J, Powell F, Larsen NA, Lai Z, Byth KF, Read J, et al. Mechanism and in vitro
pharmacology of TAK1 inhibition by (5Z)-7-Oxozeaenol. *ACS chemical biology.*
2013;8(3):643-50.

40.   Zhang D, Yan H, Li H, Hao S, Zhuang Z, Liu M, et al. TGFbeta-activated Kinase 1 (TAK1)
Inhibition by 5Z-7-Oxozeaenol Attenuates Early Brain Injury after Experimental Subarachnoid
Hemorrhage. *The Journal of biological chemistry.* 2015;290(32):19900-9.

41.   Lu L, Zhang X, Tong H, Zhang W, Xu P, and Qu S. Central Administration of 5Z-7-Oxozeaenol
Protects Experimental Autoimmune Encephalomyelitis Mice by Inhibiting Microglia
Activation. *Front Pharmacol.* 2017;8:789.

42.   Merson TD, Binder MD, and Kilpatrick TJ. Role of cytokines as mediators and regulators of
microglial activity in inflammatory demyelination of the CNS. *Neuromolecular Med.*
2010;12(2):99-132.

43.   White BJ, Tarabishy S, Venna VR, Manwani B, Benashski S, McCullough LD, et al. Protection
from cerebral ischemia by inhibition of TGFbeta-activated kinase. *Exp Neurol.*

2012;237(1):238-45.

44.   Ivshina M, Alexandrov IM, Vertii A, Doxsey S, and Richter JD. CPEB regulation of TAK1
638   synthesis mediates cytokine production and the inflammatory immune response. *Mol Cell Biol.*
639   2015;35(3):610-8.

640   45.   Ireland DJ, Kemp MW, Miura Y, Saito M, Newnham JP, and Keelan JA. Intra-amniotic
641   pharmacological blockade of inflammatory signalling pathways in an ovine chorioamnionitis
642   model. *Mol Hum Reprod.* 2015;21(5):479-89.

643   46.   Neubert M, Ridder DA, Bargiotas P, Akira S, and Schwaninger M. Acute inhibition of TAK1
644   protects against neuronal death in cerebral ischemia. *Cell death and differentiation.*
645   2011;18(9):1521-30.

646   47.   Bettermann K, Vucur M, Haybaeck J, Koppe C, Janssen J, Heymann F, et al. TAK1 suppresses
647   a NEMO-dependent but NF-kappaB-independent pathway to liver cancer. *Cancer Cell.*
648   2010;17(5):481-96.

649   48.   Omori E, Morioka S, Matsumoto K, and Ninomiya-Tsuji J. TAK1 regulates reactive oxygen
650   species and cell death in keratinocytes, which is essential for skin integrity. *J Biol Chem.*
651   2008;283(38):26161-8.

652   49.   Bui BV, and Fortune B. Ganglion cell contributions to the rat full-field electroretinogram. *J
653   Physiol.* 2004;555(Pt 1):153-73.

654   50.   Cao H, Lu J, Du J, Xia F, Wei S, Liu X, et al. TAK1 inhibition prevents the development of
655   autoimmune diabetes in NOD mice. *Sci Rep.* 2015;5:14593.

656   51.   Tan L, Gurbani D, Weisberg EL, Jones DS, Rao S, Singer WD, et al. Studies of TAK1-centered
657   polypharmacology with novel covalent TAK1 inhibitors. *Bioorg Med Chem.* 2017;25(4):1320-
658   8.

659   52.   Butler A, Hoffman P, Smibert P, Papalexi E, and Satija R. Integrating single-cell transcriptomic
660   data across different conditions, technologies, and species. *Nat Biotechnol.* 2018;36(5):411-20.

661   53.   Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, 3rd, et al.
662   Comprehensive Integration of Single-Cell Data. *Cell.* 2019;177(7):1888-902 e21.

663   54.   Penn JS, Tolman BL, and Henry MM. Oxygen-induced retinopathy in the rat: relationship of
664   retinal nonperfusion to subsequent neovascularization. *Invest Ophthalmol Vis Sci.*
665   1994;35(9):3429-35.

666   55.   Bee YS, Tu L, Sheu SJ, Lin HC, Tang JH, Wang JH, et al. Gene Delivery of Calreticulin Anti-
667   Angiogenic Domain Attenuates the Development of Choroidal Neovascularization in Rats.
668   *Hum Gene Ther.* 2017;28(5):403-14.

669   56.   Connor KM, Krah NM, Dennison RJ, Aderman CM, Chen J, Guerin KI, et al. Quantification
670   of oxygen-induced retinopathy in the mouse: a model of vessel loss, vessel regrowth and
671   pathological angiogenesis. *Nat Protoc.* 2009;4(11):1565-73.

672   57.   Zudaire E, Gambardella L, Kurcz C, and Vermeren S. A computational tool for quantitative
673   analysis of vascular networks. *PLoS One.* 2011;6(11):e27385.

674   58.   Zhao D, Nguyen CT, Wong VH, Lim JK, He Z, Jobling AI, et al. Characterization of the
675   Circumlimbal Suture Model of Chronic IOP Elevation in Mice and Assessment of Changes in
676   Gene Expression of Stretch Sensitive Channels. *Front Neurosci.* 2017;11:41.
677

680　**Fig. 1 Characterization of TAK1 expression in patients with retinal neovascularization**
681　**associated with proliferative diabetic retinopathy.**

(A) Single nuclei RNAseq (snRNAseq) characterization of TAK1 (MAP3K7) expression in normal human retina. TAK1 and TAK1-associated proteins (TAB1-3) were expressed in most neural (red) and non-neural (blue) cells in the retina, including cell associated with the retinal vasculature. (B) Characterization of TAK1 expression in the normal human retina. Endothelial cells were labelled with CD31 (red) and counterstained with TAK1 (green) in retinal section. Apart from neural cells, TAK1 expression was also found in vascular endothelial cells, especially in retinal capillaries (3 and 4). Scale bar: 100 μm. (C) Gene-set enrichment analysis (GSEA) indicated that TAK1 upstream pathways including IL1, TNF, TCR, BCR, TLR, TNFR, or TGFβ signaling (Reactome Pathway Database) were positively enriched in retinal samples from patients with retinal neovascularization due to proliferative diabetic retinopathy. Engagement of TAK1-activated NFκB, p38 MAPK and JNK pathways as well as negative impact on retinal neural functions were also identified. (D) GSEA Enrichment plots illustration of gene set involved in TAK1-NFκB, TAK1-p38 MAPK and TAK1-JNK pathways in the retina of patients with proliferative diabetic retinopathy. GCL: ganglion cell layer; INL: inner nuclear layer; ONL: outer nuclear layer; RPE: retinal pigment epithelium; NSE: normalized enrichment score; FDR: false discovery rate.

Fig. 2 Involvement of TAK1 in pathological retinal angiogenesis in a rat model of oxygen-induced retinopathy. (A) Schematic diagram of the Sprague-Dawley rat model of oxygen-induced retinopathy (OIR). Neonatal rats were exposed to 80% of oxygen for 21 hours each day from postnatal day 0 (P0) to P14. Rats were returned to room air from P14 to P18. Vessel loss was induced during oxygen exposure and retinal angiogenesis could be induced after rats returned to room air. (B) qPCR analysis of *TAK1* expression in OIR at P18 as compared with the normoxic control (Nor) group (n = 6-7). (C) Western blot analysis of TAK1 expression in OIR at P18 compared with normoxic control group (n = 10-12). Representative immunoblotting from 4 independent samples were shown for normoxic control (lanes 1-4) and OIR group (lanes 5-8). (D) Immunobiological staining indicated expression of TAK1 in the pathological vessel. Vasculature were visualized by isolectin GS-IB4 labelling (green) and counterstained with TAK1 (red) in retinal whole mount. Arrowheads indicate vessel tufts. Scale bar: 100 μm. (E) GSEA indicated that most of TAK1 upstream pathways including TNF, IL1, TCR, BCR, TLR, and VEGF pathways (from Reactome Pathway Database) were positively enriched in OIR at P18. Engagement of TAK1-activated NFκB, p38 MAPK and JNK pathways as well as negative impact on retinal neural functions were also identified. Group data are shown as means ± SEM. Statistical analysis was undertaken with two-tailed unpaired t-test; ***$P < 0.001$, ****$P < 0.0001$. NSE: normalized enrichment score; FDR: false discovery rate.

Fig. 3 TAK1 Inhibition by 5Z-7-oxozeaenol suppressed retinal angiogenesis in OIR rats.

(A) Experimental protocol for 5Z-7-oxozeaenol treatment in the OIR rat model. A single intravitreal injection of vehicle, low (18 ng) or high (90 ng) dose of 5Z-7-oxozeaenol was given to Sprague-

Dawley OIR rats at P14, followed by quantitative analysis of retinal neovascularization of flat mounted
retinae at P18. (B) Retinal neovascularization is highlighted in white, and insets show selected areas
at high magnification. Arrowheads indicate vessel tufts. Retinal neovascularization was quantified
from 10 individual retinas per group. Scale bar: 500 µm. (C) Vaso-obliteration was simultaneously
evaluated with the same samples analyzed in Fig. 3B. The white areas represent the avascular area in
the retina. No significant difference was observed between groups (n = 10). Scale bar: 1 mm. (D)
Vessel area of superficial and deep vascular layers was quantified in P18 control (n = 4) and OIR rats
following a single intravitreal injection of vehicle, low or high dose of 5Z-7-oxozeaenol at P14 (n = 5
each group). Scale bar: 100 µm. Group data are shown as means ± SEM. Statistical analysis was
undertaken with one-way ANOVA and Tukey's multiple comparison test; ****$P < 0.0001$.
731

Fig. 4 TAK1 deletion inhibits the expression of cytokines and cell adhesion molecules induced by TNFα in human microvascular endothelial cells.

(A) Principal components analysis of RNA-seq data of wild-type (WT) endothelial cells (TIMEs), WT TIMEs treated with TNFα, TAK1 knockout (TAK1-KO) TIMEs, and TAK1-KO TIMEs treated with TNFα. Each dot represents an experimental replicate. (A) Volcano plots show genes that were significantly changed in both WT and TAK1-KO TIMEs when stimulated by TNFα. Red dots refer to genes showing a log2|fold change| > 1 and FDR < 0.05. Labels indicate significantly up-regulated genes in each data set. (C) GSEA of the top 20 KEGG gene sets in the two groups. Significantly enriched gene sets in WT TNF set (WT TNF vs. WT) (positive NES) and less enriched ones in TAK1-KO set (TAK1-KO TNF vs. WT TNF) (negative NES) are highlighted in red. False discovery rate (FDR) q-value < 0.05 was considered to be statistically significant. (D) GSEA plots for gene sets with Gene Ontology (GO) term "CYTOKINE AND CYTOKINE RECEPTOR INTERACTION" (related to cytokines) in the two groups. The Venn diagram shows the number of genes from the cytokine gene sets in the respective and overlapping areas of the two groups. (E) GSEA plots for gene sets with GO term "CELL ADHESION MOLECULES-CAMS" (related to adhesion molecules) in the two groups. The Venn diagram shows the number of genes from the chemokine gene sets in respective and overlapping areas of the two groups. (F) Heat map generated from the identified overlapping areas of "cytokines (n = 59)" "adhesion molecules (n = 23)" gene sets based on RNA-Seq read counts.

Fig. 5 TAK1 deletion prevents TNFα-induced inflammatory cascade through the NFκB and MAPK pathway in human microvascular endothelial cells.

(A) Schematic diagram of TNFα-TAK1 signaling. Upon TNFα stimulation, TAK1 binds to the polyubiquitin chain though TAB2, and activates the IKK complex, leading to the activation of NF-κB.

TAK1 also activates MAPKs cascades. NFκB and MAPKs/AP-1 induce expression of inflammatory
cytokines and cell adhesion proteins. (B) The Venn diagram respectively shows the number of genes
from the "cytokine" and "adhesion molecule" gene sets, and the overlap indicates the number of genes
directly targeted by NFκB or AP-1. (C) Western blot characterization of the TNFα-induced
phosphorylation of TAK1, NFκB, and MAPKs proteins in TAK1 knockout (TAK1-KO) TIMEs. Wild-
type (WT) and TAK1-KO TIMEs were stimulated by TNFα (10 ng/ml) for 10 minutes. Activation of
TAK1, NFκB p65, IκB, JNK, p38, and ERK assessed using western blot (n = 3-4). Genetic TAK1
deletion reduced TAK1 expression (D) and impaired NFκB (NFκB p65 and IκB) (E) and MAPK (p38,
JNK, and ERK) (F) signaling in TIMEs treated with TNFα. (G) qPCR results revealed that TAK1
deletion suppressed the expression of selected cytokines (*CXCL8* and *VEGFA)* and cell adhesion
molecules (*ICAM-1* and *VCAM-1)* in TIMEs stimulated with TNFα for 24 hours (n = 3-4). Group data
are shown as means ± SEM. Statistical analysis was undertaken with one-way ANOVA and Tukey's
multiple comparison test; *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$.
769

Fig. 6 TAK1 inhibition by 5Z-7-oxozeaenol decreased inflammatory gene expression and adhesion of microglia to vessels in OIR.

(A) Experimental protocol for 5Z-7-oxozeaenol treatment in the OIR rat model. A single intravitreal injection of vehicle, low (18 ng) or high (90 ng) dose of 5Z-7-oxozeaenol was given to Sprague-Dawley OIR rats at P14, and retinae were collected for qPCR and immunostaining at P16 and P18. (B) Intravitreal injection of 5Z-7-oxozeaenol in OIR rats at P14 inhibited *TNFα*, *VEGFA* and *ICAM-1* gene expression assessed using qPCR (n = 5-7). (C) Representative images of microglial adhesion to the vascular surface in a retina from a control rat and OIR rats that had a single intravitreal injection of vehicle, low or high dose of 5Z-7-oxozeaenol (Oxo). Vessels were visualized with isolectin GS-IB4

staining (green), and distribution of microglia was immunolabeled with Iba-1 (red). Scale bar: 50 µm.
(D) Quantitative analysis of microglial number relative to retinal vessel area in vehicle or 5Z-7-
oxozeaenol treated OIR rats (n = 7-10; 10 images from each retina were shown as separated dot plot).
(E) Schematic diagram for the role of TAK1 signaling in microvascular endothelial cells.
Proinflammatory cytokines such as TNFα can activate TAK1 in microvascular endothelial cells, and
TAK1/TAB1/TAB2-3 complex can lead to activation of NFκB and MAPKs/AP-1 cascades. NFκB and
MAPKs/AP-1 induce expression of "cytokines" and "adhesion proteins", thereby triggering the
"inflammatory cycle" and "leukostasis", and leading to retinal neovascularization. Group data are
shown as means ± SEM. Statistical analysis was undertaken with one-way ANOVA and Tukey's
multiple comparison test; *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$.
790

 SUPPLEMENTARY MATERIALS

792 **Table S1** Effects on the expression of enriched genes mediated by NFκB or AP-1 signaling in "TAK1-

793 KO TNF *vs.* WT TNF" data set compared to "WT TNF *vs.* WT" data set.

Gene name	WT TNF vs WT		TAK1-KO TNF vs TAK1-KO TNF		NFkB or AP-1 targeted gene
	log2FoldChange	FDR q-value	log2FoldChange	FDR q-value	
Cytokines-Cytokine Receptors (n=58)					
IL1B	4.35670	3.94E-11	-4.41950	2.40E-05	NFκB
CXCL1	1.83710	7.24E-11	-2.34930	3.41E-04	NFκB
IL1A	3.29810	5.17E-10	-1.87860	6.40E-03	NFκB
VEGFC	1.72730	1.06E-09	-0.81200	1.29E-02	NFκB
TNFRSF9	6.36650	1.62E-09	-5.03510	2.97E-05	None
TNFRSF4	4.86660	1.03E-08	-2.56130	9.43E-03	None
CCL2	4.79430	2.35E-08	-3.10600	2.47E-02	NFκB
CCL5	6.50100	1.47E-07	-6.05740	5.44E-07	NFκB
CXCL2	2.68970	8.93E-07	-2.39200	1.56E-02	NFκB
CD70	1.74470	1.94E-06	-8.39220	4.56E-13	NFκB
PDGFB	1.05430	2.71E-06	-0.63160	4.20E-04	NFκB
LIF	2.06620	8.24E-06	-2.46700	2.83E-10	NFκB
CXCL5	4.02040	9.25E-06	-1.43094	3.26E-01	None
CXCL6	3.20050	1.13E-05	-3.24340	1.99E-05	NFκB
CXCL8	3.40100	2.62E-05	-4.62360	1.53E-07	NFκB
CXCL3	2.83370	3.02E-05	-1.95808	2.94E-01	NA
CCR10	1.38350	5.53E-05	-1.94630	6.67E-09	AP-1
CXCL10	9.88480	6.15E-05	-5.92910	5.15E-03	NFκB
CSF1	1.54050	9.12E-05	-1.47410	2.91E-09	NFκB
CXCL11	8.86230	1.07E-04	-5.92440	9.63E-03	NFκB
IL12RB1	2.38450	1.60E-04	-2.56180	6.75E-08	NFκB
IFNGR2	1.06720	2.21E-04	-1.20550	3.02E-09	NFκB
VEGFA	0.97810	2.91E-04	-0.96540	5.89E-07	NFκB
CX3CL1	4.93410	3.00E-04	-5.75310	1.09E-05	NFκB
CCL20	5.98070	3.08E-04	-6.66700	2.81E-04	NFκB
TNFRSF11B	2.27490	5.60E-04	-1.22159	2.09E-01	None
LTB	5.02310	8.03E-04	-5.28530	4.51E-07	NFκB
IL15RA	0.95000	1.28E-03	-1.26430	8.62E-09	NFκB
TNFSF9	1.00000	1.36E-03	-2.03650	8.17E-15	NFκB
TNFSF15	1.19150	2.94E-03	-2.26180	4.67E-18	None

BMP2	1.54300	3.27E-03	-1.69090	8.02E-03	NFκB
TGFBR1	0.70040	2.20E-02	-0.82610	2.84E-05	NFκB
IL12A	1.68170	3.07E-02	-1.53970	1.53E-02	NFκB
PRLR	2.91580	3.15E-02	-3.27440	4.52E-03	AP-1
IL15	0.85040	3.16E-02	-1.26610	1.07E-06	NFκB
IL23A	1.82120	5.59E-02	-1.58160	1.52E-02	NFκB/AP-1
IL2RG	0.95541	1.01E-01	-1.82390	5.10E-01	None
TNFRSF12A	0.71783	1.18E-01	-0.43630	6.51E-03	AP-1
TNFRSF14	0.46530	1.53E-01	-0.48970	2.73E-03	None
TNFRSF10B	0.52018	1.97E-01	-0.35370	4.89E-02	NFκB
BMPR1B	1.02127	1.98E-01	-2.11260	2.99E-04	None
HGF	2.08440	2.92E-01	-2.90270	1.53E-02	NFκB
PDGFC	0.42220	2.94E-01	-0.84640	1.00E-05	NFκB
IL22RA1	3.47750	3.63E-01	-5.93710	5.43E-03	NFκB
CCR4	2.87390	4.03E-01	-3.34307	3.91E-01	NA
CSF2	1.64173	4.61E-01	-0.17474	5.04E-01	NFκB
OSMR	0.40440	5.10E-01	-0.24603	3.82E-01	None
TNFSF10	1.37807	5.88E-01	-0.74565	3.64E-01	None
PDGFRB	0.42136	7.46E-01	-1.75040	7.60E-05	AP-1
TNFSF13B	3.28969	NA	-3.31250	3.42E-02	NFκB
TNFRSF18	2.87284	NA	-2.91772	1.48E-01	AP-1
IL7R	3.67340	NA	-3.50272	NA	NFκB/AP-1
CX3CR1	2.38491	NA	-2.44076	NA	NFκB
TNF	5.49810	NA	-5.56714	NA	NFκB
CXCL9	3.32350	NA	-3.39533	NA	NFκB
IFNAR2	0.34410	NA	-0.39382	9.84E-02	AP-1
INHBA	0.83100	NA	-3.20137	2.55E-01	None
CCL8	2.91000	NA	-2.98153	NA	None
CSF2RB	1.54048	NA	-2.04434	6.12E-02	None
Cell Adhesion Molecules (n=23)					
HLA-B	1.54750	1.24E-10	-2.12490	4.67E-29	NFκB
NRCAM	2.08430	1.99E-10	-3.42680	3.44E-08	None
CLDN14	2.72790	8.91E-10	-1.73390	1.61E-02	None
HLA-F	2.57600	2.25E-07	-4.58300	1.43E-14	NFκB
ICAM1	3.58140	4.18E-07	-3.17350	3.58E-04	NFκB
VCAM1	10.39070	6.70E-07	-7.44920	3.86E-04	NFκB
CNTNAP1	1.13470	8.81E-06	-0.87440	3.03E-07	None

ICOSLG	3.32720	1.44E-05	-3.15500	4.04E-03	None
CLDN1	2.09970	2.08E-05	-4.77710	1.22E-12	None
ITGAV	1.70150	6.11E-05	-1.20700	8.66E-03	None
HLA-C	0.85300	2.49E-04	-1.31420	9.92E-15	None
HLA-A	0.83990	3.40E-04	-0.81560	2.38E-06	NFκB
SDC4	0.91350	5.62E-04	-0.79390	6.91E-05	None
PDCD1LG2	0.98570	2.49E-03	-0.96010	1.02E-04	NFκB
SDC1	0.90460	1.09E-02	-1.18610	1.06E-06	None
SDC3	0.41120	4.08E-02	-1.86010	7.91E-51	None
CLDN23	1.91240	1.40E-01	-4.50220	1.21E-05	None
SELE	4.44980	1.87E-01	-9.34310	4.74E-05	NFκB
ITGB8	0.61910	2.80E-01	-2.01250	1.03E-10	NFκB
HLA-G	0.77070	3.23E-01	-1.45580	2.09E-03	None
MADCAM1	0.83220	7.39E-01	-0.84027	NA	None
HLA-DQA1	2.94940	NA	-3.01068	NA	None
CD6	1.94430	NA	-2.00757	NA	None

796 FDR: False Discovery Rate, NA: Not Available, TAK1-KO: TAK1 knockout, TNF: TNFα, WT: Wild-

797 type.

 Table S2. Key Resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
IκBα (L35A5) mouse mAb	Cell Signaling Technology	4814
Phospho-NFκB p65 (Ser536) (93H1) rabbit mAb	Cell Signaling Technology	3033
NFκB p65 (C22B4) rabbit mAb	Cell Signaling Technology	4764
Phospho-p44/42 MAPK (Erk1/2) (Thr202/Tyr204) rabbit pAb	Cell Signaling Technology	9101
p44/42 MAPK (Erk1/2) (137F5) rabbit mAb	Cell Signaling Technology	4695
Phospho-TAK1 (Ser412) rabbit pAb	Cell Signaling Technology	9339
TAK1 rabbit pAb	Cell Signaling Technology	4505
Phospho-SAPK/JNK (Thr183/Tyr185) (G9) mouse mAb	Cell Signaling Technology	9255
SAPK/JNK rabbit pAb	Cell Signaling Technology	9252
Phospho-p38 MAPK (Thr180/Tyr182) rabbit pAb	Cell Signaling Technology	9211
p38 MAPK (D13E1) XP® rabbit mAb	Cell Signaling Technology	8690
Actin (clone C4) mouse mAb	Merck Millipore	MAB1501
TAK1 rabbit pAb	Novus Biologicals	NBP1-87819
Human CD31 mouse mAb	DAKO	M0823
Iba1 rabbit pAb	FUJIFILM Wako Chemicals	019-19741
Isolectin GS-IB4 from Griffonia simplicifolia, Alexa Fluor™ 488 conjugate	Life Technologies Australia	I21411
Goat anti-mouse IgG HRP-conjugated secondary Ab	Merck Millipore	12-349
Goat anti-rabbit Alexa Fluor 488 plus secondary Ab	Invitrogen	A32731

Goat anti-mouse Alexa Fluor 594 plus secondary Ab	Invitrogen	A32742
NucBlue™ Reagent (Hoechst 33342)	Life Technologies Australia	R37605
Chemicals		
5Z-7-oxozeaenol	Tocris Bioscience	3604/1
3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-2H-tetrazolium bromide (MTT)	Sigma-Aldrich	M2128
DMSO	Sigma-Aldrich	D8418
Sodium citrate dihydrate	Sigma-Aldrich	W302600
Triton X100	Sigma-Aldrich	T8787
Ampicillin	Sigma-Aldrich	A0166
Polybrene	Sigma-Aldrich	H9268
Puromycin	Sigma-Aldrich	P8833
Agarose	Sigma-Aldrich	A6361
Critical Commercial Assays		
Pierce™ BCA assay kit	Life Technologies Australia	23227
Amersham ECL Prime Western Blotting Detection Kit	GE Healthcare	RPN2232
Q5 Hot Start High-Fidelity 2X Master Mix	New England Biolabs	M0494
High-capacity cDNA reverse transcription Kit	Life Technologies Australia	4368814
TaqMan™ Fast Advanced Master Mix	Life Technologies Australia	4444553
Quick-RNA MiniPrep kit	Zymo Research	R1055
Agencourt CleanSeq Kit	Beckman Coulter	A29151
ZymoPURE Plasmid Miniprep Kit	Zymo Research	D4210
Quick-DNA Microprep Kit	Zymo Research	D3021
Experimental Models: Cell Lines		
Telomerase-immortalized human microvascular endothelial cells (TIMEs)	American Type Culture Collection	ATCC® CRL-4025™
Human retinal microvascular endothelial cells (HRMECs)	Cell Systems	ACBRI 181 V
Human embryonic kidney 293FT	Life Technologies Australia	R70007
Experimental Models: Organisms/Strains		

Sprague-Dawley rat and C57BL/6 mouse	St Vincent's Hospital	004/16
Brown Norway rat	University of Melbourne	13-044UM
Sprague-Dawley rat	University of Tasmania	A0017311
Oligonucleotides		
TAK1 sgRNA#1 Forward	Integrated DNA Technologies	CACCGTAGACCAACAACGAGTCATC
TAK1 sgRNA#1 Reverse	Integrated DNA Technologies	AAACGATGACTCGTTGTTGGTCTAC
TAK1 sgRNA#2 Forward	Integrated DNA Technologies	CACCGGAGTTGTTTGCAAAGCTAAG
TAK1 sgRNA#2 Reverse	Integrated DNA Technologies	AAACCTTAGCTTTGCAAACAACTCC
TAK1 sgRNA#3 Forward	Integrated DNA Technologies	CACCGGAGGAAAGCGTTTATTGTAG
TAK1 sgRNA#3 Reverse	Integrated DNA Technologies	AAACCTACAATAAACGCTTTCCTCC
TAK1 sgRNA cassettes PCR Forward	Integrated DNA Technologies	TGAGTATTACGGCATGTGAGGGC
TAK1 sgRNA cassettes PCR Reverse	Integrated DNA Technologies	TCAATGTATCTTATCATGTCTGCTCGA
Human MAP3K7 siRNA (Smartpool)	Dharmacon	n/a
Human Control siRNA	Dharmacon	n/a
qPCR Probes		
Rat Map3k7 TaqMan probe	Applied Biosystems	Rn01437015_m1
Rat Tnf alpha TaqMan probe	Applied Biosystems	Rn01525859_g1
Rat Vegf-a TaqMan probe	Applied Biosystems	Rn01511602_m1
Rat Icam-1 TaqMan probe	Applied Biosystems	Rn00564227
Rat beta actin TaqMan probe	Applied Biosystems	Rn00821065_g1
Human MAP3K7 TaqMan probe	Applied Biosystems	Hs00177373_m1
Human VEGF-A TaqMan probe	Applied Biosystems	Hs00900055_m1
Human ICAM-1 TaqMan probe	Applied Biosystems	Hs00164932_m1
Human VCAM-1 TaqMan probe	Applied Biosystems	Hs01003372_m1
Human CXCL8 TaqMan probe	Applied Biosystems	Hs00174103_m1
Human GAPDH TaqMan probe	Applied Biosystems	Hs99999905_m1
Software and Algorithms		

Image Pro Plus	Media Cybernetics	n/a
Incucyte® Live-Cell Imaging and Analysis System	Sartorius	n/a
ImageJ version 1.48	Schneider et al., 2012 https://imagej.nih.gov/ij	n/a
Adobe Photoshop	Adobe. Connor et al., 2009	n/a
AngioTool Software	NIH. Zudaire et al., 2011	n/a
FIJI software	https://fiji.sc/	n/a
Prism 7	GraphPad Software	n/a
R (v4.0.2)	Bioconductor	n/a
Trimgalore (0.4.4)	Babraham Institute	n/a
STAR (2.5.3)	Alexander Dobin	n/a
FeatureCounts (subread 1.6.4)	Liao Y, Smyth GK and Shi W., 2014	n/a
Other		
Pierce RIPA buffer	Life Technologies Australia	89900
Protease inhibitor cocktail	Roche Diagnostics	11697498001
Recombinant Human TNF-α protein	Gibco™	PHC3015
Bolt™ 4-12% Bis-Tris Plus Gel	Invitrogen	NW04120BOX
Immun-Blot® PVDF Membrane	Bio-Rad Laboratories	1620177
Human VEGF protein	R&D Systems	293-VE-010
Matrigel™ Basement Membrane Matrix	Corning	356234
Normal donkey serum	Sigma-Aldrich	D9663
PBS	Gibco™	14190-144
Antibody Diluent	DAKO	S0809
Fluorescence Mounting Medium	DAKO	S3023
BsmBI	New England Biolabs	R0580S
lentiCRISPRv2 plasmid	Addgene	52961
NEB Stable Competent E. coli cells	New England Biolabs	C3040I
pCMV-dR8.91 (plasmid)	RNAiCore; Academia Sinica, Taipei, Taiwan	n/a
pMD2.G (plamid)	Addgene	12259

Lipofectamine 2000	Life Technologies Australia	11668019
0.45µm PDVF syringe filter	Merck Millipore	SLHV033RS
5x PEG-it	System Biosciences	LV810A-1
Opti-MEM	Life Technologies Australia	31985070
EBMTM Plus Basal Medium supplied with EGMTM Plus SingleQuotsTM supplements	Lonza	CC-5035
MycoAlert™ Mycoplasma Detection Kit	Lonza	LT07
Dulbecco's modified Eagle's medium	Life Technologies Australia	11965092
Glutamine	Life Technologies Australia	2503008
Penicillin-streptomycin	Life Technologies Australia	15070063
Fetal bovine serum	Sigma-Aldrich	F9423
TrypLE Express Enzyme (1X)	Life Technologies Australia	12605028

SUPPLEMENTARY METHODS

Generation of TAK1-KO TIME cell line

Three sgRNAs targeting the exonic regions of the human TAK1 gene (**Table S2**) were designed on CHOPCHOP (https://chopchop.cbu.uib.no). The annealed oligos ordered from the Integrated DNA Technologies (IDT; Coralville, Iowa, USA) representing the various sgRNAs were cloned into the BsmBI sites (R0580S; New England Biolabs) of the lentiCRISPRv2 plasmid (52961; Addgene). The plasmid was validated by Sanger sequencing with 3500 Series Genetic Analyzers (Thermo Fisher Scientific; Waltham, MA). HEK293FT cells (R70007; Life Technologies) were plated at 1×10^7 cells per T75 flask one day prior to transfection. Lentivirus was made by co-transfecting lentiCRISPRv2 (15µg), pCMV Δ8.91 (15µg) and pMDG plasmids (1.5µg) (RNAiCore; Academia Sinica, Taipei, Taiwan) in HEK293FT cells using Lipofectamine 2000 (11668019; Life Technologies). Media was changed 5 hours post transfection. The virus-containing media was harvested and filtered through a 0.45 µm PDVF syringe filter (SLHV033RS; Merck Millipore) to remove cell debris 48 hours post transfection. Lentivirus aliquots were precipitated by 5x PEG-it (LV810A-1; System Biosciences) at 4°C overnight and centrifugated at 1500g at 4°C for 30 minutes. After discarding the supernatant, the lentivirus were resuspend within opti-MEM ready for use. Wild type TIME cells (7.5×10^5) were plated at T75 flask before transduction. When grow to 50% confluence, cells were transduced by TAK1 sgRNA lentiCRISPRv2 lentivirus with polybrene (8µg/mL) (H9268; Sigma-Aldrich) for 72 hours. Medium containing 0.5 µg/mL puromycin (P8833; Sigma-Aldrich) was then replaced for another 72 hours to select stable clones which were subsequently cultured as the stable TAK1 KO cell lines.

Single colony selection of TAK1-KO cell line

Once the selected TAK1 KO cell lines reached 80% confluence, cells were trypsinized from the stable cell pool using TrypLE Express Enzyme (1X) (12605028; Life Technologies) to break up any clumps into individual cells. TAK1-KO cells were then diluted to a density of 5 cells/mL and resuspended with 100 ul placed in each well in 96-well plate. After expanding for 7-14 days, colonies of single cells were subsequently cultured as the single colony selected TAK1-KO cell line in culture dishes by trypsinization.

Identification of TAK1 indels in TAK1-KO line

Genomic DNA was isolated from wild type and TAK1-KO TIME cell lines using Quick-DNA Microprep Kit (D3021; Zymo Research) according to the manual. Extracted genomic DNA was used as a PCR template to amplify the target of the TAK1 sgRNA using sequence specific primers (**Table S2**). Briefly, PCR reactions were made up to 25 µL comprising 12.5 µL Q5 Hot Start High-Fidelity

2X Master Mix (M0494; New England Biolabs), 1.25 µL of forward and reverse primers and the

diluted gDNA, under thermocycling conditions of 98°C initial denaturation for 30 seconds, and 30

cycles of 98°C denaturation for 10 seconds, 65°C annealing for 30 seconds, and 72°C extension for 30

seconds with a 72°C final extension for 2 minutes. The PCR amplification product was validated using

electrophoresis using 1% agarose gel (A6361; Sigma-Aldrich). The gDNA was then cleaned by using

a Quick-DNA Microprep Kit (D3021; Zymo Research). Following the first-round of PCR

amplification, a second round of sequencing PCR was performed. The second-round PCR products

were subsequently purified using Agencourt CleanSeq Kit (A29151; Beckman Coulter). The PCR

samples were denatured at 98°C for 5 minutes and sequenced by Sanger sequencing.

Electroretinography (ERG) in rat

Rats were dark-adapted overnight prior to the ERG assessment under fully dark-adapted conditions.

Details for functional assessment are as previously described (58). Briefly, a pair of custom-made

chloride silver active and ring-shaped reference electrodes (99.9%, A&E Metal Merchants, Sydney,

NSW, AU), connected to platinum leads (F-E-30, Grass Telefactor, West Warwick, RI), was place on

the central cornea and around the sclera, respectively. A stainless-steel needle electrode (F-E2-30,

Grass Telefactor) was inserted subcutaneously into the tail of the animal to act as the ground.

Measurements were recorded simultaneously for both eyes. ERG analysis was undertaken as

previously described to return measures for three major waveforms, which are known to reflects the

function of key retinal cells, specifically photoreceptors (a-wave), bipolar cells (b-wave), and ganglion

cells (scotopic threshold response, STR).

Spectral-domain optical coherence tomography (OCT) in rat

Following ERG measurement, rat eyes were imaged using Envisu R2200 VHR spectral domain-OCT

(Bioptigen, Inc., Morrisville, NC, USA). Volume scans consisting of 1000 A-scans per 100 B-scans

(100 horizontal B-scans evenly spaced in the 1.4 mm vertical dimension) centered on the optic nerve

head (ONH; 1.4 × 1.4 × 1.57 mm) were obtained from both eyes. Four B-scans that crossed the ONH

were analyzed using FIJI software (https://fiji.sc/). In each B-scan the inner limiting membrane, retinal

nerve fiber layer, inner plexiform layer and Bruch's membrane were manually segmented by a masked

observer as previously described (58). Total retinal thickness was measured from the inner limiting to

Bruch's membrane. Retinal nerve fiber layer thickness was measured from the inner limiting

membrane to the inner aspect of the inner plexiform layer. Outer retinal thickness was measured from

Bruch's membrane to the outer plexiform layer.

SUPPLEMENTARY FIGURES

Fig. S1 Uniform Manifold Approximation and Projection (UMAP) clustering of the integrated datasets from snRNA-seq data (A) representing the expressions of TAK1 (MAP3K7) and TAK1-associated proteins (TAB1-3) in identified retinal and non-retinal cellular clusters (B).

Fig. S2 (A) Gene set enrichment analysis (GSEA) revealed positive enrichment of TAK1 upstream
pathways in the retinae from patients with retinal neovascularization due to proliferative diabetic
retinopathy. Gene set database includes Biocarta, Pathway Interaction Database (PID), Kyoto
Encyclopedia of Genes and Genomes (KEGG) and WikiPathways (WP). (B) Readcounts of selected
genes related to TAK1 in the retinae from normal control and patients with retinal neovascularization
due to diabetic retinopathy. False discovery rate (FDR) was set at a q value <0.05.

895
896 **Fig. S3** Heat maps generated from unsupervised clustering of genes from Reactome Pathway Database
897 of (A) "TAK1 Activates NFκB by Phosphorylation and Activation of IKKS Complex", (B) "Activated
898 TAK1-mediated p38 MAPK Activation, and (C) "JNK kinases phosphorylation and activation
899 mediated by activated TAK1". False discovery rate (FDR) was set at a q value <0.05.
900

908
Fig. S4 Gene-set enrichment analysis from multiple databases indicated the positive impact of TAK1
upstream pathways in rat OIR retinas. Genome-wide transcriptional analysis of rat OIR retina samples
indicated the differential gene expression associated with positive impact in TAK1 upstream pathways.
Biocarta, Pathway Interaction Database (PID), Kyoto Encyclopedia of Genes and Genomes (KEGG),
and WikiPathways (WP) gene sets were used in GSEA. False discovery rate (FDR) was set at a q value
<0.05.
915

Neovascular area

928
929 **Fig. S5** Uncropped retinal flat mounts. The region indicated by the dashed line was used to show retinal
930 neovascularization (highlighted in white) in Fig. 3B. Oxo: 5Z-7-oxozeaenol. Scale bar: 1 mm.
931
932

933
934
935

936
937 **Fig. S6** Inhibition of TAK1 did not cause retinal vessel remodeling. Retinal flat mount was labelled
938 with IB4 28 days after a single intravitreal injection of vehicle, low (18 ng) or high (90 ng) dose of 5Z-
939 7-oxozeaenol (Oxo) in normoxic rats to examine drug effects on normal retinal vasculature. Scale bar:
940 1 mm.
941
942
943

954
Fig. S7 Generation of CRISPR/Cas9-mediated MAP3K7 knockout (TAK1-KO) in human endothelial cells. (A) Genome of TAK1 (MAP3K7). A sgRNA was designed to target exon 2 of the TAK1 gene. (B) As compared with wild-type (WT) TIMEs, genomic DNA sequences revealed a heterogeneous MAP3K7-KO (M) and a single homogenous MAP3K7-KO (S) TIMEs were established after CRISPR/Cas9 gene editing. (C) Sanger sequencing confirmed that MAP3K7-KO (M) TIMEs had mixed indel profiles, and MAP3K7-KO (S) TIMEs was a single nucleotide insertion, which created as premature Stop Condon in TAK1 exon 2 region. (D) Western blot characterization of TAK1 knockout in MAP3K7-KO (M) and MAP3K7-KO (S) TIMEs.

 Fig. S8 (A) Principal components analysis of RNA-seq data of wild type cells, wild-type (WT) cells and TAK1 knockout (TAK1-KO) TIMEs. Each dot represents an experimental replicate. (A) Volcano plots showing genes that were significantly different between WT and TAK1-KO TIMEs. Red dots refer to genes showing a log2|fold change| > 1 and FDR < 0.05. Labels indicate the significantly up-regulated genes in each data set. (C) Gene set enrichment analysis (GSEA) revealed the top 10 Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways that were positively or negatively enriched in TAK1-KO TIMEs. False discovery rate (FDR) was set at a q value <0.05.

Fig. S9 TAK1 deletion reduced the expression of *TAK1, ICAM-1, CXCL8 and VEGF* genes in human microvascular endothelial cells. qPCR results revealed that TAK1 deletion suppressed the expression of selected cytokines (*CXCL8* and *VEGFA)* and adhesion molecules (*ICAM-1)* in TIMEs stimulated with IL1β for 24 hours (n = 4; technical duplicate from each biological repeat was shown as separated dot plot). Group data are shown as means ± SEM. Statistical analysis was undertaken with one-way ANOVA and Tukey's multiple comparison test; *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$.

1002
Fig. S10 TAK1 knockdown by siRNA inhibited the expression of *TAK1, ICAM-1, CXCL8 and VEGF*
genes in human primary retinal microvascular endothelial cells (HRMECs). qPCR results revealed that
TAK1 knockdown by siRNA targeting TAK1 (siTAK1) suppressed the expression of selected
cytokines (*CXCL8* and *VEGFA)* and adhesion molecules (*ICAM-1)* in HRMECs stimulated with TNFα
for 24 hours (n = 3). Group data are shown as means ± SEM. Statistical analysis was undertaken with
one-way ANOVA and Tukey's multiple comparison test; **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$.
1009
1010

Fig. S11 TAK1 inhibition by 5Z-7-oxozeaenol inhibited the expression of *TAK1, ICAM-1, CXCL8 and VEGF* genes in primary human retinal microvascular endothelial cells (HRMECs). qPCR results revealed that pharmacological inhibition of TAK1 by 5Z-7-oxozeaenol (Oxo) suppressed *ICAM-1*, *PTGS2*, *CXCL8* and *VEGFA* in HRMECs stimulated with TNFα in a dose-dependent manner (n = 3; technical duplicate from each biological repeat was shown as separated dot plot). Group data are shown as means ± SEM. Statistical analysis was undertaken with one-way ANOVA and Tukey's multiple comparison test; *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$.

photoreceptoral and bipolar cell responses (B) or ganglion cell responses (C) in 5Z-7-oxozeaenol-injected eyes (n = 11, red traces) and corresponding vehicle-injected eyes (n = 12, black traces). Group average (± SEM) photoreceptor (D. $P = 0.71$) bipolar cell (E. $P = 0.63$) and ganglion cell (F, $P = 0.41$). Response amplitude were not different between 5Z-7-oxozeaenol-injected eyes and corresponding vehicle-injected eyes. (G) Representative OCT images from a 5Z-7-oxozeaenol- and vehicle-injected eye. (H) Total retinal thickness (TRT), outer retina, and retinal nerve fiber layer (RNFL) thickness were not different between groups ($P = 0.0639$ of TRT, $P=0.0502$ of outer retina, $P = 0.064$ of RNFL). All data are shown expressed as means ± SEM. Statistical analysis was undertaken with two-tailed Student's *t*-test.