Natural loss of function of ephrin-B3 shapes spinal flight circuitry in birds

Baruch Haimson[1], Oren Meir[1], Reut Sudakevitz-Merzbach[1], Gerard Elberg[1], Samantha Friedrich[3],

Peter V. Lovell[3], Sónia Paixão[2], Rüdiger Klein[2], Claudio V. Mello[3*] and Avihu Klar[1*]

1 - Department of Medical Neurobiology, IMRIC, Hebrew University-Hadassah Medical School,

Jerusalem, 91120, Israel

2 - Department 'Molecules – Signals – Development, Max Planck Institute of Neurobiology, Am

Klopferspitz 18, 82152 Martinsried, Germany

3- Department of Behavioral Neuroscience, Oregon Health and Science University, Portland,

Oregon, USA.

* - Corresponding authors

Avihu Klar
Department of medical Neurobiology
Hadassah Medical School
The Hebrew University, Jerusalem
91120
Israel
avihu.klar@mail.huji.ac.il

Claudio V. Mello
Department of Behavioral Neuroscience
Oregon Health and Science University
Portland, Oregon
USA
melloc@ohsu.edu

Abstract

Flight in birds evolved through patterning of the wings from forelimbs and transition from

alternating gait to synchronous flapping. In mammals, the spinal midline guidance molecule

ephrin-B3 instructs the wiring that enables limb alternation, and its deletion leads to synchronous

hopping gait. Here we show that the ephrin-B3 protein in birds lacks several motifs present in

other vertebrates, diminishing its affinity for the EphA4 receptor. The avian *ephrin-B3* gene lacks

an enhancer that drives midline expression, and is missing in Galliformes. The morphology and

wiring at brachial levels of the chick spinal cord resemble those of *ephrin-B3* null mice.

Importantly, dorsal midline decussation, evident in the mutant mouse, is apparent at the chick

brachial level, and is prevented by expression of exogenous *ephrin-B3* at the roof plate. Our

findings support a role for loss of ephrin-B3 function in shaping the avian brachial spinal cord

circuitry and facilitating synchronous wing flapping.

Teaser:

Walking vs flying: Deciphering the organization and evolution of the neuronal network that

controls wing flapping in birds.

Introduction

A critical aspect of locomotion is left-right coordination. Limb alternation is the predominant gait

mode in some amphibians (salamanders), reptiles and terrestrial mammals, while synchronous

movements are manifested in the legs of other amphibians (frogs) and some marsupials, and in

the wings of bats and birds. Based on classical electrophysiological studies in cats and rodents, and

genetic manipulations of cell fate and axonal trajectories in mice, a modified "half-center

oscillator" model has been proposed that explains limb alternations in mammals (*1, 2*). The core

elements are mutually inhibitory neuronal centers on each side of the spinal cord that produce

oscillatory excitatory output. This output is simultaneously transmitted to interneurons that

innervate motoneurons, termed pre-motoneurons (pre-MNs). These include ipsilateral excitatory

pre-MNs that activate motor activity on one side, and commissural inhibitory pre-MNs that inhibit

motor output on the other side (Fig. 1A) (*3-6*).

The axon guidance molecule ephrin-B3 and its cognate receptor EphA4 are key regulators of

alternating gait. In this family, ligand binding to the receptor typically results in axonal repulsion

(forward signaling), but ephrin receptors can also act as ligands for ephrins (reverse signaling),

prompting repulsion or adhesion (*7*). In wt mice, ephrin-B3 serves as a midline repulsive cue,

preventing EphA4-expressing axons of excitatory pre-MNs from aberrantly crossing the ventral

and dorsal midlines (Fig. 1A) (*8-12*). Transition from left-right alternation to synchrony occurs in

mice mutated in the genes encoding ephrin-B3, EphA4*,* or the downstream signaling molecule α2-

chimaerin, referred to as *hop* mutants. These mice show ectopic commissural projections of

otherwise ipsilaterally-projecting excitatory pre-MNs (Fig. 1B) and walk with synchronous

movements of the hind- and forelimbs, producing a hopping gait. Furthermore, impairment of the

neuroepithelial organization of the roof plate (RP), which leads to partial absence of ephrin-B3

expression, results in rabbit-like hopping (*13, 14*).

Birds have evolved from dinosaurs that use alternating gait (*15, 16*). Their adaptation for

flight includes substantial structural specializations of the forelimb and of the spinal cord circuits

that enable synchronous limb movements. Classical transplantation experiments in chick support

a role for neuronal networks at the lumbar and brachial spinal levels in the control of the

locomotion pattern of the hind- and forelimbs, respectively. In chicks where brachial segments are

surgically replaced by lumbosacral segments, the movement of the wings become alternation,

whereas coordinated flapping movements are not observed (*17*). In the complementary

experiment, transplantation of brachial segments to the lumbar cord result in synchronous,

hopping-like leg movements (*18*). These findings support the hypothesis that local circuits in the

lumbar and brachial spinal cord instruct level-specific alternate or synchronous movements,

respectively.

Flight requires spinal circuits that enable the bilateral coordination of forelimb movements.

It thus seems reasonable to hypothesize that the emergence of synchronous wing flapping in birds

involved a transition from ipsilateral excitatory/contralateral inhibitory inputs to a pattern of

bilateral co-excitatory input to wing motoneurons. Such a transition likely required changes in the

pre-motor spinal wiring. Notably, mutations in the midline guidance molecules Netrin1 or ephrin-

B3 in mice lead, respectively, to a reduction in cross inhibition and an increase in cross excitation

(*9, 19, 20*). Netrins were initially described in chick, and have well characterized activity and roles

in commissural axonal guidance (*21, 22*). However, little is known about ephrin-B3 in chick, and

more generally in birds.

To explore a possible role of ephrin-B3 in locomotion patterns in birds, we have conducted a

set of genomic, molecular and neuroanatomical studies, using both chick and zebra finch as

experimental organisms. We found an ephrin-B3 gene ortholog (EFNB3) in several bird species and

lineages, however their open reading frames (ORFs) were considerably shorter, and some domains

critical for forward and reverse signaling were absent or heavily modified compared to non-avian

tetrapods. We also observed marked loss of dorsal midline ephrin-B3 expression in the brachial

spinal cord of zebra finch, likely due to an avian loss of a critical enhancer element. An EFNB3

ortholog could not be found in chicken and related species, suggesting a gene loss in Galliformes.

Spinal cord interneurons in chick showed extensive dorsal midline crossing at brachial levels,

similar to ephrin-B3 null mice. In contrast, we observed little dorsal crossing at lumbar levels due

to the presence of the glycogen body, which serves as a physical barrier to neurite crossing. The

crossing at brachial levels was largely prevented by dorsal midline expression of mouse ephrin-B3.

These findings are consistent with a role of ephrin-related mechanisms in spinal wiring patterns

that may subserve limb alternation. They also support the notion that diminished barriers to

dorsal midline crossing may have favored synchronous limb movements and facilitated the origin

and/or maintenance of coordinated flight in birds.

Results

EFNB3 gene in birds

We have conducted an extensive and thorough analysis of the genomic region containing

EFNB3 and syntenic genes in birds and non-avian outgroups. Our approach required extensive

curation of gene predictions in annotated databases, permissive BLAST searches of genomes, WGS

databases and p-reads (from Pacbio assemblies) using avian EFNB3 and syntenic gene queries, and

RefSeq alignment searches of unplaced genomic scaffolds and reads (details in Methods). We

found EFNB3 in a broad range of bird species and orders, including several songbirds, some eagles,

an owl, a few cormorants, as well as the seriema, a hummingbird, a ruff, a crane, a stork, and

some Anseriformes (duck and goose). WRAP53 was immediately upstream of EFNB3 in all birds

where synteny could be examined and in all non-avian organisms studied, including mammals,

non-avian sauropsids and amphibians (examples in Fig. 1C,D), providing strong support for EFNB3

orthology. A gene prediction was present in only few species, whereas in several others EFNB3

was partial, possibly due to incomplete sequences in Illumina assemblies.

Notably, we could not identify EFNB3 in any Galliformes (Fig. 1D, red square), even in

species with high quality Pacbio assemblies and no local gaps. In chicken, the genes immediately

syntenic to EFNB3 in other birds (WRAP53 and KDM6B) were on different chromosomes,

suggestive of a local inter-chromosomal rearrangement in this species, with no trace of EFNB3

(Fig. 1D, chicken). EFNB3 was also absent in the chicken WGS and Pacbio p-reads, and in numerous

RNAseq databases. In quail, a WRAP53 prediction and an unannotated KDM6B (found by BLAST)

were found on the same incomplete scaffold. Several Pacbio p-reads from the current assembly

(*23*) bridged the local gaps but revealed no trace of EFNB3 (Fig. 1D, quail). EFNB3 was also not

detected in the assemblies or WGS of other Galliformes (turkey, pheasant, guinea fowl), even

though the syntenic genes were sometimes present in unplaced scaffolds (not shown). A scaffold

in the unannotated sage grouse genome contained traces of WRAP53 and KDM6B but no evidence

of EFNB3 sequences (Fig. 1D, sage grouse). We also could not find EFNB3 in ratites, but the data

were too sparse to conclusively prove or disprove an EFNB3 loss in this basal group

Importantly, in most birds where EFNB3 synteny could be verified KDM6B was immediately

downstream, whereas in non-avian sauropsids and mammals DNAH2 was interposed between

EFNB3 and KDM6B (Fig. 1D, blue). Given there were no gaps in several avian assemblies between

EFNB3 and KDM6B, this observation supports an avian-specific loss of DNAH2, as previously

indicated (Lovell et al., 2014). As we show later, this finding has important implications for EFNB3

expression regulation. The exception among birds was the kakapo, where the receptor gene

EphB4 was downstream of EFNB3, the only such case we are aware of. In sum, the EFNB3 gene is

clearly present in a broad range of Neoaves, as well as in Anseriformes, but seems absent in

Galliformes as a group (Fig. 1C,D). We suggest this is possibly due to a loss that occurred after their

split from Anseriformes, an estimated 46 million years ago (*16*), but prior to their later

diversification.

EFNB3 gene and protein structure in songbirds

Songbirds were the avian group with the most complete information on EFNB3. In zebra finches,

EFNB3 is largely conserved compared to non-avian tetrapods, consisting of 5 exons and a large 3'-

UTR (Fig. S1A), but predicted avian ephrin-B3 proteins are shorter compared to amphibian,

reptilian and mammalian EFNB3 (247-267 vs 332-340 aminoacid residues; Fig. S2). The

intracellular domain is shorter in birds, and while the PDZ motif is intact, the entire SH2/3 binding

motif, which is required for reverse signaling (*24, 25*), as well as the tyrosine residues that are

phosphorylated upon activation of the reverse signaling in other organisms, are missing (Fig. 1E;

Fig. S2), suggesting that reverse signaling is impaired. Birds also lack an extracellular heparin

binding motif within the region juxtaposed to the transmembrane (TM) domain (*26, 27*), and

several residues in the tetramerization interface within the receptor binding domain (RBD),

required for normal signaling through the formation of receptor-ligand (Eph–ephrin) complexes

(*28*) (Fig. 1E; Fig. S2). The region in the dimerization interface where an L to P (L111P) substitution

leads to a hopping mouse phenotype (*29*) is conserved in birds and shows other substitutions

(DLDL_108-111_QRDV) compared to mammals (Fig. S2). Unlike ephrin-B3, the predicted avian

ephrin-B1 and ephrin-B2 proteins are highly similar to the mammalian proteins in terms of size,

sequence and sub-domain partitioning.

To test whether these structural differences affect cell trafficking of the avian ephrin-B3,

we expressed an amino-myc tagged isoform of the mouse and zebra finch genes in COS-7 cells.

Surface staining revealed that both proteins are presented on the cell membrane (Fig. 2A). Next,

we used an assay where a chimeric protein composed of the extracellular domain of chicken

EphA4 fused to placental alkaline phosphatase is tested for binding to different EFNB3 orthologs

expressed in transiently transfected COS-7 cells, noting that chick and the zebra finch EphA4

proteins share 98.4% identity. We found that chicken EphA4 binds mouse EFNB3, but not zebra

finch EFNB3 (Fig. 2B,C).

Expression of ephrin-B3 in zebra finch

The presence of EFNB3 gene in songbirds prompted us to examine its expression pattern in the

zebra finch spinal cord. *In situ* hybridization with antisense riboprobes revealed ephrin-B3 mRNA

expression in both the floor plate (FP) and roof plate (RP) of the brachial spinal cord in P0 zebra

finches, with apparent lower expression in the RP (Fig. 2D). Quantification confirmed that ephrin-

B3 expression is higher in the FP and declines along the RP in the ventral-to-dorsal direction (Fig.

2D'). This contrasted with mouse at a comparable developmental stage, where expression is

similarly high in both the FP and RP (Fig. 2E), with no dorsal decline (Fig. 2E').

We reasoned that the modified ephrin-B3 expression seen in zebra finch compared to

mouse might relate to changes in regulatory elements. Using UCSC's genome browser, we

searched for enhancer hallmarks in mouse EFNB3 and its flanking genes. We found that mouse

DNAH2, located immediately downstream of EFNB3 (Fig. 1D, blue), contains several features

consistent with presence of an enhancer element in its 1^{st} intron (Fig. S3). Because the DNAH2

gene is not present in birds (Fig. 1D blue; Lovell et al., 2014), birds may lack an important EFNB3

enhancer element. To test whether this DNAH2-embedded region would be sufficient to confer

midline RP expression in the spinal cord of a bird species, a reporter gene cloned downstream of

the putative enhancer element was electroporated into the chick spinal cord at E3. Expression of

the reporter in both the FP and RP was evident at E9 and E15 (Fig. 2F,G), similar to mouse where

endogenous ephrin-B3 midline expression occurs during embryogenesis and early post gestation

(E15 chick is equivalent to mouse P2), and contrasting with the more ubiquitous expression of the

co-electroporated GFP under a ubiquitous promoter (Fig. 2F,G, insets). Notably, *DNAH2* is missing

in all bird species examined to date but it is present in reptiles (Fig. 1D; (*30*)), suggesting the loss of

the EFNB3 regulatory element is specific to the avian lineage.

Expression of ephrin-B family ligands and EphA4 in chick spinal cord

We next used *in situ* hybridization to examine expression of ephrin-B1 and -B2 at the chick

brachial cord, as both can mediate axonal repulsion through the EphA4 receptor and could thus

potentially compensate for an EFNB3 loss. Both ephrin-B1 and B2 were expressed in the FP and RP

at E10 (Fig. 3A,C). At E16, when the spinal circuitry elicits simultaneous wing flapping (*31*), ephrin-

B1 expression persisted in the FP but dorsally it became restricted to the ventral aspect of the RP

(Fig. 3B), whereas ephrin-B2 expression was detected only at the ventral tip of the RP (Fig. 3D).

Expression of ephrin-B genes in chick spinal cord is thus largely restricted to the FP, low in ventral

RP, and absent in the dorsal RP.

Next, we studied the expression of the receptor EphA4. In mouse, interneurons that

express EphA4 are excluded from the dorsal midline, but many of these cells invade the dorsal

midline in *EphA4* and *ephrin-B3* null mutants (*32*). In the chick, expression is detected throughout

the spinal cord gray matter at E10, with high levels in scattered interneurons, in the lateral

subdivision of the lateral motor column (LMC$_l$), and in the medial motor column (MMC), but

absent in the dorsal midline (Fig. 3E). At E16, EphA4 shows high expression in discrete cells

distributed all over the gray matter, including adjacent to and within the dorsal midline (Fig. 3F,

white arrow in inset).

When we assigned x-y coordinates to EphA4-expressing cells, we observed their midline

invasion at E16 compared to E10 (Fig. 3E',F',G). Thus, the presence of EphA4-expressing neurons in

the dorsal midline correlates with the temporal reduction of ephrin-B1-2 levels. Taken together

with the lack of ephrin-B3, we conclude that a molecular barrier that prevents midline crossing is

missing in the chick dorsal spinal cord. This contrasts markedly with mouse, where midline

expression of ephrin-B3 persists at high levels both ventrally and dorsally at post gestation stages

(*9*).

Extended roof plate of the chick brachial cord

To further investigate how the natural loss of EFNB3 might have affected the organization of the

avian spinal cord, we examined several parameters that differ significantly between wt and *hop*

mutants, namely the size and axonal decussation at the RP, and the somata distribution and

neurotransmitter phenotype of commissural pre-MNs (Figs. 4-6).

In wt mice, the length of the FP was relatively constant along the longitudinal axis, but the

RP was substantially shorter at limb levels, resulting in low RP:FP ratios at those levels (Fig. S4).

Thus, the RP:FP ratio can serve as a reasonable proxy for the size of the RP. The length of the RP at

limb levels was markedly enlarged in EphA4 knockout mice (Fig. 4B,D) in comparison to control

mice (Fig. 4A,C), with a significantly higher RP:FP ratio in the homozygous mutant (1.68±0.22,

mean of RP:FP ratios ± SD) than in control mice (0.34±0.13). In birds, an elongated RP was evident

at brachial levels in E15 chick (Fig. 4E) and P8 zebra finch (not shown), with RP:FP ratios of

2.18±0.2 and 1.53±0.08, respectively, thus comparable to EphA4 knockout mice (Fig. 4G). Hence,

the morphology of the avian brachial RP is similar to that of *hop* mutants. At the lumbar sciatic

level, the dorsal midline in birds is occupied by an ovoid gelatinous mass - the glycogen body, an

avian specific organ (GB; Fig. 4F) (*33-35*). These features differ from those in alternating limbed

reptiles (lizard, turtle) where the RP is short, as in wt mice (*36-38*)

Patterns of axonal midline crossing in the chick spinal cord

Axon fibers that cross the RP in rodents originate mostly from sensory neurons and inhibitory

interneurons (*39, 40*). To further study the extent of dorsal midline crossing in the chick embryonic

spinal cord, we examined immunoreactivity for neurofilament (3A10) and for sensory axons

(TAG1) at E15 and E17. We observed neurofilament-positive axons crossing the brachial FP in a

tight bundle (Fig. 5A, yellow arrows in insets), whereas those crossing the dorsal midline were less

fasciculated and sparsely distributed along the dorsal RP (Fig. 5A, white arrows in insets).

Quantification showed that 44.8±23% of midline crossing axons at the brachial level were in the

RP (Fig. 5G, brachial). TAG1-expressing sensory axons did not appear to cross the midline at this

level (Fig. 5B). To further interrogate midline axonal crossing, a membrane-tethered GFP was

electroporated at E3 into one side of the spinal cord. At E15, GFP-labeled axons crossing the

ventral and dorsal midline were evident (Fig. 5C, yellow and white arrows in insets). Interestingly,

a few labeled cells were apparent on the contralateral side dorsally, suggesting that some cells

may migrate through the RP at the brachial level (Fig. 5C, white arrowheads in insets).

At the lumbar sciatic level, axonal decussation was apparent only through the FP (Fig. 5D-

E), with no axons crossing the GB. At the lumbar crural level, which is devoid of GB, dorsal midline

crossing was apparent, with 31±12% of crossing fibers occurring through the RP (Fig. 5F,G, crural).

Notably, the sciatic nerve innervates muscles of the shank and foot, while nerves originating from

the crural plexus innervate the thigh muscles (*41, 42*). Importantly, in the swing phase of birds, the

ankle flexion leads to the elevation of the feet, while the knee is relatively stable. Hence, the

sciatic nerve mediates the flexion and extension of the ankle and foot joints during stepping.

Significant dorsal midline decussation was also apparent at the cervical, brachial and thoracic

levels (Fig. 5G), as well as sacral levels (not shown), suggesting that the GB serves as a physical

barrier for axonal decussation at the sciatic plexus level, thus allowing hindlimb alternation. In

contrast, dorsal midline crossing at brachial levels correlates with synchronous wing (forelimb)

movements, similar to that seen in *hop* mice.

Connectivity of pre-MNs in the chick spinal cord

The synchronous gait of *hop* mice results from ectopic midline crossing of excitatory pre-MN

through the FP and RP (Fig. 1B) (*9, 43*). To analyze the patterns of connectivity of pre-MNs at the

lumbar and brachial levels of the chick, we injected the trans-synaptic PRV virus (*44, 45*) into the

leg, the distal wing musculature, and the main flight muscle, the pectoralis major. We observed

substantial differences in the densities and laminar distributions of commissural pre-MNs (cpre-

MNs) at lumbar versus brachial levels (Fig. 6A-C, right side). cpre-MNs accounted for 7.81% of

labeled cells at the sciatic level, and for 25.3% and 34.8% of brachial level cells labeled from the distal wing and the pectoral muscle, respectively.

The distribution of ipsilateral pre-MNs (ipre-MNs) was similar at lumbar and brachial levels, occupying the medial spinal cord (Fig. 6A-C, left side; Fig. S6). In contrast, the distribution of cpre-MNs was markedly different across levels. At lumbar levels (both sciatic and crural) we observed cpre-MNs ventro-medially, with most cells ventral to the central canal (Fig. 6A, right side; Fig. S6). At the brachial level, cpre-MNs of the distal wing musculature and the pectoral muscle were distributed ventrally and dorsally to the central canal (Fig. 6B-C, right side). Likewise, decussation of pre-MNs differed markedly between lumbar and brachial levels, with axons crossing only the FP at the lumbar level (Fig. 6A'), but both the RP and the FP at the brachial level (Fig. 6B'-C').

Synchronous gait in mice *hop* mutants is also associated with pre-MNs that innervate motoneurons bilaterally (*43*). To directly test whether the chick brachial cord contains such cells, we injected PRV-GFP and PRV-cherry into the left and right pectoral muscles, respectively (Fig. 6D, left). 10.9% of pectoral cpre-MNs were double-labelled, most located above the central canal and adjacent to the dorsal midline (Fig. 6D), with neurite processes oriented toward the dorsal midline (Fig. 6D'). Hence, like in *hop* mice, a subpopulation of the brachial pre-MNs innervate motoneurons bilaterally.

To examine the neurotransmitter phenotype of spinal pre-MNs, we processed sections from hindlimb and wing PRV-injected embryos for *in situ* hybridization with a marker for excitatory neurons (vGlut2). At the lumbar level, 46.5% of the ipre-MNs and 38.6% of the cpre-MNs were excitatory. At the brachial level, 35% of the ipre-MNs and 38% of the ventral cpre-MNs were excitatory. Of the brachial-specific cpre-MNs that reside dorsal the central canal, 47% were excitatory (Table S1). The chick brachial-specific pre-MNs are thus comparable to the ectopic cpre-MNs in *hop* mutants in location, wiring and excitatory phenotype (*14, 43*).

Exogenous ephrin-B3 expression in the chick RP prevents midline axonal crossing

Our observations so far suggest that dorsal midline crossing and the wiring pattern of brachial pre-

MNs in chick spinal cord might be linked to a lack of ephrin-B3 expression due to the genomic loss.

To test this hypothesis, we examined whether expression of mouse ephrin-B3 in the chick would

affect RP neurite crossing. Mouse ephrin-B3 or GFP, cloned into a Cre-dependent expression

vector under the RP-specific Wnt1 enhancer element (*46*), were co-electroporated into the E3

chick spinal cord (Fig. 7A), and the extent of RP neurite crossing analyzed at E15. To exclude

possible effects of exogenous ephrin-B3 in reverse signaling, we also used an ephrin-B3 isoform

that lacks the intracellular domain (ephrin-B3-ΔC). Ephrin-B3 expression was confined to the RP

(Fig. 7B), and the number of midline crossing neurites was significantly lower for ephrin-B3 (7.1 ±

3.1) or ephrin-B3-ΔC (7.1 ± 2.6) than for GFP (16.5 ± 4.6) (Fig. 7C-E). Hence, as in mammals,

midline ephrin-B3 expression is sufficient to prevent midline crossing.

Next, after targeting ephrin-B3 expression to the RP at E3, PRV was injected into the wing

musculature at E14 (Fig. 8A), and the numbers and distribution of cpre-MNs examined at E15.5. In

controls, both ipre-MNs and cpre-MNs were seen, the latter both dorsal and ventral to the central

canal (Fig. 8B), and their neurites crossed at both the FP and RP (Fig. 8B'). In ephrin-B3-expressing

embryos, fewer cpre-MNs were seen, most ventral to the central canal (Fig. 8C,C',C''), and most

neurites were observed to cross the FP (Fig. 8C,C',C'', right insets). Short cell extensions, possibly

dendrites originating from ipre-MNs, did cross the RP (Fig. 8C, C', right insets). We note that *in ovo*

electroporation results in mosaics where only a portion of the cells express the transgene. Our

efficiencies varied from 5% to 64% (sections with midline expression of ephrin-B3), thus some

axons may circumvent ephrin-B3 and cross the midline close to non-EFNB3 expressing RP cells

(Supp movie).

Next, we analyzed the spatial distribution of cpre-MNs in five embryos with variable

efficiencies of wnt1::ephrin-B3 expression (Fig. 8D). A dorsal to ventral shift in the location of cpre-

MNs is apparent when comparing the embryos with lower vs higher exogenous ephrin-B3

expression in the RP (1-3 vs 4-5 in Fig. 8D), suggesting a reduction in the size of the brachial-

specific dorsal population of cpre-MNs. To further quantify this observation, we subdivided cpre-

MNs into ventral and dorsal groups, based on location relative to the central canal, and examined

their numbers compared to ipre-MNs. The ratios between pre-MNs populating the different zones

were calculated and correlated with the efficiencies of ectopic expression of ephrin-B3, using

linear regression (Fig. 8E-H). The ratio of ventral cpre-MNs to ipre-MNs was similar in controls and

all experimental embryos (Fig. 8E), thus ephrin-B3 RP expression did not affect this population.

The ratio of dorsal cpre-MNs to ipre-MNs was reduced in proportion to higher ephrin-B3

expression (Fig. 8F), as was their ratio to ventral cpre-MNs (Fig. 8G). The ratio of all cpre-MNs to

the ipre-MNs was also negatively correlated with increasing ephrin-B3 expression (Fig. 8H). We

conclude that robust ephrin-B3 RP expression results in fewer cpre-MNs that cross through the RP.

Discussion

During vertebrate evolution, most terrestrial species adopted alternating limb motion as an effective gait strategy, while adaptation to powered flight in birds emerged via morphological changes that patterned wings from forelimbs, and required transition from alternate gait to synchronous flapping. Our genomic analysis indicates that the gene encoding the midline axonal repellent ephrin-B3 is missing in chicken and other Galliformes. In songbirds, the EFNB3 gene encodes an ephrin-B3 protein that does not bind to the EphA4 receptor, and whose expression in the spinal cord is reduced along the roof plate, probably due to the loss of a downstream enhancer. Morphological and circuit analyses of the chick spinal cord revealed similarities at the brachial level with mutant mice for components of the ephrin-B3/EphA4 pathway, while at the lumbar level it resembled wt mice (Fig. 8I,J). Importantly, dorsal midline expression of exogenous (rodent) ephrin-B3 in the chick brachial spinal cord prevents crossing of cpre-MN axons. Altogether, our findings support the notion that a reduction in ephrin-B3 expression and/or function plays an important role in shaping the wing level circuitry in the avian spinal cord.

Loss of ephrin-B3 function in birds

Changes in gene copy number, enhancers, and protein sequence, arguably some of the foremost molecular events in evolution, are thought to have influenced locomotion patterns in vertebrates. For example, a point mutation that leads to a premature stop codon in *DMRT3* enabled Icelandic horses to develop their characteristic Tölt ambling gait (*47*). A modification and subsequent loss of a regulatory element of *GDF6* during the transition to bipedalism in humans is thought to have played a role in foot digit shortening (*48*). Limb loss in snakes has been shown to be a result of mutations in a limb-specific enhancer of Sonic hedgehog (*49*). Loss of function for ephrin-B3 leads to midline crossing in the spinal cord of rodents, resulting in a remarkable hopping gait (*9, 10*).

Our findings now indicate that the ephrin-B3 gene has undergone significant modifications in birds

compared to non-avian groups, likely impacting flight control circuits in the avian spinal cord.

Absence of *DNAH2* is seen in all birds where this genomic region could be assessed,

suggesting that the loss of a putative ephrin-B3 enhancer was an early event in avian evolution, or

possibly even in dinosaurs. The apparent loss of EFNB3 in chick and other Galliformes was likely a

more recent event specific to that basal clade, since EFNB3 is present in Anseriformes and in

several Neoaves, including the more recently evolved Passeriformes. It is hard to estimate the

timing of mutations in the coding region, since high quality sequence is limited to a few species.

Nonetheless, similar protein changes compared to mammals and non-avian saurospids are present

in eagles and in songbirds (Fig. S2), which originated ~65 and ~45 million years ago, respectively

(*16*). High-quality sequences of this region in the most basal avian clade, Palaeognathae, currently

not available, will be critical for better understanding the timing and extent of avian EFNB3

changes.

In zebra finch, besides a predicted lack of reverse signaling due to loss of specific domains,

we provide clear evidence that forward signaling through the EphA4 receptor is impaired. The lack

of binding to avian EphA4, for example, is likely related to substitutions in residues 108-111 (in the

mouse protein). This region is within the G -strand which forms a loop with the H -strand, and is

part of the ligand-receptor dimerization interphase (*50*). The D108Q, L109R and L111V

substitutions may therefore affect receptor binding by disrupting the formation of the G-H loop.

Interestingly, ephrin-B3 is also a known receptor for the Nipah paramyxovirus (*Henipavirus* group)

and the G-H loop is critical for infectivity (*51*), as shown by mutations in the H -strand of ephrin-

B3 (*52*), suggesting possible related changes in spinal circuitry and viral susceptibility mechanisms.

Phenotype of avian and *hop* mouse spinal cords

Our data show that the brachial spinal cord of the chick, which lacks an EFNB3 gene, shares

similarities with mouse *hop* mutants in its morphology, and in the wiring and excitatory/inhibitory

balance of cpre-MNs (Fig. 1B, 8J). The significance of the dorsal midline expansion, anatomically

the most profound alteration in *hop* mutants, is unclear. While synchronous motoneuron activity

is seen in the spinal cord of *hop* mutants using an *in vitro* fictive locomotion assay, severing of the

dorsal midline does not restore left-right alternation (*8, 53*), arguing that bilateral coupling is

facilitated by ventral midline crossing. However, *in vivo* disruption of the roof plate integrity is

sufficient to induce synchronous gait in mice (*13, 14*). It is conceivable that in *hop* mutants,

proprioceptive signals from weight-load receptors, which are used only *in vivo*, inhibit the bilateral

coupling mediated by the dorsally crossing axons. In support of this hypothesis, removal of EphA4

from a subpopulation of dorsal interneurons results in ectopic dorsal midline crossing.

Furthermore, the over-ground gait of these mice is normal, while during air-walking and swimming

their limbs move in synchrony (*43*). In birds, dorsal midline crossing might be sufficient to instruct

the bilateral coupling of the wings motoneurons, since weight load without ground support during

flight does not affect the wings.

The importance of excitatory neurons for synchronous gait in mouse *hop* mutants is shown

by conditional targeting of EphA4 in vGlut2 neurons, which results in hopping gait (*8*).

Furthermore, a rather small shift in the excitatory/inhibitory balance in cpre-MNs, from 21%

excitatory and 46% inhibitory in wt to 26% and 35% in the EphA4 null mouse, is sufficient to

induce synchronous gait (*9, 53*). We found that the brachial dorsal population of cpre-MNs in chick

is moderately but significantly more enriched (–47%) with excitatory neurons than in wt mice.

Furthermore, the difference between this population and the brachial ventral and lumbar cpre-

MNs, 38% and 35% of excitatory neurons, respectively (Fig. 6, Table S1), is similar to the difference

between wt and *hop* mutant mouse. These observations support the notion that the brachial-

specific population of cpre-MNs is sufficient to support synchronous activation of motoneurons

involved in wing flapping.

Intriguingly, the glycogen body (GB) (Fig. 8I) is a prominent but poorly understood dorsal

lumbar gelatinous structure that arises from roof plate cells (*33, 34*) and is present across birds

(*54, 55*). Our study has uncovered evidence that the GB serves as a major physical barrier to dorsal

midline crossing at sciatic lumbar levels in the chick. We thus suggest that the GB is likely a major

contributor to the alternating gait in birds, analogous to the molecular barriers to midline crossing

in mice.

In summary, we provide evidence linking loss of ephrin-B3/ EphA4 signaling to the

emergence of synchrony-mediating circuitry in the avian brachial spinal cord. While our findings

do not exclude the importance of other molecular pathways, they suggest that modulation of

ephrin-related mechanisms may have been an important contributor to the origin of flight in birds.

	Materials and Methods

	Genomics

	The EFNB3 genomic region in birds has presented several difficulties for sequencing and

	assembling, likely due to very high GC content and density of poorly characterized repeat

	elements. The gene and its syntenic region are not present, or present only in partial form, in most

	Sanger- and Illumina-based avian assemblies, but are better represented in the more recent

	PacBio assemblies (Korlach et al., 2017). Furthermore, this region is likely part of a short avian

	micro-chromosome, which until the recent Swainson's thrush genome had not been assembled

	and/or annotated in any species. Our current study of EFNB3 in birds has therefore required

	considerable search, annotation and curation efforts.

	Briefly, we initially examined the annotated databases (NCBI's RefSeq, Ensembl) available

	for >50 avian species, searching for gene predictions annotated as EFNB3 or ephrin B3-like, and

	using reciprocal BLAT alignments and verification of conserved synteny of the predictions to

	confirm their orthology with humans and non-avian vertebrates. We were able to identify EFNB3

	and to verify its orthology in seven songbird species (including zebra finch), an eagle, and a parrot

	(kakapo). Notably, in the Swainson's thrush EFNB3 and syntenic genes have been assigned to

	chr37, the first case any genes have been assigned to that avian micro-chromosome. We also note

	that two apparently identical zebra finch EFNB3 models (discontinued Gene IDs: 115492780 and

	115492629) were present in an unplaced scaffold (NW_022045341.1) of an early PacBio assembly

	(bTaeGut1_v1.p; (*56*), but have been recently discontinued as this entire region is not present in

	the latest zebra finch Pacbio assemblies in NCBI. We used the discontinued XM_030261704.1

	transcript prediction in our study, noting that it is well supported by ESTs and transcriptome data

	(Fig. S1A,B). Ensembl has EFNB3 predictions for the zebra finch, kakapo, and sparrowhawk, the

	latter in a short segment and right upstream of KDM6B. We also BLAST searched avian and non-

avian RefSeq databases using as queries the genomic regions just downstream of avian gene

predictions annotated as WRAP53 (or WD repeat containing antisense to TP53, a.k.a. telomerase

Cajal body protein 1), or just upstream of avian gene predictions for KDM6B (or lysine specific

demethylase 6B). This effort identified EFNB3 in an owl and in another songbird species, even

though no EFNB3 predictions were present. We also BLAST searched all existing avian genomes

(May 2020) as well as the respective WGS databases, using as queries the existing avian EFNB3

predictions, and manually examined all hits with scores >50, excluding cases that did not show top

reciprocal alignments (typically other EFNB family members). This effort identified EFNB3 in a

cormorant, and in a few other passerines and Anseriformes with limited synteny information, as

well as in several species with incomplete data for synteny verification (a duck, a goose, several

other passerines, eagles and cormorants, the seriema, a ruff, a crane, and a stork). As an EFNB3

was not present in the current Anna's hummingbird PacBio assembly (the only such case for a

non-Galliformes avian PacBio assembly), we also BLAST searched the pre-assembled reads (p-

reads) for this assembly, and obtained evidence for an ortholog with similar synteny as in other

bird groups. We also conducted BLAST searches for EFNB3 in the current assemblies of chicken

(Pacbio-based GRCg6a) and quail (hybrid Coturnix japonica 2.1), as well as in the corresponding

Pacbio p-reads and previous assemblies. These searches were extensive and permissive (>30 hit

scores), followed by manual verification of detected hits. We also searched the BBSRC ChickEST

Database and several brain transcriptome databases at E10, E17, P0 and 10 weeks old chick

available in NCBI (ERR2576391, ERR2576392, ERR2576465, ERR2576464, ERR2576488,

ERR2576584, ERR2576585, ERS2480079), but found no evidence of EFNB3.

Animals

Fertilized White Leghorn chicken eggs (Gil-Guy Farm, Israel) were incubated in standard conditions

at 38 °C. All experiments involved with animals were conducted in accordance with the designated

Experiments in Animals Ethic Committee policies and under its approval. Experiments were

performed on adult male and female mice. EphA4^{lox} mice (*57*) were crossed to PGK-Cre mice (*58*)

to generate full knock-out of the EphA4 gene. Animals were kept and used in accordance with

regulations from the government of Upper Bavaria.

DNA

For the binding assay in COS cells, the mouse ephrin-B3 was cloned into pSecTag-B plasmid (Termo

Fisher Scientific, Waltham, USA). Six repeats of the myc epitope were cloned at the carboxyl end

of ephrin-B3. For cell surface detection of the mouse ephrin-B3, the kappa chain signal sequence

plus 3 repeats of the myc epitope were cloned upstream to amino acid 30 of the mouse gene. The

sequence of the full-length zebra finch ephrin-B3 transcript was reconstructed from RNA-seq

reads (Fig. S1B,C) using the Trinity platform (*59*). The sequence matched the exonic sequence of

the EFNB3 transcript prediction (XM_030261704.1). For cloning the zebra finch ephrin-B3, an

optimized sequence (available upon request) of residues 21-171 (Fig. S2), predicted to encode the

extracellular domain, was synthesized (ITD, Coralville, USA) and cloned downstream to an Ig kappa

chain signal sequence in the pSecTag-B plasmid. The transmembrane and intracellular portions of

the zebra finch transcript were obtained by fully sequencing the EST DV945855 cDNA clone. The

synthetic 5' portion of the gene plus the 3' DV945855-derived sequence were fused using a PCR-

generated EcoRV site at the synthetic 5' part and an internal SmaI site in the 3' DV945855-derived

sequence. For detection of the *in vitro* transcribed protein, 4 repeats of the myc epitope were

inserted between the Ig kappa chain signal sequence and predicted residue 21 of zebra finch

ephrin-B3. The EphA4-AP fused gene was generated by cloning amino acids 1-543 of the chick

EphA4 (*60*) into the pSecTag-B plasmid. The SEAP gene was cloned downstream to EphA4.

Cell culture and transfection

COS-7 cells (ATCC) were cultured in DMEM media (Sigma, Saint Louis, USA) supplemented with

10% fetal bovine serum 100U/ml penicillin, 100 mg/ml streptomycin and 2 mM glutamine

(Biological industries, Beit HaEmek, Israel). COS-7 cells were transiently transfected with

appropriate cDNA using PEI (Polyethylenimine, linear, MW 25000; Polysciences, Inc).

AP-fusion protein binding experiments

To produce secreted EphA4-AP (alkaline phosphatase, placental) fusion protein, COS-7 cells were

transiently transfected with appropriate cDNA. Post-transfection cells were supplemented with

Opti-MEM (Gibco), and media collected after 48 hr. The media was concentrated through a 30K

ultracell filter (Merck Millipore, Tullagreen, Ireland). The amount of the AP fusion protein was

quantified by SDS-PAGE analysis and Coomassie staining. The amount of EphA4-AP was calibrated

by comparison to known amounts of EphA4-Fc fusion protein (R&D, Indinapolis, USA). AP activity

was measured using para-nitrophenylphosphate substrate (pNPP, Sigma).

Binding assays were carried in COS-7 cells, transiently transfected with the mouse or zebra

finch ephrin-B3. Following over-night (O/N) incubation, cells were incubated with varying amount

of EphA4-AP in binding buffer (Hanks buffer saline supplemented with 20 mM Hepes, PH 7.3,

0.05% BSA, 5 mM $CaCl_2$, 1 mM $MgCl_2$ and 10 µg/ml heparin sodium salt (Sigma) at room

temperature (RT) for 70 min. For visualization of AP staining, cells were plated on glass coverslip

coated with poly-lysine (0.005%, Sigma), and following AP-ligand binding as described above, cells

were washed with-binding buffer, fixed in 4% paraformaldehyde (PFA) in PBS for 15 min at RT, and

then washed with HBS solution (20 mM HEPES pH 7.3, 150 mM NaCl). Endogenous AP activity was

inactivated by incubation at 65 °C for 100 min. For color reaction, cells were incubated with

NBT/BCIP substrates (Roche, Basel, Switzerland) in AP buffer (100 mM Tris pH 9.5, 100 mM NaCl,

5 mM MgCl$_2$), O/N. After washing in HBS and fixation for 15 min in 4% PFA, and an additional

washing in HBS, coverslips were mounted under mounting medium (Thermo Scientific, USA).

For colorimetric assays, following AP-ligand binding and washing, cells were incubated O/N

at 4°C in lysis buffer (10 mM Tris, ph 8.0, with 1% Triton X100). The supernatant was subjected to

heat inactivation for 15 min at 65°C. Lysates were transferred to 96-well high protein absorbance

plate (Thermo Scientific, Nunc International USA) and the level of AP activity was determined

following addition of equal volume of 2 mg/ml pNPP substrate in reaction buffer (200 mM Tris, pH

9.5, 10 mM MgCl2, 200 mM NaCl). Plates were incubated at RT for 30 min to 1 hr and optical

density was determined at 405 nM using an ELISA plate reader.

Ephrin-B3 surface staining

COS-7 cells were transiently transfected with the mouse or zebra finch myc-EphrinB3 expression

vectors. 24 hr post-transfection, cells were incubated with mouse anti-myc (9E10) antibody in

phosphate buffered saline (PBS), washed with PBS, and subsequently incubated with Rhodamine

Red X (RRX) labeled anti-mouse antibody and washed again. Cells were fixed in 4%

paraformaldehyde/0.1 M phosphate buffer and mounted before imaging.

In-ovo electroporation

A DNA solution of 5 mg/mL was injected into the lumen of the neural tube at HH stage 17– 18

(E2.75-E3). Electroporation was performed using 3 × 50 ms pulses at 25–30 V, applied across the

embryo using a 0.5 mm tungsten wire and a BTX electroporator (ECM 830). Following

electroporation, 150–300 µL of antibiotic solution, containing 100 unit/mL penicillin in Hank's

Balanced Salt Solution (Biological Industry, Beit-Haemek) was added on top of the embryos.

Embryos were incubated for 3–19 days prior to further treatment or analysis. Roof plate-specific

expression was achieved utilizing the Wnt1 or DNAH2 enhancers, coupled with a dorsal electrode

positioning.

Immunohistochemistry and *in-situ* hybridization

Embryos were fixed overnight at 4°C in 4% paraformaldehyde/0.1 M phosphate buffer, washed

twice with phosphate buffered saline (PBS), incubated in 30% sucrose/PBS for 24 h, and

embedded in OCT (Scigen, Grandad, USA). Cryostat sections (20 µm) were collected on Superfrost

Plus slides and kept at −20°C. For thicker sections, spinal cords were isolated from the fixed

embryos and subsequently embedded in warm 5% agar (in PBS), and 100 µm sections (E12–E17)

were cut with a Vibratome. Sections were collected in wells (free-floating technique) and

processed for immunolabeling. The following primary antibodies were used: rabbit polyclonal anti-

GFP 1:1000 (Molecular Probes, Eugene, Oregon, USA), mouse anti-GFP 1:100, goat anti-GFP 1:300

(Abcam, Cambridge, UK), rabbit anti-RFP 1:1000 (Acris, Hiddenhausen, Germany), goat anti-ChAT

1:300 (Cemicon, Temecula, CA, USA), mouse anti-neurofilament 1:100 (3A10), mouse anti-myc

1:10 (9E10), rabbit anti-myc 1:100 (Abcam). The following secondary antibodies were used: Alexa

Fluor 488/647-AffiniPure Donkey anti-mouse, -rabbit and -goat (Jackson), and Rhodamin Red-X

Donkey anti-mouse and -rabbit (Jackson). Images were taken under a microscope (Eclipse Ni;

Nikon) with a digital camera (Zyla sCMOS; Andor) or a confocal microscope (FV1000; Olympus).

In situ hybridization was performed as described (Avraham et al., 2010). The following probes

were employed: chick ephrin-B1, ephrin-B2, EphA4 and Vglut2, zebra finch ephrin-B3 and mouse

ephrin-B3. Probes were PCR-amplified from brain cDNAs of E6 chick embryo, P4 zebra finch, and

E15 mouse, respectively, using the following primers. PCR products were verified by sequencing

prior to use.

Name	Sequence
cEFNB1-F	GAAAAAGGACCAGGCAGATGC
cEFNB1-R-T7	AGCTAATACGACTCACTATAGTTCTCTGTAGTCCGTAAGGG
cEFNB2-F	GAGGGGTGTGCCAGACAAAA
cEFNB2-R-T7	AGCTAATACGACTCACTATAGCACTGTCTGCAGTCCTTAGAG
cEPHA4-F	ACAGGCCCAATGGAGTCATC
cEPHA4-T7	AGCTAATACGACTCACTATAGGCATCCAAGGAGCCATTCTC
mEFNB3-F2	CCAGAACAGGAAGTCGCACT
mEFNB3-T7	AGCTAATACGACTCACTATAGCAAGACAAATTGGGCCTGCC
vGlut2-F	GGAAGATGGGAAGCCCATGG
vGlut2-T7	GAAGTCGGCAATTTGTCCCC
zfEFNB3_3'-F-T3	GCAATTAACCCTCACTAAAGGCACGCCCACAAAGGGAAGGA
zfEFNB3_3'-R-T7	AGCTAATACGACTCACTATAGTCTAATCAGCAGCAGCCGCG
zfEFNB3_5'-F-T3	GCAATTAACCCTCACTAAAGGTTACGTGCTGTACCCGCAGAT
zfEFNB3_5'-R-T7	AGCTAATACGACTCACTATAGCTCGTCTGATTGGCTGAACCT

PRV injection

We used two isogenic recombinants of an attenuated PRV strain (PRV Bartha) that express

enhanced GFP (PRV152) and monomeric red fluorescent protein (PRV614). The viruses were

harvested from Vero cell cultures at titers 4×10^8, 7×10^8 and 1×10^9 plaque forming units

(pfu/mL), respectively. Viral stocks were stored at −80°C. For pre-motoneuron infection, injections

of 3 µL of PRV152 or PRV614 were made into the thigh or wing musculature of E13 or E14 chick

embryos, using Hamilton syringes (Hamilton; Reno, NV, USA) equipped with a 33-gauge needle.

The embryos were incubated for 36–40 hr and sacrificed for analysis. For bilateral pre-

motoneurons infection, PRV152 and PRV614 were each injected into the pectoralis musculature

on opposite sides of the embryo midline.

Cell distribution and density plots construction

The codes for the cell distribution and density plots imaging and analysis were written in Matlab.

The distribution plots were constructed by calculating the relative midline expression, after

background subtraction and a subsequent length normalization between all analyzed sections. The

density plots were generated based on cross section images transformed to a standard form. The

background was subtracted, and the cells were filtered automatically based on their soma area or

using a manual approach. Subsequently, the neurons were depicted by semi-transparent circles or

their distribution was visualized by two-dimensional kernel density estimation, using the MATLAB

function "kde2d". Unless indicated otherwise, a contour plot was drawn for density values

between 20% and 80% of the estimated density range, in six contour lines. An additional script

was used for pre-motoneuron classification based on their neurotransmitter identity and laminar

location.

A similar density visualization procedure was done for the EphA4 density plots (Fig. 3E' and F'), but

was preceded by signal binarization and then an averaging filter (5 cycles) was used to achieve a

smoother density signal. The intensity of EphA4 signal as a function of distance from the midline

was obtained by summation of the binary signal over the horizontal axis, followed by a rescaling

step for each image. Statistical analysis was done after subdividing the distance around the dorsal

midline spanning from -15 to 15 (the relative distance between the central canal – 0, and the

white/gray matter boundary – 100 and -100). into 6 equal intervals, and t-test was performed for

the grouped data of each interval, originating from E10 and E16 data.

Statistical Analysis

Fig. 3I.
599

Distance from midline	-15 to -10	-10 to -5	-5 to -1	0 to 5	5 to 10	10 to 15
p-value (t-test)	1.3E-24	3.29E-44	4.38E-26	1.07E-98	1.1E-121	4.27E-26

601 Fig. 4G.

603 **Quantiles**

Level	Minimum	10%	25%	Median	75%	90%	Maximum
1_Mouse EphA4 Het (n=3)	0.1875	0.2	0.266667	0.296703	0.450321	0.535417	0.75
2_Mouse EphA4 KO(n=3)	1.304348	1.396667	1.508333	1.699095	1.857143	1.979861	2.25
3_Chicken (n=2)	1.76087	1.917544	2.06875	2.132456	2.385346	2.482353	2.578947
4_ZF (n=1)	1.333333	1.405848	1.491071	1.536585	1.604167	1.622017	1.681818

606 **Means and Std Deviations**

Level	Number	Mean	Std Dev	Std Err Mean	Lower 95%	Upper 95%
1_Mouse EphA4 Het (n=3)	36	0.3476214	0.1355198	0.0225866	0.3017681	0.3934747
2_Mouse EphA4 KO(n=3)	36	1.6895941	0.2226598	0.03711	1.6142568	1.7649313
3_Chicken (n=2)	22	2.1805534	0.2087605	0.0445079	2.0879941	2.2731126
4_ZF (n=1)	25	1.5334836	0.0831688	0.0166338	1.4991531	1.567814

609 **Means Comparisons**

610 **Comparisons with a control using Dunnett's Method**

611 Control Group = 1_Mouse EphA4 Het (n=3)

612 **Confidence Quantile**

|d|	Alpha
2.39255	0.05

615 **LSD Threshold Matrix**

Level	Abs(Dif)-LSD	p-Value
3_Chicken (n=2)	1.721	<.0001*
2_Mouse EphA4 KO(n=3)	1.244	<.0001*
4_ZF (n=1)	1.078	<.0001*
1_Mouse EphA4 Het (n=3)	-0.1	1.0000

618 Positive values show pairs of means that are significantly different.

620   <u>Fig. 5G</u>

621   **Quantiles**

Level	Minimum	10%	25%	Median	75%	90%	Maximum
Cervical	0.166667	0.251282	0.5	0.583333	0.75	0.9	1
Brachial	0.076923	0.214907	0.294118	0.384615	0.545455	0.822222	1.2
Thorax	0.304348	0.377778	0.470588	0.555556	0.727273	1	1.166667
Crural	0.125	0.131594	0.1875	0.315789	0.4	0.477778	0.571429
Sciatic	0	0	0	0	0	0	0

624   **Means and Std Deviations**

Level	Number	Mean	Std Dev	Std Err Mean	Lower 95%	Upper 95%
Cervical	15	0.6103012	0.2033028	0.0524926	0.4977158	0.7228865
Brachial	31	0.4487473	0.2349142	0.0421918	0.36258	0.5349145
Thorax	27	0.6096124	0.2135042	0.0410889	0.525153	0.6940719
Crural	23	0.311106	0.1260886	0.0262913	0.2565812	0.3656308
Sciatic	14	0	0	0	0	0

626
627
628

629 <u>Fig. 7E.</u>
630 **Quantiles**

Level	Minimum	10%	25%	Median	75%	90%	Maximum
GFP	8	9.9	13.25	16	20	23	29
mEFNB3	2	4	4	7	10	12	13
mEFNB3-S	3	4	5	7	9	11	13
zf-EFNB3	9	9	11	13	16	21	21

631
632 **Means and Std Deviations**

Level	Number	Mean	Std Dev	Std Err Mean	Lower 95%	Upper 95%
GFP	48	16.5208	4.72202	0.68156	15.150	17.892
mEFNB3	42	7.1667	3.21518	0.49611	6.165	8.169
mEFNB3-S	47	7.1277	2.66717	0.38905	6.345	7.911
zf-EFNB3	39	13.9744	3.72394	0.59631	12.767	15.182

633
634 **Means Comparisons**
635 **Comparisons with a control using Dunnett's Method**
636 Control Group =
637 GFP
638
639 **LSD Threshold Matrix**

Level	Abs(Dif)-LSD	p-Value
GFP	-1.78	1.0000
zf-EFNB3	0.663	0.0045*
mEFNB3	7.509	<.0001*
mEFNB3-S	7.601	<.0001*

640
641 Positive values show pairs of means that are significantly different.
642
643
644

<u>Fig. 8E-8H</u>

R squared values:

E. $R^2 = 0.086$

F. $R^2 = 0.9066$

G. $R^2 = 0.7469$

H. $R^2 = 0.7511$

References

1. T. G. Brown, On the nature of the fundamental activity of the nervous centres; together with
an analysis of the conditioning of rhythmic activity in progression, and a theory of the
evolution of function in the nervous system. *J Physiol* **48**, 18-46 (1914).

2. I. A. Rybak, N. A. Shevtsova, O. Kiehn, Modelling genetic reorganization in the mouse
spinal cord affecting left-right coordination during locomotion. *J Physiol* **591**, 5491-5508
(2013).

3. W. A. Alaynick, T. M. Jessell, S. L. Pfaff, SnapShot: spinal cord development. *Cell* **146**,
178-178 e171 (2011).

4. M. Goulding, Circuits controlling vertebrate locomotion: moving in a new direction. *Nature*
reviews. Neuroscience **10**, 507-518 (2009).

5. S. Grillner, A. El Manira, Current Principles of Motor Control, with Special Reference to
Vertebrate Locomotion. *Physiol Rev* **100**, 271-320 (2020).

6. S. Grillner, T. M. Jessell, Measured motion: searching for simplicity in spinal locomotor
networks. *Current opinion in neurobiology* **19**, 572-586 (2009).

7. A. Kania, R. Klein, Mechanisms of ephrin-Eph signalling in development, physiology and
disease. *Nat Rev Mol Cell Biol* **17**, 240-256 (2016).

8. L. Borgius *et al.*, Spinal glutamatergic neurons defined by EphA4 signaling are essential
components of normal locomotor circuits. *J Neurosci* **34**, 3841-3853 (2014).

9. K. Kullander *et al.*, Role of EphA4 and EphrinB3 in local neuronal circuits that control
walking. *Science* **299**, 1889-1892 (2003).

10. K. Kullander *et al.*, Ephrin-B3 is the midline barrier that prevents corticospinal tract axons
from recrossing, allowing for unilateral motor control. *Genes Dev* **15**, 877-888 (2001).

11. H. Wegmeyer *et al.*, EphA4-dependent axon guidance is mediated by the RacGAP alpha2-
chimaerin. *Neuron* **55**, 756-767 (2007).

12. N. Yokoyama *et al.*, Forward signaling mediated by ephrin-B3 prevents contralateral
corticospinal axons from recrossing the spinal cord midline. *Neuron* **29**, 85-97 (2001).

13. K. Katayama, J. R. Leslie, R. A. Lang, Y. Zheng, Y. Yoshida, Left-right locomotor circuitry
depends on RhoA-driven organization of the neuroepithelium in the developing spinal cord.
J Neurosci **32**, 10396-10407 (2012).

14. S. Skarlatou *et al.*, Afadin Signaling at the Spinal Neuroepithelium Regulates Central Canal
Formation and Gait Selection. *Cell Rep* **31**, 107741 (2020).

15. K. Padian, L. M. Chiappe, The origin of birds and their flight. *Sci Am* **278**, 38-47 (1998).

16. R. O. Prum *et al.*, A comprehensive phylogeny of birds (Aves) using targeted next-
generation DNA sequencing. *Nature* **526**, 569-573 (2015).

17. K. Straznicky, Function of Heterotopic Spinal Cord Segments Investigation in the Chick.
Acta Biol Acad Sci Hung **14**, 143-153 (1963).

18. C. H. Narayanan, V. Hamburger, Motility in chick embryos with substitution of lumbosacral
by brachial and brachial by lumbosacral spinal cord segments. *J Exp Zool* **178**, 415-431
(1971).

19. N. Rabe Bernhardt *et al.*, DCC mediated axon guidance of spinal interneurons is essential
for normal locomotor central pattern generator function. *Developmental biology* **366**, 279-
289 (2012).

20. N. Rabe, H. Gezelius, A. Vallstedt, F. Memic, K. Kullander, Netrin-1-dependent spinal
interneuron subtypes are required for the formation of left-right alternating locomotor
circuitry. *J Neurosci* **29**, 15642-15649 (2009).

21. T. E. Kennedy, T. Serafini, J. R. de la Torre, M. Tessier-Lavigne, Netrins are diffusible
chemotropic factors for commissural axons in the embryonic spinal cord. *Cell* **78**, 425-435
(1994).

22. T. Serafini *et al.*, The netrins define a family of axon outgrowth-promoting
proteinshomologous to C. elegans UNC-6. *Cell* **78**, 409-424 (1994).
23. K. M. Morris *et al.*, The quail genome: insights into social behaviour, seasonal biology and
infectious disease response. *BMC Biol* **18**, 14 (2020).
24. C. A. Cowan, M. Henkemeyer, The SH2/SH3 adaptor Grb4 transduces B-ephrin reverse
signals. *Nature* **413**, 174-179 (2001).
25. Q. Lu, E. E. Sun, R. S. Klein, J. G. Flanagan, Ephrin-B reverse signaling is mediated by a
novel PDZ-RGS protein and selectively inhibits G protein-coupled chemoattraction. *Cell*
105, 69-79 (2001).
26. H. L. Holen *et al.*, Ephrin-B3 binds to a sulfated cell-surface receptor. *Biochem J* **433**, 215-
223 (2011).
27. H. L. Holen *et al.*, Ephrin-B3 binds specifically to B lymphocytes in blood and induces
migration. *Scand J Immunol* **74**, 144-154 (2011).
28. J. P. Himanen, N. Saha, D. B. Nikolov, Cell-cell signaling via Eph receptors and ephrins.
Curr Opin Cell Biol **19**, 534-542 (2007).
29. C. N. Arnold *et al.*, ENU-induced phenovariance in mice: inferences from 587 mutations.
BMC Res Notes **5**, 577 (2012).
30. P. V. Lovell *et al.*, Conserved syntenic clusters of protein coding genes are missing in birds.
Genome Biol **15**, 565 (2014).
31. V. Hamburger, R. Oppenheim, Prehatching motility and hatching behavior in the chick. *J*
Exp Zool **166**, 171-203 (1967).
32. S. Paixao *et al.*, EphrinB3/EphA4-mediated guidance of ascending and descending spinal
tracts. *Neuron* **80**, 1407-1420 (2013).
33. L. D. De Gennaro, Origin of the avian glycogen body: I. Effects of tail bud removal in the
chick embryo. *Growth Dev Aging* **55**, 19-26 (1991).
34. L. D. De Gennaro, Origin of the avian glycogen body. II. Observations in support of a glial
nature in the chick embryo. *Growth Dev Aging* **57**, 275-281 (1993).
35. R. Necker, The structure and development of avian lumbosacral specializations of the
vertebral canal and the spinal cord with special reference to a possible function as a sense
organ of equilibrium. *Anat Embryol (Berl)* **210**, 59-74 (2005).
36. G. Fabbiani, M. I. Rehermann, C. Aldecosea, O. Trujillo-Cenoz, R. E. Russo, Emergence of
Serotonergic Neurons After Spinal Cord Injury in Turtles. *Front Neural Circuits* **12**, 20
(2018).
37. A. Kusuma, H. J. ten Donkelaar, Propriospinal fibers interconnecting the spinal
enlargements in some quadrupedal reptiles. *J Comp Neurol* **193**, 871-891 (1980).
38. R. F. Samara, S. N. Currie, Crossed commissural pathways in the spinal hindlimb
enlargement are not necessary for right left hindlimb alternation during turtle swimming. *J*
Neurophysiol **98**, 2223-2231 (2007).
39. J. D. Comer *et al.*, Sensory and spinal inhibitory dorsal midline crossing is independent of
Robo3. *Front Neural Circuits* **9**, 36 (2015).
40. E. N. Orlino, Jr., C. M. Wong, P. E. Phelps, L1 and GAD65 are expressed on dorsal
commissural axons in embryonic rat spinal cord. *Brain Res Dev Brain Res* **125**, 117-130
(2000).
41. G. Kardon, Muscle and tendon morphogenesis in the avian hind limb. *Development* **125**,
4019-4032 (1998).
42. C. Lance-Jones, L. Landmesser, Pathway selection by chick lumbosacral motoneurons
during normal development. *Proc R Soc Lond B Biol Sci* **214**, 1-18 (1981).
43. D. Satoh, C. Pudenz, S. Arber, Context-Dependent Gait Choice Elicited by EphA4 Mutation
in Lbx1 Spinal Interneurons. *Neuron* **89**, 1046-1058 (2016).

44. I. B. Hogue *et al.*, Fluorescent Protein Approaches in Alpha Herpesvirus Research. *Viruses*
7, 5933-5961 (2015).

45. Y. Hadas *et al.*, A 'tool box' for deciphering neuronal circuits in the developing chick spinal
cord. *Nucleic Acids Res* **42**, e148 (2014).

46. D. H. Rowitch *et al.*, Identification of an evolutionarily conserved 110 base-pair cis-acting
regulatory sequence that governs Wnt-1 expression in the murine neural plate. *Development*
125, 2735-2746 (1998).

47. L. S. Andersson *et al.*, Mutations in DMRT3 affect locomotion in horses and spinal circuit
function in mice. *Nature* **488**, 642-646 (2012).

48. V. B. Indjeian *et al.*, Evolving New Skeletal Traits by cis-Regulatory Changes in Bone
Morphogenetic Proteins. *Cell* **164**, 45-56 (2016).

49. E. Z. Kvon *et al.*, Progressive Loss of Function in a Limb Enhancer during Snake Evolution.
Cell **167**, 633-642 e611 (2016).

50. J. P. Himanen *et al.*, Crystal structure of an Eph receptor-ephrin complex. *Nature* **414**, 933-
938 (2001).

51. O. Pernet, Y. E. Wang, B. Lee, Henipavirus receptor usage and tropism. *Curr Top Microbiol*
Immunol **359**, 59-78 (2012).

52. O. A. Negrete *et al.*, Two key residues in ephrinB3 are critical for its use as an alternative
receptor for Nipah virus. *PLoS Pathog* **2**, e7 (2006).

53. C. E. Restrepo *et al.*, Change in the balance of excitatory and inhibitory midline fiber
crossing as an explanation for the hopping phenotype in EphA4 knockout mice. *Eur J*
Neurosci **34**, 1102-1112 (2011).

54. R. L. Watterson, Development of the glycogen body of the chick spinal cord; normal
morphogenesis, vasculogenesis and anatomical relationships. *J Morphol* **85**, 337-389
(1949).

55. R. L. Watterson, B. E. Spiroff, Development of the glycogen body of the chick spinal cord;
effects of unilateral and bilateral leg-bud extirpation. *Physiol Zool* **22**, 318-337, pl (1949).

56. J. Korlach *et al.*, De novo PacBio long-read and phased avian genome assemblies correct
and add to reference genes generated with intermediate and short reads. *Gigascience* **6**, 1-16
(2017).

57. A. Filosa *et al.*, Neuron-glia communication via EphA4/ephrin-A3 modulates LTP through
glial glutamate transport. *Nature neuroscience* **12**, 1285-1292 (2009).

58. Y. Lallemand, V. Luria, R. Haffner-Krausz, P. Lonai, Maternally expressed PGK-Cre
transgene as a tool for early and uniform activation of the Cre site-specific recombinase.
Transgenic Research **7**, 105-112 (1998).

59. M. G. Grabherr *et al.*, Full-length transcriptome assembly from RNA-Seq data without a
reference genome. *Nat Biotechnol* **29**, 644-652 (2011).

60. A. Kania, T. M. Jessell, Topographic motor projections in the limb imposed by LIM
homeodomain protein regulation of ephrin-A:EphA interactions. *Neuron* **38**, 581-596
(2003).

61. W. Halfter, Y. P. Yip, J. W. Yip, Axonin 1 is expressed primarily in subclasses of avian
sensory neurons during outgrowth. *Brain Res Dev Brain Res* **78**, 87-101 (1994).

62. F. Tie *et al.*, CBP-mediated acetylation of histone H3 lysine 27 antagonizes Drosophila
Polycomb silencing. *Development* **136**, 3131-3141 (2009).

63. W. A. Whyte *et al.*, Master transcription factors and mediator establish super-enhancers at
key cell identity genes. *Cell* **153**, 307-319 (2013).

Acknowledgments

The authors thank Tamir Lederman for technical assistance; Jerome Gros for providing the Pacbio

p-reads of the quail genome, Wes Warren for providing the Pacbio p-reads of the chicken genome,

and Anat Barnea for providing zebra finch hatchlings.

Funding:

This work was supported by a grant to AK and CVM from the US-Israel Binational Science

Foundation (grant No. 2017/172), an NSF graduate fellowship to SF, and grants to AK from the

Israel Science Foundation (grant No. 1400/16) and the Avraham and Ida Baruch endowment fund.

Author contributions:

AK, CVM, BH: research design, data analysis; BH: circuit analysis, in situ hybridization; OM: circuit

analysis; RSM: enhancer analysis; JE: protein studies; SF, PVL: genomics and gene structure

analysis; SP, RK: generated conditional ephrin-B3 knockout mouse; AK, CVM: wrote manuscript.

Competing interests:

The authors declare no competing of interest

Data and materials availability:

The data was deposited at any archive.

 Figures and Tables

 Figures legends

Figure 1: Spinal gait circuitry and EFNB3 genomics

A,B. A scheme illustrating the morphology and wiring of the spinal cord in wt mouse (A) and in

ephrin-B3/EphA4/ 2-chimaerin mice collectively referred to as *hop* mutants (B); pre-MN: pre-

motoneurons, RP: roof plate. **C.** Simplified cladogram depicting the relations across representative

organisms examined in this study. Branch lengths are arbitrary and not calibrated for time, red

means EFNB3 is absent in chicken. **D.** Comparison of EFNB3 (red) synteny across organisms where

synteny could be examined, with chromosomal location or data source indicated. EFNB3 is absent

in Galliformes (red square) and DNAH2 (blue) is absent in birds. Non-avian sauropsids: lizard and

alligator; wgs: whole-genome shotgun contigs; HB: hummingbird. **E.** Schematic structure of the

predicted mouse and avian ephrin-B3 proteins. RBD: receptor binding domain; TM:

transmembrane domain.

Figure. 2: Zebra finch ephrin-B3 protein and mRNA expression

A. The mouse and zebra finch proteins are presented on the cell membrane. Anti myc antibody

(9E10) was used for surface staining of COS-7 cells transiently transfected with amino-terminal

myc-tagged ephrin-B3 isoforms. The sequence of the full length zebra finch ephrin-B3 transcript

was reconstructed from RNA-seq reads using the Trinity platform (*59*) (Supp. Fig. S1). **B.** Binding

assay shows EphA4-AP binding to mouse ephrin-B3 protein but not to GFP or to zebra finch

ephrin-B3. **C.** Representative EphA4-AP/ephrin-B3 binding curves assessed by AP activity in a

colometric assay. Apparent KD value for mouse ephrin-B3 was 40.15 nM (n=3).**D.** *In situ*

hybridization for ephrin-B3 mRNA in P0 zebra finch spinal cord. High expression is evident in the

floor plate (FP), but declines along the roof plate (RP). **D'.** Normalized optical density (OD) profile

of zebra finch ephrin-B3 midline expression; shown is the means from 58 cross sections. **E.** *In situ*

hybridization for ephrin-B3 mRNA in E15.5 mouse spinal cord. High expression is evident in both

the FP and RP. **E'.** Normalized optical density (OD) profile of mouse ephrin-B3 midline expression;

shown is the means from 160 cross sections. **F,G.** Analysis of the activity of the ephrin-B3

enhancer element, at E9 (F) and E15 (G). The cherry reporter gene was cloned downstream to the

putative enhancer and electroporated into the chick spinal cord. GFP, under the control of the

ubiquitous CAGG enhancer-promoter, was co-electroporated. The reporter is detected in both FP

and RP (F,G), similar to the pattern of ephrin-B3 expression in mice, while GFP is expressed

ubiquitously.

Figure 3: Expression of ephrin-B1*, *ephrin-B2 and EphA4 mRNAs in the chick spinal cord.

In situ hybridization for ephrin-B1 (A,B)*, *ephrin-B2 (C,D) and EphA4 (E,F) mRNAs at the brachial

level in E10 (A,C,E) and E16 (B,D,F) cords. The floor plate (FP) is indicated by black arrows and the

roof plate (RP) by magenta arrows. **A,B**. Ephrin-B1 at E10 is high in the FP and low in the RP at E16

it is high in the FP and low in ventral RP. **C,D.** Ephrin-B2 at E10 is low in the FP and RP; at E16 it is

low in ventral RP. **E,F.** EphA4 expression at E10 is widespread in the gray matter, with higher levels

in scattered interneurons the lateral subdivision of the lateral motor columns (LMC_l) and the

medial motor columns (MMC). At E16, expression is detected in scattered interneurons and

motoneurons. Interneurons invading the dorsal midline are apparent (white arrow in inset in F).

Insets in E and F are enlargements of the areas boxed in dashed yellow. **E',F'.** Cell density

distribution, extracted from 10 cross sections from the regions flanking the dorsal midline at E10

and E16, respectively (indicated by white dashed boxes in E and F). **G.** Summation of cell *in situ*

hybridization relative intensity in the area indicated in E' (E10, blue) and F' (E16, red). The area

along the X was subdivided to 100 epochs. N=10 sections for each experiment. Values are

significantly different around the dorsal midline (spanning from -15 to 15) p<0.0001, t-test, see

Materials and methods and supplementary statistics).

Figure 4: Spinal cord morphology in *hop* mutants and chick show enlarged brachial RPs.

A-D, Dark field microphotographs of transverse sections of adult mouse spinal cord at lumbar and

brachial levels. Compared are control heterozygous (EphA4$^{lx/lx-}$; A,C) and homozygous null mice

(EphA4$^{lx/lx-}$/PGK-Cre$^+$; B,D) (*32*). Right insets (enlargements of the boxed areas) depict the central

canal and the RP and FP.

E,F. Transverse sections at the brachial (E) and lumbar (F) levels of E15 chick spinal cord. The white

matter is labelled with the anti-neurofilament antibody 3A10, the gray matter with nuclear DAPI

staining. Right inset (in E) is an enlargement of the boxed area.

G. Box plot representation of the roof plate (RP) to floor plate (FP) length ratios at the brachial

level in adult EphA4+/- mouse (N= 3, 36 cross sections), adult EphA4-/- mouse (N= 3, 36 sections),

E15 chick (N= 2, 22 sections) and P0 zebra finch (N= 1, 25 sections). The difference between the

heterozygous mouse, chick and zebra finch groups and the control homozygous null mouse group

is statistically significantly (See supplementary statistics). RP: roof plate; FP: floor plate.

Figure 5: Midline crossing at the brachial and lumbar levels of chick spinal cord

A. Dorsal midline crossing at brachial level in E15 chick demonstrated by neurofilament (3A10)

immunostaining in transverse sections. Detail views of three examples in insets, left inset is an

enlargement of the boxed area in A. Axons cross at the FP (yellow arrows) and the dorsal part of

the RP (white arrows). **B.** Dorsal midline crossing at brachial level in E17 chick; the boxed area is

enlarged in B' and B''. Anti-3A10 labels all axons, and anti-TAG1 labels sensory axons (*61*). **C.**

Dorsal midline crossing is evident following unilateral expression of a reporter gene. GFP under

the control of the CAGG enhancer/promoter construct was electroporated at E3 into the left side

of the spinal cord. At E15, midline-crossing axons are evident in the brachial FP (yellow arrows)

and RP (white arrows) . Detail views of three examples in insets, left inset is an enlargement of the

boxed area in C. **D,E.** Transverse sections at the sciatic plexus level at E15 (D) and E17 (E). Axons

cross only at the FP (yellow arrows). The glycogen body (GB) lacks axonal labelling. **F.** Transverse

section at the crural plexus level at E17. Right inset is an enlargement of the boxed area in F.

Axons cross at the FP (yellow arrow) and RP (white arrow). **G.** Box plot of the ratio between axons

crossing the RP and the FP at different axial levels in E15 chick. Numbers of midline-crossing axons

were scored at cervical (15 sections), brachial (31 sections), thoraxic (27 sections), lumbar crural

(23 sections) and lumbar sciatic (14 sections) levels of one embryo. At the sciatic level axons did

not cross the glycogen body. (See supplementary statistics). RP: roof plate; FP: floor plate.

Figure 6: Patterns of pre-MNs in the brachial and lumbar chick spinal cord

A-C. Cell distribution maps in the brachial and lumbar segments of E15 spinal cord sections of

chicks in which the hindlimb (A), distal wing (B), or pectoral (C) musculature were injected with

PRV-cherry at E14. Data are from 1038 ipre-MNs and 88 cpre-MNs in A (3 embryos), 620 ipre-MNs

and 210 cpre-MNs in B (4 embryos) and 734 ipre-MNs and 392 cpre-MNs in C (2 embryos). Ellipses

in dashed magenta represent the areas occupied by motoneurons. A'-C'. Images of PRV-positive

neurons supplemented with immunostaining for motoneurons (ChAT); right insets are

enlargements of the midline boxed area showing axonal decussation of pre-MNs at the FP (yellow

arrows) and RP (white arrows); white ellipses represent the central canal.

D. Identification of a subpopulation of pre-MNs that innervate pectoral muscle motoneurons

bilaterally. PRV-GFP and PRV-cherry were injected bilaterally at E14 into the pectoral muscles. Cell

distribution map of bilaterally innervating pectoral pre-MNs seen at E15 is shown in gray. D'.

Examples of bilaterally innervating pre-MNs; left panels: merged images; middle panels: PRV-

cherry; right panels: PRV-GFP. Out of 500 and 730 cpre-MNs on each side (18 sections), 11.9% ±

0.05 and 10.9% ± 0.04 were double labeled.

Figure 7: Expression of exogenous ephrin-B3 in the chick RP prevents midline crossing

A. Strategy for ectopic expression of the mouse ephrin-B3 at the roof plate. **B.** Cross-section from

brachial spinal cord of E17 embryo expressing ephrin-B3 (magenta) at the dorsal midline,

immunostained for neurofilament (cyan). **C,D.** Magnifications of 3 examples of dorsal midline

region of spinal cords electroporated with GFP (C,C',C") or ephrin-B3 (D,D',D") (both in magenta),

immunostained for neurofilament (cyan). The black and white images are the neurofilament

(3A10) staining. Some neurites cross the GFP-expressing RP (C) but not the ephrin-B3-expressing

RP. **E.** Quantification of dorsal midline crossing in spinal cords expressing GFP (N=43 sections, 3

embryos), mouse ephrin-B3 (N=71 sections, 3 embryos) or ephrin-B3-ΔC (N=59 sections, 3

embryos). The numbers of neurites crossing ephrin-B3- or ephrin-B3-ΔC-expressing RPs were

significantly lower than in GFP-expressing controls (See supplementary statistics).

Figure 8: Expression of exogenous ephrin-B3 in the chick RP prevents pre-MN crossing

A. Schematic of the experimental setup. GFP or ephrin-B3 were expressed in the midline at E3. At

E14, PRV-cherry was unilaterally injected into the wing musculature, and embryos fixed and

processed 38 hr later. ChAT-immunostained (cyan) motoneurons (MN)and PRV-labeled (magenta)

pre-MNs are indicated; dashed line represents cpre-MN axons that cross the RP. **B,C.** Transverse

sections of control (B) and Wnt1::ephrin-B3-expressing E15 embryos (3 examples in C,C'C"). The

black and white images are in B' and the insets in C,C',C" are PRV staining. In the control (B), cpre-

MNs somata are located ventral and dorsal to the central canal and neurites of cpre-MNs are

apparent at the FP (yellow arrows in B') and RP (white arrows in B'). In Wnt1::mEFNB3 embryos

(C, C', C''), the somata of pre-MNs are mainly located ventral to the central canal. Axons crossing

the FP (yellow arrows) and few short neurites (white arrowheads) crossing the dorsal midline are

shown in the right insets. **D.** Density maps of brachial cpre-MNs in control (GFP, N = 880 cells from

six embryos) and five ephrin-B3-expressing embryos (N = 242, 467, 366, 49 and 67 cells). The

fraction of sections containing ephrin-B3 expression along the entire dorsal midline is indicated at

the upper part of each spinal cord. A ventral shift of the distribution is apparent in the high

efficiency electroporated spinal cords of wnt1-B3-4 and wnt-1-B3-5. **E-H.** Plots demonstrating the

correlation between ventral commissural/ipsilateral, commissural/Ipsilateral, dorsal

commissural/ipsilateral and dorsal commissural/ ventral commissural pre-MNs ratios to the

electroporation efficiency (as described in D), respectively. **I,J.** Schematic illustration of the

circuitry at the lumbar (I) and brachial (J) avian spinal cord. We propose that dorsal midline

crossing is prevented at the lumbar level by the glycogen body, while dorsal midline crossing at

the brachial level is enabled due to the loss of ephrin-B3 activity. The brachial avian spinal cord

shares significant morphological and wiring similarities with mouse *hop* mutants.

Fig. 1

Figure 2

Fig. 3

E10 brachial	E16 brachial

Ephrin-B1

A

B

100μm

100μm

Ephrin-B2

C

D

100μm

100μm

EphA4

E

100μm

LMCi   MMC

100μm

F

100μm

100μm

E'

F'

gray/white
matter boundary

central
canal

gray/white
matter boundary

G

Relative intensity

E10
E16

0.8

0.6

0.4

0.2

-100   -15   0   15   100

Distance from the midline

Fig. 4

A
EphA4^lox/-
lumbar
500μm
250μm

B
PGK-Cre;EphA4^lox/-
lumbar
500μm
250μm

C
EphA4^lox/-
brachial
500μm
250μm

D
PGK-Cre;EphA4^lox/-
brachial
500μm
250μm

E
Chick E15
brachial
Lorem ipsum
100μm
100μm
3A10 DAPI

F
GB
Chick E15
lumbar
100μm
3A10
DAPI

G

RP/FP ratio

mouse
EphA4+/-
mouse
EphA4-/-
chick
zebra
finch

Fig. 5

Fig. 6

A

Wnt1 | Cre

CAG | PBase

BParms | CAG | STOP | ephrin-B3 | 6xmyc | BParms

B

EFNB3
3A10

200μm

GFP

C **C'** GFP 3A10 | 3A10 **C"** 200μm

ephrin-B3

D **D'** ephrin-B3 3A10 | 3A10 **D"** 200μm

E

# of axons crossing the RP

GFP ephrin-B3 ephrin-B3-ΔC

Fig. 8

964
965

Supplementary Materials for

Natural loss of function of ephrin-B3 shapes spinal flight circuitry in birds

Baruch Haimson[1], Oren Meir[1], Reut Sudakevitz-Merzbach[1], Gerard Elberg[1], Samantha Friedrich[3],

Peter V. Lovell[3], Sónia Paixão[2], Rüdiger Klein[2], Claudio V. Mello[3*] and Avihu Klar[1*]

* Avihu Klar. Email: avihu@mail.huji.ac.il
* Claudio V. Mello. Email: melloc@ohsu.edu

This PDF file includes:

      Supplementary Text
      Figs. S1 to S6
      Tables S1
      References (59,62,63)

989 Fig. S1. Ephrin-B3 gene and transcripts in zebra finch

C

```
GGCGGCGGCGGCGGCGGCGGCGGCGGCGGCGGAGCGGGGGGCGGCCGGGGCCCACCGGGAGGGGCCCAGAGCGGC
CGGGGGGCGCCGGGGCCCGGCGGGGCGGGGGGCGATGGCGGGGCGGGCGGCGGCGCTGCTGGTGGCCGTGCTGGG
CGCGGGGGCCGCGGGGCTGAGCCTGGAGCCGGTGGTCTGGCACACGGGGAACCGGCGGTTCCTGGAGGCCGGCGG
TTACGTGCTGTACCCGCAGATCGGTGACCGCCTGGACCTGGTGTGCCCGGGGGGCGCGGGGTACGAGTACTACAA
GCTGTACCTGGTGGGGGGGGGCGCAGGCGCGGCGCTGCCAGGTGCCCCCCGCGCCCACCCTGCTGCTGACCTGCGA
CCGCCCCCAGCGCGACGTGCGCTTCACCATCAAGTTCCAGGAGTTCAGCCCCAACCTCTGGGGCCACGAGTTCCG
CCGCCAGCACGACTATTACATCATCACCACCTCGGACGGGACCCCCGAGGGGCTGGAGAACCGCCAGGGCGGGGC
CTGCCTGACCCGGGCCATGCGGGTCACCCTGCCGCTGGGGCAGCGCCCCGGGGCGGAGACACCGCCGGAGCAGGA
TGGGACAGGTTCAGCCAATCAGACGAGCCCGCCCGGGCCGGGCCCGCCCCCCGCGCGGGGAGGGGCGGAGCTGTG
GGGCGCGGGGGGAGGGGCGGCGCTGCTGGCGCTGGGGGGGGCGGGGGGGGCGCTCTGGTGGCGGCGCCGCCCCCG
GAGACCCCCAAAAACTCCACGGGGTGGGGGAGGGGCGGGGCCAACCCGGCCACGCCCCCACGGGCCCGGCCCCGC
CCACCCCCTGCGCGAGCTGCCCCTCCCCCCCGCCGCCCCTCCCCCCCACCTACTACAAGGTGTGAGGGGTGGGGGA
GGGGCGGGACACCCAAACTGGGACGGGCCACGCCCACAAAGGGAAGGACACGCCCCCACCCTGAAGCTCCGCCCCC
TCGGGGCAAAGGGGAATGACGGACCCAAAAAGGGAAGAATTTGCCCCAAATTGACCCCCCAGGTCTGGATGGGTG
TGGCTTAACCGGGAGGCGTGGCCTGAATTGGGCGTGGTCACTAAAGGGGTGTGGCTGTGATGTCATATGGTGCCG
CCCAGTGGGCGTGGTCAAATATGGGCGTGGTCTCCATAAATGGGCGTGGCTTTACCTGGGGAGGCGGAGCTTAAA
TGGCTTTGAGGCGTCACCCCTGGGAAAGGGGCGGGGCTCGCTCACGGCTCCCTCCCATTGGTCACGATGAAGCCC
CGCCCACACAAAGGGGGCGGAGCTTTCCCGGCGCAGTAACATGGCCGCCGCCACAGCGGATGAACAGGAAGTGAC
GTCACGGCGACAGAGAGGCGGGACTTCCTGTGTGATGTCGCGTGGGGGCGGGGCCTGCGCGCGCGGCTGCTGCTG
ATTAGACAGTTTTGTGCAAAAAAAAAAAAAAAAA
```

990
A. Zebra finch ephrin-B3 gene, predicted transcript, predicted protein, and transcript variants. The
two EST clones that include coding exons are DV945855 and DV950988. Both include frameshifts
that result in premature *termination (indicated by asterisks)*.

B. RNAseq reads of Ephrin-B3 in zebra finch. RNAseq data in zebra finches provide expression

support for this structure and for expression of all five exons, and is consistent with the predicted

polyA site based on multiple ESTs.

C. The sequence of the full length zebra finch ephrin-B3 transcript was reconstructed from RNA-

seq reads using the Trinity platform *(59)*. The coding exons are indicated in alternating blue and

yellow colors. The 5'UTR (black) is included in the 1st exon and the 3'UTR (black) is included in the

5th exon.

.

Fig. S2. Phylogeny of the ephrin-B3 protein

1003
Sequence alignment of the predicted ephrin-B3 proteins from different species. The T-Coffee
algorithm was used for alignment of multiple sequences. Avian-specific deletions were detected in
the heparin binding, tetramerization interface and SH2/3-binding domains (indicated by blue -
with asterisks, yellow and gray rectangles, respectively), and specific mutations observed in the
dimerization interface domain (indicated by red rectangles and horizontal bars in residues 108-111
of mouse). The open box indicates the transmembrane domain.
1010

1011   Fig. S3. A putative EFNB3 enhancer in mouse DNAH2

1012
1013

A display of ENCODE analysis of candidate cis-Regulatory elements in mouse chr11:69,546,195-
69,547,544; shown is the region that includes EFNB3 (exons 3-5) and DNAH2 (exons 1-2) genes on
the minus strand. A DNaseI hypersensitive (HS) site is in the 1st intron of DNAH2
(chr11:69,546,195-69,547,544 in GRCm38/mm10). This motif binds modified histones (H3K4me3,
H3K4me1 and H3K27ac) that are associated with active enhancers; histone H3K27me3 is
associated with repression of developmental genes, as well as CTCF – a nuclear factor associated
with active enhancers (61,62). Binding of these proteins is apparent in the mouse embryonic
cerebellum and brain. The data was obtained from UCSC genome browser
(https://genome.ucsc.edu/index.html).
1023

1024   Fig. S4. RP/FP ratio along the longitudinal axis of wt mice

1025

1026
1027

1028   The ratio between the roof plate and floor plate lengths at different segmental levels. The limb-

1029   innervating segments exhibit a lower RP/FP ratio compared to the thoracic and sacral levels.

1030   Analysis was done on images from the Mouse spinal cord atlas (Allen Institute for Brain Science; N

1031   = 17 mice, 15-16 section per mouse). Box plot representation of the roof plate (RP) to floor plate

1032   (FP) length ratios at the cervical, thoracic, lumbar and sacral levels at P4.

1033
1034
1035

1036 **Fig. S5. Distribution of pre-MNs at the spinal crural plexus level**

1037

1038

1039 pre-MN map at the crural lumbar level; shown are 361 ipre-MNs and 192 cpre-MNs, respectively (N=3 embryos).

1040

1041 Fig. S6. Brachial pre-MN neurotransmitter identity

1042

1043 Experimental design for brachial pre-MN labeling and *in situ* hybridization for VGlut2. Density
1044 maps of ipre-MNs and cpre-MNs are shown.

1045
1046

1047 Table S1. The ratio of excitatory pre-MNs in the lumbar and brachial levels of the chick spinal cord.

1048

level	side	% vGlut2	#
Lumbar sciatic	ipsi	46.46	145
	comm	38.62	145
brachial	ipsi	35	274
	Ventral comm	38	29
	Dorsal comm	47	51

1049
1050

Supplementary References

59   Grabherr, M. G. *et al.* Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat Biotechnol* **29**, 644-652, doi:10.1038/nbt.1883 (2011).

62   Tie, F. *et al.* CBP-mediated acetylation of histone H3 lysine 27 antagonizes Drosophila Polycomb silencing. *Development* **136**, 3131-3141, doi:10.1242/dev.037127 (2009).

63   Whyte, W. A. *et al.* Master transcription factors and mediator establish super-enhancers at key cell identity genes. *Cell* **153**, 307-319, doi:10.1016/j.cell.2013.03.035 (2013).