Acidovorax pan-genome reveals specific functional traits for plant beneficial and pathogenic

plant-associations

Roberto Siani[1,2], Georg Stable[3], Caroline Gutjahr[3], Michael Schloter[1,2], Viviane Radl[1*]

[1] Helmholtz Center for Environmental Health, Institute for Comparative Microbiome Analysis;

Ingolstaedter Landstr. 1; 85758 Oberschleissheim, Germany

[2] Technical University of Munich, School of Life Sciences, Chair for Soil Science; Emil-Ramann-

Strasse 2; 85354 Freising, Germany

[3] Technical University of Munich, School of Life Sciences, Chair for Plant Genetics; Emil-

Ramann-Strasse 2; 85354 Freising, Germany

*Corresponding author:

viviane.radl@helmholtz-muenchen.de

phone +49 89 3187 2903

Keywords:

Acidovorax, plant growth promotion, plant pathogens, *Lotus japonicus*, comparative genome

analysis

Abstract

Beta-proteobacteria belonging to the genus *Acidovorax* have been described from various

environments. Many strains can interact with a range of hosts, including humans and plants,

forming neutral, beneficial or detrimental associations. In the frame of this study we investigated

genomic properties of 52 bacterial strains of the genus *Acidovorax*, isolated from healthy roots

of *Lotus japonicus,* with the intent of identifying traits important for effective plant-growth

promotion. Based on single-strain inoculation bioassays with *L. japonicus*, performed in a

gnotobiotic system, we could distinguish 7 robust plant-growth promoting strains from strains

with no significant effects on plant-growth. We could show that the genomes of the two groups

differed prominently in protein families linked to sensing and transport of organic acids,

production of phytohormones, as well as resistance and production of compounds with

antimicrobial properties. In a second step, we compared the genomes of the tested isolates with

those of plant pathogens and free-living strains of the genus *Acidovorax* sourced from public

repositories. Our pan-genomics comparison revealed features correlated with commensal and

pathogenic lifestyle. We could show that commensals and pathogens differ mostly in their ability

to use plant-derived lipids and in the type of secretion-systems being present. Most free-living

Acidovorax strains did not harbour any secretion-systems. Overall, our data indicated that

Acidovorax strains undergo extensive adaptations to their particular lifestyle by horizontal

uptake of novel genetic information and loss of unnecessary genes.

Introduction

Several soil-derived bacteria, which can colonize plant roots, directly or indirectly promote plants' health by facilitating nutrient mobilization and uptake [1], altering plant hormone levels hormones [2] and competing with pathogens [3, 4]. At the same time, soils also harbour a large repertoire of plant pathogens. It is not entirely clear what distinguishes non-harmful root-associated commensals from pathogens, as even closely related strains have been proven to promote plant-growth or, contrariwise, to negatively affect it [5]. Several studies [6, 7] link this to loss or acquisition of genomic features [e.g., virulence factor, pathogenicity islands], resulting from mutations and horizontal gene transfer [8]. Our existing knowledge in this field is still scarce and inconsistent, as in most cases only single strains were compared, which does not allow to understand evolutionary patterns and to define a generalized model.

Bacteria of the genus *Acidovorax* belong to the group of Gram-negative beta-proteobacteria [9] and form associations with a wide range of monocotyledonous and dicotyledonous plants [10]. *Acidovorax* has been mostly considered as biotrophic pathogen [11]. However, bacteria of this genus are also known as commensal species or plant beneficial bacteria, which produce secondary metabolites and hormones promoting plant growth, as well as competing with pathogens [12]. Most common species are *A. delafieldii* and *A. facilis*, whilst the most studied pathogenic species is *A. avenae*, the agent of corn (*Zea mays)*, sugar cane (*Saccharum officinarum)* and rice (*Oryza sativa)* leaf blight, orchids brown spot disease and watermelon (*Citrullus lanatus)* fruit blotch. Thus, *Acidovorax* is a valuable model organism to investigate evolutionary patterns of plant-associated bacteria and to study which genes differentiate bacteria acting as biotrophic pathogens, as commensals or plant growth-promoting bacteria [13].

In the frame of this study, we compared the genomes of isolates classified as *Acidovorax* which were obtained from roots of healthy *Lotus japonicus* ecotype Gifu plants. We tested the strains for their ability to promote plant growth of *L. japonicus* in sterilized sand and correlated differences in the genomes of the strains to the observed effects on plant growth. For a broader comparative analysis, we included additional genomes, available in public databases, belonging

to strains from 15 different species of *Acidovorax* from different environments, including all

major groups of plant pathogens from this genus. We studied the diversity in the pan-genome

and critically evaluated enriched gene functions, secondary metabolites biosynthetic clusters in

the different functional groups of *Acidovorax*.

Material and methods

Origin of strains and genomes

Fifty-two strains of endophytic *Acidovorax* were isolated from the root systems of healthy *Lotus*

japonicus ecotype Gifu B-129 (subsequently called *L. japonicus)* grown in natural soil in

Cologne, Germany. DNA from all strains was extracted and sequenced using an Illumina HiSeq

2500 device (Illumina, USA). The isolation and sequencing procedure is described in detail

elsewhere [14]. Genomes assemblies are available in the European Nucleotide Archive under

the accession number PRJEB37696.

In addition, 54 other genomes where obtained from NCBI. The collection includes 15 species:

A. anthurii, *A. avenae*, *A. carolinensis*, *A. cattleyae*, *A. citrulli*, *A. delafieldii*, *A. defluvii*, *A.*

ebreus, *A. facilis*, *A. kalamii*, *A. konjaci*, *A. monticola*, *A. oryzae*, *A. radicis, A. varianellae* and

A. spp. According to the database, all the selected bacterial genomes originate from plant, soil

and water samples. Details on the selected genomes are summarized in Supplemental Material

1. The mean level of completion for all 106 genomes was calculated to 0.986.

Based on the available metadata, we assigned the genomes to 3 behavioural phenotypes: 62

genomes belong to commensal or beneficial plant-associated strains (thereby termed

"commensals"), 21 to plant pathogenic strains (thereby termed "pathogens") and 23 to free-

living strains (thereby termed "free-living"). According to the results of our *in planta* bioassays

(see below), we further classified the *Lotus*-isolated strains by their ability to promote *Lotus*

japonicus growth.

Linear discriminant analysis effect sizes

We studied genomic features across the strains isolated from *L. japonicus* and correlated the

obtained data with the effects detected by the *in planta* bioassays (see below). Therefore, we

annotated the genomes using Anvi'o 6.2 microbial multi-omics platform [15] which computes *k*-

mer frequencies and identifies coding sequences using Prodigal [16]. Afterwards, HMMER3 [17]

was used to profile the genomes with hidden Markov models and to find homologous protein

families in the Pfam database v33.1 [18]. Using the features occurrence frequency matrix

generated by Anvi'o 6.2, we calculated the effect sizes of the features over a linear discriminant

analysis (LEfSe [19]) between the groups of interest. We chose the default alpha values of 0.05

for the factorial Kruskal-Wallis test (KW) and the pairwise Wilcoxon test (W) and a threshold of

2.0 for the Linear Discriminant Analysis (LDA) logarithmic scores.

Pan-genome analysis

We used Anvi'o 6.2 further to pre-process the genomes and reconstruct *Acidovorax* pan-

genome [20]. We inferred the pan-genome with the following options: DIAMOND in sensitive

mode to calculate amino-acids sequences similarity, Minbit parameter to 0.5, minimum

occurrence of gene clusters to 2, MCL inflation parameter to 7. We separated gene clusters into

occurrence frequency classes. Kernel gene clusters were defined as present in 106 genomes.

Core gene clusters were defined as present in 100 to 105 genomes. Shell gene clusters were

defined as present in 16 to 99 genomes. Cloud gene clusters were defined as present in 2 to 15

genomes. Singleton gene clusters were defined as present in only 1 genome and were

excluded from the downstream processing to reduce computational load. We calculated

functional and geometrical homogeneity for each gene cluster and genome average nucleotide

identity using pyANI [21]. We calculated a functions occurrence frequency matrix and functions

enrichment scores for gene clusters' functions by assigning each gene cluster to the closest

Pfam.

Comparative genomics

We performed all downstream analysis in R (v4.02, code available on https://github.com/rsiani/

AVX_PGC). We calculated total length, GC%, number of genes, gene clusters and singleton

gene clusters, average gene length and number of genes per kb (gene density) for each of the

genomes. We assessed the significance of differences by Pairwise Student's t-test with Holm-

Bonferroni's correction for multiple testing ($p < 0.05$).

We converted the function frequency occurrence matrix (3036 Pfams) to a binary

presence/absence matrix and filtered zero variance (757 Pfams), near zero variance (1668

Pfams) and correlated variables (0 Pfams) to remove redundancy (resulting in 1368 Pfams). We

performed a Principal Component Analysis (PCA, R package: "FactoMineR", v2.3) to ordinate

the strains based on their functional similarity and tested the significance of the clustering by

PERMANOVA (package: "vegan", v2.5-7). We isolated the features responsible for most of the

variance explained by extracting the highest contributors for each of the first 3 components.

From the enriched feature profiles predicted by Anvi'o 6,2, we manually curated a list of

features putatively correlated with plant-association. Enrichment score and adjusted

significance value were used to assess the relevance of the features across the groups.

We retrieved secondary metabolites biosynthesis gene clusters using AntiSMASH 5.0 [22]. We

launched a local instance of the program with no extra features at a "relaxed" strictness level,

running only the core detection modules. At this strictness level, well-defined clusters and

partial clusters are detected and annotated against the antiSMASH database.

We built a classification model using neural networks with feature extraction (R package "caret",

v6.0-86), a model employing a preliminary feature extraction step to reduce redundancy, while

preserving information and decreasing computational load [23]. We trained the model over 90

% of the strains with "leave-one-out" cross-validation over 1000 repeats, using accuracy scores

to select the optimal parameters. The fit of the model was evaluated by predicting labels for the

remaining 10 % of the strains and again against the whole dataset Results were assessed by

calculating confusions matrix for both predictions.

In planta bioassays

We prepared seeds of *L. japonicus* by sandpaper scarification and surface sterilization with a

10% DanKlorix Original (CP GABA GmbH, Germany) and 0.1% sodium dodecyl sulfate (SDS)

solution. We germinated the seeds on a 0,8 % water-agar plate for 3 days at 22 °C in the dark

followed by 4 days at 22 °C in the light at 210 µM/m²s intensity (growth cabinet PK 520-LED,

Polyklima).

We transferred plantlets to pots (Göttinger 7 x 7 x 8 cm, Hermann Meyer KG, Germany) filled

with washed and autoclave sterilized quartz-sand (Casafino Quarzsand, fire-dried, 0,7 − 1,2

mm, BayWa AG, Germany) at a density of 5 plants per pot. Pots were supplied with a thin layer

of synthetic cotton at the bottom to prevent the sand running out through the holes.

44 of the *Acidovorax* strains isolated from healthy *L. japonicus* plants were individually pre-

grown in 50 % TSB media at 30°C overnight and diluted to an OD_{600} of 0,001 using 1/3 BNS-

AM medium (Basal Nutrient Solution [24] modified for arbuscular mycorrhiza (0.025 mM

$KH_2PO_{4)}$). 25 ml of this bacterial suspension was applied to the pots. Finally, 20 ml of the sterile

1/3 BNS-AM-bacterial solution was added. Plants were grown at 60 % air humidity and light

intensity of 150 µM/m²s. The photo-period was set to 16 h light/ 8 h dark long day with a

temperature cycle of 24/22 °C. For each bacterial strain tested, four replicate pots were

prepared.

After 9 weeks of growth, we harvested the plants and determined root and shoot length, wet

weight and dry weight after freeze drying (Alpha 1-2, Martin Christ Gefriertrocknungsanlagen

GmbH, Germany). Statistical significance was assessed by Pairwise Student's *t*-test adjusted

for multiple comparisons using the Holm-Bonferroni method ($p < 0.05$).

Results

Acidovorax strains isolated from healthy L. japonicus roots diversely affect their host

We compared growth metrics of *L. japonicus* inoculated with different strains to evaluate the

outcomes of the plant-association. Overall, 34 out of 44 tested strains showed significant effects

($p < 0.05$; Figure 1) when considering all the collected measurements of plants' growth (n = 25).

Twenty-one strains displayed a positive effect on at least one metric. Nine strains displayed a

negative effect on plant growth, markedly on fresh weight measurements. Four strains

displayed contrasting effects on different metrics and 10 had no significant effect on growth.

From the 21 strains that resulted in improved plant-growth, 7 strains displayed a more

consistent effect across all the replicates on at least one of the considered metrics ($p < 0.001$).

We selected these robust growth-promoters for a follow-up genomics comparison.

Nineteen Pfams discriminate the robust plant growth-promoting *Acidovorax* strains

To isolate genomic differences correlated with the outcomes of the *in planta* bioassays, we

calculated linear discriminant analysis effect sizes on the function occurrence frequency table

and identified 19 Pfams discriminating the 7 robust growth-promoters from the remainder of the

Lotus-isolated strains (KW $p < 0.05$, W $p < 0.05$, logarithmic LDA score > 2.0, Figure 2; Table

1). Out of those, 15 Pfams were enriched in the genomes of robust growth-promoters, related to

4 broad functional categories: chemotaxis by sensing and uptake of organic acids and sugars

(tripartite tricarboxylate transporters, TTTs, and periplasmic binding proteins, PBPs),

colonization through synthesis and detoxification of plant secondary metabolites (aldolase, aldo-

keto reductase, rhodanase, amidase and mannosyltransferase), competition by synthesis,

secretion and detoxification of antimicrobial compounds (beta-lactamase, polyketide cyclase,

tetrapyrrole corrin-porphyrin methlyase, dynein-related domain and dienelactone hydrolase) and

transcriptional regulation (FCD domain, IclR domain and sigma 70 factor). We found 4 Pfams

which were less abundant in the genomes of robust growth-promoters, also implicated in

chemotaxis (Tripartite ATP-independent periplasmic transporters, TRAPs), colonization (FliO

and the FG-GAP repeat) and regulation (metallo-peptidase M90).

Acidovorax has an open pan-genome

We reconstructed the *Acidovorax* pan-genome from 106 genomes (Figure 3), representing 15

out of the 23 classified *Acidovorax* species [51], along with several yet unclassified strains, and

capturing a large share of the genomic variability in the genus. The pan-genome contains a total

of 523555 genes grouped in 34146 gene clusters, of which 2 % belonged to the kernel (present

in all of the genomes), 3% belonged to the core (present in 100 to 105 genomes), 13%

belonged to the shell (present in 16 to 99 of the genomes), 45 % belonged to the cloud (present

in 2 to 15 genomes) and 37 % were singleton (present in one genome only). Ninety-five percent

of the gene clusters were classified as accessory, while 5 % are shared by all or most of the

genomes. By fitting our results to a Heaps' Power-law regression model [52] we obtained a ɣ <

1 of 0.48, which defines *Acidovorax* pan-genome as "open", with positive rates of novel genes

discovery for every additional genome included.

Pathogenic *Acidovorax* strains have longer genomes but less genes

We calculated total length, GC%, number of genes, gene clusters and singleton gene clusters,

average gene length and number of genes per kb (gene density) for each of the genomes

(Figure 4a, Supplemental Material 2). On average pathogens had longer genomes (5.38 mb,

SD = 0.27) than commensals (5.21 mb, SD = 1.03, $p < 0.01$) and free-living (4.72 mb, SD =

0.53, $p < 0.0001$, Figure 4b). However, when testing for differences in gene density, pathogens

had less genes per kb (0.86, SD = 0.09) than both commensals (1.01, SD = 0.05, $p < 0.0001$)

and free-living (0.94, SD = 0.03, $p < 0.001$, Figure 4c).

Strains separate according to behavioural phenotype

We performed a PCA on the function presence/absence matrix and found that the first 3

dimensions explained 47 % of the variance in the matrix (24.3 %, 12.4 % and 10.3 %). The

individual strains clustered according to their observed behavioural phenotype ($p < 0.05$), with

pathogens grouping in a neatly separated cluster and a partial overlap between free-living and

commensal strains (Figure 5ab). Notably, no clustering was visible when considering the

original habitat of the strains. We extracted the highest contributing variables for the first 3

components (Figure 5c, Table 2).

This approach retrieved a total of 14 Pfams displaying a strong variance across the groups.

Two of these, the C-terminal domain of a secreted lipases and the mutagenesis inducer HIM1,

contributed the most to the first component and were found in all of the commensals, 35 % of

the free-living and none of the pathogens. A third Pfam, characterized by a WG repeat motif,

was only retrieved in 40 % of the commensals and accounted for a high proportion of the

second component explanatory power. Finally, the highest contributor to the third component

was a putative genomic island, encoding 11 elements of the type VI secretion-system, present

in 95 % of the pathogens, 61 % of the commensals and only 13 % of the free-living strains.

Pathogenic and commensal *Acidovorax* strains have a distinctive set of features

We performed a function enrichment analysis to isolate features unique or more represented in

the different groups (Supplemental Material 3). 1243 out of 3036 Pfams were significantly

enriched in 1 or 2 groups (adjusted $q < 0.05$). We further investigated enriched functions from

plant-associated strains only: 371 Pfams strongly associated with commensals ($q < 0.05$) and

303 Pfams strongly associated with pathogens ($q < 0.05$) (Figure 6a). From these, we manually

curated a list of 54 features (Table 3) putatively involved in plant-microbe and microbe-microbe

interactions (Figure 6b). We assigned them to 4 families based on known performed function:

22 effectors, 13 hydrolytic enzymes , 12 motility functions and 7 secondary metabolites .

Eighteen effectors and known virulence factors were enriched in pathogens, among which,

notably, members of the HrpB gene of the type III secretion system, and its regulatory element

HpaP. Four effectors were also identified in commensals, including a type II secretion-system

protein.

Among the hydrolytic enzymes, 9 different Pfams were found to be enriched in pathogens, all

related to plant and fungal polysaccharides degradation, including components of pectate

lyases, amylases, chitinases, cellulases and glucanases. Four hydrolytic enzymes Pfams were

enriched in commensals, but mainly responsible for the degradation of complex bioactive

compounds, such as flavonoids, chlorophyll and aromatic-ring backbones.

Interestingly, 10 Pfams related to diverse motility structures, namely flagella and pili, and

chemotaxis were predominantly found in commensals. For comparison, just 2 motility related

Pfams were enriched in pathogens.

Among those enriched in commensals, we also found 6 Pfams related to bacterial cellulose

biosynthesis and 1 polyketide-synthetase Pfams enriched in pathogens.

Using AntiSMASH 5.0, we retrieved a repertoire of 15 biosynthesis gene clusters (BGCs)

among the bacterial genomes (Figure 7). The most commonly retrieved BGCs were those for

bacteriocin and terpene production. Interestingly, phosphonate and all non-ribosomal peptide-

synthetase BGCs (NRPS, NRPS-like, NRPS-T1PKS) were almost unique to pathogens and

retrieved in more than half of the pathogenic strains.

Genotypes can predict the behaviour of the strains

We tested whether a model could predict a strain behavioural phenotype from its gene

repertoire. We trained a neural network with feature extraction classifier on the gene function's

presence/absence profiles of 90% of our strains (n = 96). The optimal model, which we selected

according to accuracy value, registered a mean balanced accuracy = 0.93 and an area under

the curve (AUC) = 0.97, with the selected final values of size = 3 and decay = 0.1. We tested

the predictive power of the model against the remainder 10% of the strains (n = 10) and

registered accuracy = 1, sensitivity = 1, specificity = 1, kappa = 1, 95% confidence interval =

0.69 to 1 and p = 0.006. When testing against the whole collection (n = 106), all metrics

remained unchanged but for the 95% confidence interval = 0.97 to 1 and p = 2.2e-16.

Discussion

Robust growth-promoting strains share traits which allow a better use of phytochemicals

Our *in planta* bioassays showed a wide range of effects associated with the presence of diverse

Acidovorax isolates obtained from healthy roots of *L. japonicus*. We compared the genomes of

7 robust growth-promoters against the remainder of the isolates and retrieved 19 discriminant

Pfams related to different aspects of plant-microbe and microbe-microbe interactions:

chemotaxis towards root exudates, metabolism of plant secondary metabolites, antagonistic

competition and transcriptional regulation.

Organic acids are a major component of root exudates [110]. Microbial transporters, such as

ABC-type transport systems, TRAPs and TTTs, are essential to make use of these carbon rich

compounds and a number of studies confirmed that their presence is correlated to improved

colonization in both plant-growth promoters [111] and pathogens [32]. Many transporters are

highly specific for individual substrates. For example, TTTs, which are enriched in robust-growth

promoters in our study, show a higher affinity towards citrate compared to TRAPs. As shown by

metabolic profiling [112], citrate is more abundant than other organic acids in *L. japonicus* root

tissue and improved sensing and uptake of citrate could translate in an evolutionary advantage

of microbiota towards the colonization of this plant. Our robust growth-promoters also shared an

enrichment of several features related to the metabolism of various plant-derived compounds,

such as benzoic acids, carbonyls, cyanide and rare sugars. Some of these traits have been

already correlated to improved plant-association in taxonomically diverse bacteria [13, 113,

114]. Furthermore, we observed the enrichment of an amidase involved in the biosynthesis of

indole-3-acetic acid and of a known regulator of GABA uptake [115]. The role of auxin in plant-

growth promotion by bacteria has been thoroughly characterized [116] and GABA has been

more recently discovered to act as signal from plants to their associated microbiota [117, 118].

Finally, we found enriched components of diverse regulatory pathways and of both antibiotic

synthesis and degradation, which could help the robust growth-promoters to not only better

colonize the plant by degrading plant-derived phytochemicals with antimicrobial properties, but

also to gain competitive advantage over other microbial populations.

Overall, we found evidence of genomic differences among robust growth-promoting bacteria

and the remainder of *Acidovorax* isolates, suggesting improved chemotaxis, competitive traits

and interaction with the plant metabolism and hormonal balance, possibly leading to a stronger

association between host and colonizers and explaining the better growth outcomes observed

in planta.

The evolutionary trajectory towards pathogenic plant-association

Acidovorax strains are naturally found to occupy widely diverse niches. In the frame of this
study, we reconstructed their pan-genome to explore the importance of genomic features
across the whole behavioural spectrum. Pan-genome analyses have become a staple to
estimate the complete gene repertoire accessible to an organism and to understand genotype
variations and evolution in a broader context [119, 120]. We found that the *Acidovorax* pan-
genome is largely made up by accessory or unique gene clusters (95 %) and predisposed to a
continuous inflation, a predictive feature of functionally flexible organisms [121], with each
occupied niche being a potential source of novel genomic traits.

Specialization has been associated with smaller genomes and previously reported in
pathogenic strains [121, 122]. Therefore, we studied *Acidovorax* genome size and density,
expecting to observe evidence of reductive evolution. For example, Merhej and colleagues
analysed a comprehensive collection of 317 bacterial genomes to trace the evolutionary path
from free-living to specialized intracellular strains [123], uncovering "massive gene loss" as a
consequence of more gene loss than horizontal gene transfer (HGT) events. Mainly in isolated
niches like the interior of roots, HGT rates are considered as low due to the missing interaction
with other microbiota. Surprisingly, in our study, the genomes of pathogenic *Acidovorax* were
the largest, yet showed a significantly lower gene density than both free-living and commensals
strains. Similar data was also observed for *Rickettsia prowazekii* [124], which features more
non-coding sequences than any of its close, non-pathogenic, relatives. In their review on
pathogenomics, Georgiades and Raoult argued that true bacterial speciation is only observed in
the contest of segregated niche, such in the case of obligated parasitism [125]. For Acidovorax,
Fegan [10] listed 28 known Graminaceae hosts for *Acidovorax avenae* subsp. a*venae* and 10
Cucurbitaceae natural hosts and 2 Solenaceae for *Acidovorax avenae* subsp. c*itrulli*. Thus,
Acidovorax pathogens seems still far from true speciation, as suggested by their wide choice of
hosts, and, likely, have already endured evolutionary driven gene losses, whereas extensive
genome reductions still have to occur.

Major functional differences across the groups

Principal component analysis revealed that pathogens differ in their genomes from commensal

and free-living strains. Among the differences that contributed the most to the separation, we

found that the lipase C-terminal domain and the mutagenesis inducer HIM1 were both enriched

in commensal strains. Assis and colleagues investigated the phylogeny of lipases in bacteria

and found orthologous of the same secreted lipase ubiquitous among plant-associated strains,

including *Acidovorax* [126]. Pathogens may be less dependent on lipases, as, during infection,

plant-derived lipids may be of minor value to them, as they preferentially hydrolyse

carbohydrates to cover their nutritional needs.

The presence of type VI secretion-systems has been strongly correlated with plant-association

in Gram-negative bacteria [127] and has been in observed in pathogenic and commensal

strains alike. Similarly, in *Acidovorax*, we retrieved a putative type VI secretion-system genomic

island in both groups of plant-associated bacteria (commensals and pathogens, but only in 13

% of the free-living strains). For the pathogens, we also identified an operon encoding

components of the type III secretion-system, absent from the other groups.

The distribution of these Pfams across the groups suggests the recruitment of commensals

from the larger pool of free-living strains through horizontal gene transfer of a single genomic

island encoding elements of the type VI secretion-system and further specialization of

commensals into pathogenic strains, through gene losses and acquisition of type III secretion-

systems.

We observed in the commensals an enrichment of several motility-associated domains. Pallen

and Wren [128] postulated the loss of motility as a common side-effect of intracellular

endosymbiosis, a phenomena recently witnessed in several emerging pathogens. It stands to

be clarified whether this adaptation occurs for disuse, to prevent recognition by the host

immune system or both. For commensals however, which are depending on chemotaxis to

make use of the plant-derived phytochemicals, motility is an essential function.

Finally, pathogens showed an enrichment of polyketide synthases and non-ribosomal peptide

synthases, which have been both extensively researched due to their promising role in

pharmaceutical applications for the production of diverse antimicrobial, immunosuppressive and

cytostatic compounds [129, 130]. Pathogenic *Acidovorax*, thus, have access to a wider array of

secondary metabolites, likely necessary for microbe-microbe competition [131] and possibly to

interact with the host.

Conclusions

The focus of this research study has been to assess the role of genomic traits across the

Acidovorax behavioural spectrum and to evaluate which genomic features can discriminate

plant-pathogenic from commensal and plant-growth promoting strains. We have reported many

discriminant traits through association of plant growth data with *in silico* pan-genome-wide

comparisons. The robustness of our analysis was also supported by our neural network

classifier, which accurately matched each individual to its observed phenotype by using the

information encoded in the pan-genome, validating our hypothesis. Researchers in the field of

genomics and microbiology have been investigating the potential of machine learning

applications to extract meaningful patterns from the high-dimensional data generated by Next-

Generation Sequencing [132, 133, 134, 135]. Even though our method was tested on a single

genus, it shows promises for generalization and the advancement of genotype-based, high-

throughput phenotyping and monitoring of emerging pathogens.

However, our data is based on genomic predictions and the importance of the described Pfams,

both for plant-growth promoting commensals as well as plant-pathogens, needs to be verified

firstly by analysing bacterial transcriptomes during the interaction with the plant, followed by

targeted mutagenesis and analysis of the impact of bacterial mutants on survival in the

rhizosphere and plant growth.

Authors statement

The authors declare no financial conflicts of interest.

Acknowledgement

Michael Schloter and Caroline Gutjahr were supported by grants from the Deutsche

Forschungsgemeinschaft [DFG; SCHL446/38-1 and GU1423/3-1] respectively the frame of the

SPP2125 "Deconstruction and reconstruction of the plant microbiota [DECRyPT]". We thank

Ruben Garrido-Oter for making the isolates available for the *L. japonicus* bioassays and the

respective sequenced genomes for the genomics comparison We highly appreciate the

discussions with Martin Parniske and Corinna Dawid. We thank M.H.H. Nguyen for help with

the *L. japonicus* bioassays.

Bibliography

1. Harbort CJ, Hashimoto M, Inoue H, Niu Y, Guan R, et al. Root-Secreted Coumarins and the Microbiota Interact to Improve Iron Nutrition in Arabidopsis. Cell Host Microbe 2020;28:825-837.e6. doi: 10.1016/j.chom.2020.09.006

2. Finkel OM, Salas-González I, Castrillo G, Conway JM, Law TF, et al. A single bacterial genus maintains root growth in a complex microbiome. Nature 2020;587:103–108. doi: 10.1038/s41586-020-2778-7

3. Berg G, Grube M, Schloter M, Smalla K. The plant microbiome and its importance for plant and human health. Front Microbiol 2014;5:1. doi: 10.3389/fmicb.2014.00491

4. Durán P, Thiergart T, Garrido-Oter R, Agler M, Kemen E, et al. Microbial Interkingdom Interactions in Roots Promote Arabidopsis Survival. Cell 2018;175:973-983.e14. doi: 10.1016/j.cell.2018.10.020

5. Rodriguez PA, Rothballer M, Chowdhury SP, Nussbaumer T, Gutjahr C, et al. Systems Biology of Plant-Microbiome Interactions. Mol Plant 2019;12:804–821. doi: 10.1016/j.molp.2019.05.006

6. Idnurm A, Howlett BJ. Analysis of loss of pathogenicity mutants reveals that repeat-induced point mutations can occur in the Dothideomycete Leptosphaeria maculans. Fungal Genet Biol 2003;39:31–37. doi: 10.1016/S1087-1845(02)00588-1

7. Kanda A, Ohnishi S, Tomiyama H, Hasegawa H, Yasukohchi M, et al. Type III secretion machinery-deficient mutants of Ralstonia solanacearum lose their ability to colonize resulting in loss of pathogenicity. J Gen Plant Pathol 2003;69:250–257. doi: 10.1016/S1087-1845(02)00588-1

8. Melnyk RA, Hossain SS, Haney CH. Convergent gain and loss of genomic islands drive lifestyle changes in plant-associated Pseudomonas. ISME J 2019;13:1575–1588. doi: 10.1016/S1087-1845(02)00588-1

9. Willems A, Gillis M. Acidovorax . In: Bergey's Manual of Systematics of Archaea and Bacteria. John Wiley & Sons, Ltd; 2015. pp. 1–16. doi: 10.1099/00207713-42-1-107

10. Fegan M. Plant pathogenic members of the genera Acidovorax and Herbaspirillum. In: Plant-Associated Bacteria. Springer Netherlands; 2006. pp. 671–702. doi: 10.1007/978-1-4020-4538-7_18

11. Giordano PR, Chaves AM, Mitkowski NA, Vargas JM. Identification, characterization, and

distribution of Acidovorax avenae subsp. avenae associated with creeping bentgrass etiolation

and decline. Plant Dis 2012;96:1736–1742. doi: 10.1007/978-1-4020-4538-7_18

12. Han S, Li D, Trost E, Mayer KF, Corina V, et al. Systemic responses of barley to the 3-

hydroxy-decanoyl-homoserine lactone producing plant beneficial endophyte acidovorax radicis

N35. Front Plant Sci 2016;7:1868. doi: 10.3389/fpls.2016.01868

13. Levy A, Salas Gonzalez I, Mittelviefhaus M, Clingenpeel S, Herrera Paredes S, et al. Genomic

features of bacterial adaptation to plants. Nat Genet 2018;50:138–150. doi: 10.1038/s41588-017-

0012-9

14. Wippel K, Tao K, Niu Y, Zgadzaj R, Guan R, et al. Host preference and invasiveness of

commensals in the Lotus and Arabidopsis root microbiota. BioRxiv 2021;2021.01.12.426357. doi:

10.1101/2021.01.12.426357

15. Eren AM, Esen OC, Quince C, Vineis JH, Morrison HG, et al. Anvi'o: An advanced analysis

and visualization platformfor 'omics data. PeerJ 2015;2015:e1319. doi: 10.7717/peerj.1319

16. Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, et al. Prodigal: Prokaryotic gene

recognition and translation initiation site identification. BMC Bioinformatics 2010;11:119. doi:

10.1186/1471-2105-11-119

17. Eddy SR. A new generation of homology search tools based on probabilistic inference.

Genome Inform 2009;23:205–211. doi: 10.1142/9781848165632_0019

18. El-Gebali S, Mistry J, Bateman A, Eddy SR, Luciani A, et al. The Pfam protein families

database in 2019. Nucleic Acids Res 2019;47:D427–D432. doi: 10.1093/nar/gky995

19. Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, et al. Metagenomic biomarker

discovery and explanation. Genome Biol 2011;12:R60. doi: 10.1186/gb-2011-12-6-r60

20. Delmont TO, Eren EM. Linking pangenomes and metagenomes: The Prochlorococcus

metapangenome. PeerJ 2018;2018:e4320. doi: 10.7717/peerj.4320

21. Pritchard L, Cock P, Esen Ö. pyani v0.2.8: average nucleotide identity (ANI) and related

measures for whole genome comparisons. Epub ahead of print 5 March 2019. doi:

10.5281/ZENODO.2584238.

22. Blin K, Shaw S, Steinke K, Villebro R, Ziemert N, et al. antiSMASH 5.0: updates to the

secondary metabolite genome mining pipeline. Nucleic Acids Res 2019;47:W81–W87. doi:

10.1093/nar/gkz310

23. Khalid S, Khalil T, Nasreen S. A survey of feature selection and feature extraction techniques

in machine learning. In: Proceedings of 2014 Science and Information Conference, SAI 2014.

Institute of Electrical and Electronics Engineers Inc.; 2014. pp. 372–378. doi:

10.1109/SAI.2014.6918213

24. Conn SJ, Hocking B, Dayod M, Xu B, Athman A, et al. Protocol: Optimising hydroponic growth

systems for nutritional and physiological analysis of Arabidopsis thaliana and other plants. Plant

Methods 2013;9:1–11. doi: 10.1186/1746-4811-9-4

25. Kelly DJ, Thomas GH. The tripartite ATP-independent periplasmic (TRAP) transporters of

bacteria and archaea. FEMS Microbiol Rev 2001;25:405–424. doi: 10.1111/j.1574-

6976.2001.tb00584.x

26. Simpson DA, Ramphal R, Lory S. Characterization of Pseudomonas aeruginosa fliO, a gene

involved in flagellar biosynthesis and adherence. Infect Immun 1995;63:2950–2957. doi:

10.1128/iai.63.8.2950-2957.1995

27. Barker CS, Meshcheryakova I V., Kostyukova AS, Samatey FA. FliO Regulation of FliP in the

Formation of the Salmonella enterica Flagellum. PLoS Genet 2010;6:e1001143. doi:

10.1371/journal.pgen.1001143

28. Absalon C, Van Dellen K, Watnick PI. A Communal Bacterial Adhesin Anchors Biofilm and

Bystander Cells to Surfaces. PLoS Pathog 2011;7:e1002210. doi: 10.1371/journal.ppat.1002210

29. Göhler AK, Staab A, Gabor E, Homann K, Klang E, et al. Characterization of MtfA, a novel

regulatory output signal protein of the glucose-phosphotransferase system in Escherichia coli K-

12. J Bacteriol 2012;194:1024–1035. doi: 10.1128/JB.06387-11

30. Xu Q, Göhler AK, Kosfeld A, Carlton D, Chiu HJ, et al. The structure of MLC titration factor a

(MtfA/YeeI) reveals a Prototypical zinc metallopeptidase related to anthrax lethal factor. J Bacteriol

2012;194:2987–2999. doi: 10.1128/JB.00038-12

31. Rosa LT, Bianconi ME, Thomas GH, Kelly DJ. Tripartite ATP-independent periplasmic (TRAP)

transporters and Tripartite Tricarboxylate Transporters (TTT): From uptake to pathogenicity.

Frontiers in Cellular and Infection Microbiology 2018;8:33. doi: 10.3389/fcimb.2018.00033

32. Cangelosi GA, Ankenbauer RG, Nestert EW. Sugars induce the Agrobacterium virulence

genes through a periplasmic binding protein and a transmembrane signal protein (crown

gapl/signal   transduction/ChvE/VirA/galactose-binding   proteins).   1990;87:6708-6712   doi:

http://www.jstor.org/stable/2355369.

33. Rodionova IA, Li X, Thiel V, Stolyar S, Stanton K, et al. Comparative genomics and functional

analysis of rhamnose catabolic pathways and regulons in bacteria. Front Microbiol 2013;4:407.

doi: 10.3389/fmicb.2013.00407

34. Yamauchi Y, Hasegawa A, Taninaka A, Mizutani M, Sugimoto Y. NADPH-dependent

reductases involved in the detoxification of reactive carbonyls in plants. J Biol Chem

2011;286:6999–7009. doi: 10.1074/jbc.M110.202226

35. Cipollone R, Ascenzi P, Tomao P, Imperi F, Visca P. Enzymatic detoxification of cyanide:

Clues from Pseudomonas aeruginosa rhodanese. J Mol Microbiol Biotechnol 2008;15:199–211.

doi: 10.1159/000121331

36. Pollmann S, Müller A, Weiler EW. Many Roads Lead to 'Auxin': of Nitrilases, Synthases, and

Amidases. Plant Biol 2006;8:326–333. doi: 10.1055/s-2006-924075

37. Lengeler KB, Tielker D, Ernst JF. Protein-O-mannosyltransferases in virulence and

development. Cellular and Molecular Life Sciences 2008;65:528–544. doi: 10.1007/s00018-007-

7409-z

38. Zowawi HM, Harris PNA, Roberts MJ, Tambyah PA, Schembri MA, et al. The emerging threat

of multidrug-resistant Gram-negative bacteria in urology. Nat Rev Urol 2015;12:570–584. doi:

10.1038/nrurol.2015.199

39. van den Bosch TJM, Tan K, Joachimiak A, Welte CU. Functional profiling and crystal

structures of isothiocyanate hydrolases found in gut-associated and plant-pathogenic bacteria.

Appl Environ Microbiol 2018;84:2020. doi: 10.1128/AEM.00478-18

40. Sultana A, Kallio P, Jansson A, Wang JS, Niemi J, et al. Structure of the polyketide cyclase

SnoaL reveals a novel mechanism for enzymatic aldol condensation. EMBO J 2004;23:1911–

1921. doi: 10.1038/sj.emboj.7600201

41. Raux E, Schubert HL, Warren MJ. Biosynthesis of cobalamin (vitamin B12): A bacterial

conundrum. 2000. Epub ahead of print 2000. doi: 10.1007/PL00000670.

42. Schoenhofen IC, Li G, Strozen TG, Howard SP. Purification and characterization of the N-

terminal domain of ExeA: A novel ATPase involved in the type II secretion pathway of Aeromonas

hydrophila. J Bacteriol 2005;187:6370–6378. doi: 10.1128/JB.187.18.6370-6378.2005

43. Schlomann M, Schmidt E, Knackmuss HJ. Different types of dienelactone hydrolase in 4-

fluorobenzoate-utilizing bacteria. J Bacteriol 1990;172:5112–5118. doi: 10.1128/jb.172.9.5112-

5118.1990

44. Schlomann M, Ngai KL, Ornston LN, Knackmuss HJ. Dienelactone hydrolase from

Pseudomonas cepacia. J Bacteriol 1993;175:2994–3001. doi: 10.1128/jb.175.10.2994-3001.1993

45. Lord DM, Uzgoren Baran A, Soo VWC, Wood TK, Peti W, et al. McbR/YncC: Implications for

the mechanism of ligand and DNA binding by a bacterial gntr transcriptional regulator involved in

biofilm formation. Biochemistry 2014;53:7223–7231. doi: 10.1021/bi500871a

46. Planamente S, Mondy S, Hommais F, Vigouroux A, Moréra S, et al. Structural basis for

selective GABA binding in bacterial pathogens. Mol Microbiol 2012;86:1085–1099. doi:

10.1111/mmi.12043

47. Thomson NR, Nasser W, McGowan S, Sebaihia M, Salmond GPC. Erwinia carotovora has two

KdgR-like proteins belonging to the IclR family of transcriptional regulators: Identification and

characterization of the RexZ activator and the KdgR repressor of pathogenesis. Microbiology

1999;145:1531–1545. doi: 10.1099/13500872-145-7-1531

48. Molina-Henares AJ, Krell T, Eugenia Guazzaroni M, Segura A, Ramos JL. Members of the

IclR family of bacterial transcriptional regulators function as activatorsand/or repressors. FEMS

Microbiology Reviews 2006;30:157–186. doi: 10.1111/j.1574-6976.2005.00008.x

49. Paget MSB, Helmann JD. The σ70 family of sigma factors. Genome Biology 2003;4:203. doi:

10.1186/gb-2003-4-1-203

50. Kazmierczak MJ, Wiedmann M, Boor KJ. Alternative Sigma Factors and Their Roles in

Bacterial Virulence. Microbiol Mol Biol Rev 2005;69:527–543. doi: 10.1128/mmbr.69.4.527-

543.2005

51. https://www.ncbi.nlm.nih.gov/taxonomy

52. Tettelin H, Riley D, Cattuto C, Medini D. Comparative genomics: the bacterial pan-genome.

Current Opinion in Microbiology 2008;11:472–477. doi: 10.1016/j.mib.2008.09.006

53. Chen CKM, Lee GC, Ko TP, Guo RT, Huang LM, et al. Structure of the

Alkalohyperthermophilic Archaeoglobus fulgidus Lipase Contains a Unique C-Terminal Domain

Essential for Long-Chain Substrate Binding. J Mol Biol 2009;390:672–685. doi:

10.1016/j.jmb.2009.05.017

54. Flaugnatti N, Le TTH, Canaan S, Aschtgen MS, Nguyen VS, et al. A phospholipase A1

antibacterial Type VI secretion effector interacts directly with the C-terminal domain of the VgrG

spike protein for delivery. Mol Microbiol 2016;99:1099–1118. doi: 10.1111/mmi.13292

55. Kelberg EP, Kovaltsova S V., Alekseev SY, Fedorova I V., Gracheva LM, et al. HIM1, a new

yeast Saccharomyces cerevisiae gene playing a role in control of spontaneous and induced

mutagenesis. Mutat Res - Fundam Mol Mech Mutagen 2005;578:64–78. doi:

10.1016/j.mrfmmm.2005.03.003

56. Karlowski WM, Zielezinski A, Carrère J, Pontier D, Lagrange T, et al. Genome-wide

computational identification of WG/GW Argonaute-binding proteins in Arabidopsis. Nucleic Acids

Res 2010;38:4231–4245. doi: 10.1093/nar/gkq162

57. Filloux A, Hachani A, Bleves S. The bacterial type VI secretion machine: yet another player for

protein transport across membranes. Microbiology 2008;154:1570–1583. doi:

10.1099/mic.0.2008/016840-0

58. Kapitein N, Bönemann G, Pietrosiuk A, Seyffer F, Hausser I, et al. ClpV recycles VipA/VipB

tubules and prevents non-productive tubule formation to ensure efficient type VI protein secretion.

Mol Microbiol 2013;87:1013–1028. doi: 10.1111/mmi.12147

59. English G, Byron O, Cianfanelli FR, Prescott AR, Coulthurst SJ. Biochemical analysis of TssK,

a core component of the bacterial Type VI secretion system, reveals distinct oligomeric states of

TssK and identifies a TssK-TssFG subcomplex. Biochem J 2014;461:291–304. doi:

10.1042/BJ20131426

60. Bingle LE, Bailey CM, Pallen MJ. Type VI secretion: a beginner's guide. Current Opinion in

Microbiology 2008;11:3–8. doi: 10.1016/j.mib.2008.01.006

61. Mintz KP, Fives-Taylor PM. impA, a gene coding for an inner membrane protein, influences

colonial morphology of Actinobacillus actinomycetemcomitans. Infect Immun 2000;68:6580–6586.

doi: 10.1128/IAI.68.12.6580-6586.2000

62. Suarez G, Sierra JC, Sha J, Wang S, Erova TE, et al. Molecular characterization of a

functional type VI secretion system from a clinical isolate of Aeromonas hydrophila. Microb Pathog

2008;44:344–361. doi: 10.1016/j.micpath.2007.10.005

63. Bladergroen MR, Badelt K, Spaink HP. Infection-blocking genes of a symbiotic Rhizobium

leguminosarum strain that are involved in temperature-dependent protein secretion. Mol Plant-

Microbe Interact 2003;16:53–64. doi: 10.1094/MPMI.2003.16.1.53

63. Zhou Y, Tao J, Yu H, Ni J, Zeng L, et al. Hcp family proteins secreted via the type VI secretion

system coordinately regulate Escherichia coli K1 interaction with human brain microvascular

endothelial cells. Infect Immun 2012;80:1243–1251. doi: 10.1128/IAI.05994-11

64. Cunnac S, Boucher C, Genin S. Characterization of the cis-Acting Regulatory Element

Controlling HrpB-Mediated Activation of the Type III Secretion System and Effector Genes in

Ralstonia solanacearum. J Bacteriol 2004;186:2309–2318. doi: 10.1128/JB.186.8.2309-2318.2004

65. Furutani A, Takaoka M, Sanada H, Noguchi Y, Oku T, et al. Identification of novel type III

secretion effectors in Xanthomonas oryzae pv. oryzae. Mol Plant-Microbe Interact 2009;22:96–

106. doi: 10.1094/MPMI-22-1-0096

66. van Gijsegem F, Gough C, Zischek C, Niqueux E, Arlat M, et al. The hrp gene locus of

Pseudomonas solanacearum, which controls the production of a type III secretion system,

encodes eight proteins related to components of the bacterial flagellar biogenesis complex. Mol

Microbiol 1995;15:1095–1114. doi: 10.1111/j.1365-2958.1995.tb02284.x

67. Baruch K, Gur-Arie L, Nadler C, Koby S, Yerushalmi G, et al. Metalloprotease type III effectors

that specifically cleave JNK and NF-κB. EMBO J 2011;30:221–231. doi: 10.1038/emboj.2010.297

68. Zhang H, Gao ZQ, Wang WJ, Liu GF, Xu JH, et al. Structure of the type vi effector-immunity

complex (Tae4-Tai4) provides novel insights into the inhibition mechanism of the effector by its

immunity protein. J Biol Chem 2013;288:5928–5939. doi: 10.1074/jbc.M112.434357

69. López-Lara IM, Kafetzopoulos D, Spaink HP, Thomas-Oates JE. Rhizobial NodL O-acetyl

transferase and nodS N-methyl transferase functionally interfere in production of modified nod

factors. J Bacteriol 2001;183:3408–3416. doi: 10.1128/JB.183.11.3408-3416.2001

70. Müller MG, Moe NE, Richards PQ, Moe GR. Resistance of Neisseria meningitidis to human

serum depends on T and B cell stimulating protein B. Infect Immun 2015;83:1257–1264. doi:

10.1128/IAI.03134-14

71. Carr S, Walker D, James R, Kleanthous C, Hemmings AM. Inhibition of a ribosome-

inactivating ribonuclease: The crystal structure of the cytotoxic domain of colicin E3 in complex

with its immunity protein. Structure 2000;8:949–960. doi: 10.1016/S0969-2126(00)00186-6

72. Krishnan HB. NolX of Sinorhizobium fredii USDA257, a type III-secreted protein involved in

host range determination, is localized in the infection threads of cowpea (Vigna unguiculata [L.]

Walp) and soybean (Glycine max [L.] Merr.) nodules. J Bacteriol 2002;184:831–839. doi: 10.1128/

JB.184.3.831-839.2002

73. El Ghachi M, Bouhss A, Barreteau H, Touzé T, Auger G, et al. Colicin M exerts its bacteriolytic

effect via enzymatic degradation of undecaprenyl phosphate-linked peptidoglycan precursors. J

Biol Chem 2006;281:22761–22772. doi: 10.1074/jbc.M602834200

74. Hattori Y, Iwata K, Suzuki K, Uraji M, Ohta N, et al. Sequence characterization of the vir region

of a nopaline type Ti plasmid, pTi-SAKURA. Genes Genet Syst 2001;76:121–130. doi:

10.1266/ggs.76.121

75. Fields KA, Nilles ML, Cowan C, Straley SC. Virulence Role of V Antigen of Yersinia pestis at

the Bacterial Surface. Infect Immun 1999;67:5395–5408. doi: 0019-9567/99/$04.000

76. Zhang D, de Souza RF, Anantharaman V, Iyer LM, Aravind L. Polymorphic toxin systems:

Comprehensive characterization of trafficking modes, processing, mechanisms of action, immunity

and ecology using comparative genomics. Biol Direct;7. Epub ahead of print 25 June 2012. doi:

10.1186/1745-6150-7-18.

77. Patzer SI, Albrecht R, Braun V, Zeth K. Structural and mechanistic studies of pesticin, a

bacterial homolog of phage lysozymes. J Biol Chem 2012;287:23381–23396. doi:

10.1074/jbc.M112.362913

78. Guyon K, Balagué C, Roby D, Raffaele S. Secretome analysis reveals effector candidates

associated with broad host range necrotrophy in the fungal plant pathogen Sclerotinia

sclerotiorum. BMC Genomics 2014;15:1–19. doi: 10.1186/1471-2164-15-336

79. Korotkov K V., Sandkvist M, Hol WGJ. The type II secretion system: Biogenesis, molecular

architecture and mechanism. Nat Rev Microbiol 2012;10:336–351. doi: 10.1038/nrmicro2762

80. Lohrke SM, Nechaev S, Yang H, Severinov K, Jin SJ. Transcriptional activation of

Agrobacterium tumefaciens virulence gene promoters in Escherichia coli requires the A.

tumefaciens rpoA gene, encoding the alpha subunit of RNA polymerase. J Bacteriol

1999;181:4533–4539. doi: 10.1128/jb.181.15.4533-4539.1999

81. Kadioglu A, Weiser JN, Paton JC, Andrew PW. The role of Streptococcus pneumoniae

virulence factors in host respiratory colonization and disease. Nat Rev Microbiol 2008;6:288–301.

doi: 10.1038/nrmicro1871

82. Charkowski AO, Alfano JR, Preston G, Yuan J, He SY, et al. The Pseudomonas syringae pv.

tomato HrpW protein has domains similar to harpins and pectate lyases and can elicit the plant

hypersensitive response and bind to pectate. J Bacteriol 1998;180:5211–5217. doi:

10.1128/jb.180.19.5211-5217.1998

83. Kotrba P, Inui M, Yukawa H. Bacterial phosphotransferase system (PTS) in carbohydrate

uptake and control of carbon metabolism. Journal of Bioscience and Bioengineering 2001;92:502–

517. doi: 10.1016/S1389-1723(01)80308-X

84. Johnson PE, Joshi MD, Tomme P, Kilburn DG, McIntosh LP. Structure of the N-terminal

cellulose-binding domain of Cellulomonas fimi CenC determined by nuclear magnetic resonance

spectroscopy. Biochemistry 1996;35:14381–14394. doi: 10.1021/bi961612s

85. Larson SB, Greenwood A, Cascio D, Day J, McPherson A. Refined molecular structure of pig

pancreatic α-amylase at 2.1 Å resolution. J Mol Biol 1994;235:1560–1584. doi:

10.1006/jmbi.1994.1107

86. Rodríguez-Sanoja R, Oviedo N, Sánchez S. Microbial starch-binding domain. Current Opinion

in Microbiology 2005;8:260–267. doi: 10.1016/j.mib.2005.04.013

87. Gijzen M, Kuflu K, Qutob D, Chernys JT. A class I chitinase from soybean seed coat. J Exp

Bot 2001;52:2283–2289. doi: 10.1093/jexbot/52.365.2283

88. Itoh T, Hibi T, Suzuki F, Sugimoto I, Fujiwara A, et al. Crystal structure of chitinase ChiW from

Paenibacillus sp. str. FPU-7 reveals a novel type of bacterial cell-surface-expressed multi-modular

enzyme machinery. PLoS One;11. Epub ahead of print 1 December 2016. doi:

10.1371/journal.pone.0167310.

89. Py B, Isabelle BG, Haiech J, Chippaux M, Barras F. Cellulase egz of Erwinia chrysanthemi:

Structural organization and importance of his98 and glu133 residues for catalysis. Protein Eng

Des Sel 1991;4:325–333. doi: 10.1093/protein/4.3.325

90. Palumbo JD, Sullivan RF, Kobayashi DY. Molecular characterization and expression in

Escherichia coli of three β-1,3-glucanase genes from Lysobacter enzymogenes strain N4-7. J

Bacteriol 2003;185:4362–4370. doi: 10.1128/JB.185.15.4362-4370.2003

91. Sugimoto K, Senda T, Aoshima H, Masai E, Fukuda M, et al. Crystal structure of an aromatic

ring opening dioxygenase LigAB, a protocatechuate 4,5-dioxygenase, under aerobic conditions.

Structure 1999;7:953–965. doi: 10.1016/S0969-2126(99)80122-1

92. Stehr M, Diekmann H, Smau L, Seth O, Ghisla S, et al. A hydrophobic sequence motif

common to N-hydroxylating enzymes. Trends Biochem Sci 1998;23:56–57. doi: 10.1016/S0968-

0004(97)01166-3

93. Kauppi B, Lee K, Carredano E, Parales RE, Gibson DT, et al. Structure of an aromatic-ring-

hydroxylating dioxygenasenaphthalene 1,2-dioxygenase. Structure 1998;6:571–586. doi: 10.1016/

S0969-2126(98)00059-8

94. Tsuchiya T, Ohta H, Okawa K, Iwamatsu A, Shimada H, et al. Cloning of chlorophyllase, the

key enzyme in chlorophyll degradation: Finding of a lipase motif and the induction by methyl

jasmonate. Proc Natl Acad Sci U S A 1999;96:15362–15367. doi: 10.1073/pnas.96.26.15362

95. Park SY, Chao X, Gonzalez-Bonet G, Beel BD, Bilwes AM, et al. Structure and function of an

unusual family of protein phosphatases: The bacterial chemotaxis proteins CheC and CheX. Mol

Cell 2004;16:563–574. doi: 10.1016/j.molcel.2004.10.018

96. Holmgren A, Brändén CL. Crystal structure of chaperone protein PapD reveals an

immunoglobulin fold. Nature 1989;342:248–251. doi: 10.1038/342248a0

97. Draper J, Karplus K, Ottemann KM. Identification of a chemoreceptor zinc-binding domain

common to cytoplasmic bacterial chemoreceptors. J Bacteriol 2011;193:4338–4345. doi: 10.1128/

JB.05140-11

98. Szurmant H, Ordal GW. Diversity in Chemotaxis Mechanisms among the Bacteria and

Archaea. Microbiol Mol Biol Rev 2004;68:301–319. doi: 10.1128/mmbr.68.2.301-319.2004

99. Ide N, Kutsukake K. Identification of a novel Escherichia coli gene whose expression is

dependent on the flagellum-specific sigma factor, FliA, but dispensable for motility development.

Gene 1997;199:19–23. doi: 10.1016/S0378-1119(97)00233-3

100. Pelicic V. Type IV pili: e pluribus unum? Mol Microbiol 2008;68:827–837. doi: 10.1111/j.1365-

2958.2008.06197.x

101. Serizawa M, Yamamoto H, Yamaguchi H, Fujita Y, Kobayashi K, et al. Systematic analysis of

SigD-regulated genes in Bacillus subtilis by DNA microarray and Northern blotting analyses. Gene

2004;329:125–136. doi: 10.1016/j.gene.2003.12.024

102. Martinez RM, Dharmasena MN, Kirn TJ, Taylor RK. Characterization of two outer membrane

proteins, FlgO and FlgP, that influence Vibrio cholerae motility. J Bacteriol 2009;191:5669–5679.

doi: 10.1128/JB.00632-09

103. Bäckman M, Källström H, Jonsson AB. The phase-variable pilus-associated protein PilC is

commonly expressed in clinical isolates of Neisseria gonorrhoeae, and shows sequence variability

among strains. Microbiology 1998;144:149–156. doi: 10.1099/00221287-144-1-149

104. Kawamoto T, Noshiro M, Shen M, Nakamasu K, Hashimoto K, et al. Structural and

phylogenetic analyses of RGD-CAP/βig-h3, a fasciclin- like adhesion protein expressed in chick

chondrocytes. Biochim Biophys Acta - Gene Struct Expr 1998;1395:288–292. doi: 10.1016/S0167-

4781(97)00172-3

105. Alm RA, Hallinan JP, Watson AA, Mattick JS. Fimbrial biogenesis genes of Pseudomonas

aeruginosa: pilW and pilX increase the similarity of type 4 fimbriae to the GSP protein-secretion

systems and pilY1 encodes a gonococcal PilC homologue. Mol Microbiol 1996;22:161–173. doi:

10.1111/j.1365-2958.1996.tb02665.x

106. Nambu T, Kutsukake K. The Salmonella FlgA protein, a putative periplasmic chaperone

essential for flagellar P ring formation. Microbiology 2000;146:1171–1178. doi: 10.1099/00221287-

146-5-1171

107. Funa N, Ohnishi Y, Ebizuka Y, Horinouchi S. Alteration of reaction and substrate specificity of

a bacterial type III polyketide synthase by site-directed mutagenesis. Biochem J 2002;367:781–

789. doi: 10.1042/BJ20020953

108. Ross P, Mayer R, Benziman M. Cellulose biosynthesis and function in bacteria. Microbiol

Rev;55. Epub ahead of print 1991. doi: 10.1128/mmbr.55.1.35-58.1991.

109. Matthysse AG. Role of bacterial cellulose fibrils in Agrobacterium tumefaciens infection. J

Bacteriol 1983;154:906–915. doi: 10.1128/jb.154.2.906-915.1983

110. Jones DL. Organic acids in the rhizosphere - A critical review. Plant Soil 1998;205:25–44.

doi: 10.1023/A:1004356007312

111. de Souza RSC, Armanhi JSL, Damasceno N de B, Imperial J, Arruda P. Genome Sequences

of a Plant Beneficial Synthetic Bacterial Community Reveal Genetic Features for Successful Plant

Colonization. Front Microbiol 2019;10:1779. doi: 10.3389/fmicb.2019.01779

112. Desbrosses GG, Kopka J, Udvardi MK. Lotus japonicus metabolic profiling. Development of

gas chromatography-mass spectrometry resources for the study of plant-microbe interactions. In:

Plant Physiology. American Society of Plant Biologists; 2005. pp. 1302–1318. doi:

10.1104/pp.104.054957

113. Liu CF, Tonini L, Malaga W, Beau M, Stella A, et al. Bacterial protein-O-mannosylating

enzyme is crucial for virulence of Mycobacterium tuberculosis. Proc Natl Acad Sci U S A

2013;110:6560–6565. doi: 10.1073/pnas.1219704110

114. Fernández-Álvarez A, Elias-Villalobos A, Ibeas JI. The O-mannosyltransferase PMT4 is

essential for normal appressorium formation and penetration in Ustilago maydis. Plant Cell

2009;21:3397–3412. doi: 10.1105/tpc.109.065839

115. Planamente S, Mondy S, Hommais F, Vigouroux A, Moréra S, et al. Structural basis for

selective GABA binding in bacterial pathogens. Mol Microbiol 2012;86:1085–1099. doi:

10.1111/mmi.12043

116. Li M, Guo R, Yu F, Chen X, Zhao H, et al. Indole-3-acetic acid biosynthesis pathways in the

plant-beneficial bacterium arthrobacter pascens zz21. Int J Mol Sci 2018;19:443. doi:

10.3390/ijms19020443

117. Chevrot R, Rosen R, Haudecoeur E, Cirou A, Shelp BJ, et al. GABA controls the level of

quorum-sensing signal in Agrobacterium tumefaciens. Proc Natl Acad Sci U S A 2006;103:7460–

7464. 10.1073/pnas.0600313103

118. Park DH, Mirabella R, Bronstein PA, Preston GM, Haring MA, et al. Mutations in γ-

aminobutyric acid (GABA) transaminase genes in plants or Pseudomonas syringae reduce

bacterial virulence. Plant J 2010;64:318–330. doi: 10.1111/j.1365-313X.2010.04327.x

119. Vernikos G, Medini D, Riley DR, Tettelin H. Ten years of pan-genome analyses. Current

Opinion in Microbiology 2015;23:148–154. doi: 10.1016/j.mib.2014.11.016

120. Tettelin H, Riley D, Cattuto C, Medini D. Comparative genomics: the bacterial pan-genome.

Current Opinion in Microbiology 2008;11:472–477. doi: 10.1016/j.gde.2005.09.006

121. Moran NA. Microbial minimalism: Genome reduction in bacterial pathogens. Cell

2002;108:583–586. doi: 10.1016/S0092-8674(02)00665-7

122. Wixon J. Reductive evolution in bacteria: Buchnera sp., Rickettsia prowazekii and

Mycobacterium leprae. 2001. Epub ahead of print 2001. doi: 10.1002/cfg.70.

123. Merhej V, Royer-Carenzi M, Pontarotti P, Raoult D. Massive comparative genomic analysis

reveals convergent evolution of specialized bacteria. Biol Direct 2009;4:13. doi: 10.1186/1745-

6150-4-13

124. Andersson SGE, Zomorodipour A, Andersson JO, Sicheritz-Ponte T, Cecilia U, et al. The

genome sequence of Rickettsia prowazekii and the origin of mitochondria. 1998;396. doi:

10.1038/2409

125. Georgiades K, Raoult D. Defining pathogenic bacterial species in the genomic era. Front

Microbiol 2011;1:151. doi: 10.3389/fmicb.2010.00151

126. Assis RDAB, Polloni LC, Patané JSL, Thakur S, Felestrino ÉB, et al. Identification and

analysis of seven effector protein families with different adaptive and evolutionary histories in

plant-associated members of the Xanthomonadaceae. Sci Rep;7. Epub ahead of print 1

December 2017. doi: 10.1038/s41598-017-16325-1.

127. Bernal P, Llamas MA, Filloux A. Type VI secretion systems in plant-associated bacteria.

Environ Microbiol 2018;20:1–15. doi: 10.1111/1462-2920.13956

128. Pallen MJ, Wren BW. Bacterial pathogenomics. Nature 2007;449:835–842. doi:

10.1038/nature06248

129. Weissman KJ, Leadlay PF. Combinatorial biosynthesis of reduced polyketides. Nature

Reviews Microbiology 2005;3:925–936. doi: 10.1038/nrmicro1287

130. Grünewald J, Marahiel MA. Chemoenzymatic and Template-Directed Synthesis of Bioactive

Macrocyclic Peptides. Microbiol Mol Biol Rev 2006;70:121–146. doi: 10.1128/mmbr.70.1.121-

146.2006

131. Khalid S, Baccile JA, Spraker JE, Tannous J, Imran M, et al. NRPS-Derived Isoquinolines

and Lipopetides Mediate Antagonism between Plant Pathogenic Fungi and Bacteria. ACS Chem

Biol 2018;13:171–179. doi: 10.1021/acschembio.7b00731

132. Dias R, Torkamani A. Artificial intelligence in clinical and genomic diagnostics. Genome

Medicine 2019;11:70. doi: 10.1186/s13073-019-0689-8

133. Iversen KH, Rasmussen LH, Al-Nakeeb K, Armenteros JJA, Jensen CS, et al. Similar

genomic patterns of clinical infective endocarditis and oral isolates of Streptococcus sanguinis and

Streptococcus gordonii. Sci Rep 2020;10:1–11. doi: 10.1038/s41598-020-59549-4

134. Qu K, Guo F, Liu X, Lin Y, Zou Q. Application of machine learning in microbiology. Front

Microbiol 2019;10:827. doi: 10.3389/fmicb.2019.00827

135. Tarca AL, Carey VJ, Chen X wen, Romero R, Drăghici S. Machine learning and its

applications to biology. PLoS Comput Biol 2007;3:e116. doi: 10.1371/journal.pcbi.0030116

pathogen (21)   free-living (23)   commensal (62)

FIGURES:

Fig.1: Heat-map showing the effects of single-strains on *Lotus japonicus* growth parameters as observed in the *L. japonicus* growth bioassays. Measures are shown as the median, for each observed group, of the percentage change from the control.

Fig.2: Bar-plot of the logarithmic scores from the Linear Discriminant analysis for the discriminant Pfams. Pfams are assigned to 4, color-coded, functional groups.

Fig.3: Anvi'o 6.2 circular representation of *Acidovorax* pan-genome and heat-map of average nucleotide identity across the genomes.

Fig.4: a) Dot-chart of individuals logarithmic genome length. b) box-plot of groups logarithmic genome length. c) box-plot of groups gene density expressed as number of genes per 1000 base-pairs. Significance was assessed by Student's t-test with Holm's correction ($p < 0.05$).

Fig.5: a) Ordination plot for the first and second component of the principal component analysis (PCA) b) Ordination for the first and third component of the PCA. c) Bar-plot of the Pfams with the highest contribution for first, second and third component of the PCA.

Fig.6: a) Volcano plot displaying Pfams enriched in commensal (left) and pathogenic strains (right). On the x axis is represented logarithmic fold change and on the y axis adjusted *q*-value. Dots are represented in red when crossing the significance threshold of 0.05. b) Dot-plot of the curated list of enriched Pfams related to host-microbe or microbe-microbe interaction. Each dot is color coded based on the functional family of assignment.

Fig.7: Bar-plot of the percentage of genomes per group featuring each of the retrieved biosynthesis gene clusters.