Effects of land-use intensification and environmental stressors on intraspecific functional traits variability in bumblebees

Nicola Tommasi [1], Emiliano Pioltelli [1], Paolo Biella [1], Massimo Labra [1, 2], Maurizio Casiraghi [1], Andrea Galimberti [1, 2]*

1 -ZooplantLab, Department of Biotechnology and Biosciences, University of Milano-Bicocca, Milano, Italy

2 -INFN Sezione di Milano Bicocca, Milano, Italy

*Correspondence: andrea.galimberti@unimib.it

AUTHOR CONTRIBUTION

NT, EP and AG conceived the ideas and designed methodology; NT, EP and PB collected the data; NT and EP analysed the data, PB contributed to analyses checking/development and checked the final model outcomes; EP, NT and PB led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

ABSTRACT

Landscape anthropization replaces natural areas with agricultural or urban covers, leading to land-use intensification. This worldwide phenomenon affects biodiversity, but little is known about the effects on the intraspecific variation of functional traits related to ecosystem services. By combining field data collection, remote sensing and land cover analysis we investigated how land use intensification affects functional traits in two bumblebee species. Namely, the impact of different land use cover (i.e., semi-natural, urban, and agricultural) and of multiple biotic and abiotic stressors (i.e., temperature, resource availability, and air pollutants) was investigated. Along a gradient of landscape anthropization, we sampled populations of two European bumblebee species (*Bombus terrestris* and *B. pascuorum*) at 37 sites in Northern Italy. Through geometric morphometrics we investigated the variation of morphological traits related to flight performance (i.e., wing centroid size and shape and size fluctuating asymmetry FA), previously used as indicators of stress during insect development. Our results point out an idiosyncratic response of the two species to landscape anthropization. Smaller individuals of *B. pascuorum* were observed in response to increased impervious cover and temperature. No similar patterns were noticed in *B. terrestris*, which was characterized by larger individuals in response to floral resource availability. Wing size FA was positively associated with warmer temperatures and increased levels of NO_2 only in *B. terrestris*. Overall, this study found taxon-specific functional trait variation at the intraspecific level in syntopic organisms, expanding our understanding about the effects of land-use intensification on the ecological activity of pollinator model species.

Keywords: Functional diversity, heat islands, pollination, urban ecosystems, wing asymmetry

INTRODUCTION

Widespread phenomena, such as the progressive expansion of agricultural and urban lands are driving deep changes on ecosystems, among which the replacement of natural areas and the creation of novel ecosystem conditions Plants and animals respond to these marked environmental variations by shifting their distribution (Colla et al., 2012) and phenology (Huchler et al., 2020), and/or shaping some morphological traits considered "functional", i.e., relevant for their ecology, fitness and behaviour (Alberti et al., 2017; Eggenberger et al., 2019; Nooten & Rehan, 2020). Trait variation due to environmental alteration is particularly relevant in those organisms responsible for ecosystem services, such as pollinators, since changes in their traits could jeopardize the service provided (Buchholz & Egerer, 2020). For instance, land-use intensification and degradation (i.e., the growing proportion of anthropized or productive surfaces across landscapes) could lead to limited floral resource availability to pollinators (Steffan-Dewenter et al., 2001). This scenario, in turn, could impose changes in pollinator functional traits, such as body size decline due to less food supplied to larvae. Furthermore, anthropization of landscapes changes the local microclimate, thus determining an alteration of pollinator insects ecology, development and activities (Radmacher & Strohm, 2010). Specifically, higher degrees

of impervious land cover are often associated with increasing temperatures (Chun &

Guldmann, 2018), a phenomenon known as "heat island effect", that characterizes

both urban and suburban areas. Observations from previous studies have strengthened

the hypothesis that pollinator insects could face a shift towards smaller body size as an

adaptation to reduce risk of overheating in presence of warmer conditions (Peters et

al., 2016; Gérard et al., 2018a). Considering that the steady growth of the human

population is driving a dramatic increase in the area allocated to intensive agriculture

and urban expansion worldwide, new insights in the study of pollinator response are

necessary to predict which scenarios would likely occur for urban biodiversity and its

mediated services.

Previous studies investigated the morphological responses of pollinators to

anthropogenic pressures, mainly focusing on key functional traits, such as body size

(e.g., Chown & Gaston, 2010; Eggenberger et al., 2019; Theodorou et al., 2020). In

bees, this character responds rapidly to environmental changes (Chown & Gaston,

2010), it shows little heritability and its variation mainly depends on the amount of

food received during the larval development (Couvillon et al., 2010). Bee size directly

influences their mobility and so their foraging range (Greenleaf et al., 2007), but also

determines the metabolism rate and resource needs of adult imagos, with larger bees

having higher metabolism rate (Kelemen et al., 2019) and thus potentially being more

susceptible to variation in floral resource availability leading to starvation (Couvillon

& Dornhaus, 2010). However, to date, the investigation of pollinator body size

variation in anthropogenic habitats yielded heterogeneous results. A recent study on

bumblebees found bigger workers in cities (Theodorou et al., 2020). This pattern has

been interpreted as an adaptation to longer flights for collecting resources, due to severe habitat fragmentation and dispersion caused by land use intensification (Greenleaf et al., 2007). Conversely, a study by Eggenberger et al. (2019) found smaller bumblebee foragers in cities, interpreted as an effect of both limited local resource availability and warmer temperature in urban areas.

A promising research field in the study of morphological traits is to identify possible stress biomarkers of habitat quality and population health status (Adams et al., 2001). The advantage of using trait variation to measure stress resides in the fact that changes of phenotypes are detectable before a decrease in population viability occurs (Hoffmann et al., 2005). Therefore, quantifying traits variation could become an essential practice when evaluating local- and landscape-level stressors. A metric that has grown in popularity is the fluctuating asymmetry (FA) (Klingenberg, 2001; Beasley et al., 2013; Alves-Silva et al., 2018), defined as the presence of small, randomly placed deviations from perfect bilateral symmetry due to the occurrence of developmental instability, driven by exogenous environmental conditions (Klingenberg, 2015). FA differs from another type of bilateral asymmetry, the directional asymmetry (DA), that occurs when one specific side tends to be steadily larger than the other. While DA have a genetic basis and therefore could be less impacted by the environment (Palmer & Strobeck, 2003), the FA is considered a valid proxy of stress exposure to conditions that typically occur in anthropized environments (e.g., temperature, air pollutants, and pesticides) (Beasley et al., 2013). For instance, laboratory-based studies have demonstrated that CO_2 or low temperature leads to wing FA, supporting the possible role of traffic pollutant in determining

developmental instability (Klingenberg et al., 2001; Hoffmann et al., 2002). Interesting, however, these studies also indicated how wing shape and wing size asymmetries had different responses to the same stressor type. This aspect is ecologically very relevant. Both wing size and shape are important functional traits in pollinators. This is because wing size is believed to be related to flight length and it influences metabolic costs (Fernandez et al., 2017; Soule et al., 2020), while shape is considered as important for flight maneuverability (Kolliker et al., 2003, Grilli et al., 2017). Thus, their asymmetry could reduce flight performance ability and thus could reduce fitness and even impact on species interactions and pollination efficiency.

In order to characterize the effects of landscape anthropization and related pressures on pollinator insects, we quantified the morphological variation in two species of bumblebee (i.e., *Bombus pascuorum* and *B. terrestris*), by investigating different populations across a gradient of growing land-use alteration, from semi-natural areas to rural sites and urban places. Specifically, we chose the region surrounding Milan (Northern Italy). This is one of the major European cities and lies on a territory that experienced a strong anthropogenic footprint and ongoing efforts are trying to mitigate it. We expected to find quantitative variation in bumblebee functional traits (i.e.body size and wing FA) in response to increased land-use intensification. Our hypothesis was that in anthropized areas, pressures eliciting a decrease in body size (e.g., high temperature and low floral resource availability) would be prevailing and would lead to smaller foragers in urbanized and agricultural habitats. We also expected to find increased wing size and shape FA in response to higher levels of biotic and abiotic stressors (e.g., floral resources abundance,

temperature, and air pollutants). Hence, our ultimate goal was to determine which type

of response prevails and the putative pressures that are determining it to better

understand the impact of landscape anthropization on the ecology of pollinator insects.

MATERIALS AND METHODS

Study species

This study was focused on two co-occurring species of bumblebee: *Bombus*

terrestris (Linnaeus 1758) and *B. pascuorum* (Scopoli 1763). Both species are

efficient pollinators, common in Europe, and can be found in different habitats from

the natural to the agricultural and urban ones (Polce et al., 2018). Given these

characteristics, these species are reliable models to investigate responses by

pollinating insects to landscape anthropization (Eggenberger et al., 2019; Theodorou

et al., 2020). The two selected species have slightly different foraging ranges, with

estimated maximum dispersal range of 449 and 758 m for *B. pascuorum* and *B.*

terrestris, respectively (Knight et al., 2005). Nesting habits are also dissimilar as *B.*

terrestris builds its nest in subterranean holes, while *B. pascuorum* on top of or

slightly beneath the soil surface (Goulson, 2010). Another important difference is

represented by their dietary regimes with the *B. pascuorum* usually having a narrower

trophic niche and a preference for deep-corolla flowers (Harder, 1985).

Study design and sampling

Samplings were conducted at 37 sites (Fig. 1), mainly lying within the

metropolitan area of Milan in July 2019, only on days with sunny and not windy

weather conditions.

The study sites were distributed in an area of about 1800 km^2 covering four

administrative provinces, Milano, Monza e della Brianza, Lecco and Como. A

minimum distance between sites of 1 km was imposed to avoid the non-independence

of sites (Phillips et al., 2019) since it is above the mean foraging range observed for

the two species (Knight et al., 2005). A Moran test was also applied to confirm the

absence of spatial auto-correlation within sites (P>0.05).

Study sites spread along a gradient of landscape anthropization, ranging from areas

dominated by semi-natural hay meadows close to forest, to agriculture dominated

landscapes (i.e., sites at the edges of crop fields of mainly large monoculture) and sites

characterized by a high degree of impervious surface (i.e., concrete, building, asphalt).

For each species, five to six specimens, excluding queens and males, were captured in

a plot of about 50 m x 50 m at each site using an entomological net. After collection,

the insects were stored at -80 °C until further analysis.

Landscape and environmental variables

In order to assess landscape anthropization, land-use data were obtained from

the latest version of land cover database provided by Lombardy region (2018-DUSAF

6.0;   https://www.dati.lombardia.it/Territorio/DUSAF-6-0-Uso-del-suolo-

[2018/7rae198](2018/7rae198) fng6). Using QGIS 3.10.11, a 1 km radius buffer area was created

around each site where landscape composition was evaluated arranging DUSAF

original level and sub-level of land-use classification into three macro categories:

impervious (i.e., buildings, infrastructures, roads, and cemented surface), agricultural

(i.e., arable land devoted to crop production) and semi-natural land (i.e., meadows,

forest and urban green spaces) (Online resource 1 for a list of DUSAF codes assigned

to each grouping). For each site, the ratio between impervious and semi-natural land

(Impervious / Natural) as well as the ratio between agricultural and semi-natural land

(Agricultural / Natural) was computed to describe the urbanization and agricultural

intensification processes, respectively. Habitat fragmentation was quantified by

computing the edge density (ED), as the ratio of edge length of green and semi-natural

patches over their total area (Wang et al., 2014).

Other environmental biotic and abiotic features were considered to test for their

potential effects on altering body size and wing size/shape FA.Specifically, land

surface temperature was calculated as the mean value in the period June-July using

data retrieved through remote sensing imaging spectroradiometer (MODIS)

MOD11A2   from   the   NASA   database

(https://modis.gsfc.nasa.gov/data/dataprod/mod11.php) with a resolution of 1 km. The

choice of considering temperature for these two months was dictated from the

supposition that our sampled specimens developed during that period, according to the

biology of two selected species (Goulson, 2010). Air pollution was estimated as the

mean of daily concentrations of NO_2 over two months (June and July) registered by

Regional agency for environmental protection (ARPA) through monitoring stations

located nearby our sampling sites (https://www.arpalombardia.it/Pages/Aria/qualita-

aria.aspx).

Floral resources at each site (i.e., the total number of flowers) was estimated in

sampling plots by using six randomly placed quadrats 1 m x 1 m (covering a

proportion of sampling area similar to that reported in Fisher et al., 2017) and

summing the number of flowers found within each quadrat.

Specimens imaging and wings measurement

The forewings of all individuals were detached at the base and scanned at high

resolution. The obtained images were converted into TPS files using tps-UTIL 1.74

and digitised using the tps-Dig 2.31 software (Rohlf, 2015), with two-dimensional

cartesian coordinates of 15 landmarks positioned at wing vein junction (Fig. 2) (as in

Aytekin et al., 2007; Klingenberg et al., 2001). Bumblebees with damaged or badly

worn wings were excluded from further analyses.

The analysis of landmark configuration was conducted in MorphoJ 1.07 software

(Klingenberg, 2011). To remove all differences unrelated to shape (i.e., rotation,

translation, and scale) a generalized least square Procrustes superimposition was

applied (Klingenberg, 2011). The output of this procedure is a new set of

superimposed landmark coordinates, called 'Procrustes shape coordinates', that

contains all the shape information. Wing size was estimated as the centroid size (i.e.,

the square root of the sum of squared distances from the centroid of each landmark

configuration) and used as a proxy of body size (hereafter "body size", as in

Outomuro & Johansson, 2011 and Dellicour et al., 2017). Wing size asymmetry was

computed by dividing the absolute difference between left and right centroid sizes by

the mean centroid size and multiplying by 100 (Leonard et al., 2018). To estimate

wing shape variation, Procrustes distances were computed for each individual

(Klingenberg, 2015). These represent the measure of an individual's overall

asymmetry (i.e., sum of DA and FA components), obtained by taking the square root

of the sum of squared distances between corresponding right and left Procrustes'

coordinates.

Statistical analysis

Since the levels of asymmetry in bilateral traits are subtle, we estimated the

measurement error, which could possibly cause considerable variation in the

assessment of asymmetry levels, by double-scanning wings and digitizing their

landmarks for a subset of 50 specimens, in MorphoJ with the Procrustes ANOVA

(Klingenberg, 2001), in order to evaluate if measurement error was negligible. The

presence of directional asymmetry (DA) was also tested with the same approach, and

if DA occurred, the total asymmetry measures were corrected subtracting the mean

DA from the overall individual asymmetry, thus isolating the FA component as in

Costa et al. (2015).

To investigate the relationship between morphological traits and covariates (i.e., ratio

between impervious and natural cover, ratio between agricultural and natural cover,

edge density, temperature, NO_2, and floral resources) a series of linear mixed models

were used. In order to improve the fit between the predictors and the response

variable, mathematical transformations were applied on the covariates as reported in

Table 1. The goodness of these transformations was assessed through the AIC

criterion. (Δ AIC > 2).

The response of the two species was assessed separately, furthermore different models

were used to evaluate the effects of highly correlated variables (r>0.7). Specifically,

the response of centroid size was tested in two different models to account for

covariates collinearity. The first one (Table 1, model id A) tested the effect of i) edge

density of green areas, ii) ratio between impervious and semi-natural land cover, iii)

ratio between agricultural and semi-natural land-cover, and iv) floral resources.

Conversely, the second model (Table 1, model id B) evaluated separately the impact

of temperature because of its correlation with the ratio between impervious and semi-

natural land cover. Regarding FA, the effects of covariates known to influence the

developmental stability as temperature (Table 1, model id C), and NO_2 (Table 1,

model id D) were tested individually both for wing size FA and wing shape FA

because of their correlation. In all models, the sampling site was included as the

random effect. For all models we applied a stepwise model selection based on AIC

values removing variables that did not improve the model fit (Zuur et al., 2009).

Statistical analyses were conducted using MorphoJ 1.07 and JMP 14.2.0 (© SAS

Institute Inc. Cary, North Carolina, US).

RESULTS

Overall, the sampling dataset was similar in the two bumblebee species,

encompassing a total of 179 *B. pascuorum* (mean per site = 4.8±0.4) and 169 *B.*

terrestris (mean per site = 4.5±0.3) that were considered for morphometric analyses.

The measurement error was negligible because not significant for wing size (df = 79,

F = 2.67 p = 0.4578) and shape (df = 2054, F = 0.51, p = 0.9976). Different patterns of

size variation were found in the two bumblebee species. *B. pascuorum* decreased its

body size in response to both the increasing temperature (Fig. 3a; Table 1: model id B)

and the higher ratio of impervious on semi-natural land cover (Online resource 2 - Fig.

S1 ; Table 1: model id A). Conversely, no significant effect of these two predictor

variables was found on *B. terrestris*, that increased body size in response to the

availability of floral resources (Fig. 3b; Table 1: model id A).

Concerning wing asymmetry, both species showed a significant level of shape DA (*B.*

pascuorum df = 26, F = 4.66; p < 0.0001; *B. terrestris* df = 26, F = 5.60; p < 0.0001),

while size DA was statistically significant only in *B. pascuorum* (df = 1, F = 29.77; p

< 0.0001; in *B. terrestris* df = 1, F = 0.51; p = 0.4779). Measures of FA were then

obtained correcting for directional components.

B. terrestris size FA was positively correlated to temperature (Fig. 4; Table 1: model

id C) and concentration of NO_2 (Online resource2 - Fig. S2; Table 1: model id D), but

no similar response was observed in *B. pascuorum.* None of the predictor variables

(i.e., temperature, NO_2, resource abundance) showed any significant effect on wing

shape asymmetry in both bumblebee species.

305 Table 1: Results of the regression analysis of centroid size and fluctuating size asymmetry as a function of

306 landscape and biotic and abiotic covariates. Only final models selected through AIC criterion with significant

307 covariates are reported. β_i: regression coefficient; F: F-value; P: p-values. Models and results of size FA for *B.*

308 *pascuorum* and of shape FA for both species are not reported in the table as they were inconsistent.

Response variable	Model ID	Covariates	Bi	Df; F	P > F
B. pascuorum Centroid size	A	Impervious/Natural (Log)	-0.00869	1; 4.2661	0.0462*
	B	Temperature	-0.00341	1; 7.7271	0.0088*
B. terrestris Centroid size	A	Floral resources (Log)	0.0725	1; 11.0222	0.0026*
B. terrestris Fluctuating size asymmetry	C	Temperature	0.05259	1; 11.5565	0.0019*
	D	NO$_2$	0.01433	1; 7.2304	0.0118*

DISCUSSION

In this study, we quantified the spatial intraspecific variation of functional traits

that can shape the ecology of two common bumblebee species, possibly altering their

dispersion ability, and pollination efficiency. We focused on their variation along the

main trajectories of land use alteration, namely urbanization and agriculture

intensification of landscapes, by relating variation of body size and wing asymmetry

with landscape features and environmental stressors.

Landscape anthropization was found to be associated with intraspecific

variation of body size (i.e., estimated as wing centroid size) across bumblebees

populations. Specifically, foragers of one of the two investigated species, *B.*

pascuorum, showed a shift toward smaller body size in response to increasing

proportion of impervious surfaces and temperature, two conditions that characterize

urban landscapes and emerged as highly correlated from our analysis. A similar

pattern of body size reduction in urban bumblebees was reported by Eggenberger et al.

(2019) and it agrees with observations showing a general decrease of body size in

urban insect communities (Merckx et al., 2018). These works suggest that a potential

driver of size reduction in bumblebees is represented by the diminished, and spatially

dispersed, availability of floral resources that is often associated with higher

impervious cover (Eggenberger et al., 2019). However, in our study, neither the

estimation of floral resources, nor the fragmentation of green areas (i.e., the edge

density), correlated with size variation in *B. pascuorum*, the species that responded to

urbanization. Hence, a more reliable driver of urban-induced size shrinkage could be

warmer temperatures linked to the urban "heat island effect" (Merckx et al., 2018),

and this is also supported by the results of our study. Indeed, temperature has a strong

effect on insect body size, with multiple historical series collections-based and

experimental studies that revealed how higher environmental temperatures represent a

driver of body size reduction in different species of bees (e.g., Nooten & Rehan, 2020;

Theodorou et al., 2020). Higher temperature accelerates larval development, which

likely result in smaller adults (Sibly & Atkinson, 1994). Furthermore, smaller sizes in

warmer areas could also be a strategy for reducing overheating risks while foraging,

due to an increased convective heat loss in smaller bees (de Farias-Silva & Freitas,

2020). Functionally, smaller foragers could travel shorter foraging distances

(Greenleaf et al., 2007) and could also load less pollen and nectar (Goulson et al.,

2002). As a consequence, the shift towards smaller body size in *B. pascuorum* in more

urbanized environments could imply that it will pollinate less plants or handle flowers

less efficiently (Földesi et al., 2020), a concerning aspect in view of colony provision

and pollination.

In our study *B. terrestris* body size was found to respond only to the abundance of

flower resources, with larger individuals observed where potentially more food is

available. This is in accordance with the observation that adult size is strictly

correlated with the amount of food received during larval development (Couvillon &

Dornhaus 2009). However, this trend seems to be not clearly confirmed by *B.*
pascuorum probably due to the higher specialization of this species (Harder, 1985)
that may prevent it from benefiting from higher availability of flowers, since this not
necessarily results in higher possibility to exploit adequate resources for their diet.

Despite *B. pascuorum* and *B. terrestris* belong to the same genus, the responses
of these two species were not equal. Similarly, idiosyncratic responses were also
observed in other bumblebee species, where if the body size of some decreased over
warming decades, others responded in the opposite way (Gérard et al., 2020). In our
study the invariant size of *B. terrestris* in warmer conditions could be explained by the
high heat tolerance of this species (Martinet et al., 2020). In addition, *B. terrestris*
nests further underground compared to the other bumblebee, and might be less
exposed to warm, urban air temperatures during larval stages. These aspects
strengthen the hypothesis that temperature is a major determinant of pollinator size
reduction in cities because they affected the body size of the more temperature-
sensitive species, but not the heat-tolerant one. These species-level, idiosyncratic
responses are very relevant for our understanding of the potential mechanism of
intraspecific trait variation associated with landscape anthropization. This supports the
need to consider a wider panel of species when investigating the impact of landscape
anthropization on functional biodiversity.

Flight performance does not only depend on body size, but it is also affected by
asymmetries in shape and size between wings (Grilli et al., 2017; Soule et al., 2020).
Here, we found that wing size FA was positively correlated with environmental
stressors such as increased temperatures and NO_2 concentration in *B. terrestris*. The

absence of variations in shape asymmetry registered for both the bumblebee species

could confirm the results from other studies that have indicated shape variation as less

susceptible to stressors than size asymmetry (e.g., Gérard et al., 2018b). Variation in

wing asymmetry was observed in other insect taxa; the effect of temperature was

previously investigated under controlled laboratory conditions (Mpho et al., 2002)

while the role of pollutants was discussed interpreting the results of a field

experiment evaluating the impact of road traffic exposure (Leonard et al., 2018).

Studies associated the increased wing FA to environmental stressors, indicating that

impairment of developmental processes might take place (e.g., Klingenberg et al.,

2001; Kerr et al., 2013). Concerning pollutants, Klingenberg et al. (2001)

hypothesized that they could alter development through mechanisms related to gas

exchange. Trait variation does not only show developmental instability, but it also has

ecological implications. Wing size FA impacts the management of lengthy flights

(Fernandez et al., 2017, Soule et al., 2020), compared to wing shape which is often

associated with flight maneuverability. Thus, size asymmetry could specifically

impact flight performance, flight length in time and space, and even affect bumblebee

interactions with plants, the provisioning of larvae with food and their pollination rate.

CONCLUSIONS

Overall, the experimental results obtained in this study suggest that landscape

anthropization could affect different functional traits, and it occurs heterogeneously in

different pollinator species. Eventually, these responses could bring to the alarming

outcome of decreased pollination efficiency, because the studied traits are often

involved in flying abilities. Temperature emerged as one of the main drivers of these

phenotypic variations, affecting the two species in a different manner. The multiple

responses to the same stressor, observed for the two species, underline the necessity

for future studies to consider a wider panel of taxa instead of single model species.

From a conservation perspective, the comprehension of how pollinators cope with the

challenging conditions occurring in novel anthropogenic habitats, plays a key role in

informing suitable policy efforts to conserve their biodiversity and the ecosystem

service they provide. In the future, cities are predicted to expand constantly and thus

designing of urban landscapes will become a fundamental step for achieving

sustainability outcomes. The creation and the wise management of urban green spaces

will allow not only to create conditions that could support pollinators species

abundance and diversity but also sustain the biological interactions that are the basis

of a correct ecosystem functioning. At the same time, urban forestry and greenery

practices (e.g., plantation of street and residential trees and the creation of urban

greenbelts or greenways) could represent a valid solution to mitigate stressful

conditions related to the urban environment, such as the "heat island effect" (Chun &

Guldmann, 2018) and the high level of air pollution (Ottosen & Kumar, 2020) that

here were found to influence functional diversity issues.

ACKNOWLEDGMENTS:

The authors thank Amelia Pioltelli for artworks, Carola Miuccio for her technical

help, Giulia Masoero and Lorenzo Guzzetti for useful advice during manuscript

preparation.

DECLARATIONS:

Funding: This research was partially supported by the PIGNOLETTO project, co-

financed with the resources of POR FESR 2014-2020, European regional

development fund with the contribution of resources from the European Union, Italy

and the Lombardy Region.

Conflicts of interest/Competing interests: The authors declare that no competing

interests exist.

Ethics approval: Sampling permits were obtained when needed from local authorities

Availability of data and material: All relevant data are within the paper or stored in

a public repository (https://doi.org/10.6084/m9.figshare.13637594).

Authors' contributions: NT, EP and AG conceived the ideas and designed

methodology; NT, EP and PB collected the data; NT and EP analysed the data, PB

contributed to analyses checking/development and checked the final model outcomes;

EP, NT and PB led the writing of the manuscript. All authors contributed critically to

the drafts and gave final approval for publication.

REFERENCES

Adams, S. M., Giesy, J. P., Tremblay, L. A., & Eason, C. T. (2001). The use of biomarkers in ecological risk assessment: Recommendations from the Christchurch conference on Biomarkers in Ecotoxicology. *Biomarkers*, *6*(1), 1–6. https://doi.org/10.1080/135475001452724

Alberti, M., Correa, C., Marzluff, J. M., Hendry, A. P., Palkovacs, E. P., Gotanda, K. M., Hunt, V. M., Apgar, T. M., & Zhou, Y. (2017). Global urban signatures of phenotypic change in animal and plant populations. *Proceedings of the National Academy of Sciences*, *114*(34), 8951–8956. https://doi.org/10.1073/pnas.1606034114

Alves-Silva, E., Santos, J. C., & Cornelissen, T. G. (2018). How many leaves are enough? The influence of sample size on estimates of plant developmental instability and leaf asymmetry. *Ecological Indicators*, *89*, 912–924. https://doi.org/10.1016/j.ecolind.2017.12.060

Aytekin, M. A., Terzo, M., Rasmont, P., & Çağatay, N. (2007). Landmark based geometric morphometric analysis of wing shape in Sibiricobombus Vogt (Hymenoptera: Apidae: Bombus Latreille). *Annales de la Société entomologique de France (N.S.)*, *43*(1), 95–102. https://doi.org/10.1080/00379271.2007.10697499

Beasley, D. A. E., Bonisoli-Alquati, A., & Mousseau, T. A. (2013). The use of fluctuating asymmetry as a measure of environmentally induced developmental instability: A meta-analysis. *Ecological Indicators*, *30*, 218–226. https://doi.org/10.1016/j.ecolind.2013.02.024

Chown, S. L., & Gaston, K. J. (2010). Body size variation in insects: a macroecological perspective. Biological Reviews, 85(1), 139-169. https://doi.org/10.1111/j.1469-185X.2009.00097.x

Buchholz, S., & Egerer, M. H. (2020). Functional ecology of wild bees in cities: Towards a

better understanding of trait-urbanization relationships. *Biodiversity and*

Conservation, *29*(9), 2779–2801. https://doi.org/10.1007/s10531-020-02003-8

Chun, B., & Guldmann, J.-M. (2018). Impact of greening on the urban heat island: Seasonal

variations and mitigation strategies. *Computers, Environment and Urban Systems*, *71*,

165–176. https://doi.org/10.1016/j.compenvurbsys.2018.05.006

Colla, S. R., Gadallah, F., Richardson, L., Wagner, D., & Gall, L. (2012). Assessing declines

of North American bumble bees (Bombus spp.) using museum specimens.

Biodiversity and Conservation, *21*(14), 3585–3595. https://doi.org/10.1007/s10531-

012-0383-2

Costa, M., Mateus, R. P., & Moura, M. O. (2015). Constant fluctuating asymmetry but not

directional asymmetry along the geographic distribution of *Drosophila antonietae*

(Diptera, Drosophilidae). *Revista Brasileira de Entomologia*, *59*(4), 337–342.

https://doi.org/10.1016/j.rbe.2015.09.004

Couvillon, M. J., & Dornhaus, A. (2010). Small worker bumble bees (*Bombus impatiens*) are

hardier against starvation than their larger sisters. *Insectes Sociaux*, *57*(2), 193–197.

https://doi.org/10.1007/s00040-010-0064-7

Couvillon, Margaret J., Jandt, J. M., Duong, N., & Dornhaus, A. (2010). Ontogeny of worker

body size distribution in bumble bee (Bombus impatiens) colonies. *Ecological*

Entomology, *35*(4), 424–435. https://doi.org/10.1111/j.1365-2311.2010.01198.x

Couvillon, Margaret J. & Dornhaus, A. (2009) Location, location, location: larvae position

inside the nest is correlated with adult body size in worker bumble bees (*Bombus*

impatiens). Proceedings of the Royal Society B, 276, 2411–2418.

de Farias-Silva, F. J., & Freitas, B. M. (2020). Thermoregulation in the large carpenter bee

Xylocopa frontalis in the face of climate change in the Neotropics. *Apidologie*.

https://doi.org/10.1007/s13592-020-00824-8

Dellicour, S., Gerard, M., Prunier, J. G., Dewulf, A., Kuhlmann, M., & Michez, D. (2017).

Distribution and predictors of wing shape and size variability in three sister species of

solitary bees. *PLOS ONE*, *12*(3), e0173109.

https://doi.org/10.1371/journal.pone.0173109

Eggenberger, H., Frey, D., Pellissier, L., Ghazoul, J., Fontana, S., & Moretti, M. (2019).

Urban bumblebees are smaller and more phenotypically diverse than their rural

counterparts. *Journal of Animal Ecology*, *88*(10), 1522–1533.

https://doi.org/10.1111/1365-2656.13051

Fernández, M. J., Driver, M. E., & Hedrick, T. L. (2017). Asymmetry costs: Effects of wing

damage on hovering flight performance in the hawkmoth *Manduca sexta*. *The Journal*

of Experimental Biology, *220*(20), 3649–3656. https://doi.org/10.1242/jeb.153494

Fisher, K., Gonthier, D. J., Ennis, K. K., & Perfecto, I. (2017). Floral resource availability

from groundcover promotes bee abundance in coffee agroecosystems. Ecological

Applications, 27(6), 1815-1826.

Földesi, R., Howlett, B. G., Grass, I., & Batáry, P. (2020). Larger pollinators deposit more

pollen on stigmas across multiple plant species—A meta□analysis. Journal of

Applied Ecology. https://doi.org/10.1111/1365-2664.13798

Gérard, M., Martinet, B., Maebe, K., Marshall, L., Smagghe, G., Vereecken, N. J., Vray, S.,

Rasmont, P., & Michez, D. (2020). Shift in size of bumblebee queens over the last

century. *Global Change Biology*, *26*(3), 1185–1195.

https://doi.org/10.1111/gcb.14890

Gérard, M., Vanderplanck, M., Franzen, M., Kuhlmann, M., Potts, S. G., Rasmont, P., Schweiger, O., & Michez, D. (2018a). Patterns of size variation in bees at a continental scale: Does Bergmann's rule apply? *Oikos*, *127*(8), 1095–1103. https://doi.org/10.1111/oik.05260

Gérard, M., Michez, D., Debat, V., Fullgrabe, L., Meeus, I., Piot, N., Sculfort, O., Vastrade, M., Smagghe, G., & Vanderplanck, M. (2018b). Stressful conditions reveal decrease in size, modification of shape but relatively stable asymmetry in bumblebee wings. *Scientific Reports*, *8*(1), 15169. https://doi.org/10.1038/s41598-018-33429-4

Combined effects of global change pressures on animal-mediated pollination. *Trends in Ecology & Evolution*, *28*(9), 524–530. https://doi.org/10.1016/j.tree.2013.05.008

Goulson, D. (2010). *Bumblebees: Behaviour, Ecology, and Conservation*. OUP Oxford.

Goulson, D., Peat, J., Stout, J. C., Tucker, J., Darvill, B., Derwent, L. C., & Hughes, W. O. H. (2002). Can alloethism in workers of the bumblebee, Bombus terrestris, be explained in terms of foraging efficiency? *Animal Chown, S. L., & Gaston, K. J. (2010). Body size variation in insects: a macroecological perspective. Biological Reviews, 85(1), 139-169.Behaviour*, *64*(1), 123–130. https://doi.org/10.1006/anbe.2002.3041

Greenleaf, S. S., Williams, N. M., Winfree, R., & Kremen, C. (2007). Bee foraging ranges and their relationship to body size. *Oecologia*, *153*(3), 589–596. https://doi.org/10.1007/s00442-007-0752-9

Grilli, M. G., Lambertucci, S. A., Therrien, J.-F., & Bildstein, K. L. (2017). Wing size but not wing shape is related to migratory behavior in a soaring bird. *Journal of Avian Biology*, *48*(5), 669–678. https://doi.org/10.1111/jav.01220

Harder, L. D. (1985). Morphology as a Predictor of Flower Choice by Bumble Bees. *Ecology*, *66*(1), 198–210. https://doi.org/10.2307/1941320

Hoffmann, A. A., Collins, E., & Woods, R. (2002). Wing Shape and Wing Size Changes as

Indicators of Environmental Stress in Helicoverpa punctigera (Lepidoptera:

Noctuidae) Moths: Comparing Shifts in Means, Variances, and Asymmetries.

Environmental Entomology, *31*(6), 965–971. https://doi.org/10.1603/0046-225X-

31.6.965

Hoffmann, A. A., Woods, R. E., Collins, E., Wallin, K., White, A., & McKenzie, J. A.

(2005). Wing shape versus asymmetry as an indicator of changing environmental

conditions in insects. *Australian Journal of Entomology*, *44*(3), 233–243.

https://doi.org/10.1111/j.1440-6055.2005.00469.x

Huchler, K., Schulze, C. H., Gamauf, A., & Sumasgutner, P. (2020). Shifting Breeding

Phenology in Eurasian Kestrels Falco tinnunculus: Effects of Weather and

Urbanization. *Frontiers in Ecology and Evolution*, *8*.

https://doi.org/10.3389/fevo.2020.00247

Karra, S., Malki-Epshtein, L., & Neophytou, M. (2011). The Dispersion of Traffic Related

Pollutants Across a Non-Homogeneous Street Canyon. *Procedia Environmental*

Sciences, *4*, 25–34. https://doi.org/10.1016/j.proenv.2011.03.004

Kelemen, E. P., Cao, N., Cao, T., Davidowitz, G., & Dornhaus, A. (2019). Metabolic rate

predicts the lifespan of workers in the bumble bee Bombus impatiens. *Apidologie*,

50(2), 195–203. https://doi.org/10.1007/s13592-018-0630-y

Kerr, E. D., Phelan, C., & Woods, H. A. (2013). Subtle direct effects of rising atmospheric

CO_2 on insect eggs. *Physiological Entomology*, *38*(4), 302–305.

https://doi.org/10.1111/phen.12034

Klingenberg, C. P. (2011). MorphoJ: An integrated software package for geometric

morphometrics. *Molecular Ecology Resources*, *11*(2), 353–357.

https://doi.org/10.1111/j.1755-0998.2010.02924.x

Klingenberg, C. P. (2015). Analyzing Fluctuating Asymmetry with Geometric

Morphometrics: Concepts, Methods, and Applications. *Symmetry*, *7*(2), 843–934.

https://doi.org/10.3390/sym7020843

Klingenberg, C. P., Badyaev, A. V., Sowry, S. M., & Beckwith, N. J. (2001). Inferring

Developmental Modularity from Morphological Integration: Analysis of Individual

Variation and Asymmetry in Bumblebee Wings. *The American Naturalist*, *157*(1),

11–23. https://doi.org/10.1086/317002

Knight, M. E., Martin, A. P., Bishop, S., Osborne, J. L., Hale, R. J., Sanderson, R. A., &

Goulson, D. (2005). An interspecific comparison of foraging range and nest density of

four bumblebee (Bombus) species. *Molecular Ecology*, *14*(6), 1811–1820.

https://doi.org/10.1111/j.1365-294X.2005.02540.x

Kölliker☐Ott, U. M., Blows, M. W., & Hoffmann, A. A. (2003). Are wing size, wing shape

and asymmetry related to field fitness of Trichogramma egg parasitoids? *Oikos*,

100(3), 563–573. https://doi.org/10.1034/j.1600-0706.2003.12063.x

Leonard, R. J., Wat, K. K. Y., McArthur, C., & Hochuli, D. F. (2018). Urbanisation and wing

asymmetry in the western honey bee (Apis mellifera, Linnaeus 1758) at multiple

scales. *PeerJ*, *6*, e5940. https://doi.org/10.7717/peerj.5940

Martinet, B., Dellicour, S., Ghisbain, G., Przybyla, K., Zambra, E., Lecocq, T., Boustani, M.,

Baghirov, R., Michez, D., & Rasmont, P. (2020). Global effects of extreme

temperatures on wild bumblebees. *Conservation Biology*.

https://doi.org/10.1111/cobi.13685

Merckx, T., Souffreau, C., Kaiser, A., Baardsen, L. F., Backeljau, T., Bonte, D., Brans, K. I.,

Cours, M., Dahirel, M., Debortoli, N., De Wolf, K., Engelen, J. M. T., Fontaneto, D.,

Gianuca, A. T., Govaert, L., Hendrickx, F., Higuti, J., Lens, L., Martens, K., … Van

Dyck, H. (2018). Body-size shifts in aquatic and terrestrial urban communities. *Nature*, *558*(7708), 113–116. https://doi.org/10.1038/s41586-018-0140-0

Mpho, M., Callaghan, A., & Holloway, G. J. (2002). Temperature and genotypic effects on life history and fluctuating asymmetry in a field strain of Culex pipiens. *Heredity*, *88*(4), 307–312. https://doi.org/10.1038/sj.hdy.6800045

Nooten, S. S., & Rehan, S. M. (2020). Historical changes in bumble bee body size and range shift of declining species. *Biodiversity and Conservation*, *29*(2), 451–467. https://doi.org/10.1007/s10531-019-01893-7

Ottosen, T.-B., & Kumar, P. (2020). The influence of the vegetation cycle on the mitigation of air pollution by a deciduous roadside hedge. *Sustainable Cities and Society*, *53*, 101919. https://doi.org/10.1016/j.scs.2019.101919

Outomuro, D., & Johansson, F. (2011). The effects of latitude, body size, and sexual selection on wing shape in a damselfly. Biological Journal of the Linnean Society, 102(2), 263-274. https://doi.org/10.1111/j.1095-8312.2010.01591.x

Palmer, A. R., & Strobeck, C. (2003). CH 17. Fluctuating asymmetry analyses revisited. *Developmental Instability: Causes and Consequences*, Oxford University Press, Oxford, 279-319.

Peters, M. K., Peisker, J., Steffan☐Dewenter, I., & Hoiss, B. (2016). Morphological traits are linked to the cold performance and distribution of bees along elevational gradients. *Journal of Biogeography*, *43*(10), 2040–2049. https://doi.org/10.1111/jbi.12768

Phillips, B. B., Gaston, K. J., Bullock, J. M., & Osborne, J. L. (2019). Road verges support pollinators in agricultural landscapes, but are diminished by heavy traffic and summer cutting. *Journal of Applied Ecology*, *56*(10), 2316–2327. https://doi.org/10.1111/1365-2664.13470

Polce, C., Maes, J., Rotllan-Puig, X., Michez, D., Castro, L., Cederberg, B., Dvorak, L.,

Fitzpatrick, Ú., Francis, F., Neumayer, J., Manino, A., Paukkunen, J., Pawlikowski,

T., Roberts, S., Straka, J., & Rasmont, P. (2018). Distribution of bumblebees across

Europe. *One Ecosystem*, 1. http://dx.doi.org/10.3897/oneeco.3.e28143

Radmacher, S., & Strohm, E. (2010b). Factors affecting offspring body size in the solitary

bee Osmia bicornis (Hymenoptera, Megachilidae). *Apidologie*, *41*(2), 169–177.

https://doi.org/10.1051/apido/2009064

Rohlf, F. (2015). The tps series of software. *Hystrix, the Italian Journal of Mammalogy*,

26(1). https://doi.org/10.4404/hystrix-26.1-11264

Sascha, B., & Egerer, M. H. (2020). Functional ecology of wild bees in cities: Towards a

better understanding of trait-urbanization relationships. *Biodiversity & Conservation*,

29(9–10), 2779–2801. http://dx.doi.org/10.1007/s10531-020-02003-8

Schneider, S. S., Leamy, L. J., Lewis, L. A., & DeGrandi☐Hoffman, G. (2003). The

Influence of Hybridization Between African and European Honeybees, Apis

Mellifera, on Asymmetries in Wing Size and Shape. *Evolution*, *57*(10), 2350–2364.

https://doi.org/10.1111/j.0014-3820.2003.tb00247.x

Sibly, R. M., & Atkinson, D. (1994). How Rearing Temperature Affects Optimal Adult Size

in Ectotherms. *Functional Ecology*, *8*(4), 486–493. https://doi.org/10.2307/2390073

Soule, A. J., Decker, L. E., & Hunter, M. D. (2020). Effects of diet and temperature on

monarch butterfly wing morphology and flight ability. *Journal of Insect*

Conservation, *24*(6), 961–975. https://doi.org/10.1007/s10841-020-00267-7

Steffan-Dewenter, I., Münzenberg, U., & Tscharntke, T. (2001). Pollination, seed set and

seed predation on a landscape scale. *Proceedings of the Royal Society of London.*

Series   B: Biological   Sciences,   *268*(1477),   1685–1690.

https://doi.org/10.1098/rspb.2001.1737

Theodorou, P., Baltz, L. M., Paxton, R. J., & Soro, A. (2020). Urbanization is associated with

shifts in bumblebee body size, with cascading effects on pollination. *Evolutionary*

Applications, *14*(1), 53–68. https://doi.org/10.1111/eva.13087

Wang, X., Blanchet, F. G., & Koper, N. (2014). Measuring habitat fragmentation: an

evaluation of landscape pattern metrics. Methods in Ecology and Evolution, 5(7),

634-646. https://doi.org/10.1111/2041-210X.12198

Zuur, A., Ieno, E. N., Walker, N., Saveliev, A. A., & Smith, G. M. (2009). *Mixed Effects*

Models and Extensions in Ecology with R. Springer Science & Business Media.

FIGURE LEGENDS

Fig.1: Map showing the distribution of the sampling sites and the border of the different provinces included in

the study area.

Fig.2: Right forewing of *B. terrestris* with landmark locations

Fig. 3: Regression analysis of centroid size as a function of (a) temperature, and (b) floral resources . *B.*

pascuorum is represented by black dots and lines, *B. terrestris* is represented by grey triangles and lines.

Continuous lines indicate the significant relationships while dashed lines represent the non-significant ones.

Dots and triangles represent observed values. The two species were tested in separate models but represented

together to facilitate the comparison.

Fig. 4: Regression analysis of fluctuating size asymmetry as a function temperature . *B. pascuorum* is

represented by black dots and lines, *B. terrestris* is represented by grey triangles and lines. Continuous lines

indicate the significant relationships, while dashed lines represent the non-significant ones.

Fig.1

Fig.2

Fig.3

Fig.4