Effect of urbanization and its environmental stressors on the intraspecific

variation of flight functional traits in two bumblebee species

Nicola Tommasi [1], Emiliano Pioltelli [1], Paolo Biella [1], Massimo Labra [1, 2], Maurizio Casiraghi [1],

Andrea Galimberti [1, 2]*

1 -ZooplantLab, Department of Biotechnology and Biosciences, University of Milano-Bicocca,

Milano, Italy

2 -INFN Sezione di Milano Bicocca, Milano, Italy

*Correspondence: andrea.galimberti@unimib.it

15   **AUTHOR CONTRIBUTION**

17   NT, EP and AG conceived the ideas and designed methodology; NT, EP collected and

18   analysed the data; EP, NT and PB led the writing of the manuscript. All authors

19   contributed critically to the drafts and gave final approval for publication.

ABSTRACT

The way urbanization shapes the intraspecific variation of pollinator functional traits is little understood. However, this topic is relevant for investigating ecosystem services and pollinator health. Here, we studied how urbanization affects the functional traits of workers in two bumblebee species (*Bombus terrestris* and *B. pascuorum*) sampled in 37 sites along a gradient of urbanization in North Italy (an area of 1800 km^2 including the metropolitan context of Milan and other surrounding capital districts). Namely, we investigated the role played by land-use composition, configuration, temperature, flower resource abundance, and air pollutants on the variation of traits related to flight performance (i.e., body size, wing shape and size fluctuating asymmetry). These traits are relevant because they are commonly used as indicators of stress during insect development. The functional traits of the two bumblebees responded idiosyncratically to urbanization. Urban temperatures were associated with smaller wing sizes in *B. pascuorum* and with more accentuated fluctuating asymmetry of wing size in *B. terrestris*. Moreover, flower abundance correlated with bigger wings in *B. terrestris* and with less asymmetric wing sizes in *B. pascuorum*. Other traits did not vary significantly and other urban variables played minor effects. These patterns highlight that environmental stressors linked to urbanization negatively impact traits related to flight performance and development stability of these species with possible consequences on the pollination service they provide.

Overall, this study found species-specific variation patterns in syntopic taxa,

expanding our understanding about the effects of anthropic disturbance in shaping

relevant functional traits of pollinator model species.

Keywords: Functional diversity, urban heat island, pollination, urban ecosystems,

wing asymmetry, geometric morphometrics

INTRODUCTION

Widespread phenomena of urbanization are driving deep changes on landscape

features, their temperatures and pollutants, creating novel ecosystem conditions that

impact biodiversity (Foley et al., 2005; Weng et al., 2007; Wenzel et al., 2020) . Plants

and animals respond to these environmental variations by shifting their distribution

(Colla et al., 2012), phenology (Huchler et al., 2020), and/or shaping some

morphological traits considered "functional", *i.e.* relevant for their ecology, fitness

and behaviour (Alberti et al., 2017; Eggenberger et al., 2019; Nooten & Rehan, 2020).

In bees, trait variation due to environmental alteration could affect the efficiency of

the ecosystem services they provide though impacting the way they interact with

plants (Buchholz & Egerer, 2020, Biella et al., 2019). Environmental alteration could

also impact pollinator development, for example by limiting the abundance of floral

resources due to the increasing proportion of anthropized surfaces (Steffan-Dewenter

et al., 2001). This scenario, in turn, could trigger body size declines due to less food

supplied to larvae (Couvillon et al., 2010). Furthermore, landscape anthropization

could change the local climate, thus altering pollinator ecology, development and

foraging (Radmacher & Strohm, 2010). Specifically, the higher degree of cemented

"impervious" land cover that characterizes urban areas is often associated with

increasing temperatures , a phenomenon known as the "heat island effect" (Chun &

Guldmann, 2018). Observations from previous studies have strengthened the

hypothesis that pollinator insects could face a shift towards smaller body size as an

adaptation to reduce the risk of overheating while foraging in warmer conditions

(Peters et al., 2016; Gérard et al., 2018a). Considering that the worldwide steady

growth of the human population is driving a dramatic urban sprawl, new insights on

pollinator responses are necessary.

Previous studies investigated the morphological responses of pollinators to

anthropogenic pressures, mainly focusing on body size (e.g., Chown & Gaston, 2010;

Eggenberger et al., 2019; Theodorou et al., 2021). In bees, this character responds

rapidly to environmental changes (Chown & Gaston, 2010), it shows little heritability

and its variation mainly depends on the amount of food received during the larval

development (Couvillon et al., 2010). Bee size is positively correlated with the

foraging range (Greenleaf et al., 2007). Generally, larger bees show more efficient

flight performances (Harrison & Roberts, 2000), since flight muscle ratio is known to

increase with body size in flying insects (Samejima & Tsubaki, 2010). Size also

determines the metabolic rate and resource needs of adult imagos, with larger bees

having higher metabolic rate (Kelemen et al., 2019) and thus potentially being more

susceptible to shortage in floral resource availability (Couvillon & Dornhaus, 2010).

However, to date, the investigation of pollinators body size variation in anthropogenic

habitats yielded heterogeneous results. A recent study on bumblebees found bigger

workers in cities (Theodorou et al., 2021). This study speculated that such pattern is

an adaptation to longer flights for collecting resources, particularly in view of the

severe green patches fragmentation of urban landscapes (Greenleaf et al., 2007).

Conversely, a study by Eggenberger et al. (2019) found smaller bumblebee foragers in

cities. This was interpreted as an effect of both limited local resource abundance and

warmer temperature in urban areas, but the effect of these variables was not directly

tested. Given these contradicting results and different interpretations, more studies are

needed for clarifying the existing patterns of pollinator morphological responses to

urbanization.

Morphometric studies are gaining in importance for quantifying even subtle

variations in morphological traits. These variations are usually informative of stress

exposure, and thus provide information about animal population health status (Adams

et al., 2001). One of the advantages of using trait variation to measure stress is that

changes of phenotypes are detectable before an overall decrease in population

viability (Hoffmann et al., 2005). Therefore, quantifying traits variation could become

an essential practice when evaluating local and landscape-level stressors. A metric that

has grown in popularity is the fluctuating asymmetry (FA) (Klingenberg, 2001;

Beasley et al., 2013; Alves-Silva et al., 2018), defined as the presence of small,

randomly placed deviations from perfect bilateral symmetry due to the occurrence of

developmental instability, driven by exogenous environmental conditions

(Klingenberg, 2015). FA differs from another type of bilateral asymmetry, the

directional asymmetry (DA), that occurs when the two sides are steadily different with

a predictable direction to this difference. While DA has a genetic basis and therefore

could be less impacted by the environment (Palmer & Strobeck, 2003), the FA is

considered a valid proxy of stress exposure to conditions that typically occur in urban

environments (e.g., higher temperature and air pollutants) (Beasley et al., 2013). For

instance, laboratory-based studies have demonstrated that higher CO_2 level or low

temperature lead to an increase in wing FA, supporting the possible role of traffic

pollutants and climatic variation in determining developmental instability

(Klingenberg et al., 2001; Hoffmann, Collins & Woods., 2002). However,

asymmetries could be found in wing shape and/or in wing size and they even have

different responses to the same stressor type. For instance, in a recent study by Gerard

et al., 2018, variations in wing size asymmetry were observed in response to thermic

and parasitic stress while these same stressors caused no alteration in wing shape

asymmetry level. Both wing size and shape are important functional traits in

pollinators. This is because wing size is believed to be related to flight length and it

influences metabolic costs (Fernandez et al., 2017; Soule et al., 2020), while shape is

considered important for flight maneuverability (Kolliker et al., 2003, Grilli et al.,

2017).

In order to characterize the effects of urbanization and of the related

environmental stressors on pollinator insects, we quantified the morphological

variation in two species of bumblebee (*i.e.*, *Bombus pascuorum* and *B. terrestris*). We

sampled foraging workers from populations spanned across a gradient of growing

urbanization (from seminatural areas to highly urbanized sites) in Northern Italy, a

region that experienced a strong anthropogenic footprint (Perini and Magliocco 2014;

Salata 2017). The two species were selected as they are among the most common and

widespread pollinators in Europe (Pekkarinen & Teräs, 1993; Rasmont et al., 2008),

and have been largely used as model species in many studies related to the effects of

urbanization or other stressors (Eggenberger et al., 2019; Theodorou et al., 2021).

We expected to find quantitative variation in bumblebee functional traits of

body size and wing FA in response to several facets of urbanization. Firstly, we tested

associations with increased fragmentation of green patches that is often found in urban

landscapes (Li et al., 2019). We also tested responses due to environmental stressors

amplified by urbanization, such as increased temperatures (Feng et al., 2014) and

pollutants (Salahodjaev 2014), and decreased floral resource abundance (Ushimaru

2014). Regarding body size we based our survey on two alternative expectations that

emerged from previous studies, depending on the prevalent pressure acting on this

trait. First, one could expect to observe an increase in body size if green patches

fragmentation triggered an adaptation to increase foraging ranges, as suggested by

(Warzecha et al., 2016). Alternatively, a reduction in body size could arise as a way to

reduce the risk of overheating in warmer urban habitats (Maebe et al., 2021;

Pereboom & Biesmeijer, 2003) or as a consequence of limited floral resources

(Chown & Gaston, 2010). Regarding wing FA in shape and size, we expected to find

increased FA in response to higher levels of biotic and abiotic stressors that are

expected to occur in more urbanized landscapes, such as limited floral resources,

temperature, and air pollutants.

MATERIALS AND METHODS

Study species

This study was focused on two co-occurring species of bumblebee: *Bombus*

terrestris (Linnaeus 1758) and *B. pascuorum* (Scopoli 1763). Both species are

pollinators common in Europe, and can be easily found while foraging in different

habitats (Polce et al., 2018), even in urban areas (Meeus et al. 2021, Banaszak-

Cibicka, and Żmihorski 2012), including the surveyed region (personal observation of

the authors). Given these characteristics, these species are reliable models to

investigate responses by pollinating insects to landscape anthropization (Eggenberger

et al., 2019; Theodorou et al., 2021). Using two cases of different, albeit related,

model species could even clarify if the observed patterns are general or rather shaped

by different life histories. The two selected species, in fact, have slightly different

foraging ranges, with an estimated maximum of 449 and 758 m for *B. pascuorum* and

B. terrestris, respectively (Knight et al., 2005). Nesting habits are also dissimilar as *B.*

terrestris builds its nest in subterranean holes, while *B. pascuorum* on top of or

slightly beneath the soil surface (Goulson, 2010). Another important difference is

represented by their dietary regimes, since *B. pascuorum* usually have a narrower

trophic niche and a preference for deep-corolla flowers (Harder, 1985) while *B.*

terrestris is highly polylectic (Dafni et al., 2010, Biella et al., 2019).

Study design and sampling

183   Samplings were conducted at 37 sites (Fig. 1), in July 2019, between 9:00 and

184 12:00 only on days with sunny and windless weather conditions. The study sites were

185 distributed within an area of about 1800 km^2 covering four administrative provinces,

186 Milano, Monza e della Brianza, Lecco and Como in northern Italy. A minimum

187 distance between sites of 1 km was imposed to avoid the non-independence of sites

188 (Phillips et al., 2019) since it is above the maximum foraging range observed for the

189 two species (Knight et al., 2005). Study sites were selected along a gradient of

190 growing urbanization, ranging from areas highly dominated by seminatural hay

191 meadows close to forest with little urban areas nearby, to sites characterized by a high

192 degree of impervious surface (i.e., concrete, building, and asphalt). To select sampling

193 sites, impervious surfaces were mapped in a GIS software based on a regional land

194 use cartography (2018-DUSAF 6.0; https://www.dati.lombardia.it/Territorio/Dusaf-6-

195 0-Uso-del-suolo-2018/7rae-fng6) This land use cover map is available at the scale of

196 1: 10000 and was developed from AGEA orthophotos and SPOT 6/7 satellite images..

197 Sites were chosen on a visible gradient of growing impervious cover. Once identified

198 as suitable, the land use composition in the surrounding of the candidate sampling

199 sites was confirmed using satellite images. For each species, five to six worker

200 specimens were captured while foraging inside a plot of about 50 m x 50 m at each

201 site using an entomological net. After collection, the insects were stored at -80 °C

202 until further analyses.

204 **Landscape and environmental variables description**

The previously mentioned land use cartography was used to quantify landscape

urbanization around sampling sites. Through QGIS 3.10.11, a 1 km radius buffer area

was created around each site where landscape composition was evaluated arranging

DUSAF original level and sub-level of land-use classification into two macro

categories: impervious (i.e., buildings, infrastructures, roads, and cemented surface),

and seminatural land (i.e., meadows, forest and urban green spaces) (see Online

resources, Additional information 2 for a list of DUSAF codes assigned to each

grouping).

For each site the ratio between impervious and green land was computed to

quantify the intensity of urbanization. The gradient of urbanization was also described

by green habitat fragmentation, a measure of landscape configuration, that was

quantified by computing the edge density (ED), as the ratio of edge length of green

and seminatural patches over their total area (Wang et al., 2014). Other environmental

biotic and abiotic features, possibly influenced by the urbanization degree, were

considered to test for their potential effects on altering body size and wing size/shape

FA. Specifically, land surface temperature was calculated as the mean value in the

period June-July using data retrieved through remote sensing imaging

spectroradiometer (MODIS) MOD11A2 from the NASA database

(https://modis.gsfc.nasa.gov/data/dataprod/mod11.php) with a resolution of 1 km. As

the two species of bumblebee studied are characterized by a life-cycle from eggs to

adults of about two months (Goulson, 2010), the time frame for which these data were

taken into consideration should well represent the mean temperature experienced by

larvae during their development. It is important to notice that at the resolution used

here it is not possible to properly describe microclimatic variation, but rather broader

temperature variation covering the landscape scale and the foraging range of the two

bumblebee species (Knight et al., 2005). A map reporting the variation of mean

temperatures along the investigated landscape is reported in Online resources, Figure

S1. Air pollution was estimated as the mean of daily concentrations of NO_2 over two

months (June and July) registered by Regional agency for environmental protection

(ARPA), data taken from monitoring stations located nearby our sampling sites

(https://www.arpalombardia.it/Pages/Aria/qualita-aria.aspx). A map reporting the

location of monitoring stations along the investigated landscape is provided in Online

resources, Figure S1.

An expeditive estimation of floral resources at each site (i.e., the total number of

flowers) was performed by using six quadrats 1 m x 1 m (covering a proportion of

sampling area similar to that reported in Fisher et al., 2017) randomly placed in the

flowering green spaces within or closest to the sampling area, and counted the number

of flowers found there (as in Ushimaru 2014). Flowers were counted considering

single or composed inflorescences as units: for *Myosotis* sp., *Galium* sp., and *Capsella*

bursa-pastoris, and all Asteraceae the number of inflorescences was counted. The

values of the listed landscape and environmental variables in all the sampling sites are

reported in Online resources, Table S1 along with histograms showing their variation

along the sites (Additional information 1)..

Specimens imaging and wings measurement

The forewings of all individuals were detached at the base and scanned at high

resolution (i.e., 600 dpi). The obtained images were converted into TPS files using

tps-UTIL 1.74. This file format follows the standard formats for geometric

morphometrics (Rohlf, 2015). TPS file can contain two or three dimensional landmark

data and the information about the scale factor applied to each specimen is also

provided. Once converted into TPS, images were digitised using the tps-Dig 2.31

software (Rohlf, 2015), with two-dimensional cartesian coordinates of 15 landmarks

positioned at wing vein junction (Fig. 2) (as in Aytekin et al., 2007; Klingenberg et al.,

2001). Bumblebees with damaged or badly worn wings were excluded from further

analyses.

The analysis of landmark configuration was conducted in MorphoJ 1.07 software

(Klingenberg, 2011). To remove all the effects of scale, rotation and position, a

standard protocol based on a generalized least square Procrustes superimposition was

applied (Klingenberg, 2011). This strategy permits to obtain a new set of

superimposed landmark coordinates (i.e., 'Procrustes shape coordinates') describing

the wing shape and size features. Wing size was estimated as the centroid size: *i.e.*, the

square root of the sum of squared distances from the centroid of each landmark

configuration, and used as a proxy of body size (hereafter "body size", as in Outomuro

& Johansson, 2011 and Dellicour et al., 2017). To confirm the positive relation

between centroid size and body size the inter tegular distance (IT), another measure of

body size (Warzecha et al., 2016), was retrieved from a subset of 50 individuals of

each species. Afterwards, the correlation between IT and centroid size was calculated

(*B. terrestris* r = 0.7, p <0.001; *B. pascuorum* r = 0.7, p <0.001). Wing size asymmetry

was computed by dividing the absolute difference between left and right centroid sizes

by the mean centroid size and multiplying by 100 (Leonard et al., 2018). To estimate

wing shape variation, Procrustes distances were computed for each individual

(Klingenberg, 2015). These represent the measure of an individual's overall

asymmetry (*i.e.*, combining DA and FA components), obtained by taking the square

root of the sum of squared distances between corresponding right and left Procrustes'

coordinates (Klingenberg, 2015).

Statistical analysis

According to the protocol outlined in Klingenberg 2015, we first estimated the

entity of the measurement error because the levels of asymmetry in bilateral traits are

subtle and it could possibly cause considerable variation in the assessment of

asymmetry levels. This was performed by double-scanning the wings and digitizing

their landmarks for a subset of 40 specimens, with the Procrustes ANOVA in

MorphoJ (Klingenberg 2001, Klingenberg 2015) in order to evaluate the measurement

error. Afterwards, following Costa et al. (2015), to isolate the FA component, we

calculated the amount of directional asymmetry (DA) and tested its entity again with

the Procrustes ANOVA in MorphoJ considering all the measured specimens in a

single analysis. Only if DA occurred significantly, the individual asymmetry measures

were corrected by subtracting the mean DA, thus isolating the FA component as in

Costa et al. (2015).

To investigate the relationship between morphological traits and covariates, linear

mixed models were used. The responses of the two species were assessed separately.

In all the models, the ratio between impervious and seminatural surfaces was initially

included as a predictor with the other variables. However, Variance inflation factor

(VIF) criteria was used to assess the absence of collinearity among model variables

and it indicated that the ratio between impervious and seminatural surfaces was highly

collinear with the other variables, see also the correlation matrix reported in Online

resources, Table S2. Thus, we decided to exclude the ratio between impervious and

seminatural surfaces from subsequent models. The other variables, describing

landscape configuration, biotic, and abiotic features, as well as the interaction between

all these variables, were included in the models following the ecological expectations

of our hypothesis. Specifically, changes in body size were evaluated in response to the

edge density of green area, temperature, and floral resource availability because they

could directly influence body size, with bigger sizes in increasingly fragmented green

areas, and/or with more flower resources, and/or less temperatures (Warzecha et al.,

2016; Pereboom & Biesmeijer, 2003; Chown & Gaston, 2010). Concerning wing FA,

the temperature, concentration of NO_2, and flower resources limitation were included

in the models following our hypothesis that they could be stressors expected to

increase asymmetry (Hoffmann, Collins & Woods, 2002; Klingenberg et al., 2001;

Leonard et al.,2018) Sampling site was included as a random effect in all the models.

A backward stepwise model selection based on AIC was used to remove variables and

their combinations that did not improve the model fit and thus to obtain reliable final

models (Zuur et al., 2009). In order to improve the fit between the predictors and the

response variable, mathematical transformations were applied on some of the

covariates as reported in Table 1. All the analyses were performed using R (version

3.6.1; R CoreTeam 2019).

RESULTS

After excluding queen, males, and specimens presenting damaged wings, 179

B. pascuorum (mean per site = 4.8 ± 0.4) and 169 *B. terrestris* (mean per site = 4.5 ±

0.3) were subjected to morphometric analyses.

The measurement error was negligible because it was not significant for wing size (df

= 79, F = 2.67 p = 0.4578, R^2 = 0.0009) and shape (df = 2054, F = 0.51, p = 0.9976, R^2

= 0.07). Different patterns of size variation were found in the two bumblebee species.

B. terrestris body size was found to increase in response to floral resource abundance

(Fig. 3a, Table1) while *B. pascuorum* body size decreased in response to the

increasing temperature (Fig. 3 b, Table 1).

Concerning wing asymmetry, both species showed a significant level of shape DA (*B.*

pascuorum df = 26, F = 4.66; p < 0.0001, R^2 = 0.008; *B. terrestris* df = 26, F = 5.60; p

< 0.0001, R^2 = 0.009), while size DA was statistically significant only in *B.*

pascuorum (df = 1, F = 29.77; p < 0.0001, R^2 = 0.0007; in *B. terrestris* df = 1, F =

0.51; p = 0.4779, R^2 = 0.000008). Measures of FA were then obtained by correcting

for directional components.

B. terrestris size FA was positively correlated to temperature (Fig. 4 a; Table 1) while

B. pascuorum size FA decreased with higher floral resource availability (Fig 4 b;

Table 1). None of the predictor variables (i.e. temperature, NO_2, resource abundance,

and the interaction among these variables) showed a significant effect on wing shape

asymmetry in both bumblebee species (output of non significant regression models are

available in Online resources, Table S3).

DISCUSSION

In this study, we quantified the spatial intraspecific variation of functional

traits in two common bumblebee species, to infer the possible alteration of their

ecological features, such as the dispersion ability, with consequences on different

aspects of their life history.

We focused on the morphological variations of *B. terrestris* and *B. pascuorum* along

an urbanization gradient within an area of 1800 km^2 including the metropolitan

context of Milan and other surrounding capital districts. Our results highlighted some

correlations between stressors related to urbanization, and traits as body size and wing

size asymmetry. Specifically, the landscape temperature and the abundance of floral

resources, two environmental features influenced by the degree of urbanization (See

online resources Table S1, Additional information 1, and Table S2), emerged as

candidate drivers of intraspecific variation of body size across bumblebee populations,

acting differently on the two investigated species . Foragers of *B. pascuorum* showed

a shift towards smaller body size in response to increasing temperature, a condition

often associated with deeply urbanized landscapes, generally referred to as the heat

island effect. Although a similar pattern of body size reduction in urban bumblebees

has previously been reported by Eggenberger et al. (2019), they did not evaluate the

effect of temperature but only proposed it as a possible driver of the observed decrease

of body size. This relationship has been investigated on multiple historical series

collections-based and experimental studies that revealed how higher environmental

temperatures represent a driver of body size reduction in different species of bees

(e.g., Nooten & Rehan, 2020; Theodorou et al., 2021). Higher temperature accelerates

larval development, which likely results in smaller adults (Sibly & Atkinson, 1994).

Furthermore, smaller sizes in warmer areas could also be a strategy for reducing

overheating risks while foraging, due to an increased convective heat loss in smaller

bees (de Farias-Silva & Freitas, 2020). Functionally, smaller foragers could travel

shorter foraging distances (Greenleaf et al., 2007) and could also load less pollen and

nectar (Goulson et al., 2002). As a consequence, the shift towards smaller body size in

B. pascuorum could imply that it will pollinate less plants or handle flowers less

efficiently (Földesi et al., 2020), a concerning aspect in view of colony provision and

pollination. It is not known if microclimatic conditions could mitigate the effects we

observed, as we used temperature measured at a broader scale. Furthermore, it is

important to underline that other landscape variables could act synergically with urban

temperature. However, these synergic effects could be difficult to disentangle in the

field, due to the high correlation of these variables with the amount of impervious

land. Further investigations considering microclimatic variations (e.g., by using data

loggers at each sampling site) and field experiments pointing at cause-effect

relationship between temperature and body size will be required to exclude the

possible role of other urban related stressors. The role of different categories of

impervious surface (i.e., concrete, building, and asphalt) in contributing to

temperature increase should also be addressed in future research, to better inform

mitigation strategies in urban contexts.

Body size reduction was also previously explained by the decrease in floral

resource abundance in urban landscapes, possibly as a consequence of reduction of

green areas (Merckx et al., 2018; Eggenberger et al., 2019). According to this

evidence, we found a correlation between *B. terrestris* body size and the abundance of

flower resources, with larger individuals observed where more food is potentially

available. This is in accordance with the observation that adult size is strictly

correlated with the amount of food received during larval development (Couvillon &

Dornhaus 2009). However, this trend seems to be not clearly confirmed by *B.*

pascuorum probably due to a possibly higher flower specialization of this species

(Harder, 1985). Indeed the narrower diet of *B. pascuorum* could result in the inability

of this species in exploiting abundant flowering plant species that do not represent its

host plants. It seems a fruitful avenue of future research to investigate other important

features such as the nutritional quality of available resources, their diversity, and

changes along landscape variation (Vaudo et al., 2015; Vaudo et al., 2016). Here, we

considered floral resource abundance at the local scale. In spite of this limitation, we

could observe some relationships between traits and flower resources. However,

additional data will further illuminate how resources shape urban pollinator traits.

Despite *B. pascuorum* and *B. terrestris* belong to the same genus, they showed

a different susceptibility towards the investigated stressors. This suggests that

different responses are likely to come from different habits and behaviour.

Idiosyncratic responses were also observed in other bumblebee species, where body

size decreased over warming decades, but others responded in the opposite way

(Gérard et al., 2020). In our study the invariant size of *B. terrestris* in warmer

conditions could be explained by its higher heat tolerance (Martinet et al., 2020). In

addition, *B. terrestris* nests further underground compared to *B. pascuorum*, and it

might be less exposed to warm air temperatures during larval development. These

aspects strengthen the hypothesis that temperature could be a major determinant of

pollinator size reduction in cities because they affected the body size of the more

temperature-sensitive species, but not the heat-tolerant one. These idiosyncratic

species-specific responses are very relevant for understanding the potential

mechanism of intraspecific trait variation associated with urbanization and supports

the need to consider a wider panel of species in this kind of studies.

Regarding wing asymmetry we found that size FA was positively correlated

with increased temperatures in *B. terrestris*. Variation in both wing size and wing

shape asymmetry was observed in other insect taxa and the effect of temperature was

previously investigated under controlled laboratory conditions (Mpho et al., 2002).

Studies associated the increased wing size and shape FA to environmental stressors,

indicating that impairment of developmental processes might take place (e.g.,

Klingenberg et al., 2001; Kerr et al., 2013). The absence of variation in shape

asymmetry registered for both the bumblebee species could confirm the results from

other studies that have indicated shape variation as less susceptible to stressors than

size asymmetry (e.g., Gérard et al., 2018b). Importantly, floral diet could represent a

possible mitigation of environmental stressors during development (Archer et al.,

2014). Here, this view is supported by the negative correlation found between

resource abundance and wing size FA, although only in *B. pascuorum*.

433   Flight performance largely depends on body size, and is also affected by

asymmetries in shape and size between wings (Grilli et al., 2017; Soule et al., 2020).

Variation in these traits does not only show developmental instability, but also has

deep ecological implications. Indeed, body size is determinant in predicting dispersal

ability of insects, thus determining their foraging range (Greenleaf et al. 2007).

Similarly, wing size FA impacts the management of lengthy flights (Fernandez et al.,

2017, Soule et al., 2020), while wing shape FA is often associated with flight

maneuverability. The combination of these morphological changes could deeply

impact flight performance, flight length in time and space, and consequently

bumblebee foraging (Kenna, Pawar & Gill., 2021), with potential consequences on

their pollination efficiency. However, an important aspect to consider is that wing size

and its asymmetry could even determine behavioural changes. For example, insects

could increase visitation rates at closer distances to colonies, and even spend a higher

time on the available resources instead of flying at a broader distance (Andrieu et al.,

2009). Such changes, albeit difficult to quantify in the field, could merit further

investigation when trying to forecast the impact of functional trait changes in response

to urbanization.

CONCLUSIONS

This study suggests that the environmental changes associated with urbanization could

affect different functional traits of pollinators, and that their impact occurs

heterogeneously on different insect species. Eventually, as the studied traits are often

involved in flying abilities, these responses could bring to the alarming outcome of

decreased foraging efficiency and pollination effectiveness. Furthermore, the different

responses to the same stressor among the two bumblebees underline the necessity for

future studies to consider a wider panel of taxa instead of single model species in

order to draw conclusions that could be applied to the whole pollinator insects

community.

From a conservation perspective, the comprehension of how pollinators cope with the

challenging conditions occurring in novel anthropogenic habitats, plays a key role in

informing suitable policy efforts to conserve their biodiversity and the ecosystem

service they provide. In the future, cities are predicted to expand constantly and thus

designing of urban landscapes will become a fundamental step for achieving

sustainability outcomes. The pollinator-friendly design and management of urban

green spaces will possibly create suitable conditions for pollinators and thus for the

ecosystem services they provide (Guenat et al., 2019; Tommasi et al., 2021). At the

same time, urban forestry and greenery practices (e.g., plantation of street and

residential trees and the creation of urban greenbelts or greenways) could represent a

valid solution to mitigate stressful conditions related to the urban environment, such

as the lack of floral resources and the heat island effect (Chun & Guldmann, 2018)

that here were found to influence functional traits.

ACKNOWLEDGMENTS:

The authors are grateful to Amelia Pioltelli for artworks, Giulia Masoero, Lorenzo

Guzzetti, Carota, Sam and Jemy for useful advice and support during manuscript

preparation.

DECLARATIONS:

Funding: This research was partially supported by the PIGNOLETTO project, co-

financed with the resources of POR FESR 2014-2020, European regional

development fund with the contribution of resources from the European Union, Italy

and the Lombardy Region.

Conflicts of interest/Competing interests: The authors declare that no competing

interests exist.

Ethics approval: Sampling permits were obtained when needed from local authorities

Availability of data and material: All relevant data are within the paper or stored in

a public repository (https://doi.org/10.6084/m9.figshare.13637594).

REFERENCES

Adams, S. M., Giesy, J. P., Tremblay, L. A., & Eason, C. T. (2001). The use of biomarkers in

ecological risk assessment: Recommendations from the Christchurch conference on

Biomarkers in Ecotoxicology. *Biomarkers*, *6*(1), 1–6.

https://doi.org/10.1080/135475001452724

Alberti, M., Correa, C., Marzluff, J. M., Hendry, A. P., Palkovacs, E. P., Gotanda, K. M.,

Hunt, V. M., Apgar, T. M., & Zhou, Y. (2017). Global urban signatures of phenotypic

change in animal and plant populations. *Proceedings of the National Academy of Sciences*, *114*(34), 8951–8956. https://doi.org/10.1073/pnas.1606034114

Alves-Silva, E., Santos, J. C., & Cornelissen, T. G. (2018). How many leaves are enough? The influence of sample size on estimates of plant developmental instability and leaf asymmetry. *Ecological Indicators*, *89*, 912–924. https://doi.org/10.1016/j.ecolind.2017.12.060

Andrieu, E., Dornier, A., Rouifed, S., Schatz, B., & Cheptou, P. O. (2009). The town Crepis and the country Crepis: How does fragmentation affect a plant–pollinator interaction?. acta oecologica, 35(1), 1-7.

Archer, C. R., Pirk, C. W., Wright, G. A., & Nicolson, S. W. (2014). Nutrition affects survival in A frican honeybees exposed to interacting stressors. Functional ecology, 28(4), 913-923.

Aytekin, M. A., Terzo, M., Rasmont, P., & Çağatay, N. (2007). Landmark based geometric morphometric analysis of wing shape in Sibiricobombus Vogt (Hymenoptera: Apidae: Bombus Latreille). *Annales de la Société entomologique de France (N.S.)*, *43*(1), 95–102. https://doi.org/10.1080/00379271.2007.10697499

Banaszak-Cibicka, W., & Żmihorski, M. (2012). Wild bees along an urban gradient: winners and losers. Journal of Insect Conservation, 16(3), 331-343.

Beasley, D. A. E., Bonisoli-Alquati, A., & Mousseau, T. A. (2013). The use of fluctuating asymmetry as a measure of environmentally induced developmental instability: A meta-analysis. *Ecological Indicators*, *30*, 218–226. https://doi.org/10.1016/j.ecolind.2013.02.024

Chown, S. L., & Gaston, K. J. (2010). Body size variation in insects: a macroecological perspective. Biological Reviews, 85(1), 139-169. https://doi.org/10.1111/j.1469-185X.2009.00097.x

Biella, P., Akter, A., Ollerton, J., Tarrant, S., Janeček, Š., Jersáková, J., & Klecka, J. (2019).

Experimental loss of generalist plants reveals alterations in plant-pollinator

interactions and a constrained flexibility of foraging. Scientific reports, 9(1), 1-13.

Biella, P., Tommasi, N., Akter, A., Guzzetti, L., Klecka, J., Sandionigi, A., ... & Galimberti,

A. (2019). Foraging strategies are maintained despite workforce reduction: A

multidisciplinary survey on the pollen collected by a social pollinator. PloS one,

14(11), e0224037.

Buchholz, S., & Egerer, M. H. (2020). Functional ecology of wild bees in cities: Towards a

better understanding of trait-urbanization relationships. *Biodiversity and*

Conservation, *29*(9), 2779–2801. https://doi.org/10.1007/s10531-020-02003-8

Chown, S. L., & Gaston, K. J. (2010). Body size variation in insects: a macroecological

perspective. Biological Reviews, 85(1), 139-169.

Chun, B., & Guldmann, J.-M. (2018). Impact of greening on the urban heat island: Seasonal

variations and mitigation strategies. *Computers, Environment and Urban Systems*, *71*,

165–176. https://doi.org/10.1016/j.compenvurbsys.2018.05.006

Colla, S. R., Gadallah, F., Richardson, L., Wagner, D., & Gall, L. (2012). Assessing declines

of North American bumble bees (Bombus spp.) using museum specimens.

Biodiversity and Conservation, *21*(14), 3585–3595. https://doi.org/10.1007/s10531-

012-0383-2

Costa, M., Mateus, R. P., & Moura, M. O. (2015). Constant fluctuating asymmetry but not

directional asymmetry along the geographic distribution of *Drosophila antonietae*

(Diptera, Drosophilidae). *Revista Brasileira de Entomologia*, *59*(4), 337–342.

https://doi.org/10.1016/j.rbe.2015.09.004

Couvillon, M. J., & Dornhaus, A. (2010). Small worker bumble bees (*Bombus impatiens*) are

hardier against starvation than their larger sisters. *Insectes Sociaux*, *57*(2), 193–197.

https://doi.org/10.1007/s00040-010-0064-7

Couvillon, Margaret J., Jandt, J. M., Duong, N., & Dornhaus, A. (2010). Ontogeny of worker

body size distribution in bumble bee (Bombus impatiens) colonies. *Ecological*

Entomology, *35*(4), 424–435. https://doi.org/10.1111/j.1365-2311.2010.01198.x

Couvillon, Margaret J. & Dornhaus, A. (2009) Location, location, location: larvae position

inside the nest is correlated with adult body size in worker bumble bees (*Bombus*

impatiens). Proceedings of the Royal Society B, 276, 2411–2418.

Dafni, A., Kevan, P., Gross, C. L., & Goka, K. (2010). Bombus terrestris, pollinator, invasive

and pest: An assessment of problems associated with its widespread introductions for

commercial purposes. Applied Entomology and Zoology, 45(1), 101-113.

de Farias-Silva, F. J., & Freitas, B. M. (2020). Thermoregulation in the large carpenter bee

Xylocopa frontalis in the face of climate change in the Neotropics. *Apidologie*.

https://doi.org/10.1007/s13592-020-00824-8

Dellicour, S., Gerard, M., Prunier, J. G., Dewulf, A., Kuhlmann, M., & Michez, D. (2017).

Distribution and predictors of wing shape and size variability in three sister species of

solitary   bees.   *PLOS*   *ONE*,   *12*(3),   e0173109.

https://doi.org/10.1371/journal.pone.0173109

Eggenberger, H., Frey, D., Pellissier, L., Ghazoul, J., Fontana, S., & Moretti, M. (2019).

Urban bumblebees are smaller and more phenotypically diverse than their rural

counterparts.   *Journal*   *of*   *Animal*   *Ecology*,   *88*(10),   1522–1533.

https://doi.org/10.1111/1365-2656.13051

Fernández, M. J., Driver, M. E., & Hedrick, T. L. (2017). Asymmetry costs: Effects of wing damage on hovering flight performance in the hawkmoth *Manduca sexta*. *The Journal of Experimental Biology*, *220*(20), 3649–3656. https://doi.org/10.1242/jeb.153494

Feng, H., Zhao, X., Chen, F., & Wu, L. (2014). Using land use change trajectories to quantify the effects of urbanization on urban heat island. Advances in Space Research, 53(3), 463-473.

Fisher, K., Gonthier, D. J., Ennis, K. K., & Perfecto, I. (2017). Floral resource availability from groundcover promotes bee abundance in coffee agroecosystems. Ecological Applications, 27(6), 1815-1826.

Földesi, R., Howlett, B. G., Grass, I., & Batáry, P. (2020). Larger pollinators deposit more pollen on stigmas across multiple plant species—A meta□analysis. Journal of Applied Ecology. https://doi.org/10.1111/1365-2664.13798

Foley, J. A., DeFries, R., Asner, G. P., Barford, C., Bonan, G., Carpenter, S. R., ... & Snyder, P. K. (2005). Global consequences of land use. science, 309(5734), 570-574.

Gérard, M., Martinet, B., Maebe, K., Marshall, L., Smagghe, G., Vereecken, N. J., Vray, S., Rasmont, P., & Michez, D. (2020). Shift in size of bumblebee queens over the last century. *Global Change Biology*, *26*(3), 1185–1195. https://doi.org/10.1111/gcb.14890

Gérard, M., Vanderplanck, M., Franzen, M., Kuhlmann, M., Potts, S. G., Rasmont, P., Schweiger, O., & Michez, D. (2018a). Patterns of size variation in bees at a continental scale: Does Bergmann's rule apply? *Oikos*, *127*(8), 1095–1103. https://doi.org/10.1111/oik.05260

Gérard, M., Michez, D., Debat, V., Fullgrabe, L., Meeus, I., Piot, N., Sculfort, O., Vastrade, M., Smagghe, G., & Vanderplanck, M. (2018b). Stressful conditions reveal decrease

in size, modification of shape but relatively stable asymmetry in bumblebee wings.

Scientific Reports, *8*(1), 15169. https://doi.org/10.1038/s41598-018-33429-4

Goulson, D. (2010). *Bumblebees: Behaviour, Ecology, and Conservation*. OUP Oxford.

Goulson, D., Peat, J., Stout, J. C., Tucker, J., Darvill, B., Derwent, L. C., & Hughes, W. O. H.

(2002). Can alloethism in workers of the bumblebee, Bombus terrestris, be explained

in terms of foraging efficiency? *Animal Chown, S. L., & Gaston, K. J. (2010). Body*

size variation in insects: a macroecological perspective. Biological Reviews, 85(1),

139-169.Behaviour, *64*(1), 123–130. https://doi.org/10.1006/anbe.2002.3041

Greenleaf, S. S., Williams, N. M., Winfree, R., & Kremen, C. (2007). Bee foraging ranges

and their relationship to body size. *Oecologia*, *153*(3), 589–596.

https://doi.org/10.1007/s00442-007-0752-9

Grilli, M. G., Lambertucci, S. A., Therrien, J.-F., & Bildstein, K. L. (2017). Wing size but not

wing shape is related to migratory behavior in a soaring bird. *Journal of Avian*

Biology, *48*(5), 669–678. https://doi.org/10.1111/jav.01220

Guenat, S., Kunin, W. E., Dougill, A. J., & Dallimer, M. (2019). Effects of urbanisation and

management practices on pollinators in tropical Africa. Journal of Applied Ecology,

56(1), 214-224.

Harder, L. D. (1985). Morphology as a Predictor of Flower Choice by Bumble Bees. *Ecology*,

66(1), 198–210. https://doi.org/10.2307/1941320

Harrison, J. F., & Roberts, S. P. (2000). Flight respiration and energetics. Annual review of

physiology, 62(1), 179-205.

Hoffmann, A. A., Collins, E., & Woods, R. (2002). Wing Shape and Wing Size Changes as

Indicators of Environmental Stress in Helicoverpa punctigera (Lepidoptera:

Noctuidae) Moths: Comparing Shifts in Means, Variances, and Asymmetries.

Environmental Entomology, *31*(6), 965–971. https://doi.org/10.1603/0046-225X-

31.6.965

Hoffmann, A. A., Woods, R. E., Collins, E., Wallin, K., White, A., & McKenzie, J. A.

(2005). Wing shape versus asymmetry as an indicator of changing environmental

conditions in insects. *Australian Journal of Entomology*, *44*(3), 233–243.

https://doi.org/10.1111/j.1440-6055.2005.00469.x

Huchler, K., Schulze, C. H., Gamauf, A., & Sumasgutner, P. (2020). Shifting Breeding

Phenology in Eurasian Kestrels Falco tinnunculus: Effects of Weather and

Urbanization. *Frontiers in Ecology and Evolution*, *8*.

https://doi.org/10.3389/fevo.2020.00247

Karra, S., Malki-Epshtein, L., & Neophytou, M. (2011). The Dispersion of Traffic Related

Pollutants Across a Non-Homogeneous Street Canyon. *Procedia Environmental*

Sciences, *4*, 25–34. https://doi.org/10.1016/j.proenv.2011.03.004

Kelemen, E. P., Cao, N., Cao, T., Davidowitz, G., & Dornhaus, A. (2019). Metabolic rate

predicts the lifespan of workers in the bumble bee Bombus impatiens. *Apidologie*,

50(2), 195–203. https://doi.org/10.1007/s13592-018-0630-y

Kenna, D., Pawar, S., & Gill, R. J. (2021). Thermal flight performance reveals impact of

warming on bumblebee foraging potential. Functional Ecology.

Kerr, E. D., Phelan, C., & Woods, H. A. (2013). Subtle direct effects of rising atmospheric

CO_2 on insect eggs. *Physiological Entomology*, *38*(4), 302–305.

https://doi.org/10.1111/phen.12034

Klingenberg, C. P. (2011). MorphoJ: An integrated software package for geometric

morphometrics. *Molecular Ecology Resources*, *11*(2), 353–357.

https://doi.org/10.1111/j.1755-0998.2010.02924.x

Klingenberg, C. P. (2015). Analyzing Fluctuating Asymmetry with Geometric

Morphometrics: Concepts, Methods, and Applications. *Symmetry*, *7*(2), 843–934.

https://doi.org/10.3390/sym7020843

Klingenberg, C. P., Badyaev, A. V., Sowry, S. M., & Beckwith, N. J. (2001). Inferring

Developmental Modularity from Morphological Integration: Analysis of Individual

Variation and Asymmetry in Bumblebee Wings. *The American Naturalist*, *157*(1),

11–23. https://doi.org/10.1086/317002

Knight, M. E., Martin, A. P., Bishop, S., Osborne, J. L., Hale, R. J., Sanderson, R. A., &

Goulson, D. (2005). An interspecific comparison of foraging range and nest density of

four bumblebee (Bombus) species. *Molecular Ecology*, *14*(6), 1811–1820.

https://doi.org/10.1111/j.1365-294X.2005.02540.x

Kölliker☐Ott, U. M., Blows, M. W., & Hoffmann, A. A. (2003). Are wing size, wing shape

and asymmetry related to field fitness of Trichogramma egg parasitoids? *Oikos*,

100(3), 563–573. https://doi.org/10.1034/j.1600-0706.2003.12063.x

Leonard, R. J., Wat, K. K. Y., McArthur, C., & Hochuli, D. F. (2018). Urbanisation and wing

asymmetry in the western honey bee (Apis mellifera, Linnaeus 1758) at multiple

scales. *PeerJ*, *6*, e5940. https://doi.org/10.7717/peerj.5940

Li, F., Zheng, W., Wang, Y., Liang, J., Xie, S., Guo, S., ... & Yu, C. (2019). Urban green

space fragmentation and urbanization: A spatiotemporal perspective. Forests, 10(4),

333.

Maebe, K., Hart, A. F., Marshall, L., Vandamme, P., Vereecken, N. J., Michez, D., &

Smagghe, G. (2021). Bumblebee resilience to climate change, through plastic and

adaptive responses. Global Change Biology.

Martinet, B., Dellicour, S., Ghisbain, G., Przybyla, K., Zambra, E., Lecocq, T., Boustani, M.,

Baghirov, R., Michez, D., & Rasmont, P. (2020). Global effects of extreme

temperatures on wild bumblebees. *Conservation Biology*.

https://doi.org/10.1111/cobi.13685

Meeus, I., Parmentier, L., Pisman, M., de Graaf, D. C., & Smagghe, G. (2021). Reduced nest

development of reared Bombus terrestris within apiary dense human-modified

landscapes. Scientific reports, 11(1), 1-9.

Merckx, T., Souffreau, C., Kaiser, A., Baardsen, L. F., Backeljau, T., Bonte, D., Brans, K. I.,

Cours, M., Dahirel, M., Debortoli, N., De Wolf, K., Engelen, J. M. T., Fontaneto, D.,

Gianuca, A. T., Govaert, L., Hendrickx, F., Higuti, J., Lens, L., Martens, K., … Van

Dyck, H. (2018). Body-size shifts in aquatic and terrestrial urban communities.

Nature, *558*(7708), 113–116. https://doi.org/10.1038/s41586-018-0140-0

Mpho, M., Callaghan, A., & Holloway, G. J. (2002). Temperature and genotypic effects on

life history and fluctuating asymmetry in a field strain of Culex pipiens. *Heredity*,

88(4), 307–312. https://doi.org/10.1038/sj.hdy.6800045

Nooten, S. S., & Rehan, S. M. (2020). Historical changes in bumble bee body size and range

shift of declining species. *Biodiversity and Conservation*, *29*(2), 451–467.

https://doi.org/10.1007/s10531-019-01893-7

Ottosen, T.-B., & Kumar, P. (2020). The influence of the vegetation cycle on the mitigation

of air pollution by a deciduous roadside hedge. *Sustainable Cities and Society*, *53*,

101919. https://doi.org/10.1016/j.scs.2019.101919

Outomuro, D., & Johansson, F. (2011). The effects of latitude, body size, and sexual selection

on wing shape in a damselfly. Biological Journal of the Linnean Society, 102(2), 263-

274. https://doi.org/10.1111/j.1095-8312.2010.01591.x

Palmer, A. R., & Strobeck, C. (2003). CH 17. Fluctuating asymmetry analyses

revisited. *Developmental Instability: Causes and Consequences*, Oxford University

Press, Oxford, 279-319.

Pekkarinen, A., & Teräs, I. (1993, January). Zoogeography of Bombus and Psithyrus in northwestern Europe (Hymenoptera, Apidae). In Annales Zoologici Fennici (pp. 187-208). Finnish Zoological Publishing Board, formed by the Finnish Academy of Sciences, Societas Biologica Fennica Vanamo, Societas pro Fauna et Flora Fennica, and Societas Scientiarum Fennica.

Pereboom, J. J. M., & Biesmeijer, J. C. (2003). Thermal constraints for stingless bee foragers: the importance of body size and coloration. Oecologia, 137(1), 42-50.

Perini, K., & Magliocco, A. (2014). Effects of vegetation, urban density, building height, and atmospheric conditions on local temperatures and thermal comfort. Urban Forestry & Urban Greening, 13(3), 495-506.

Peters, M. K., Peisker, J., Steffan□Dewenter, I., & Hoiss, B. (2016). Morphological traits are linked to the cold performance and distribution of bees along elevational gradients. *Journal of Biogeography*, *43*(10), 2040–2049. https://doi.org/10.1111/jbi.12768

Phillips, B. B., Gaston, K. J., Bullock, J. M., & Osborne, J. L. (2019). Road verges support pollinators in agricultural landscapes, but are diminished by heavy traffic and summer cutting. *Journal of Applied Ecology*, *56*(10), 2316–2327. https://doi.org/10.1111/1365-2664.13470

Polce, C., Maes, J., Rotllan-Puig, X., Michez, D., Castro, L., Cederberg, B., Dvorak, L., Fitzpatrick, Ú., Francis, F., Neumayer, J., Manino, A., Paukkunen, J., Pawlikowski, T., Roberts, S., Straka, J., & Rasmont, P. (2018). Distribution of bumblebees across Europe. *One Ecosystem*, 1. http://dx.doi.org/10.3897/oneeco.3.e28143

Radmacher, S., & Strohm, E. (2010b). Factors affecting offspring body size in the solitary bee Osmia bicornis (Hymenoptera, Megachilidae). *Apidologie*, *41*(2), 169–177. https://doi.org/10.1051/apido/2009064

Rasmont, P., Coppée, A., Michez, D., & De Meulemeester, T. (2008, January). An overview

of the Bombus terrestris (L. 1758) subspecies (Hymenoptera: Apidae). In Annales de

la Société entomologique de France (Vol. 44, No. 2, pp. 243-250). Taylor & Francis

Group.

Rohlf, F. (2015). The tps series of software. *Hystrix, the Italian Journal of Mammalogy*,

26(1). https://doi.org/10.4404/hystrix-26.1-11264

Salahodjaev, R. (2014). Over-urbanization and air pollution: a cross-country analysis.

International Economics Letters, 3(2), 69-75.

Salata, S. (2017). Land use change analysis in the urban region of Milan. Management of

Environmental Quality: An International Journal.

Samejima, Y., & Tsubaki, Y. (2010). Body temperature and body size affect flight

performance in a damselfly. Behavioral Ecology and Sociobiology, 64(4), 685-692.

Sascha, B., & Egerer, M. H. (2020). Functional ecology of wild bees in cities: Towards a

better understanding of trait-urbanization relationships. *Biodiversity & Conservation*,

29(9–10), 2779–2801. http://dx.doi.org/10.1007/s10531-020-02003-8

Schneider, S. S., Leamy, L. J., Lewis, L. A., & DeGrandi☐Hoffman, G. (2003). The

Influence of Hybridization Between African and European Honeybees, Apis

Mellifera, on Asymmetries in Wing Size and Shape. *Evolution*, *57*(10), 2350–2364.

https://doi.org/10.1111/j.0014-3820.2003.tb00247.x

Sibly, R. M., & Atkinson, D. (1994). How Rearing Temperature Affects Optimal Adult Size

in Ectotherms. *Functional Ecology*, *8*(4), 486–493. https://doi.org/10.2307/2390073

Soule, A. J., Decker, L. E., & Hunter, M. D. (2020). Effects of diet and temperature on

monarch butterfly wing morphology and flight ability. *Journal of Insect*

Conservation, *24*(6), 961–975. https://doi.org/10.1007/s10841-020-00267-7

Steffan-Dewenter, I., Münzenberg, U., & Tscharntke, T. (2001). Pollination, seed set and seed predation on a landscape scale. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *268*(1477), 1685–1690. https://doi.org/10.1098/rspb.2001.1737

Theodorou, P., Baltz, L. M., Paxton, R. J., & Soro, A. (2021). Urbanization is associated with shifts in bumblebee body size, with cascading effects on pollination. *Evolutionary Applications*, *14*(1), 53–68. https://doi.org/10.1111/eva.13087

Tommasi, N., Biella, P., Guzzetti, L., Lasway, J. V., Njovu, H. K., Tapparo, A., ... & Galimberti, A. (2021). Impact of land use intensification and local features on plants and pollinators in Sub-Saharan smallholder farms. Agriculture, Ecosystems & Environment, 319, 107560.

Ushimaru, A., Kobayashi, A., & Dohzono, I. (2014). Does urbanization promote floral diversification? Implications from changes in herkogamy with pollinator availability in an urban-rural area. The American Naturalist, 184(2), 258-267.

Vaudo, A. D., Tooker, J. F., Grozinger, C. M., & Patch, H. M. (2015). Bee nutrition and floral resource restoration. Current opinion in insect science, 10, 133-141.

Vaudo, A. D., Patch, H. M., Mortensen, D. A., Tooker, J. F., & Grozinger, C. M. (2016). Macronutrient ratios in pollen shape bumble bee (Bombus impatiens) foraging strategies and floral preferences. Proceedings of the National Academy of Sciences, 113(28), E4035-E4042.

Wang, X., Blanchet, F. G., & Koper, N. (2014). Measuring habitat fragmentation: an evaluation of landscape pattern metrics. Methods in Ecology and Evolution, 5(7), 634-646. https://doi.org/10.1111/2041-210X.12198

Warzecha, D., Diekötter, T., Wolters, V., & Jauker, F. (2016). Intraspecific body size

increases with habitat fragmentation in wild bee pollinators. Landscape Ecology,

31(7), 1449-1455.

Weng, Y. C. (2007). Spatiotemporal changes of landscape pattern in response to

urbanization. Landscape and urban planning, 81(4), 341-353.

Wenzel, A., Grass, I., Belavadi, V. V., & Tscharntke, T. (2020). How urbanization is driving

pollinator diversity and pollination–A systematic review. Biological Conservation,

241, 108321.

Zuur, A., Ieno, E. N., Walker, N., Saveliev, A. A., & Smith, G. M. (2009). *Mixed Effects*

Models and Extensions in Ecology with R. Springer Science & Business Media.

Table 1: Output of Linear mixed models of body size (N= 348) and fluctuating size asymmetry (size

FA) (N= 347) of each species as a function of biotic and abiotic covariates of urbanization, with site

identity as random factor. Final models were selected through backward stepwise selection using AIC

criterion. ΔAIC reports the difference in AIC values between full and final models. βi: regression

coefficient; χ2: chi square values; df: degrees of freedom. Models and results of shape FA are reported

in Online resources, Table S3 as they were non significant.

Species	Response variable	Full model covariates	Final model covariates	ΔAIC	Bi	χ2; df	p value
Bombus terrestris	Body size	Temperature T Edge density ED log (Floral resources FL) Interaction T x ED x FL	log (Floral resources)	16.5	0.025	6.610;1	**0.010**
B. pascuorum	Body size	Temperature T Edge density ED log (Floral resources FL) Interaction T x ED x FL	Temperature	17.9	-0.003	7.403;1	**0.006**
B. terrestris	Size FA	Temperature T log (Floral resources FL) NO₂ N Interaction Tx FL x N	Temperature	23.5	0.052	7.183	**0.007**
B. pascuorum	Size FA	Temperature T log (Floral resources FL) NO₂ N Interaction Tx FL x N	log (Floral resources)	30.3	-0.161	6.118;1	**0.013**

FIGURE LEGENDS

Fig.1: Map showing the distribution of the sampling sites along the urbanization gradient. White areas

correspond to cemented surfaces.

Fig.2: Right forewing of *B. terrestris* with landmark locations used in this study. Details on the formulas applied

to calculate centroid size and consequently fluctuating asymmetry from these landmarks are reported in the

manuscript section "Specimens imaging and wings measurement" and in the references within.

Fig. 3: Body size variations as a function of (a) Floral resource abundance in *B.terrestris* and (b) Summer

temperature in *B. pascuorum*. The black line and grey areas indicate the relationship and its confidence intervals

as estimated with Linear mixed models, see methods for further details.

Fig. 4: Variations in wing size Fluctuating Asymmetry (FA) as a function of (a) Summer temperature in

B.terrestris and (b) Floral resource abundance in *B. pascuorum*. The black line and grey areas indicate the

relationship and its confidence intervals as estimated with Linear mixed models, see methods for further details.

Fig.1

Fig.2

Fig.3

Fig. 4