Pseudomonas can prevent the parasitic fungus, while keeping the crop fungus unaffected, in

the gardens of _Odontotermes obesus_.

Renuka Agarwal[1], Manisha Gupta[1], Abin Antony[1], Ruchira Sen[2], Rhitoban Raychoudhury[1*]

1- Department of Biological Sciences, Indian Institute of Science and Education Research Mohali

(IISER Mohali). Knowledge City, Sector 81, SAS Nagar, Manauli PO 140306, Punjab, India.

2- Sri Guru Gobind Singh College, Sector 26, Chandigarh 160019, India.

* Corresponding Author: rhitoban@iisermohali.ac.in

Keywords: symbiosis, _Termitomyces_, _Pseudoxylaria_, bacteria, pan-microbiome, core-microbiome

Abstract

Insects that farm monocultures of fungi are canonical examples of nutritional symbiosis as well as

independent evolution of agriculture in non-human animals. But just like in human agriculture,

these fungal crops face constant threat of invasion by weeds which, if unchecked, takes over the

crop fungus. In fungus-growing termites, the crop fungus (*Termitomyces*) faces such challenges

from the parasitic fungus *Pseudoxylaria*. The mechanism by which *Pseudoxylaria* is suppressed

is not known. However, evidence suggests that some bacterial secondary symbionts can serve as

defensive mutualists by preventing the growth of *Pseudoxylaria*. However, such secondary

symbionts must possess the dual, yet contrasting, capabilities of suppressing the weedy fungus

while keeping the growth of the crop fungus unaffected. This study describes the isolation,

identification and culture-dependent estimation of the roles of several such putative defensive

mutualists from the colonies of the wide-spread fungus-growing termite from

India, *Odontotermes obesus*. From the 38 bacterial cultures tested, a strain of *Pseudomonas*

showed significantly greater suppression of the weedy fungus than the crop fungus. Moreover, a

16S rRNA pan-microbiome survey, using the Nanopore platform, revealed *Pseudomonas* to be a

part of the core microbiota of *Odontotermes obesus*. A meta-analysis of microbiota composition

across different species of *Odontotermes* also confirms the wide-spread prevalence

of *Pseudomonas* within this termite. These evidence indicate that *Pseudomonas* could be playing

the role of defensive mutualist within *Odontotermes*.

Introduction

Domestication of specific fungal cultivars by insects, for food, has evolved independently over the

past 50 million years in three major insect lineages of ants, termites, and beetles [1-3]. Curiously,

fungus-growing ants and termites show parallel symbiotic evolution in two distinct geographical

locations. The former is found in the new world, whereas, the latter (Subfamily: Macrotermitinae)

are found in tropical Africa and Asia [4]. However, both systems face the invasion of parasitic

fungi taking over their crop. In fungus-growing termites, the most prominent of these parasites is

Pseudoxylaria (Phylum: Ascomycota) which invade the monoculture of the crop fungus,

Termitomyces (Phylum: Basidiomycota) [2, 5-7]. Termites have been reported to regulate this

invasion by actively grooming out any unwanted spores [8], selectively burying the parasitic fungi

[9] and applying chemical secretions with specific anti-fungal properties [10]. However, the

effectiveness of these measures in controlling *Pseudoxylaria* remains unknown [4, 11].

The presence of fungicide-producing mutualistic microbes in fungus-growing ants, which

arguably suppresses the growth of parasitic fungi in their colonies [12-15], points to a similar role

of secondary bacterial symbionts in fungus-growing termites as well [16, 17]. This conjecture is

strengthened by the fact that even after cultivating very divergent crop fungi, the microbiota across

all insect-fungal systems shows a remarkable phylogenetic and compositional similarity [18, 19].

This multi-level convergent evolution suggests the importance of such secondary microbes in

maintaining successful nutritional symbiosis across all fungus-growing insects [18, 19]. The

evolutionary logic for the presence of such mutualistic microbes within these symbioses is

inescapable. Since, the termites cannot subsist on any fungi other than *Termitomyces* [20, 21] and

Pseudoxylaria has been found in almost all fungus-growing termites [22], colonies with such

mutualistic microbes would have a strong selective advantage. Consequently, many strains of

Bacillus and *Streptomyces* have been found from termite colonies which can act against

Pseudoxylaria [11, 23-26]. But, most of these strains also inhibit the growth of the cultivar fungus

[23, 24].

This indicates that such a secondary symbiont cannot be a generic anti-fungal agent as it will also

adversely affect the crop fungus on which the termites subsist. Thus, a potential mutualist must

have the dual, but contrasting, capability of acting as an antagonist towards *Pseudoxylaria*, but, as

a mutualist towards *Termitomyces*. Unfortunately, there is no clear and unambiguous evidence of

such a microbe from fungus-growing termites. Moreover, the presence of such microbes has not

been demonstrated across the core microbiota of individual termite populations as these surveys

are either from the gut or from the comb [11, 27]. But if these mutualists are indeed essential then

they must be present within the different castes as well as the fungus comb of the termite. This is

especially true for winged reproductives (alates) which are the mobile castes of termites that move

out of the natal colony to find new ones. Alates are presumed to be equipped with the full

complement of the microbiota required for the successful establishment of new colonies and would

be selected against if they lack such critical secondary mutualists [28, 29]. Thus, the presence of

any such a secondary symbiont should permeate across the different castes as well as the comb.

However, since no such analysis of caste-specific microbiota has been reported from any fungus-

growing termites, it is not possible to ascertain whether the previously reported microbes are an
essential part of the symbiosis.

In this study, we present evidence of the presence of several potential mutualistic bacteria within
the core microbial community of the fungus-growing termite *Odontotermes obesus*. We isolated
38 different bacterial cultures from different castes and the comb and tested their potential to be
an antagonist towards *Pseudoxylaria.* Then we tested some of these antagonists to ascertain
whether they also impede the growth of the crop fungus. A particular strain of *Pseudomonas*
appears from these two contrasting screens to be an ideal mutualist as it strongly impedes the
growth of *Pseudoxylaria* but show almost negligible inhibition of *Termitomyces.* To test whether
Pseudomonas is a part of the symbiotic community, we also enumerated the pan-microbiome of
O. obesus by obtaining Nanopore reads of 16S rRNA gene fragments (V3-V4 regions) from six
different samples (comb, both male and female alates, both major and minor workers and nymphal
castes). From this pan-microbiome, we then established the presence of *Pseudomonas* within the
core microbiota of *O. obesus*. Finally, we performed a meta-analysis of microbiota across many
different *Odontotermes* sp. and ascertain the ubiquity *Pseudomonas* in this termite genus.

Materials and Methods

Source of termites, combs and their DNA extraction

Different castes, nymphs and fungus combs of *O. obesus* were collected from two different
mounds of the IISER Mohali campus (Fig. S1-S3). The combs were collected in sterile plastic
bags, termites were removed and then crushed within them. These were divided into 0.5 g portions
in sterile glass containers. To minimize environmental contamination and degradation, DNA was
extracted from some of these portions within 3 hours of the collection while others were used for
microbial isolation. Major workers, minor workers and nymphs collected from these combs were
also kept in separate sterile vials. Male and female alates were collected during swarming, which
began after the first rains during June 2018. DNA was extracted from individual termite using the
CTAB (Cetyl Trimethyl Ammonium Bromide) method [30] (supplementary methods S1).

DNA extractions from fungus combs often have extensive humic acid contamination which can
hinder downstream PCR reactions. To remove this humic acid, the precipitation step with Phenol:
Chloroform: Isoamyl alcohol were repeated until the final pellet obtained was white in color [11].

Isolation and identification of bacterial strains

Workers, nymphs, and alates were washed in the sterilized water to clean them of soil particles and then homogenized in 500 µl of sterile 1X PBS buffer with sterile plastic pestles. 0.5 gm portions of comb were also similarly homogenized. A dilution series ($1x - 10^{-6}x$) of these homogenates were then plated on Luria Bertani Agar (LBA) and Tryptic Soya Agar (TSA) for any bacterial growth at 30°C for 24 hrs. For strain identification, DNA from individual bacterial colonies was extracted with CTAB buffer and amplified with the primer set 341F/806R [31] for a portion of the 16S rRNA gene (supplementary methods S1). The sequences thus obtained were used to identify the bacterial strains by a BLAST search in NCBI. These were submitted to GenBank (accession number MN908295- MN908332). Since, multiple strains within each bacterial genus were obtained, the nomenclature of these strains were modified to read as 'Genera-NCBI Accession Number' (Table S1).

Bacterial- fungal interactions

Bacterial growth curves for all the 38 isolated strains were determined at 30°C using Freedom EVO (Tecan Life Sciences) to ensure that interaction with the fungi was done when they were in their respective log phases. To identify whether these bacteria prevent the growth of the parasitic fungus, paper disc diffusion assay was used in 90 mm diameter Petri dishes containing Potato Dextrose Agar media (PDA) [23]. Paper discs soaked with 10 µl of bacterial cultures were inoculated at the center of the Petri dish along with actively growing *Pseudoxylaria* in two plugs. Control growth plates were set up for both the bacteria and fungus and incubated in dark at 30°C in a Heratherm™ Compact Microbiological Incubator (Thermo Fisher Scientific). The magnitude of inhibition by bacteria was evaluated by using the following formula from Royse and Ries [32]:

Magnitude of Inhibition = (Total Area of fungal growth in the control in mm^2) – (Total Area of fungal growth in the interaction in mm^2) / Total Area of fungal growth in the control in mm^2

This assay was slightly modified for interaction against *Termitomyces*, as it grows much more slowly than *Pseudoxylaria*. While *Pseudoxylaria* plugs can fill up a control Petri plate within 5 days, *Termitomyces* plugs can take over 30 days to do so (Fig. S4). To account for this disparity, a liquid suspension of *Termitomyces* nodules and mycelia was prepared by homogenizing a 4 cm^2 block of actively growing culture in sterile 1X PBS and spread on plates with Potato-Dextrose

with Yeast Malt (PYME) Agar [33]. These plates were first incubated for 36 hours at 30˚C, before

adding bacterial discs [24]. Another modification was the use of four bacterial discs against

Termitomyces compared to a single bacterial disc against *Pseudoxylaria*. This enabled the

establishment of contact between *Termitomyces* and the bacterial strains much faster than would

have been possible with the use of single discs. These two modifications standardized the growing

conditions of the two fungi so that final data could be taken on the 7th day after inoculation. All

the interaction plates were evaluated for any type of inhibition every 24 hours for 7 days post-

inoculation by taking photographs using the Panasonic Lumix G2 camera (Fig. S5). Photographs

from the 7th day were counted as the final data and were analyzed in Adobe Photoshop CS6 by

measuring the area of fungal growth in pixels and converting it into millimeters (100 pixels = 1

mm^2). All interaction assays were repeated with sample sizes ranging from 3-5 plates.

Representative figures for all the bacterial-fungal interaction assays are given in supplementary

figures (Fig. S6-S8).

Sample preparation, running the Nanopore platform and obtaining sequences

To identify the microbiota present within termite castes and fungus combs, the V3-V4 region of

the 16S rRNA gene was amplified using primers 341F/806R [31]. Six different DNA samples

(major worker, minor worker, nymph, male alate, female alate and comb) were prepared for pan-

microbiome analysis on the Nanopore platform (supplementary methods S1).

Samples were sequenced on a MinION platform (MinION Mk1B) on FLO-MIN106 flowcells for

48 hours using MinKNOW software with the protocol *NC_48Hr_sequencing_FLO-*

MIN106_SQK-LSK108_plus_Basecaller. After the completion of the run, sequences were

separated and trimmed according to their barcodes and quality (supplementary methods S1).

Cleaned sequences were deposited to NCBI Sequence Read Archive (SRA) under the BioProject

PRJNA608773 (BioSample accessions: SAMN14208512 - SAMN14208517).

Taxonomic identification of the pan-microbiome

A customized microbial repository was made by combining all the available 16S rRNA gene

fragments from NCBI FTP site and the DictDb v 3.0 database [34]. LAST v 973 [35], was used to

identify the sequences with the following parameters: match score of 1, gap opening penalty of 1,

and gap extension penalty of 1 [36]. To estimate the sequencing depth rarefaction curves were

generated with the Vegan package v 2.5-4 in R v 3.6.2. The estimated community richness (Chao1)

and diversity indices (Evenness, Shannon, Simpson, and Inv Simpson) [37-39] were also
calculated in R.

We compared the gut and comb microbiota from the other known *Odontotermes* sp. [31, 40, 41]
with the results obtained from this study using weighted principle coordinate analysis (PCoA) in
phyloseq package of R. Nineteen different microbiota were used for the similarity analysis as these
also amplified the same 16S rRNA gene region and used the same reference database.

Results

Eight different bacterial strains prevent the growth of the *Pseudoxylaria*

38 different bacterial cultures were obtained from the termites and the comb. These were then
tested for their ability to prevent the growth of *Pseudoxylaria*. Four different kinds of interactions
were found: clear zone of inhibition, reduced growth of fungus near the bacteria, contact inhibition,
and negligible inhibition [13]. As fig. 1 (left panel) indicates, eight different bacteria showed
significant capability to prevent the growth of *Pseudoxylaria*.

A clear zone of inhibition was seen in the interaction between *Pseudoxylaria* and both the strains
of *Pseudomonas*, as well as *Bacillus*-MN908304 and *Streptomyces*-MN908327 (Fig. 2, Fig. S6).
Whereas, against *Bacillus*-MN908305, *Pseudoxylaria* showed reduced growth near the bacteria
(Fig. 2 and Fig. S6). Against three other bacteria (*Bacillus*-MN908297, *Bacillus*-MN908298 and
Burkholderia-MN908310) *Pseudoxylaria* showed contact inhibition (Fig. 2, Fig. S6). For all the
rest of the 30 bacterial cultures, no discernible reduction of fungus growth was observed as it grew
over the bacterial colonies (Fig. S7).

Thus, the eight strains of bacteria were found to be capable of preventing the growth of the parasitic
fungus in culture. As fig. 1 (right panel) indicates there is hardly any phylogenetic similarity across
these bacterial strains with this ability, indicating, that this capability might have evolved
independently or different modes of prevention of fungal growth are employed by these bacteria.

Five bacterial cultures show greater inhibition of the parasitic fungus than the crop fungus

These eight strains were then tested against the crop fungus, to test whether these show generic
anti-fungal properties and also impedes the growth *Termitomyces* (Fig. S8). As fig. 2 indicates
these bacteria show variable degrees of inhibition of *Termitomyces*. However, when compared

with the magnitude of inhibition against both the crop and parasitic fungi, it was clear that *Bacillus*-

MN908297, *Bacillus*-MN908304 and *Burkholderia*-MN908310 behaved as a generalist fungal

growth inhibitor (Fig. 2) and affected the crop fungus to a greater degree than the parasitic fungus.

The remaining five bacterial strains showed variable levels of inhibition against the crop fungus

but the magnitude of the inhibition was lower as compared to *Pseudoxylari*a. Of particular interest

were the two *Pseudomonas* strains obtained as *Pseudomonas*-MN908321 showed the highest

magnitude of inhibition against *Pseudoxylaria* but also showed inhibition, albeit to a lesser degree,

against *Termitomyces*. But *Pseudomonas*-MN908322 showed high levels of inhibition of

Pseudoxylaria as well as almost negligible levels of inhibition against *Termitomyces*. Thus,

Pseudomonas-MN908322 emerged as an excellent candidate for the role of a mutualist. The

inhibition by bacterial strains was not due to lack of nutrition and this was confirmed by re-growing

Pseudoxylaria and *Termitomyces* on used media (supplementary methods S1).

Thus, these two rounds of selection proved that the microbial communities of *O. obesus* can harbor

multiple bacteria which have the potential to be a defensive mutualist and *Pseudomonas*-

MN908322 emerged as the best candidate.

Pseudomonas is a part of the core microbiota of *O. obesus*

The pan-microbiome of *O. obesus* was established from two types of workers (major and minor),

nymphal castes, alates (male and female) and the comb. Over 2.6×10^5 high-quality Nanopore

reads were obtained from the six different samples (Table 1) with the highest number of reads

from male alates (~1.1×10^5 reads) and the lowest from minor workers (~0.03×10^5 reads). The

rarefaction curves (Fig. 3) showed exhaustive sampling of three different samples (comb, major

worker and male alate) indicating sufficient coverage. However, the number of reads obtained for

minor workers and nymph samples were low (Fig. 3). The number of bacterial OTUs identified

ranged between 975 (in comb) to 212 (minor worker). The entire pan-microbiome of *O. obesus*

had 1045 unique genera (Fig. 4, Table S2).

To identify the core microbiota, the list of all the identified bacterial genera were parsed to find

how many of them occur in at least five out of the six different samples. We used this criterion as

little is known about caste-specific microbial variation in fungus-growing termites, especially,

whether difference in caste roles are also reflected in their microbial composition. As fig. 4

indicates the core microbiota consists of 95 different bacterial genera. *Pseudomonas* was found

across five of the six samples and was absent only from minor workers. As minor workers and
nymph samples also had the least number of reads (0.03×10^5 and 0.1×10^5 reads, respectively), the
absence of *Pseudomonas* in the former can probably be explained by the low depth of sequence
coverage (Fig. 3).

Discussion

The major goal of this study was to find whether secondary symbionts with mutualistic capabilities
are present within the bacterial community of *O. obesus* which can help in preventing parasitic
fungal invasion. We hypothesized that as the crop fungus was the only source of nutrition for these
termites, any mutualist present in the community would therefore be selected to be present in any
successful colonies of *O. obesus*. However, the critical requisite for any such putative bacterial
mutualist must be their dual, but contrasting, capability of preventing the parasitic fungus but not
the crop fungus. The 38 different bacterial cultures obtained were therefore subjected to these two
rounds of selection against the two fungi. *Pseudomonas*-MN908322 emerged as the best candidate
as it strongly inhibits *Pseudoxylaria* but not *Termitomyces*. We further show that *Pseudomonas* is
also widely present in the caste members of *O. obesus* by constructing the core microbiota
indicating that these are indeed key microbes in this symbiosis.

This is the first study reporting *Pseudomonas* as a potential mutualist in a fungus-growing termite.
This is surprising, since, *Pseudomonas* is widely known to prevent many different kinds of fungi
and is even used as a commercial antifungal agent [42]. Moreover, they are often one of the
dominant microbial genera within fungus-growing insects [19]. This study has uncovered two
strains of *Pseudomonas* which have strong inhibitory effects on *Pseudoxylaria*. However,
Pseudomonas-MN908321 also shows some inhibition of *Termitomyces*, but still less than against
Pseudoxylaria. But *Pseudomonas*-MN908322 shows negligible levels of inhibition of
Termitomyces and remains the best candidate for a mutualist. As the phylogenetic analysis shows
(Fig. 5), *Pseudomonas*-MN908321 is a close relative of *P. aeruginosa* which has known
constitutive antifungal activities [43]. However, *Pseudomonas*-MN908322 clusters with *P.*
plecoglossicida, *P. fluorescens* and *P. putida*. Curiously, *P. putida* can act as a positive enhancer
of fungal growth, where it can help augment the mycelial growth of many different fungi within
Basidiomycota [43]. It is significant to note that the crop fungus, *Termitomyces*, also belongs to
the phylum Basidiomycota.

Our study also confirms the importance of *Bacillus* and *Streptomyces* as potential defensive

mutualists in fungus-growing termites. The role of *Bacillus* is well documented in fungus-growing

termites [11, 23] where some bacillaene- producing strains can act as a selective inhibitor of

Pseudoxylaria [23]. Four strains of *Bacillus* (MN908297, MN908298, MN908304 and

MN908305) obtained in this study show substantial growth limiting capabilities against

Pseudoxylaria but what is also significant is the other nine different *Bacillus* strains do not. Thus,

even within a specific genus of bacteria this capability to prevent the parasitic fungus shows

phenotypic variability. This finding agrees with previous studies [11, 23] where different *Bacillus*

strains showed a wide variance in their ability to prevent the parasitic fungus.

Streptomyces can also provide chemical defense by producing specific inducible secondary

metabolites in fungus-growing insects [44, 45]. A wide range of chemicals produced by

Streptomyces can not only act as antifungal agents but can also show antibacterial properties in

some insects [46].

The genus *Odontotermes* is one of the most speciose and wide-spread of all fungus-growing

termites with around 390 species spread across Africa and Tropical Asia [47]. Therefore,

identifying the roles of the different secondary symbionts within *Odontotermes* colonies can be a

key to understand the evolutionary origins of this symbiosis. This study presents an extensive

analysis of the different bacterial genera found within O. *obesus* colonies. To test how

representative these communities are across different species of *Odontotermes*, a comparison was

done with the data obtained in this study with previously published data from nineteen different

species of *Odontotermes* [31, 40, 41]. The microbial abundance datasets obtained from this study

were combined from all the six different samples and a consolidated list of bacterial genera was

made for comparison. As fig. 6 shows, *O. obesus* microbial composition differs substantially from

other *Odontotermes* species. A weighted PCoA analysis of the Unifrac distances among these

samples also shows a limited clustering (Fig. S10). However, the presence of *Pseudomonas* could

be detected in fourteen of the nineteen samples (Fig. 6). This presence of *Pseudomonas* in

Odontotermes colonies, across their range, further underscores their importance to this symbiosis.

The results from this study also raise several important questions about microbial community

ecology and termite biology. First, how many different mutualistic bacterial strains are there in *O.*

obesus colonies and what are their specific roles? We obtained 38 different cultures but high-

throughput methods employed (Nanopore platform) identified at least 1045 different bacterial
genera. Thus, the existence of other bacterial mutualists cannot be ruled out. Another confounding
factor is the difficulty in identifying the number of strains present from any one particular bacterial
genus. This question is important as our results indicate that there are at least two different cultures
of *Pseudomonas*, showing varying degrees of mutualistic capabilities. Until and unless
unambiguous enumeration of the number of strains is done, it is difficult to identify specific
functions with specific secondary symbionts.

The other major question that emerges from this study is, what roles, if any, are the termites playing
in preventing the parasitic fungus. Specifically, are termites using any of these identified mutualists
in preventing *Pseudoxylaria*, or are the mutualists doing the prevention independent of the
termites? The answer must await future experiments which include both termites as well as the
identified candidate mutualists.

Acknowledgement

302 We thank Dr. Nivedita Saha, Zoological Survey of India, for identifying different termite species
303 and Nimisha E.S. for help with the fieldwork. We also thank Paras Verma for assistance with the
304 bioinformatics analysis. We acknowledge all EVOGEN lab members for their valuable comments
305 on manuscript.

Declarations

Funding

308 This work was supported by the funds from the Indian Institute of Science and Education Research
309 (IISER) Mohali. RA was supported by a graduate fellowship from IISER Mohali.

Conflicts of interest

311 The authors declare that they have no conflicts of interest.

Availability of data and material

313 The microbial dataset generated is available as supplementary material to this article. 16S rRNA
314 fragments obtained have been submitted to GenBank (accession number MN908295–MN908332)

315 and Nanopore sequencing generated have been submitted to NCBI Sequence Read Archive (SRA)

316 under the BioProject PRJNA608773 (BioSample accessions: SAMN14208512-

317 SAMN14208517).

Code availability

319 Not applicable

Authors' Contributions

321 RA and RR conceived and planned the study. RA performed field collections, sequencing,

322 microbial culturing, data collection and analysis. AA helped in fieldwork and pilot experiments.

323 Nanopore Sequencing and its analysis were done by RA and MG. The statistical analysis of

324 interaction experiments was done by RS and RA. RR and RA wrote the paper.

Figure and table legends

Fig. 1 Magnitude of inhibition (left panel) against the parasitic fungus, *Pseudoxylaria*, of the 38
bacterial cultures (right panel) obtained from *O. obesus* colonies. Whisker over horizontal bars
represent standard deviation and number indicates the type of inhibition shown by the bacterial
strains (1= clear zone of inhibition, 2= reduced growth near bacteria, 3= contact inhibition, 4=
negligible inhibition). The phylogenetic analysis was run on MEGAX with the K2+g substitution
model using *Acidobacterium* as the outgroup. ◘ represents > 75 bootstrap value. Bacterial names
in bold indicate significant inhibition of *Pseudoxylaria*. Numbers following the genera names
indicate NCBI accession number.

Fig. 2 A comparison of the magnitude of inhibition against the crop fungus (*Termitomyces*) and
the parasitic fungus (*Pseudoxylaria*) for the eight selected bacterial strains. The data for
Pseudoxylaria is same as in fig. 1 and is represented here for comparison. The right panel shows
representative pictures of *Termitomyces* and *Pseudoxylaria* interactions with the candidate
bacterial cultures with the top panel showing growths in control plates. The bacterial strains are
arranged on the basis of the magnitude of inhibition against the crop fungus with the top most
strain showing the least inhibition. Numbers on *Termitomyces* bars represent the type of inhibition
shown by the bacterial strains (1= clear zone of inhibition, 2= reduced growth near bacteria and
4= negligible inhibition).

Fig. 3 Rarefaction analysis of bacterial 16S rRNA OTUs from six different sources from the
colony of *O. obesus.*

Fig 4 Core-microbiota of *O. obesus*. Bacterial strains that were present in at least five samples of
this study are shown in the heatmap. The genera mentioned in blue were also obtained in culture.
† are the genera classified according to DictDB database.

Fig 5 A maximum likelihood phylogenetic tree of *Pseudomonas* obtained in this study (blue) with other related strains. Taxa in yellow are found in fungus-growing insects. The phylogenetic tree was constructed using partial 16S rRNA gene sequences in MEGAX with 1000 bootstrap replicates using K2+g model. The sequences from other fungus growing insects were obtained from Cardoza et al. (2009) [48], Muwawa et al. (2016) [49]. *Cellvibrio japonicus* was used as the outgroup.

Fig 6 Comparison of *O. obesus* core-microbiota with the microbiota of 19 other *Odontotermes* species to evaluate the presence of *Pseudomonas*. *Pseudomonas* was found in 15 samples. The data from 19 different *Odontotermes* sp. were compiled from Dietrich et al., 2014 [40]; Otani et al., 2014 [31] and Otani et al., 2016 [41] (numbering of the *Odontotermes* sp. reflects their original nomenclature from the studies cited). † are the genera classified according to DictDB database.

Table 1 Number of reads obtained to identify the pan-microbiome of *O. obesus*. The table also shows different diversity indices for all the samples. The details of the taxonomic distribution, as well as abundance, for all the identified reads are given in supplementary material, table S2.

References

1.   Batra SW, Batra LR (1967) The fungus gardens of insects. Scientific American. 217(5): 112-124.
2.   Mueller UG, Gerardo NM, Aanen DK, Six DL, Schultz TR (2005) The evolution of agriculture in insects. Annu. Rev. Ecol. Evol. Syst. 36: 563-595.
3.   Schultz TR, Brady SG (2008) Major evolutionary transitions in ant agriculture. Proceedings of the National Academy of Sciences. 105(14): 5435-5440.
4.   Mueller UG, Gerardo N (2002) Fungus-farming insects: multiple origins and diverse evolutionary histories. Proceedings of the National Academy of Sciences. 99(24): 15247-15249.
5.   Aanen DK, Eggleton P (2005) Fungus-growing termites originated in African rain forest. Current Biology. 15(9): 851-855.
6.   Aanen DK (2006) As you reap, so shall you sow: coupling of harvesting and inoculating stabilizes the mutualism between termites and fungi. Biology Letters. 2(2): 209-212.
7.   Batra LR, Batra SW (1966) Fungus-growing termites of tropical India and associated fungi. Journal of the Kansas Entomological Society: 725-738.
8.   Currie CR, Stuart AE (2001) Weeding and grooming of pathogens in agriculture by ants. Proceedings of the Royal Society of London. Series B: Biological Sciences. 268(1471): 1033-1039.
9.   Katariya L, Ramesh PB, Gopalappa T, Desireddy S, Bessière JM, Borges RM (2017) Fungus-farming termites selectively bury weedy fungi that smell different from crop fungi. Journal of Chemical Ecology. 43(10): 986-995.
10.   Rosengaus RB, Traniello JF, Lefebvre ML, Maxmen AB (2004) Fungistatic activity of the sternal gland secretion of the dampwood termite *Zootermopsis angusticollis*. Insectes Sociaux. 51(3): 259-264.
11.   Mathew GM, Ju YM, Lai CY, Mathew DC, Huang CC (2012) Microbial community analysis in the termite gut and fungus comb of *Odontotermes formosanus*: the implication of *Bacillus* as mutualists. FEMS Microbiology Ecology. 79(2): 504-517.
12.   Currie CR, Scott JA, Summerbell RC, Malloch D (1999) Fungus-growing ants use antibiotic-producing bacteria to control garden parasites. Nature. 398(6729): 701-704.

13. Sen R, Ishak HD, Estrada D, Dowd SE, Hong E, Mueller UG (2009) Generalized antifungal activity and 454-screening of *Pseudonocardia* and *Amycolatopsis* bacteria in nests of fungus-growing ants. Proceedings of the National Academy of Sciences. 106(42): 17805-17810.

14. Francoeur CB, May DS, Thairu M, Hoang DQ, Panthofer O, Bugni TS, Pupo MT, Clardy J, Pinto-Tomás AA, Currie CR (2021) *Burkholderia* from fungus gardens of fungus-growing ants produce antifungals that inhibit the specialized parasite *Escovopsis*. bioRxiv.

15. Santos AV, Dillon RJ, Dillon VM, Reynolds SE, Samuels RI (2004) Ocurrence of the antibiotic producing bacterium *Burkholderia* sp. in colonies of the leaf-cutting ant *Atta sexdens rubropilosa*. FEMS Microbiology Letters. 239(2): 319-323.

16. Benndorf R, Guo H, Sommerwerk E, Weigel C, Garcia-Altares M, Martin K, Hu H, Küfner M, De Beer ZW, Poulsen M Beemelmanns C (2018) Natural products from Actinobacteria associated with fungus-growing termites. Antibiotics. **7**(3): 83.

17. Kim KH, Ramadhar TR, Beemelmanns C, Cao S, Poulsen M, Currie CR, Clardy J (2014) Natalamycin A, an ansamycin from a termite-associated *Streptomyces* sp. Chemical science. 5(11): 4333-4338.

18. Aylward FO, Suen G, Biedermann PH, Adams AS, Scott JJ, Malfatti SA, del Rio TG, Tringe SG, Poulsen M, Raffa KF, Klepzig KD (2014) Convergent bacterial microbiotas in the fungal agricultural systems of insects. MBio. 5(6).

19. Barcoto MO, Carlos-Shanley C, Fan H, Ferro M, Nagamoto NS, Bacci M, Currie CR, Rodrigues A (2020) Fungus-growing insects host a distinctive microbiota apparently adapted to the fungiculture environment. Scientific Reports. 10(1): 1-13.

20. Aanen DK and Boomsma JJ (2005) Evolutionary dynamics of the mutualistic symbiosis between fungus-growing termites and *Termitomyces* fungi. Insect-fungal associations: ecology and evolution. 191-210.

21. Aanen DK, Ros VI, de Fine Licht HH, Mitchell J, De Beer ZW, Slippers B, Rouland-LeFèvre C, Boomsma JJ (2007) Patterns of interaction specificity of fungus-growing termites and *Termitomyces* symbionts in South Africa. BMC Evolutionary Biology. **7**(1): 1-11.

22. Visser AA, Kooij PW, Debets AJ, Kuyper TW, Aanen DK (2011) *Pseudoxylaria* as stowaway of the fungus-growing termite nest: Interaction asymmetry between *Pseudoxylaria*, *Termitomyces* and free-living relatives. Fungal Ecology. 4(5): 322-332.

23. Um S, Fraimout A, Sapountzis P, Oh DC, Poulsen M (2013) The fungus-growing termite *Macrotermes natalensis* harbors bacillaene-producing *Bacillus* sp. that inhibit potentially antagonistic fungi. Scientific Reports. 3(1): 1-7.

24. Visser AA, Nobre T, Currie CR, Aanen DK, Poulsen M (2012) Exploring the potential for actinobacteria as defensive symbionts in fungus-growing termites. Microbial Ecology. 63(4): 975-985.

25. Carr G, Poulsen M, Klassen JL, Hou Y, Wyche TP, Bugni TS, Currie CR, Clardy J (2012) Microtermolides A and B from termite-associated *Streptomyces* sp. and structural revision of vinylamycin. Organic letters. 14(11): 2822-2825.

26. Schwitalla JW, Benndorf R, Martin K, Vollmers J, Kaster AK, de Beer ZW, Poulsen M, Beemelmanns C (2020) *Streptomyces smaragdinus* sp. nov., isolated from the gut of the fungus growing-termite *Macrotermes natalensis*. International Journal of Systematic and Evolutionary Microbiology. 70(11): 5806.

27. Hongoh Y, Ekpornprasit L, Inoue T, Moriya S, Trakulnaleamsai S, Ohkuma M, Noparatnaraporn N, Kudo T (2006) Intracolony variation of bacterial gut microbiota among castes and ages in the fungus-growing termite *Macrotermes gilvus*. Molecular Ecology. 15(2): 505-516.

28. Diouf M, Hervé V, Mora P, Robert A, Frechault S, Rouland-Lefèvre C and Miambi E (2018) Evidence from the gut microbiota of swarming alates of a vertical transmission of the bacterial symbionts in *Nasutitermes arborum* (Termitidae, Nasutitermitinae). Antonie van Leeuwenhoek. 111(4): 573-587.

29. Meirelles LA, McFrederick QS, Rodrigues A, Mantovani JD, de Melo Rodovalho C, Ferreira H,
Bacci Jr M and Mueller UG (2016) Bacterial microbiomes from vertically transmitted fungal
inocula of the leaf-cutting ant *Atta texana*. Environmental Microbiology Reports. 8(5): 630-640.
30. Rouland-Lefevre C, Diouf MN, Brauman A, Neyra M (2002) Phylogenetic relationships in
Termitomyces (Family Agaricaceae) based on the nucleotide sequence of ITS: a first approach to
elucidate the evolutionary history of the symbiosis between fungus-growing termites and their
fungi. Molecular Phylogenetics and Evolution. 22(3): 423-429.
31. Otani S, Mikaelyan A, Nobre T, Hansen LH, Koné NG, Sørensen SJ, Aanen DK, Boomsma JJ,
Brune A, Poulsen M (2014) Identifying the core microbial community in the gut of fungus-growing
termites. Molecular Ecology. 23(18): 4631-4644.
32. Royse D, Ries S (1978) The influence of fungi isolated from peach twigs on the pathogenicity of
Cytospora cincta. Phytopathology. 68(4): 603-607.
33. Thomas R (1985) Selective medium for isolation of *Termitomyces* from termite nests. Transactions
of the British Mycological Society. 84(3): 519-526.
34. Mikaelyan A, Dietrich C, Köhler T, Poulsen M, Sillam-Dussès D, Brune A (2015) Diet is the
primary determinant of bacterial community structure in the guts of higher termites. Molecular
Ecology. 24(20): 5284-5295.
35. Frith MC, Hamada M, Horton P (2010) Parameters for accurate genome alignment. BMC
Bioinformatics. *11*(1): 1-14.
36. Shin J, Lee S, Go MJ, Lee SY, Kim SC, Lee CH, Cho BK (2016) Analysis of the mouse gut
microbiome using full-length 16S rRNA amplicon sequencing. Scientific Reports. 6: 29681.
37. Pielou EC (1966) The measurement of diversity in different types of biological collections. Journal
of Theoretical Biology. 13: 131-144.
38. Shannon CE, Weaver W (1962) The mathematical theory of communication: University of Illinois
Press.
39. Simpson EH (1949) Measurement of diversity. Nature. 163(4148): 688-688.
40. Dietrich C, Köhler T, Brune A (2014) The cockroach origin of the termite gut microbiota: patterns
in bacterial community structure reflect major evolutionary events. Applied and Environmental
Microbiology. 80(7): 2261-2269.
41. Otani S, Hansen LH, Sørensen SJ, Poulsen M (2016) Bacterial communities in termite fungus
combs are comprised of consistent gut deposits and contributions from the environment. Microbial
Ecology. 71(1): 207-220.
42. Walsh UF, Morrissey JP, and O'Gara F (2001) *Pseudomonas* for biocontrol of phytopathogens:
from functional genomics to commercial exploitation. Current Opinion in Biotechnology. 12(3):
289-295.
43. Rainey P (1991) Effect of *Pseudomonas putida* on hyphal growth of *Agaricus bisporus*.
Mycological Research. 95(6): 699-704.
44. Chevrette MG, Carlson CM, Ortega HE, Thomas C, Ananiev GE, Barns KJ, Book AJ, Cagnazzo
J, Carlos C, Flanigan W, Grubbs KJ (2019) The antimicrobial potential of *Streptomyces* from insect
microbiomes. Nature Communications. 10(1): 1-11.
45. Hulcr J, Adams AS, Raffa K, Hofstetter RW, Klepzig KD, Currie CR (2011) Presence and diversity
of *Streptomyces* in *Dendroctonus* and sympatric bark beetle galleries across North America.
Microbial Ecology. 61(4): 759-768.
46. Poulsen M, Oh DC, Clardy J, Currie CR (2011) Chemical analyses of wasp-associated
Streptomyces bacteria reveal a prolific potential for natural products discovery. PloS One. 6(2):
e16763.
47. Krishna K, Grimaldi DA, Krishna V, Engel MS (2013) Treatise on the Isoptera of the World:
Termitidae (Part One). Bulletin of the American Museum of Natural History. 2013(377): 973-1495.
48. Cardoza YJ, Vasanthakumar A, Suazo A, Raffa KF (2009) Survey and phylogenetic analysis of
culturable microbes in the oral secretions of three bark beetle species. Entomologia experimentalis
et applicata. 131(2): 138-147.

49.   Muwawa EM, Budambula N, Osiemo ZL, Boga HI, Makonde HM (2016) Isolation and
498   characterization of some gut microbial symbionts from fungus-cultivating termites (*Macrotermes*
499   and *Odontotermes* spp.). African Journal of Microbiology Research.

Magnitude of inhibition

Bacterial strains tested against *Pseudoxylaria*

Magnitude of inhibition

Against *Termitomyces* (Crop)   Against *Pseudoxylaria* (Weed)

Control *Termitomyces*   Control *Pseudoxylaria*

Against *Termitomyces*   Against *Pseudoxylaria*

Pseudomonas MN908322

Streptomyces MN908327

Bacillus MN908297

Bacillus MN908305

Bacillus MN908298

Bacillus MN908304

Pseudomonas MN908321

Burkholderia MN908310

Table 1: Number of reads obtained to identify the pan-microbiome of *O. obesus*. The table also shows different diversity indices for all the samples. The details of the taxonomic distribution, as well as abundance, for all the identified reads are given in supplementary information, table S2.

Sample source	Total reads (initial)	Total reads (post-cleanup)	Reads identified	Bacterial genera identified	Chao1	Shannon	Simpson	InvSimspon
Male Alate	1,31,070	1,11,703	87,153 (78.0%)	450	780.9	2.9	0.9	7.7
Major Worker	86,642	80,852	78,991 (97.7%)	705	1073.6	3.9	1.0	24.1
Comb	36,371	29,758	28,065 (94.3%)	975	1418.8	4.6	1.0	33.6
Female Alate	32,719	27,885	24,589 (88.2%)	368	698.1	3.4	0.9	13.8
Nymph	12,499	10,174	7,757 (76.2%)	238	376.6	2.5	0.8	4.4
Minor Worker	3,517	3,135	2,965 (94.6%)	212	422.4	4.0	1.0	30.8
	3,02,818	2,63,507		1,045 unique bacterial genera				