Innate visual attraction in wood ants is a hardwired behaviour seen across different motivational and ecological contexts

Cornelia Buehlmann & Paul Graham

School of Life Sciences, University of Sussex, Brighton BN1 9QG, UK

ABSTRACT

Wood ant foragers show an innate attraction to conspicuous visual cues. These foragers inhabit cluttered woodland habitat and feed on honeydew from aphids on trees, hence, the attraction to 'tree-like' objects might be an ecologically relevant behaviour that is tailored to wood ant foraging ecology. Foragers from other ant species with different foraging ecologies show innate attractions different to wood ants. What we do not know, however, is whether these innate visual responses vary with the ants' motivational states or caste. We thus recorded the innate visual response of wood ant foragers with different motivaional states, i.e. unfed or fed, as well as males that have a short life span and show no foraging activity. Our results show that ants from all three groups orient towards the visual cue, i.e. the wood ants' innate visual attraction is not context dependent, but a hardwired behaviour seen across different motivational and ecological contexts.

KEY WORDS

Innate visual behaviour, foraging ecology, wood ants, *Formica rufa*

INTRODUCTION

Ants cleverly combine innate and learnt navigational strategies to travel between their nest and feeding sites (Knaden & Graham 2016; Buehlmann *et al.* 2020b). Innate strategies such as path integration (e.g. (Collett *et al.* 2001; Mueller & Wehner 2010)), pheromone trails (e.g. (Harrison *et al.* 1989)), attractive food odours (Buehlmann *et al.* 2014), or innate visual responses (e.g. (Graham *et al.* 2003; Collett 2010)) are key when unfamiliar with an environment. These innate responses can structure the ants' paths, hence, act as a scaffold and facilitate the learning of information relevant for navigation.

Wood ant foragers show an innate attraction to large and conspicuous objects (Voss 1967; Graham *et al.* 2003; Buehlmann *et al.* 2020a; Buehlmann *et al.* 2020c). These foragers feed on honeydew from aphids on trees (Domisch *et al.* 2016), hence, the attraction to 'tree-like' objects might be an ecologically relevant behaviour that is tailored to wood ant foraging ecology. Foragers from other ant species with different foraging ecologies show innate attractions different to wood ants (e.g. (Heusser & Wehner 2002; Collett 2010)). Furthermore, behavioural experiments in *Drosophila* have shown that fruit flies have simple innate visual responses that seem to be tuned to the flies' behaviour requirements (Maimon *et al.* 2008; Grabowska *et al.* 2018), with small objects being aversive and large

thin objects being attractive. Aversive behaviour towards small objects potentially helps flies to avoid

collision with other flying insects or predators, whereas bar-like objects may represent attractive feeding

sites. Taken together, these different examples raise the question as to whether innate visual responses

in insects are ecologically tuned.

We investigate here if the innate visual response seen in wood ant foragers is a hardwired sensori-

motor reflex behaviour or a context dependent behaviour tuned to foraging ecology. To do so, we recorded

the innate visual response of wood ant foragers with different motivational states, i.e. unfed or fed, as well

as males that have a short life span and show no foraging activity. We found that ants from all three groups

orient towards the visual cue, i.e. the wood ants' innate visual attraction is not context dependent, but a

hardwired behaviour seen across different motivational and ecological contexts.

METHODS

Ants

Experiments were performed with laboratory kept wood ants *Formica rufa L.* collected from

Ashdown forest, East Sussex, UK. Ants were kept in the laboratory under a 12 h light: 12 h darkness

cycle at 25-27° C. Ants were fed *ad libitum* with sucrose, dead crickets and water. During the

experiments, food was limited to a minimum to increase the ants' foraging motivation, but water was

permanently available.

We recorded the innate visual response from naïve wood ants from three different groups: unfed

female foragers, fed female foragers and unfed males. To get fed foragers, foragers were fed with

sucrose before the behavioural recording. We only recorded males that walked on the platform. Males

are winged, however, they are not good flyers, i.e. most of the released males walked.

Experimental setup

Ants from these three groups were released in the centre of a circular platform (120 cm in

diameter) surrounded by a cylinder (diameter 3 m, height 1.8 m) with white walls (Figure 1A). A 20°

wide black rectangle (height: 90 cm, width: 52 cm) was placed on the inner wall of the surrounding

cylinder. As a control, additional ants from the three groups were recorded when the visual cue was

absent. Ants were only recorded once. To remove possible olfactory traces, the surface of the platform

was covered with white paper which was rotated between recordings. Further, the visual cue was rotated

between recordings to avoid the use of cues other than the black rectangle. The centre of the platform

consisted of a cylindrical holding chamber of 6.5 cm diameter, which was remotely lowered to release

the ant onto the platform. The ants' position was recorded every 20 ms using a tracking video camera

(Trackit, SciTrackS GmbH). Path were analysed in Matlab with similar analyses as done previously

(Buehlmann *et al.* 2020c).

RESULTS

Paths from unfed female foragers, fed female foragers and unfed males that show no foraging activity were recorded in the behavioural arena in the presence of a conspicuous visual object (Figure 1A). Ants from all three groups were directed (Rayleigh test; all $p \ll 0.001$) and the ants walked towards the visual cue (Figure 1B). The 95% CI for the ants' final headings overlaid with the visual cue and the groups did not differ from each other (Watson Williams tests; all $p > 0.05$). Whereas ants from all three groups approached the visual cue (Figure 1B), we observed differences in the ants' walking speed (Figure 2A) and path straightness (Figure 2B). Males walked significantly faster and straighter than fed and unfed foragers. Unfed and fed foragers differed in their walking speed but not in the path straightness (Figure 2).

In the absence of the visual cue (Figure 1C), unfed foragers were not directed (Rayleigh test; $p > 0.05$), whereas the other two groups were directed (Rayleigh test; males, $p < 0.01$; fed foragers, $p < 0.05$) but showed a very large spread in directions.

DISCUSSION

Conspicuous objects are attractive for many insects, including ants (fruit flies: (Wehner 1972; Gotz 1987; Strauss & Pichler 1998), locusts: (Wallace 1962), ladybirds: (Collett 1988), mantids: (Poteser & Kral 1995), ants: (Voss 1967; Heusser & Wehner 2002; Graham *et al.* 2003; Collett 2010; Buehlmann *et al.* 2020a; Buehlmann *et al.* 2020c)). Moreover, studies on innate visual responses of different ant species have revealed that innate visual preferences are tailored to the ants' habitat. Whereas wood ant foragers are attracted to large conspicuous objects (Voss 1967; Graham *et al.* 2003; Buehlmann *et al.* 2020a; Buehlmann *et al.* 2020c), desert ants avoid them (Heusser & Wehner 2002; Collett 2010). Wood ants (*Formica rufa*) inhabit cluttered woodland habitat where they predominantly feed on honeydew from aphids on trees (Domisch *et al.* 2016). *Cataglyphis fortis* desert ants forage on dead arthropods that are unpredictably distributed on the ground in the food-scarce terrain of the Saharan salt pans (Buehlmann *et al.* 2014). *C. fortis* avoid large objects such as bushes, potentially to avoid predators. Hence, the differences in the innate bias of wood and desert ant foragers make sense from an ecological point of view and seems to be tuned to their foraging ecology. What we do not know, however, is whether these innate visual responses vary with an ants' motivational state or caste. To test this, we recorded the innate visual behaviour of unfed foragers, fed foragers and males that have a short life span and show no foraging activity. We show here that the innate visual attraction to conspicuous objects in wood ants is not a context dependent behaviour, but a hardwired sensori-motor behaviour seen across different motivational and ecological contexts. It is possible that males are attracted to conspicuous objects in order to gain elevation to assist in dispersal. However, any role in innate visual orientation that assists in foraging would expect to be modulated by the ants' motivational state.

There are several examples of flexible innate behaviours in insects. For example, it was shown in parasitoid wasps that the insects' individual feeding state controls their innate behaviour (Waeckers 1994). Unfed wasps are attracted by flower odours and yellow targets that indicate food while fed wasps

are attracted to host odours and are not attracted by yellow colours (Waeckers 1994). Furthermore,
experiments with hawkmoths have shown that the moths' innate colour preference depends on ambient
light conditions (Kuenzinger *et al.* 2019). These moths are crepuscular and their colour preferences are
tuned to illuminance and background. This flexible behaviour allows them to successfully forage under
different light conditions. These examples show that insects are equipped with innate visual preferences,
but they keep the necessary behavioural flexibility and their behaviour is tuned to their ecology and
foraging ecology.

Given that there are many examples of flexibility, there is the question why wood ants do not
show any flexibility in their innate visual orientation. Perhaps, ant foragers rely on their multimodal
navigation toolkit to overcome potential problems with an inflexible innate visual reflex. Fed foragers
will have path integration information and potentially odour cues and visual information (Buehlmann
et al. 2020b) as they attempt to return to their nest and these sources of orientation information may be
essential in overcoming innate visual attractions that may otherwise disrupt a homeward journey.

FIGURES

Figure 1: Innate visual attraction in wood ants is seen across different motivational and ecological
contexts. (Ai) The experimental arena in which naïve ants were recorded. Circular white platform
(radius: 60 cm) is located in the centre of a cylinder (radius: 1.5 m, height: 1.8 m). A 20° wide black
rectangle (height: 90 cm, width: 52 cm) is mounted at the inner wall of the surrounding cylinder. A
camera recorded the ants' paths from above. A small door permitted access to the arena shown here
open and larger for clarity. **(Aii)** A top-down view of the arena shown in Ai. **(B)** Paths of ants released

at the centre of the arena are shown as black lines. If the data is directed, dotted arcs show 95%

confidence intervals (CIs) of the heading directions. The visual cue is shown at the platform edge instead

of on the cylinder wall. Bi: unfed female foragers; Bii: fed female foragers; Biii: unfed males. **(C)** As

in B but ants were recorded without the visual cue.

Figure 2: Walking speed and path straightness of ants approaching a visual cue with different

motivational and ecological contexts. (A) Walking speed of ants differed significantly between the

three groups (left: unfed foragers; middle: fed foragers; right: males). Kruskal Wallis with Mann

Whitney test and Bonferroni correction; unfed vs fed, $p < 0.001$; unfed vs males, $p < 0.05$; fed vs males,

$p < 0.001$. **(B)** Path straightness of males was significantly higher than of fed and unfed foragers

(Kruskal Wallis with Mann Whitney test and Bonferroni correction; unfed vs males, $p < 0.001$; fed vs

males, $p < 0.001$). There was no difference between the paths of fed and unfed foragers (Kruskal Wallis

with Mann Whitney test and Bonferroni correction; unfed vs fed, $p > 0.05$). Unfed, n = 90 ants; fed, n

= 50 ants; males, n = 108 ants. Boxplots: median, 25th and 75th percentiles (edges of the boxes) and

whiskers for extreme values not considered as outliers (circles).

ACKNOWLEDGEMENTS

The research was funded by the Biotechnology and Biological Sciences Research Council

(grant number: BB/R005036/1).

REFERENCES

1. Buehlmann, C., Aussel, A. & Graham, P. (2020a). Dynamic multimodal interactions in navigating

wood ants: What do path details tell us about cue integration? *J. Exp. Biol.*, 223, jeb221036.

2. Buehlmann, C., Graham, P., Hansson, B.S. & Knaden, M. (2014). Desert ants locate food by
combining high sensitivity to food odors with extensive crosswind runs. *Curr. Biol.*, 24, 960-
964.

3. Buehlmann, C., Mangan, M. & Graham, P. (2020b). Multimodal interactions in insect navigation.
Anim. Cogn., 23, 1129–1141.

4. Buehlmann, C., Wozniak, B., Goulard, R., Webb, B., Graham, P. & Niven, J. (2020c). Mushroom
bodies are required for learnt visual navigation but not for innate visual behaviour in ants. *Curr.*
Biol., 30, 3438-3443.

5. Collett, M. (2010). How desert ants use a visual landmark for guidance along a habitual route. *Proc.*
Natl. Acad. Sci. U. S. A., 107, 11638-11643.

6. Collett, T. (1988). How ladybirds approach nearby stalks: a study of visual selectivity and attention.
J. Comp. Physiol. A, 163, 355-363.

7. Collett, T.S., Collett, M. & Wehner, R. (2001). The guidance of desert ants by extended landmarks.
J. Exp. Biol., 204, 1635-1639.

8. Domisch, T., Risch, A.C. & Robonson, E.J.H. (2016). Wood ant foraging and mutulaism with aphids.
In: *Wood Ant Ecology and Conservation* (ed. Stockan, J, & Robinson, E. (Eds.)). Cambridge:
Cambridge University Press.

9. Gotz, K.G. (1987). Course-control, metabolism and wing interference during ultralong tethered flight
in *Drosophila melanogaster*. *J. Exp. Biol.*, 128, 35-46.

10. Grabowska, M.J., Steeves, J., Alpay, J., Van De Poll, M., Ertekin, D. & van Swinderen, B. (2018).
Innate visual preferences and behavioral flexibility in *Drosophila*. *J. Exp. Biol.*, 221,
jeb185918.

11. Graham, P., Fauria, K. & Collett, T.S. (2003). The influence of beacon-aiming on the routes of
wood ants. *J. Exp. Biol.*, 206, 535-541.

12. Harrison, J.F., Fewell, J.H., Stiller, T.M. & Breed, M.D. (1989). Effects of experience on use of
orientation cues in the giant tropical ant. *Anim. Behav.*, 37, 869-871.

13. Heusser, D. & Wehner, R. (2002). The visual centring response in desert ants, *Cataglyphis fortis*. *J.*
Exp. Biol., 205, 585-590.

14. Knaden, M. & Graham, P. (2016). The Sensory Ecology of Ant Navigation: From Natural
Environments to Neural Mechanisms. *Annu. Rev. Entomol.*, 61, 63-76.

15. Kuenzinger, W., Kelber, A., Weesner, J., Travis, J., Raguso, R.A. & Goyret, J. (2019). Innate colour
preferences of a hawkmoth depend on visual context. *Biol. Lett.*, 15, 20180886.

16. Maimon, G., Straw, A.D. & Dickinson, M.H. (2008). A simple vision-based algorithm for decision
making in flying *Drosophila*. *Curr. Biol.*, 18, 464-470.

17. Mueller, M. & Wehner, R. (2010). Path Integration Provides a Scaffold for Landmark Learning in
Desert Ants. *Curr. Biol.*, 20, 1368-1371.

18. Poteser, M. & Kral, K. (1995). Visual distance discrimination between stationary targets in praying
194   mantids: an index of the use of motion parallax. *J. Exp. Biol.*, 189, 2127-2137.

195   19. Strauss, R. & Pichler, J. (1998). Persistence of orientation toward a temporarily invisible landmark
196   in *Drosophila melanogaster*. *J. Comp. Physiol. A*, 182, 411-423.

197   20. Voss, C. (1967). Uber das Formensehen der roten Waldameise (*Formica rufa* Gruppe). *Zeitschrift
198   Fur Vergleichende Physiologie*, 55, 225-254.

199   21. Waeckers, F.L. (1994). The effect of food deprivation on the innate visual and olfactory preferences
200   in the parasitoid *Cotesia rubecula. J. Insect Physiol.*, 40, 641-649.

201   22. Wallace, G.K. (1962). Experiments on visually controlled orientation in the desert locust,
202   *Schistocerca gregaria* (Forskål). *Anim. Behav.*, 10, 361-369.

203   23. Wehner, R. (1972). Spontaneous pattern preferences of *Drosophila melanogaster* to black areas in
204   various parts of the visual field. *J. Insect Physiol.*, 18, 1531–1543.