Experimental sexual selection reveals rapid divergence in male and female reproductive

transcriptomes and their interactions

Paris Veltsos[1], Damiano Porcelli[2], Yongxiang Fang[3], Andrew R. Cossins[4], Michael G. Ritchie[5,]

Rhonda R. Snook[6*]

[1]Department of Ecology and Evolutionary Biology, University of Kansas, Lawrence, KS, USA. Orcid

0000-0002-8872-6281

[2]Department of Animal and Plant Sciences, University of Sheffield, Sheffield UK. Orcid: 0000-0002-

9019-5758

[3]CGR, Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool, Liverpool,

L69 7ZB, UK. Orcid 0000-0002-4514-5292

[4]Centre for Genomic Research, Institute for Integrative Biology, University of Liverpool, Liverpool

UK. Orcid 0000-0002-0813-5212

[5]Centre for Biological Diversity, University of St Andrews, St Andrews, Fife, Scotland KY16 9TH, UK.

Orcid 0000-0001-7913-8675

[6]Department of Zoology, Stockholm University, Stockholm 106 91 Sweden. Orcid 0000-0003-1852-

1448

*corresponding author

Abstract

Mating causes substantial changes in females, altering male and female reproductive fitness. Some postmating effects are hypothesized to be at least partially mediated by gene expression changes, driven by postcopulatory sexual selection, which results in population divergence of reproductive proteins that could generate reproductive isolation. However, understanding of the direct role of sexual selection on gene expression divergence along with the subsequent molecular mismatches that could occur between diverging populations is limited. Here, we analyze gene expression divergence following over 150 generations of experimental evolution in which *Drosophila pseudoobscura* evolved under either elevated polyandry or enforced monogamy. We find that sexual selection rapidly impacted sex-, tissue-, and mating-specific responses, and not always in the predicted direction. Postmating female responses are either unique to each sexual selection treatment or exhibit asymmetric non-congruence, in which monogamy females upregulate and polyandry females downregulate the same genes following mating. This substantial population divergence of gene expression also gives rise to either unique or mismatched gene expression patterns in crosses between treatments. Many of these genes are involved in immune and stress responses, and non-congruent responses are particularly prevalent in the female reproductive tract, the main arena for postcopulatory sexual selection. In summary, we show that sexual selection has pervasive impacts on gene expression divergence acting both differentially between reproductive tissues of the same sex and asymmetrically in postmating female responses, and this divergence is highest in the female reproductive tract, the main arena for postcopulatory sexual selection.

Keywords: postmating response, postmating prezygotic reproductive isolation, RNA-seq, testes, accessory glands, female reproductive tract.

Introduction

Sexual reproduction involves pre-ejaculatory and post-ejaculatory interactions between the sexes with sexual selection influencing male and female traits that mediate the fitness outcome of these interactions. While aspects of reproduction can be cooperative, the sexes also diverge over the optima of reproductive traits, such as courtship signals, fertilization and offspring production. The intensity of sexual selection is linked to the extent to which reproductive fitness optima differ between the sexes and can generate sexual antagonism, in which selection acts in opposing directions on the sexes (Rice, 1996; Holland and Rice, 1999). As the sexes share the majority of their genome, intra-locus sexual conflict can occur over phenotypes that are influenced by these shared genes. Resolving this conflict can occur by the evolution of sexual dimorphism, mediated by changes in the shared genome, such as sex-specific regulation of gene expression (Mank 2017). Comparative genomic studies have found that genes with rapid divergence and that show stronger signatures of positive divergent selection are often sex-biased or sex-limited in expression (e.g. Cheng *et al.* 2016; Pröschel *et al.*, 2006; Ellegren & Parsch, 2007; Zhang *et al.*, 2007). This pattern is sometimes stronger in species that have traits associated with indicators of sexual selection, such as increased sexual dimorphism (Harrison et al. 2015; Wright *et al.* 2019).

In internally fertilizing species, the main arena for post-ejaculatory molecular interactions is the female reproductive tract which includes sites of sperm transfer, storage and subsequent fertilization of eggs transiting from the ovaries. Postmating female responses are extensive, influencing female behaviour, morphology and physiology. These responses are mediated by interactions between components of the male ejaculate, including sperm produced in the testes, rapidly evolving seminal fluid proteins (Sfps) produced by accessory glands, and female reproductive proteins in the female reproductive tract and ovaries (Sirot *et al.* 2015; Oku *et al.* 2019; Wolfner 2009). Postmating changes in females include altering investment in oogenesis (Wolfner 2009), remating propensity (Chapman *et al.* 2003), and sperm storage and usage (Avila *et al.* 2010; Avila *et al.* 2015). Other aspects of physiology are also altered, for example, female hunger (Carvalho *et al.* 2006), female aggression (Bath *et al.* 2017), and homeostasis (Ribeiro and Dickson 2010; Cognigni *et al.* 2011). Genes associated with immunity and stress have altered gene expression upon mating in females, which may impact susceptibility or resistance to pathogens and/or parasites (Oku *et al.* 2019; Zhong *et al.* 2013).

An increasing number of studies have characterized transcriptomic postmating changes and used gene ontology to determine the associated gene functions, although many studies typically examine either whole bodies or abdomens of females (Lawniczak and Begun 2004; McGraw *et al.* 2004, 2008, Innocenti and Morrow 2009; Innocenti *et al.* 2014; Hollis *et al.* 2014, 2016; Delbare *et al.* 2017; Fowler *et al.* 2019) or examine only one component of the female reproductive tract ("lower" reproductive tract, defined by the female sperm storage organs, Mack *et al.* 2006,

Prokupek *et al.* 2008; or the "upper" reproductive tract, defined by the oviducts, Kapelnikov *et al.*

2008). No study has simultaneously assessed postmating effects on the sperm receiving

components of the female reproductive tract (the uterus and sperm storage organs) and the egg

production components (oviducts and ovaries). Receipt of the male ejaculate effects sperm

storage dynamics, oogenesis and oviposition (Sirot *et al.* 2015) so all are subject to sexual

selection (and sexual conflict). Likewise, for males, the main focus on the role of sexual selection

and sexual conflict has been on Sfp evolution given that they are among the most rapidly evolved

proteins known (Ahmed-Braimah *et al.* 2017; Ellegren and Parsch 2007), although sperm and the

cellular architecture of the testes also can be subject to rapid morphological evolution and sexual

selection (Lüpold *et al.* 2009).

The reproductive interactions between the sexes, by generating strong postmating sexual selection

and sexual conflict, also cause rapid evolutionary changes between lineages which may lead to

reproductive isolation (Markow 1997; Manier *et al.* 2013; Ahmed-Braimah *et al.* 2020). In particular,

postmating prezygotic (PMPZ) reproductive isolation in which gametes do not interact properly

prior to fertilization (Ahmed-Braimah *et al.* 2017; Garlovsky *et al.* 2020) or affect egg production

(Matute and Coyne 2010) may evolve due to mismatches between lineages in male ejaculate-

female reproductive tissue interactions (Ahmed-Braimah *et al.* 2020). Direct evidence of any

consequences of rapid evolutionary divergence between male and female reproductive processes

and the generation of PMPZ isolation is lacking. Some studies have examined the postmating

gene expression responses in females mating to either con- or hetero- specific males to assess

evidence for mismatches to understand how PMPZ isolation occurs (Bono *et al.* 2011; Ahmed-

Braimah *et al.* 2020). However, in most cases, these lineages are quite distinct, representing

different species, and the sexual selection history is unknown.

Experimental sexual selection, in which populations are subjected to either polyandrous conditions

that facilitates sexual selection or enforced monandrous conditions that eliminate sexual selection,

has been used to examine the evolution of gene expression responses to sexual selection, and

link these to macroevolutionary patterns of sex-biased and sex-limited gene expression evolution

(Hollis *et al.* 2014, 2016; Immonen *et al.* 2014; Veltsos *et al.* 2017). Previous studies supported the

role of sexual selection in divergent sex-biased gene expression, but whether male- or female-

biased genes responded the most depended on species (Hollis *et al.* 2014; Immonen *et al.* 2014;

Veltsos *et al.* 2017; Parker *et al.* 2019; Hollis *et al.* 2019). Additionally, these studies were limited in

that postmating responses and/or sex-specific tissue responses were rarely examined.

Consequently, the understanding of how sexual selection impacts sex-limited and reproductive

tissue-specific gene expression, the consequences of this divergence on postmating responses,

and the opportunity for this divergence to trigger barriers to reproduction is limited.

In this study, we use replicate populations of *D. pseudoobscura* following over 150 generations of

experimental evolution in which either monandry is enforced (referred to as M), eliminating sexual

selection and conflict, or elevating the opportunity for polyandry (referred to as E). We take a

quantitative transcriptomics approach to investigate the impact of sexual selection on gene

expression divergence. We separate the effects between different reproductive tissues of males

(testes and accessory glands) and females (female reproductive tract and ovaries), and examine

the postmating response within the separate female reproductive tissues to determine whether

sexual selection acts on these similarly or differently. This approach also allows us to polarize the

direction of postmating gene expression changes arising from variation in sexual selection intensity

to then compare the female postmating response when mating occurs between individuals of the

different sexual selection treatments. Previous work on these populations has documented the

action of sexual selection and sexual conflict on phenotypic reproductive interactions between the

sexes (Crudgington *et al.* 2005; 2009; 2010; Debelle *et al.* 2014; Debelle *et al.* 2016; Snook *et al.*

2005), including those that could influence postmating responses. For example, changes in

accessory gland investment (Crudgington *et al.* 2009) of males and ovariole number and

subsequent offspring production in females (Crudgington *et al.* 2010; Immonen *et al.* 2014). Sex-

biased gene expression evolution has also occurred but previous studies of expression divergence

either examined whole bodies of only one sex (Immonen *et al.* 2014) or in heads and abdomens of

each sex, but not following mating (Veltsos *et al.* 2017).

Results

Gene expression divergence in male tissues

Previous work has hypothesized that monandry selects for downregulation and polyandry selects

for upregulation of genes expressed in male reproductive tissues in the abdomen of *D.*

melanogaster (Hollis *et al.* 2016). We found that virgin male testes had 359 differentially expressed

genes, of which 161 were upregulated in polyandry (or downregulated in monandry) and 198

upregulated in monandry (or downregulated in polyandry; Figure 1a). While marginally significantly

different (χ^2 = 3.81, df = 1, p = 0.051) the gene expression pattern is in the opposite direction to

that predicted. The genes upregulated under polyandry were enriched for biological processes

(BP) related to stress responses, such as double strand break repair, cellular response to UV, and

terpenoid metabolic process (Supplementary File 1). In contrast, genes upregulated in monandry

males were related to biological processes (BP) of proteolysis, digestion and the innate immune

response (Supplementary File 1).

In male accessory glands, 80 genes were differentially expressed between sexual selection

treatments, with 34 genes upregulated in polyandry males (Figure 1a) and 46 upregulated in

monandrous males. While the number of DE genes did not differ between sexual selection

treatments (χ^2=1.8, p=0.18), the proportion of DE genes was higher among all genes expressed in

accessory glands than all genes expressed in testes (0.031 vs 0.007; x2=179.32, df=1, p<0.001).

The 34 accessory gland genes with higher expression under polyandry were enriched for BP terms

related to eggshell chorion assembly, development, and neuropeptide signaling (Supplementary

File 1). The 46 genes upregulated in monandry males were enriched for the BP term "detection of

chemical stimulus involved in sensory perception" (Supplementary File 1).

Sfps are some of the most rapidly evolving proteins, hypothesized to be in response to sexual

selection. To examine whether Sfps were more likely to be upregulated in polyandrous males

(Hollis *et al.* 2016), we compared expression levels between the sexual selection treatments of

different sets of accessory gland proteins recently described for *D. pseudoobscura* (Karr *et al.*

2019). Karr and colleagues identified 3281 proteins produced in the accessory gland (the

accessory gland "proteome"), of which 528 had protein secretory signals (the "secretome") and

163 were orthologous to *D. melanogaster* seminal fluid proteins (putative Sfps; also referred to as

the "exoproteome"). In contrast to the prediction, we found that monandry males had higher overall

expression of secretome proteins compared to polyandry males (W=21345000, p<0.001; Figure

2a) and there were no significant treatment effects on the proteome or Sfps. However, of the 80

accessory gland genes that were differentially expressed between treatments, we found a non-

significant trend towards higher expression in polyandry males (Figure 2b).

Overall, while relatively more gene expression changes were found in the accessory glands than

testes, we found a weak trend for monandry males to have higher expression in both the testes

and accessory glands relative to polyandry males, including accessory gland genes with secretory

function.

Gene expression divergence in virgin female tissues

Virgin female reproductive tracts had 120 differentially expressed genes, 66 (54) of which were

upregulated in polyandry (monoandry) females (Figure 1b), a nonsignificant difference (χ^2 = 1.2, df

= 1, p = 0.27). The 66 genes upregulated under polyandry were enriched in GO terms associated

with the immune system and egg production, such as egg activation whereas the 54 genes

upregulated under monandry were enriched for multicellular reproduction biological processes

(Supplementary File 1).

Virgin ovaries had 417 differentially expressed genes between the sexual selection treatments

(Figure 1b), with significantly more genes upregulated in polyandry (299) than in monandry (118)

(χ^2 = 78.6, df = 1, p < 0.001). The proportion of DE genes in ovaries was significantly greater than

that in FRT (0.040 vs 0.014; x2=120.1, df=1, p<0.001). Upregulated polyandry genes were

enriched in BP terms associated with eggshell chorion assembly whereas upregulated genes in

monandry ovaries were enriched for negative regulation of BMP signaling pathway

(Supplementary File 1).

Thus, we find that the female reproductive tissues do not respond similarly to the sexual selection

treatment, with the ovaries showing more gene regulation divergence in response to sexual

selection treatment than the reproductive tract. The biological processes associated with

differentially expressed genes also differed between sexual selection treatment and tissues.

Sexual selection causes divergence in the female postmating response

Sexual selection is thought to result in rapid divergence of the postmating female response. Above,

we show that, in virgins, different male and female reproductive tissues have different responses to

sexual selection which sets the stage for divergence in postmating responses. Sexual selection

treatment did not affect the total number of genes involved in the female postmating response in

the reproductive tract (n= 400 for polyandry females, n=357 for monandry females; χ^2=2.4425,

p=0.12) (Figure S1). However, polyandry females had significantly more postmating responsive

genes showing downregulation (n=246) than upregulation (n=154) after mating (χ^2=21.16,

p<0.001), whereas monandry females showed the opposite pattern, with significantly more genes

upregulated (n=233) than downregulated (n=124) after mating (χ^2=33.28, p<0.001) (Figure S1). In

contrast to the large number of genes involved in the postmating response of the female

reproductive tract, the ovaries had relatively few genes showing expression divergence upon

mating (n=23; Figure S1). Polyandry females showed significantly more differentially expressed

genes (n=22) than monandry females (n=1) (χ^2=19.174, p<0.001). Of the 22 genes involved in

postmating divergence of polyandry ovaries, 14 were upregulated after mating and 8

downregulated, not significantly different (χ^2=1.64, p=0.2; Supplementary File 1).

For the reproductive tract, genes downregulated in polyandry females after mating were

significantly enriched for metabolic related processes, eggshell chorion assembly, and immune

response, whereas downregulated genes in postmating monandry females were related to terms

associated with development (Supplementary File 1). Postmating upregulated genes in the

polyandry female reproductive tract were significantly enriched for immune and stress responses,

whereas monandry females upregulated genes related to immune response and chorion-related

functions (Supplementary File 1). For the ovaries, downregulated genes in polyandry females after

mating were significantly enriched for cold acclimation and synaptic target inhibition whereas the

one monandry downregulated gene was related to synaptic target inhibition and motor neuron

axon guidance (Supplementary File 1). Postmating upregulated genes in the polyandry female

ovaries were significantly enriched for processes involved with egg production and monandry

females did not significantly upregulate any ovarian genes after mating (Supplementary File 1).

Unique and shared postmating responses between sexual selection treatments

We compared shared and divergent postmating responses between the selection lines in the

female reproductive tract (Figure 3a) and the ovaries (Figure 3d). In the female reproductive tract,

many of the genes whose expression was altered upon mating were unique to each treatment

(blue (E) and yellow (M) dots, Figure 3a). However, there was a subset of genes that altered

expression in both sexual selection treatments (green diamonds, Figure 3a). 86 of these genes

responded congruently; that is mating resulted in both monandry and polyandry females either

upregulating (n = 67) or downregulating (n = 19) those genes (purple congruent line, Fig. 3a).

Shared upregulated genes were enriched for defense responses whereas downregulated genes

were enriched for fatty acid elongation and chitin-based cuticle development (Supplementary File

1). Intriguingly, 57 genes responding to mating were regulated non-congruently between the

sexual selection treatments; that is, gene expression was altered upon mating in both monandry

and polyandry females but the direction of the expression change was opposite in the sexual

selection treatments (red non-congruent line, Fig. 3a). Moreover, this non-congruence was

asymmetrical; only genes that were significantly upregulated after mating in monandry females, but

were downregulated after mating in polyandry females, were observed. These non-congruent

genes were related to chitin and melanin catabolic processes, chorion related processes, and

immune responses (Supplementary File 1). In contrast, the postmating response in ovaries

predominantly showed the upregulation of genes only in polyandry females and, while monandry

females did not significantly alter these genes, these E genes are congruent (Figure 3d).

Matings with non-coevolved individuals affected different genes than those of the coevolved female postmating response

Same female, different male

We use the coevolved female postmating response contrasts (Figure 3a and 3d) to polarize the

direction of change and assess whether gene expression differs in non-coevolved crosses. We first

contrast the effect of mating on females within a sexual treatment when mated to a coevolved vs

non-coevolved male ("same female, different male contrast": MM v ME and EE v EM). In the

female reproductive tract, only monogamy females show differential regulation of gene expression

dependent on male partner (number of DE genes = 19; Figure 3b, blue crosses; Figure S2). 18 of

these genes are downregulated after monandry females mate with polyandry males. However, 17

of these genes are on the non-congruent axis (Figure 3b), indicating that these genes are normally

relatively downregulated after EE mating but upregulated after an MM mating (compare blue

pluses on Fig. 3b with data in the lower right quadrant of Fig. 3a). Thus, a M female mating with an

E male downregulates postmating response genes, which is similar to the effect of E male mating

with E females. Even though this is the female reproductive tract, these genes are significantly

enriched for egg production (Supplementary file 1).

The ovarian gene expression pattern as a result of mating with a non-coevolved male (Figure 3e)

differs from the response in the female reproductive tract (Figure 3b). In the ovaries, gene

expression changes when mated to non-coevolved males occur only for E females (n=88 genes;

Figure S2). Nearly 100% of these genes were not differentially expressed upon mating in EE

crosses (n=87; compare Figure 3d with Figure 3e). 12 genes are downregulated (Figure 3e, purple

triangles), and 76 genes are upregulated (Figure 3e, yellow triangles), after an E female mates

with a non-coevolved M male. Most of the genes that are upregulated in E females after mating

with a M male are non-congruent. This means that the coevolved postmating response tends

towards upregulation in MM and downregulation in EE. Thus, mating with M males, makes E

females more M-, and less E-like. Non-coevolved upregulated genes are enriched for a variety of

disparate biological processes with molecular functions of DNA and mRNA binding

(Supplementary file 1). Non-coevolved downregulated genes are significantly enriched for rRNA

transcription and purine ribonucleoside biosynthetic processes, both of which are involved in cell

growth and proliferation (Supplementary file 1). The enriched molecular functions of these genes

are associated with structural constituents of chorion (Supplementary file 1).

Different female, same male

We then contrast the impact of male sexual selection treatment on the postmating female

response ("different female, same male contrast"; here we compare the effect of a M female

mating with an M male (MM) with a E female mating with an M male (EM); likewise we compare a

E female mating with an E male (EE) with a M female mating with an E male (ME)). A large

number of genes are differentially regulated in these contrasts but the effect of sexual selection

treatment of the male varies across female tissues. In the female reproductive tract, mating with

monandry males led to nearly 8x as many genes changing expression than mating with polyandry

males (431 vs. 58; χ^2=284.52, p<0.001; Figure S3). Ovarian responses were in the opposite

direction such that mating with polyandry males changed expression of about 2x more genes than

mating with monandry males (310 vs. 148; χ^2=57.3, p<0.001; Fig. S3). Regardless of tissue or

male sexual selection treatment, the majority of genes that changed in expression following a non-

coevolved mating were not differentially expressed in coevolved crosses.

In the female reproductive tract contrast between a coevolved MM and non-coevolved EM mating,

106 of 196 genes that showed downregulation when a E female mated to a M male (Figure 3c,

purple pluses) were not DE in coevolved postmating responses (Figure 3a). These downregulated

genes were significantly enriched for various metabolic processes, proteolysis and antimicrobial

humoral response. 213 of the 235 genes that showed relatively higher expression in E females

after mating with M males (Figure 3c, orange crosses) did not change expression in coevolved

crosses (Figure 3a). These upregulated genes were enriched for 68 different BP terms, and

subsequent analysis of cellular components and molecular functions suggest changes in genes
related to DNA replication. In the opposite cross, comparing the effects of a M female mating to an
E male relative to a EE mating, 31 genes were downregulated and 27 genes were upregulated.
Upregulated genes are enriched for immune responses and downregulated genes are enriched for
ecdysone biosynthetic process and midgut development.

In the ovaries, because there were so few coevolved changes (Figure 3d), nearly all the ovarian
gene expression changes following a non-coevolved mating with a different sexual selection line
male are uniquely altered (Figure 3f). Genes downregulated in the ovary after a E female mates
with a M male are enriched for redox process, vitellogeneis and tracheal system whereas genes
upregulated are enriched for BP processes that have cellular components associated with the P
granule and chorion. The majority of change in the ovaries were following a noncoevolved mating
between a M female and E male. The 196 genes downregulated after mating are enriched for a
small number of BP terms (eggshell chorion assembly, male meiosis 1, and rRNA transcription)
whereas the 114 genes upregulated after mating are enriched for immunity and stress related
terms.

While most non-coevolved gene expression changes were unique, the direction of change was
sometimes along the congruent axis of coevolved responses and for other genes differential
expression occurred along the non-congruent axis (Figure 3c; Figure 3f). In the female
reproductive tract, significantly more genes are DE along the non-congruent axis than the
congruent axis when M males mate with E females ($X2 = 44.73$, df = 1, P < 0.001) whereas there
is no difference between non-congruent and congruent responses in the female reproductive tract
when M females mate with E males. For the ovaries, non-coevolved matings with either a M or E
male result in more genes with differential expression that are along the congruent axis compared
to the non-congruent axis (M: $x2=10.28$, df=1, p<0.001; E: $x2=10.4$, df=1, p=0.0012).

Because non-congruent changes may be more likely to generate incompatibilities, we note
enrichment for DE genes in that axis. In the female reproductive tract, non-congruent changes
when M males mated with E females were the large number of BP terms related to DNA replication
that were upregulated after mating and for E males mated with M females associated with immune
responses, also upregulated after mating (Supplementary file 1). In the ovaries, non-congruent DE
genes when mating with M males were enriched for only two seemingly disparate biological
processes and two cellular components, 3 of which were unique terms not seen in other contrasts
with EM data. Non-congruent DE genes when mating with E males are enriched for eggshell
chorion assembly and rRNA transcription which were downregulated after mating.

Discussion

Understanding how sexual selection impacts the molecular basis of sexual interactions is important
because it can have profound effects on sex-specific fitness and is predicted to influence the
evolution of postmating prezygotic reproductive isolation. Here we combined transcriptomics with
experimental evolution to determine how sexual selection affects gene regulation simultaneously in
multiple sex-specific tissues in male and female virgins and in the postmating female response
when mating with either males that evolved in the same or different intensity of sexual selection as
the females. Our results reveal substantial gene expression divergence after only 150 generations
of altered sexual selection intensity, with sexual selection affecting components of male and
female reproductive tissues differentially, as well as alterations in gene expression specific to
mating with non-coevolved partners.

An evolutionary history of polyandry previously has been suggested to select for higher expression
of genes affecting sperm competition and/or showing antagonistic effects on female reproductive
behavior. This hypothesis has been examined in male abdomens of *D. melanogaster* testing the
effects of experimental sexual selection on gene expression of 138 Sfps (Hollis *et al.* 2016). That
study found an overall lower gene expression across all Sfps in monandry-selected males, while
no individual Sfp was differentially expressed. Here we separate the effects of sexual selection on
evolutionary divergence in both accessory gland and testes. Testes are also subject to sexual
selection via increased sperm number for male-male competition and/or in testicular architecture to
support rapid sperm production (Lüpold *et al.* 2009; Pitnick *et al.* 2008). Experimental sexual
selection can alter relative testes size and accessory gland size, although this varies between
species, between studies of the same species, and has different fitness outcomes (Hosken *et al.*
2001; Hosken and Ward 2001; Pitnick *et al.* 2001; Wigby and Chapman 2004; Linklater *et al.* 2007;
Reuter *et al.* 2008; Crudgington *et al.* 2009; Hollis *et al.* 2019). Previously we found that *D.*
pseudoobscura polyandry males had evolved larger accessory glands but not testes, and
polyandry males had greater mating capacity (Crudgington *et al.* 2009). Although we found that
accessory glands had proportionally more DE genes than testes, contrary to predictions, virgin
monandry males had more upregulated genes both in the accessory glands and testes than
polyandry males. A similar result was observed when we limited our analysis to putative *D.*
pseudoobscura Sfps (either the secretome or Sfps with *D. melanogaster* orthologs; Karr *et al.*
2019).

The male genes that changed in expression in response to sexual selection treatment were
associated with different biological processes. In the accessory glands, polyandry males had
higher expression of neuropeptides, which are diverse neuronal signaling molecules that regulate
physiological processes and behaviour in animals (Elphick *et al.* 2018). Given that postmating
female responses frequently involve changes in female reproductive physiology and behavior
(Sirot *et al.* 2015), polyandry may generally increase investment by males in the expression of
genes that profoundly alter female reproductive fitness. In contrast, monandry male accessory

gland genes were enriched for "detection of chemical stimulus involved in sensory perception" and,
while genes encompassed by this term can evolve rapidly, they have weak fitness effects (Librado
and Rozas 2016). Thus, for male responses, we see that monandry males have increased gene
expression in testes and accessory glands, including putative Sfps, although distinctly enriched
gene functions between sexual selection treatments suggests differences in the potential
consequences on female mates.

In females, the reproductive tract is considered to be the main arena for postmating sexual
interactions. However, the ovaries may also be subject to sexual antagonism given that they are
the egg-producing structure, and the sexes can be in conflict over fecundity schedules (Arnqvist
and Rowe 2005). We have previously shown that polyandry *D. pseudoobscura* females have more
ovarioles than monandry females (Immonen *et al.* 2014). Using whole body microarrays, we found
that genes upregulated in polyandry females are enriched in the ovary and associated with
oogenesis whereas monandry females upregulated genes associated with somatic tissues and
metabolism (Immonen *et al.* 2014). In virgin abdomens, RNAseq data showed sexual selection
treatment affected sex-biased gene expression (Veltsos *et al.* 2017). Here we decompose
responses between the female reproductive tract and the ovaries in virgin and mated monandry
and polyandry females. The reproductive tract showed a similar number of differentially expressed
genes between virgin polyandry and monandry females while in the ovaries more genes were
upregulated under polyandry than monandry. Genes overexpressed in virgin polyandry
reproductive tissues were associated with late stages of egg production (eggshell chorion
assembly, egg activation), whereas the genes over-expressed in monandry reproductive tissues
were associated with earlier egg production (BMP signalling pathway, involved in patterning the
Drosophila eggshell; Niepielko *et al*. 2012). Thus polyandry females appear ready ("primed") for
fast reproduction.

Related to this, previous work in *D. melanogaster* based on whole body gene expression has
suggested that polyandry females are poised for receipt of a manipulative ejaculate (Heifetz and
Wolfner 2004; McGraw *et al.* 2004), and therefore do not have as elevated a postmating response
as monandry females (Hollis *et al.* 2016). Our results partially support this prediction, dependent
on female reproductive tissue. In the female reproductive tract, polyandry females downregulated
more genes following mating than were upregulated, whereas monandry females showed the
opposite pattern. However, in the ovaries, only polyandry females changed expression with more
upregulated, than downregulated, genes and enriched for genes associated with egg production.
Overall our results support an interpretation in which polyandry females are poised to receive the
ejaculate and are more reproductively mature with respect to egg production.

Many of the postmating differentially expressed genes in the female reproductive tract are distinct
between the polyandry and monandry treatments. However, some were shared across sexual

selection treatments. Differential expression of these genes could be either congruent (both sexual selection treatments upregulated or downregulated these genes) or non-congruent, in which monandry upregulated the postmating gene expression, whereas the same gene in polyandry females was downregulated. Both upregulated and downregulated congruent genes were related to fatty acid elongation and some upregulated congruent genes were related to stress/immune responses. Fatty acids are critical to signaling pathways and have been associated with sex pheromone synthesis and with maturation of egg follicles in *D. melanogaster* (Vrablik and Watts 2013). The shared responses between sexual selection treatments suggests a shared critical postmating reproductive function. Non-congruent genes were enriched for immune response (see later) and egg production.

In polyandrous species, interactions between the sexes that impact fertility are thought to be particularly dynamic and coevolve, perhaps antagonistically. In turn, such coevolution may generate postmating prezygotic (PMPZ) incompatibilities between populations, contributing to reproductive isolation. Standard approaches in speciation research to identify reproductive isolation use crosses between males and females within and between populations. Several studies have compared postmating transcriptomic responses to test the hypothesis that misregulation between components in the male ejaculate and female reproductive tissues could generate PMPZ incompatibilities (Bono *et al.* 2011; Ahmed-Braimah *et al.* 2020). However, it is very difficult to infer the historical role of different evolutionary processes from patterns of contemporary divergence between species. Experimental evolution introduces fewer confounding variables but lacks the full realism of natural conditions. Previous experimental evolution work has been criticized for crossing different populations to infer the direction of sexual conflict (Pizzari and Snook, 2005; Rowe *et al.* 2005). Here we can use known coevolved population differentiation in gene expression to examine the consequences on mismatches in gene expression when crossed with a non-coevolved individual. The effect of mating with a non-coevolved individual depended on whether we were contrasting the effect of either the male or female treatment, what female reproductive tissue was being analyzed, and the sexual selection treatment itself. Regardless of these factors, most of the postmating genes that were differentially expressed in non-coevolved crosses were distinct from the postmating differentially expressed genes within a coevolved cross. This pattern indicates much of the non-coevolved postmating response is unique, supporting substantial and rapid coevolution between the male ejaculate and female reproductive tissues within each treatment that differs between sexual selection treatments. Such novel mismatches may have negative fitness consequences that could potentially contribute to the evolution of PMPZ reproductive isolation.

When contrasting the effect of the same female treatment when mated to different males (MMvME and EEvEM), differential expression in the female reproductive tract only occurred with M females mated to E males whereas in the ovaries, differential expression only occurred with E females mated to M males. In both cases, the differentially expressed genes were mainly non-congruent

and in such a way that the postmating response was more similar to one of the co-evolved crosses. We have previously found that when M females mate with E males, they increase the number of eggs laid earlier in their lifespan relative to E females mated to E males (Crudgington *et al.* 2010). Whether the gene expression changes in the female reproductive tract at least partially explains sexual conflict over oviposition patterns remains to be determined. In the ovaries, mismatched gene expression was related to a variety of different biological processes, including oocyte development and regulation of antimicrobial peptide biosynthesis. To get a better idea of the overall impact of this diverse set of responses, we examined the cellular component enrichment, which included P body which is involved in mRNA metabolism and in Drosophila oogenesis (Lin *et al.,* 2008). Enrichment of molecular function was of mRNA 3'-UTR binding which may be involved in post-transcriptional gene expression regulation mediated by *trans*-acting factors and in which microRNAs guide-associated proteins towards the 3' UTRs of mRNAs to repress expression (Mayya and Duchaine 2019). miRNAs have been implicated in regulation of male and female fertility and ovary morphology (Chen *et al.,* 2014) and were recently shown to be involved in a number of postmating responses in *D. melanogaster* females (Fowler *et al.,* 2019). That these show mismatched expression changes in the ovary, potentially affecting female fertility, suggests that these alterations could negatively impact fitness resulting in postmating prezygotic reproductive isolation.

When contrasting the effect of the sexual selection treatment of the male-on-female postmating responses (MMvEM; EEvME) we again found differences between the female tissues. In the female reproductive tract, the main arena for postcopulatory sexual selection, the number of mismatched differentially expressed genes was 8x greater after mating with monandry males than polyandry males whereas in the ovaries, polyandry males caused 2x more differentially expressed genes than monandry males. In the female reproductive tract, more genes were upregulated after non-coevolved mating (M males with E females) than downregulated and in ovaries more genes were downregulated after a non-coevolved mating (E males with M females) than upregulated. However, in both the female reproductive tract and ovaries, genes upregulated when a E female mates with a M male are related to a large number of biological processes influencing DNA replication, and in the female reproductive tract these genes are non-congruent. This indicates that, while these responses are unique and mismatched in non-coevolved crosses, M males tend to cause postmating responses that makes the E female more M-like, although many of these genes do not show extreme changes in gene expression. In the ovaries, while E males mated to M females resulted in a large number of differentially expressed genes, the number of biological processes was limited, impacting eggshell assembly and immunity.

We often found changes to immune-related genes. Differential expression of Drosophila immune-related genes in response to mating in *D. melanogaster* have been shown previously (Lawniczak and Begun 2004; McGraw *et al.* 2004; 2008; Mack *et al.* 2006). Innoncenti and Morrow (2009)

have suggested that upregulation of immune genes after mating indicates males are immunogenic, arising from sexually antagonistic sexual interactions. Here we found that some of these genes are upregulated after mating, no matter what the sexual selection treatment. We also found that polyandry females had significant upregulation of some immune genes when virgin which were downregulated after mating. The upregulation of immune genes in virgin polyandry females may be in preparation for a manipulative or pathogenic ejaculate, and upregulation of some genes during courtship provides immunogenic benefits (Zhong et al. 2013). However, such priming does not always appear to benefit *D. melanogaster* females (McKean and Nunney 2005). In contrast, monandry females, who do not experience sexual conflict, upregulated immune genes only after mating. Thus some postmating expression changes in immune related genes are not immunogenic *sensu* Innocenti and Morrow (2009). Previous work examining the postmating response of females mated to heterospecific males identified immunogenic effects (Ahmed-Braimah *et al.* 2020). We found a consistent pattern in which gene expression of immune and stress related genes were either uniquely altered in non-coevolved matings or showed non-congruent changes. While we show immune-related genes frequently change expression during mating, alter expression patterns in response to sexual selection, and can be mismatched during non-coevolved crosses, how these gene expression changes affect fitness remains to be determined (Oku *et al.* 2019).

Changes in gene expression we identified here and in sex-biased gene expression in response to sexual selection (Veltsos et al. 2017) have recently been shown to associate with genomic divergence in these lines (Wiberg *et al.* in review). Sexual selection has been implicated in gene expression and genomic divergence in natural populations and our experimental evolution work supports rapid molecular evolution in response to sexual selection. Rapid coevolution of the molecular interactions, particularly in the female reproductive tract, as we see here, and the subsequent effect on the postmating response is thought to generate mismatches between male and female proteins necessary for successful fertilization (e.g., Ahmed-Braimah *et al.* 2020). Consequently, we found gene expression mismatches in matings between diverged populations, with the female reproductive tract being particularly sensitive. Such mismatches may form the basis of subsequent postmating prezygotic reproductive isolation. Our results, simultaneously comparing gene expression responses in multiple reproductive tissues and under mating system manipulation, highlight the complexity and rapid evolution of ejaculate-female interactions and their potential to influence population divergence.

Material and methods

Experimental evolution lines

The origin, establishment, and maintenance of the selection lines are described in detail elsewhere

(Crudgington *et al.* 2005). Briefly, 50 wild-caught females of *D. pseudoobscura* from a population in

Tucson, AZ USA were brought into the laboratory and reared for three generations, then four

replicate lines of two different sexual selection treatments were established. We modified the

opportunity for sexual selection by manipulating the adult sex ratio in food vials (2.5 x 80 mm) by

either confining one female with a single male (enforced monogamy treatment; M) or one female

with six males ("elevated" polyandry treatment; E). This species is naturally polyandrous with wild-

caught females frequently being inseminated by at least two males at any given time (Anderson

1974). We successfully equalized effective population sizes between the treatments (Snook *et al.*

2009). At each generation, offspring were collected and pooled together within each replicate line

for each treatment, and a sample from this pool was used to start the next non-overlapping

generation in the appropriate sex ratios. Thus, this proportionally reflected the differential offspring

production across families within a replicate and treatment. Generation time was 28 days and all

populations were kept at $22^{o}C$ on a 12L;12D cycle, with standard food media and added live yeast.

Note that 'monandry' versus 'polyandry' as used here refers to the evolutionary history under which

the individuals have evolved, not their current reproductive status.

Sample preparation

To generate experimental males and females, parents were collected from each replicate at

generation 157-158. We standardized for maternal and larval environments as previously

described (Crudgington *et al.* 2010). Briefly, parents were mated en masse in food bottles,

transferred to containers with oviposition plates, allowed to oviposit for 24 h, and then 48 hr later,

100 first instar larvae were seeded in standard food vials (Crudgington *et al.* 2010). Virgin males

and females were collected under light CO_2 anesthesia on the day of eclosion and kept in vials of

10 individuals for 5 days to ensure reproductive maturity (Snook and Markow 2001). On Day 5,

within a 2 h window after lights turned on, one virgin female was placed in a food vial with one

virgin male that was from either the same experimental replicate ("coevolved"; MM, EE where the

first letter is the female) or the other treatment ("non-coevolved"; ME, EM). We dissected age- and

circadian rhythm-matched virgin males and females from the same collections. Each treatment

used 100 individuals, the tissues of which were equally split into 4 separate tubes, for easy

pooling. For the mating treatments, males were put first in individual vials with fly food and allowed

to settle. Females were then added, and were dissected 6 h after the first couple mated, in the

order of mating, within a 2 h block. Dissections were performed under ether anaesthesia in RNA

later (Ambion) on ice blocks. We separated the ovaries and the remainder of the female

reproductive tract, including the sperm storage organs (seminal receptacle and spermathecae).

We refer to these different female tissue sets as ovaries and the female reproductive tract. The

male accessory glands and testes were dissected separately (ejaculatory bulbs were not included).

All tissues were left at $4^{o}C$ in RNA later for one day and then transferred to $-80^{o}C$ until RNA

extraction. Pools of 4 replicates of the E and M treatments, each containing tissues from 25

individuals per replicate, were processed for RNA extraction using Trizol (Ambion) following the

manufacturer's instructions. RNA extractions were cleaned up in Qiagen RNeasy kit columns

according to the manufacturer's protocol, including the 15 min DNase treatment. The quality of

RNA extractions was checked with Nanodrop and Bioanalyser.

Sequencing and mapping

The sample libraries were sequenced using an Illumina HiSeq 2000. Reads were mapped to the *D.*

pseudoobscura genome v3.1, and indexed using bowtie2 (Langmead *et al.* 2009). Paired-end

reads were aligned using option "-g 1 –library-type fr-secondstrand" with TopHat2.0.8b (which calls

bowtie2.1.0; Kim *et al.* 2013). The option "-g 1" instructs TopHat2 to report the best alignment to

the reference for a given read. Exon features were counted using HTSeq-count (Anders *et al.*

2015) and the reads of all exons of each gene were combined to provide overall measures of gene

expression.

Statistical analyses

We analysed the count data using edgeR v3.18.1 (Robinson *et al.* 2010) running in R v.3.4.0 (R

Development Core Team, 2007). Libraries were normalised with the default edgeR normalisation

procedure (TMM) and only genes with at least 3 counts per million across all libraries used in each

analysis were retained. Dispersion was measured with default parameters using a negative

binomial model using only the directly contrasted libraries. We performed different contrasts, but

within each we tested for mean expression differences between the significantly upregulated and

downregulated genes using Mann-Whitney rank tests using R v3.4.0 (R Development Core Team

2007). We considered genes to be differentially expressed if they were below the 5% false

discovery rate (FDR) threshold (Benjamini and Hochberg, 1995). We did not employ a $\log_2$FC

threshold because allometry is unlikely to influence results obtained from specific tissues

(Montgomery and Mank 2016).

We performed several different contrasts including between the lines for divergence in virgin gene

expression in testes, accessory glands, ovaries, and female reproductive tract separately. Note

that differential gene expression results are relative. For simplicity, figures report upregulated

genes next to the name of each contrast; they can also be interpreted as downregulated genes for

the opposite contrast. For example, 161 genes were upregulated in the testes of E males, which

could also be interpreted as 161 genes downregulated in M testes. We also assessed divergence

in the female postmating response by contrasting, within each sexual selection treatment,

differential expression between virgin and mated ovaries and the female reproductive tract,

separately. We consider the virgin gene expression status to be the baseline, so we can categorize

changes as either upregulated after mating (gene expression is higher in mated compared to

virgin) or downregulated after mating (gene expression is higher in virgin compared to mated). We also investigated the effect of non-coevolved matings relative to coevolved matings, either focusing on the same female treatment but different male treatment (e.g. a contrast between either M̲M v M̲E or E̲E v E̲M, where the focal sex is unscored and females are always listed first) or on the same male treatment but different female treatment (i.e. MM̲ v EM̲ or EE̲ v ME̲). These were conducted for both components of the female reproductive tissue. Here we consider the coevolved mating as a baseline, so we can categorize changes in gene expression in non-coevolved matings relative to the coevolved. Notation then is either that the non-coevolved postmating response has relatively higher or lower expression (the former indicated by, e.g., "EM up" and the later indicated by, e.g., "EM down").

We identified gene subsets based on congruent and non-congruent response to sexual selection between the selection lines using venn diagrams, constructed using Venny 2.1 (Oliveros 2015). We performed GO enrichment analysis for all DE genes from all contrasts, and those udentified from the venn diagrams, using topGO v2.22.0 with the weight01 algorithm option to account for GO topology (Alexa and Rahnenfuhrer 2010). Results with $p < 0.05$ on Fisher's exact tests, corrected for topology, were retained (Supplementary File 1).

For analysis of Sfps, we contrasted the distribution of the change in expression ($\log_2$FC) of all genes, and differentially expressed genes between the selection lines from accessory glands using density plots. We made contrasts to the distribution of 3 Sfp-related gene subsets, identified from *D. pseudoobscura* proteomics (Karr *et al.* 2019). The largest subset was 3281 proteins produced in the accessory gland ("proteome"). Of these, 528 had protein secretory signals ("secretome") and 163 were also orthologous to *D. melanogaster* seminal fluid proteins (putative Sfps or "exoproteome"). Differences in the median expression differences between these sets were analysed with Wilcoxon tests, conducted in R v3.4.0 (R Development Core Team 2007).

Acknowledgements

This work was supported by the Natural Environment Research Council (NE/I014632/1 to M.G.R., A.R.C., and R.R.S) and the Natural Environment Research Council Biomolecular Analysis Facility (NBAF654 to M.G.R.). The RNAseq data underlying this article have been submitted to ArrayExpress under accession number E-MTAB-10047. The scripts to reproduce the analysis are available at https://osf.io/z7fm9/?view_only=054171ba3f534f839a0814fa1b8f9f61. We thank the many people that have contributed to the generation and maintenance of the experimental sexual selection lines in RRS's lab and R Axel W Wiberg for help with dissections.

Figures

Figure 1. Differential gene expression (absolute $\log_2$FC changes, y axis) and number of

significantly upregulated genes (numbers to the right of the ridge plot) when comparing monoandry

(M) and polyandry (E) tissues of a) males and b) females.

Figure 2. Effect of sexual selection on expression of genes coding for a) all and b) differentially

expressed male accessory gland genes between polyandry (positive x axis values) and monandry

(negative x-axis values) treatments. The line colours differentiate between independently

categorized parts of the transcriptome as proteome, secretome or seminal fluid proteins (Karr *et al.*

2019).

 Figure 3. Postmating responses of differentially expressed genes (log$_2$FC) in the female

reproductive tracts (top; a-c) and ovaries (bottom; d-f). a,c) Within line mating contrasts showing

either upregulation (positive values) or downregulation (negative values) of genes following mating

within lines, either in monandry (x axis) or polyandry (y axis) females. Gene expression response

direction is defined relative to virgins. In panel a, a similar response (congruent) in both selection

lines is shown with a purple diagonal line, and a non-congruent response is shown with a red

diagonal line. Panel a and c legends indicate genes uniquely differentially expressed in the

monandry line (M) in yellow circles, those in the polyandry line (E) in blue circles and genes

differentially expressed in both selection lines (M+E) in green diamonds. Grey circles are genes

that are not differentially expressed. b,d) Same female, different male contrast showing either

upregulation (yellow triangle, red cross) or downregulation (purple triangle, blue plus) of genes

following mating with a non-coevolved male, focusing on the response within each female line.

Panel b and d legends show the female line in the first letter and the male line the second letter

(i.e., ME = monandry (M) female mated to polyandry (E) male; EM = polyandry female (E) mated

to monandry (M) male). Contrast made relative to relevant coevolved cross (i.e., for ME, contrast

with MM; for EM, contrast with EE). c,f) Different male, same female contrast showing either

upregulation (red triangle, yellow cross) or downregulation (blue triangle, purple plus) of genes

following mating with a non-coevolved male, focusing on the response of each male line. Panel c

and e legends show the female line in the first letter and the male line the second letter. Contrast

made relative to relevant coevolved cross (i.e., for ME, contrast with EE; for EM, contrast with

MM).

References

Ahmed-Braimah YH, Unckless RL, Clark AG. 2017. Evolutionary dynamics of male reproductive
651   genes in the *Drosophila virilis* subgroup. *G3* 7:3145-3155.
652   Ahmed-Braimah YH, Wolfner MF, Clark AG. 2020. Differences in post-mating transcriptional
653   responses between conspecific and heterospecific matings in *Drosophila*. *Mol Biol Evol.*
654   doi:10.1093/molbev/msaa264
655   Anders S, Pyl PT, Huber W. 2015. HTSeq-a Python framework to work with high-throughput
656   sequencing data. *Bioinformatics.* 31:166-169.
657   Anderson WW. 1974. Frequent multiple insemination in a natural population of *Drosophila*
658   *pseudoobscura*. *Am Nat.* 108:709-711.
659   Alexa A, Rahnenfuhrer J. 2010. topGO: enrichment analysis for gene ontology. R package version.
660   2
661   Arnqvist G, Rowe L. 2005. Sexual Conflict. New Jersey: Princeton University Press.
662   Avila FW, Ram KR, Qazi MCB, Wolfner MF. 2010. Sex peptide is required for the efficient release
663   of stored sperm in mated *Drosophila females*. *Genetics.*186:595-600.
664   Avila FW, Mattei AL, Wolfner MF. 2015. Sex peptide receptor is required for the release of stored
665   sperm by mated *Drosophila melanogaster* females. *J Insect Physiol.* 76:1-6.
666   Bath E, Bowden S, Peters C, Reddy A, Tobias JA, Easton-Calabria E, Seddon N, Goodwin SF,
667   Wigby S. 2017. Sperm and sex peptide stimulate aggression in female *Drosophila*. *Nat
668   Ecol Evol.* 1:0154.
669   Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful
670   approach to multiple testing. *J Roy Stat Soc B.* 57:289-300.
671   Bono JM, Matzkin LM, Kelleher ES, Markow TA. 2011. Postmating transcriptional changes in
672   reproductive tracts of con- and heterospecifically mated Drosophila mojavensis females.
673   *Proc Natl Acad Sci U S A.* 108:7878-7883.
674   Carvalho GB, Kapahi P, Anderson DJ, Benzer S. 2006. Allocrine modulation of feeding behavior
675   by the sex peptide of *Drosophila*. *Curr Biol.* 16:692-696.
676   Chapman T, Bangham J, Vinti G, Seifried B, Lung O, Wolfner MF, Smith HK, Partridge L. 2003.
677   The sex peptide of *Drosophila melanogaster*: female post-mating responses analyzed by
678   using RNA interference. *Proc Natl Acad Sci U S A.* 100:9923-9928.
679   Chen YW, Song S, Weng R, Verma P, Kugler JM, Buescher M, Rouam S, Cohen SM. 2014.
680   Systematic study of *Drosophila* microRNA functions using a collection of targeted knockout
681   mutations. *Dev Cell.* 31:784-800.
682   Cheng C, Kirkpatrick M. 2016. Sex-specific selection and sex-biased gene expression in humans
683   and flies. *PLoS Genet.* 12:e1006170.
684   Cognigni P, Bailey AP, Miguel-Aliaga I. 2011. Enteric neurons and systemic signals couple
685   nutritional and reproductive status with intestinal homeostasis. *Cell Metab.* 13:92-104.

Coyne JA, Orr HA. 2004. Speciation. Massachusetts: Sinauer Associates.

Crudgington HS, Beckerman AP, Brüstle L, Green K, Snook RR. 2005. Experimental removal and
elevation of sexual selection: does sexual selection generate manipulative males and
resistant females? *Am Nat.* 165:S72-S87.

Crudgington HS, Fellows S, Badcock NS, Snook RR. 2009. Experimental manipulation of sexual
selection promotes greater male mating capacity but does not alter sperm investment.
Evolution. 63:926-938.

Crudgington HS, Fellows S, Snook RR. 2010. Increased opportunity for sexual conflict promotes
harmful males with elevated courtship frequencies. *J Evol Biol.* 23:440-446.

Debelle A, Ritchie MG, Snook RR. 2014. Evolution of divergent female mating preference in
response to experimental sexual selection. *Evolution.* 68:2524-2533.

Debelle A, Ritchie MG, Snook RR. 2016. Sexual selection and assortative mating: an experimental
test. *J Evol Biol.* 29:1307-1316.

Delbare SYN, Chow CY, Wolfner MF, Clark AG. 2017. Roles of female and male genotype in post-
mating responses in *Drosophila melanogaster*. *J Hered.* 108:740-753.

Ellegren H, Parsch J. 2007. The evolution of sex-biased genes and sex-biased gene expression.
Nat Rev Genet. 8:689-698.

Elphick MR, Mirabeau O, Larhammar D. 2018. Evolution of neuropeptide signalling systems. *J Exp
Biol.* 221:jeb.151092

Fowler EK, Bradley T, Moxon S, Chapman T. 2019. Divergence in transcriptional and regulatory
responses to mating in male and female fruitflies. *Scientific Reports.* 9:1-15.

Garlovsky MD, Yusuf LH, Ritchie MG, Snook RR. 2020. Within-population sperm competition
intensity does not predict asymmetry in conpopulation sperm precedence. *Phil Trans Roy
Soc B.* 375:20200071.

Harrison PW, Wright AE, Zimmer F, Dean R, Montgomery SH, Pointer MA, Mank JE. 2015. Sexual
selection drives evolution and rapid turnover of male gene expression. *Proc Natl Acad Sci
U S A.* 112:4393-4398.

Heifetz Y, Wolfner MF. 2004. Mating, seminal fluid components, and sperm cause changes in
vesicle release in the *Drosophila* female reproductive tract. *Proc Natl Acad Sci U S A.*
101:6261-6266.

Holland B, Rice WR. 1999. Experimental removal of sexual selection reverses intersexual
antagonistic coevolution and removes a reproductive load. *Proc Natl Acad Sci U S A.*
96:5083-5088.

Hollis B, Houle D, Yan Z, Kawecki TJ, Keller L. 2014. Evolution under monogamy feminizes gene
expression in *Drosophila melanogaster*. *Nat Commun.* 5:3482.

Hollis B, Houle D, Kawecki TJ. 2016. Evolution of reduced post-copulatory molecular interactions
in *Drosophila* populations lacking sperm competition. *J Evol Biol.* 29:77-85.

Hollis B, Koppik M, Wensing KU, Ruhmann H, Genzoni E, Erkosar B, Kawecki TJ, Fricke C, Keller
L. 2019. Sexual conflict drives male manipulation of female postmating responses in
Drosophila melanogaster. Proc Natl Acad Sci U S A. 116:8437-8444.

Hosken DJ, Ward PI. 2001. Experimental evidence for testis size evolution via sperm competition.
Ecol Lett. 4:10-13.

Hosken DJ, Garner TWJ, Ward PI. 2001. Sexual conflict selects for male and female reproductive
characters. *Curr Biol.* 11:489-493.

Immonen E, Snook RR, Ritchie MG. 2014. Mating system variation drives rapid evolution of the
female transcriptome in *Drosophila pseudoobscura. Ecol Evol.* 4:2186-2201.

Innocenti P, Morrow EH. 2009. Immunogenic males: a genome-wide analysis of reproduction and
the cost of mating in *Drosophila melanogaster* females. *J Evol Biol.* 22:964-973.

Innocenti P, Flis I, Morrow EH. 2014. Female responses to experimental removal of sexual
selection components in *Drosophila melanogaster. BMC Evol Biol.* 14:239.

Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. 2013. TopHat2: accurate
alignment of transcriptomes in the presence of insertions, deletions and gene fusions.
Genome Biol. 14:R36.

Kapelnikov A, Zelinger E, Gottlieb Y, Rhrissorrakrai K, Gunsalus KC, Heifetz Y. 2008. Mating
induces an immune response and developmental switch in the *Drosophila* oviduct. *Proc*
Natl Acad Sci U S A. 105:13912-13917.

Karr TL, Southern H, Rosenow MA, Gossmann TI, Snook RR. 2019. The old and the new:
Discovery proteomics identifies putative novel seminal fluid proteins in *Drosophila. Mol Cell*
Proteo. 18:S23-S33.

Langmead B, Trapnell C, Pop M, Salzberg SL. 2009. Ultrafast and memory-efficient alignment of
short DNA sequences to the human genome. *Genome Biol.* 10:R25.

Lawniczak MK, Begun DJ. 2004. A genome-wide analysis of courting and mating responses in
Drosophila melanogaster females. *Genome.* 47:900-910.

Librado P, Rozas J. 2016. Weak polygenic selection drives the rapid adaptation of the
chemosensory system: Lessons from the upstream regions of the major gene families.
Genome Biol Evol. 8:2493-2504.

Lin MD, Jiao X, Grima D, Newbury SF, Kiledjian M, Chou TB. 2008. *Drosophila* processing bodies
in oogenesis. *Dev Biol.* 322:276-288.

Linklater JR, Wertheim B, Wigby S, Chapman T. 2007. Ejaculate depletion patterns evolve in
response to experimental manipulation of sex ratio in *Drosophila melanogaster. Evolution.*
61:2027-2034

Lüpold S, Linz GM, Rivers JW, Westneat DF, Birkhead TR. 2009. Sperm competition selects
beyond relative testes size in birds. *Evolution.* 63:391-402.

Mack PD, Kapelnikov A, Heifetz Y, Bender M. 2006. Mating-responsive genes in reproductive
tissues of female *Drosophila melanogaster. Proc Natl Acad Sci U S A.* 103:10358-10363.

Manier MK, Lüpold S, Belote JM, Starmer WT, Berben KS, Ala-Honkola O, Collins WF, Pitnick S.
2013. Postcopulatory sexual selection generates speciation phenotypes in Drosophila. *Curr*
Biol. 23:1853-1862.

Mank JE. 2017. The transcriptional architecture of phenotypic dimorphism. *Nat Ecol Evol.* 1:0006.

Markow TA. 1997. Assortative fertilization in *Drosophila. Proc Natl Acad Sci U S A.* 94:7756-7760.

Matute DR, Coyne JA. 2010. Intrinsic reproductive isolation between two sister species of
Drosophila. Evolution. 64:903-920.

Mayya VK, Duchaine TF. 2019. Ciphers and executioners: How 3'-untranslated regions determine
the fate of messenger RNAs. *Front Genet.* 10:6.

McGraw LA, Clark AG, Wolfner MF. 2008. Post-mating gene expression profiles of female
Drosophila melanogaster in response to time and to four male accessory gland proteins.
Genetics. 179:1395-1408.

McGraw LA, Gibson G, Clark AG, Wolfner MF. 2004. Genes regulated by mating, sperm, or
seminal proteins in mated female *Drosophila melanogaster. Curr Biol.* 14:1509-1514.

McKean KA, Nunney L. 2005. Bateman's principle and immunity: phenotypically plastic
reproductive strategies predict changes in immunological sex differences. *Evolution.*
59:1510-1517.

Montgomery SH, Mank JE. 2016. Inferring regulatory change from gene expression: the
confounding effects of tissue scaling. *Mol Ecol.* 25:5114-5128.

Niepielko MG, Ip K, Kanodia JS, Lun DS, Yakoby N. 2012. Evolution of BMP signaling in
Drosophila oogenesis: a receptor-based mechanism. *Biophys J.* 102:1722-1730.

Oku K, Price TAR, Wedell N. 2019. Does mating negatively affect female immune defences in
insects? *Ani Biol.* 69:117-136.

Oliveros, J.C. 2007-2015 Venny. An interactive tool for comparing lists with Venn's
diagrams. http://bioinfogp.cnb.csic.es/tools/venny/index.html

Parker DJ, Bast J, Jalvingh K, Dumas Z, Robinson-Rechavi M, Schwander T. 2019. Sex-biased
gene expression is repeatedly masculinized in asexual females. *Nature Commun.* 10:1-11.

Pitnick S, Hosken DJ, Birkhead TR. 2008. Sperm biology: an evolutionary perspective.
Massachusetts: Academic Press.

Pitnick S, Miller GT, Reagan J, Holland B. 2001. Males' evolutionary responses to experimental
removal of sexual selection. *Proc Roy Soci Biol B.* 268:1071-1080.

Pizzari T, Snook RR. 2003. Perspective: sexual conflict and sexual selection: chasing away
paradigm shifts. *Evolution.* 57:1223-1236.

Prokupek AM, Kachman SD, Ladunga I, Harshman LG. 2009. Transcriptional profiling of the sperm
storage organs of *Drosophila melanogaster. Insect Mol Biol.* 18:465-475.

Pröschel M, Zhang Z, Parsch J. 2006. Widespread adaptive evolution of *Drosophila* genes with
sex-biased expression. *Genetics.* 174:893.

R Development Core Team. 2007 R: A language and environment for statistical computing,

Vienna, Austria. ISBN 3-900051-07-0, URL http://www.R-project.org. R-project.org

Reuter M, Linklater JR, Lehmann L, Fowler K, Chapman T, Hurst GD. 2008. Adaptation to

experimental alterations of the operational sex ratio in populations of *Drosophila*

melanogaster. *Evolution*. 62:401-412.

Rice WR. 1996. Sexually antagonistic male adaptation triggered by experimental arrest of female

evolution. *Nature*. 381:232-234.

Ribeiro C, Dickson BJ. 2010. Sex peptide receptor and neuronal TOR/S6K signaling modulate

nutrient balancing in *Drosophila*. *Curr Biol*. 20:1000-1005.

Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for differential

expression analysis of digital gene expression data. *Bioinformatics*. 26:139-140.

Rowe L, Cameron E, Day T. 2003. Detecting sexually antagonistic coevolution with population

crosses. *Proc Roy Soc Biol B*. 270:2009-2016.

Sirot LK, Wong A, Chapman T, Wolfner MF. 2015. Sexual conflict and seminal fluid proteins: a

dynamic landscape of sexual interactions. *Cold Spring Harbor Perspectives in Biology*.

7:a017533.

Snook RR, Markow TA. 2000. Mating system evolution in sperm-heteromorphic *Drosophila*. *J*

Insect Physiol. 47:957-964.

Snook RR, Robertson A, Crudgington HS, Ritchie MG. 2005. Experimental manipulation of sexual

selection and the evolution of courtship song in *Drosophila pseudoobscura*. *Behavior*

Genetics. 35:245-255.

Snook RR, Brüstle L, Slate J. 2009. A test and review of the role of effective population size on

experimental sexual selection patterns. *Evolution*. 63:1923-1933.

Snook RR, Markow TA, Karr TL. 1994. Functional nonequivalence of sperm in *Drosophila*

pseudoobscura. *Proc Natl Acad Sci U S A*. 91:11222-11226.

Veltsos P, Fang Y, Cossins AR, Snook RR, Ritchie MG. 2017. Mating system manipulation and the

evolution of sex-biased gene expression in *Drosophila*. *Nat Commun*. 8:2072.

Vrablik TL, Watts JL. 2013. Polyunsaturated fatty acid derived signaling in reproduction and

development: Insights from *Caenorhabditis elegans* and *Drosophila melanogaster*. *Mol*

Reprod Dev. 80:244-259.

Wolfner MF. 2009. Battle and ballet: molecular interactions between the sexes in *Drosophila*. *J*

Hered. 100:399-410.

Wiberg RAW, Veltsos P, Snook RR, Ritchie, MG. In review. Experimental evolution supports

signatures of sexual selection in genomic divergence. *Evol. Lett.* (a previous version on

biorxiv: https://doi.org/10.1101/2020.09.07.285650)

Wigby S, Chapman T. 2004. Female resistance to male harm evolves in response to manipulation

of sexual conflict. *Evolution*. 58:1028-1037

Wright AE, Rogers TF, Fumagalli M, Cooney CR, Mank JE. 2019. Phenotypic sexual dimorphism

is associated with genomic signatures of resolved sexual conflict. *Mol Ecol.* 28:2860-2871.

Zhang Y, Sturgill D, Parisi M, Kumar S, Oliver B. 2007. Constraint and turnover in sex-biased gene

expression in the genus *Drosophila*. *Nature.* 450:233-237.

Zhong, W., McClure, C.D., Evans, C.R., Miynski, D.T., Immonen, E., Ritchie, M.G. & Priest, N.K.

2013. Immune anticipation in *Drosophila*: Turandot M promotes immunity against sexually

transmitted fungal infections. *Proc Roy Soc Biol B* 280: 20132018

Supplementary Files

Supplementary File 1. Summary of gene ontology enrichment analysis for genes and all contrasts, shown in Figures 1 and 3 (and Figures S1-3). The order of the tabs in the file corresponds to the order of appearance of the genes in the Figures. Each tab shows all DE genes, and separately upregulated and downregulated genes. Tab names refer to Figure, tissue and contrast. Contrast names refer to the polyandry (E) and monogamous (M) lines. The virgin treatment is represented by the letter indicating the selection line, while the mated treatment is represented by two letters, starting with the sexual selection line of the female and followed by the male partner (i.e., "EM" indicates the E females mated with M males).

Figure S1. Differential gene expression (absolute log$_2$FC changes, y axis) and number of
significantly upregulated (Up) or downregulated (Down) genes (numbers to the right of the ridge
plot) when comparing the postmating response in the female reproductive tract (FRT) or ovaries
(Ovary) in monandry (M) crosses and in polyandry (E) crosses. The direction of gene expression is
defined relative to mated females (e.g. Down means the mated female had significantly lower
expression than in the virgin status; Up means the mated female had significantly higher
expression than in the virgin status).

Figure S3. Differential gene expression (absolute log$_2$FC changes, y axis) and number of
significantly upregulated (Up) or downregulated (Down) genes (numbers to the right of the ridge
plot) when comparing the postmating response in the female reproductive tract (FRT) or ovaries
(Ovary) for the male effect (i.e., when a polyandry (E) female is mated to a monandry (M) male
(Effect of M male), or when a monandry female (M) is mated to a polyandry (E) male (Effect of E
male)). The direction of gene expression is defined relative to the relevant coevolved cross (e.g.
Down means a female mated to the non-coevolved male (i.e., either EM or ME) had significantly
lower expression than when mated to a coevolved male (i.e., MM or EE, respectively); Up means
the female mated to non-coevolved male had significantly higher expression than when mated to a
coevolved male).