Development of first linkage map for *Silphium integrifolium* (*Asteraceae*) enables identification of sporophytic self-incompatibility locus

John H. Price[1], Andrew R. Raduski[2], Yaniv Brandvain[2], David L. Van Tassel[3] & Kevin P. Smith[1]

1. Department of Agronomy and Plant Genetics, University of Minnesota Twin Cities, 411 Borlaug Hall, 1991 Upper Buford Circle, St. Paul, MN 55108

2. Department of Plant and Microbial Biology, University of Minnesota Twin Cities, 140 Gortner Laboratory, 1479 Gortner Ave, St. Paul, MN 55108

3. The Land Institute, 2440 E. Water Well Road, Salina, KS 67401

Corresponding author: Kevin P. Smith, +1 612 624 1211, smith376@umn.edu

16 SUMMARY

- *Silphium integrifolium* (*Asteraceae*) has been identified as a candidate for domestication
18 as a perennial oilseed crop and has a sporophytic self-incompatibility system—the
genetic basis of which is not well understood in the *Asteraceae*. To address this gap, we
20 sought to map the genomic location of the self-recognition locus (S-locus) in this species.

- We used a biparental population and genotyping-by-sequencing to create the first genetic
22 linkage map for this species. Then we developed a novel crossing scheme and set of
analysis methods in order to infer S-locus genotypes for a subset of these individuals,
24 allowing us to map the trait. Finally, we identified potential gene candidates using
synteny analysis with the annual sunflower (*Helianthus annuus*) genome.

26 - Our linkage map contains 198 SNP markers and resolved into the correct number of
linkage groups. We were able to successfully map the S-locus and identify several
28 potential gene candidates in the sunflower syntenic region.

- Our method is effective and efficient, allowed us to map the *S. integrifolium* S-locus
30 using fewer resources than previous studies, and could be readily be applied to other
species. Our best gene candidate appears to be worthy of future work in *S. integrifolium*
32 and other *Asteraceae* species.

INTRODUCTION

38   *Silphium integrifolium* (Michx.) (wholeleaf rosinweed or silflower) is a member of the

Asteraceae family native to prairies throughout the central United States. In the early 2000s, *S.*

40   *integrifolium* was selected to be a candidate for domestication as a perennial oilseed crop by the

Land Institute in Salina, Kansas (Van Tassel *et al.,* 2017), attracting attention for its tolerance to

42   drought, upright growth habit, and large seeds (DeHaan *et al.*, 2016). Subsequently, *S.*

integrifolium has been found to have a seed oil composition similar to landrace sunflower

44   (Reinert et al., 2019) and good winter survival and persistence in a range of climates (J.H. Price

& D.L. Van Tassel, pers. obs.). In addition, the yield potential of *S. integrifolium* populations

46   that have undergone relatively little selection is approximately 60% the yield of advanced

sunflower hybrids (Kandel *et al.*, 2019; Schiffner *et al.*, 2020), indicating that significant

48   improvement is likely with continued breeding efforts. These characteristics further encourage

the domestication of this species as a new crop. In the past, domestication occurred over long

50   periods of time, largely due to the largely unintentional nature of early selection (Rindos, 1984).

With the advantage of contemporary knowledge of genetics, genomics, and breeding techniques,

52   the amount of time necessary to domesticate a new crop could be drastically reduced. Therefore,

the development of genomic resources is a crucial step in this process (Sedbrook *et al.*, 2014). To

54   this end, we have developed the first genetic linkage map for *S. integrifolium*.

56   Among the traits for which better genetic knowledge will accelerate the domestication of *S.*

integrifolium is self-incompatibility. Although occasional *S. integrifolium* individuals have been

58   observed to produce at least some seed when self-pollinated (Reinert *et al.*, 2020), *S.*

integrifolium is self-incompatible, and as a member of the *Asteraceae* family is assumed to have

60   a sporophytic self-incompatibility (SSI) system (Hiscock, 2000). In sporophytic systems, self-

recognition is typically controlled by a single multi-allelic locus, known as the "S-locus", with

62   rejection of self-pollen caused by stigma recognition of S-locus gene products found in or on the

pollen. Because these products are produced in the anther, pollen acceptance or rejection is

64   determined by the diploid genotype of the male parent, rather than the haploid genotype of a

given pollen grain (Hiscock & Tabah, 2003). SSI alleles are also expected to display complex

66   dominance patterns, and dominance relationships between alleles may differ from the anther to

the stigma (Hiscock & Tabah, 2003).

The molecular mechanisms that underlie SSI are best described in the *Brassicaceae*, where the female S-phenotype is determined by a receptor kinase complex known as SRK. When pollen of the same S-phenotype lands on the stigma, this kinase binds a cysteine-rich protein (CRP) found in the pollen coat, known as SP_{11}/SCR, initiating the pollen rejection response (Fujii & Takayama, 2018). These two genes are tightly linked and rarely recombine; thus, they combine to form the S-locus, which is more properly thought of as an S-haplotype (Edh *et al.*, 2009). SRK is unable to bind SP_{11}/SCR proteins produced by different haplotypes, resulting in the acceptance of non-self pollen (Fujii & Takayama, 2018). Although the identity of the receptor protein varies, secreted CRPs are also considered a promising candidate for the male determinant of SSI in the *Convolvulaceae* (morning glory) family (Rahman *et al*., 2007), and serve as the female determinant of gametophytic self-incompatibility (GSI) in the genus *Papaver* (poppy) (Marshall *et al*., 2011),.

Asteraceae systems are less well understood. In *Senecio squalidis*, and subsequently in other *Asteraceae*, SRK-like sequences have been identified and cloned. However, results from *S. squalidis* and chicory (*Cichorium intybus*) indicate that they likely are not integral to S-genotype determination (Hiscock & Tabah, 2003 ; Gonthier *et al*., 2013), and that the molecular control of *Asteraceae* SSI is quite different from the *Brassicaceae* system (Allen *et al.* 2011). Efforts to map the S-locus in chicory provided a 1.8 cM QTL region but have not yet determined a molecular basis for self-incompatibility (Gonthier *et al.*, 2013). Although efforts have been undertaken in other species, chicory represents perhaps the only example where a true *Asteraceae* S-locus has been definitively mapped, as opposed to other loci contributing to breakdowns in self-incompatibility (Gandhi *et al*., 2005; Koseva *et al*., 2017).

Mapping the S-locus is important for breeding efforts and for our understanding of the genetics and evolution of this critical locus. For example, the "collaborative nonself recognition" system identified in the *Solanaceae* (Kubo *et al.* 2011) has revealed that the mechanism underlying self-incompatibility can change how new S-alleles originate (Bod'ová *et al*., 2018, Harkness *et al*., 2021) and migrate (Harkness & Brandvain, 2020) across populations. Identifying the basis (or

98   bases) of SSI in the *Asteraceae* would help us better understand and predict features of its

evolution, including its maintenance in populations with very few (two to six) S-alleles (Brennan

100   *et al*. 2006).

102   To map the S-locus, researchers must determine the S-locus genotype for individuals in a

population large enough to conduct linkage mapping. In other species, S-genotype determination

104   required mating large numbers of full-sibling individuals to one or a few "tester" genotypes with

a known S-genotype (Camargo *et al*., 1997; Tomita *et al*., 2004). Testers may be a parent of the

106   population, or may be obtained by mating siblings in a diallel design, grouping individuals based

on their compatibility (Hiscock, 2000), and then selecting one or several of these individuals as a

108   tester for their siblings (Gonthier *et al.*, 2013). However, this process has several limitations. For

some species, pollen availability may limit the number of testcrosses that can be made, especially

110   if clonal propagation is not used to multiply tester individuals. Additionally, because the tester is

typically used as the male parent, differentiation between alleles may be difficult for SSI as

112   dominance relationship between alleles may differ from anthers to stigma. Finally, this process

often requires a large number of crosses, especially if the parents are not available. A diallel

114   requires n(n-1) crosses, where **n** is the number of individuals to be analyzed, and multiple

crosses are likely required for each individual being tested. Even selecting a tester from amongst

116   a conservative number of siblings may require a prohibitively high number of crosses. Methods

of S-allele determination that address these issues may make S-locus mapping feasible for a

118   greater number of species.

120   In this study, we present a putative location for the *S. integrifolium* S-locus. To achieve this, we

developed a novel framework for inferring the S-genotypes of individuals within a population

122   large enough for mapping. Our method does not require tester individuals with known S-

genotypes, requires only three to four crosses per individual, and identifies alleles with sex-

124   specific dominance interactions. We used these inferred genotypes, in conjunction with the

aforementioned linkage map, to identify a QTL likely containing the S-locus. We then identify

126   regions in the sunflower (*Helianthus annuus*) genome assembly syntenic with our putative S-

locus and use this information to identify potential gene candidates for S-allele determination.

MATERIALS AND METHODS

Population development: A linkage map was constructed using an F_1 population of 265

132 *Silphium integrifolium* (Michx.) individuals, derived from the crossing of two genotypes known

as "965" and "1767", selected at the Land Institute. These parents were chosen because they

134 expressed agronomically favorable morphologies and differed for several phenological traits.

Seventy-four seeds from this cross were grown at the Land Institute, and 191 were grown at the

136 University of Minnesota. All 265 individuals were used for map construction, but only a subset

of the 191 Minnesota individuals were used for S-allele determination. Following approximately

138 two months of growth in a greenhouse, the Minnesota seedlings were transplanted to a field at

the Minnesota Agricultural Experiment Station at the University of Minnesota St. Paul campus

140 with 1.2 meters between each plant.

142 **Genotyping and Variant Detection:** Tissue was collected from seedling leaves of the 74 Land

Institute progeny, and adult leaf tissue was collected from the parents and all other progeny, with

144 all tissue lyophilized prior to DNA extraction. Dual indexed genotyping-by-sequencing (GBS)

libraries were created from genomic DNA using the restriction enzymes *SbfI* and *TaqI* and

146 sequenced on 1.5 Illumina NovaSeq 6000 lanes (1x100 single-end reds). All extraction, library

preparation, and sequencing occurred at the University of Minnesota Genomics Center.

Demultiplexed reads were mapped to *S. integrifolium* genomic contigs using BWA-MEM (Y.

150 Brandvain, unpublished; Li, 2013). The "ref_map" pipeline of Version 2.5 of the "Stacks"

variant calling software was then used for the detection of SNP loci (Catchen *et al.*, 2013). Loci

152 that were missing in at least 30% of the population were excluded, as were loci missing in either

parent. This resulted in 935 SNP markers.

Linkage Map Construction: Linkage analysis and map construction were carried out in

156 *JoinMap5* (Stam, 1993). First, 715 SNP markers showing significant (Chi-square test, $P > 0.01$)

segregation distortion were removed. Markers were then grouped by LOD independence score,

158 using a threshold score of five. Any groups that contained fewer than four markers were

discarded. Map order and distances were then estimated using the maximum likelihood mapping

160   algorithm, with default settings. One marker was removed from the end of linkage group three

because it created a gap of more than 50 cM. Linkage group names were assigned based on

162   estimated centimorgan length, with group one being the longest.

164   **Crossing design to determine self-incompatibility genotype:** Of the 191 members of this

mapping population planted in Minnesota, 84 were intermated for S-allele determination. To

166   cross, capitula (compound flower heads) to be used as a female, or both as a male and female,

were covered with a cotton bag prior to anthesis to prevent pollination. Capitula to be used

168   exclusively as a male parent were covered with mesh bags at least one day prior to crossing to

reduce contamination by insects depositing pollen from other plants. After stigma emergence,

170   pollen from a male parent capitulum was collected into a container and dusted onto stigma with a

pipe cleaner. Alternatively, on occasion a capitulum was removed and used to brush pollen

172   directly onto a female parent. Each female capitulum was mated with only one male parent. The

bag was then reclosed, harvested after senescence, and dried. For each capitula, the number of

174   filled seeds and total number of seeds were then counted, with filled and unfilled seeds

differentiated by visual and manual assessment. Seed set was then calculated as the ratio of filled

176   seeds to total seeds.

178   Individuals were crossed in a structured design, which we have named the "connected small

diallel design", illustrated in Figure 1. This structure is based on groupings of four individuals

180   that are mated in a diallel. These small diallels are then linked together through reciprocally

mating single individuals. The purpose of this design was to maximize the amount of

182   information that could be derived from the mating behavior of any given individual, while

minimizing the number of matings for which it would need to be used. The design ensures that

184   all individuals may be connected to one another through pairs of matings, allowing for the entire

population to be used to predict the mating behavior each individual.

The design for this experiment included 20 small diallels. The design was implemented

188   incompletely, with many of the recommended crosses not completed due to time or pollen

availability constraints. To ensure that all individuals could be connected, more crosses

190   connecting diallels were conducted than indicated in the design. Finally, several additional

individuals were included in the study with only reciprocal crosses to one other individual. In total 268 crosses were performed between the 84 individuals, covering 138 different combinations of parents. Of these crosses, 126 combinations were crossed reciprocally, with both individuals used as male and female. One hundred and eighty-six crosses were conducted in 2018, and an additional 82 were conducted in 2019, primarily to increase the number of reciprocal crosses. In seven cases, multiple replications of a cross were performed. These observations were combined by summing the total number of filled and total seeds from each replication, and then calculating a seed set from the sums.

Three distinct but complementary methods were used to translate the results of the connected small diallel experiment into an S-locus map position. In all three methods, each mated pair of individuals is first determined to be compatible or incompatible, based on a threshold seed set value. A threshold value of 20% was selected for this experiment, based on previous observations of seed set in manually self-pollinated individuals (Reinert *et al*., 2020). Matings that resulted in a seed set value above 20% were considered compatible. These methods, along with their relative strengths and weaknesses, are described below. All methods were developed using the R language (R Core Team, 2019).

Direct mapping with single-marker regression: The simplest method attempted to directly associate variation for compatibility with an SNP marker, without inferring the S-genotype of any particular individual. To accomplish this, a logistic regression was conducted following the formula $S_i = M_{ij} + P_{ij} + M_{ij}:P_{ij}$, where the success **S** of the *i*th mating was predicted by the maternal genotype **M**, the paternal genotype **P**, and the interaction of those genotypes, all for the *j*th marker. The calculated P-value of the maternal by paternal genotype interaction was then used to determine association with cross incompatibility, and thus the S-locus. Only biallelic markers were used in this approach, with homozygotes recoded as -1 or 1 and heterozygotes as 0. This approach only makes limited use of available genotype data, as each SNP marker is considered individually. Additionally, this approach could not be applied to species without genetic marker data. Finally, because it does not make inferences about the S-genotype carried by any particular individual or take potential dominance relationships into account, it is not able

to fully leverage the advantages of the connected crossing design used in this study. Thus, this method's usefulness is likely limited to confirming the results of other methods.

Inference of self-incompatibility genotype for all individuals followed by QTL mapping of the inferred S-genotype: As an alternative approach to mapping the S-locus, we develop two methods to infer every individual's S-locus genotype from its crossing behavior, then map this inferred phenotype with traditional QTL mapping software. Both methods assumed that the two parents of the population were each heterozygous at the S-locus and did not share any S-alleles with each other, resulting in a population with four distinct S-alleles and therefore four distinct S-genotypes. Each method is described in detail below. Code to replicate all methods may be found on GitHub (see data availability statement).

Approach 1: A hill-climbing algorithm: The first S-allele determination method used a simple hill-climbing algorithm to fit genotypes, given a user-generated set of dominance relationships among alleles. To start, the user hypothesizes a set of crossing relationships, dictating which of the sixteen possible pairings of genotypes will and will not be able to successfully mate. A random genotype is then assigned to each individual in the population. One arbitrarily selected individual is always set to a predetermined genotype to enable comparison of different runs of the algorithm. Each mated pair in the dataset is then scored using the user-supplied crossing relationships. A mismatch between the observed and predicted outcomes results in a score between 0.5 and 1, determined by the user. A match results in a score of one minus the previously mentioned score. Scores for the entire dataset are then summed. Next, the genotype of one individual is randomly changed, and the dataset rescored. If the total score decreases, then that set of genotypes is kept; if not, the algorithm returns to the previous position.

The cycle is repeated, stopping if 1,000 changes are attempted without decreasing the score. For this experiment, the algorithm was run 4,000 times for each set of dominance relationships, with the solution to the lowest scoring run considered optimal genotype assignments for a given dominance relationship. The lowest-scoring genotype assignments across all dominance relationships were then used for QTL mapping. We compared the results of 48 dominance

relationships, each of which met three criteria: 1) individuals sharing both alleles must be

252   incompatible, 2) individuals sharing no alleles must be compatible, and 3) there must be at least

one combination of genotypes that was asymmetrical (compatible when one individual was used

254   as a female and incompatible when the other was used as a female), as this was observed in the

crossing data.

This method has both the advantage and disadvantage of being highly parameterizable. This

258   makes it flexible, and gives the user a high degree of control, but may take many attempts to find

the right combination of parameters to develop a solution. This method may be employed using a

260   personal computer, however the reduction in parallelization necessary to achieve this may make

it take too long to be useful. In addition, as a hill-climbing method there is no guarantee that this

262   approach will find a global maximum.

264   *Approach 2: A Markov Chain Monte Carlo algorithm:* The second method employed sets of

extreme gradient boosting decision trees with Markov chain Monte Carlo (MCMC) algorithms to

266   infer S-locus genotypes. For each MCMC chain, an initial set of genotypes was created by first

assigning a random heterozygous S-genotype to one random individual. The genotype of an

268   individual that had been mated to the initial individual was then determined. If the cross was

incompatible, the second individual's genotype could share one or two alleles with the initial

270   individual's genotype, with equal probability. If the cross was compatible, the second

individual's genotype could share either zero or one allele with the initial plant's genotype, with

272   equal probability. The procedure was then repeated by choosing, at random, an individual that

did not yet have a genotype assigned and that was crossed to the most recent individual assigned

274   a genotype. If all individuals that were crossed to the most recently assigned individual had an

assigned genotype, a random individual that did not yet have an assigned genotype was selected

276   and the process started again until all individual were assigned genotypes.

278   Maternal and paternal S-locus genotypes were then treated as predictive variables with cross

success used as a binary response variable. Constraints were placed on predictive variables so

280   that matching maternal and paternal genotypes were not allowed to interact with one another, as

crosses between individuals with matching genotypes should always be incompatible. At each

282   step of an MCMC chain, model performance was measured as error of a logistic regression for
        classification using an extreme gradient boosting model with four-fold cross validation with 200
284   iterations and maximum tree depth of four splits, using the R package "xgboost" (Chen &
        Guestrin, 2016). A single genotype was altered at each MCMC chain step, using the same sets
286   of probabilities used to construct the initial set. The newly proposed set of genotypes was
        accepted if the ratio of errors from the proposed genotype set to the former genotype set was
288   greater than a randomly drawn number bound by zero and one.

290   Each MCMC chain, starting from a unique set of genotypes, was allowed to explore parameter
        space for 96 hours (~ 1 million steps). The set of genotypes that produced the smallest error from
292   each chain was then used as the starting condition for hill-climbing algorithms. We used the
        same extreme gradient boosting model conditions described above, however at each step
294   proposed genotypes were only accepted if they produced an error that was less than the previous
        set of genotypes. The hill-climbing algorithms each ran for 96 hours. For each chain, the set of
296   genotypes that produced the smallest error, calculated as the percentage of crosses classified
        incorrectly, were recorded. A cross was predicted to be a success if the estimated logistic
298   regression probability was greater than 0.5. The lowest error models were considered the best
        candidates for S-genotype determination. This method has the advantage of less dependence on
300   user decisions, making it more repeatable and facilitating the exploration of a wider space of
        possible solutions. However, the resources required by this method mean that it generally cannot
302   be employed on a personal computer.

304   *Mapping the S-locus from inferred S-locus genotypes:* Assigned alleles from the two inference
        methods were then used as phenotypes for QTL mapping, using the "qtl" R package (Broman *et
306   al*., 2003), with the population treated as a four-way testcross. Marker genotypes were imputed
        using the "sim.geno" function, and QTL were identified using the "scanone" function to conduct
308   interval mapping using the EM algorithm. Self-incompatibility allele was treated as a binary
        trait, with one allele from each parent arbitrarily coded as "1" and the other as "0", and the two
310   parental alleles mapped separately. For the MCMC method, all 96 of the inferred genotype sets
        were mapped, with the maximum LOD score produced by each set serving as a criterion to
312   differentiate the sets. Eight genotype sets produced by the hill-climbing method were also

mapped. Significance thresholds were determined independently for each set of genotypes used

314   for mapping; only QTL with an error probability less than 5% were considered significant. This

threshold ranged from a LOD score of 3.6 to 4.2. Finally, a Bayesian credible interval (similar to

316   a confidence interval) around any identified QTL was calculated, using the "bayesint" function.

318   **Synteny with related species:** To help elucidate the relationship between the *S. integrifolium* S-

locus and other *Asteraceae*, the genomic contigs associated with the loci that comprise this

320   linkage map were aligned to the annual sunflower (*Helianthus annuus*, variety 'HA412', version

HOv1.1) (Badouin *et al.*, 2017) and lettuce (*Lactuca sativa*, variety 'Salinas' , version 7) (Reyes-

322   Chin-Wo *et al.*, 2017) genome assemblies using BLASTN (Altschul *et al.*, 1990). Syntenic

regions were then identified using the R package "syntR" (Ostevik *et al.*, 2020). For synteny

324   analysis, all BLASTN alignments for a given *S. integrifolium* query sequence that had a bitscore

greater than 90% of the maximum bitscore for that query sequence were used.

Identification of candidate genes: For regions identified as syntenic to the *S. integrifolium* S-

328   locus, gene annotations for sunflower were obtained from the genome assembly HA412v1

(Badouin *et al.*, 2017). InterPro taxonomy was searched for each gene in the region to identify

330   plausible candidates by shared terms with S-genes in other species. The male and female

determinants from *Brassica oleracea* (*Brassicaceae*) and *Petunia hybrida* (*Solanaceae*), as well

332   as the female determinant of *Papaver rhoeas* (*Ranunculales*) chosen as representatives for

comparison (The UniProt Consortium, 2019). In addition, defensin-like proteins were considered

334   a proxy for the *Ipomoea trifida* (*Convolvulaceae*) male determinant (Rahman *et al.*,

2007). These species were selected because their self-recognition genes are relatively well

336   characterized and they represent a broad range of Eudicot diversity, including both SSI and GSI

systems. Any genes homologous to known CRPs were also considered potential candidates

338   (Marshall *et al.*, 2011). The best homologs for a subset of these candidates were then identified

in the *S. integrifolium* reference transcriptome (Raduski *et al.*, 2021), and their expression was

340   measured in a previously-published set of whole-plant seedling transcriptomes from a diversity

panel of 73 wild collected *S. integrifolium* accessions (Raduski *et al.*, 2021). Expression was

342   quantified in total transcripts per million (TPM), TPM as a percentile of total expressed genes,

and TPM as a percentile of a reference set of conserved single copy genes, known as BUSCO

344   (Simão *et al.*, 2015). We hypothesized that strong S-locus gene candidates would not be highly expressed in seedling vegetative tissue (Williams *et al*., 2014).

Additionally, the coding region sequence for the *B. oleracea* S-receptor kinase protein (Stein *et*
348   *al*., 1991) was aligned to the lettuce and sunflower genomes using BLAST. These alignments were then compared to the lettuce and sunflower regions syntenic to the *S. integrifolium* S-locus.

RESULTS

Linkage Map: The linkage map contains 198 markers, spanning 1,049 centimorgans and
divided into 7 linkage groups (Fig. 2, Table S1). This is consistent with the observation that *S. integrifolium* has seven chromosomes (Settle, 1967). The average distance between markers is
5.5 cM, with 29 gaps greater than 10 cM.

S-locus mapping: Of the 268 crosses completed, 53% were incompatible, with a seed set value less than 20%, and 47% were compatible. These frequencies were not significantly different
from an equal occurrence of compatible and incompatible crosses (df = 1, χ^2 = 0.956, P = 0.328). This indicates there is a hierarchy of dominance between alleles in this population—if all alleles
were codominant a ratio of 25% compatible to 75% incompatible would be expected. Of the 126 reciprocal pairs of genotypes crossed, 30% were compatible in both crossing directions (both
individuals could be used as male and female), 34% were incompatible, and 36% were asymmetrical, or compatible when one individual was used as a female and incompatible when
the other was used as a female. The presence of asymmetrical crosses indicates that the dominance relationship between some allele pairs differs from anthers to stigma. These
observations confirm that *S. integrifolium* has a sporophytic SI system, as dominance relationships are not observed in gametophytic systems (Breton *et al*., 2014). Figure 3 shows the
distribution of seed set values for reciprocal crosses, numerical data for all crosses may be found in Table S2.

Single marker regression for male/ female interaction: The interaction of male and female
374   marker genotype was found to be a significant (P > 0.001) predictor of crossing success for four

markers, located at two regions on the linkage map: markers "272299" and "120685" located at

376 18.9 and 9 cM, respectively, on linkage group six (P = 7.57 x 10^{-8}, P = 1.46 x 10^{-4} df = 259 for both), and markers "185443" and "118778", located at 126.33 cM and 120.7 cM, respectively,

378 on linkage group two (P = 3.22 x 10^{-5}, P = 3.31 x 10^{-4}, df = 259 for both).

380 *Hill-climbing algorithm method:*

Of the 48 dominance relationships tested using the hill-climbing algorithm, nine were used in

382 QTL mapping. Of these, seven produced at least one significant LOD peak, all located on

linkage group 6. The highest of these (LOD = 5.3) was located at 18.9 cM, and all but one of the

384 other six were located between 9-31 cM.

386 *MCMC method:* Of the 192 genotype sets produced by the MCMC method (96 models, with a

separate genotype set for each of the two parental alleles), 30 produced at least one significant

388 LOD peak when used for QTL mapping, covering five linkage groups. Eleven of these genotype

sets produce a significant LOD peak at either 18.9 or 27.7 cM on linkage group 6, and a further

390 five were located elsewhere between 0-31 cM on linkage group 6. No other linkage group

contained more than five of the 30 significant LOD peaks. The highest LOD peak produced by a

392 single MCMC model was 7.56, located at 18.9 cM on linkage group six. This model also had the

lowest error, and so we have selected it as the best result from the MCMC method. This set of

394 genotype assignments is illustrated in Figure 4.

396 *Consensus map location:* Of the three methods used to map the S-locus, the MCMC method

produced the strongest association between a genomic region and S-allele. The peak of this QTL

398 was located at 18.9 cM on linkage group six, with a 95% probability Bayes credible interval

from 0 to 27.7 cM. Broadly speaking, the results of the hill-climbing method agree with this

400 region, although the credible interval for several of those results extend to 31 cM. Additionally,

the most significant single-marker regression association between the interaction of male and

402 female genotype with cross success was found at 18.9 cM. Taken together, we conclude that the

S. integrifolium S-locus is located between 0 and 31 cM on linkage group six, with the marker

closest to the locus likely located at 18.9 cM.

S-locus synteny with related species: Synteny was found between this the putative *S. integrifolium* S-locus region and both the sunflower and lettuce genomes. In sunflower, synteny

was found with two chromosomes—the *S. integrifolium* 0-18.9 cM region was found to align with chromosome 17, from 230.7 Mb to 262.67 Mb, and the 18.9- 31cM region was found to

align with chromosome 3, from 133.21 Mb to 149.9 Mb. In lettuce, the 0-9 cM region aligned with chromosome 4, from 18.13 Mb to 27.29 Mb, and the 18.9-31 cM region aligned with

chromosome 9, from 144.19 Mb to 147.34 Mb. The syntenic region on sunflower chromosome 17 does not appear to overlap with the putative self-incompatibility breakdown QTL found on

that chromosome by Gandhi *et al.* (2005), which we estimate was somewhere between 42 and 98 Mb, based on the alignment of SSR marker primer sequences to the sunflower genome

assembly.

Identification of candidate genes: Of the 642 genes located within the sunflower genomic regions syntenic with the *S. integrifolium* S-locus, 42 shared at least one InterPro term with

either the male or female determinants of self-incompatibility in *B. oleracea* or *P. hybrida,* the female determinant of *P. rhoeas*, or the male determinant of *I. trifida*. Thirty of these genes were,

like *B. oleracea* SRK, protein kinases, which is not a specific enough homology to infer a gene candidate. The other 12 genes were, like the female determinant in *P. hybrida*, F-box genes. In

addition, one gene was similar to "STIG1", a CRP. The expression of this gene, as well as two of the 12 F-box genes, were then measured in the 73 diversity panel seedling transcriptomes. One

of these F-box genes was expressed in 65 of the seedling transcriptomes, with TPM as a percentile of BUSCO genes ranging from 5.4 to 79.5. The other was expressed in 42 of the

seedlings, with TPM as a percentile of BUSCO genes ranging from 0.09 to 41.2. The STIG1-like gene was expressed in five of the seedling transcriptomes; expression varied widely but was

always relatively low (Table 1).

The best alignment for *B. oleracea* SRK in sunflower was found on chromosome 11, while the best alignment in lettuce was found on chromosome 7. Neither of these alignments are near any

regions syntenic with the putative *S. integrifolium* S-locus in either species.

DISCUSSION

The described S-locus mapping methods are effective and efficient: This study represents one of the few successful identifications of a genomic region containing the S-locus in an *Asteraceae* species. This result is valuable for its own sake, as the comparison of syntenic genomic regions between *S. integrifolium* and other species may help to answer questions about the evolution of SSI systems in other *Asteraceae*. In addition, this study lays the groundwork for future efforts to clone the *S. integrifolium* S-locus.

In this experiment, we were able to predict the S-locus genotype for 84 individuals by conducting 268 crosses, with an average of 3.2 crosses per individual. This method did not require all 84 individuals to be crossed with a single tester, and so could be applied to a pre-existing full-sibling family without necessitating clonal propagation of any individuals. As a point of comparison, we estimate that the mapping of the S-locus in chicory, which combined a twenty-two plant diallel with testcrossing, required approximately 3,000 crosses (Gonthier *et al.*, 2013). That approach, which also required the clonal propagation of tester genotypes, was used to assign an S-genotype to approximately 350 individuals (multiple replications of each cross were performed). The chicory mapping effort produced a more accurate determination of S-genotype and thus a narrower S-locus QTL region than our study, but with the trade-off of requiring more crosses per individual and the clonal replication of tester genotypes. Additionally, our method requires less time to complete, as testcrossing cannot be conducted until the diallel is completed, the results interpreted, and selected testers clonally propagated. Finally, our method does not require every cross to be replicated, as the analysis relies on inference methods that are relatively robust to error in the data.

Our method should be readily applicable to any other species that meet three criteria: 1) They possess an SSI system, 2) large full sibling families can be produced, and 3) individuals can readily be used as both a male and female parent in three to five matings. Among numerous other species, self-incompatible members of the genus *Helianthus* could be excellent candidates for this method and could help to confirm our results. Applying this method to phylogenetically diverse *Asteraceae* species could evaluate the conservation of the method of SSI in the family. Additionally, with some modifications, our framework could be used for GSI species.

Evaluation of potential S-locus candidate genes: Of the 642 annotated or predicted genes

470   present in the sunflower genomic regions syntenic to the putative S-locus, 43 show potential

similarity to known S-genes. Of these, two groups of genes were considered particularly

472   interesting, and are detailed below. It is important to note that because virtually nothing is known

about the molecular mechanisms that underlie SSI in the *Asteraceae*, any discussion of potential

474   candidate genes based on annotations or similarity to S-related genes in other systems is at best

informed speculation. However, we see this as an important step towards identifying the genes

476   that control SSI in the *Asteraceae*, and therefore believe it is worth pursuing.

478   One potential candidate is the sunflower gene Ha3_00036110, which is annotated as a putative

member of the "Stigma Specific Protein 1" like, or STIG1-like, group of genes. In sunflower,

480   this gene is found on the chromosome 3 region syntenic with the 18.9-31 cM segment of the

putative *S. integrifolium* S-locus. In lettuce, the best BLAST alignment for Ha3_00036110 is on

482   chromosome 9, within the region syntenic to the putative *S. integrifolium* S-locus, implying that

this region may be conserved between the three species.

The best characterized member of this family, STIG1, encodes a small, cysteine-rich protein

486   (CRP). In tomato (*Solanum lycopersicum*), this protein is primarily found in stigma exudate, and

binds a pollen-specific kinase to promote pollen tube growth (Huang *et al*., 2014). In tomato,

488   silencing either STIG1 or the pollen kinase resulted in slowed pollen tube growth and reduced

seed set (Huang *et al*., 2014). More broadly speaking, as CRPs, STIG1-like genes are members

490   of a gene class that also includes the male determinant of SSI in the *Brassicaceae* and the

Convolvulaceae, and the female determinant of GSI in the genus *Papaver*. Based on this

492   similarity to other SI genes, the function of STIG1, and the location of this gene within the

sunflower genomic region syntenic to the *S. integrifolium* S-locus, we speculate that

494   Ha3_00036110 is a signaling protein that controls the female determinant of SSI in *S.

integrifolium*, with differential binding between different alleles of the STIG1-like protein and an

496   unidentified pollen kinase resulting in compatibility or incompatibility. If this is correct, a

possible candidate for this pollen kinase may be the gene Ha3_00036112, a serine/threonine

protein kinase located 109 Kbp downstream from Ha3_00036110.

The best *S. integrifolium* homolog for Ha3_00036110 was only expressed in five of 73 seedling transcriptomes, and Ha3_00036112 was found to be expressed in four of 73 seedlings. The two

genes were never expressed in the same individual, and expression was relatively low, never exceeding the 33rd percentile of BUSCO genes (Table 1). The low level of seedling expression

for these genes further supports their candidacy, as S-gene expression is expected to be primarily limited to floral tissue (Williams *et al.*, 2014). Expression in other tissues may still occur at low

levels, either due to gene expression noise or as-yet uncharacterized uses for the gene product. For example, in tea (*Camellia sinensis*), an S-gene candidate was found to be expressed in leaf

tissue at approximately 20% the level it was expressed in style tissue (Zhang *et al.*, 2016).

Another set of potential candidate genes are found on the sunflower chromosome 17 region syntenic with the 0-18.9 cM segment of the putative *S. integrifolium* S-locus. This region

contains 12 F-box genes, resembling known gametophytic self-incompatibility (GSI) systems. For example, in *Petunia*, 17 tightly linked F-box genes serve as the male determinant of GSI

(Williams *et al.*, 2015). However, no genes in this sunflower region share any annotation terms with known S-RNAse genes (The UniProt Consortium, 2019), which serve as the female

determinant in GSI systems. In addition, two of these genes that were arbitrarily selected as representatives were found to be expressed in more than half of available seedling

transcriptomes, further diminishing the likelihood that they control the self-incompatibility response. Because so little is known about the molecular basis of the *Asteraceae* SSI response,

these genes cannot be completely discounted, but they appear to be less likely candidates than the STIG1-like gene.

In addition to providing candidates for genes involved in *S. integrifolium* self-recognition, this

524   study suggests that SRK-like genes are not involved in *S. integrifolium* SSI, as the best homologues for the *B. oleracea* SRK in the lettuce and sunflower genomes are not found in or

526   near the regions of those genomes syntenic with the putative *S. integrifolium* S-locus. This finding adds evidence to the theory that specific genes that underlie *Asteraceae* SSI are different

528   than those in the *Brassicaceae* (Allen & Hiscock, 2008).

Implications for domestication and breeding: We expect the results of this study to facilitate the domestication of *S. integrifolium*. The availability of a genetic map, associated with particular restriction enzymes, supports the relatively inexpensive GBS genotyping of large numbers of progeny; both because these enzymes are now known to be useable in *S. integrifolium*, and because it is reasonably likely that some of the same loci would be recovered if these enzymes were applied to a different population. This may support the implementation of marker-based selection, marker-based pedigree development, and genome-wide selection. Additionally, this genetic map will assist in the anchoring and orientation of future genome assemblies.

Identifying a map location for the S-locus may also contribute to practical breeding efforts. If molecular marker data is routinely available for individuals within a breeding program, it may be possible to predict whether any two plants will be able to successfully mate, through the direct identification of particular S-alleles or ancestral haplotype blocks in the S-region. This information could save effort by excluding crosses that would not be successful. Perhaps more importantly, this would allow for the S-allele diversity of a given population to be monitored and maximized. It is likely that at least some *S. integrifolium* cultivars will take the form of synthetic populations, where a set of superior genotypes are intermated and their progeny form a distinct variety that may be reproduced for several generations. If a synthetic population is released that contains a low number of S-alleles, its long-term fecundity may be adversely affected by the limited number of individuals that are able to intermate. Alternatively, as *S. integrifolium* is known to express moderate to severe inbreeding depression for a number of traits (Price *et al.*, 2021), it is possible that S-allele characterization information could be used increase long-term productivity by limited mating among relatives for several generations within synthetic populations.

Overall, we anticipate that the availability of a genetic map and identification of the self-incompatibility locus will support efforts to domesticate *S. integrifolium* as a crop that will help to enable sustainable agricultural systems.

ACKNOWLEDGMENTS

562 Funding for this work was provided by The Perennial Agriculture project in conjunction with
The Land Institute and the Malone Family Land Preservation Fund, the United States

564 Department of Agriculture's National Institute of Food and Agriculture Grant no. 2019-67011-
29607 to JHP, the Minnesota Department of Agriculture - Forever Green Agricultural Initiative,

566 and NSF grant #1737827 Dimensions US-China to YB. The authors thank Shannon Lee
Anderson, Karen Beaubein, and Jill Ekar for their help in completing the controlled crosses for

568 this experiment. In addition, the authors thank Dr. Kevin Dorn for assistance in developing a
GBS protocol, Dr. Adam Herman for bioinformatics advice, and Dr. Owen Beisel for inspiring

570 the hill-climbing algorithm method. Finally, the authors acknowledge the Minnesota
Supercomputing Institute (MSI) at the University of Minnesota for providing computing

572 resources that contributed to the analysis of this study.

574 AUTHOR CONTRIBUTIONS

The effort to develop a linkage map was initiated by KPS, DLVT, and YB, who also secured

576 initial funding. DLVT created the mapping population, contributed text and visualization, and
provided feedback in writing. JHP constructed the linkage map, initiated and designed the S-

578 locus mapping work, and developed the HC method. ARR developed the MCMC method. JHP
and ARR conducted data analysis and wrote the manuscript. KPS and YB provided supervision,

580 and feedback on analysis, writing, and visualization.

582 DATA AVAILABILITY

All code to replicate analyses may be found on GitHub, at https://github.com/UMN-

584 BarleyOatSilphium/SilphiumSLocus. All sequence data may be accessed under BioProject
PRJNA695552.

REFERENCES

Allen AM, Hiscock SJ. 2008. Evolution and phylogeny of self-incompatibility systems in

594 angiosperms. In: Franklin-Tong VE, ed. *Self-incompatibility in flowering plants: Evolution, Diversity and Mechanisms*. Heidelberg, DE: Springer-Verlag, 73-101.

Allen AM, Thorogood CJ, Hegarty MJ, Lexer C, Hiscock SJ. 2011. Pollen-pistil interactions

598 and self-incompatibility in the Asteraceae: new insights from studies of *Senecio squalidus* (Oxford ragwort). *Annals of Botay* **108(4):** 687-698.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search

602 tool. *J. Mol. Biol.* **215**:403-410

604 **Badouin H, Gouzy J, Grassa CJ, Murat F, Staton SE, Cottret L, Lelandais-Brière C., Owens GL, Carrère S, Mayjonade B et al. 2017.** The sunflower genome provides insights into

606 oil metabolism, flowering and Asterid evolution. *Nature* **546(7656)**: 148-152.

608 **Bod'ová K, Priklopil T, Field DL, Barton NH, Pickup M. 2018.** Evolutionary pathways for the generation of new self-incompatibility haplotypes in a nonself-recognition system. *Genetics*

610 **209(3):** 861-883.

Brennan AC, Harris SA, Hiscock SJ. 2006. The population genetics of sporophytic self-

612 incompatibility in Senecio squalidus L. (Asteraceae): the number, frequency, and dominance interactions of S alleles across its British range. *Evolution* **60(2):** 213-224.

Breton CM, Farinelli D, Shafiq S, Heslop-Harrison JS, Sedgley M. , Bervillé AJ. 2014. The

616 self-incompatibility mating system of the olive (*Olea europaea* L.) functions with dominance between S-alleles. *Tree Genetics & Genomes* **10(4)**: 1055-1067.

Broman KW, Wu H, Sen Ś, Churchill GA. 2003. R/qtl: QTL mapping in experimental crosses.
620 *Bioinformatics* **19**:889-890

622 **Camargo LEA, Savides L, Jung G, Nienhuis J, Osborn TC. 1997.** Location of the self-
incompatibility locus in an RFLP and RAPD map of *Brassica oleracea*. *Journal of Heredity*
624 **88(1**): 57-60.

626 **Catchen J, Hohenlohe P, Bassham S, Amores A, Cresko W. 2013.** Stacks: an analysis tool set
for population genomics. *Molecular Ecology* **22(11)**:3124-40
628

Chen T, Guestrin C. 2016. XGBoost: A Scalable Tree Boosting System. In: *Proceedings of the*
630 *22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining* (*KDD '16*). New York, NY, USA: Association for Computing Machinery, 785–794.
632

DeHaan LR, Van Tassel DL, Anderson JA, Asselin SR, Barnes R, Baute GJ, Cattani DJ,
634 **Culman SW, Dorn KM, Hulke BS et al. 2016.** A pipeline strategy for grain crop
domestication. *Crop Science* **56(3)**: 917-930.
636

Edh K, Widén B, Ceplitis A. 2009. Molecular population genetics of the SRK and SCR self-
638 incompatibility genes in the wild plant species *Brassica cretica* (*Brassicaceae*). *Genetics* **181(3)**: 985-995.
640

Gonthier L, Blassiau C, Mörchen M, Cadalen T, Poiret M, Hendriks T, Quillet MC. 2013.
642 High-density genetic maps for loci involved in nuclear male sterility (NMS1) and sporophytic
self-incompatibility (S-locus) in chicory (*Cichorium intybus* L., *Asteraceae*). *Theoretical and*
644 *applied genetics* **126(8)**: 2103-2121.

646   **Harkness A, Brandvain Y. 2020.** Nonself-recognition-based self-incompatibility can alternatively promote or prevent introgression. *bioRxiv.* doi: 10.1101/2020.09.29.318790

Harkness A, Goldberg EE, Brandvain Y. 2021. Diversification or collapse of self-
650   incompatibility haplotypes as a rescue process. *The American Naturalist*. doi:10.1086/712424

652   **Hiscock SJ. 2000.** Genetic control of self-incompatibility in *Senecio squalidus* L.(*Asteraceae*): a successful colonizing species. *Heredity* **85(1)**: 10-19.

Hiscock SJ, Tabah DA. 2003. The different mechanisms of sporophytic self-incompatibility.
656   *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* 358(1434): 1037–1045.

Huang WJ, Liu HK, McCormick S, Tang WH. 2014. Tomato Pistil Factor STIG1 Promotes in
660   Vivo Pollen Tube Growth by Binding to Phosphatidylinositol 3-Phosphate and the Extracellular Domain of the Pollen Receptor Kinase LePRK2. *The Plant Cell* **26(6)**: 2505–2523.

Fujii S, Takayama S. 2018. Multilayered dominance hierarchy in plant self-incompatibility.
664   *Plant reproduction* 31(1): 15-19.

666   **Gandhi SD, Heesacker AF, Freeman CA, Argyris J, Bradford K, Knapp SJ. 2005.** The self-incompatibility locus (S) and quantitative trait loci for self-pollination and seed dormancy in
668   sunflower. *Theoretical and Applied Genetics* **111(4)**: 619-629.

670   **Larson S, DeHaan L, Poland J, Zhang X, Dorn K, Kantarski T, Anderson J, Schmutz J, Grimwood J, Jenkins J et al. 2019.** Genome mapping of quantitative trait loci (QTL)
672   controlling domestication traits of intermediate wheatgrass *(Thinopyrum intermedium)*. *Theoretical and Applied Genetics* **132(8)**: 2325-2351.

Marshall E, Costa LM, Gutierrez-Marcos J. 2011. Cysteine-rich peptides (CRPs) mediate

676 diverse aspects of cell–cell communication in plant reproduction and development. *Journal of Experimental Botany* **62(5)**: 1677-1686.

Kandel H, Hulke B, Ostlie M, Schatz B, Aberle E, Bjerke K, Eriksmoen E, Kraklau A,

680 **Effertz J, Rickertsen J. et. al. 2019.** *North Dakota Sunflower Variety Trial Results for 2019 and Selection Guide.* Fargo, ND, USA: North Dakota State University Extension.

682 **Koseva B, Crawford DJ, Brown KE, Mort ME, Kelly JK. 2017.** The genetic breakdown of sporophytic self-incompatibility in *Tolpis coronopifolia* (*Asteraceae*). *New Phytologist* **216(4)**:

684 1256-1267.

686 **Kubo K, Entani T, Takara A, Wang N, Fields AM, Hua Z. 2010.** Collaborative non-self recognition system in S-RNase-based self-incompatibility. *Science* **330(6005):** 796-799.

Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

arXiv:1303.3997 [q-bio.GN].

Price JH, Brandvain Y, Smith KP. 2021. Measurements of lethal and non-lethal inbreeding depression inform the *de novo* domestication of *Silphium integrifolium*. *American Journal of*

Botany, in press.

Ostevik KL, Samuk K, Rieseberg LH. 2020. Ancestral Reconstruction of Karyotypes Reveals an Exceptional Rate of Nonrandom Chromosomal Evolution in Sunflower. *Genetics* **214(4)**:

1031-1045.

Ouellette LA, Reid RW, Blanchard SG, Brouwer CR. 2018. LinkageMapView—rendering high-resolution linkage and QTL maps. *Bioinformatics* **34(2)**: 306-307.

R Core Team. 2019. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

706   **Raduski AR, Herman A, Pogoda C, Dorn KM, Van Tassel DL, Kane N, Brandvain Y. 2021.**
Patterns of genetic variation in a prairie wildflower, *Silphium integrifolium*, suggest a non-prairie
708   origin and untapped variation available for improved breeding. *American Journal of Botany*
108(1): 145-158.

710   **Rahman MH, Uchiyama M, Kuno M, Hirashima N, Suwabe K, Tsuchiya T, Kagaya Y,
Kobayashi I, Kakeda K, Kowyama Y. 2007.** Expression of stigma-and anther-specific genes
712   located in the S locus region of *Ipomoea trifida*. *Sexual Plant Reproduction* **20(2)**: 73-85.

714   **Reinert S, Van Tassel D, Schlautman B, Kane N, Hulke B. 2019.** Assessment of the
biogeographical variation of seed size and seed oil traits in wild *Silphium integrifolium* Michx.
716   genotypes. *Plant Genetic Resources: Characterization and Utilization* **17(5)**: 427-436.

718   **Reinert S, Price JH, Smart BC, Pogoda CS, Kane NC, Van Tassel D, Hulke B. 2020.** Mating
compatibility and fertility studies in an herbaceous perennial Aster undergoing de novo
720   domestication to enhance agroecosystems. *Agronomy for Sustainable Development* **40**:27.

722   **Reyes-Chin-Wo S, Wang Z, Yang X, Kozik A, Arikit S, Song C, Xia L, Froenicke L,
Lavelle DO, Truco MJ et al. 2017.** Genome assembly with in vitro proximity ligation data and
724   whole-genome triplication in lettuce. *Nature Communications* **8(1)**: 1-11.

726   **Rindos D. 1984.** *The Origins of Agriculture: An Evolutionary Perspective*. Orlando, FL, USA:
Academic Press.

Schiffner S, Jungers JM, Hulke BS, Van Tassel DL, Smith KP, Sheaffer CC. 2020. Silflower

730   seed and biomass responses to plant density and nitrogen fertilization. *Agrosystems, Geosciences & Environment* 3:e20118.

Sedbrook JC, Phippen WB, Marks MD. 2014. New approaches to facilitate rapid

734   domestication of a wild plant to an oilseed crop: example pennycress (*Thlaspi arvense* L.). *Plant Science* **227**:122-132.

Settle WJ. 1967. The chromosome morphology in the genus *Silphium* (*Compositae*). *The Ohio*

738   *Journal of Science* **67(1)**: 10-19.

740   **Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015.** BUSCO:

assessing genome assembly and annotation completeness with single-copy orthologs.

742   *Bioinformatics* **31(19)**: 3210-3212.

Stam P. 1993. Construction of integrated genetic linkage maps by means of a new computer

744   package: JoinMap. *The Plant Journal* **3**: 739-744.

Stein JC, Howlett B, Boyes DC, Nasrallah ME, Nasrallah JB. 1991. Molecular cloning of a

746   putative receptor protein kinase gene encoded at the self-incompatibility locus of *Brassica oleracea*. *Proceedings of the National Academy of Sciences* **88(19)**: 8816-8820.

748   **Tomita RN, Fukami K, Takayama S, Kowyama, Y. 2004.** Genetic mapping of AFLP/AMF-

derived DNA markers in the vicinity of the self-incompatibility locus in *Ipomoea trifida*. *Sexual*

750   *Plant Reproduction* **16(6)**: 265-272.

The UniProt Consortium. 2019. UniProt: a worldwide hub of protein knowledge. *Nucleic Acids*

752   *Research* 47(D1): D506–D515

754   **Van Tassel DL, DeHaan LR, Cox TS. 2010.** Missing domesticated plant forms: Can artificial

selection fill the gap? *Evolutionary Applications* **3(5–6)**: 434–452.

756 **Van Tassel DL, Albrecht KA, Bever JD, Boe AA, Brandvain Y, Crews TE, Gansberger M, Gerstberger P, González-Paleo L, Hulke BS. et al. 2017.** Accelerating Silphium

758 Domestication: An Opportunity to Develop New Crop Ideotypes and Breeding Strategies Informed by Multiple Disciplines. *Crop Science* **57(3)**: 1274–1284.

760 **Williams JS, Der JP, dePamphilis CW, Kao TH. 2014.** Transcriptome analysis reveals the same 17 S-locus F-box genes in two haplotypes of the self-incompatibility locus of *Petunia*

762 *inflata*. *The Plant Cell*, **26(7)**: 2873-2888.

Williams JS, Wu L, Li S, Sun P, Kao TH. 2015. Insight into S-RNase-based self-

764 incompatibility in *Petunia*: recent findings and future directions. *Frontiers in Plant Science* **6**:41.

Zhang CC, Wang LY, Wei K, Wu LY, Li HL, Zhang F, Cheng H, Ni DJ. 2016.

766 Transcriptome analysis reveals self-incompatibility in the tea plant (*Camellia sinensis*) might be under gametophytic control. *BMC Genomics* **17:**359.

768 SUPPORTING INFORMATION

Table S1: The linkage map in tabular form.

770 Table S2: All crossing data used in this experiment.

Table 1: Expression of two S-gene candidates in a set of whole-plant seedling transcriptomes from a diversity panel of 73 wild collected *Silphium integrifolium* accessions. Expression is for *S. integrifolium* homologs of the named sunflower genes.

Sunflower gene	Description	Plant ID	Transcripts per million (TPM)	TPM as percentile of all genes	TPM as a percentile of BUSCO genes
Ha3_00036110	"Stig1-like" gene	C7	0.13	0.076	0.052
		D8	2.26	0.352	0.332
		E4	0.57	0.256	0.206
		E5	0.57	0.264	0.237
		H2	0.2	0.130	0.096
Ha3_00036112	serine/threonine protein kinase	B3	0.27	0.177	0.130
		B9	1.14	0.315	0.305
		E8	1.37	0.340	0.329
		H7	0.07	0.027	0.019

Figure 1: Visualization of the crossing design used in this experiment. Each circle represents one plant, with black circles representing plants that were used for crosses, and gray plants representing other members of the mapping population that were not selected. The dashed box represents one of the small diallels that formed the basis for the design—twenty of these were used for this experiment. Arrows represent matings, with each arrow pointing from the male parent to the female parent.

Figure 2: Linkage map for *Silphium integrifolium*. The black bar on linkage group 6 represents the putative S-locus QTL, with the circle showing the LOD peak and the extent of the bar showing a composite of the 95% Bayes credible intervals for several mapping methods. Map visualization was performed using "LinkageMapView" (Ouellette *et al.*, 2018).

Figure 3: Distribution of seed set values for reciprocal matings (pairs of individuals mated, with each individual used as both a male and female). Each dot represents the seed set value of one plant, with lines connecting mated pairs. The individuals with the higher seed set value in the pair are placed on the left side of the chart.

Figure 4: Representation of all possible matings that could have been completed for this study, with each square in the heatmap representing one pairing. Red or blue squares represent the crosses that were actually made, with red representing incompatible crosses, and blue representing compatible. Individuals are grouped by S-locus genotype, as assigned by the best MCMC genotype inference model. "A" and "B" refer to the S-alleles from one parent of the population, and "C" and "D" to the alleles from the other.