Adaptive Behavior and the Role of Primary Somatosensory Cortex

Christian Waiblinger[1], Peter Y Borden[1], Garrett B Stanley[1]

[1]Wallace H Coulter Department of Biomedical Engineering, Georgia Institute of Technology and Emory University, Atlanta, GA, USA

With 33 text pages, 5 main figures, 4 supplemental figures
Word counts: Abstract 138, Intro 503, Results: 3382, Discussion 1149

*Correspondence: Garrett B Stanley

Coulter Department of Biomedical Engineering
Georgia Institute of Technology & Emory University
313 Ferst Drive
Atlanta GA 30332-0535
USA

Phone: 404-385-5037
Fax: 404-385-5044
Email: garrett.stanley@bme.gatech.edu

Acknowledgements

C. W. was supported by a fellowship from the Deutsche Forschungsgemeinschaft (GZ: WA 3862/1-1) and NIH grant U01NS094302. P.Y.B. was supported by an NIH NRSA pre-doctoral fellowship F31NS09869. G. B. S. was supported by NIH Brain Grants R01NS104928 and U01NS094302. We thank Cornelius Schwarz for helpful feedback on the manuscript. We also acknowledge April Reedy with the Emory University Integrated Cellular Imaging Core.

ABSTRACT

Behavioral experience and flexibility are crucial for survival in a constantly changing environment.

What are the neuronal processes that selectively transform dynamic sensory information into an

appropriate behavioral response, and how do these processes adapt to changes in the

environment? Here, we use voltage imaging to measure signals in primary somatosensory cortex

(S1) during sensory learning and behavioral adaptation in the mouse. We found that in response

to changing sensory stimulus statistics, mice adopt a task strategy that modifies their detection

behavior in a context dependent manner as to maintain reward expectation. Correspondingly,

neuronal activity in S1 shifts from simply representing stimulus properties to adaptively

representing stimulus context in an experience dependent manner. Our results suggest that

neuronal signals in S1 are part of an adaptive and dynamic framework that facilitates flexible

behavior as an individual gains experience.

INTRODUCTION

Survival in a constantly changing sensory environment requires a high degree of behavioral

flexibility and experience. While much is known about how and where in the brain of human and

non-human animals sensory signals are processed and where decision signals are accumulated

[1–3], far less is known about how behavioral strategies are formed with practice and experience

and whether primary sensory areas are involved in this dynamic process. At the core of any

learning process is the detection of sensory stimuli, which requires efficient neuronal coding of

sensory features in the early sensory pathway. However, studies investigating the role of primary

sensory cortex in visual[4–6], auditory[7–10] and somatosensory behaviors[11,12] are in complete

disagreement about its function. This discrepancy could be a result of the complexity of the

chosen behavioral paradigm or it could be because cortical signals are highly dynamic and

context driven. More recent findings have shown that signals in primary somatosensory cortex

can enhance stimulus selectivity with behavioral training[13], fluctuate according to the behavioral

state[14], or even remap depending on downstream signals[15]. Together, these findings have

opened up many questions and they motivate us to believe that neuronal signals in primary

sensory areas may be highly dynamic, context or experience dependent, and part of an adaptive

framework.

Here, we investigate perceptual capabilities of the mouse vibrissa system during learning and

behavioral adaptation. We hypothesize that signals in primary somatosensory cortex (S1) not only

represent the strength of a sensory input but also play a key role in the transformation of context

dependent behaviors. To test this hypothesis, we designed a series of psychophysical

experiments evaluating behavioral performance and neuronal activity at different training stages.

The training stages include gradual learning of a basic detection task and an advanced stage with

changing sensory contingencies[16]. To repeatedly measure signals of large neuronal pools across

training stages, we performed chronic wide-field imaging of S1 activity with the genetically

encoded voltage indicator (GEVI) 'ArcLight' [17,18] in behaving mice.

We found that in response to changing statistical properties of the sensory stimulus, mice adopt

a strategy that modifies their behavior in a way as to maintain reward in the face of these changes.

Our results further reveal that S1 activity correlates with behavioral changes in a way that depends

upon experience. During learning of the basic task, S1 sensitivity is mostly stimulus driven and

uncorrelated with gradual changes in behavioral performance. However, once an animal reaches

expert level and is trained to adapt to a change in stimulus statistics, neuronal activity dynamically

shifts between changes in S1 sensitivity and decision criterion downstream. The change in S1

sensitivity is more pronounced at a later training stage when the animal has already experienced

the task modulation before. Our findings suggest a translation of these context dependent

changes between different brain structures along the hierarchy, where S1 is not simply

representing sensory stimulus properties, but instead reflecting an adaptive process as part of a

behavioral strategy to engage changing stimulus contingencies in a changing sensory

environment.

MATERIALS AND METHODS

Animals, surgery, and general procedures for behavioral testing.

All experimental and surgical procedures were approved by the Georgia Institute of Technology Institutional Animal Care and Use Committee and were in agreement with guidelines established by the NIH. Subjects were seven male mice (C57BL/6, Jackson Laboratories), aged 4-6 weeks at time of implantation. The basic procedures of virus delivery, head-plate preparation and cortical imaging exactly followed the ones published in a recent paper[17]. In the following text, only procedures pertaining to the specific procedures established here are described in detail.

Virus delivery

At least four weeks prior to experimentation, mice were anesthetized using isoflurane, 3% to 5% in a small induction chamber, and then placed on a heated platform (FHC, Inc.) to maintain body temperature with a stereotaxic nose cone to maintain anesthesia. During the surgery, the anesthesia levels were adjusted to 1–1.5% to achieve ∼1/second breathing rate in mice. For virus delivery, 3 small craniotomies (burr holes of 0.7 mm diameter) were created over the barrel field of the primary somatosensory cortex (S1) according to stereotaxic measurements taken from the bregma ([1 x 3 mm, 3 x 3 mm, 3 x 1 mm] bregma x lateral). One additional craniotomy and Injection was performed over motor cortex (M1, -1 x 1 mm bregma x lateral). The virus was loaded into a neural syringe (Hamilton Neuros Syringe 700/1700). The injection needle was initially lowered to 1000 μm below the pia surface for pre-penetration and then retracted to the target depth of 500 μm, using a 10-μm resolution stereotaxic arm (Kopf, Ltd.). Following a 1-min delay to allow for tissue relaxation, each animal was injected with 2 μl of adeno-associated virus (AAV)1-hsyn1-ArcLight-D-WPRESV40 (UPenn Viral Vector Core, AV-1-36857P) at a flow rate of 0.05-μL∕min (0.5 μl each for four injections). After injection, the needle remained in place for an additional 5 min before slowly being removed from the brain. The craniotomies were left to close naturally. In all cases, the skull was sealed by suturing the skin. Throughout the experiment, sterile techniques

were used to keep the injection area clean and free from infection. Additionally, opioid and non-

steroidal anti-inflammatory analgesic were administered (SR-Buprenorphine 0.8 - 1 mg/kg, SC,

pre-operatively and Ketoprofen 5-10 mg/kg, IP, post-operatively).

Head plate Implantation

After at least four weeks post injection, a metal head-plate was secured to the skull in order to

reduce vibration and allow head-fixation during imaging and behavior experiments. Following

anesthetization and analgesia, a large incision was made over the skull. The connective tissue

and muscles surrounding the skull were removed using a fine scalpel blade (Henry Schein #10).

The custom titanium head-plate formed an open ring (half-moon shape with an inner radius of 5

mm) and was placed on top of the two hemispheres with an extended bar above the cerebellum

(~10 mm, perpendicular to the midline of the skull). The extended bar was designed to attach to

a stainless steel holder, serving the purpose of stable head-fixation. The head-plate was attached

to the bone using a three stage dental acrylic, Metabond (Parkell, Inc.). The Metabond was chilled

using ice, slowly applied to the surface of the skull, and allowed to cure for 5 to 10 min. After

securing the head-plate, the skull was cleaned and covered with a thin layer of transparent

Metabond. During preparation for histological validation, the head-plate could not be separated

from the attached skull and the brain was extracted by removing the lower jaw. The final head-

plate and dental acrylic structure additionally created a well for mineral oil that helped maintain

skull transparency for the upcoming imaging sessions. Mice were allowed to recover for at least

7 days before habituation training. After wound healing, subjects were housed together with a

maximum number of three in one group cage and kept under a 12/12 h inverted light / dark cycle.

Whisker Stimulation.

For whisker stimulation a galvo-motor (galvanometer optical scanner model 6210H, Cambridge

Technology) as described in a previous study[19] was used. The rotating arm of the galvo-motor

contacted a single whisker on the right of the mouse's face at 5 mm (±1 mm tolerance) distance

from the skin, and thus, directly engaged the proximal whisker shaft, largely overriding bioelastic

whisker properties. All the remaining whiskers were trimmed to prevent them from being touched

by the rotating arm. Across mice, different whiskers were chosen (M01: C1, M02: D2, M03: C1,

M04: E2, M05: D1, M06: C1, M07: D1). Voltage commands for the actuator were programmed in

Matlab and Simulink (Ver. 2015b; The MathWorks, Natick, Massachusetts, USA). A stimulus

consisted in a single event, a sinusoidal pulse (half period of a 100 Hz sine wave, starting at one

minimum and ending at the next maximum). The pulse amplitudes used (A = [0, 1, 2, 4, 8, 16]°,

correspond to maximal velocities: V_{max}= [0 314 628 1256 2512 5023°/s) or mean velocities: V

$_{mean}$= [0 204 408 816 1631 3262°/s) and were well within the range reported for frictional slips

observed in natural whisker movement[20,21].

Cortical GEVI Imaging.

ArcLight transfected mice were chronically imaged through the intact skull using a wide-field

fluorescence imaging system to measure cortical spatial activity (MiCAM05-N256 Scimedia, Ltd.).

Figure 1a shows the experimental apparatus and schematically describes the wide-field

fluorescence microscope. During all imaging experiments, mice were awake and head-fixed. The

head-plate was used as a well for mineral oil in order to keep the bone surface wet and maintain

skull transparency. The skull was covered with a silicon elastomer (Kwik-Cast Sealant World

Precision Instruments) between imaging sessions for protection. The barrel cortex was imaged

using a 256 x 256-pixel CMOS camera (Scimedia Model N256CM) at 200 Hz with a pixel size of

69µm x 69µm and an active imaging area of 11.1 x 11.1 mm, given a magnification of 0.63. Note,

this resolution does not consider the scattering of the light in the tissue. During experimental

imaging, the illumination excitation light was left continuously on. The entire cortical area was

illuminated at 465 nm with a 400-mW/cm^2 LED system (Scimedia, Ltd.) to excite the ArcLight

fluorophore. The excitation light was further filtered (cutoff: 472-30-nm bandpass filter, Semrock,

Inc.) and projected onto the cortical surface using a dichroic mirror (cutoff: 495 nm, Semrock,

Inc.). Collected light was filtered with a bandpass emission filter between wavelengths of 520-35

nm (Semrock, Inc.). The imaging system was focused at ~300 μm below the cortical surface to

target cortical layer 2/3. The first imaging session was used for identifying the barrel field in the

awake and naïve animal at least four weeks after ArcLight viral injection. The barrel field was

mapped by imaging the rapid response to a sensory stimulus given to a single whisker (A = 4-16

°, or mean velocities respectively: V = 816 - 3262°/s). We used two criteria to localize and isolate

the barrel filed: standard stereotaxic localization (~3 mm lateral, 0.5 to 1.5 mm caudal from

bregma) and relative evoked spatial and temporal response (visible evoked activity 20 to 25 ms

after stimulation). A single whisker was chosen if it elicited a clear response within the barrel field.

All subsequent imaging experiments were centered on the same exact location and the same

whisker was chosen in repeated sessions for a given animal. Figure 1b shows the characteristic

spread of ArcLight expression in an example coronal brain section. The same hemisphere is

shown *in vivo* in Figure 1c with spontaneous fluorescence activity in S1 at the beginning of a

behavioral session. Spontaneous and sensory evoked activity in S1 was imaged every trial, 2 s

before and 2 s after the punctate whisker. Figure 1d shows frames with typical fluorescence

activity patterns recorded at a framerate of 200 Hz and shown separately for three different

stimulus amplitudes (rows). The calculation of the $\Delta F/F_0$ metric, region of interest (ROI) and data

processing is described under 'Imaging analysis').

Behavioral paradigm and training

During successive days of behavioral testing, water intake was restricted to the experimental

sessions where animals were given the opportunity to earn water to satiety. Testing was paused

and water was available *ad libitum* during 2 days a week. Body weight was monitored daily, and

was typically observed to increase or remain constant during training. In some cases, the body

weight dropped slightly across successive training days due to a higher task difficulty. If the weight

dropped for more than ~5g, supplementary water was delivered outside training sessions to

maintain the animal's weight. 1-2 experiments were usually conducted per day comprising 50-

250 trials. During behavioral testing a constant auditory white background noise (70 dB) was

produced by an arbitrary waveform generator to mask any sound emission of the galvo-motor-

based whisker actuator. All seven mice were trained on a standard Go/No-Go detection task (Fig.

1e) employing a similar protocol as described before [16,22–25]. In this task, the whisker is deflected

at intervals of 4-10 s (flat probability distribution) with a single pulse (detection target). A trial was

categorized as a 'hit' (H) if the animal generated the 'Go' indicator response, a lick at a water

spout within 1000 ms of target onset. If no lick was emitted the trial counted as a 'miss' (M). In

addition, catch trials were included, in which no deflection of the whisker occurred (A = 0°) and a

trial was categorized as a 'correct rejection' (CR) if licking was withheld ('No-Go'). However, a trial

was categorized as a 'false alarm' (FA) if random licks occurred within 1000 ms of catch onset.

Premature licking in a 2 s period before the stimulus was mildly punished by resetting time ('time-

out') and starting a new inter-trial interval of 4-10 s duration, drawn at random from a flat

probability distribution. Note these trial types were excluded from the main data analysis.

The first step of behavioral training was systematic habituation to head-fixation and experimental

chamber lasting for about one week. During the second training phase, a single whisker deflection

with fixed amplitude was presented interspersed by catch trials (P_{stim}=0.8, P_{catch}=0.2). Immediately

following stimulus offset, a droplet of water became available at the waterspout to condition the

animal's lick response thereby shaping the stimulus-reward association. Once subjects showed

stable and immediate consumption behavior (usually within 1-2 sessions), water was only

delivered after an indicator lick of the spout within 1000 ms, turning the task into an operant

conditioning paradigm in which the response is only reinforced by reward if it is correctly emitted

after the stimulus. Subsequent experiments were performed systematically and the behavioral

performance was measured with simultaneous GEVI imaging. The different experiments are

described in detail in the following section.

Basic learning: Learning was studied once an animal had entered the operant phase of training

after the basic habituation procedure. From this point forward, experiments were conducted with

equal conditions across sessions and without manual interference by the experimenter. To assess

differences in learning based on stimulus strength, animals were separated into two groups: group

1 (M01-M03) receiving only one low amplitude stimulus and catch trials (A = [4 0]°) and group 2

(M04-M06) receiving only one high amplitude stimulus and catch trials (A = [16 0]°). Performance

metrics are described under *data analysis and statistics*.

Adaptive behavior: After mice had learned the basic detection task, the psychometric curve was

measured in a subgroup of animals (mouse 4-7) using the method of constant stimuli, which

entails the presentation of repeated stimulus blocks containing multiple stimulus amplitudes. On

a single trial, one out of multiple possible stimulus amplitudes was presented after a variable time

interval (4-10s), each with equal probability (uniform distribution, p = 0.2). A stimulus block

consisted of a trial sequence comprising all stimuli and a catch trial in pseudorandom order (e.g.,

each type once per block). A behavioral session consisted of repeated stimulus blocks until the

animal disengaged from the task, i.e. when it did not generate lick responses for at least an entire

stimulus block. Therefore, the chosen stimulus occurred repetitively but randomly within a

session. An experiment consisted of multiple sessions comprising two conditions, where each

condition is defined by a different stimulus distribution. A condition was always kept constant

within and across multiple behavioral sessions before the task was change. In the 'high range'

condition, four stimulus amplitudes plus catch trial were used (A = [0, 2, 4, 8, 16] °) and presented

in multiple successive sessions. Following this, four new stimulus amplitudes were presented (A

= [0, 1, 2, 4, 8] °) forming the 'low range' condition. Both stimulus distributions shared two of the

three stimulus amplitudes; however, the largest stimulus amplitude of the high range condition

(16°) was not part of the low range, and vice versa, the smallest amplitude of the low range

condition (1°) was not part of the high range. In order to test the reversibility of potential behavioral

and neuronal changes, another high range condition was presented after the animals had

completed the low range condition. This reversal of conditions also served the purpose of testing

an animal's experience level at multiple transitions (first, second etc.). Each condition consisted

of 8-10 sessions performed over approximately 5 days. Each animal performed 1-2 sessions per

day with a minimal break of 3 hours in between.

Data analysis and statistics.

Behavior. The learning curve was measured by calculating a dprime, $'_{behav}$, from the observed

hit rate and false alarm rate of each training session

$$d'_{behav} = Z(hit) - Z(fa)$$

where the function Z(p), p $\in$ [0,1], is the inverse of the cumulative distribution function of the

Gaussian distribution. A criterion of d' = 2.3 (calculated with $p(hit)$ = 0.95 and $p(fa)$ = 0.25) was

used to determine the end of the basic learning period and learning progress was separated into

three equally distributed stages, naive (d' = 0-0.8), intermediate (d' = 0.8-1.5) and expert level (d'

= 1.5-2.3). Psychometric data were assessed as response-probabilities averaged across

sessions within a given stimulus condition. This was done separately for each of the seven

animals. Psychometric curves were fit using Psignifit[26–28]. Briefly, a constrained maximum

likelihood method was used to fit a modified logistic function with 4 parameters: α (the

displacement of the curve), β (related to the inverse of slope of the curve), γ (the lower asymptote

or guess rate), and λ (the higher asymptote or lapse rate) as follows:

$$P(GO|s_i) = \gamma + (1 - \gamma - \lambda) \frac{1}{1 + \exp(-z(s_i))}$$

$$z(s_i) = \frac{s_i - \alpha}{\beta}$$

where s_i is the stimulus on the ith trial. Response thresholds were calculated from the average

psychometric function for a given experimental condition using Psignifit. The term "response

threshold" refers to the inverse of the psychometric function at some particular performance level

with respect to the stimulus dimension. Throughout this study, we use a performance level of 50%

(probability of detection = 0.5). Statistical differences between psychophysical curves were

assessed using bootstrapped estimates of 95% confidence limits for the response thresholds

provided by the Psignifit toolbox.

Reward accumulation. Let the stimulus amplitude delivered on the ith trial be denoted as s_i, the

corresponding reward as r_i, (since r_i is a fixed value, it can just be termed r) and the accumulated

reward for N trials as R_N. Over N trials, the expected accumulated reward is

$$E\{R_N\} = \sum_{i=1}^{N} P(s_i)P(GO|s_i)r_i$$

where $P(s_i)$ comes from the experimentally controlled stimulus distribution, $P(GO|s_i)$ is the

probability of a positive response (or "Go") for the given stimulus amplitude, and $E\{\}$ denotes

statistical expectation.

We considered the null hypothesis of this behavioral paradigm to be that animals do not adapt

their behavior in response to an experimentally forced change in stimulus distribution and thus

operate from the same psychometric curve (represented as dotted curves in Figure 3b). Note, this

corresponds to the same curve but for a different range of stimuli ($P(GO|s_i)$ across different s_i.

For example, in moving from the high range to the low range stimulus condition, this would result

in a decrease in the total accumulated reward for the same number of trials.

As an alternative hypothesis, one possible strategy the animal could take in response to a change

in the stimulus distribution would be to adjust behavior to maintain the same amount of

accumulated reward during a session. For example, in moving from high range stimuli to low

range stimuli, the accumulated reward would be assumed fixed, and we can determine a new set

of probabilities $P(GO|s_i)$ that define an adapted psychometric function. Note that there is not a

unique solution, but one simple possibility is that the original psychometric function maintains the

same asymptotes (γ and λ) and false alarm rate but is compressed, with a decrease in response

threshold and an increase in slope to maintain the same total accumulated reward. We denote

this situation as our hypothetical psychometric function, represented as dashed curves in Figure

3b.

Imaging analysis. All voltage imaging data was analyzed using custom written image-analysis

software (MATLAB 2015a, Mathworks, Inc.). The specific methods of processing the ArcLight raw

fluorescence signal and basic data analysis followed those of a recent study from our laboratory[17].

Briefly, raw images were loaded and converted from the proprietary file format of the imaging

system using custom scripts. Due to the natural decay of the fluorescent signal caused by photo

bleaching, each trial was first normalized to a baseline and reported as a percent change in

fluorescent activity (%$\Delta F/F_0$). The $\Delta F/F_0$ measurement was calculated by subtracting and dividing

each trials fluorescence *F(x, y, t)* by the frame preceding the stimulus delivery:

263
$$\frac{\Delta F}{F_0} = \frac{F - F_0}{F_0}$$

where $F_0(x,y)$ is the frame of stimulus delivery (F_0= F at t=0). Note, an extended analysis was

performed with different normalization methods by subtracting and dividing each trials

fluorescence F by the fluorescence averaged across different time windows before stimulus onset

(t=-50-0ms, -100-0ms, -200-0ms, etc.). Increasing normalization windows slightly altered the

change in fluorescence magnitude and variance of the evoked response; however, varying the

normalization window did not affect the adaptive cortical response reported in this study (Suppl.

Fig. 3).

A single region of interest (ROI) was identified using the largest 10 x 10 pixel (434 x 434 μm) area

response 25 ms following stimulus onset. The average activity within this region was extracted

across all frames to compute the temporal dynamics of the fluorescent signal. Note, due to the

fluorophore[18], positive changes in membrane potential correspond to a decrease in ArcLight

fluorescent activity. In line with our previous study[17] all traces have been inverted to show a

positive increase in fluorescence. Fluorescent voltage traces and behavioral lick responses were

acquired within the same time window and aligned with regard to stimulus onset (Fig. 1f).

Ideal observer analysis. To quantify the fluorescence signal over the course of learning a metric

d'_{neuro} was computed. For a given day or session, single trial distributions of evoked signal

peaks (maximum %$\Delta F/F_0$ within 100 ms post stimulus) are compared to the corresponding noise

distributions when no stimulus was present. d'_{neuro} is then defined as:

282
$$d'_{neuro} = \frac{\mu S - \mu N}{\sqrt{\frac{1}{2}(vS + vN)}}$$

where μS and μN are the mean and vS and vN are the variance of the signal and noise distribution.

d'_{neuro} was then directly compared with the behavioral learning curve derived from d'_{behav}.

The same analysis was repeated for the data acquired with changing stimulus statistics, with the

exception that only trials with an intermediate stimulus amplitude shared between the high and

the low range condition were used. Note, the chosen stimulus of 8-degree whisker angle

represents the midpoint of the high range and the upper limit of the low range condition.

Distributions consisting of all evoked trials termed 'signal' (peak $\Delta F/F_0$ with 8°) and catch trials

termed 'noise' peak ($\Delta F/F_0$ with 0°) were computed separately for the high and low range condition

with one switch from high to low or vice versa. Within a given condition, trials were concatenated

for consecutive sessions and data from all four mice were used to calculate d'_{neuro} and d'_{behav}.

Statistical differences between distributions were assessed using bootstrapped estimates of 95%

confidence limits for the d' metrics. This means resampling the d' from a given session with

replacement 1000 times, taking the average of each resampled dataset, and then taking the

interval that spans the central 95% of this distribution of averages across resampled datasets.

Significance values were further estimated with a non-parametric Wilcoxon rank-sum or Kruskal-

Wallis test. Throughout this manuscript, '*' indicates p < 0.05; '**' indicates p < 0.01; '***' indicates

p < 0.001; and 'n.s.' indicates 'not significant'.

Cascade framework. This analytical framework was created to describe the correspondence

between the sensory input distribution, the neuronal response function derived from the GEVI

signal in S1, and the behavioral readout. The experimentally controlled stimulus amplitude on a

given trial s_i is drawn randomly from the input distribution. The evoked GEVI signal in S1 can then

be expressed as a stimulus response function $G(s_i)$. To establish a link between $G(s_i)$ and the

behavior, a mathematical function $f(.)$ was created that transforms $G(s_i)$ to a probability of a lick

response $P(GO|s_i)$, i.e. the psychometric function, as illustrated in Figure 4a. In other words $f(.)$

represents a matching of the evoked fluorescence to the lick response, given a particular stimulus.

Combining both $G(s_i)$ and $f(.)$ results in a function $f[G(s_i)]$, as an estimate of the psychometric

curve, $\hat{P}(GO|s_i)$, that can then be directly compared to the actual psychometric curve $P(GO|s_i)$.

This approach can be applied across the different stimulus conditions (high versus low), and

comparisons made across the corresponding $G(s_i)$ and $f(.)$ functions.

To quantitatively estimate how much of the behavioral variation can be explained by S1 activity

versus downstream, the following control was performed: $G(s_i)$ is considered to change between

the high and the low range condition, as observed experimentally, from $G_{High}(s_i)$ to $G_{Low}(s_i)$. As

a null test for the transition from the high to low condition, to capture how much of the observed

changes in behavior is explained solely by the changes in $G(s_i)$, $f(.)$ is held constant, operating

from a function $f_{High}(.)$, that only reflects the high range condition. The combination of $G_{Low}(s_i)$

and $f_{High}(.)$ then produces an estimated psychometric function $\hat{P}_{Null}(GO|s_i)$ that is only

influenced by changes in $G(s_i)$, and thus serves as a null test for the prediction based on changes

in neural activity in S1 alone. The fraction explained by $G(s_i)$ and thus S1, can be estimated by

comparing the area under the curve of $\hat{P}_{Null}(GO|s_i)$ with the area under the curve of

$f_{Low}[G_{Low}(s_i)]$. The remaining is that explained downstream.

All curves were fit using the psignifit toolbox[26] and the goodness of fit was assessed by calculating

metrics of deviance (D) as well as the corresponding cumulative probability distribution (CPE). To

rule out the possibility of poor fitting in the cascade framework, we inspected the goodness-of-fit

metric of deviance (D) as well as estimates of where the goodness-of-fit lay in bootstrapped

cumulative probability distributions of this error metric (CPE) using the psignifit toolbox. Due to

the steep increase in $\Delta F/F_0$ at lower stimulus amplitudes we find that a Weibull function provides

the best fit for $G(s_i)$ (example $G(s_i)$ in Fig. 4b, D_{high} = 0.12, CPE_{high} = 0.06; D_{low} = 1.03, CPE_{low} =

0.46). Both, the linker function f(.) and the psychometric function are best fit by a logistic function

due to the sigmoid configuration of the data (example f(.) in Fig. 4c, D_{high} = 0.6, CPE_{high} = 0.54;

D_{low} = 1.57, CPE_{low} = 0.67).

RESULTS

The current study investigates learning and experience dependent adaptation in the mouse

vibrissa system. Our main goal here is to establish a relationship between controlled whisker

inputs, S1 activity, and behavior output, firstly, during basic detection learning and secondly,

during flexible adaptation to changing sensory contingencies. To repeatedly measure signals of

large neuronal pools at different training stages, we performed chronic wide-field imaging of S1

activity with the genetically encoded voltage indicator (GEVI) 'ArcLight' in behaving mice.

Figure 1 outlines the experimental design and summarizes the basic neuronal and behavioral

metrics of this study. Seven mice were transfected with the GEVI ArcLight before being imaged

through the skull using a wide-field fluorescence microscope (Fig. 1a). We have previously

described this imaging technique in detail[17]. Figure 1b shows the characteristic spread of ArcLight

expression in an example coronal brain section after a subject had undergone all tests and its

brain was extracted for histological validation. The same hemisphere is shown *in vivo* in Figure

1C with spontaneous fluorescence activity in S1 at the beginning of a behavioral session. Figure

1D shows sequential frames of typical fluorescence activity patterns of a trained mouse recorded

at a framerate of 200 Hz and separated into catch trials (top row) and stimulation trials (bottom

row) in response to a single whisker deflection (see methods for details about GEVI imaging).

All transfected mice were first trained on a tactile Go/No-Go detection task as described

previously[16,22,23,29,30]. Briefly, this task requires animals to detect pulse-shaped deflections of a

single whisker and report the decision by either generating a lick on a waterspout ('Go') or by

withholding licking ('No-Go') if no stimulus is present (Fig. 1e). The trained mouse shows a

stereotypical lick response pattern and a clear cortical fluorescence response to stimulation trials

(Fig. 1f). Note, the temporal resolution of ArcLight allows the disentangling of the sensory

response from potential motor related signals caused by licking (at 200-300 ms post stimulus).

To investigate behavior and neuronal signals at different training stages, we split the experiments

into two main parts (Fig. 1g). The first part was designed to evaluate learning from one session

to another, covering both behavior and neuronal S1 data in mice acquiring the basic principles of

the Go/No-Go task, which we refer to here as "basic learning". The second part describes S1

activity when mice are challenged with a change in sensory statistics. We refer to this part as

"adaptive behavior". In contrast to the first part, these experiments employed multiple whisker

deflection amplitudes presented randomly and we systematically manipulated the statistical

distribution of the stimulus.

S1 responses during basic learning

Naive subjects received single whisker stimulation at intervals of 4-10 s with a single pulse or

catch trial. Learning was measured by calculating the hit $p(hit)$ and false alarm rate $p(fa)$ of

successive training sessions with one session performed each day. A criterion of $p(hit)$ = 0.95

and $p(fa)$ = 0.25 was used to determine successful acquisition of the task. Figure 2a shows hit

and false alarm rates for three mice trained on a weak stimulus (A = 4°). The choice of a weak

stimulus is anticipated to make the task difficult requiring more training and facilitating sufficient

data collection especially during early learning. Indeed, with this training protocol, subjects

required extended exposure to the task with more than 30 sessions to achieve successful

performance. Figure 2b shows frames of cortical fluorescence activity from an example mouse

over the course of learning. Each square represents a top view of the same cortical hemisphere

on selected training days with the average fluorescent activity ($\Delta F/F_0$ in %) at the peak of the

sensory evoked signal (A = 4°, top row) or during catch trials (A = 0°, bottom row). Qualitative

inspection of the fluorescence signal regarding its magnitude and spread in cortical space reveals

a high level of variability from one day to another, but a systematic change with learning progress

(e.g. signal in- or decrease) cannot be identified. We further quantitatively evaluated the

fluorescence signal over the course of learning by computing a neuromeric sensitivity measure

d'_{neuro}. Single trial distributions of evoked signal peaks are compared to the corresponding noise

distributions when no stimulus is present. The d'_{neuro} for a given day or session is then calculated

by subtracting the means of the distributions and dividing it by the variance. To compare this

metric to the behavior, a d'_{behav} was calculated from the observed behavioral hit and false alarm

rate of each training session (see methods). Data from all three mice were used for this analysis.

Figure 2c shows the d'_{neuro} (orange) along with the behavioral learning curve d'_{behav} (black), across

all sessions and mice. The dashed lines separate performance into 'naive' (d'<0.8), and 'acquired'

(d'>1.5). In contrast to the behavior, the average neuromeric sensitivity remains at a relatively

constant level, representing a stable signal to noise relationship independent of the continuing

learning progress. By separating the data into naive and acquired (Fig. 2c, Inset) across the group

of mice, this finding can further be confirmed statistically (d'_{neuro} naive vs. acquired, p>0.1,

Wilcoxon rank sum test). Note, a second group of animals was trained on a much stronger

stimulus for comparison (A = 16°). Those mice achieved much higher hit rates at the beginning

of training and reached successful task acquisition in half the time (Suppl. Fig. 1a); however, the

neuromeric sensitivity was also orthogonal to learning progress (Suppl. Fig. 1b-c). We conclude

that neuronal sensitivity in S1 measured from a large population of neurons does not change

during basic learning.

Adaptive behavior and changes in S1 in the highly trained animal

To investigate behavioral and neuronal dynamics with regard to changing context, we performed

experiments in which we systematically manipulated stimulus statistics. First, we present an

analysis of data from highly trained animals only, which we refer to here as "experienced". Those

animals had successfully acquired the basic training and first time adaptation to changing stimulus

statistics. The psychophysical techniques were based on a behavioral paradigm we recently

developed in the rat[16]. Figure 3a shows the manipulation of the stimulus distribution range; that

is, the upper and lower limits of the statistical distribution of whisker deflection amplitudes

presented in a psychophysical test. The first distribution consists of four different stimulus

amplitudes and a catch trial (A = [0, 2, 4, 8, 16] °, magenta) which we refer to as the 'high range'

condition. The second distribution consists of four new stimulus amplitudes and a catch trial (A =

[0, 1, 2, 4, 8] °, green), which we refer to as the 'low range' condition. Each stimulus or catch trial

was presented with equal probability (uniform distribution). Importantly, the experimental design

involves amplitudes common to both high-range and low-range conditions. Hence, three of the

four stimulus amplitudes are shared between conditions. However, the largest amplitude of the

high range (16°) is not part of the low range, and vice versa, the smallest amplitude of the low

range (1°) is not part of the high range. Figure 3b depicts typical psychometric curves from an

example mouse, performing the task first under the high range (magenta) and under the low range

condition (green). There is an obvious and consistent shift of the psychometric curve in response

to the changing stimulus statistics, which we refer to here as "adaptive behavior".

In response to a shift in stimulus distribution, we consider two extreme hypotheses as established

previously by a simple reward expectation model [16]. The null hypothesis (H0) asserts that the

animal does not adjust its behavior, and thus operates from the same psychometric function

(black dotted curve on top of magenta curve). In moving from the high range to the low range

condition, this would result in a decreased reward rate for the same number of trials. The

alternative hypothesis (H1) predicts that the animal adapts its behavior to maintain accumulated

reward in the face of a changing stimulus distribution. The black dashed curve denotes the

hypothetical psychometric function with the same lapse and guess rate (see methods) as the

original curve from the mouse, but allowing it to shift to the left such that the expected reward per

trial remains constant. The experimentally measured psychometric function in the low range

condition (green) comes quite close to the hypothetical performance level based on the

assumption of maintained reward expectation. The observed shift results in a significant decrease

in response threshold for all mice (T_{high} = 4.83 ± 0.33, T_{low} = 2.93 ± 0.34, Mean and SD, $p<0.05$,

Kruskal-Wallis test). Figure 3c depicts the actual trial-by-trial reward accumulation by the example

mouse. Overlaid are results for n=11 sessions with the high range distribution (magenta) and

n=10 sessions with the low range distribution (green). The slope of reward accumulation in the

low range condition nearly matches that of the high range condition, and the slope for the low

case (green) is close to the prediction from the maintenance of accumulated reward hypothesis,

H1, while being clearly separable from the slope representing the null hypothesis (dotted line).

The total number of rewards acquired on average per session and across all mice further confirms

this (total # high range = 44.5 ± 8.6, total # low range = 42.0 ± 10.6, Mean and SD, Fig. 3c inset),

whereas there was no evidence for an alternative strategy to maintain the total number of rewards

by working substantially more trials.

This finding shows that detection behavior is highly flexible in the face of a changing stimulus

distribution and it further supports the concept of statistical integration and probabilistic

computations in the mouse brain. The stability of the stimulus representation during basic learning

might suggest that this occurs downstream of S1. Figure 3d shows frames of evoked S1 activity

of the same mouse from Figure 3b and c engaged in two example sessions, one presenting the

high range (top) and the other presenting the low range distribution (bottom). The frames are

aligned for stimuli common to both datasets. Qualitatively, both examples show a higher change

and spatial spread in fluorescence with increasing stimulus amplitude. Note, there is a fixed linear

relationship between the percent change in fluorescence magnitude (% DF/F_0) and the activated

cortical area, as both scale with stimulus strength in a highly correlated fashion (Suppl. Fig. 2).

For simplicity, we use percent change in fluorescence magnitude as a metric for cortical

activation.

Interestingly, the evoked activity for the same stimulus is clearly higher in the low range compared

to the high range condition. To further investigate this modulated activity, we focus our analysis

first on trials with an intermediate stimulus shared between both datasets (outlined box). The

chosen deflection angle of 8 degrees represents the midpoint of the high range distribution and

the upper limit of the low range distribution. According to the psychophysical results, this particular

stimulus is considered a 'supra-threshold' stimulus and therefore within a detectable range. Figure

3e depicts the temporal fluorescent signal averaged across sessions from all mice at the

experienced level. Histograms of behavioral lick responses are shown on top. As we suspected

from qualitative assessment of the cortical response, the evoked response to the same 8-degree

deflection is higher on average if presented within the low range compared to the high range

distribution.

To compare the modulated S1 activity with the adapted behavior, we used classical signal

detection theory[31,32]. Figure 3f outlines the approach schematically. Multiple scenarios are

considered to trigger a change within sensory and higher order processing stages; behavioral

adaptation can either be induced by changes in sensitivity (d') intrinsic to S1, changes in decision

criterion (also called bias or decision boundary) by a downstream observer, or both. Sensitivity in

S1 improves through reduction in the overlap between sensory signal and noise distributions (left

panel). In addition, the downstream observer may value hits and false alarms differently by

altering the decision criterion (right panel). To test these predictions, distributions consisting of

many evoked trials termed 'signal' ($\Delta F/F_0$ with 8°) and catch trials termed 'noise' ($\Delta F/F_0$ with 0°)

were computed separately for the high and low range condition (Fig. 3g). Trials were

concatenated for consecutive sessions with one switch from high to low or vice versa and data

from all experienced mice were used for this analysis. The noise distribution (grey) is comparable

for the high and the low range condition with its means being identical (ΔF_{high}=0%, ΔF_{low}=0%).

However, the signal distribution for the low range (green) is shifted towards higher fluorescence

changes as compared to the signal distribution of the high range (magenta) with a clear difference

in its mean (ΔF_{high}=0.35%, ΔF_{low}=0.52%). A significant difference in neuronal sensitivity was

confirmed by calculating the neuromeric d' as introduced earlier in this study (d'$_{high}$=0.69,

d'$_{low}$=1.04, p<0.01, Kruskal-Wallis test). Note that while the neuronal d' was altered slightly by

different normalization methods of the $\Delta F/F_0$ metric, an extended analysis revealed that the

difference in d' across conditions persisted (Suppl. Fig. 3a-c). In addition, we created receiver

operating characteristic (ROC) curves by varying a criterion threshold across the $\Delta F/F_0$ signal and

noise distributions and plotting the hit rate (signal detected) against the false alarm rate (incorrect

guess) (Suppl. Fig. 3d). The ROC curve for the low range condition was higher than that for the

high range condition, quantified by a larger area under the low range ROC curve (AUC_{low} = 0.77)

than for the ROC curve for the high range condition (AUC_{high} = 0.69), thereby confirming a change

in S1 sensitivity. This approach further allowed us to infer changes in criterion by comparing the

hit rate in ROC space (neurometric) with the average hit rate measured from the behavior

(psychometric). The criterion (expressed as % $\Delta F/F_0$,) shows a slight, yet non-significant decrease

when operating from the low range compared to the high range condition (C_{high} = 0.17, C_{low} = 0.08

p>0.05, Kruskal-Wallis test, Fig. 3g blue). Note, the S1 sensitivity and downstream criterion

change in opposite direction, e.g. an increase in hit rate can be caused by an increase in S1

sensitivity and/or a decrease in criterion. We conclude that, in experienced animals, within the

signal detection framework, there is evidence of adaptive sensitivity in S1 yet comparatively

smaller adaptive changes in criterion by a downstream observer.

Changes across training stages

The data described thus far was derived from highly trained ("experienced") subjects in response

to a stimulus that is close to the perceptual threshold. For this case, the signal detection approach

hints at the fact that while changes in S1 dominate, both S1 sensitivity and downstream criterion

may be altered in a complementary or constructive way. However, it is unclear whether such a

co-modulation is dependent on the individual's level of experience. To answer this question, we

created an analytic, cascade framework that enables us to evaluate the relative contributions from

activity in S1 and downstream of S1 in predicting the observed behavior (see methods for details).

Briefly, the framework establishes a link between the sensory input distribution s_i, the S1 response

function $G(s_i)$, and the psychometric curve $P(GO|s_i)$, as shown in Figure 4a. The function $f(.)$

represents a downstream process, matching the evoked fluorescence to the lick response, given

a particular stimulus, and is referred to as the "linker" function, as it links S1 activity to behavior.

This framework can then be used to describe the changes underlying the adaptive behavior.

Figure 4b shows an example function $G(s_i)$ of an experienced mouse challenged with a change

in stimulus statistics. The function $G(s_i)$ is represented by a curve fitted to the data, and separately

shown for each condition, $G_{High}(s_i)$ (magenta) and $G_{Low}(s_i)$ (green). As we expected based on

the qualitative observations of the GEVI response and the ROC analysis, a difference between

$G_{High}(s_i)$ and $G_{Low}(s_i)$ is clearly visible. This is particularly obvious for larger stimulus strengths,

but it is also true for smaller stimulus strengths that are less obvious on the logarithmic plot. Figure

4c shows the linker function $f(.)$ computed separately for the high and low range condition,

resulting in $f_{High}(.)$ (magenta) and $f_{Low}(.)$ (green). Another critical observation is that $f_{High}(.)$

and $f_{Low}(.)$ are different, confirming our previous finding that a change in S1 sensitivity cannot

fully explain the observed differences in behavior. Figure 4d shows the combined functions

$f_{High}[G_{High}(s_i)]$ and $f_{Low}[G_{Low}(s_i)]$ (dotted curves, magenta and green, respectively) which serve

as predictors of the psychometric curve, along with the original psychometric curves $P_{High}(GO|s_i)$

and $P_{Low}(GO|s_i)$ (solid curves, magenta and green, respectively). The predictions and actual data

match well, by construction, with minor differences due to the fitting accuracy when generating

both, $G(s_i)$ and $f(.)$.

This framework enables us now to assess the relative roles of the two stages of the model in the

context of the high and low range conditions imposed during the behavior. Specifically, as a next

step we sought to quantitatively estimate how much of the behavioral adaptation can be explained

by S1 activity versus other processes downstream. To capture how much of the observed

changes in behavior is explained solely by the changes in $G(s_i)$ in the transition from the high to

the low condition, $f(.)$ is held constant, operating from a function $f_{High}(.)$, that only reflects the

high range condition. The combination of $G_{Low}(s_i)$ and $f_{High}(.)$ then produces an estimated

psychometric function $\hat{P}_{Null}(GO|s_i)$ that is only influenced by changes in $G(s_i)$, and thus serves

as a null test for the prediction based on changes in neural activity in S1 alone, as shown in Figure

4d (black dashed curve). The fraction explained by $G(s_i)$, can be estimated by comparing the

area under the curve of $\hat{P}_{Null}(GO|s_i)$ with the area under the curve of $f_{Low}[G_{Low}(s_i)]$. In this

example of an experienced mouse, changes of the S1 neuronal response function $G(s_i)$ explain

67% of the behavioral adaptation, whereas the remaining 33% would have to be explained

downstream, by changes in $f(.)$.

To investigate the effect of experience, we applied the cascade model separately for each mouse

and at different training stages. So far, we presented data from highly trained ("experienced")

animals, but now we are including data from the same animals at the first exposure to changing

stimulus statistics, which we refer to here as "novel". Figure 5a and b shows in the face of

changing stimulus statistics, all animals adapted their behavior equally well across training stages

as revealed by their similar performance metrics (threshold, Figure 5a) and their adaptive change

in psychometric thresholds for the high range and the low range condition (Figure 5b, Novel [T_{high}

= 4.78 ± 1, T_{low} = 2.93 ± 0.34], Experienced [T_{high} = 4.83 ± 0.33, T_{low} = 2.93 ± 0.34], Mean and SD,

p<0.05, Kruskal-Wallis test). However, by assessing the S1 contribution at different training

stages, an interesting pattern emerged. Figure 5c shows the relative explanatory power of S1

versus downstream, quantified for all mice undergoing repeated task changes. At the first

transition only two out of four individuals show a small change in $G(s_i)$ (M04 = 21%, M05 = 0%,

M06= 16%, M07 = 0%), even though all animals show behavioral flexibility. This indicates that

the associated neuronal processes that are predictive of the behavior could primarily be

performed outside S1. However, at the experienced level (bottom), all four mice show a

substantial change in S1 activity or $G(s_i)$, each explaining a significant fraction of the behavior

(M04 = 67%, M05 = 67%, M06= 45%, M07 = 82%, p < 0.05, Wilcoxon rank-sum test, Fig. 5d). To

rule out that this effect is due to the direction of task manipulation, we performed a control

experiment with multiple changes occurring with alternating conditions (high-low-high-low),

showing that S1 sensitivity closely followed the direction of changes in a way that increasingly

explained the behavioral adaptation (Suppl. Fig. 4). Again, the psychometric threshold itself is

consistently changing once contextual changes are introduced. In other words, animals performed

equally well when challenged with a change whether they had experienced it before or not. This

suggests that the first time behavioral adaptation is required, areas outside of S1 could be

primarily involved in driving this kind of behavior. However, as animals gain experience, adaptive

responses seem to emerge in S1. We conclude that neuronal computations associated with

behavioral flexibility can be found in S1 at an advanced training stage and propose that such

computations might shift across the cortical hierarchy in an experience dependent manner.

DISCUSSION

In this study, we have investigated learning and experience dependent behavior in the mouse

somatosensory system. Our findings provide evidence that activity in primary somatosensory

cortex can be highly dynamic in support of flexible sensory processing and experience dependent

behavioral adaptation. We present the following novel aspects. First, S1 population activity is

stable and mostly stimulus driven during early learning despite changes in behavioral

performance. Second, detection behavior can be modified in a way as to maintain reward in the

face of changing statistical properties of the stimulus. Third, S1 activity is highly dynamic in the

face of a changing sensory environment predicting behavioral adaptation as individuals gain

experience.

Learning occurs when an individual forms an association based on a new stimulus or context.

This process provides obvious benefits such as flexible hunting, optimal foraging, and social

communication, especially in environments that tend to change frequently and unpredictably.

There is no doubt that associative learning can occur in animals without cortex, including all

classes of vertebrates[33] and a large number of invertebrate species[34]. Several studies

elucidating the role of S1 with classical or operant conditioning have found that lesions of S1 do

not affect learning[12,35]. Hong and colleagues showed that mice can learn to detect objects even

after chronic inactivation of primary somatosensory cortex. This finding seems in direct contrast

to the large body of work that argues the contrary[36–38]. However, this disagreement can

presumably be resolved by considering the possibility that neuronal signals in these areas are

actually highly dynamic, context dependent, and part of a flexible or adaptive framework.

When probing S1 wide field activity during learning of the basic detection task we found that

neuronal sensitivity is mostly stimulus driven and does not change during basic learning. This

finding suggests that S1 provides a stable sensory representation over time as an individual

learns, but it does not correlate with the increase in performance or the formation of a basic

association.

However, we extended our training program and challenged subjects with a change in stimulus

statistics once they had learned the basics of the task. Our results show that mice can develop a

flexible behavioral strategy that is closely modulated by the change in sensory statistics allowing

them to maintain a constant payoff. When probing S1 wide field activity during this behavioral

adaptation, we find that neuronal sensitivity changes in an experience dependent manner; the

change in S1 sensitivity emerges at a later training stage when subjects had already experienced

the task modulation before.

The general behavioral question posed here relates to how the animal responds to changes in

the sensory environment. The behavioral paradigm was designed as a highly simplified but

carefully controlled manipulation of the statistical distribution of the magnitude of whisker

deflections experienced by the animal. Aside from matching amplitudes and velocities of the

whisker movement that have been described in a range of studies[20,21,39–41], the current study does

not attempt to place this in the context of the natural sensory environment for the animal, and the

passive stimulus paradigm does not speak to active sensing. Importantly, due to practical

considerations, the primary behavioral experiments in this study are admittedly limited to a fairly

narrow view of the adaptive behavior of the animal, with blocks of behavioral sessions with a

single controlled switch in stimulus statistics, as opposed to presumably a more complex evolution

of stimulus statistics in the natural environment. However, we performed a control with multiple

changes occurring with alternating conditions, showing that S1 sensitivity closely followed the

direction of changes in a way that increasingly explained the behavioral adaptation (Suppl. Fig.

4), suggesting that the findings here likely generalize to more complex scenarios. Furthermore,

our previous study employing a similar psychophysical approach in the rat shows that behavioral

adaptation takes place independent of the direction of task manipulation, and that the adaptive

behavior generalizes to changes in other aspects of the statistics of the sensory stimulus[16].   How

the behavior, and corresponding sensory representations change in more complex, naturalistic

settings, and at a finer time resolution, are important, and need further investigation in future

studies.

Nevertheless, our finding of context dependent adaptive responses in S1 is surprising as it

suggests that reward based choice signals might shift across the cortical network and can appear

and influence sensory representation in S1 once an individual has successfully adapted its

behavioral strategy. In this context it is important to note that the behavioral adaptation itself is

similar at different experience levels, showing consistent changes in performance already before

the adaptive response even appears in S1 (novel performer). This is important, as the change in

S1 we observe through GEVI imaging averaged across trials is therefore not simply reflecting a

difference in performance (e.g. ratio of hits to misses) across the levels of experience, but instead

a true change in the S1 response with experience in the context of this task.

So, what are the downstream targets that drive early behavior and ultimately change the stimulus

representation of primary sensory areas? A recent study probing flexible decision making in the

somatosensory pathway of mice found that orbitofrontal cortex dynamically interacts with sensory

cortex triggering plasticity based on value signals[15]. However, other important reward based

choice signals have been reported to influence neuronal signaling throughout cortex[42].

Furthermore, studies of visual attention in primates have distinguished changes in neuronal

sensitivity from an observer's response criterion in extra striate cortex, superior colliculus, as well

as lateral prefrontal cortex[43–45]. Hence, there is a large number of downstream candidates

potentially modulating the sensory signal stream. Admittedly, in the current study we did not

record in any other area outside S1, but the effects described by those other studies collectively

point to the interesting idea that the neuronal signal transfer identified by our study could be a

common principle across neocortex with "earlier" stages emerging with experience. Moreover, we

propose that this shift in cognitive signaling could even affect subcortical structures such as

primary thalamus as we have shown recently in highly trained animals[25].

Arguably, our current study was not specifically designed to identify attention as a driver for

adaptive behavior, even though the measured changes in performance are indicative of a

somewhat conscious change in the subjects' behavioral strategy. Other effects related to the

subjects' thirst, satiation, or arousal level within this behavioral framework can further be

excluded, as we have shown previously[16]. However, although speculative, the experience

dependent change in S1 sensitivity and the corresponding change in downstream criterion may

be signatures of a higher order cognitive process that could be described as an "attentional

spotlight" or gain control moving down the cortical hierarchy. Such a brain-wide dynamic process

would facilitate selective and efficient cognitive processing as an individual is adapting to changes

in the environment. This transfer could explain some of the disparities in the literature and

therefore needs to be further investigated in future studies.

References

1. Heekeren, H. R., Marrett, S. & Ungerleider, L. G. The neural systems that mediate human perceptual decision making. *Nature Reviews Neuroscience* **9,** 467–479 (2008).

2. Shadlen, M. N., Britten, K. H., Newsome, W. T. & Movshon, J. A. A computational analysis of the relationship between neuronal and behavioral responses to visual motion. *J. Neurosci.* **16,** 1486–1510 (1996).

3. De Lafuente, V. & Romo, R. Neuronal correlates of subjective sensory experience. *Nat. Neurosci.* **8,** 1698–1703 (2005).

4. Glickfeld, L. L., Histed, M. H. & Maunsell, J. H. R. Mouse primary visual cortex is used to detect both orientation and contrast changes. *J. Neurosci.* **33,** 19416–19422 (2013).

5. Petruno, S. K., Clark, R. E. & Reinagel, P. Evidence That Primary Visual Cortex Is Required for Image, Orientation, and Motion Discrimination by Rats. *PLoS One* **8,** (2013).

6. Lashley, K. S. The mechanism of vision IV. The cerebral areas necessary for pattern vision in the rat. *J. Comp. Neurol.* **53,** 419–478 (1931).

7. Kato, H. K., Gillet, S. N. & Isaacson, J. S. Flexible Sensory Representations in Auditory Cortex Driven by Behavioral Relevance. *Neuron* **88,** 1027–1039 (2015).

8. Kelly, J. B. & Glazier, S. J. Auditory cortex lesions and discrimination of spatial location by the rat. *Brain Res.* **145,** 315–321 (1978).

9. Jaramillo, S., Borges, K. & Zador, A. M. Auditory Thalamus and Auditory Cortex Are Equally Modulated by Context during Flexible Categorization of Sounds. *J. Neurosci.* **34,** 5291–5301 (2014).

10. Talwar, S. K., Musial, P. G. & Gerstein, G. L. Role of mammalian auditory cortex in the perception of elementary sound properties. *J. Neurophysiol.* **85,** 2350–2358 (2001).

11. Stüttgen, M. C. & Schwarz, C. Barrel cortex: What is it good for? *Neuroscience* **368,** 3–16 (2018).

12. Hong, Y. K., Lacefield, C. O., Rodgers, C. C. & Bruno, R. M. Sensation, movement and learning in the absence of barrel cortex. *Nature* **561,** 542–546 (2018).

13. Kim, J., Erskine, A., Cheung, J. A. & Hires, S. A. Behavioral and Neural Bases of Tactile Shape Discrimination Learning in Head-Fixed Mice. *Neuron* (2020). doi:10.1016/j.neuron.2020.09.012

14. Lee, C. C. Y., Kheradpezhouh, E., Diamond, M. E. & Arabzadeh, E. State-Dependent Changes in Perception and Coding in the Mouse Somatosensory Cortex. *Cell Rep.* **32,** (2020).

15. Banerjee, A. *et al.* Value-guided remapping of sensory cortex by lateral orbitofrontal cortex. *Nature* **585,** 245–250 (2020).

16. Waiblinger, C., Wu, C. M., Bolus, M. F., Borden, P. Y. & Stanley, G. B. Stimulus context and reward contingency induce behavioral adaptation in a rodent tactile detection task. *J.*

697   *Neurosci.* **39,** 1088–1099 (2019).

698   17.   Borden, P. Y. *et al.* Genetically expressed voltage sensor ArcLight for imaging large scale
699   cortical activity in the anesthetized and awake mouse (erratum). *Neurophotonics* **4,**
700   39801 (2017).

701   18.   Jin, L. *et al.* Single Action Potentials and Subthreshold Electrical Events Imaged in
702   Neurons with a Fluorescent Protein Voltage Probe. *Neuron* **75,** 779–785 (2012).

703   19.   Chagas, A. M. *et al.* Functional analysis of ultra high information rates conveyed by rat
704   vibrissal primary afferents. *Front. Neural Circuits* **7,** 190 (2013).

705   20.   Ritt, J. T., Andermann, M. L. & Moore, C. I. Embodied Information Processing: Vibrissa
706   Mechanics and Texture Features Shape Micromotions in Actively Sensing Rats. *Neuron*
707   **57,** 599–613 (2008).

708   21.   Wolfe, J. *et al.* Texture Coding in the Rat Whisker System: Slip-Stick Versus Differential
709   Resonance. *PLoS Biol.* **6,** e215 (2008).

710   22.   Stüttgen, M. C., Rüter, J. & Schwarz, C. Two psychophysical channels of whisker
711   deflection in rats align with two neuronal classes of primary afferents. *J. Neurosci.* **26,**
712   7933–41 (2006).

713   23.   Ollerenshaw, D. R. *et al.* Detection of tactile inputs in the rat vibrissa pathway. *J.*
714   *Neurophysiol.* **108,** 479–490 (2012).

715   24.   Ollerenshaw, D. R., Zheng, H. J. V, Millard, D. C., Wang, Q. & Stanley, G. B. The
716   Adaptive Trade-Off between Detection and Discrimination in Cortical Representations
717   and Behavior. *Neuron* **81,** 1152–1164 (2014).

718   25.   Waiblinger, C., Whitmire, C. J., Sederberg, A., Stanley, G. B. & Schwarz, C. Primary
719   Tactile Thalamus Spiking Reflects Cognitive Signals. *J. Neurosci.* **38,** 4870–4885 (2018).

720   26.   F.A., W. & N.J., H. The psychometric function: I. Fitting, sampling, and goodness of fit.
721   *Percept. Psychophys.* **63,** 1293–1313 (2001).

722   27.   Wichmann, F. A. & Hill, N. J. The psychometric function: II. Bootstrap-based confidence
723   intervals and sampling. *Percept. Psychophys.* **63,** 1314–1329 (2001).

724   28.   Frund, I., Haenel, N. V & Wichmann, F. A. Inference for psychometric functions in the
725   presence of nonstationary behavior. *J. Vis.* **11,** 16 (2011).

726   29.   Cook, E. P. & Maunsell, J. H. R. Dynamics of neuronal responses in macaque MT and
727   VIP during motion detection. *Nat. Neurosci.* **5,** 985–994 (2002).

728   30.   Waiblinger, C., Whitmire, C. J., Sederberg, A., Stanley, G. B. & Schwarz, C. Primary
729   tactile thalamus spiking reflects cognitive signals. *J. Neurosci.* **38,** 4870–4885 (2018).

730   31.   Green, D. M. & Swets, J. A. Signal detection theory and psychophysics. *Society* **1,** 521
731   (1966).

732   32.   Luo, T. Z. & Maunsell, J. H. R. Attentional Changes in Either Criterion or Sensitivity Are
733   Associated with Robust Modulations in Lateral Prefrontal Cortex. *Neuron* **97,** 1382–

1393.e7 (2018).

33.   Jerison, H. J. & Macphail, E. M. Brain and Intelligence in Vertebrates. *Am. J. Psychol.* **97,**
138 (1984).

34.   Giurfa, M. Learning and cognition in insects. *Wiley Interdisciplinary Reviews: Cognitive*
Science **6,** 383–395 (2015).

35.   Hutson, K. A. & Masterton, R. B. The sensory contribution of a single vibrissa's cortical
barrel. *J. Neurophysiol.* **56,** 1196–1223 (1986).

36.   Galvez, R., Weible, A. P. & Disterhoft, J. F. Cortical barrel lesions impair whisker-CS
trace eyeblink conditioning. *Learn. Mem.* **14,** 94–100 (2007).

37.   Miyashita, T. & Feldman, D. E. Behavioral detection of passive whisker stimuli requires
somatosensory cortex. *Cereb. Cortex* **23,** 1655–1662 (2013).

38.   Guo, Z. V *et al.* Flow of cortical activity underlying a tactile decision in mice. *Neuron* **81,**
179–194 (2014).

39.   Waiblinger, C., Brugger, D., Whitmire, C. J., Stanley, G. B. & Schwarz, C. Support for the
slip hypothesis from whisker-related tactile perception of rats in a noisy environment.
Front. Integr. Neurosci. **9,** (2015).

40.   Hartmann, M. Vibrissa mechanical properties. *Scholarpedia* **10,** 6636 (2015).

41.   O'Connor, D. H. *et al.* Vibrissa-based object localization in head-fixed mice. *J. Neurosci.*
30, 1947–1967 (2010).

42.   Gold, J. I. & Shadlen, M. N. The neural basis of decision making. *Annual Review of*
Neuroscience **30,** 535–574 (2007).

43.   Luo, T. Z. & Maunsell, J. H. R. Attentional Changes in Either Criterion or Sensitivity Are
Associated with Robust Modulations in Lateral Prefrontal Cortex. *Neuron* **97,** 1382–
1393.e7 (2018).

44.   Crapse, T. B., Lau, H. & Basso, M. A. A Role for the Superior Colliculus in Decision
Criteria. *Neuron* **97,** 181–194.e6 (2018).

45.   Luo, T. Z. & Maunsell, J. H. R. Neuronal Modulations in Visual Cortex Are Associated
with Only One of Multiple Components of Attention. *Neuron* **86,** 1182–1188 (2015).

Figure 1. Imaging and behavior methods. a, Top: Schematic of the imaging system. The GEVI 'ArcLight' is expressed in superficial layers of S1. Bottom: Schematic of the behavior setup. **b,** Confocal image of a coronal mouse brain section (100 µm, right hemisphere) showing the characteristic spread of ArcLight in S1. Blue: DAPI staining, Green: Arclight fluorescence. **c,** Top view of the same hemisphere in vivo showing ArcLight fluorescence in S1. **d,** Images of the same hemisphere in a trained mouse with ArcLight fluorescence color coded. Punctate stimuli (16 degree whisker angle) or catch trials (0 degree) were presented. Frames were captured at 200 Hz and depicted from stimulus onset onward. Each frame is normalized to the frame at stimulus delivery ($\Delta F/F_0$ = F-F_0/F_0). Shown are fluorescence responses averaged over an example imaging session (n = 24 trials per condition). A region of interest (red box, 434 x 434 µm) is located with its center at the peak fluorescence to extract and average voltage traces for further analysis. Scale bar = 1 mm. **e,** Go/No-Go detection task. A punctate stimulus (10 ms) has to be detected by the mouse with an indicator lick to receive reward. Reward is only delivered in hit trials. Impulsive licks in a 2 s period before trial onset are mildly punished by a time-out triggering a new inter-trial-interval (4 –10 s, gray arrow). **f,** Example traces of Go and No-Go trials (n = 284 each). Top: Readouts from the stimulator (Stim = 16 degree whisker angle or catch trial = 0 degree). Middle: Lick response histograms from a trained mouse. Bottom: Average voltage response from the region of interest of the same mouse. **g,** Different task versions under investigation. 1. Basic learning: Learning of the Go/No-Go task. 2. Adaptive behavior: Once animals have learned the basic task, they are challenged with multiple stimulus amplitudes and changes in the statistics of the stimulus distribution.

Figure 2. Basic Learning and sensory signals in S1. **a,** Learning curve for 3 mice trained on the basic Go/No-Go detection task with a weak stimulus (4 degree). Performance is expressed as the probability of a hit (lick after stimulus) and false alarm (lick after no stimulus) for a given session. Individual mice are represented by symbols, and the mean by bold lines. **b,** Fluorescent activity in S1 from an example mouse during learning of the task. Shown are frames at the peak response to the 4 degree stimulus, catch trials are shown below. Scale bar = 1 mm. **c,** Dprime metrics for both behavioral and neuronal data during learning of the task. The d'_{behav} was calculated from the observed hit rate and false alarm rate for a given training session. The d'_{neuro} was calculated by comparing single trial distributions of all evoked signal peaks (maximum $\Delta F/F_0$ (%) within 100 ms post stimulus) with the corresponding noise distributions when no stimulus was present. Shown are means across all data for a given session, errorbars represent bootstrapped estimates of 95% confidence limits. The dotted lines separate performance into naive ($d'<0.8$), and acquired ($d' > 1.5$). The right panel shows the same data separated for individuals (symbols) before (naive) and after learning (acquired). Bars represent means across mice, errorbars represent SD. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; 'n.s.' not significant, two-sided Wilcoxon rank-sum test or Kruskal-Wallis test.

Figure 3. Adaptive behavior and S1 responses in the experienced mouse. a, Manipulation of stimulus distribution range. Every stimulus and catch trial (not shown) is presented with equal probability (p = 0.2). The design involves amplitudes common to both high-range (magenta) and low-range (green) conditions. **b,** Psychometric curves and response thresholds for an example animal (M04) working in both conditions. Each dot corresponds to response probabilities from a single session. Solid curves are logistic fits to the average data (n=10-11 sessions). Dotted line is a hypothetical curve assuming no change in performance (H_0). Dashed line is a hypothetical curve assuming a change in performance to maintain reward (H_1) when switching from high range to low range stimulus distributions. Response thresholds are shown as vertical lines with 95% confidence limits. Inset. Response thresholds of all mice (grey symbols). Bars represent means across mice with SD. **c,** Number of rewards (correct trials) accumulated by the same animal from b. Each line corresponds to one session. Inset. Average total reward number per session for each mouse. The average number of trials is shown on top. Figure conventions are the same as in b. **d,** Frames of evoked cortical fluorescence activity from two example sessions (M04), one with a high range stimulus distribution (top) and the other with a low range stimulus distribution (bottom). The frames are aligned for amplitudes common to both datasets. Data with a deflection angle of 8 degrees was chosen for further analysis (outlined box). **e,** Temporal fluorescent signal in response to 8 degree stimulation, extracted from the region of interest and averaged across sessions and mice. PSTHs of behavioral lick responses are shown on top. **f,** Observer model based on signal detection theory. Behavioral adaptation can either be induced by changes in sensitivity (d') intrinsic to S1, changes in decision criterion by a downstream observer, or both. **g,** Distributions of evoked trials (signal) and catch trials (noise) computed separately for the high and low range condition. Dashed lines indicate mean ΔF/F$_0$, black numbers indicate dprime metrics. Blue indicates criterion metrics as computed from ROC curves. Statistical difference was assessed using bootstrapped estimates of 95% confidence limits for the dprime and criterion. * P < 0.05; ** P < 0.01; *** P < 0.001; 'n.s.' not significant, two-sided Wilcoxon rank-sum test or Kruskal-Wallis test).

Figure 4. Linking neuronal activity and behavior
a, Model incorporating the sensory input distribution, the neuronal response function derived from the GEVI signal in S1, and the behavioral readout. The experimentally controlled stimulus amplitude on a given trial s_i is drawn randomly from the input distribution. The evoked GEVI signal in S1 can then be expressed as a stimulus response function $G(s_i)$. To establish a link between $G(s_i)$ and the behavior, a mathematical function $f(.)$ is created that transforms $G(s_i)$ to a probability of a lick response $P(GO|s_i)$, i.e. the psychometric function. Combining both $G(s_i)$ and $f(.)$ results in a function $f[G(s_i)]$, as an estimate of the psychometric curve, $\hat{P}(GO|s_i)$, that can then be directly compared to the actual psychometric curve $P(GO|s_i)$. **b,** Example functions $G_{High}(s_i)$ and $G_{Low}(s_i)$ of an experienced mouse challenged with a change in stimulus statistics. Data points represent fluorescence changes ($\Delta F/F_0$) in response to different stimulus amplitudes averaged across sessions. The functions $G_{High}(s_i)$ and $G_{Low}(s_i)$ are represented by curves fitted to the data. **c,** Linker function computed separately for both conditions $f_{High}(.)$ and $f_{Low}(.)$. **d,** Combination of $G(s_i)$ and $f(.)$. The predictions from the model $f_{High}[G_{High}(s_i)]$ and $f_{Low}[G_{Low}(s_i)]$ (dotted curves) are superimposed on the measured psychometric functions $P_{High}(GO|s_i)$ and $P_{Low}GO|s_i)$ (solid curves). $\hat{P}_{Null}(GO|s_i)$ is only influenced by changes in $G(s_i)$, and thus serves as a null test for the prediction based on changes in neural activity in S1 alone. In going from the high to the low range condition, the fraction explained by $G(s_i)$, can be estimated by comparing the area under the curve of $\hat{P}_{Null}(GO|s_i)$ with the area under the curve of $f_{Low}[G_{Low}(s_i)]$.

Figure 5. S1 exhibits signature of experience. a, Psychometric thresholds for different experience levels. 'Novel' refers to animals that experience the first switch in stimulus statistics (left panel). 'Experienced' refers to the second switch or later (right panel). **b,** Behavioral adaptation is calculated as the change in psychometric threshold (Δ threshold) across mice and experience level. Bars represent means across mice, errorbars represent SD. **c,** Explanatory power of S1 versus downstream quantified for the behavior of all mice undergoing repeated task changes. Pie plots depict the fraction explained by S1 activity $G(s_i)$ versus downstream $f(.)$. Columns depict different mice, rows correspond to training stages. **d,** Percent explained by S1 versus downstream on average, errorbars represent SD across mice. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; 'n.s.' not significant, two-sided Wilcoxon rank-sum test.

Waiblinger et al., Adaptive Behavior and the Role of Primary Somatosensory Cortex

Supplemental Figures

Supplemental Figure 1. **Basic Learning with a different stimulus. a,** Learning curve for 3 mice trained on the basic Go/No-Go detection task with a strong stimulus (16 degrees). **b,** Fluorescent activity in S1 from an example mouse during learning. Shown are frames at the peak response to a 16 degree stimulus, catch trials are shown below. Scale bar = 1 mm. **c,** Dprime metrics for both behavioral and neuronal data during learning of the task. Note, mice achieved higher hit rates at the beginning of training (interim) and reached successful task acquisition in half the time if compared to mice detecting weak stimuli (Fig 2). The dotted lines separate performance into interim (intermediate performance, d' = 0.8-1.5), and acquired (d' > 1.5). The right panel shows the same data separated for individuals (symbols) before (interim) and after learning (acquired). Other figure conventions are the same as in Figure 2.

Supplemental Figure 2. The relationship of magnitude and activated area in GEVI imaging of sensory evoked response. a, Fluorescence activity in response to different stimuli. Each frame is normalized to the frame at stimulus delivery ($\Delta F/F_0$ = F-F_0/F_0). Shown are images at the response peak. The dotted line represents a slice through the images aligned with the maximum fluorescence. Scale-bar = 1 mm. **b,** Magnitude (response amplitude $\Delta F/F_0$) versus cortical area extracted from the slice in a. Activity patterns are shown in different shades of grey for different stimulus amplitudes. **c,** Relationship between fluorescence magnitude and width of activated area, both scale with stimulus strength in a highly correlated fashion. The width is derived from a threshold ($\Delta F/F_0$ = 0.25%, dashed line in b). Data points correspond to different stimulus amplitudes fitted with a linear regression. Plots show average data across mice (n = 4) and sessions (n = 40).

Supplemental Figure 3. Analysis to test stability of adaptive S1 response in the experienced mouse. a, Different normalization methods. F_0 was varied by using the mean fluorescence across different time windows before stimulus onset. The $\Delta F/F_0$ measurement is then calculated by $F-F_0/F_0$. Mean fluorescence traces (n=4 mice, 831-870 trials) in response to the high (magenta) and the low range (green) stimulus condition. The grey box depicts the window for calculating F_0. From left to right: F_0=0 ms (frame at stimulus delivery), F_0 from [-100,0] ms, and F_0 from [-200,0] ms. **b,** Single-trial signal-peak and noise distributions from the data analysis as in A. Red and green numbers are mean fluorescent values in % $\Delta F/F_0$ for the high and low range condition. Black numbers represent dprime metrics. **c,** Dprime and variance for different F_0 calculations. **d,** Receiver operating characteristic (ROC) curves created by shifting the criterion across the $\Delta F/F_0$ signal and noise distributions of the high and low range condition. The downstream criterion can be inferred by comparing the hit rate in ROC space with the average behavioral hit rate (dashed lines).

Supplemental Figure 4. Relative explanatory power of S1 versus downstream for multiple task changes. Pie plots depict the fraction explained by S1 [$G(s_i)$] versus downstream [$f(.)$] quantified for the behavior of a control animal undergoing four changes (high-low-high-low).