Emerging experience-dependent dynamics in primary somatosensory cortex reflect behavioral adaptation

Christian Waiblinger[1], Megan E McDonnell [1], Peter Y Borden[1], Garrett B Stanley[1*]

[1]Wallace H Coulter Department of Biomedical Engineering, Georgia Institute of Technology and Emory University, Atlanta, GA, USA

With 41 text pages, 6 main figures, 4 supplemental figures
Word counts: Abstract 148, Intro 448, Results: 3852, Discussion 1593

*Correspondence: Garrett B Stanley

Coulter Department of Biomedical Engineering
Georgia Institute of Technology & Emory University
313 Ferst Drive
Atlanta GA 30332-0535
USA

Phone: 404-385-5037
Fax: 404-385-5044
Email: garrett.stanley@bme.gatech.edu

ABSTRACT

Behavioral experience and flexibility are crucial for survival in a constantly changing environment.

Despite evolutionary pressures to develop adaptive behavioral strategies in a dynamically

changing sensory landscape, the underlying neural correlates have not been well explored. Here,

we use genetically encoded voltage imaging to measure signals in primary somatosensory cortex

(S1) during sensory learning and behavioral adaptation in the mouse. In response to changing

stimulus statistics, mice adopt a strategy that modifies their detection behavior in a context

dependent manner as to maintain reward expectation. Surprisingly, neuronal activity in S1 shifts

from simply representing stimulus properties to transducing signals necessary for adaptive

behavior in an experience dependent manner. Our results suggest that neuronal signals in S1 are

part of an adaptive framework that facilitates flexible behavior as individuals gain experience,

which could be part of a general scheme that dynamically distributes the neural correlates of

behavior during learning.

INTRODUCTION

Survival in a dynamically changing sensory environment requires a high degree of behavioral

flexibility and experience. While much is known about the origin and processing of sensory signals

in mammalian brains[1–3], far less is understood about how resultant behavioral strategies are

formed with practice and experience, and the role of primary sensory areas in this output process.

Studies have investigated the role of the primary sensory cortex in visual[4–6], auditory[7–10] and

somatosensory behaviors[11,12]. There are varying views on whether primary sensory cortex simply

relays an ascending sensory signal or plays a deeper role in decision making and behavior.

It is likely that primary cortical signals are highly dynamic and context driven, depending on the

implemented behavioral paradigm. Signals in primary somatosensory cortex can: enhance

stimulus selectivity with behavioral training[13], fluctuate according to the behavioral state[14], or even

remap depending on downstream signals[15]. Neuronal signals in primary sensory areas may be

highly dynamic, context or experience dependent, and part of an adaptive framework.

Here, we investigate perceptual capabilities of the primary somatosensory cortex (S1) during

learning and behavioral adaptation, using the highly conserved mouse vibrissa system. We

hypothesize that signals in S1 not only represent the strength of a sensory input but also play a

key role in the transformation of context dependent behaviors. To test this, we designed a series

of psychophysical experiments evaluating behavioral performance and neuronal activity during:

1) gradual learning of a basic detection task and 2) adaptation to repetitive changes in sensory

contingencies. To repeatedly measure signals of large neuronal pools across both training stages,

we performed chronic wide-field imaging of S1 activity with the genetically encoded voltage

indicator (GEVI) "ArcLight" [16,17] in behaving mice.

During learning of the basic task, S1 sensitivity is mostly stimulus driven and uncorrelated with

gradual changes in behavioral performance. Interestingly, basic detection was not abrogated by

S1 inactivation through lesioning. In contrast, our results further reveal that S1 activity correlates

with behavioral adaptation, following long-term exposure to changing sensory stimulus statistics.

Mice adopt a strategy that modifies their behavior in a way as to maintain reward in the face of

these changes: once an animal is trained to adapt to a change in stimulus statistics, neuronal

activity dynamically shifts between changes in S1 sensitivity and decision criterion downstream.

S1 inactivation through lesioning disrupted the adaptive behavior, suggesting the S1 primary

cortex is necessary for an adaptive response to dynamic stimuli.

Our findings suggest a translation of these context dependent changes between different brain

structures along the hierarchy, where S1 is not only producing primary neuronal signals in

response to tactile input, but also transducing signals necessary for adaptive behavior strategies

in a dynamically changing environment.

MATERIALS AND METHODS

Animals, surgery, and general procedures for behavioral testing.

All experimental and surgical procedures were approved by the Georgia Institute of Technology

Institutional Animal Care and Use Committee and were in agreement with guidelines established

by the NIH. Subjects were 13 male mice (C57BL/6, Jackson Laboratories), aged 4-6 weeks at

time of implantation. The basic procedures of virus delivery, head-plate preparation and cortical

imaging exactly followed the ones published in a recent paper[16]. In the following text, only

procedures pertaining to the specific procedures established here are described in detail.

Virus delivery

At least 4 weeks prior to experimentation, mice were anesthetized using isoflurane, 3-5 % in a

small induction chamber, and then placed on a heated platform (FHC, Inc.) to maintain body

temperature with a stereotaxic nose cone to maintain anesthesia. During the surgery, the

anesthesia levels were adjusted to 1-1.5 % to achieve ~1/s breathing rate in mice. For virus

delivery, 3 small craniotomies (burr holes of 0.7 mm diameter) were created over the barrel field

of S1 according to stereotaxic measurements taken from the bregma ([1 x 3 mm, 3 x 3 mm, 3 x 1

mm] bregma x lateral). The virus was loaded into a neural syringe (Hamilton Neuros Syringe

700/1700). The injection needle was initially lowered to 1000 µm below the pia surface for pre-

penetration and then retracted to the target depth of 500 µm, using a 10 µm resolution stereotaxic

arm (Kopf, Ltd.). Following a 1-minute delay to allow for tissue relaxation, each animal was

injected with 1.5 µl of adeno-associated virus (AAV)1-hsyn1-ArcLight-D-WPRESV40 (UPenn

Viral Vector Core, AV-1-36857P) at a flow rate of 0.05 µL∕min (0.5 µl each, for three injections).

After injection, the needle remained in place for an additional 5 minutes before slowly being

removed from the brain. The craniotomies were left to close naturally. The skull was sealed by

suturing the skin. Throughout the experiment, sterile techniques were used to keep the injection

area clean and free from infection. Additionally, opioid and non-steroidal anti-inflammatory

analgesic were administered (SR-Buprenorphine 0.8-1 mg/kg, SC, pre-operatively and

Ketoprofen 5-10 mg/kg, IP, post-operatively).

Head plate Implantation

After at least 4 weeks post injection, a metal head-plate was secured to the skull in order to reduce

vibration and allow head-fixation during imaging and behavior experiments. Following

anesthetization and analgesia, a large incision was made over the skull. The connective tissue

and muscles surrounding the skull were removed using a fine scalpel blade (Henry Schein #10).

The custom titanium head-plate formed an open ring (half-moon shape with an inner radius of 5

mm) and was placed on top of the two hemispheres with an extended bar above the cerebellum

(~10 mm, perpendicular to the midline of the skull). The extended bar was designed to attach to

a stainless steel holder, serving the purpose of stable head-fixation. The head-plate was attached

to the bone using a three stage dental acrylic, Metabond (Parkell, Inc.). The Metabond was chilled

using ice, slowly applied to the surface of the skull, and allowed to cure for 5-10 minutes. After

securing the head-plate, the skull was cleaned and covered with a thin layer of transparent

Metabond. During preparation for histological validation, the head-plate could not be separated

from the attached skull and the brain was extracted by removing the lower jaw. The final head-

plate and dental acrylic structure additionally created a well for mineral oil that helped maintain

skull transparency for the upcoming imaging sessions. Mice were allowed to recover for at least

7 days before habituation training. After wound healing, subjects were housed together with a

maximum number of three in one group cage and kept under a 12/12 h inverted light/dark cycle.

Cortical Lesions.

After the collection of at least 5 sessions of psychometric data and the mapping of spatial

activation using GEVI imaging, lesions of the barrel field in S1 were performed as a control

experiment (n = 4 mice). Animals were given water *ad libitum* for the day preceding the procedure.

Following anesthetization, a small craniotomy (burr hole of 0.7 mm diameter) was created over

the area in S1 with the highest extent of activation from the GEVI map. The neurotoxin ibotenic

acid[18] (aablocks, AA003BBF) at a concentration of 10 mg/mL was loaded into a neural syringe

(Hamilton Neuros Syringe 700/1700). The injection needle was lowered to 1000 µm depth for pre-

penetration, then retracted to a depth of 500 µm, using a 10 µm resolution stereotaxic arm (Kopf,

Ltd.). Following a delay of 5 minutes for tissue relaxation, animals were injected with 0.25 µL of

ibotenic acid at a flow rate of 0.05 µL/min. After injection, the needle remained in place for 5

additional minutes before being slowly retracted from the brain. The craniotomy was sealed using

the dental acrylic Metabond (Parkell, Inc.), and the silicone elastomer Kwik-Cast (Kwik-Cast

Sealant World Precision Instruments), for protection. Animals were allowed to recover for 1-3

days until activity and body weight were normal, and then behavioral testing and GEVI imaging

continued.

Whisker Stimulation.

Precise whisker deflections were performed using a calibrated galvo-motor (galvanometer optical

scanner model 6210H, Cambridge Technology) as described in a previous study[19]. The opening

of the rotating arm was narrowed with dental cement to prevent whisker motion at the point of

insertion. The rotating arm contacted a single whisker on the right of the mouse's face at 5 mm

(±1 mm tolerance) distance from the skin, and thus, directly engaged the proximal whisker shaft,

largely overriding bioelastic whisker properties. Distance and angle between stimulator and

whisker were systematically measured with a camera before each session to ensure that

stimulation was consistent. All the remaining whiskers were trimmed to prevent them from being

touched by the rotating arm. Across mice, different whiskers were chosen (C1, D1, D2 or E2), but

the same whisker was used throughout sessions within each mouse Voltage commands for the

actuator were programmed in Matlab and Simulink (Ver. 2015b; The MathWorks, Natick,

Massachusetts, USA). A stimulus consisted in a single event, a sinusoidal pulse (half period of a

100 Hz sine wave, starting at one minimum and ending at the next maximum). The pulse

amplitudes used (A = [0, 1, 2, 4, 8, 16] °, correspond to maximal velocities: V_{max} = [0 314 628

1256 2512 5023] °/s) or mean velocities: V_{mean} = [0 204 408 816 1631 3262] °/s) and were well

within the range reported for frictional slips observed in natural whisker movement[20,21].

Cortical GEVI Imaging.

ArcLight transfected mice were chronically imaged through the intact skull using a wide-field

fluorescence imaging system to measure cortical spatial activity (MiCAM05-N256 Scimedia, Ltd.).

Figure 1a shows the experimental apparatus and schematically describes the wide-field

fluorescence microscope. During all imaging experiments, mice were awake and head-fixed. The

head-plate was used as a well for mineral oil in order to keep the bone surface wet and maintain

skull transparency. The skull was covered with a silicone elastomer (Kwik-Cast Sealant World

Precision Instruments) between imaging sessions for protection. The barrel cortex was imaged

using a 256 x 256 pixel CMOS camera (Scimedia Model N256CM) at 200 Hz with a pixel size of

69 μm x 69 μm and an active imaging area of 11.1 x 11.1 mm, given a magnification of 0.63. Note,

this resolution does not consider the scattering of the light in the tissue. During experimental

imaging, the illumination excitation light was left continuously on. The entire cortical area was

illuminated at 465 nm with a 400 mW/cm^2 LED system (Scimedia, Ltd.) to excite the ArcLight

fluorophore. The excitation light was further filtered (cutoff: 472-30 nm bandpass filter, Semrock,

Inc.) and projected onto the cortical surface using a dichroic mirror (cutoff: 495 nm, Semrock,

Inc.). Collected light was filtered with a bandpass emission filter between wavelengths of 520-35

nm (Semrock, Inc.). The imaging system was focused at ∼300 μm below the cortical surface to

target cortical layer 2/3. The first imaging session was used for identifying the barrel field in the

awake and naïve animal at least four weeks after ArcLight viral injection. The barrel field was

mapped by imaging the rapid response to a sensory stimulus given to a single whisker (A = 4-

16°, or mean velocities respectively: V = 816- 3262°/s). We used two criteria to localize and isolate

the barrel field: standard stereotaxic localization (∼3 mm lateral, 0.5-1.5 mm caudal from bregma)

and relative evoked spatial and temporal response (visible evoked activity 20-25 ms after

stimulation). A single whisker was chosen if it elicited a clear response within the barrel field. All

subsequent imaging experiments were centered on the same exact location and the same

whisker was chosen in repeated sessions for a given animal. Figure 1b shows the characteristic

spread of ArcLight expression in an example coronal brain section. The same hemisphere is

shown *in vivo* in Figure 1c with spontaneous fluorescence activity in S1 at the beginning of a

behavioral session. Spontaneous and sensory evoked activity in S1 was imaged every trial, 2 s

before and 2 s after the punctate whisker. Figure 1d shows frames with typical fluorescence

activity patterns recorded at a framerate of 200 Hz and shown separately for three different

stimulus amplitudes (rows). The calculation of the $\Delta F/F_0$ metric, region of interest (ROI) and data

processing is described under "Imaging analysis".

Behavioral paradigm and training

During successive days of behavioral testing, water intake was restricted to the experimental

sessions where animals were given the opportunity to earn water to satiety. Testing was paused

and water was available *ad libitum* during 2 days a week. Body weight was monitored daily, and

was typically observed to increase or remain constant during training. In some cases, the body

weight dropped slightly across successive training days due to a higher task difficulty. If the weight

dropped for more than ~5g, supplementary water was delivered outside training sessions to

maintain the animal's weight. 1-2 training sessions were usually conducted per day comprising

50-250 trials. During behavioral testing a constant auditory white background noise (70 dB) was

produced by an arbitrary waveform generator to mask any sound emission of the galvo-motor-

based whisker actuator. All mice were trained on a standard Go/No-Go detection task (Fig. 1e)

employing a similar protocol as described before [22–26]. In this task, the whisker is deflected at

intervals of 4-10 s (flat probability distribution) with a single pulse (detection target). A trial was

categorized as a "hit" if the animal generated the "Go" indicator response, a lick at a water spout

within 1000 ms of target onset. If no lick was emitted the trial counted as a "miss". In addition,

catch trials were included, in which no deflection of the whisker occurred (A = 0°) and a trial was

categorized as a "correct rejection" if licking was withheld ("No-Go"). However, a trial was

categorized as a "false alarm" if random licks occurred within 1000 ms of catch onset. Premature

licking in a 2 s period before the stimulus was mildly punished by resetting time ("time-out") and

starting a new inter-trial interval of 4-10 s duration, drawn at random from a flat probability

distribution. Note these trial types were excluded from the main data analysis.

The first step of behavioral training was systematic habituation to head-fixation and experimental

chamber lasting for about one week. During the second training phase, a single whisker deflection

with fixed amplitude was presented interspersed by catch trials (P_{stim} = 0.8, P_{catch} = 0.2).

Immediately following stimulus offset, a droplet of water became available at the waterspout to

condition the animal's lick response thereby shaping the stimulus-reward association. Once

subjects showed stable and immediate consumption behavior (usually within 1-2 sessions), water

was only delivered after an indicator lick of the spout within 1000 ms, turning the task into an

operant conditioning paradigm in which the response is only reinforced by reward if it is correctly

emitted after the stimulus. Subsequent experiments were performed systematically and the

behavioral performance was measured with simultaneous GEVI imaging. The different

experiments are described in detail in the following section.

Basic learning: Learning was studied once an animal had entered the operant phase of training

after the basic habituation procedure. From this point forward, experiments were conducted with

equal conditions across sessions and without manual interference by the experimenter. To assess

differences in learning based on stimulus strength, animals were separated into two groups: group

1 (n = 3) receiving only one low amplitude stimulus and catch trials (A = [4 0] °, Fig. 2) and group

2 (n = 3) receiving only one high amplitude stimulus and catch trials (A = [16 0] °, Suppl. Fig. 2).

Performance metrics are described under *data analysis and statistics*.

Adaptive behavior: After mice had learned the basic detection task, the psychometric curve was

measured in a subgroup of animals (n = 10) using the method of constant stimuli, which entails

the presentation of repeated stimulus blocks containing multiple stimulus amplitudes. On a single

trial, one out of multiple possible stimulus amplitudes was presented after a variable time interval

(4-10 s), each with equal probability (uniform distribution, p = 0.2). A stimulus block consisted of

a trial sequence comprising all stimuli and a catch trial in pseudorandom order (e.g., each type

once per block). A behavioral session consisted of repeated stimulus blocks until the animal

disengaged from the task, i.e. when it did not generate lick responses for at least an entire

stimulus block. Therefore, the chosen stimulus occurred repetitively but randomly within a

session. A condition was always kept constant within and across multiple behavioral sessions

before the task was changed. A "switch" was defined as the transition between two sessions with

different conditions. In the "high range" condition, four stimulus amplitudes plus catch trial were

used (A = [0, 2, 4, 8, 16] °) and presented in multiple successive sessions. Following this, four

new stimulus amplitudes were presented (A = [0, 1, 2, 4, 8] °) forming the "low range" condition.

Both stimulus distributions shared two of the three stimulus amplitudes; however, the largest

stimulus amplitude of the high range condition (A = 16 °) was not part of the low range, and vice

versa, the smallest amplitude of the low range condition (A = 1 °) was not part of the high range.

Conditions were always changed in alternating fashion to test the reversibility of adaptation (e.g.

high-low-high or vice versa) and multiple switches were performed to test adaptation at different

training stages. The different timescales are summarized as follows:

Number of trials: the number of trials were variable since each session was stopped after an

animal missed an entire set of stimuli (i.e. A = [0, 2, 4, 8, 16] ° for the high range). The reason for

this implementation is to ensure that the data are not affected by the animal's potential satiation

on any given day. As shown in Figure 3c, the number of trials per session averages out to

approximately one hundred trials (n = 101 for high range and n = 104 for low range).

Number of sessions: Within a block there are 5-11 daily sessions depending on the animal.

However, each animal experienced a matched number of sessions across blocks (e.g. n = 5 for

high range and n = 5 for low range).

Number of blocks: Depending on the animal, 3-6 blocks were performed with alternating

conditions either starting with the high range or the low range condition (resulting in 2-5 switches).

Histology.

Upon completion of experiments, transcardiac perfusion was performed on all animals with a 4 %

paraformaldehyde solution in phosphate-buffered saline. The head plates were carefully removed

from the skulls, and the brains were extracted and post-fixed for 2-24 hours depending on the

success of the perfusion. Brains were then sectioned at 100µm thickness on a vibratome (Leica

VT1000S). The brain sections were stained with Cytochrome C oxidase and DAPI for histological

analysis of lesion extent or ArcLight expression. Sections were mounted and imaged using a wide

field color microscope (Zeiss Axio Observer Z1).

Data analysis and statistics.

Behavior. The learning curve was measured by calculating a dprime, d'_{behav}, from the observed

hit rate and false alarm rate of each training session

243
$$d'_{behav} = Z(hit) - Z(fa)$$

where the function Z(p), p ∈ [0,1], is the inverse of the cumulative distribution function of the

Gaussian distribution. A criterion of d' = 2.3 (calculated with $p(hit)$ = 0.95 and $p(fa)$ = 0.25) was

used to determine the end of the basic learning period and learning progress was separated into

three equally distributed stages, "detect-naive" (d' = 0-0.8), "detect-interim" (d' = 0.8-1.5) and

"detect-experienced" level (d' = 1.5-2.3). Psychometric data were assessed as response-

probabilities averaged across sessions within a given stimulus condition.

Psychometric curves were fit using Psignifit[27–29]. Briefly, a constrained maximum likelihood

method was used to fit a modified logistic function with 4 parameters: α (the displacement of the

curve), β (related to the inverse of slope of the curve), γ (the lower asymptote or guess rate), and

λ (the higher asymptote or lapse rate) as follows:

$$P(GO|s_i) \; = \gamma \; + \; (1 \; - \; \gamma \; - \; \lambda) \; \frac{1}{1 + \exp{(-z(s_i))}}$$

$$z(s_i) \; = \; \frac{s_i - \alpha}{\beta}$$

where s_i is the stimulus on the i^{th} trial. Response thresholds were calculated from the average

psychometric function for a given experimental condition using Psignifit. The term "response

threshold" refers to the inverse of the psychometric function at some particular performance level

with respect to the stimulus dimension. Throughout this study, we use a performance level of 50%

(probability of detection of 0.5). Statistical differences between psychophysical curves were

assessed using bootstrapped estimates of 95% confidence limits for the response thresholds

provided by the Psignifit toolbox. To assess the effects of the lesion on behavior, the psychometric

curves and response thresholds were compared in lesion animals, from before and after the

lesion, and between high- and low-range stimuli. Kruskal-Wallis or rank sum tests were used to

compare significance of the lesion effects on response threshold in lesioned versus healthy

animals.

Reward accumulation. Let the stimulus amplitude delivered on the i^{th} trial be denoted as s_i, the

corresponding reward as r_i, (since r_i is a fixed value, it can just be termed r) and the accumulated

reward for N trials as R_N. Over N trials, the expected accumulated reward is

$$E\{R_N\} = \sum_{i=1}^{N} P(s_i)P(GO|s_i)r_i$$

where $P(s_i)$ comes from the experimentally controlled stimulus distribution, $P(GO|s_i)$ is the

probability of a positive response (or "Go") for the given stimulus amplitude, and $E\{\}$ denotes

statistical expectation.

We considered the null hypothesis of this behavioral paradigm to be that animals do not adapt

their behavior in response to an experimentally forced change in stimulus distribution and thus

operate from the same psychometric curve (represented as dotted curves in Fig. 3b). Note, this

corresponds to the same curve but for a different range of stimuli ($P(GO|s_i)$ across different s_i.

For example, in moving from the high range to the low range stimulus condition, this would result

in a decrease in the total accumulated reward for the same number of trials.

As an alternative hypothesis, one possible strategy the animal could take in response to a change

in the stimulus distribution would be to adjust behavior to maintain the same amount of

accumulated reward during a session. For example, in moving from high range to low range

stimuli, the accumulated reward would be assumed fixed, and we can determine a new set of

probabilities $P(GO|s_i)$ that define an adapted psychometric function. Note that there is not a

unique solution, but one simple possibility is that the original psychometric function maintains the

same asymptotes (γ and λ) and false alarm rate but is compressed, with a decrease in response

threshold and an increase in slope to maintain the same total accumulated reward. We denote

this situation as our hypothetical psychometric function, represented as dashed curves in Figure

3b.

Imaging analysis. All voltage imaging data was analyzed using custom written image-analysis

software (MATLAB 2015a, Mathworks, Inc.). The specific methods of processing the ArcLight raw

fluorescence signal and basic data analysis followed those of a recent study from our laboratory[16].

Briefly, raw images were loaded and converted from the proprietary file format of the imaging

system using custom scripts. Due to the natural decay of the fluorescent signal caused by photo

bleaching, each trial was first normalized to a baseline and reported as a percent change in

fluorescent activity (% $\Delta F/F_0$). The $\Delta F/F_0$ measurement was calculated by subtracting and dividing

each trials fluorescence *F(x, y, t)* by the frame preceding the stimulus delivery:

298
$$\frac{\Delta F}{F_0} = \frac{F - F_0}{F_0}$$

where $F_0(x, y)$ is the frame of stimulus delivery ($F_0 = F$ at $t = 0$). Note, an extended analysis was

performed with different normalization methods by subtracting and dividing each trials

fluorescence F by the fluorescence averaged across different time windows before stimulus onset

($t = 0$ ms, t from [-100,0] ms, t from [-200,0] ms). Increasing normalization windows slightly altered

the change in fluorescence magnitude and variance of the evoked response; however, varying

the normalization window did not affect the adaptive cortical response reported in this study

(Suppl. Fig. 3).

A single region of interest (ROI) was identified using the largest 10 x 10 pixel (434 x 434 μm) area

response 20-25 ms following stimulus onset. The average activity within this region was extracted

across all frames to compute the temporal dynamics of the fluorescent signal. Note, due to the

fluorophore[17], positive changes in membrane potential correspond to a decrease in ArcLight

fluorescent activity. In line with our previous study[16] all traces have been inverted to show a

positive increase in fluorescence. Fluorescent voltage traces and behavioral lick responses were

acquired within the same time window and aligned with regard to stimulus onset (Fig. 1f). After

an animal was lesioned, GEVI responses were qualitatively compared to the responses recorded

before lesion. Temporal dynamics and overall fluorescence were assessed.

Ideal observer analysis. To quantify the fluorescence signal over the course of learning a metric

d'_{neuro} was computed. For a given day or session, single trial distributions of evoked signal

peaks (maximum % $\Delta F/F_0$ within 100 ms post stimulus) were compared to the corresponding

noise distributions when no stimulus was present. d'_{neuro} is then defined as:

319
$$d'_{neuro} = \frac{\mu S - \mu N}{\sqrt{\frac{1}{2}(vS + vN)}}$$

where μS and μN are the mean and vS and vN are the variance of the signal and noise distribution.

d'_{neuro} was then directly compared with the behavioral learning curve derived from d'_{behav}.

The same analysis was repeated for the data acquired with changing stimulus statistics, with the

exception that only trials with an intermediate stimulus amplitude shared between the high and

the low range condition were used. Note, the chosen stimulus of 8-degree whisker angle

represents the midpoint of the high range and the upper limit of the low range condition.

Distributions consisting of all evoked trials termed "signal" (peak $\Delta F/F_0$ with 8 °) and catch trials

termed "noise" peak ($\Delta F/F_0$ with 0 °) were computed separately for the high and low range

condition with one switch from high to low or vice versa. Statistical differences between

distributions were assessed using bootstrapped estimates of 95 % confidence limits for the d'

metrics. This means resampling the d' from a given session with replacement 1000 times, taking

the average of each resampled dataset, and then taking the interval that spans the central 95 %

of this distribution of averages across resampled datasets. Significance values were further

estimated with a non-parametric Wilcoxon rank-sum or Kruskal-Wallis test. Throughout this

manuscript, * indicates $p < 0.05$; ** indicates $p < 0.01$; *** indicates $p < 0.001$; and "n.s." indicates

"not significant".

Cascade framework. This analytical framework was created to describe the correspondence

between the sensory input distribution, the neuronal response function derived from the GEVI

signal in S1, and the behavioral readout. The experimentally controlled stimulus amplitude on a

given trial s_i is drawn randomly from the input distribution. The evoked GEVI signal in S1 can then

be expressed as a stimulus response function $G(s_i)$. To establish a link between $G(s_i)$ and the

behavior, a mathematical function $f(.)$ was created that transforms $G(s_i)$ to a probability of a lick

response $P(GO|s_i)$, i.e. the psychometric function, as illustrated in Figure 5a and b. In other words

$f(.)$ represents a matching of the evoked fluorescence to the lick response, given a particular

stimulus. Combining both $G(s_i)$ and $f(.)$ results in a function $f[G(s_i)]$, as an estimate of the

psychometric curve, $\hat{P}(GO|s_i)$, that can then be directly compared to the actual psychometric

curve $P(GO|s_i)$. This approach can be applied across the different stimulus conditions (high

versus low), and comparisons made across the corresponding $G(s_i)$ and $f(.)$ functions.

To quantitatively estimate how much of the behavioral variation can be explained by S1 activity

versus downstream, the following control was performed: $G(s_i)$ is considered to change between

the high and the low range condition, as observed experimentally, from $G_{High}(s_i)$ to $G_{Low}(s_i)$. As

a null test for the transition from the high to low condition, to capture how much of the observed

changes in behavior is explained solely by the changes in $G(s_i)$, $f(.)$ is held constant, operating

from a function $f_{High}(.)$, that only reflects the high range condition. The combination of $G_{Low}(s_i)$

and $f_{High}(.)$ then produces an estimated psychometric function $\hat{P}_{Null}(GO|s_i)$ that is only

influenced by changes in $G(s_i)$, and thus serves as a null test for the prediction based on changes

in neural activity in S1 alone.

Because experiments were designed to present a range of stimulus amplitudes, comparisons of

the fluorescence (Fig. 4c) and behavioral (Fig. 5a, b) responses involved comparing curves. One

simple way to do this is to compare the area under the curves (AUCs) by taking the difference

between the area under the curve across the conditions. For the observed GEVI fluorescence

signals for the high and low range stimulus conditions (Fig. 4c), this difference was:

$$\Delta AUC(Low - High) = AUC\big(G_{Low}(s_i)\big) - AUC\big(G_{High}(s_i)\big)$$

where the area was computed across the "overlap range" of stimulus amplitudes.

Similarly, to quantify the relative explanatory power of S1 versus downstream processing (Fig.

5c, d, e), we implemented the following. The behavioral adaptation was quantified by subtracting

the areas of the model fits for the high and low range conditions:

367
$$\Delta AUC(Low - High) = \ AUC(f_{Low}[G_{Low}(s_i)]) - \ AUC(f_{High}[G_{High}(s_i)])$$

Examples of data and model fits are in the right column of Figure 5a and b. Since, by construction,

this analysis involved components from both S1 and downstream of S1, this allowed us to

evaluate the relative contributions of each. Assuming that $G(s_i)$ changed from $G_{High}(s_i)$ to

$G_{Low}(s_i)$ but $f(.)$ is constant ($f_{High}(.)$), this resulted in a model prediction $\hat{P}_{Null}(GO|s_i)$ for the

behavior performance, if the effect were due to S1 alone. The prediction for the behavioral

adaptation based on a change in $G(s_i)$ (and thus S1) alone was then calculated as follows:

374
$$\Delta AUC(Low - High)_{S1} = \ AUC(\hat{P}_{Null}(GO|s_i) - \ AUC(f_{High}[G_{High}(s_i)])$$

375   Finally, the fraction explained by S1 was then calculated from $\Delta AUC(Low - High)$ and

376    $\Delta AUC(Low - High)_{S1}$ as follows:

377
$$\% \ Explained \ by \ S1 = \frac{\Delta AUC(Low - High)_{S1}}{\Delta AUC(Low - High)} \times 100$$

and the remaining was explained downstream of S1, or $\% \ Explained \ downstream = 100 -$

$\% \ Explained \ by \ S1$.

All curves were fit using the psignifit toolbox[27] and the goodness of fit was assessed by calculating

metrics of deviance (D) as well as the corresponding cumulative probability distribution (CPE). To

rule out the possibility of poor fitting in the cascade framework, we inspected the goodness-of-fit

metric of deviance (D) as well as estimates of where the goodness-of-fit lay in bootstrapped

cumulative probability distributions of this error metric (CPE) using the psignifit toolbox. Due to

the steep increase in $\Delta F/F_0$ at lower stimulus amplitudes we find that a Weibull function provides

the best fit for $G(s_i)$. Both, the linker function $f(.)$ and the psychometric function are best fit by a

logistic function due to the sigmoid configuration of the data.

RESULTS

The current study investigates learning and experience dependent adaptation in the mouse

vibrissa system. Our main goal is to establish a relationship between controlled whisker inputs

(stimulus), S1 activity, and licking response (behavioral output). We manipulated stimulus inputs

and observed S1 activity during early basic detection learning. We then observed these animals

longitudinally for flexible adaptation in a changing environment. To repeatedly measure signals of

large neuronal pools at different training stages, we performed chronic wide-field imaging of S1

activity with the genetically encoded voltage indicator (GEVI) "ArcLight" in behaving mice.

Figure 1 outlines the experimental design and summarizes the basic neuronal and behavioral

metrics of this study. Seven mice were transfected with the GEVI ArcLight before being imaged

through the skull using a wide-field fluorescence microscope (Fig. 1a). We have previously

described this imaging technique in detail[16]. Figure 1b shows the characteristic spread of ArcLight

expression in an example coronal brain section. The same hemisphere is shown *in vivo* in Figure

1c with spontaneous fluorescence activity in S1 at the beginning of a behavioral session. Figure

1d shows sequential frames of typical fluorescence activity patterns in a trained mouse recorded

in response to deflections of a single whisker.

Transfected mice were first trained on a tactile Go/No-Go detection task [22,23,26,30,31]. Animals detect

pulse-shaped deflections of a single whisker and generate a lick on a waterspout ("Go") or, if no

stimulus is present, animals withhold licking ("No-Go") (Fig. 1e). In response to stimulation, trained

mice first show a clear cortical fluorescence response within 100 ms, then stereotypical lick

response approximately 200 ms later (Fig. 1f). Note, the temporal resolution of ArcLight allows us

to identify the fluorescence response peak in S1 (maximum % $\Delta F/F0$) within the 100 ms window

post stimulus which clearly precedes the typical behavioral lick response. Hence, the sensory

signal in S1 can be disentangled from potentially interfering motor related signals caused by

licking. For early "basic learning" experiments the neuronal S1 signal was observed in mice

acquiring the basic principles of the Go/No-Go task, as demonstrated by learned lick responses

(Fig. 1g, left). For later "adaptive behavior" experiments (Fig. 1g, right), S1 was imaged in mice

challenged with randomly presented whisker deflection amplitudes and systematically

manipulated statistical distributions.

S1 responses during basic learning

All subjects received uncued single whisker stimulations at random intervals of 4-10 s with a single

pulse or no stimulation (catch trial). Learning was measured by calculating the hit $p(hit)$ and false

alarm rate $p(fa)$ of successive daily trainings. A criterion of $p(hit)$ = 0.95 and $p(fa)$ = 0.25 was

used to determine successful acquisition of the task. Figure 2a shows hit and false alarm rates

for three mice trained on the weak stimulus (A = 4°). With this weak stimulus, subjects required

extended training for more than 30 sessions to achieve successful performance.

Figure 2b shows frames of cortical S1 fluorescence activity from an example mouse over the

course of learning. Each square represents a top view of the same cortical hemisphere on

selected training days with the average fluorescent activity ($\Delta F/F_0$ in %) at the frame

corresponding to the peak of the sensory evoked signal (A = 4°, top row) or during catch trials (A

= 0°, bottom row). The magnitude and spread of fluorescence is highly variable throughout the

"basic learning" daily sessions, and a significant trend cannot be identified with learning progress.

To further quantitatively evaluate the fluorescence signal over the course of learning, we used

classical signal detection theory[32,33] with a computed neuromeric sensitivity measure d'$_{neuro}$. Single

trial distributions of evoked signal peaks were compared to the corresponding noise distributions,

when no stimulus is present. The d'$_{neuro}$ for a given day or session was then calculated by

subtracting the means of the distributions and dividing by the variance. To compare this metric to

the behavior, a d'_{behav} was calculated from the observed behavioral hit and false alarm rate of

each training session (see methods). Figure 2c shows the d'_{neuro} (orange) along with the

behavioral learning curve d'_{behav} (gray), across all sessions and mice. The dashed lines separate

performance into "detect-naive" (d'<0.8), and "detect-experienced" (d'>1.5). In contrast to the

behavior, the average neuromeric sensitivity remains at a relatively constant level, representing

a stable signal to noise relationship independent of the continuing learning progress. Note, a

second group of animals (n = 3) was trained on a much stronger stimulus for comparison (A =

16°). Those mice achieved much higher hit rates at the beginning of training and reached

successful task acquisition in half the time (Suppl. Fig. 1a); however, the neuromeric sensitivity

was also orthogonal to the learning progress (Suppl. Fig. 1b-c). We conclude that neuronal

sensitivity in S1 measured from a large population of neurons does not change during basic

learning.

Adaptive behavior and changes in S1

To investigate behavioral and neuronal dynamics with regard to changing context, we performed

experiments in which we systematically manipulated stimulus statistics (Fig. 3a). The

psychophysical techniques were adapted from a behavioral paradigm we recently developed in

the rat[26] which are described in materials and methods. The first distribution consists of four

different stimulus amplitudes of whisker deflection, as well a catch trial (A = [0, 2, 4, 8, 16] °,

magenta) which we refer to as the "high range" condition. The second distribution consists of four

new stimulus amplitudes of whisker deflection, and a catch trial (A = [0, 1, 2, 4, 8] °, green), which

we refer to as the "low range" condition. Each stimulus or catch trial was presented with equal

probability (uniform distribution). Importantly, the experimental design involves amplitudes

common to both high-range and low-range conditions, three of the four stimulus amplitudes were

shared between conditions. Figure 3b depicts typical psychometric curves from an example

mouse, performing the task first under the high range (magenta) and low range condition (green).

There is a consistent shift of the psychometric curve in response to the changing stimulus

statistics, which we refer to here as "adaptive behavior". All animals experienced both a switch

from high range to low range, as well as low range to high range. Adaptation was reversible,

meaning that over multiple switches, the animal's performance changed, indicating that they are

able to apply a reciprocal task strategy, modulated by the shift in stimulus distribution.

The simple reward expectation model tests if the animal adapts its behavior to maintain

accumulated reward in the face of a changing stimulus distribution. If the animal does not adjust

its behavior the predicted low range psychometric function would align with the high range

psychometric function (Fig. 3b black dotted curve under magenta curve; H_0). In moving from the

high range to the low range condition, this would result in a decreased reward rate for the same

number of trials. However, if the animal adjusts its behavior, the model predicts that the

psychometric function shifts to the left such that the expected reward per trial remains constant

(Fig. 3b black dashed curve; H_1).

We observed a shift of the psychometric curve when going from high range to low range

performance. The experimentally measured psychometric function in the low range condition

(green) comes quite close to the hypothetical performance, suggesting that the animal adapts its

behavior to maintain reward. The psychometric shift results in a significant decrease in response

threshold for all mice (Fig. 3b inset). Figure 3c depicts the actual trial-by-trial reward accumulation

by the example mouse. Overlaid are results for n = 11 sessions with the high range distribution

(magenta) and n = 10 sessions with the low range distribution (green). The slope of reward

accumulation in the low range condition nearly matches that of the high range condition, and the

slope for the low case (green) is close to the prediction from the maintenance of accumulated

reward hypothesis, H1 (dashed line), while being clearly separable from the slope representing

the null hypothesis (dotted line). The total number of rewards acquired on average per session

and across all mice further confirms this (total # high range = 44.5 ± 8.6, total # low range = 42.0

± 10.6, Mean and SD, Fig. 3c inset), whereas there was no evidence for an alternative strategy

to maintain the total number of rewards by working substantially more trials. The total number of

trials per session was nearly the same for each condition on average (high range, n = 101 and

low range, n = 104, Fig. 3c inset). These findings show that detection behavior is highly flexible

in the face of a changing stimulus distribution.

Figure 3d shows frames of evoked S1 fluorescent activity in a mouse exposed to high range

stimuli (top) or low range stimuli (bottom). In both conditions, we observed an increase in

fluorescence amplitude and spatial spread, with increasing stimulus amplitude. Note, there is a

relatively consistent linear relationship between the percent change in fluorescence magnitude

and the activated cortical area, as both scale with stimulus strength in a highly correlated fashion

(Suppl. Fig. 2). For simplicity, we use percent change in fluorescence magnitude as a metric for

cortical activation. The evoked activity for the same stimulus is clearly higher in the low range

compared to the high range condition.

To further dissect this change, we focused our analysis on trials with an intermediate stimulus

shared between both datasets (outlined box in Fig. 3d). The chosen whisker deflection amplitude

of 8 degrees represents the midpoint of the high range distribution and the upper limit of the low

range distribution. Figure 3e depicts the temporal fluorescent signal in response to an 8°

amplitude, averaged across sessions from all four mice. These mice have been exposed to both

high and low range conditions as well as a switch between conditions. Histograms of behavioral

lick responses are shown on top – the lick response occurs approximately 200 ms window after

the measured fluorescence peak. We did not observe a significant difference in behavioral

response latencies between the high-range and the low range condition. However, the evoked

S1 response to the same 8° deflection was higher in animals challenged with the low range vs

the high range stimuli.

Figure 3f outlines the observer model we used based on signal detection theory, to interpret and

analyze data from the S1 neural signals, as well as a presumed downstream decision criterion.

Multiple scenarios are considered to trigger a change within sensory and higher order processing

stages; behavioral adaptation can be induced by changes in sensitivity (d') intrinsic to S1,

changes in decision criterion (c) by a downstream observer, or both. Sensitivity in S1 improves

through reduction in the overlap between sensory signal and noise distributions (top panel). In

addition, the downstream observer may value hits and false alarms differently by altering the

decision criterion (bottom panel). To test these predictions, distributions consisting of many

evoked trials termed "signal" ($\Delta F/F_0$ with 8 °) and catch trials termed "noise" ($\Delta F/F_0$ with 0 °) were

computed separately for the high and low range condition (Fig. 3g, top). The noise distribution

(grey) is comparable for the high and the low range condition with its means being identical

(dashed vertical black lines). However, the signal distribution for the low range (green) is shifted

towards higher fluorescence changes as compared to the signal distribution of the high range

(magenta) with a clear difference in its mean (dashed vertical magenta and green lines). As shown

in the bottom of Figure 3g, a significant difference in neuronal sensitivity was confirmed by

calculating the neuromeric d' as introduced earlier in this study. While the neuronal d' was altered

slightly by different normalization methods of the $\Delta F/F_0$ metric, an extended analysis revealed that

the difference in d' across conditions persisted (Suppl. Fig. 3a-c). In addition, we created receiver

operating characteristic (ROC) curves by varying a criterion threshold across the $\Delta F/F_0$ signal and

noise distributions and plotting the hit rate (signal detected) against the false alarm rate (incorrect

guess) (Suppl. Fig. 3d). The ROC curve for the low range condition was higher than that for the

high range condition, quantified by a larger area under the low range ROC curve (AUC_{low} = 0.77)

than for the ROC curve for the high range condition (AUC_{high} = 0.69), thereby confirming a change

in S1 sensitivity.

Changes in criterion were inferred by comparing the hit rate in ROC space (neurometric) with the

average hit rate measured from the behavior (psychometric). The criterion shows a slight, yet

non-significant decrease when operating from the low range compared to the high range condition

(Fig. 3g, bottom left, blue). Note, the S1 sensitivity and downstream criterion change in opposite

direction, e.g. an increase in hit rate can be caused by an increase in S1 sensitivity and/or a

decrease in criterion. We conclude that, in highly trained animals there is evidence of adaptive

sensitivity in S1, yet comparatively smaller adaptive changes in criterion by a downstream

observer.

Changes across training stages

The data described thus far, comparing the adaptive behavior and cortical activity across the high

and low range conditions was derived from highly trained subjects. To better understand the

development of this phenomenon over time we re-analyzed the data from highly trained subjects

at different training stages. Conditions were always changed in alternating fashion to test the

reversibility of adaptation (high-low-high etc.). Figure 4a shows the psychometric threshold for

mice undergoing repetitive task changes starting with either the high range or low range condition.

We observed that the order of stimulus conditions does not affect behavioral adaptation and

response thresholds closely follow the direction of task manipulation. This suggests that mice are

able to apply a flexible and reciprocal task strategy modulated by the shift in stimulus distribution.

Behavioral adaptation was stable across multiple training stages, as revealed by consistently

modulated thresholds.

Since individual mice experienced different numbers of switches (n = 2-5) and different numbers

of sessions for a given condition (n = 5-11 per condition), we asked how many sessions or trials

are generally required to adapt after a switch? We focused our analysis on the transition from one

condition to another. Figure 4b shows psychometric thresholds extracted from consecutive

sessions at the high- to low range transition (left) and at the low- to high range transition (right).

The data were normalized with regard to the transition and data from all switches were included

in this analysis. Across mice, the target threshold for the high or low range condition (average

threshold per condition) was reached within approximately three sessions, indicating that

behavioral adaptation to changing stimulus statistics required repeated exposure to

approximately 300 trials within a given condition.

In order to investigate S1 dynamics during long-term adaptive behavior, GEVI data were acquired

and analyzed across different training stages. Figure 4c shows neuronal stimulus response as a

function of stimulus amplitude derived from the $\Delta F/F_0$ measurement in S1 of three mice that were

challenged with a minimum of four switches. The functions are represented by a curve fitted to

the average data (see methods), and separately shown for each high range and low range

condition. Comparing the S1 response curves across different training stages revealed an

interesting pattern; at the first switch, high- and low range curves are similar despite differences

in the corresponding stimulus distributions. However, high- and low range curves progressively

separate as a function of switches and training experience. Since the fitting procedure is

constrained by data collected across different stimulus ranges, we focused our analysis on the

"overlap" range which is defined as the stimulus range common to both high- and low range

datasets (A = [2, 4, 8] degrees, dashed vertical grey lines). This range was then used to calculate

the area under the curve (AUC) for each condition, as a way to compare the two curves (see

methods). Figure 4d shows the difference between high- and low range S1 responses, calculated

as the difference in AUCs of the low and high range response curves ($\Delta AUC\ (low - high)$). S1

responses adapt progressively and systematically as a function of training experience. To

determine whether this neuronal effect was dependent upon the animals' choice, we conducted

a separate analysis by tracking the stimulus evoked S1 signal across different training stages,

separately for HIT and MISS trials (Suppl. Fig. 4). The results from the parsed trials for either HITs

or MISSes (Suppl. Fig. 4a-b) are very similar to the combined trials (Suppl. Fig. 4c), confirming

that the emerging difference in S1 activity is not biased by the choice element.

S1 contribution to adaptive behavior

The core elements offered through the behavioral and neurometric measures analyzed thus far

provide somewhat limited "slices" of the underlying phenomena. We thus sought to develop a

more comprehensive analytic, cascade framework that enables us to evaluate the relative

contributions from activity in S1 and downstream of S1 in predicting the observed behavior across

the full range of stimuli. The framework establishes a link between the sensory input distribution

s_i, the S1 response function $G(s_i)$, and the psychometric curve $P(GO|s_i)$ (Fig. 5). The function $f(.)$

represents a downstream process, matching the evoked fluorescence to the lick response, given

a particular stimulus, and is referred to as the "linker" function, as it links S1 activity to behavior.

This framework can then be used to describe the changes underlying the adaptive behavior and

assess the relative roles of the two stages, S1 and downstream (see methods for details). Animals

that were challenged with a single switch were defined as "switch-naive"; animals that were

challenged with a minimum of 4 switches were defined as "switch-experienced". Figure 5a and b

show the different elements of the cascade framework applied to data from multiple training

stages, allowing us to separately investigate switch-naive (Fig. 5a) and switch-experienced (Fig.

5b) transformations.

S1 response curves are shown separately for the high- and low range conditions (Fig. 5 a, b; left

panels), revealing a strong separation at the switch-experienced level. In contrast, the behavior

data (Fig. 5 a, b; right panels) are consistently modulated by changes in the stimulus distribution

at both training stages. This asymmetric relationship of S1 activity and the behavioral output has

an impact on the downstream linker function (Fig. 5a, b; middle panel). At the switch-naive training

stage (Fig. 5a; middle panel), high and low range linker functions appear different, indicating that

the associated neuronal processes that are predictive of behavior, are not S1 based. In contrast,

at the switch-experienced stage (Fig. 5b; middle panel) high- and low range linker functions are

relatively congruent, transforming the modulated S1 response functions directly into an adapted

behavioral response. All predictions from combining S1 response functions and linker functions

(Fig. 5a, b; right panel, dashed curves) match well with the measured psychometric curves (Fig.

5a, b; right panel, solid curves), with minor differences due to the fitting accuracy when generating

both S1 and linker functions. To capture how much of the observed changes in behavior is

explained solely by changes in S1, we used a null test approach assuming a fixed linker function

(e.g. high range only) and predicting the change in psychometric curve (e.g. low range) based

only on changes in S1 response (e.g. from high to low range). The behavioral adaptation was

estimated by subtracting the areas under the curve of the low- and high range psychometric model

predictions, again as a way of comparing the two curves, and the fraction explained by S1 was

then estimated by comparing this difference with the null test results (see methods for details).

Figure 5c shows the relative explanatory power summarized for mice at the switch-naive versus

switch-experienced training stages and separately for the overlap range (left) and full range

(right). The results of both analyses reveal that changes in S1 activity cannot explain behavioral

adaptation at the switch-naive stage (0 %), whereas they can explain a large proportion (85 %) at

the switch-experienced level. The primary behavioral adaptation occurs in regions outside of S1.

However, as animals gain experience in a changing sensory environment, adaptive responses

emerge in S1.

In addition to a changing environment, switch-experience also incorporates increased exposure

to both stimulus conditions and can be described as session-experience. We applied the cascade

model to each animal individually to test for the effect of session experience vs switch experience.

Figure 5d depicts the relative explanatory power of S1 versus downstream, quantified for all mice

at each transition. The first group of mice (M04-M06) experienced only two switches but a higher

number of sessions within a condition (n = 10 sessions), whereas the second group (M07-M09)

experienced four to five switches but a smaller number of sessions within a condition (n = 5

sessions). Despite the variability across mice, the S1 explanatory power increased collectively

with training, again suggesting that the adaptive changes are increasingly emerging in S1 with

experience to the point where a large percentage of the behavioral changes are reflected in the

sensory evoked S1 responses. Figure 5e presents the explanatory power of S1 as a function of

the number of sessions across animals, showing a progression across sessions, reaching 50 %

explanatory power between 20-30 sessions. Interestingly, mice challenged with a smaller number

of sessions per block required more switches for the explanatory power to emerge from S1,

suggesting the number of switches and number of sessions, are both important determinants of

this experience dependent effect.

To causally test whether context-dependent representation in S1 ultimately contributes to

behavioral adaptation, we chronically inactivated S1 in a subgroup of trained mice with the

neurotoxin ibotenic acid, where GEVI activity was used to facilitate targeted acid injections (Fig.

6). Both, simple detection and adaptation to changing conditions were tested in healthy (n = 6),

lesioned (n = 4), and sham lesioned (n = 2) mice, and compared either within the same group or

across groups. From the four lesioned mice, two were switch-naive and two were switch-

experienced. Figure 6a shows a wide-field image of a coronal mouse brain section (100 µm, right

hemisphere) with cytochrome oxidase staining, showing an example S1 lesion. Figure 6b shows

frames of cortical fluorescence activity from an example mouse working on the detection task pre-

and post-lesion with a clear absence of evoked cortical fluorescence after the lesion (bottom row).

Interestingly, mice could still detect stimuli quite well, reaching a level of performance comparable

to pre-lesion levels after one day of recovery, as indicated by the psychometric threshold (Fig.

6c). These results agree with recent work showing that S1 is not required for simple detection[12].

However, if challenged with changing stimulus conditions (high- versus low range), the adaptive

behavior was clearly compromised following the lesion as shown by the psychometric curves and

corresponding psychometric thresholds in Figure 6d. We observed a significant decrease in

switch-behavioral adaptation (Δ threshold) in all animals (Fig. 6e), demonstrating that S1 is

necessary for behavioral adaptation. There were no differences in animals that were lesioned at

the "switch-naive" stage versus "switch-experienced" stage.

Our results suggest cumulatively that the S1 cortical region is not necessary for simple stimulus

detection; rather, it plays an important function in experience dependent stimulus adaptation. This

highly conserved brain region is likely to be important for an animals' survival strategies in a

dynamically challenging sensory environment.

DISCUSSION

In this study, we have investigated learning and experience dependent behavior in the mouse

somatosensory system. Our findings provide evidence that activity in the highly conserved

primary somatosensory cortex can be remarkably dynamic in support of flexible sensory

processing and experience dependent behavioral adaptation.

We present the following novel aspects. First, S1 population activity reflects a reliable signal to

noise relationship that is driven by the sensory input and does not change during the basic

learning process. Second, mice can modify detection behavior in a way as to maintain reward in

the face of changing statistical properties of the stimulus. Third, S1 activity is highly dynamic in

the face of a changing sensory environment, predicting behavioral adaptation as individuals gain

experience. Fourth, S1 is required for this adaptive behavior.

Learning occurs when an individual forms an association based on a new stimulus or context.

This process provides obvious benefits such as flexible hunting, optimal foraging, and social

communication, especially in environments that tend to change frequently and unpredictably.

There is no doubt that associative learning can occur in animals without cortex, including all

classes of vertebrates[34] and a large number of invertebrate species[35]. Further, several studies

elucidating the role of S1 with classical or operant conditioning have found that chronic lesioning

of S1 does not affect basic detection[12,36], a finding we corroborate here. It is simultaneously the

case that other studies demonstrate the opposite effect with acute inactivation of S1[37,38],

suggesting that S1 is normally central to this behavior. Although this is a complex issue, these

studies together seem to suggest that S1 is nominally used for these behaviors, but that given

sufficient time for remapping/learning, other pathways are sufficient, and thus S1 may be viewed

as involved in but not critical for this simple behavior. A central finding of the current study,

however, is that S1 is important for the experience-dependent adaptive behavior *for the exact*

same task and does not recover following lesioning of S1, pointing to a critical role for S1 in

context dependent, adaptive strategies in a changing environment that could underlie a range of

behaviors in more natural settings.

The general behavioral question posed here relates to how the animal responds to changes in

the sensory environment. The behavioral paradigm was designed as a highly simplified but

carefully controlled manipulation of the statistical distribution of the magnitude of whisker

deflections experienced by the animal. Aside from matching amplitudes and velocities of the

whisker movement that have been described in a range of studies[20,21,39–41], the current study does

not attempt to place this in the context of the natural sensory environment for the animal, and the

passive stimulus paradigm does not speak to active sensing that relies on the animal's own

movement in acquiring sensory information. Importantly, although centered around a simple

switching in the stimulus statistics, these switches were implemented with multiple changes

occurring with alternating conditions, showing that S1 sensitivity closely followed the direction of

changes in a way that increasingly explained the behavioral adaptation (Fig. 5), suggesting that

the findings here likely generalize to more complex scenarios in the dynamic landscape of the

natural sensory environment. The relative role of S1 in the adaptive behavior was found to be

strongly experience dependent, where experience is likely to be a complex function of time and

exposure to specific aspects of the environment (here number of sessions), as well as the nature,

degree, and frequency of change in the environment (here number of switches), which remains

to be explored in more detail in naturalistic settings. Overall, the behavior here reveals an active

strategy by the animal that robustly maintains reward in the face of a changing environment. The

behavior is reversible and independent of the direction of switching, and the animals do not just

remain at a higher level of behavioral performance once achieving this or simply work longer

sessions to accumulate the same amount of reward. This demonstrates that the phenomenon

here is part of an active strategy that involves a cognitive cost to the effort, be it effort on a trial-

by-trial basis or the cumulative effort across a session. The changes in strategy we observed

were purely reflected in performance, and not in any changes in peripheral sensing such as

whisker position or movement given the high degree of control exerted by our passive stimulation

paradigm. However, these elements likely play a key role in the overall behavioral strategy in

more ethological behaviors. How the behavior, and corresponding sensory representations,

change in more complex, naturalistic settings, and at a finer time resolution, are important, and

need further investigation in future studies.

While the spatial and temporal resolution of the GEVI imaging here does seem to capture the

appropriate scale for the simple detection task, it targets the aggregate activity of large neuronal

pools restricted to approximately cortical layer 2/3, and thus does not capture the details of

individual neurons of specific cell sub-type across layers that conventional electrophysiological

approaches would provide. However, given the long literature implicating the role of inhibitory

interneurons in functional plasticity in cortex even on short time scales[42,43], we would predict that

the adaptive changes in S1 described here would correspond to a differential change in inhibitory

drive, and that cortical layer 4 and deeper cortical layers would exhibit less adaptive properties

as compared to more superficial layers of cortex, due to the differential dependence upon direct

thalamic drive reported across cortical layers[44]. How the behavioral phenomenon we describe

here is ultimately driven by population activity across the cortical network is left to more detailed

electrophysiological studies in the future. Nevertheless, our finding of context dependent adaptive

responses in S1 is surprising as it suggests that reward based choice signals might shift across

the cortical network and can appear and influence sensory representation in S1 once an individual

has successfully adapted its behavioral strategy. In this context it is important to note that the

behavioral adaptation itself is similar at different experience levels, showing consistent changes

in performance already before the adaptive response even appears in S1 (switch-naive

performer). This is important, as the change in S1 we observe through GEVI imaging averaged

across trials is therefore not simply reflecting a difference in performance/choice (e.g. ratio of hits

to misses) across the levels of experience, but instead a true change in the S1 response with

experience in the context of this task, further reinforced by a direct analysis of the choice signaling

(Suppl. Fig. 4). The findings here are thus not in conflict with the previous reports of choice

signaling in S1, but instead represent a context dependent element related to more complex

behaviors that are likely part of a larger framework of decision making that can only be precisely

identified through causal manipulations in future studies.

So, what are the downstream targets that drive early behavior and ultimately change the stimulus

representation of primary sensory areas? A recent study found that the perirhinal cortex, the last

station in the medial-temporal loop projecting to S1, acts as a gate for the enhancement of cortico-

cortical inputs, which are necessary for basic detection learning[45]. Another recent study probing

flexible decision making in the somatosensory pathway of mice found that orbitofrontal cortex

dynamically interacts with S1 triggering plasticity based on value signals[15]. In addition, other

important reward based choice signals have been reported to influence neuronal signaling

throughout cortex[46]. Furthermore, studies of visual attention in primates have distinguished

changes in neuronal sensitivity from an observer's response criterion in extra striate cortex,

superior colliculus, as well as lateral prefrontal cortex[47–49]. Hence, there is a large number of

downstream candidates potentially modulating the sensory signal stream. Admittedly, in the

current study we did not record in any other area outside S1, but the effects described by those

other studies collectively point to the interesting idea that the neuronal signal transfer identified

by our study could be a common principle across neocortex with "earlier" stages emerging with

experience. Moreover, we propose that this shift in cognitive signaling could even affect

subcortical structures such as primary thalamus, which we have recently shown to exhibit

cognitive signatures in highly trained animals[25].

Arguably, our current study was not specifically designed to identify attention as a driver for

adaptive behavior, even though the measured changes in performance are indicative of a

somewhat conscious change in the subjects' behavioral strategy. Other effects related to the

subjects' thirst, satiation, or arousal level within this behavioral framework can further be

excluded, as we have shown previously[26]. However, although speculative, the experience

dependent change in S1 sensitivity and the corresponding change in downstream criterion may

be signatures of a higher order cognitive process that could be described as an "attentional

spotlight" or gain control[50] moving down the cortical hierarchy. Such a brain-wide dynamic process

would facilitate selective and efficient cognitive processing as an individual is adapting to changes

in the environment. This transfer could explain some of the disparities in the literature and

therefore needs to be further investigated in future studies.

The primary sensory cortex is classically described as the origin of sensory signals in response

to stimuli. We have described here a novel role for S1 in long term adaptive behavior. We propose

that this highly conserved region has a dual function that is temporally regulated, relaying sensory

signals in an acute reactionary fashion vs processing and incorporating dynamic signals into an

adaptive behavioral response longitudinally. This suggests primary sensory cortex may not only

be involved in fundamental sensory representations, but also long term adaptive strategies which

could be necessary for survival.

Data availability

The data that support the findings of this study are available in "Dryad" with the identifier
https://doi.org/10.5061/dryad.h18931zmm.

Code availability

All computer code are available in "Dryad" with the identifier
https://doi.org/10.5061/dryad.h18931zmm.

References

1. Heekeren, H. R., Marrett, S. & Ungerleider, L. G. The neural systems that mediate human perceptual decision making. *Nature Reviews Neuroscience* **9,** 467–479 (2008).

2. Shadlen, M. N., Britten, K. H., Newsome, W. T. & Movshon, J. A. A computational analysis of the relationship between neuronal and behavioral responses to visual motion. *J. Neurosci.* **16,** 1486–1510 (1996).

3. De Lafuente, V. & Romo, R. Neuronal correlates of subjective sensory experience. *Nat. Neurosci.* **8,** 1698–1703 (2005).

4. Glickfeld, L. L., Histed, M. H. & Maunsell, J. H. R. Mouse primary visual cortex is used to detect both orientation and contrast changes. *J. Neurosci.* **33,** 19416–19422 (2013).

5. Petruno, S. K., Clark, R. E. & Reinagel, P. Evidence That Primary Visual Cortex Is Required for Image, Orientation, and Motion Discrimination by Rats. *PLoS One* **8,** (2013).

6. Lashley, K. S. The mechanism of vision IV. The cerebral areas necessary for pattern vision in the rat. *J. Comp. Neurol.* **53,** 419–478 (1931).

7. Kato, H. K., Gillet, S. N. & Isaacson, J. S. Flexible Sensory Representations in Auditory Cortex Driven by Behavioral Relevance. *Neuron* **88,** 1027–1039 (2015).

8. Kelly, J. B. & Glazier, S. J. Auditory cortex lesions and discrimination of spatial location by the rat. *Brain Res.* **145,** 315–321 (1978).

9. Jaramillo, S., Borges, K. & Zador, A. M. Auditory Thalamus and Auditory Cortex Are Equally Modulated by Context during Flexible Categorization of Sounds. *J. Neurosci.* **34,** 5291–5301 (2014).

10. Talwar, S. K., Musial, P. G. & Gerstein, G. L. Role of mammalian auditory cortex in the perception of elementary sound properties. *J. Neurophysiol.* **85,** 2350–2358 (2001).

11. Stüttgen, M. C. & Schwarz, C. Barrel cortex: What is it good for? *Neuroscience* **368,** 3–16 (2018).

12. Hong, Y. K., Lacefield, C. O., Rodgers, C. C. & Bruno, R. M. Sensation, movement and

learning in the absence of barrel cortex. *Nature* **561,** 542–546 (2018).

13.   Kim, J., Erskine, A., Cheung, J. A. & Hires, S. A. Behavioral and Neural Bases of Tactile
Shape Discrimination Learning in Head-Fixed Mice. *Neuron* (2020).
doi:10.1016/j.neuron.2020.09.012

14.   Lee, C. C. Y., Kheradpezhouh, E., Diamond, M. E. & Arabzadeh, E. State-Dependent
Changes in Perception and Coding in the Mouse Somatosensory Cortex. *Cell Rep.* **32,**
(2020).

15.   Banerjee, A. *et al.* Value-guided remapping of sensory cortex by lateral orbitofrontal
cortex. *Nature* **585,** 245–250 (2020).

16.   Borden, P. Y. *et al.* Genetically expressed voltage sensor ArcLight for imaging large scale
cortical activity in the anesthetized and awake mouse (erratum). *Neurophotonics* **4,**
39801 (2017).

17.   Jin, L. *et al.* Single Action Potentials and Subthreshold Electrical Events Imaged in
Neurons with a Fluorescent Protein Voltage Probe. *Neuron* **75,** 779–785 (2012).

18.   Schwarcz, R. *et al.* Ibotenic acid-induced neuronal degeneration: A morphological and
neurochemical study. *Exp. Brain Res.* **37,** 199–216 (1979).

19.   Chagas, A. M. *et al.* Functional analysis of ultra high information rates conveyed by rat
vibrissal primary afferents. *Front. Neural Circuits* **7,** 190 (2013).

20.   Ritt, J. T., Andermann, M. L. & Moore, C. I. Embodied Information Processing: Vibrissa
Mechanics and Texture Features Shape Micromotions in Actively Sensing Rats. *Neuron*
57, 599–613 (2008).

21.   Wolfe, J. *et al.* Texture Coding in the Rat Whisker System: Slip-Stick Versus Differential
Resonance. *PLoS Biol.* **6,** e215 (2008).

22.   Stüttgen, M. C., Rüter, J. & Schwarz, C. Two psychophysical channels of whisker
deflection in rats align with two neuronal classes of primary afferents. *J. Neurosci.* **26,**
7933–41 (2006).

23.   Ollerenshaw, D. R. *et al.* Detection of tactile inputs in the rat vibrissa pathway. *J.*
Neurophysiol. **108,** 479–490 (2012).

24.   Ollerenshaw, D. R., Zheng, H. J. V, Millard, D. C., Wang, Q. & Stanley, G. B. The
Adaptive Trade-Off between Detection and Discrimination in Cortical Representations
and Behavior. *Neuron* **81,** 1152–1164 (2014).

25.   Waiblinger, C., Whitmire, C. J., Sederberg, A., Stanley, G. B. & Schwarz, C. Primary
Tactile Thalamus Spiking Reflects Cognitive Signals. *J. Neurosci.* **38,** 4870–4885 (2018).

26.   Waiblinger, C., Wu, C. M., Bolus, M. F., Borden, P. Y. & Stanley, G. B. Stimulus context
and reward contingency induce behavioral adaptation in a rodent tactile detection task. *J.*
Neurosci. **39,** 1088–1099 (2019).

27.   Wichmann, F. A. & Hill, N. J. The psychometric function: I. Fitting, sampling, and
goodness of fit. *Percept. Psychophys.* **63,** 1293–1313 (2001).

28. Wichmann, F. A. & Hill, N. J. The psychometric function: II. Bootstrap-based confidence
intervals and sampling. *Percept. Psychophys.* **63,** 1314–1329 (2001).

29. Frund, I., Haenel, N. V & Wichmann, F. A. Inference for psychometric functions in the
presence of nonstationary behavior. *J. Vis.* **11,** 16 (2011).

30. Cook, E. P. & Maunsell, J. H. R. Dynamics of neuronal responses in macaque MT and
VIP during motion detection. *Nat. Neurosci.* **5,** 985–994 (2002).

31. Waiblinger, C., Whitmire, C. J., Sederberg, A., Stanley, G. B. & Schwarz, C. Primary
tactile thalamus spiking reflects cognitive signals. *J. Neurosci.* **38,** 4870–4885 (2018).

32. Green, D. M. & Swets, J. A. Signal detection theory and psychophysics. *Society* **1,** 521
(1966).

33. Luo, T. Z. & Maunsell, J. H. R. Attentional Changes in Either Criterion or Sensitivity Are
Associated with Robust Modulations in Lateral Prefrontal Cortex. *Neuron* **97,** 1382–
1393.e7 (2018).

34. Jerison, H. J. & Macphail, E. M. Brain and Intelligence in Vertebrates. *Am. J. Psychol.* **97,**
138 (1984).

35. Giurfa, M. Learning and cognition in insects. *Wiley Interdisciplinary Reviews: Cognitive
Science* **6,** 383–395 (2015).

36. Hutson, K. A. & Masterton, R. B. The sensory contribution of a single vibrissa's cortical
barrel. *J. Neurophysiol.* **56,** 1196–1223 (1986).

37. Miyashita, T. & Feldman, D. E. Behavioral detection of passive whisker stimuli requires
somatosensory cortex. *Cereb. Cortex* **23,** 1655–1662 (2013).

38. Guo, Z. V *et al.* Flow of cortical activity underlying a tactile decision in mice. *Neuron* **81,**
179–194 (2014).

39. Waiblinger, C., Brugger, D., Whitmire, C. J., Stanley, G. B. & Schwarz, C. Support for the
slip hypothesis from whisker-related tactile perception of rats in a noisy environment.
Front. Integr. Neurosci. **9,** (2015).

40. Hartmann, M. Vibrissa mechanical properties. *Scholarpedia* **10,** 6636 (2015).

41. O'Connor, D. H. *et al.* Vibrissa-based object localization in head-fixed mice. *J. Neurosci.*
30, 1947–1967 (2010).

42. Carcea, I. & Froemke, R. C. in *Progress in Brain Research* **207,** 65–90 (2013).

43. Froemke, R. C., Merzenich, M. M. & Schreiner, C. E. A synaptic memory trace for cortical
receptive field plasticity. *Nature* **450,** 425–429 (2007).

44. Radnikow, G. & Feldmeyer, D. Layer- and cell type-specific modulation of excitatory
neuronal activity in the neocortex. *Frontiers in Neuroanatomy* **12,** (2018).

45. Doron, G. *et al.* Perirhinal input to neocortical layer 1 controls learning. *Science (80-.).*
370, (2020).

46. Gold, J. I. & Shadlen, M. N. The neural basis of decision making. *Annual Review of*
Neuroscience **30,** 535–574 (2007).

47. Luo, T. Z. & Maunsell, J. H. R. Attentional Changes in Either Criterion or Sensitivity Are
Associated with Robust Modulations in Lateral Prefrontal Cortex. *Neuron* **97,** 1382–
1393.e7 (2018).

48. Crapse, T. B., Lau, H. & Basso, M. A. A Role for the Superior Colliculus in Decision
Criteria. *Neuron* **97,** 181–194.e6 (2018).

49. Luo, T. Z. & Maunsell, J. H. R. Neuronal Modulations in Visual Cortex Are Associated
with Only One of Multiple Components of Attention. *Neuron* **86,** 1182–1188 (2015).

50. Lee, C. C. Y., Diamond, M. E. & Arabzadeh, E. Sensory Prioritization in Rats: Behavioral
Performance and Neuronal Correlates. *J. Neurosci.* **36,** 3243–3253 (2016).

Acknowledgements

C. W. was supported by a fellowship from the Deutsche Forschungsgemeinschaft (GZ: WA 3862/1-1) and NIH Brain Grants R01NS104928 and U01NS094302. P.Y.B. was supported by an NIH NRSA pre-doctoral fellowship F31NS09869. M. E. D was supported by NIH Brain Grant R01NS104928. G. B. S. was supported by NIH Brain Grants R01NS104928 and U01NS094302. We thank Cornelius Schwarz and April Reedy for helpful feedback on the manuscript. We also acknowledge April Reedy with the Emory University Integrated Cellular Imaging Core for assistance with histological imaging.

Author contributions

C.W. and G.B.S conceived the project. C.W., P.Y.B. and M.E.D. designed the study. C.W. and M.E.D. carried out all experiments. C.W., P.Y.B. and M.E.D. analyzed the data, C.W. and G.B.S. wrote the manuscript with comments from P.Y.B. and M.E.D.

Competing interests

The authors declare no competing interests.

Figure 1. Imaging and behavior methods. a, Top: Schematic of the imaging system. The GEVI "ArcLight" is expressed in superficial layers of S1. Bottom: Schematic of the behavior setup. **b,** Confocal image of a coronal mouse brain section (100 μm, right hemisphere) showing the characteristic spread of ArcLight in S1. Blue: DAPI staining, Green: Arclight fluorescence. **c,** Top view of the same hemisphere in vivo showing ArcLight fluorescence in S1. **d,** Images of the same hemisphere in a trained mouse with ArcLight fluorescence color coded. Punctate stimuli (16 degree whisker angle) or catch trials (0 degree) were presented. Frames were captured at 200 Hz and depicted from stimulus onset onward. Each frame is normalized to the frame at stimulus delivery ($\Delta F/F_0 = F-F_0/F_0$). Shown are fluorescence responses averaged over an example imaging session (n=24 trials/condition). A region of interest (red box, 434 x 434 μm) is centered at the peak fluorescence to extract and average voltage traces for further analysis. Scale bar: 1 mm. **e,** Go/No-Go detection task. A punctate stimulus (10 ms) has to be detected by the mouse with an indicator lick to receive reward. Reward is only delivered in hit trials. Impulsive licks in a 2 s period before trial onset are mildly punished by a time-out triggering a new inter-trial-interval (4-10 s, gray arrow). **f,** Example traces of Go and No-Go trials (n=284 each). Top: Readouts from the galvo sensor (16 degree stimulation or 0 degree catch trial). Middle: Lick response histograms from a trained mouse. Bottom: Average voltage response from the region of interest of S1. **g,** Different task versions under investigation. 1. Basic learning: Learning of the Go/No-Go task. 2. Adaptive behavior: Once animals have learned the basic task, they are challenged with multiple stimulus amplitudes and changes in the statistics of the stimulus distribution.

Figure 2. Basic Learning and sensory signals in S1. a, Learning curve for 3 mice trained on the basic Go/No-Go detection task with a weak stimulus (4 degree). Performance is expressed as the probability of a hit (lick after stimulus) and false alarm (lick after no stimulus) for a given session. Individual mice are represented by symbols, and the mean by bold lines. **b,** Fluorescent activity in S1 from an example mouse during learning of the task. Shown are frames at the peak response to the 4 degree stimulus, catch trials are shown below. Scale bar: 1 mm. **c,** Dprime metrics for both behavioral and neuronal data during learning of the task. The d'_{behav} was calculated from the observed hit rate and false alarm rate for a given training session. The d'_{neuro} was calculated by comparing single trial distributions of all evoked signal peaks (maximum % $\Delta F/F_0$ within 100 ms post stimulus) with the corresponding noise distributions when no stimulus was present. Shown are means across mice (n=3) for a given session, errorbars represent bootstrapped estimates of 95% confidence limits (nBoot=1000 repetitions). The dotted lines separate performance into "detect-naive" ($d'<0.8$), and "detect-experienced" ($d'>1.5$). The right panel shows the same data separated for individuals (symbols) before and after learning. Bars represent means across mice, errorbars represent SD (n=3). * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; 'n.s.' not significant, two-sided Wilcoxon rank-sum test.

Figure 3. Adaptive behavior and S1 responses. a, Manipulation of stimulus distribution range. Every stimulus and catch trial (not shown) is presented with equal probability (p=0.2). The design involves amplitudes common to both high range (magenta) and low range (green) conditions. **b,** Psychometric curves and response thresholds for an example animal working on both conditions. Each dot corresponds to response probabilities from a single session. Solid curves are logistic fits to the average data (n=10-11 sessions). Dotted line is a hypothetical curve assuming no change in performance (H_0). Dashed line is a hypothetical curve assuming a change in performance to maintain reward (H_1) when switching from high range to low range stimulus distributions. Response thresholds are shown as vertical lines with 95% confidence limits. Inset. Response thresholds of all mice (grey symbols). Bars represent means across mice with SD (n=4). **c,** Number of rewards (correct trials) accumulated by the same animal from b. Each line corresponds to one session. Inset. Average total reward number per session for each mouse. The average number of trials is shown on top. Figure conventions are the same as in b. **d,** Frames of evoked cortical fluorescence activity from two example sessions, one with a high range stimulus distribution (top) and the other with a low range stimulus distribution (bottom). The frames are aligned for amplitudes common to both datasets. Data with a deflection angle of 8 degree was chosen for further analysis (outlined box). **e,**Temporal fluorescent signal in response to 8 degree stimulation, extracted from the region of interest and averaged across sessions and mice. PSTHs of behavioral lick responses are shown on top. Median reaction times (first lick post stimulus) are shown as arrows, 25-75 percentiles as horizontal lines (n=829-831 trials). **f,** Observer model based on signal detection theory. Behavioral adaptation can either be induced by changes in sensitivity (d') intrinsic to S1, changes in decision criterion by a downstream observer, or both. **g,** Top: Distributions of evoked trials (signal) and catch trials (noise) computed separately for the high (HR) and low range (LR) condition. Dashed lines indicate mean $\Delta F/F_0$, black arrows indicate dprime values. Blue lines indicate criterion values as computed from ROC curves. Bottom: Criterion and dprime metrics (neuronal and behavioral). Shown are bootstrapped estimates of means and 95% confidence limits (n=38 (LR) or n=41 (HR) sessions, n=4 mice, nBoot=1000 repetitions). * P < 0.05; ** P < 0.01; *** P < 0.001; 'n.s.' not significant, two-sided Wilcoxon rank-sum test or Kruskal-Wallis test.

Figure 4. Timescale of behavioral adaptation and related S1 dynamics. a, Long-term behavioral adaptation. Psychometric thresholds of individual mice (n=6) undergoing repetitive task changes starting with either the high range or low range condition. Symbols represent mean thresholds across sessions (n=5-11 per condition), errorbars represent 95% confidence limits. Note, conditions are changed in an alternating fashion to test reversibility of adaptation. **b.** Short-term behavioral adaptation at the transition (dashed lines) from high to low (left) and low to high (right). Symbols represent thresholds from each mouse and each session. Bold lines represent thresholds averaged across mice before and after the transition. Horizontal lines represent thresholds averaged across mice and sessions for a given condition. Note, the data is normalized with regard to the transition, with session 0 representing the last session of the preceding condition. **c,** S1 dynamics over multiple task changes (long-term). Neuronal Stimulus response curves derived from the $\Delta F/F_0$ measurement in S1. The curves are weibull fits to the data of n=3 mice undegoing repetitive changes. The "overlap range" is defined as the stimulus range common to both high range and low range datasets (2-8 degree) which is used to calculate the area under the curve (AUC). The hypothetical part (above 8 degree for the low range) is represented as dashed curve. **d,** Difference between high range and low range S1 responses, calculated as the difference in AUC space (AUC_{Low}- AUC_{High}) at different training stages. Bars represent means across mice with SD (not all mice underwent all switches; switch 1-2, n=6 mice, switch >4, n=3 mice). * P < 0.05; ** P < 0.01; *** P < 0.001; 'n.s.' not significant, Kruskal-Wallis test.

Figure 5. S1 contribution to long-term behavioral adaptation. a, Model incorporating the sensory input distribution S_i, the neuronal response function $G(s_i)$ derived from the GEVI signal in S1 and the behavioral readout $P(GO|s_i)$. The function $f(.)$ represents a downstream process, matching the evoked fluorescence to the lick response. Model predictions and measured data at the "switch-naive" training stage, when subjects are challenged with a change in stimulus satistics the first time. **b.** Model predictions and measured data at the "switch-experienced" training stage, when subjects have undergone at least 4 switches. **c,** Relative explanatory power in S1 of mice undergoing multiple changes. Model results are shown separately for the overlap range (left) and the full range (right). Bars represent means across mice with SD (n=3). **d,** Pie plots depict the fraction explained by S1 versus downstream quantified for the behavior of mice undergoing different number of switches and different number of sessions per condition (M04-M06, n=2 switches, n=10-11 sessions, M07-M09, n=4-5 switches, n=5 sessions). **e,** Timescale of S1 explanatory power. The fraction explained by S1 is plotted as a function of the number of sessions. Solid curve is a logistic fit to all data.

Figure 6. The effect of S1 inactivation on behavior. a, Wide-field image of a coronal mouse brain section (100 μm, right hemisphere) with cytochrome oxidase staining, showing a S1 lesion upon acid injection (0.3 μL ibotenic acid 10 mg/mL). **b,** Frames of cortical fluorescence activity from an example mouse before and after the lesion. Top. Example session with S1 intact, showing GEVI activity patterns in response to different stimulus amplitudes (high range, n=25 trials/stimulus). Bottom. Example session of the same mouse after the lesion (n=18 trials/stimulus). Note, the GEVI activity pattern was used for targeted acid l injections. **c,** Psychometric curves and response thresholds for an example mouse detecting high range stimuli before and after the lesion. Each dot corresponds to response probabilities from a single session. Solid curves are logistic fits. Note, pre-lesion data is shown for several sessions (n=5) whereas the post-lesion data is shown seperately for the first (red) and second session (orange) upon lesion. **d,** Psychometric curves and response thresholds for the same mouse challenged with a change in stimulus statistics (high range to low range) upon S1 inactivation. Solid curves are logistic fits to the average data (n=4-5 sessions). Otherwise, figure conventions are the same as in Figure 3. **e,** Behavioral adaptation is calculated as the change in psychometric threshold (Δ threshold) for healthy and lesioned animals. Bars represent means across mice with SD (n=6 intact, n=4 lesioned), dotted line represents mean across sham operated mice (n=2).* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; 'n.s.' not significant, two-sided Wilcoxon rank-sum test.

Supplemental Figure 1. Basic Learning with a strong stimulus. a, Learning curve for 3 mice trained on the basic Go/No-Go detection task with a strong stimulus (16 degree). **b,** Fluorescent activity in S1 from an example mouse during learning. Shown are frames at the peak response to a 16 degree stimulus, catch trials are shown below. Scale bar: 1 mm. **c,** Dprime metrics for both behavioral and neuronal data during learning of the task. Note, mice achieved higher hit rates at the beginning of training and reached successful task acquisition in half the time if compared to mice detecting a weak stimulus (Fig 2). The dotted lines separate performance into "detect-interim" (d'=0.8-1.5), and "detect-experienced" (d'>1.5). The right panel shows the same data separated for individuals (symbols) before and after learning. Otherwise, figure conventions are the same as in Figure 2.

Supplemental Figure 2. The relationship of magnitude and activated area in GEVI imaging.
a, Fluorescence activity in response to different stimuli. Each frame is normalized to the frame at stimulus delivery ($\Delta F/F_0 = F-F_0/F_0$). Shown are images at the response peak. The dotted line represents a slice through the images aligned with the maximum fluorescence. Scale bar: 1 mm.
b, Magnitude (response amplitude $\Delta F/F_0$) versus cortical area extracted from the slice in a. Activity patterns are shown in different shades of grey for different stimulus amplitudes. **c,** Relationship between fluorescence magnitude and width of activated area, both scale with stimulus strength in a highly correlated fashion. The width is derived from a threshold (dashed line in b). Data points correspond to different stimulus amplitudes fitted with a linear regression. Shown are means across mice (n=4) and sessions (n=40).

Supplemental Figure 3. Analysis to test stability of adaptive S1 response. a, Different normalization methods. F_0 was varied by using the mean fluorescence across different time windows before stimulus onset. The $\Delta F/F_0$ measurement was then calculated by $F-F_0/F_0$. Mean fluorescence traces (n=4 mice, 831-870 trials) in response to the high (magenta) and the low range (green) stimulus condition. The grey box depicts the window for calculating F_0. From left to right: F_0=0 ms (frame at stimulus delivery), F_0 from [-100,0] ms, and F_0 from [-200,0] ms. **b,** Single-trial signal-peak and noise distributions from the data in a. Magenta and green numbers are mean fluorescent values in % $\Delta F/F_0$ for the high and low range condition respectively. Black numbers represent dprime metrics. **c,** Dprime and variance for different F_0 calculations. **d,** Receiver operating characteristic (ROC) curves created by shifting the criterion across the $\Delta F/F_0$ signal and noise distributions of the high and low range condition. The downstream criterion can be inferred by comparing the hit rate in ROC space with the average behavioral hit rate (dashed lines).

Supplemental. Figure 4. Choice-related long-term S1 dynamics. a, GEVI peak responses (maximum % $\Delta F/F_0$) for correct choices (hit trials) to 8 degree stimulation. Data is shown over multiple switches for the high- and low range condition. **b,** GEVI responses for incorrect choices (miss trials). **c,** Top: Percentage of hit and miss trials for the high- and low range condition. Bottom: Combined GEVI data (hits+misses) as a function of switches. All data shown are bootstrapped estimates of means and 95% confidence limits (based on single trials from n=6 mice, nBoot=1000 repetitions).