Smoothened and ARL13B are critical in mouse for superior cerebellar peduncle targeting

Sarah K. Suciu[*,†,‡] Alyssa B. Long[†,‡] and Tamara Caspary[†]

[*]Genetics and Molecular Biology Graduate Program, [†]Department of Human Genetics, Emory University, Atlanta, GA 30322, [‡]These authors contributed equally

ORCID ID:

Suciu: 0000-0002-2416-9519

Long: 0000-0002-4467-4213

Caspary: 0000-0002-6579-7589

Summary statement
Joubert syndrome is diagnosed by the hindbrain "molar tooth sign" malformation. Using mouse models, we show loss of the ciliary GTPase ARL13B, mutations in which lead to Joubert syndrome, result in two features of the molar tooth sign: hypoplasia of the cerebellar vermis and inappropriate targeting of the superior cerebellar peduncles. Furthermore, we demonstrate that loss of vertebrate Hedgehog signaling may be the underlying disrupted mechanism as we extend its role in axon guidance to the superior cerebellar peduncles.

Running title (~35 characters, including spaces):

SMO and ARL13B in SCP targeting

Key words/phrases (up to ten):

molar tooth sign, Joubert syndrome, Smo, ARL13B, SCP targeting, cerebellar vermis

Corresponding author:

Tamara Caspary

Department of Human Genetics

615 Michael Street, Suite 301

Atlanta, GA 30322

404-727-9862

tcaspar@emory.edu

Abstract

Patients with the ciliopathy Joubert syndrome present with physical anomalies,

intellectual disability, and a hindbrain malformation described as the "molar tooth sign"

due to its appearance on an MRI. This radiological abnormality results from a

combination of hypoplasia of the cerebellar vermis and inappropriate targeting of the

white matter tracts of the superior cerebellar peduncles. ARL13B is a cilia-enriched

regulatory GTPase established to regulate cell fate, cell proliferation and axon guidance

through vertebrate Hedgehog signaling. In patients, mutations in *ARL13B* cause Joubert

syndrome. In order to understand the etiology of the molar tooth sign, we used mouse

models to investigate the role of ARL13B during cerebellar development. We found

ARL13B regulates superior cerebellar peduncle targeting and these fiber tracts require

Hedgehog signaling for proper guidance. However, in mouse the Joubert-causing R79Q

mutation in ARL13B does not disrupt Hedgehog signaling nor does it impact tract

targeting. We found a small cerebellar vermis in mice lacking ARL13B function but no

cerebellar vermis hypoplasia in mice expressing the Joubert-causing R79Q mutation.

Additionally, mice expressing a cilia-excluded variant of ARL13B that transduces

Hedgehog normally, showed normal tract targeting and vermis width. Taken together,

our data indicate that ARL13B is critical for control of cerebellar vermis width as well as

superior cerebellar peduncle axon guidance, likely via Hedgehog signaling. Thus, our

work highlights the complexity of ARL13B in molar tooth sign etiology.

INTRODUCTION

Joubert Syndrome and Related Disorders (JSRD) are recessive congenital disorders with a variety of symptoms including developmental delay, intellectual disability, abnormal respiratory rhythms, ataxia, oculomotor apraxia, polydactyly, craniofacial defects, retinal dystrophy, nephronophthisis, and hepatic fibrosis (Parisi *et al.* 2007, Bachmann-Gagescu *et al*. 2020). While the exact prevalence of JSRD is not known, published statistics range from 1:80,000 to 1:100,000, but these may be underestimates as suggested by a recent study (Brancati *et al.* 2010, Nuovo *et al.* 2020). The characteristic neuroanatomical feature of JSRD is the molar tooth sign (MTS), which is caused by hypoplasia of the cerebellar vermis and thickened, elongated superior cerebellar peduncles (SCPs) that fail to decussate (Maria *et al.* 1997;Yachnis and Rorke 1999; Poretti *et al.* 2007). However, little is known about the mechanistic etiology of this hindbrain malformation. This is especially significant as several symptoms of JSRD arise from defects in the hindbrain: cerebellar dysfunction commonly causes ataxia and hypotonia, while some patients manifest life-threatening breathing problems (Parisi 2019).

To date, mutations in over 35 genes cause JSRD, and their associated proteins almost always localize to the primary cilium or the centrosome (Parisi 2019). Thus, JSRD is classified as a ciliopathy, a category of human disease stemming from ciliary dysfunction. One of the genes implicated in JSRD is *ARL13B*, which encodes a regulatory GTPase highly enriched in cilia (Cantagrel *et al.* 2008; Bachmann-Gagescu *et al.* 2015; Thomas *et al.* 2015; Shaheen *et al.* 2016; Rafiullah *et al.* 2017). As a GTPase, ARL13B is expected to have multiple effector proteins which interact with

specific ARL13B residues. ARL13B can function as a guanine exchange factor (GEF)

for ARL3, mutations in which also lead to JSRD (Gotthardt *et al.* 2015; Ivanova *et al.*

2017). JSRD-causing mutations in either ARL3 or ARL13B can disrupt their interaction

or ARL13B's GEF activity, consistent with the notion that specific ARL13B function is

affected by JS-causing point mutations (Gotthardt *et al.* 2015; Ivanova *et al.* 2017;

Alkanderi *et al.* 2018). Most JSRD-causing ARL13B mutations cluster within the

protein's GTPase domain, although two are located in the coiled coil domains in the C

terminal half of the protein (Cantagrel *et al.* 2008; Bachmann-Gagescu *et al.* 2015;

Thomas *et al.* 2015; Shaheen *et al.* 2016; Rafiullah *et al.* 2017). ARL13B complexes

with the inositol phosphatase INPP5E, which is also implicated in causing JSRD (Bielas

et al. 2009; Humbert *et al*. 2012). ARL13B is critical for targeting INPP5E to cilia and

JSRD-causing ARL13B mutations disrupt INPP5E ciliary targeting (Humbert *et al.*

2012). INPP5E controls ciliary lipid composition through its phosphatase activity and

most JSRD-causing mutations are within its phosphatase domain (Bielas *et al.* 2009;

Chavez *et al.* 2015; Garcia-Gonzalo *et al.* 2015). Other proteins implicated in JSRD also

affect ciliary targeting with many functioning at the transition zone, supporting the notion

that abnormal ciliary traffic leading to defective signaling underlies JSRD (Arts *et al.*

2007; Delous *et al.* 2007; Garcia-Gonzalo *et al.* 2011; Hopp *et al.* 2011; Srour *et al.*

2012; Roberson *et al.* 2015).

The mechanistic connection between the cilia-related proteins implicated in

JSRD and the MTS are elusive in part because distinct biological processes are at play.

Abnormal proliferation may underlie the hypoplastic cerebellar vermis whereas defective

axonal targeting is likely involved in the abnormal SCP tracts. One signaling pathway

potentially linked to both processes is vertebrate Hedgehog (Hh) which relies on cilia

(Huangfu *et al.* 2003). Sonic hedgehog (Shh) is a mitogenic cue that controls

proliferation in the developing cerebellum so its misregulation could underlie the

cerebellar hypoplasia (Dahmane and Ruiz i Altaba 1999; Wechsler-Reya and Scott

1999; Kenney and Rowitch 2000). While the SCP tracts that normally project from the

deep cerebellar nuclei to the contralateral thalamus are guided by unknown signals,

Shh is a known commissural axon guidance cue (Charron *et al.* 2003). JSRD patients

can also display axon guidance defects in decussation of the pyramidal tracts (Yachnis

and Rorke 1999; Poretti *et al.* 1997).

ARL13B and *INPP5E,* encoding ciliary proteins linked to JSRD, are known to

regulate vertebrate Hh signaling. In mouse models, ARL13B loss disrupts cell fate

specification in the neural tube, proliferation of the cerebellar granule precursor cells in

the cerebellum and Shh-directed guidance of commissural axons in the spinal cord

(Caspary *et al.* 2007; Bay *et al.* 2018; Ferent *et al.* 2019). These data support a model

whereby disruption of Shh signaling by ARL13B mutation could provide a single

mechanism underlying the MTS. This model is bolstered by the fact that additional

phenotypes exhibited by JSRD patients, such as craniofacial defects or polydactyly, can

arise from aberrant Hh signaling (Valente *et al.* 2008; Lan and Jiang 2009; Lipinski *et al.*

2010).

As attractive as a Hh-based model for JSRD may be, not all the data support that

JSRD phenotypes result from misregulation of Hh signaling. Some features of JSRD,

such as the renal and liver anomalies, are not clearly due to misregulation of Hh

signaling (Doherty 2009; Breslow *et al.* 2018). Additionally, of over 35 genes implicated

in JSRD, only 22 play some role in Hh pathway regulation (Davey *et al.* 2006; Reiter

and Skarnes 2006; Caspary *et al*. 2007; Vierkotten *et al*. 2007; Huang *et al.* 2009;

Weatherbee *et al*. 2009; Bimonte *et al.* 2011; Dowdle *et al.* 2011; Sang *et al.* 2011; Chih

et al. 2011; Christopher *et al.* 2012; Thomas *et al.* 2012; Abdelhamed *et al*. 2013; Hynes

et al. 2014; Wu *et al.* 2014; Chavez *et al.* 2015; Garcia-Gonzalo *et al.* 2015; Asadollahi

et al. 2018; Frikstad *et al*. 2019; Munoz-Estrada and Ferland 2019). Some of these links

are tenuous. For example, mouse *Arl3* mutants mislocalize the Hh transcription factor,

GLI3, in their cilia yet do not exhibit any of the phenotypes normally displayed by

mutants in the Hh pathway (Schrick *et al.* 2006, Lai *et al.* 2011; Schwarz *et al.* 2017).

Additional signaling pathways are linked to cilia including others known to be important

in cell proliferation and axon guidance. Loss of either of the JSRD-linked genes *Ahi1* or

Cep290 in mouse leads to a small cerebellar vermis due to aberrant Wnt signaling

(Lancaster *et al.* 2011; Ramsbottom *et al.* 2020). Possible JSRD-causing mutations in

Znf423 are associated with defects in Wnt, BMP and retinoic acid signaling (Hata *et al.*

2000, Huang *et al.* 2009, Casoni *et al.* 2020, Deshpande *et al.* 2020). Conditional *Arl13b*

or *Inpp5e* deletion in the SCPs results in their disorganization and thickening through

misregulation of ciliary PI3 kinase and AKT (Guo *et al.* 2019).

Ciliopathies are well established to be genetically complex. JSRD patients with

different ARL13B mutations can display distinct phenotypes, such as obesity in an

individual expressing ARL13B[Y86C] and occipital encephalocele in a patient expressing

ARL13B[R79Q] (Cantagrel *et al.* 2008; Thomas *et al.* 2015). This is further exemplified in

cases of related individuals carrying the same mutation but exhibiting different

phenotypes and even diagnoses. For example, JSRD-causing mutations in TMEM67

(R208X) and TMEM216 (R73H) can also cause the more severe disease Meckel

syndrome (Consugar *et al.* 2007; Otto *et al.* 2009; Valente *et al.* 2010). Understanding

the genetic modifiers and environmental contribution underlying the phenotypic variation

will be key to understanding disease etiology as will understanding when and how

relevant pathways interact. In mouse models of the JSRD- and Meckel Syndrome-linked

gene *Tmem67*, two phenotypic categories emerged: one with cerebellar malformations

resembling JSRD and another with more severe CNS defects reminiscent of Meckel

syndrome (Abdelhamed *et al.* 2013). The two categories correlated with whether cilia

were retained, with the severe Meckel-like phenotype observed in animals lacking cilia.

Furthermore, the two phenotypic groups impacted Hh and Wnt signaling differently,

pointing to both pathways being critical (Abdelhamed *et al.* 2019). Importantly, it is not

simply whether cilia are present, as another JSRD mouse model, *Talpid3*, lacks cilia yet

displays a JSRD-like small cerebellar vermis (Bashford and Subramanian 2019). Thus,

mouse models are incredibly informative yet point to the enormous complexity

underlying the MTS.

Here we investigate the role of ARL13B in relation to Hh signaling in two major

features of the MTS: targeting of the SCPs to the thalamus and hypoplasia of the

cerebellar vermis. We explore these processes using a series of mouse alleles through

which we first define the roles of Hh signaling and ARL13B in SCP projections.

Subsequently, we untangle the role of ARL13B from within and outside of the cilium and

investigate a JS-causing patient allele. Taken together, our data illuminate the roles of

ARL13B in MTS etiology and the complexity in modeling aspects of the MTS in mouse.

MATERIALS & METHODS

Mouse lines

All mice were cared for in accordance with NIH guidelines and Emory

University's Institutional Animal Care and Use Committee (IACUC). Lines used were

Nex-Cre (C3H-HeJ-*Neurod6*$^{tm1(cre)Kan}$) [MGI:2668659], *Brn4-Cre* (C3H/HeJ-*Tg(Pou3f4-*

cre)32Cren) [MGI:2158470], *Smo*^{flox} (C3H/HeJ-*Smo*^{tm2AMC}) [MGI:2176256], *Arl13b*^{flox}

(C3H/HeJ-*Arl13b*^{tm1Tc}) [MGI:4948239], *Arl13b*^{V358A} (C57BL/6J-*Arl13b*^{em1Tc})

[MGI:6256969], and *Arl13b*^{R79Q} (C57BL/6J-*Arl13b*^{em2Tc}) [MGI:6279301]. Note that

Arl13b$^{\Delta}$ is the deletion allele resulting from germline deletion of the conditional *Arl13b*^{flox}

allele. Genotyping was as previously described (Heydemann *et al.* 2001; Goebbels *et*

al. 2006; Nolan-Stevaux *et al.* 2009; Su *et al.* 2012; Gigante *et al.* 2020).

To generate the R79Q mutation in *Arl13b*, a CRISPR gRNA

(ATTATTATGCTGAATCCTA*TGG*; PAM sequence is italicized) targeting exon 3 of the

Arl13b locus along with a 180bp donor oligo (5'-CTCCCACTGTTGGCTTTTCTAAAA-

TTGATCTGAGACAAGGAAAGTTCCAAGTTACCATCTTTGACTTAGGAGGTGGAAAA

AGAATTC<u>A</u>GGG<u>C</u>AT<u>A</u>TGGAAGAATTATTATGCTGAATCCTATGGGGTAATATTTGTT

GTGGATTCCAGTGATGAGGAGAGAATGGAAGAAACAAAGGAGA-3'; underlined

bases are engineered) were designed to generate a G-to-A change creating the R79Q

point mutation as well as A-to-C and T-to-A silent changes to create a *Nde*I restriction

site that could be used for genotyping (Millipore Sigma). The gRNA (50 ng/ul), ssDNA

oligo donor (50 ng/ul) and wild type *Cas9* mRNA (10 ng/ul) were injected into 1-cell

C57BL/6J zygotes and subsequently transplanted at the 2-cell stage into C57BL/6J

pseudopregnant females by the Emory Transgenic and Gene Targeting Core; all

reagents were purchased from Millipore Sigma. Genomic DNA from toes was amplified

via PCR using primers (5'-TCACTTGCAACAGAGCATCC-3') and (5'-

ACAGCTCTGCCCGTGTTTAC-3') located upstream and downstream of the donor oligo

breakpoints; products were sequenced with the forward (first) primer. A single founder

animal heterozygous for both the R79Q mutation and NdeI restriction site was identified

with no additional editing. Subsequent allele-specific genotyping of progeny was

performed on ear punch or yolk sac using the following primers: Fwd-wt primer: 5'-

GGAGGTGGAAAAAGAATaCg-3'; Fwd-mut primer: 5'-

gctctatggctgGGAGGTGGAAAAAGAATTga-3'; Rev primer: 5'-

AGTGCTAAGACACCCGAGGA-3'. PCR bands at 142bp (wild type) and/or 154bp

(mutant) were produced, due to the addition of 12 non-templated bases to the 5' end of

the Fwd-mut primer (lowercase). Note the 3' ends of the two forward primers differed in

the base that codes for the R to Q change (final nucleotide of the primer) and includes a

"wobble" base (lowercase) to provide allele-specific amplification after the first round of

PCR (Gaudet *et al.* 2009). In order to breed away any potential off-target edits, the

founder was backcrossed to C57BL/6J for three generations with at least two

independent meiotic opportunities for recombination in each generation.

Tract tracing injections and analysis

Tract tracing experiments were performed according to a protocol approved by

Emory University's Institutional Animal Care and Use Committee (IACUC). Male and

female mice at postnatal day 90 or older were used for tract tracing experiments. At

least 3 mice of each genotypic group were analyzed in experiments (exact N included in

Figures 1-3, S1). Mice were anesthetized with inhaled isoflurane and maintained under

anesthesia throughout the procedure. Animals were secured in the stereotax, and the

scalp was opened with bregma and lambda aligned to flatskull position. Dorsal thalamus

injections were targeted to the bregma (AP:-0.70, ML:+1.13, DV:-3.28, Angle:0°) and

ventral thalamus injections were targeted using coordinates to the bregma (AP:-0.70,

ML:-3.11, DV:-4.69, Angle:25°). Ventral injection targeting includes a 25° angle to avoid

pulling dye through the dorsal thalamus upon needle removal. Then, a 5 ul Hamilton

microsyringe was lowered to target and target was injected with lysine fixable dextran

tetramethylrhodamine neuroanatomical tracer (fluoro-Ruby, 10,000 MW, ThermoFisher

Scientific D1817). Animals received 0.05 - 0.5 ul injections of 10% dextran

tetramethylrhodamine in sterile phosphate buffered saline (PBS, pH = 7.25) unilaterally

at a rate of 0.1 ul/minute. The 0.5 ul injection volume initially used resulted in high

background (non-specific) fluorescence. Later surgeries were conducted with 0.05 ul of

dye, which resulted in lower background and retention of a strong signal in the DCN.

Seven days post-procedure, mice were perfused with 50mls of PBS followed by

30mls of 4% paraformaldehyde (PFA). Brains were kept in PFA overnight and

subsequently placed in 30% sucrose in 0.1M phosphate buffer (pH 7.3) for

cryoprotection for at least 48 hours. Brains were embedded in Tissue-Tek OCT

compound (Sakura) for coronal cryostat sectioning. Sections were 60 microns thick and

processed through 70% ethanol dehydration and 0.1% sudan black autofluorescence

quencher, rehydrated in PBS and treated with DAPI to stain nuclei prior to fluorescence

imaging. Images were taken on a Lionheart FX automated microscope (Biotek) or at 5x

magnification on a Leica DM6000B microscope (Leica) using SimplePCI imaging

software (Hamamatsu). The Leica images were subsequently stitched together to reveal

the entirety of the brain section in Fiji (Schindelin *et al.* 2012) or Photoshop (Adobe).

Surgical injection sites were assessed to ensure dye was present at the desired

injection site. If the injection was off-target or dye at the injection site was not seen,

samples were removed from analysis. Cerebellar images from injections were evaluated

for DCN staining with assessor blinded to genotype. The number of injections that

resulted in fluorescent DCN for each injection site (dorsal and ventral thalamus) were

compared between mutant and control genotypes using a two-sided Fisher's exact test

(PRISM 8.2.0).

Phenotypic analysis of embryos

Timed matings of mice were performed to generate somite-matched embryos at

embryonic day 10.5 (E10.5). Embryos were dissected in cold PBS and processed for

immunofluorescence staining as previously described (Constable *et al.* 2020). Primary

antibodies used were: mouse anti-Shh (5E1, 1:10), mouse anti-Pax6 (PAX6, 1:100)

(Developmental Studies Hybridoma Bank), rabbit anti-Olig2 (AB9610, 1:500, Millipore

Sigma), and mouse anti-Arl13b (N295B/66, 1:1500, NeuroMab). Multiple sections from

three embryos of each genotype were examined.

Analysis of anatomical measures

Weanling age (P20-P24) male and female mice were sacrificed and brains were

harvested and fixed in 4% PFA overnight at 4°. Ten sex-matched pairs of mouse brains

were collected for each genotype; control animals were either wild type or heterozygous

for the *Arl13b* point mutation or floxed allele (so *Arl13b*$^{R79Q/+}$, *Arl13b*$^{V358A/+}$ or *Arl13b*$^{flox/+}$)

or lacked *Brn4-Cre* (for *Arl13b*$^{Brn4-Cre}$). Brains were imaged on a tilted stage to present a

surface view of the cerebellum, with a standard ruler in frame to confirm scale, using a

dissecting microscope (Leica MZFLIII). Measurements were made in FIJI (Schindelin *et*

al. 2012) with the investigator blind to genotype. Whole cerebellar width was measured

at the widest part of the cerebellum, coinciding with lobule CI or CII. Vermis width was

calculated by measuring the widest part of lobule VII (Deshpande *et al.* 2020).

Hemisphere and vermis heights were measured at the longest rostro-caudal point of the

hemisphere or at the midline of the vermis (see Figure 5). Body and brain weights were

measured on a standard lab scale. For each sex- and age-matched pair, the ratio of

mutant to control measures were calculated and graphed. A one-sample t-test was

performed after transforming the ratio data to a log(2) scale to normalize data

distribution (PRISM 8.2.0). Using the vermis width measures from our control brains

(mean = 2.6mm, SD = 0.18), a power calculation for a one-sided t-test estimated 90%

power to detect a 10% difference with 9-10 paired samples for a nominal alpha=0.05

(https://clincalc.com/stats/samplesize.aspx and

https://www.stat.ubc.ca/~rollin/stats/ssize/n2.html).

Data availability

Mouse lines are available upon request. Supplemental material available at

figshare. Figure S1 contains diagrams illustrating the complete results of the injection

and tract tracing experiments. Figure S2 contains a schematic of the CRISPR strategy.

The authors affirm that all data necessary for confirming the conclusions of the article

are present within the article and figures.

RESULTS

SMO is required for normal SCP projection to the dorsal thalamus

In order to test whether proper projection of SCPs requires Hh signaling, we

compared the SCP tracts in mice in which we deleted the gene encoding the obligate

Hh transducer, *Smoothened* (*Smo*), to control animals. As *Smo* null embryos die during

embryogenesis, we deleted *Smo* specifically in the projection neurons by generating

Nex-Cre;Smo$^{fl/fl}$ mice, which we refer to as *Smo$^{Nex-Cre}$* (Zhang *et al.* 2001; Caspary *et al.*

2002*). Nex-Cre* initiates CRE recombinase expression at E11.5, as the precursor cells

of the deep cerebellar nuclei (DCN) begin to migrate and become specified (Fink *et al.*

2006; Goebbels *et al.* 2006). In the mature cerebellum, the SCPs project rostrally from

the DCN (illustrated in Figure 1A). After entering the midbrain, the SCPs cross the

midline and again turn rostrally to project to two positions in the rostral thalamus: one

tract takes a slight dorsal path and the other tract remains in the same plane; for

simplicity here, we term these projection sites the dorsal and ventral thalamus,

respectively (Bohne *et al.* 2019). To examine the SCP tracts, we used retrograde tract

tracing in which we performed stereotaxic injections of a lipophilic dextran dye into

either the dorsal (Figure 1B) or ventral (Figure 1C) thalamus and allowed the dye to

diffuse through the axons to the associated neuron's cell body (~7 days); we then

sacrificed the animal and examined the cerebellum for evidence of the lipophilic dye

indicating tracing.

We found that both dorsal and ventral thalamus injections resulted in visible

clusters of dye-stained cells in the contralateral DCN, and not the ipsilateral DCN, in

control animals indicating the retrograde tract tracing reliably labelled the SCPs in our

hands (Figures 1D-E, S1A-B, dorsal: 6/8; ventral: 8/11). In the $Smo^{Nex-Cre}$ mice, the

results differed depending on whether we injected in the dorsal or ventral thalamus

(Figures 1F-G, S1C-D). In the ventral thalamus injections, we detected dye-stained

clusters of cells in the contralateral DCN but not the ipsilateral DCN, indicating normal

SCP projection to the ventral thalamus (Figure 1G, 4/4 injections). This indicates that at

least some SCPs cross the midline. In the dorsal thalamus injections, we could not

detect dye-stained clusters of cells in either the contralateral or ipsilateral DCN (Figure

1F, 0/6 injections) suggesting that SCPs lacking SMO do not project to the dorsal

thalamus. These data implicate SMO as critical for proper projection of the SCPs to the

dorsal thalamus.

ARL13B is required for normal SCP projection to the dorsal thalamus

Given that ARL13B regulates vertebrate Hh signaling in a variety of contexts, we

next assessed ARL13B's role in proper SCP projection. In order to delete *ARL13B*

specifically in projection neurons, we generated *Nex-Cre;Arl13b^{fl/fl}* mice, which we refer

to as *Arl13b^{Nex-Cre}*. We performed dorsal and ventral thalamus injections for retrograde

tract tracing to examine the SCPs (Figures 1H-I, S1C-D). In the ventral thalamus

injections of *Arl13b^{Nex-Cre}* mice, we found dye-stained clusters of cells in the

contralateral DCN consistent with normal SCP projections crossing the midline and

projecting to the ventral thalamus (Figure 1I, 4/4 injections). In contrast, in the dorsal

thalamus injections of *Arl13b*$^{Nex\text{-}Cre}$ mice, we generally did not detect dye-stained

clusters of cells in either the contralateral or ipsilateral DCN, suggesting that the SCPs

lacking ARL13B do not project to the dorsal thalamus (Figure 1H, 1/7 injections). These

data link ARL13B function to normal SCP projection. Furthermore, they reveal a similar

phenotype in *Smo*$^{Nex\text{-}Cre}$ and *Arl13b*$^{Nex\text{-}Cre}$ mice.

ARL13B does not function from within cilia to mediate SCP guidance

ARL13B and the other 35 genes implicated in Joubert syndrome associate with

the cilium or centrosome leading to the assumption that protein dysfunction from these

locales underlies JSRD phenotypes (Parisi 2019). To directly ask whether ARL13B

mediates SCP guidance to the dorsal thalamus from within cilia, we examined a mouse

expressing a cilia-excluded variant of ARL13B, ARL13B^{V358A} (Figure 2) (Gigante *et al.*

2020). We previously demonstrated that ARL13B^{V358A} retains all known ARL13B

biochemical activity, is undetectable in cilia yet transduces vertebrate Hh signaling

normally (Mariani *et al.* 2016; Gigante *et al.* 2020). We found that either dorsal or ventral

thalamus injections resulted in visible clusters of dye-stained cells in the contralateral

DCN in control (Figures 2A-B, S1G-H, dorsal: 3/3; ventral: 4/4) and *Arl13b*$^{V358A/V358A}$

(Figure 2C-D, dorsal: 4/4; ventral: 3/3) animals. In the context of the previous result

showing that *Arl13b*$^{Nex\text{-}Cre}$ mice display abnormal SCP projections to the dorsal

thalamus, these data demonstrate that ARL13B does not function from within cilia to

regulate SCP projections.

SCP projection is normal in mice expressing a Joubert-causing allele, Arl13b^{R79Q}

In order to understand the relationship between ARL13B and MTS formation in

Joubert syndrome, we generated a mouse expressing the JSRD-causing R79Q

mutation (Figure 3A). We used CRISPR/Cas9 editing to change the conserved residue

in the mouse genome (Figures 3B, S2). This amino acid change disrupts ARL13B's

GEF activity for ARL3 (Gotthardt *et al.* 2015; Ivanova *et al.* 2017). We found

Arl13b$^{R79Q/R79Q}$ mice were viable and fertile. We bred the *Arl13b^{R79Q}* allele to the null

Arl13b$^{\Delta}$ allele to make *Arl13b$^{R79Q/\Delta}$* animals, which we found survived to adulthood. As

Arl13b$^{\Delta/\Delta}$ are embryonic lethal, this genetically demonstrates that *Arl13b^{R79Q}* is a

hypomorphic allele of *ARL13B* (Su *et al.* 2012).

To assess the role of ARL13B^{R79Q} in SCP guidance, we performed dorsal and

ventral thalamus dye injections in control and *Arl13b$^{R79Q/R79Q}$* mice. We identified visible

clusters of dye-stained cells in the contralateral DCN in control (Figures 3C-D, S1E-F,

dorsal: 4/4; ventral: 3/3) and *Arl13b$^{R79Q/R79Q}$* (Figure 3E-F, dorsal: 3/3; ventral: 3/4)

animals. Thus, despite the constitutive expression of the JSRD-causing allele

throughout development, we did not detect a SCP projection defect in the

Arl13b$^{R79Q/R79Q}$ mouse model. In the context of the abnormal SCP projections to the

dorsal thalamus that we identified in the *Smo$^{Nex-Cre}$* and *Arl13b$^{Nex-Cre}$* mice, this result

indicates that the *Arl13b^{R79Q}* allele does not disrupt SMO function or any ARL13B

function that regulates SMO.

Arl13b$^{R79Q/R79Q}$ mice display normal Shh signal transduction in neural tube patterning

To further investigate the role of the *Arl13b^{R79Q}* allele in Hh signaling, we

examined embryonic neural patterning as it is exquisitely sensitive to alterations in Shh

activity (Chiang *et al.* 1996; Briscoe and Ericson 1999). We generated E10.5 embryos

and stained neural tube sections with antibodies against Shh, Olig2 and Pax6 (Figure

4). As expected in wild type embryos, we observed Shh expression in the ventral-most

cells (the floorplate), Olig2 expression in lateral cells and Pax6 expression dorsally. We

also saw the established abnormal cell fates in null *Arl13b$^{\Delta/\Delta}$* embryos: loss of Shh

staining in the floorplate (Figure 4D), dorsal and ventral expansion of Olig2 expression

(Figure 4H), and a dorsal shift in Pax6 expression (Figure 4L) (Su *et al.* 2012). We

found both *Arl13b$^{R79Q/R79Q}$* (Figure 4B, F, J) and *Arl13b$^{R79Q/\Delta}$* (Figure 4C, G, K) embryos

displayed neural patterning indistinguishable from wild type embryos, indicating the

Arl13b^{R79Q} allele is not dosage-sensitive and does not disrupt Shh signaling in

determining neural cell fate. In addition, ARL13B^{R79Q} protein localized to cilia (Figure

4N), consistent with previous results (Li *et al.* 2010; Humbert *et al.* 2012; Li *et al.* 2016;

Mariani *et al.* 2016).

Arl13b$^{R79Q/R79Q}$ mice display normal cerebellar width

The lack of a SCP projection phenotype in the *Arl13b$^{R79Q/R79Q}$* mice surprised us

since JS patients display the MTS. In addition to defects in the SCPs, the MTS is due to

an underdeveloped cerebellar vermis, so we examined the width of the cerebellum and

the cerebellar vermis (Figure 5) (Aguilar *et al.* 2012). To quantify cerebellar width, we

performed analysis of surface-facing anatomical measurements validated to be

sufficiently sensitive to detect small differences in vermis width (Deshpande *et al.* 2020).

Briefly, we measured cerebellar width as well as cerebellar vermis width (widest part of

lobule VII) of fixed whole mount dissected brains. For each sex- and age-matched pair,

we calculated the ratio of the width measurements from mutant to control, which we

compared to a hypothetical value of 1 (indicating no difference between groups). We

detected no differences in the overall cerebellar width or the cerebellar vermis width

between control and $Arl13b^{R79Q/R79Q}$ mice of either sex (Figure 5B). Furthermore, we

found no difference in body or brain weight for $Arl13b^{R79Q/R79Q}$ mutants compared to

controls (Figure 5B). Thus, unlike patients carrying $ARL13B^{R79Q/R79Q}$, $Arl13b^{R79Q/R79Q}$

mice do not display a detectable growth deficit in the cerebellar vermis.

Global cerebellar hypoplasia is observed in mice lacking Ar13b in all neurons

Cerebellar size is well established to be regulated, in part, via Shh signaling

which controls proliferation of the cerebellar granule precursor cells (Kenney and

Rowitch 2000; Chizhikov *et al.* 2007). In order to better understand the vermis

hypoplasia phenotype seen in JBTS patients in relation to ARL13B, we wanted to

investigate how ARL13B regulates cerebellar width. To do so, we crossed the *Brn4-Cre*

allele into the conditional null $Arl13b^{fl/fl}$ background, called $Arl13b^{Brn4-Cre}$ (Figure 6). *Brn4-*

Cre initiates expression at E8.5 throughout the neuroectoderm so the cerebellum

develops in the absence of ARL13B (Heydemann *et al.* 2001; Hazen *et al.* 2012). We

again calculated width ratios using surface-facing anatomical measurements and found

the overall width of the cerebellum was 6% reduced in both females and males lacking

ARL13B compared to control littermates at weaning (Figure 6B, $p<0.1$). More striking, in

the cerebellar vermis we detected a 27% reduction in width in female and a 33%

reduction in male mutants compared to controls (Figure 6B, $p<0.0001$). $Arl13b^{Brn4-Cre}$

mice develop hydrocephaly just after weaning which often leads to death. This is

reflected in the body and brain weight ratios: while the mutant mice had slightly lower

body weights, they had comparatively heavier brains (Figure 6B). From these data, we

conclude that loss of *ARL13B* leads to a modest global cerebellar width deficit and a

more pronounced cerebellar vermis width reduction.

Arl13b$^{V358A/V358A}$ mice display normal cerebellar width

JS is classified as a ciliopathy due to the majority of causative genes encoding

proteins that, like ARL13B, are associated with cilia. In order to better understand the

role of ciliary ARL13B in cerebellar size, we examined cerebellar width in the mice

expressing the cilia-excluded variant ARL13B^{V358A} (Figure 7) (Gigante *et al.* 2020). We

detected no difference in the overall cerebellar width or that of the cerebellar vermis

between control and *Arl13b$^{V358A/V358A}$* mice (Figure 7B). Thus, ARL13B does not control

cerebellar width from within cilia.

DISCUSSION

Here we demonstrate that complete loss of *Arl13b* function in mouse can account

for two aspects of the MTS: aberrant SCP thalamic targeting and cerebellar vermis

hypoplasia. We expand the role of Hh signaling as a critical guidance cue by showing it

is required for proper SCP projection to the dorsal thalamus. Our finding that the SCP

phenotype is similar in *Smo$^{Nex-Cre}$* and *Arl13b$^{Nex-Cre}$* mice is consistent with a model

whereby ARL13B regulates SCP projections to the dorsal thalamus via a SMO-

dependent mechanism. In line with previous work showing that ARL13B does not

function from within cilia to regulate Shh-guided axon guidance, we found normal SCP

thalamic targeting in mice expressing only a cilia-excluded ARL13B variant (Ferent *et al.*

2019). By mutating a conserved arginine to glutamine, we generated a mouse

expressing a mutation linked to JSRD in humans and observed no change in vertebrate

Hh signaling (Cantagrel *et al.* 2008). Additionally, we identified no defects in

Arl13b$^{R79Q/R79Q}$ SCP projections. Whereas complete ARL13B deletion (*Arl13b$^{Brn4-Cre}$*) in

the cerebellum led to global hypoplasia, we show the cerebellum of *Arl13b$^{R79Q/R79Q}$* mice

is not significantly different in width compared to controls.

Overall, our data indicate ARL13B function is critical for both SCP targeting and

controlling cerebellar vermis width. At one level, our data implicate Hh signaling in the

etiology of the MTS since we show that SCP targeting requires SMO. However, at

another level, our data indicate that Hh-independent pathways are at play as we don't

observe Hh-dependent neural tube patterning defects in the presence of the JSRD-

causing *Arl13b^{R79Q}* allele. In other mouse models of JSRD where the hypoplasia is

specific to the cerebellar vermis, Wnt signaling is affected (Lancaster *et al.* 2011). Thus,

the MTS could be due to disruption of different pathways in the SCPs and the vermis.

This would imply that the 35 JSRD implicated genes all affect the distinct pathways in a

similar manner (Parisi 2019). Alternatively, the MTS may form due to alterations in any

of a few pathways – and it is even possible that alterations in one pathway could impact

other pathways – or the ability of cells to respond to those other pathways. Such a

model is hinted at by previous work showing interplay between the Hh and Wnt

pathways underlying the severity of hindbrain phenotypes (Hagemann and Scholpp

2012; Bashford and Subramanian 2019). While we haven't detected any changes in

Wnt response in the absence of ARL13B function, we may have not examined the

relevant biological process or used a sensitive enough readout (Horner and Caspary

2011). Parallel reasoning would thus suggest that while $Arl13b^{R79Q/R79Q}$ mice clearly

transduce Hh reasonably well, there may be subtle changes in Shh signaling or

changes that influence Wnt signaling. Our data are consistent with the complexity

exhibited by other JSRD mouse models examined to date (Delous *et al.* 2007; Garcia-

Gonzalo *et al.* 2011; Roberson *et al.* 2015; Bashford and Subramanian 2019).

In patients, the SCP targeting deficit appears more severe than what we

observed in the mice. Available methods for live-imaging and examination of post-

mortem tissues suggest a range of SCP decussation defects, with some tracts

appearing thickened on the ipsilateral side relative to their DCN (Yachnis and Rorke

1999; Poretti *et al.* 2007). Surprisingly, in the mutant mice we infer midline crossing of

the SCPs. In the case of the $Smo^{Nex-Cre}$ conditional mice, it is formally possible that the

SCPs do not rely on SMO for midline crossing but only for subsequent targeting to the

dorsal thalamus. The fact that the $Arl13b^{Nex-Cre}$ conditional mice display a highly similar

phenotype to the $Smo^{Nex-Cre}$ conditional mice makes this less likely, since ARL13B is

directly implicated in JSRD and regulates SMO-dependent axon guidance in other

contexts (Cantagrel *et al.* 2008; Ferent *et al.* 2019). It is also plausible that the protein

turnover driven by *Nex-Cre* completed after midline crossing occurred. *Nex-Cre*

expression initiates at E11.5 in the cells on the rhombic lip of the cerebellar anlage as

they start to migrate and be specified before occupying the deep cerebellar nuclei (Fink

et al. 2006; Goebbels *et al.* 2006). We expect deletion would occur in the precursors

and therefore the neurons of the DCN would not express protein. Finally, it is possible

that the mouse is not a valid system in which to model the SCP midline crossing defect.

This might explain why we saw no defects in the SCP targeting of the $Arl13b^{R79Q/R79Q}$

mice, as this is a constitutive mutation that requires no protein turnover, yet

homozygous expression of ARL13B^{R79Q} in patients results in the molar tooth sign, which

has been associated with failed decussation of white matter tracts (Quisling *et al.* 1999).

Indeed, other mouse mutants such as *Cep290* and *Ahi1* which recapitulate the

cerebellar vermis hypoplasia, also do not display midline crossing defects in the SCPs

(Lancaster *et al.* 2011). Whether this is due to anatomical distinctions between the

cerebellum in mouse and human or the genetic background on which these models

were examined are open questions. Recent work highlights clear molecular and

temporal differences between mouse and human cerebellar development (Haldipur *et*

al. 2019; Behesti *et al.* 2021).

Examining SCP projections is labor intensive and it has not been done

systematically among the JSRD mouse models (Bashford and Subramanian 2019; Guo

et al. 2019). While previous work showed that $Arl13b^{Nex-Cre}$ and $Inpp5e^{Nex-Cre}$ mice

exhibit SCP targeting deficits, here we pinpoint the $Arl13b^{Nex-Cre}$ defect as specific to the

projection to the dorsal thalamus (Guo *et al.* 2019). The projection to the ventral

thalamus remains intact, suggesting there is not a generalized deficit in axon outgrowth

within the tract. The work on the $Arl13b^{Nex-Cre}$ and $Inpp5e^{Nex-Cre}$ SCP targeting deficits

argue that PI3K/Akt signaling from within cilia led to the tract defects (Guo *et al.* 2019).

However, we found that cilia-excluded ARL13B mediated SCP targeting normally.

These conflicting results could be explained by differences in the experimental details

as the data supporting ciliary ARL13B function used a human ARL13B viral expression

construct to rescue conditionally-deleted mice whereas we used genetic mutations

engineered at the endogenous locus in this study. Alternatively, these data could

indicate that ARL13B plays an important cellular role in the ciliary trafficking of key

components needed for the PI3K/Akt pathways.

JSRD-causing mutations in ARL13B are generally restricted to the GTPase

domain of the protein, although two residues outside that domain are implicated in

disease (Cantagrel *et al.* 2008; Bachmann-Gagescu *et al.* 2015; Thomas *et al.* 2015;

Shaheen *et al.* 2016; Rafiullah *et al.* 2017). Based on other ARL proteins, ARL13B likely

assumes distinct conformations upon the binding either GDP or GTP, permitting

different binding partners or altering affinities for binding partners (Pasqualato *et al.*

2002; Miertzschke *et al.* 2014). None of the tested JSRD-causing mutations (R79Q,

Y86C or R200C) disrupt GTP binding or hydrolysis, however, all three mutations disrupt

ARL13B function as an ARL3 GEF (Ivanova *et al.* 2017). The arginine at this position is

located within the Switch 2 region of ARL13B and is conserved in humans, mice,

zebrafish and *Chlamydomonas* (Figure 3B). We were surprised that we were unable to

detect any defects in our mouse, as we expected to model some aspect of Joubert

syndrome. Humans with homozygous R79Q mutation exhibit motor and ocular defects

(among other symptoms), whereas *arl13b*-null zebrafish injected with human ARL13B-

R79Q mRNA only partially rescue the ciliopathy phenotypes of curved body axis and

cystic kidney (Cantagrel *et al.* 2008). These species-dependent differences emphasize

the importance of analyzing mutations expressed from the endogenous promoter over

the course of normal development. A recent study linked *arl13b* disruption during

zebrafish development to reduced granule and Purkinje cells through a down-regulation

of Wnt signaling (Zhu *et al.* 2020). Given that complete deletion of ARL13B impacts

broader biological processes in the cerebellum than the R79Q mutation and that the null

mutant misregulates Hh signaling whereas R79Q does not, we conclude that a subset

of ARL13B function is disrupted in JSRD.

Acknowledgements

We are grateful to L. Mariani for her initial work on this project, R.E. Van Sciver and E.

Gigante for critical comments on the manuscript and J.G. Mulle for the statistical

consults.

Funding

This work was supported by funding from National Institutes of Health grants

R01NS090029, R01GM110663 and R35GM122549 to T.C. and T32GM008490 and

F31NS101806 to S.K.S. with additional support from the Emory University Integrated

Cellular Imaging Microscopy Core of the Emory Neuroscience NINDS Core Facilities

grant, P30NS055077. This study was supported in part by the Mouse Transgenic and

Gene Targeting Core (TMF), which is subsidized by the Emory university School of

Medicine and is one of the Emory Integrated Core Facilities. Additional support was

provided by the National Center for Advancing Translational Sciences of the National

Institutes of Health under Award Number UL1TR000454. The content is solely the

responsibility of the authors and does not necessarily reflect the official views of the

National Institutes of Health

Conflicts of Interest

The authors have no competing interests to declare.

Author contributions statement

Author Contributions: Conceptualization T.C.; Methodology S.K.S. and A.B.L.;

Validation S.K.S. and A.B.L; Formal Analysis S.K.S. and A.B.L.; Investigation S.K.S.

and A.B.L.; Writing – Original Draft S.K.S. and T.C.; Writing – Review & Editing S.K.S.,

A.B.L. and T.C.; Visualization S.K.S. and A.B.L.; Supervision T.C.; Project

Administration T.C.; Funding Acquisition T.C.

References

Abdelhamed, Z. A., D. I. Abdelmottaleb, M. E. El-Asrag, S. Natarajan, G. Wheway et al.,

2019 The ciliary Frizzled-like receptor Tmem67 regulates canonical Wwnt/beta-

catenin signalling in the developing cerebellum via Hoxb5. *Sci Rep.* **9:** 5446.

doi:10.1038/s41598-019-41940-5

Abdelhamed, Z. A., G. Wheway, K. Szymanska, S. Natarajan, C. Toomes et al., 2013

Variable expressivity of ciliopathy neurological phenotypes that encompass

Meckel-Gruber syndrome and Joubert syndrome is caused by complex de-

regulated ciliogenesis, Shh and Wnt signalling defects. *Hum Mol Genet.* **22:**

1358-1372. doi:10.1093/hmg/dds546

Aguilar, A., A. Meunier, L. Strehl, J. Martinovic, M. Bonniere et al., 2012 Analysis of

human samples reveals impaired SHH-dependent cerebellar development in

Joubert syndrome/Meckel syndrome. *Proc Natl Acad Sci U S A.* **109:** 16951-

16956. doi:10.1073/pnas.1201408109

Alkanderi, S., E. Molinari, R. Shaheen, Y. Elmaghloob, L. A. Stephen et al., 2018 ARL3

mutations cause Joubert syndrome by disrupting ciliary protein composition. *Am*

J Hum Genet. **103:** 612-620. doi:10.1016/j.ajhg.2018.08.015

Arts, H. H., D. Doherty, S. E. C van Beersum, M. A. Parisi, S. J. F. Letteboer et al., 2007

Mutations in the gene encoding the basal body protein RPGRIP1L, a

nephrocystin-4 interactor, cause Joubert syndrome. *Nat Genet.* **39:** 882-888.

doi:10.1038/ng2069

Asadollahi, R., J. E. Strauss, M. Zenker, O. Beuing, S. Edvardson et al., 2018 Clinical

and experimental evidence suggest a link between KIF7 and C5orf42-related

ciliopathies through Sonic Hedgehog signaling. *Eur J Hum Genet.* **26:** 197-209.

doi:10.1038/s41431-017-0019-9

Bachmann-Gagescu, R., J. C. Dempsey, I. G. Phelps, B. J. O'Roak, D. M. Knutzen et

al., 2015 Joubert syndrome: A model for untangling recessive disorders with

extreme genetic heterogeneity. *J Med Genet.* **52:** 514-522.

doi:10.1136/jmedgenet-2015-103087

Bachmann-Gagescu, R., J. C. Dempsey, S. Bulgheroni, M. L. Chen, S. D'Arrigo et al.,

2020 Healthcare recommendations for Joubert syndrome. *Am J Med Genet A.*

182: 229-249. doi:10.1002/ajmg.a.61399

Bashford, A. L., and V. Subramanian, 2019 Mice with a conditional deletion of Talpid3

(KIAA0586) - a model for Joubert syndrome. *J Pathol.* **248:** 396-408.

doi:10.1002/path.5271

Bay, S.N., A. B. Long, and T. Caspary, 2018 Disruption of the ciliary GTPpase Arl13b

suppresses Sonic hedgehog overactivation and inhibits medulloblastoma

formation. *Proc Natl Acad Sci U S A.* **115:** 1570-1575.

doi:10.1073/pnas.1706977115

Behesti, H. A. Kocabas, D. E. Buchholz, T. S. Carroll, and M. E. Hatten, 2021 Altered

temporal sequence of transcriptional regulators in the generation of human

cerebellar granule cells. bioRxiv. doi:10.1101/2021.01.17.427030 (Preprint

posted January 17, 2021).

Bielas, S. L., J. L. Silhavy, F. Brancati, M. V. Kisseleva, L. Al-Gazali et al., 2009

Mutations in INPP5E, encoding inositol polyphosphate-5-phosphatase E, link

phosphatidyl inositol signaling to the ciliopathies. *Nat Genet.* **41:** 1032-1036.

doi:10.1038/ng.423

Bimonte, S., A. De Angelis, L. Quagliata, F. Giusti, R. Tammaro et al., 2011 Ofd1 is

required in limb bud patterning and endochondral bone development. *Dev Biol.*

349: 179-191. doi:10.1016/j.ydbio.2010.09.020

Bohne, P., M. K. Schwarz, S. Herlitze, and M. D. Mark, 2019 A new projection from the

deep cerebellar nuclei to the hippocampus via the ventrolateral and laterodorsal

thalamus in mice. *Front Neural Circuits.* **13:** 51. doi:10.3389/fncir.2019.00051

Brancati, F., B. Dallapiccola, and E. M. Valente, 2010 Joubert syndrome and related

disorders. *Orphanet J Rare Dis.* **5:** 20. doi:10.1186/1750-1172-5-20

Breslow, D. K., S. Hoogendoorn, A. R. Kopp, D. W. Morgens, B. K. Vu et al., 2018 A

CRISPR-based screen for Hedgehog signaling provides insights into ciliary

function and ciliopathies. *Nat Genet.* **50:** 460-471. doi:10.1038/s41588-018-0054-

7

Briscoe, J., and J. Ericson J, 1999 The specification of neuronal identity by graded

Sonic Hedgehog signalling. *Semin Cell Dev Biol.* **10:** 353-362.

doi:10.1006/scdb.1999.0295

Cantagrel, V., J. L. Silhavy, S. L. Bielas, D. Swistun, S. E. Marsh et al., 2008 Mutations

in the cilia gene ARL13B lead to the classical form of Jjoubert syndrome. *Am J*

Hum Genet. **83:** 170-179. doi:10.1016/j.ajhg.2008.06.023

Casoni, F., L. Croci, F. Vincenti, P. Podini, M. Riba et al., 2020 ZFP423 regulates early

patterning and multiciliogenesis in the hindbrain choroid plexus. *Development.*

147: dev190173. doi:10.1242/dev.190173

Caspary, T., M. J. Garcia-Garcia, D. Huangfu, J. T. Eggenschwiler, M. R. Wyler et al.,

2002 Mouse Dispatched homolog1 is required for long-range, but not juxtacrine,

Hh signaling. *Curr Biol.* **12:** 1628-1632. doi:10.1016/s)960-9822(02)01147-8

Caspary, T., C. E. Larkins, and K. V. Anderson, 2007 The graded response to Sonic

Hedgehog depends on cilia architecture. *Dev Cell.* **12:** 767-778.

doi:10.1016/j.devcel.2007.03.004

Charron, F., E. Stein, J. Jeong, A. P. McMahon, and M. Tessier-Lavigne, 2003 The

morphogen sonic hedgehog is an axonal chemoattractant that collaborates with

netrin-1 in midline axon guidance. *Cell.* **113:** 11-23. doi:10.1016/s0092-

8674(03)00199-5

Chavez, M., S. Ena, J. Van Sande, A. de Kerchove d'Exaerde, S. Schurmans et al.,

2015 Modulation of ciliary phosphoinositide content regulates trafficking and

Sonic hedgehog signaling output. *Dev Cell.* **34:** 338-350.

doi:10.1016/j.devcel.2015.06.016

Chiang, C., Y. Litingtung, E. Lee, K. E. Young, J. L. Corden et al., 1996 Cyclopia and

defective axial patterning in mice lacking Ssonic hedgehog gene function.

Nature. **383:** 407-413. Doi:10.1038/383407a0

Chih, B., P. Liu, Y. Chinn, C. Chalouni, L. G. Komuves et al., 2011 A ciliopathy complex

at the transition zone protects the cilia as a privileged membrane domain. *Nat*

Cell Biol. **14:** 61-72. doi:10.1038/ncb2410

Chizhikov, V. V., J. Davenport, Q. Zhang, E. K. Shih, O. A. Cabello et al., 2007 Cilia

proteins control cerebellar morphogenesis by promoting expansion of the granule

progenitor pool. *J Neurosci.* **27:** 9780-9789. doi:10.1523/JNEUROSCI.5586-

06.2007

Christopher, K. J., B. Wang, Y. Kong, and S. D. Weatherbee, 2012 Forward genetics

uncovers Transmembrane protein 107 as a novel factor required for ciliogenesis

and Sonic hedgehog signaling. *Dev Biol.* **368:** 382-392.

doi:10.1016/j.ydbio.2012.06.008

Constable, S., A. B. Long, K. A. Floyd, S. Schurmans, and T. Caspary, 2020 The ciliary

phosphatidylinositol phosphatase Inpp5e plays positive and negative regulatory

roles in Shh signaling. *Development.* **147:** dev183301. doi:10.1242/dev.183301

Consugar, M. B., V. J. Kubly, D. J. Lager, C. J. Hommerding, W. C. Wong et al., 2007

Molecular diagnostics of Meckel–Gruber syndrome highlights phenotypic

differences between MKS1 and MKS3. *Hum Genet.* **121:** 591-599.

doi:10.1007/s00439-007-0341-3

Dahmane, N., and A. Ruiz i Altaba, 1999 Sonic hedgehog regulates the growth and

patterning of the cerebellum. *Development.* **126:** 3089-3100.

640 Davey, M. G., I. R. Paton, Y. Yin, M. Schmidt, F. K. Bangs et al., 2006 The chicken

641   talpid3 gene encodes a novel protein essential for Hedgehog signaling. *Genes*

642   *Dev.* **20:** 1365-1377. doi:10.1101/gad.369106

643 Delous, M., L. Baala, R. Salomon, C. Laclef, J. Vierkotten et al., 2007 The ciliary gene

644   RPGRIP1L is mutated in cerebello-oculo-renal syndrome (Joubert syndrome type

645   B) and Meckel syndrome. *Nat Genet.* **39:** 875-881. doi:10.1038/ng2039

646 Deshpande, O., R. Z. Lara, O. R. Zhang, D. Concepcion, and B. A. Hamilton, 2020

647   ZNF423 patient variants, truncations, and in-frame deletions in mice define an

648   allele-dependent range of midline brain abnormalities. *PLoS Genet.* **16:**

649   e1009017. doi:10.1371/journal.pgen.1009017

650 Doherty, D., 2009 Joubert syndrome: Insights into brain development, cilium biology,

651   and complex disease. *Semin Pediatr Neurol.* **16:** 143-154.

652   doi:10.1016/j.spen.2009.06.002

653 Dowdle, W. E., J. F. Robinson, A. Kneist, M. S. Sirerol-Piquer, S. G. M. Frints et al.,

654   2011 Disruption of a ciliary B9 protein complex causes Meckel syndrome. *Am J*

655   *Hum Genet.* **89:** 94-110. doi:10.1016/j.ajhg.2011.06.003

656 Ferent, J., S. Constable, E. D. Gigante, P. T. Yam, L. E. Mariani et al., 2019 The ciliary

657   protein Arl13b functions outside of the primary cilium in Shh-mediated axon

658   guidance. *Cell Rep.* **29:** 3356-3366.e3353. doi:10.1016/j.celrep.2019.11.015

659 Fink, A. J., C. Englund, R. A. M. Daza, D. Pham, C. Lau et al., 2006 Development of

660   the deep cerebellar nuclei: Transcription factors and cell migration from the

661   rhombic lip. *J Neurosci.* **26:** 3066-3076. doi:10.1523/JNEUROSCI.5203-05.2006

Frikstad, K-A. M., E. Molinari, M. Thoresen, S. A. Ramsbottom, F. Hughes et al., 2019

A CEP104-CSPP1 complex is required for formation of primary cilia competent in

Hedgehog signaling. *Cell Rep.* **28:** 1907-1922.e6

doi:10.1016/j.celrep.2019.07.025

Garcia-Gonzalo, F. R., K. C. Corbit, M. S. Sirerol-Piquer, G. Ramaswami, E. A. Otto et

al., 2011 A transition zone complex regulates mammalian ciliogenesis and ciliary

membrane composition. *Nat Genet.* **43:** 776-784. doi:10.1038/ng.891

Garcia-Gonzalo, F. R., S. C. Phua, E. C. Roberson, G. Garcia 3rd, M. Abedin et al.,

2015 Phosphoinositides regulate ciliary protein trafficking to modulate Hedgehog

signaling. *Dev Cell.* **34:** 400-409. doi:10.1016/j.devcel.2015.08.001

Gaudet, M., A-G. Fara, I. Beritognolo, and M. Sabatti, 2009 Allele-specific PCR in SNP

genotyping. *Methods Mol Biol.* **578:** 415-424. doi:10.1007/978-1-60327-411-1_26

Gigante, E. D., M. R. Taylor, A. A. Ivanova, R. A. Kahn, and T. Caspary, 2020. ARL13B

regulates Sonic hedgehog signaling from outside primary cilia. *eLife.* **9:** e50434.

doi:10.7554/eLife.50434

Goebbels, S., I. Bormuth, U. Bode, O. Hermanson, M. H. Schwab et al., 2006 Genetic

targeting of principal neurons in neocortex and hippocampus of NEX-Cre mice.

Genesis. **44:** 611-621. doi:10.1002/dvg.20256

Gotthardt, K., M. Lokaj, C. Koerner, N. Falk, A. Giessl et al., 2015 A G-protein

activation cascade from Arl13b to Arl3 and implications for ciliary targeting of

lipidated proteins. *eLife.* **4:** e11859. doi:10.7554/eLife.11859

Guo, J., J. M. Otis, S. K. Suciu, C. Catalano, L. Xing et al., 2019 Primary cilia signaling

promotes axonal tract development and is disrupted in Joubert syndrome-related

disorders models. *Dev Cell.* **51:** 759-774 e755. doi:10.1016/j.devcel.2019.11.005

Hagemann, A. I. H., and S. Scholpp, 2012 The tale of the three brothers – Shh, Wnt,

and Fgf during development of the thalamus. *Front Neurosci.* **6:** 76.

doi:10.3389/fnins.2012.00076

Haldipur, P., K. A. Aldinger, S. Bernardo, M. Deng, A. E. Timms et al., 2019

Spatiotemporal expansion of primary progenitor zones in the developing human

cerebellum. *Science*. **366:** 454-460. Doi: 10.1126/science.aax7526

Hata, A., J. Seoane, G. Lagna, E. Montalvo, A. Hemmati-Brivanlou et al., 2000 OAZ

uses distinct DNA- and protein-binding zinc fingers in separate BMP-Smad and

Olf signaling pathways. *Cell.* **100:** 229-240. doi:10.1016/s0092-8674(00)81561-5

Hazen, V. M. , M. G. Andrews, L. Umans, E. B. Crenshaw 3rd, A. Zwijsen et al., 2012

BMP receptor-activated Smads confer diverse functions during the development

of the dorsal spinal cord. *Dev Biol.* **367:** 216-227.

doi:10.1016/j.ydbio.2012.05.014

Heydemann, A., L. C. Nguyen, and E. B. Crenshaw 3rd, 2001 Regulatory regions from

the Brn4 promoter direct LACZ expression to the developing forebrain and neural

tube. *Brain Res Dev Brain Res.* **128:** 83-90. doi:10.1016/s0165-3806(01)00137-7

Hopp, K., C. M. Heyer, C. J. Hommerding, S. A. Henke, J. L. Sundsbak et al., 2011

B9D1 is revealed as a novel Meckel syndrome (MKS) gene by targeted exon-

enriched next-generation sequencing and deletion analysis. *Hum Mol Genet.* **20:**

2524-2534. doi:10.1093/hmg/ddr151

Horner, V. L., and T. Caspary, 2011   Disrupted dorsal neural tube BMP signaling in the

cilia mutant Arl13b hnn stems from abnormal Shh signaling. *Dev Biol.* **355:** 43-

54. doi:10.1016/j.ydbio.2011.04.019

Huang, S., J. Laoukili, M. T. Epping, J. Koster, M. Hölzel et al., 2009   ZNF423 is

critically required for retinoic acid-induced differentiation and is a marker of

neuroblastoma outcome. *Cancer Cell.* **15:** 328-340.

doi:10.1016/j.ccr.2009.02.023

Huangfu, D., A. Liu, A. S. Rakeman, N. S. Murcia, L. Niswander et al., 2003   Hedgehog

signalling in the mouse requires intraflagellar transport proteins. *Nature.* **426:** 83-

87. doi:10.1038/nature02061

Humbert, M. C., K. Weihbrecht, C. C. Searby, Y. Li, R. M. Pope et al., 2012   ARL13B,

PDE6D, and CEP164 form a functional network for INPP5E ciliary targeting. *Proc*

Natl Acad Sci U S A. **109:** 19691-19696. doi:10.1073/pnas.1210916109

Hynes, A. M., R. H. Giles, S. Srivastava, L. Eley, J. Whitehead et al., 2014   Murine

Joubert syndrome reveals Hedgehog signaling defects as a potential therapeutic

target for nephronophthisis. *Proc Natl Acad Sci U S A.* **111:** 9893-9898.

doi:10.1073/pnas.1322373111

Ivanova, A. A., T. Caspary, N. T. Seyfried, D. M. Duong, A. B. West et al., 2017

Biochemical characterization of purified mammalian ARL13B protein indicates

that it is an atypical GTPase and ARL3 guanine nucleotide exchange factor

(GEF). *J Biol Chem.* **292:** 11091-11108. doi:10.1074/jbc.M117.784025

Kenney, A. M., and D. H. Rowitch, 2000   Sonic hedgehog promotes G(1) cyclin

expression and sustained cell cycle progression in mammalian neuronal

precursors. *Mol Cell Biol.* **20:** 9055-9067. doi:10.1128/mcb.20.23.9055-

9067.2000

Lai, C. K., N. Gupta, X. Wen, L. Rangell, B. Chih et al., 2011 Functional

characterization of putative cilia genes by high-content analysis. *Mol Biol Cell.*

22: 1104-1119. doi:10.1091/mbc.E10-07-0596

Lan, Y., and R. Jiang, 2009 Sonic hedgehog signaling regulates reciprocal epithelial-

mesenchymal interactions controlling palatal outgrowth. *Development*. **136:**

1387-1396. doi:10.1242/dev.028167

Lancaster, M. A., D. J. Gopal, J. Kim, S. N. Saleem, J. L. Silhavy et al., 2011 Defective

Wnt-dependent cerebellar midline fusion in a mouse model of Joubert syndrome.

Nat Med. **17:** 726-731. doi:10.1038/nm.2380

Li, Y., Q. Wei, Y. Zhang, K. Ling, and J. Hu, 2010 The small GTPases ARL-13 and

ARL-3 coordinate intraflagellar transport and ciliogenesis. *J Cell Biol.* **189:** 1039-

1051. doi:10.1083/jcb.200912001

Li, Y., X. Tian, M. Ma, S. Jerman, S. Kong et al., 2016 Deletion of ADP ribosylation

factor-like GTPase 13B leads to kidney cysts. *J Am Soc Nephrol.* **27:** 3628-3638.

doi:10.1681/ASN.2015091004

Lipinski, R. J., C. Song, K. K. Sulik, J. L. Everson, J. J. Gipp et al., 2010 Cleft lip and

palate results from Hedgehog signaling antagonism in the mouse: Phenotypic

characterization and clinical implications. *Birth Defects Res A Clin Mol Teratol.*

88: 232-240. doi:10.1002/bdra.20656

Maria, B. L., K. B. Hoang, R. J. Tusa, A. A. Mancuso, L. M. Hamed et al., 1997 "Joubert

syndrome" revisited: key ocular motor signs with magnetic resonance imaging

correlation. *J Child Neurol.* **12:** 423-430. doi:10.1177/088307389701200703

Mariani, L. E., M. F. Bijlsma, A. A. Ivanova, S. K. Suciu, R. A. Kahn et al., 2016 Arl13b

regulates Shh signaling from both inside and outside the cilium. *Mol Biol Cell.* **27:**

3780-3790. doi:10.1091/mbc.E16-03-0189

Miertzschke, M., C. Koerner, M. Spoerner, and A. Wittinghofer, 2014 Structural insights

into the small G-protein Arl13b and implications for Joubert syndrome. *Biochem*

J. **457:** 301-311. doi:10.1042/BJ20131097

Munoz-Estrada, J., and R. J. Ferland, 2019 Ahi1 promotes Arl13b ciliary recruitment,

regulates Arl13b stability and is required for normal cell migration. *J Cell Sci.*

132: jcs230680. doi:10.1242/jcs.230680

Nolan-Stevaux, O., J. Lau, M. L. Truitt, G. C. Chu, M. Hebrok et al., 2009 Gli1 is

regulated through Smoothened-independent mechanisms in neoplastic

pancreatic ducts and mediates PDAC cell survival and transformation. *Genes*

Dev. **23:** 24-36. doi:10.1101/gad.1753809

Nuovo, S., I. Bacigalupo, M. Ginevrino, R. Battini, E. Bertini et al., 2020 Age and sex

prevalence estimate of Joubert syndrome in Italy. *Neurology.* **94:** e797-e801.

doi:10/1212/WNL.0000000000008996

Onimaru, H., Y. Kumagawa, and I. Homma, 2006 Respiration-related rhythmic activity

in the rostral medulla of newborn rats. *J Neurophysiol.* **96:** 55-61.

doi:10.1152/jn.01175.2005

Otto, E. A., K. Tory, M. Attanasio, W. Zhou, M. Chaki et al, 2009 Hypomorphic

mutations in meckelin (MKS3/TMEM67) cause nephronophthisis with liver

fibrosis (NPHP11). *J Med Genet.* **46:** 663-670. doi:10.1136/jmg.2009.066613

Parisi, M. A, 2019 The molecular genetics of Joubert syndrome and related ciliopathies:

The challenges of genetic and phenotypic heterogeneity. *Transl Sci Rare Dis.* **4:**

25-49. doi:10.3233/TRD-190041

Parisi, M. A., D. Doherty, P. F. Chance, and I. A. Glass, 2007 Joubert syndrome (and

related disorders) (OMIM 213300). *Eur J Hum Genet.* **15:** 511-521.

doi:10.1038/sj.ejhg.5201648

Pasqualato, S., L. Renault, and J. Cherfils, 2002 Arf, Arl, Arp and Sar proteins: A family

of GTP-binding proteins with a structural device for 'front-back' communication.

EMBO Rep. **3:** 1035-1041. doi:10.1093/embo-reports/kvf221

Poretti, A., E. Boltshauser, T. Loenneker, E. M. Valente, F. Brancati et al., 2007

Diffusion tensor imaging in Joubert syndrome. *AJNR Am J Neuroradiol.* **28:**

1929-1933. doi:10.3174/ajnr.A0703

Quisling, R. G., A. J. Barkovich, and B. L. Maria, 1999 Magnetic resonance imaging

features and classification of central nervous system malformations in Joubert

syndrome. *J Child Neurol.* **14:** 628-635. doi:10.1177/088307389901401002

Rafiullah, R., A. B. Long, A. A. Ivanova, H. Ali, S. Berkel et al., 2017 A novel

homozygous ARL13B variant in patients with Joubert syndrome impairs its

guanine nucleotide-exchange factor activity. *Eur J Hum Genet.* **25:** 1324-1334.

doi:10.1038/s41431-017-0031-0

Ramsbottom, S. A., P. E. Thelwall, K. M. Wood, G. J. Clowry, L. A. Devlin et al., 2020

Mouse genetics reveals Barttin as a genetic modifier of Joubert syndrome. *Proc*

Natl Acad Sci U S A. **117:** 1113-1118. doi:10.1073/pnas.1912602117

Reiter, J. F., and W. C. Skarnes, 2006   Tectonic, a novel regulator of the Hedgehog

pathway required for both activation and inhibition. *Genes Dev.* **20:** 22-27.

doi:10.1101/gad.1363606

Roberson, E. C., W. E. Dowdle, A. Ozanturk, F. R. Garcia-Gonzalo, C. Li et al., 2015

TMEM231, mutated in orofaciodigital and Meckel syndromes, organizes the

ciliary transition zone. *J Cell Biol.* **209:** 129-142. doi:10.1083/jcb.201411087

Sang, L., J. J. Miller, K. C. Corbit, R. H. Giles, M. J. Brauer et al., 2011   Mapping the

NPHP-JBTS-MKS protein network reveals ciliopathy disease genes and

pathways. *Cell.* **145:** 513-528. doi:10.1016/j.cell.2011.04.019

Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair et al., 2012   Fiji: an

open-source platform for biological-image analysis. *Nat Methods.* **9:** 676-682.

doi:10.1038/nmeth.2019

Schrick, J. J., P. Vogel, A. Abuin, B. Hampton, and D. S. Rice, 2006   ADP-ribosylation

factor-like 3 is involved in kidney and photoreceptor development. *Am J Pathol.*

168: 1288-1298. doi:10.2353/ajpath.2006.050941

Schwarz, N., A. Lane, K. Jovanovic, D. A. Parfitt, M. Aguila et al., 2017   Arl3 and RP2

regulate the trafficking of ciliary tip kinesins. *Hum Mol Genet.* **26:** 2480-2492.

doi:10.1093/hmg/ddx143

Shaheen, R., K. Szymanska, B. Basu, N. Patel, N. Ewida et al., 2016 Characterizing

the morbid genome of ciliopathies. *Genome Biol.* **17:** 242. doi:10.1186/s13059-

016-1099-5

Srour, M., F. F. Hamdan, J. A. Schwartzentruber, L. Patry, L. H. Ospina et al., 2012

Mutations in TMEM231 cause Joubert syndrome in French Canadians. *J Med*

Genet. **49:** 636-641. doi:10.1136/jmedgenet-2012-101132

Su, C-Y., S. N. Bay, L. E. Mariani, M. J. Hillman, and T. Caspary, 2012 Temporal

deletion of Arl13b reveals that a mispatterned neural tube corrects cell fate over

time. *Development.* **139:** 4062-4071. doi:10.1242/dev.082321

Thomas, S., V. Cantagrel, L. Mariani, V. Serre, J-E. Lee et al., 2015 Identification of a

novel ARL13B variant in a Joubert syndrome-affected patient with retinal

impairment and obesity. *Eur J Hum Genet.* **23:** 621-627.

doi:10.1038/ejhg.2014.156

Thomas, S., M. Legendre, S. Saunier, B. Bessières, C. Alby et al., 2012 TCTN3

mutations cause Mohr-Majewski syndrome. *Am J Hum Genet.* **91:** 372-378.

doi:10.1016/j.ajhg.2012.06.017

Valente, E. M., F. Brancati, and B. Dallapiccola, 2008 Genotypes and phenotypes of

Joubert syndrome and related disorders. *Eur J Med Genet.* **51:** 1-23.

doi:10.1016/j.ejmg.2007.11.003

Valente, E. M., C. V. Logan, S. Mougou-Zerelli, J. H. Lee, J. L. Silhavy et al., 2010

Mutations in TMEM216 perturb ciliogenesis and cause Joubert, Meckel and

related syndromes. *Nat Genet.* **42:** 619-625. doi:10.1038/ng.594

Vierkotten, J., R. Dildrop, T. Peters, B. Wang, and U. Rüther, 2007 Ftm is a novel basal

body protein of cilia involved in Shh signaling. *Development.* **134:** 2569-2577.

doi:10.1242/dev.003715

Weatherbee, S. D., L. A. Niswander, and K. V. Anderson, 2009 A mouse model for

Meckel syndrome reveals Mks1 is required for ciliogenesis and Hedgehog

signaling. *Hum Mol Genet.* **18:** 4565-4575. doi:10.1093/hmg/ddp422

Wechsler-Reya, R. J., and M. P. Scott, 1999 Control of neuronal precursor proliferation

in the cerebellum by Sonic Hedgehog. *Neuron.* **22:** 103-114. doi:10.1016/s0896-

6273(00)80682-0

Wu, C., M. Yang, J. Li, C. Wang, T. Cao et al., 2014 Talpid3-binding centrosomal

protein Cep120 is required for centriole duplication and proliferation of cerebellar

granule neuron progenitors. *PLoS One.* **9:** e107943.

doi:10.1371/journal.pone.0107943

Yachnis, A. T., and L. B. Rorke, 1999 Neuropathology of Joubert syndrome. *J Child*

Neurol. **14:** 655-659. doi:10.1177/088307389901401006

Zhang, X. M., M. Ramalho-Santos, and A. P. McMahon, 2001 Smoothened mutants

reveal redundant roles for Shh and Ihh signaling including regulation of L/R

symmetry by the mouse node. *Cell.* **106:** 781-792.

Zhu, J., H.-T. Wang, Y.-R. Chen, L.-Y. Yan, Y.-Y. Han et al., 2020 The Joubert

syndrome gene arl13b is critical for early cerebellar development in zebrafish.

Neurosci Bull. **36:** 1023-1034. doi:10.1007/s12264-020-00554-y

Figure legends

Figure 1: SCPs lacking *Arl13b* or *Smo* fail to project to the dorsal thalamus. (A-C)

Schematics of injections and fluorescent tracer diffusion shown horizontally (A) or

sagittally (B-C). (A) Red dashed arrow depicts dye path in a successful injection from

injection site (red X) caudal through the brain and across the midline and into the

contralateral cerebellar DCN (red arrowhead). Grey background boxes indicate area of

subsequent images: the injection site (INJ) and cerebellum (DCN). (D-I) Representative

images of dorsal (D, F, H) or ventral (E, G, I) thalamus injection site (top panel) and

cerebellum (middle panel) with the DCN in hatched white circle and magnified (bottom

panel) with recoloring to black and white to aid visualization. The retrograde fluorescent

tracer is pink-red and sections are stained with DAPI (blue). Numbers indicate the

number of positively stained DCN clusters (DCN traced) out of the total number of

injected animals. Note that no tracing was observed on ipsilateral side to injection. (D,

E) Fluorescent tracer injection in $Smo^{fl/+}$; $Arl13b^{fl/+}$; *Nex-Cre* control animals resulted in

contralateral DCN staining in (D) 6/8 dorsal thalamus injections and (E) 8/11 ventral

thalamus injections. (F, G) Fluorescent tracer injection in $Smo^{Nex-Cre}$ animals resulted in

contralateral DCN staining in (F) 0/6 dorsal thalamus injections and (G) 4/4 ventral

thalamus injections. (H, I) Fluorescent tracer injection in $Arl13b^{Nex-Cre}$ animals resulted in

contralateral DCN staining in (H) 1/7 dorsal thalamus injections and (I) 4/4 ventral

thalamus injections. Two-tailed Fisher's exact test was performed; only the dorsal

injection results were significantly different from the controls (F: $p<0.01$; H: $p<0.05$)

Figure 2: SCPs expressing cilia-excluded ARL13B^{V358A} project normally to both

the dorsal and ventral thalamus. (A-D) Representative images of dorsal (A, C) or

ventral (B, D) thalamus injection site (top panel) and cerebellum (middle panel) with the

DCN in hatched white circle and magnified (bottom panel) with recoloring to black and

white to aid visualization. Numbers indicate the number of positively stained DCN

clusters out of the total number of injected animals. Note that no tracing was observed

on the injection's ipsilateral side. (A, B) Fluorescent tracer injection in $Arl13b^{V358A/+}$

control animals resulted in contralateral DCN staining in (A) 3/3 dorsal thalamus

injections and (B) 4/4 ventral thalamus injections. (C-D) Fluorescent tracer injection in

$Arl13b^{V358A/V358A}$ animals resulted in contralateral DCN staining in (C) 4/4 dorsal

thalamus injections (Fisher's exact test, two-tailed, ns, $p>0.9$) and (D) 3/3 ventral

thalamus injections (ns, $p>0.9$).

Figure 3: SCPs expressing JS allele *$Arl13b^{R79Q}$* project normally to both the dorsal

and ventral thalamus. (A) Schematic showing the protein domain structure of Arl13b.

The R79Q mutation occurs in a highly conserved subregion of the GTPase domain,

Switch 2 (red). (B) Alignment of protein sequence surrounding amino acid 79 of mouse

Arl13b (red highlight) showing that arginine is conserved across multiple species.

Sequences used in protein alignment: *Homo sapiens* NP_878899.1, *Mus musculus*

NP_080853.3, *Danio rerio* NP_775379.1, *Chlamydomonas reinhardtii* XP_001691430.1.

(C-F) Representative images of dorsal (C, E) or ventral (D, F) thalamus injection site

(top panel) and cerebellum (middle panel) with the DCN in hatched white circle and

magnified (bottom panel) with recoloring to black and white to aid visualization.

Numbers indicate the number of positively stained DCN clusters out of the total number

of injected animals. Note that no tracing was observed on the injection's ipsilateral side.

(C, D) Fluorescent tracer injection in *Arl13b$^{R79Q/+}$* control animals resulted in

contralateral DCN staining in (C) 4/4 dorsal thalamus injections and (D) 3/3 ventral

thalamus injections. (E-F) Fluorescent tracer injection in *Arl13b$^{R79Q/R79Q}$* animals

resulted in contralateral DCN staining in (E) 3/3 dorsal thalamus injections (Fisher's

exact test, two-tailed, ns, p>0.9) and (F) 3/4 ventral thalamus injections (ns, p>0.9).

Figure 4: Mouse embryos expressing JS allele *Arl13b^{R79Q}* display normal cell

patterning in the neural tube. Shh, Olig2, Pax6 and Arl13b staining of E10.5 *Arl13b$^{+/+}$*

(n=3)*, Arl13b$^{R79Q/R79Q}$* (n=3)*, Arl13b$^{R79Q/\Delta}$* (n=3), and *Arl13b$^{\Delta/\Delta}$* (n=3) mouse neural tubes.

Scale bar = 100 micrometers in A-L or 10 micrometers in M-P. (A-D) Shh is visible in the

notochord and floorplate of (A) *Arl13b$^{+/+}$,* (B) *Arl13b$^{R79Q/R79Q}$* and (C) *Arl13b$^{R79Q/\Delta}$* neural

tubes but absent from the floorplate of (D) *Arl13b$^{\Delta/\Delta}$* embryos. (E-H) Olig2 stains the

motor neuron precursor domain in (E) *Arl13b$^{+/+}$*, (F) *Arl13b$^{R79Q/R79Q}$* and (G) *Arl13b$^{R79Q/\Delta}$*

embryos and stains an expanded domain in (H) *Arl13b$^{\Delta/\Delta}$* embryos. (I-L) Pax6

expression is visible in the dorsal neural tube in (I) *Arl13b$^{+/+}$*, (J) *Arl13b$^{R79Q/R79Q}$* and (K)

Arl13b$^{R79Q/\Delta}$ neural tubes but shifted dorsally in (L) *Arl13b$^{\Delta/\Delta}$* neural tubes. (M-P) Arl13b

is localized to cilia visible in the ventral lumen of (M) *Arl13b$^{+/+}$,* (N) *Arl13b$^{R79Q/R79Q}$* and

(O) *Arl13b$^{R79Q/\Delta}$* neural tubes but absent from (P) *Arl13b$^{\Delta/\Delta}$* embryos.

Figure 5: *Arl13b^{R79Q}* mutation does not affect cerebellar width. (A) Representative

surface-facing images of cerebella from control and *Arl13b$^{R79Q/R79Q}$* mutant mice.

Scalebar = 1mm, black line indicates where cerebellar width was measured, red line

indicates where vermis width was measured, gray line indicates where cerebellar

hemisphere height was measured, blue line indicates where vermis height was

measured. (B) Ratios of $Arl13b^{R79Q/R79Q}$ mutant to control measurements of sex- and

age-matched pairs of mice showed no significant difference. Cerebellar and vermis

measurements were determined from surface views (symbol colors match the lines in

A), body and dissected brain weights were determined using a standard lab scale. Each

symbol represents a pair of sex- and age-matched animals: females are represented as

circles and males are represented as triangles.

Figure 6: Pan-neuronal deletion of *Arl13b* results in a small cerebellum. (A)

Representative images of cerebella from control and $Arl13b^{Brn4-Cre}$ mutant mice.

Scalebar = 1mm. (B) The ratio of $Arl13b^{Brn4-Cre}$ mutant to control cerebellar width was

reduced in both females and males (one-sample t-test: $p<0.1$, single asterisk). The ratio

of $Arl13b^{Brn4-Cre}$ mutant to control vermis width was reduced in female and male age

matched pairs (one-sample t-test: $p<0.0001$, four asterisks). While $Arl13b^{Brn4-Cre}$ mutants

had moderately lower body weights than controls (one-sample t-test: $p<0.1$, single

asterisk), their brains were slightly heavier, likely due to hydrocephaly. Each symbol

represents a pair of sex- and age-matched animals: females are represented as circles

and males are represented as triangles.

Figure 7: Mice expressing cilia-excluded *Arl13b*^{V358A} have cerebella of normal

width. (A) Representative images of cerebella from control and $Arl13b^{V358A/V358A}$ mutant

mice. Scalebar = 1mm. (B) Ratios of $Arl13b^{V358A/V358A}$ mutant to control measurements

of sex- and age-matched pairs of mice showed no significant differences (one-sample t-

test). Each dot represents a pair of sex- and age-matched animals: females are

represented as circles and males are represented as triangles.

Dorsal injection

Ventral injection

A
A'
INJ
A''
DCN

B

C

D Control
INJ
DCN
DCN traced: 6/8

E
INJ
DCN
DCN traced: 8/11

F *Smo^Nex-Cre*
INJ
DCN
0/6

G
INJ
DCN
4/4

H *Arl13b^Nex-Cre*
INJ
DCN
1/7

I
INJ
DCN
4/4

A *Arl13b^{V358A/+}*

Dorsal injection

INJ

DCN

DCN
traced: 3/3

C *Arl13b^{V358A/V358A}*

INJ

DCN

4/4

B

Ventral injection

INJ

DCN

DCN
traced: 4/4

D

INJ

DCN

3/3

A

Switch 1   Switch 2   coiled-coil   Pro-rich
N   C
                    GTPase
              R79Q   V358A

B

	Switch 1	Switch 2	
Human	PTVGFS·KINLRQGKFEVTI·FDLGGGIRIRGIWKNYYAE·SYGVIFVVDSSD		100
Mouse	PTVGFS·KIDLRQGKFQVTI·FDLGGGKRIRGIWKNYYAE·SYGVIFVVDSSD		100
Zebrafish	PTVGFS·KVDLKQGKFEVTI·FDLGGGKRIRGIWKNYYSE·SYGVVFVVDSSD		100
Chlamy	PTFGFN·STTLNEGKYKIEV·FDLGGGKNIRGVWKKYLAE·VHAIVYVVDAAD		98

R79Q

Arl13b^{R79Q/+}

Dorsal injection

INJ

DCN

DCN traced: 4/4

Arl13b^{R79Q/R79Q}

INJ

DCN

3/3

Ventral injection

INJ

DCN

DCN traced: 3/3

INJ

DCN

3/4

A Control   *Arl13b$^{R79Q/R79Q}$*

B

Arl13b^{R79Q}

Ratio (mutant/control)

Cerebellar width   Vermis width   Hemisphere height   Vermis height   Body weight   Brain weight

Circle = female
Triangle = male

A

Control   *Arl13b^{Brn4-Cre}*

B

Arl13b^{Brn4-Cre}

Circle = female
Triangle = male

A

Control   *Arl13b^{V358A/V358A}*

B

Arl13b^{V358A}

Ratio (mutant/control)

1.5

1.0

0.5

Cerebellar width | Vermis width | Hemisphere height | Vermis height | Body weight | Brain weight

Circle = female
Triangle = male