Nrg1/ErbB Signaling-Mediated Regulation of Fibrosis After Myocardial Infarction

Manabu Shiraishi MD, PhD,[1,2] Atsushi Yamaguchi MD, PhD,[1] Ken Suzuki MD, PhD[2]

[1]Department of Cardiovascular Surgery, Saitama Medical Center, Jichi Medical University,

Saitama 330-0834, Japan.

[2]William Harvey Research Institute, Barts and The London School of Medicine and Dentistry,

Queen Mary University of London, London EC1M 6BQ, United Kingdom.

Short title: Regulatory mechanism of fibrosis after MI

Address correspondence to:

Manabu Shiraishi, Department of Cardiovascular Surgery, Saitama Medical Center, Jichi

Medical University, Saitama 330-0834, Japan.

Tel: +81-48-647-2111; E-mail: manabu@omiya.jichi.ac.jp

Word count: 7526

Subject Terms

Cell signaling

Fibrosis

Ischemia

Myocardial Infarction

Abstract

RATIONALE: Appropriate fibrotic tissue formation after myocardial infarction (MI) is crucial to maintenance of the heart's structure. Reparative or M2-like macrophages play a vital role in post-MI fibrosis by activating cardiac fibroblasts. However, the mechanism by which post-MI cardiac fibrosis is regulated is not fully understood.

OBJECTIVE: We investigated the cellular and molecular mechanisms of post-MI fibrotic tissue formation, especially those related to regulation of cellular senescence and apoptosis.

METHODS AND RESULTS: *In vivo* and *in vitro* experiments were used to investigate the molecular and cellular mechanisms through which post-MI fibrosis occurs, with a focus on the role of M2-like macrophages. Microarray analysis revealed that $CD206^+F4/80^+CD11b^+$ M2-like macrophages collected from mouse hearts on post-MI day 7 showed increased expression of neuregulin 1 (Nrg1). Nrg1 receptor epidermal growth factor receptor ErbB was expressed on cardiac fibroblasts in the infarct area. In cardiac fibroblasts in which hydrogen peroxide-induced senescence, M2-like macrophage-derived Nrg1 suppressed both senescence and apoptosis of the fibroblasts, whereas blockade of ErbB function significantly accelerated. M2-like macrophage-derived Nrg1/ErbB/PI3K/Akt signaling, which was shown to be related to anti-senescence, was activated in damaged cardiac fibroblasts. Interestingly, systemic blockade of ErbB function in MI model mice enhanced senescence and apoptosis of cardiac fibroblasts and exacerbated inflammation. Further, increased accumulation of M2-like macrophages resulted in excessive progression of fibrosis in post-MI murine hearts. The molecular mechanism underlying regulation of fibrotic tissue formation in the infarcted myocardium was shown in part to be attenuation of apoptosis and senescence of cardiac fibroblasts by activation of Nrg1/ErbB/PI3K/Akt signaling.

CONCLUSIONS: M2-like macrophage-mediated regulation of Nrg1/ErbB signaling, ~~shown to~~

have a substantial effect on fibrotic tissue formation in the infarcted adult mouse heart, is critical

for suppressing the progression of senescence and apoptosis of cardiac fibroblasts.

Key Words:

apoptosis; cardiac fibroblast; fibrosis; macrophage; myocardial infarction; senescence

Nonstandard Abbreviations and Acronyms

Akt protein kinase B

αSMA alpha-smooth muscle actin

BMDM bone marrow-derived macrophage

Cdk cyclin-dependent kinase

ErbB epidermal growth factor receptor

GEO Gene Expression Omnibus

Glb1 galactosidase beta 1

H_2O_2 hydrogen peroxide

IL-6 interleukin-6

MDM2 murine double minute 2

MI myocardial infarction

mTOR mechanistic target of rapamycin

M2 M2-like macrophage

Nrg1 neuregulin 1

Pdgfa platelet derived growth factor subunit A

PI3K phosphatidylinositol 3-kinase

qRT-PCR quantitative reverse transcription-polymerase chain reaction

SA-β-gal senescence-associated β-galactosidase

SASP senescence-associated secretory phenotype

Spp1 secreted phosphoprotein 1

Tgfb transforming growth factor beta

Thy1 thymocyte antigen 1

INTRODUCTION

Myocardial infarction (MI) is a leading cause of mortality and disability worldwide. Heart failure

is common among survivors of acute MI, resulting from the adverse ventricular remodeling that

follows MI.[1,2] Because the regenerative capacity of the human heart does not fully compensate

for the loss of cardiomyocytes that occurs with MI, formation of connective tissue is essential to

maintenance of the integrity and rigidity of the heart. However, the mechanism by which cardiac

fibrosis is regulated after MI is not fully understood.

MI results in permanent loss of hundreds of millions of cardiomyocytes.[3] Studies have shown

that in the infarct area even non-cardiomyocytes, including fibroblasts, disappear in large

quantities by apoptosis and that cellular senescence-associated defects occur in cardiac repair

after MI.[4,5] Apoptosis plays an important role in the disappearance of infiltrated immune cells

and cardiac interstitial cells after MI.[5] Because senescence and apoptosis of both cardiomyocytes

and fibroblasts are deeply involved in the pathophysiology of adverse left ventricular remodeling

and myocardial rupture after MI,[6] understanding the molecular mechanisms by which senescence

and apoptosis are regulated during the post-MI tissue repair process is important. Cellular

senescence and apoptosis are processes of growth arrest that occur with age and in response to

cellular stress and damage, and they limit proliferation of cells.[7,8] Senescent cells possess a

complex phenotype and are characterized by cell cycle arrest mediated via p16 and p53/p21

pathways, with some ultimately manifesting a unique secretory phenotype known as the

senescence-associated secretory phenotype (SASP).[9] Cell cycle arrest plays a central role in the

senescent phenotype of adult cardiomyocytes, and induction of cell cycle reentry of adult

cardiomyocytes promotes post-MI cardiac repair.[10] Administration of anti-apoptotic substances

to rats after MI has been shown to suppress cardiomyocyte apoptosis, which in turn decreases

infarct size and ameliorates the cardiac dysfunction that typically follows MI.[11] Even non-

cardiomyocytes in the infarct area, including fibroblasts, undergo apoptosis.[4,5] Therefore,

senescence and/or apoptosis of cardiac fibroblasts may be involved in the tissue repair process

that follows MI. Senescent cardiac fibroblasts, in which expression of the major senescence

regulator p53 is significantly upregulated, have been shown to accumulate markedly in the

mouse heart after MI, with the upregulation of p53 shown to limit cardiac collagen production,

and conversely, the inhibition of p53 shown to increase reparative fibrosis.[1] However, the

methods used in the aforementioned studies are limited in their ability to identify the particular

subset of immune cells involved in the post-MI fibrotic process as well as the inherent cellular

and molecular mechanisms by which fibroblast senescence and collagen production are

regulated. Whereas fibrosis after MI is crucial to maintaining myocardial structure, excessive

fibrosis leads to eventual heart failure. Therefore, achieving equilibrium between profibrotic and

anti-fibrotic environments is important for a good regenerative outcome. For an understanding of

the fibrosis-based tissue repair process, intercellular communication between senescent and

apoptotic fibroblasts and surrounding immune cells must be clarified.

Investigators have shown macrophages to be essential for regeneration of the neonatal mouse

heart,[12] and we have shown that reparative or M2-like macrophages play a pivotal role in the

formation of fibrous tissue following MI by promoting proliferation and activation of cardiac

fibroblasts.[13] We hypothesized that M2-like macrophages play a vital role in attenuating

apoptosis and senescence of cardiac fibroblasts, both of which are profoundly involved in the

regulation of fibrotic processes after MI, and we investigated the molecular mechanism by which

this particular subset of immune cells effectuates this regulation.

Neuregulin 1 (*Nrg1*) is one of the neuregulin genes (*Nrg1–Nrg4*) belonging to the family of

epidermal growth factor genes,[14,15] and Nrg1/epidermal growth factor receptor (ErbB) signaling

systems play essential roles in the protection and proliferation of cardiomyocytes in response to

injury.[16-19] Although association between Nrg1 and protection of cardiomyocytes has been

studied over several decades,[16-19] the exact mechanism through which Nrg1 protects cardiac

fibroblasts and the mechanism through which Nrg1 participates in post-MI regeneration have not

been elucidated. A recent study showed that Nrg1 enhances proliferation and viability of normal

human ventricular cardiac fibroblasts, with the Nrg1 activity linked to Nrg1/ErbB signaling.[20]

Results of a study in a mouse model of myocardial hypertrophy indicated that Nrg1 has an anti-

fibrotic effect in the heart, generated by anti-inflammatory Nrg1/ErbB signaling in

macrophages.[21] Furthermore, Nrg1-loaded poly-microparticles were used in a later study to

induce macrophage polarization toward an anti-inflammatory phenotype, which prevented their

transition toward the inflammatory phenotype and enhanced cardiac repair after MI.[22] However,

whether M2-like macrophages function to ameliorate senescence and apoptosis of cardiac

fibroblasts through Nrg1/ErbB signaling activity *in vivo* under post-MI conditions is unknown.

Although the contribution of Nrg1/ErbB signaling in macrophages and macrophage polarization

toward an anti-inflammatory phenotype to cardiac tissue repair was investigated in these studies,

the significance of Nrg1/ErbB signaling activity in cardiac fibroblasts with respect to the anti-

fibrotic effect of Nrg1 has not been clarified. We established what we believe to be an

appropriate *in vitro* model for delineation of the precise roles of M2-like macrophage-derived

Nrg1 after MI in anti-senescence, anti-apoptosis, and the fibrotic phenotype of fibroblasts.

Moreover, we used this model and an *in vivo* model to investigate the molecular and cellular

mechanisms by which post-MI fibrosis occurs, with a focus on the role of M2-like macrophages.

We aimed to shed light on the pathophysiology of fibrosis after MI and provide information that

will contribute to ultimate development of new therapeutic strategies that exploit Nrg1-driven

senescence and apoptosis of cardiac fibroblasts.

METHODS

Data Availability

Microarray data: Gene Expression Omnibus GSE69879

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE69879)

Please see the Supplemental Materials for Methods and Major Resources.

RESULTS

Cardiac Fibroblasts Undergo Apoptosis and Senescence After MI

We investigated cellular senescence and apoptosis in a mouse MI model, which was established

by means of coronary artery ligation. Obvious fibrotic tissue formation and increased myocardial

expression of fibrosis-associated genes (*αSMA* [alpha-smooth muscle actin gene], *Col1a1*, and

Col3a1) were observed in the infarct area as soon as post-MI day 7 (Figure IA–B in the Data

Supplement). These changes were associated with an increase in the number of thymocyte

antigen 1 (Thy1)$^+$ fibroblasts and Thy1$^+$αSMA$^+$ myofibroblasts in the infarct area (Figure IC–D

in the Data Supplement), which peaked on post-MI day 7. Interestingly, approximately 40% of

Thy1$^+$ fibroblasts in the infarct area were positive for cleaved caspase 3 on post-MI day 7, and

approximately 15% were positive for cleaved caspase 3 on post-MI day 28, indicative of robust

apoptosis of cardiac fibroblasts (Figure 1A). Senescence-associated β-galactosidase (SA-β-gal)-

positive cells were also found in the infarct area. Numerous SA-β-gal-positive cells were spindle-

shaped and had many cytoplasmic processes and thus appeared to be fibroblasts (Figure 1B).

Additionally, myocardial expression of senescence-associated genes (*p16*, *p2*, and *Glb1* [beta 1

galactosidase gene])[1,7,8,23,24] and inflammation-associated genes (*CCL3*, *IL-6*, and *TNF*) was

upregulated in the infarct area in comparison to that in the non-infarct (remote) area (Figure 1C

and IE in the Data Supplement). Importantly, myocardial expression of reparative gene *Nrg1* was

upregulated in the infarct area, peaking on post-MI day 7 (Figure 1D). Immunohistochemistry

(IHC) showed that most Thy1[+] cardiac fibroblasts that had accumulated in both the infarct and

remote areas expressed ErbB2 and ErbB4, which are Nrg1 coreceptors[19,25,26] (Figure 1E). Taken

together, these results indicate that apoptosis and senescence of cardiac fibroblasts occur during

fibrotic tissue formation in the post-MI heart and that cardiac fibroblasts express 2 Nrg1

receptors, ErbB2 and ErbB4, in response to ischemic injury.

M2-Like Macrophages Accumulate in the Infarct Area and Express Nrg1

IHC showed that CD206[+] M2-like macrophages had accumulated in the infarct area, with peak

accumulation observed on post-MI day 7 (Figure 2A). Interestingly, this change in M2-like

macrophages corresponded to the change that occurred in cardiac fibroblasts after MI (Figure IC

in the Data Supplement). We further confirmed that CD206[+] cells were also positive for F4/80 in

both normal and infarcted hearts (Figure 2B). Microarray analysis showed that the molecular

signature of CD206[+]F4/80[+]CD11b[+] M2-like macrophages collected from infarcted hearts on

post-MI day 7 differed from that of M2-like macrophages collected from normal hearts (Figure

IIA in the Data Supplement). A range of anti-inflammatory and reparative genes was upregulated

in M2-like macrophages in comparison to expression of the same genes in M2-like macrophages

obtained from normal hearts (Figure IIB in the Data Supplement; the full dataset is available in

the Gene Expression Omnibus [GEO] database; GSE69879). Importantly, gene ontology analysis

showed association between M2-like macrophages obtained from infarcted hearts and regulation

of apoptosis and cell death (Figure IIC in the Data Supplement). By searching HomoloGene

(https://www.ncbi.nlm.nih.gov/homologene) for genes related to proliferation and viability, we

focused on the increased expression level of *Nrg1* in M2-like macrophages obtained from

infarcted hearts (Figure 2C), which was confirmed by quantitative reverse transcription-

polymerase chain reaction (qRT-PCR) and IHC (Figure 2D–E). The data indicate that M2-like

macrophages are a source of Nrg1 after MI.

Bone Marrow-Derived Macrophages Attenuate Hydrogen Peroxide (H_2O_2)-Induced

Apoptosis and Senescence of Cardiac Fibroblasts Via Nrg1 Secretion

We investigated the role of macrophages in regulating senescence and apoptosis of fibroblasts in

an *in vitro* coculture model using a Boyden Chamber system in which cells can be independently

cultured or genetically analyzed without being mixed together[13,27] (Figure IIIA in the Data

Supplement). Hydrogen peroxide (H_2O_2) was used to induce senescence of fibroblasts (Figure

IIIB–C in the Data Supplement). Bone marrow-derived macrophages (BMDMs) that were

cocultured with the senescent fibroblasts were of an M2-like macrophage phenotype (Figure

IVA–B in the Data Supplement). As observed *in vivo* after MI (Figure 2C–E and Figure 1E),

increased expression of *Nrg1* was seen in BMDMs (Figure VA in the Data Supplement) and

increased expression of *ErbB2* and *ErbB4* was seen in cardiac fibroblasts (Figure VB–C in the

Data Supplement) 48 hours after the start of coculture. Phase-contrast microscopy (Figure 3A)

showed that fibroblasts treated with H_2O_2 had an enlarged, flattened, senescent morphology, but

after coculture with BMDMs the fibroblasts took on a healthy, spindle-shaped form. Importantly,

addition of anti-ErbB antibody (Ab), which is a competitive inhibitor of Nrg1, changed the gross

morphology of fibroblasts such that they resembled the fibroblasts treated with H_2O_2. Treatment

with recombinant Nrg1 restored the senescent fibroblasts to a healthy spindle-shaped form.

Approximately 20% of fibroblasts treated with H_2O_2 were found to be positive for senescence-

associated β-galactosidase (SA-β-gal) (Figure 3B). This change was significantly attenuated by

coculture with BMDMs, whereas addition of anti-ErbB Ab eliminated the anti-senescence effect

of the BMDMs. Furthermore, treatment with recombinant Nrg1 suppressed senescence.

Expression of apoptosis-related cleaved caspase 3 (Figure 3C) was similar to that of SA-β-gal.

Conversely, expression of proliferation-related Ki-67 was significantly attenuated by H_2O_2

treatment and restored by co-culture with BMDMs. Effects of the BMDMs were diminished by

anti-ErbB Ab, whereas recombinant Nrg1 improved H_2O_2-induced functional deterioration.

Together, these results show that Nrg1 derived from BMDMs reduces both apoptosis and

senescence of cultured fibroblasts and promotes their proliferation.

BMDMs Promote Activation of Fibroblasts and Collagen Synthesis

Immunocytological staining showed that, although treatment with H_2O_2 alone did not affect

conversion of cardiac fibroblasts into αSMA⁺ myofibroblasts, the phenotypic change progressed

in coculture of cardiac fibroblasts with BMDMs. Furthermore, addition of anti-ErbB Ab

enhanced this BMDM-induced activation of fibroblasts. Addition of recombinant Nrg1 did not

affect conversion of cardiac fibroblasts (Figure 4A). Changes in synthesis of types I and III

collagen in fibroblasts in response to H_2O_2, BMDMs, and recombinant Nrg1 were similar to

changes in αSMA expression (Figure 4B–C). These results suggest that BMDMs, which are of

an M2-like phenotype (Figure IVA–B in the Data Supplement), induce activation of fibroblasts

for conversion to myofibroblasts and progression of collagen synthesis. The important take-away

point to be drawn from these results is that that cellular senescence and apoptosis alone do not

induce activation of fibroblasts for conversion to myofibroblasts and progression of collagen

synthesis; the presence of BMDMs is required. Osteopontin (secreted phosphoprotein 1 [SPP1])

is a major mediator of M2-like macrophage-induced cardiac fibroblast activation.[13] *Spp1*

expression was found to be increased in BMDMs in response to H_2O_2 (Figure VIA in the Data

Supplement). Conversely, other profibrotic factors, including transforming growth factor beta

and platelet derived growth factor subunit A,[28,29] were found not to be upregulated in M2-like

macrophages (Figure VIB in the Data Supplement).

Phosphatidylinositol 3-Kinase/Protein Kinase B Signaling is Associated with BMDM-

Attenuated Apoptosis and Senescence of Cardiac Fibroblasts Through Nrg1

We next investigated the molecular mechanism underlying attenuation of apoptosis and

senescence of cardiac fibroblasts by BMDM-derived Nrg1 in an *in vitro* setting. The

phosphatidylinositol 3-kinase (PI3K)/protein kinase B (Akt) signaling pathway is downstream of

the ErbB pathway. Western blot analysis showed that H_2O_2 treatment suppressed PI3K/Akt

activation in fibroblasts (Figure 5A). Coculture with BMDMs and treatment with recombinant

Nrg1upregulated the signaling activity, whereas addition of anti-ErbB Ab suppressed the

signaling activity. To determine the possible relation between senescence-associated p21/p16 and

Nrg1/ErbB/PI3K/Akt signaling pathways, we examined mRNA expression levels of senescence-

associated genes (*p21*, *p16*, and *Glb1)* (Figure 5B), cell cycle-associated genes (cyclin-dependent

kinase 4 [*Cdk4*], cyclin-dependent kinase 6 [*Cdk6*], cyclin-dependent kinase 2 [*Cdk2*], and *Ki-*

67) (Figure 5C), a p53 suppressor gene (murine double minute 2 [*MDM2*]) (Figure 5D), a cell

survival-associated gene (mechanistic target of rapamycin [*mTOR*]) (Figure 5E), and an SASP-

associated gene (interleukin-6 [*IL-6*]) (Figure 5F). Expression levels of senescence-associated

genes (*p16* and *p21*) and the SASP-associated gene (*IL-6*) were significantly higher and

expression levels of cell cycle-associated genes (*Cdk4*, *Cdk6*, *Cdk2*, and *Ki-67*) were markedly

lower in fibroblasts treated with H_2O_2 than in non-treated (control) fibroblasts. Importantly, these

changes in expression were reversed by coculture with BMDMs. Anti-ErbB Ab exacerbated

senescence and cell cycle arrest, whereas recombinant Nrg1 suppressed senescence and

enhanced proliferation. BMDMs enhanced or restored expression of p53 suppressor-related

MDM2 and cell survival-related *mTOR*. These results indicate that the PI3K/Akt signaling

pathway activation in cardiac fibroblasts operates downstream of the BMDM-derived Nrg1/ErbB

system and has a suppressive effect on cell cycle arrest and senescence. This signaling activity is

also likely to increase cell proliferation and survival (Figure 5G).

In Vivo Inhibition of Nrg1 Signaling Exacerbates Fibrosis

We used trastuzumab to clarify the role of Nrg1 in suppressing post-MI senescence and apoptosis

of cardiac fibroblasts *in vivo*. Trastuzumab is an anti-ErbB type 2 (ErbB2) monoclonal Ab that

binds to the extracellular juxtamembrane domain of ErbB2.[30,31] On the basis of the *in vitro*

coculture model, we hypothesized that trastuzumab administration would eliminate the anti-

apoptotic and anti-senescence effects of Nrg1 and therefore allow senescence of cardiac

fibroblasts to proceed, which would result in excessive fibrosis in the infarcted myocardium.

Intraperitoneal trastuzumab injections did not affect mRNA expression levels of genes associated

with senescence, the SASP, or activation of fibroblasts to myofibroblasts in the normal mouse

heart (Figure VIIA–C in the Data Supplement). We induced MI in mice surgically and then

injected either trastuzumab or vehicle intraperitoneally (Figure VIIIA in the Data Supplement).

Trastuzumab did not affect post-MI mortality or body weight (Figure VIIIB–C in the Data

Supplement). Gene expression profiles in cardiac fibroblasts resident in infarcted myocardium

suggested that senescence was augmented by trastuzumab administration in the infarct area

(Figure 6A). The changes in gene expression corresponded to increases in the number of

senescent cardiac fibroblasts and cardiac cells in the infarct area (Figure 6B–C). Additionally,

trastuzumab administration increased the proportion of fibrotic tissue in the infarct area (Figure

7A). In fact, the numbers of Thy-1$^+$ fibroblasts and αSMA$^+$Thy1$^+$ myofibroblasts were increased

in association with upregulation of *αSMA*, *Col1a1*, and *Col3a*1 in the infarct area (Figure 7B–C).

Considering these *in vitro* experimental results, increased inflammation (Figure 8A) and boosted

accumulation of M2-like macrophages (Figure 8B) in this area by ErbB2 blockade may explain

the increased fibroblast activation and augmented fibrosis. Interestingly, trastuzumab

administration induced expression of apoptosis- and senescence-associated genes even in the

remote area (Figure IXA–C in the Data Supplement) where there was little or no direct ischemic

damage after the MI. This corresponded to the increased inflammation (Figure XA in the Data

Supplement) and the augmented accumulation of both M2-like macrophages and cardiac

fibroblasts in the remote myocardium (Figure XB and Figure XIA in the Data Supplement).

Conversion of fibroblasts to myofibroblasts and expression of fibrosis-associated genes were

also induced (Figure XI–B in the Data Supplement). Consequently, acute exacerbation of fibrosis

occurred even in the remote area (Figure XII in the Data Supplement). Thus, administration of

trastuzumab augmented apoptosis and senescence of cardiac fibroblasts following MI, resulting

in excessive fibrosis, even in the remote non-infarct area.

DISCUSSION

Despite the clinical importance of cellular and molecular processes underlying post-MI fibrosis, the processes are not well understood. To precisely determine the roles of cardiac M2-like macrophages in apoptosis and senescence of fibroblasts, we established an *in vitro* model in which senescent cardiac fibroblasts[32] were cocultured with BMDMs. This *in vitro* coculture model reflects the *in vivo* post-MI conditions because, under pathological conditions, senescent cells attract macrophages.[33] Expression of both *Cdk4/6* and *Cdk2* should be stable or increased in senescent cells, but expression levels were decreased in the cardiac fibroblasts in which senescence was induced by H_2O_2. The mechanism is expected to involve increased expression of *p16* and *p21*, which downregulate *Cdk4/6* and *Cdk2* expression, respectively.[8,34,35] Conversely, the addition of BMDMs decreased *p16* and *p21* expression and promoted the cell cycle (as evidenced by increased expression of *Cdk 4/6* and *Cdk2*). The hypertrophic response to Nrg1 is mainly dependent on Ras, whereas the anti-apoptotic and cell proliferation effects are likely dependent on Akt.[36,37] Therefore, Nrg1 may attenuate expression of senescence-associated genes *p21*, *p16*, and *Glb1* through PI3K/Akt pathways (Figure 5G). We observed a transient increase in *p21*, *p16*, and *Glb1* expression in the infarcted myocardium and in cultured fibroblasts. Stress-induced *p53* expression increases *p21* expression in response to DNA damage and induces reversible proliferative arrest that provides time for DNA repair and facilitates survival of cells.[38,39] p21, which binds to and inhibits CDK2, is important for initiation of senescence in some settings, but *p21* expression does not persist in senescent cells.[8,34,40,41] Increased *p16* expression is found in infarcted tissue. Irreversible proliferative arrest can be induced by p16, which inhibits CDK4 and CDK6.[8,35] Therefore, the change in expression levels of these genes

(i.e., *p21* and *p16*) in both the *in vivo* and *in vitro* experiments suggests that Nrg1 is a crucial

factor that controls reversible and irreversible senescence of cardiac cells after MI. Another

possible mechanism for the temporal change in *p21* and *p16* expression in the *in vivo* experiment

is as follows: There could be multiple types of *p21*- and *p16*-positive cells. Therefore, whereas

some *p21*- and *p16*-positive cell populations disappear, other *p21*- and *p16*-positive cells may

remain. We showed a possible mechanism linking M2-like macrophages and cardiac fibroblasts,

which contributes to various processes that are critical to many aspects of cellular function,

including cell growth and survival.[42] Together, results of our *in vivo* and *in vitro* experiments

suggest that macrophage-derived Nrg1 suppresses senescence and promotes proliferation of

cardiac fibroblasts after myocardial injury by activating the ErbB/PI3K/Akt signaling pathway.

ErbB is expressed not only on the surface of fibroblasts but also on the surface of

macrophages, and myeloid-specific *ErbB* deletion exacerbates myocardial fibrosis.[21]

Furthermore, Nrg1-induced macrophage polarization from an inflammatory phenotype toward an

anti-inflammatory phenotype enhances cardiac repair after MI.[22] Therefore, the question was

raised whether ErbB signaling in BMDMs was simultaneously affected when anti-ErbB Ab was

added to the culture medium in our coculture experiments. The exacerbated phenotypic changes

characterizing senescence and apoptosis of fibroblasts might not be due to inhibition of

Nrg1/ErbB signaling in the fibroblasts. However, we showed that the phenotypic changes and

expression of genes denoting the senescence and apoptosis of H_2O_2-treated cells, with the

Nrg1/ErbB signaling-induced activation of BMDMs having been excluded, were similar to those

in the H_2O_2+BMDM+anti-ErbB Ab treated cells (Figure 3A–C and Figure 5B–C). Therefore, our

data suggest that Nrg1/ErbB signaling activity has a greater anti-senescence and anti-apoptotic

effect in fibroblasts than does the signaling activity related to macrophage polarization.

Nrg1 is a cytokine that belongs to a family of proteins structurally related to epidermal growth

factor, and it plays essential roles in protection and proliferation of cardiomyocytes in response

to injury.[16-19] Injecting Nrg1 into adult mice induces cardiomyocyte cell cycle activity and

promotes myocardial regeneration, leading to improved function after MI.[16,18] Nrg1 also

significantly decreases apoptosis of adult cardiomyocytes under hypoxia-reoxygenation

conditions.[17] However, its function in cardiac fibroblasts in this context has not been well

studied. We present new *in vivo* and *in vitro* evidence suggesting that M2-like macrophages play

a vital role in attenuating senescence and apoptosis of fibroblasts through Nrg1/ErbB/PI3K/Akt

signaling pathways. This inherent reparative function allows senescent cardiac fibroblasts to

recover to a certain degree. Conversely, incomplete rescue of fibroblasts from senescence might

lead to undesired fibrosis.

Previous studies have shown that trastuzumab efficiently stops or slows the growth of ErbB2$^+$

cells *in vitro* and inhibits the ability of cells to repair damaged DNA.[30,31] As inflammation

worsens after MI, M2-like macrophages from bone marrow accumulate at the site of damaged

tissue.[43] Our study showed that progression of senescence and apoptosis in cardiac fibroblasts

and exacerbation of inflammation induced by trastuzumab increased the accumulation of M2-

like macrophages, which in turn promoted activation of fibroblasts and excessive fibrosis. These

results are consistent with our previous report indicating that interleukin 4-mediated M2-like

macrophage activation induces conversion of fibroblasts into myofibroblasts for progression of

fibrosis.[13] Although we analyzed mRNA expression levels in sections of heart tissue and not in

single cells, increased expression levels of senescence-associated genes in the trastuzumab-

treated samples were considered to reflect progression of senescence. These results corresponded

to results obtained *in vitro* with use of anti-ErbB Ab. We administered trastuzumab systemically,

which has unclarified cardiotoxicity, to explore ErbB signaling in cardiac fibroblasts. Because

cardiotoxicity of trastuzumab is non-specific, our study was limited by the fact that the ErbB

receptor block achieved with anti-ErbB Ab is not specific to cardiac fibroblasts. Moreover, M2-

like macrophages are not the only source of Nrg1 after MI. Apart from the M2-like macrophages

we studied, endothelial cells have been identified as a potential cell type for neuregulin

production and are responsible for the healing response after myocardial injury.[16-19] Therefore,

further mechanistic studies are needed to clarify the signaling pathways downstream of these

factors and the clear role of Nrg1, i.e., studies that incorporate cardiac fibroblast-specific

conditional *ErbB2/ErbB4*-knockout mice and macrophage-specific conditional *Nrg1*-knockout

mice.

Interestingly, senescence and SASP-associated gene expression peaked a little later in the

remote area than in the infarct area. One possible pathological mechanism of the remote non-

infarct area is assumed to be indirect damage via SASP rather than direct cytotoxicity due to

ischemia. Senescent cells autonomously induce senescence-like gene expression in surrounding

non-senescent cells through the SASP.[44] Increased release of SASP-associated substances in the

infarct area might not simply induce senescence and apoptosis of cells in the infarct area; it

might also have a harmful influence on non-senescent cells in the remote non-infarct area.

Although this study focused on Nrg1-induced anti-apoptosis and anti-senescence of cardiac

fibroblasts, M2-like macrophages are likely to mediate supplementary benefits for cardiac repair

after MI. These benefits may include reduced inflammation, activation of fibroblasts, and

neovascularization, as shown in our previous study.[13] To develop potential therapies mediated by

M2-like macrophages, studies are needed to determine how the gene that encodes Nrg1 is

switched on by MI and to identify other molecules that regulate apoptosis, senescence, and

proliferation of cardiac fibroblasts.

In conclusion, our data provide evidence that the Nrg1/ErbB/PI3K/Akt signaling system

regulates senescence and apoptosis of cardiac fibroblasts in the infarcted adult murine heart. This

process might play a vital role in repair of the infarcted myocardium by regulating collagen

synthesis. Therefore, this tissue repair mechanism controls the degree of rigidity and contraction

of the infarcted heart, thereby determining the prognosis of MI. Better understanding of the

molecular mechanism at play in the healing process of MI and subsequent remodeling may

ultimately lead to new treatments. Targeted activation of M2-like macrophages might enhance

the endogenous repair mechanism in senescent cardiac fibroblasts and serve as an approach to

treatment for MI.

Acknowledgments

MS and KS conceived and designed the study. MS conducted most of the experiments and

acquired the data with support from AY and KS. All authors participated in the analysis and

interpretation of the data. MS and KS primarily wrote and edited the manuscript with input from

all other authors. We thank Ellen Knapp, PhD, and Mitchell Arico from Edanz Group (https://en-

author-services.edanz.com/ac) for editing a draft of this manuscript. We wish to thank Ms.

Wendy Alexander-Adams for her assistance in reporting our findings in English.

Sources of Funding

This project was funded by the Uehara Memorial Foundation, SENSHIN Medical Research

Foundation, a Grant-in-Aid for Scientific Research (C), the Mochida Memorial Foundation for

Medical and Pharmaceutical Research, Takeda Science Foundation, and Public Trust Surgery

Research Fund.

Disclosures

The authors have that no conflicts of interest to disclose.

SUPPLEMENTAL MATERIALS

Supplemental Methods

Online Figures I–XII

References

1. Zhu F, Li Y, Zhang J, Piao C, Liu T, Li HH, Du J. Senescent cardiac fibroblast is critical for cardiac fibrosis after myocardial infarction. *PLoS One*. 2013;8:e74535.

2. Dickstein K, Cohen-Solal A, Filippatos G, et al.; ESC Committee for Practice Guidelines (CPG). ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2008: the Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2008 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association of the ESC (HFA) and endorsed by the European Society of Intensive Care Medicine (ESICM). *Eur Heart J*. 2008;29:2388-2442.

3. Gemberling M, Karra R, Dickson AL, Poss KD. Nrg1 is an injury-induced cardiomyocyte mitogen for the endogenous heart regeneration program in zebrafish. *Elife*. 2015;4.

4. Gould KE, Taffet GE, Michael LH, et al. Heart failure and greater infarct expansion in middle-aged mice: a relevant model for postinfarction failure. *Am J Physiol Heart Circ Physiol*. 2002;282:H615-21.

5. Takemura G, Ohno M, Hayakawa Y, Misao J, Kanoh M, Ohno A, Uno Y, Minatoguchi S, Fujiwara T, Fujiwara H. Role of apoptosis in the disappearance of infiltrated and proliferated interstitial cells after myocardial infarction. *Circ Res*. 1998;82:1130-1138.

6. Shih H, Lee B, Lee RJ, Boyle AJ. The aging heart and post-infarction left ventricular remodeling. *J Am Coll Cardiol*. 2011;57:9-17.

7. Sharpless NE, Sherr CJ. Forging a signature of in vivo senescence. *Nat Rev Cancer*. 2015;15:397-408.

8. Munoz-Espin D, Serrano M. Cellular senescence: from physiology to pathology. *Nat Rev Mol Cell Biol*. 2014;15:482-496.

9.   Coppe JP, Desprez PY, Krtolica A, Campisi J. The senescence-associated secretory

phenotype: the dark side of tumor suppression. *Annu Rev Pathol*. 2010;5:99-118.

10. Alam P, Haile B, Arif M, et al. Inhibition of senescence-associated genes Rb1 and Meis2 in

adult cardiomyocytes results in cell cycle reentry and cardiac repair post-myocardial

infarction. *J Am Heart Assoc*. 2019;8:e012089.

11. Hayakawa K, Takemura G, Kanoh M, et al. Inhibition of granulation tissue cell apoptosis

during the subacute stage of myocardial infarction improves cardiac remodeling and

dysfunction at the chronic stage. *Circulation*. 2003;108:104-109.

12. Aurora AB, Porrello ER, Tan W, Mahmoud AI, Hill JA, Bassel-Duby R, Sadek HA, Olson

EN. Macrophages are required for neonatal heart regeneration. *J Clin Invest*.

2014;124:1382-1392.

13. Shiraishi M, Shintani Y, Shintani Y, Ishida H, Saba R, Yamaguchi A, Adachi H, Yashiro K,

Suzuki K. Alternatively activated macrophages determine repair of the infarcted adult

murine heart. *J Clin Invest*. 2016;126:2151-2166.

14. Meyer D, Yamaai T, Garratt A, Riethmacher-Sonnenberg E, Kane D, Theill LE, Birchmeier

C. Isoform-specific expression and function of neuregulin. *Development*. 1997;124:3575-

3586.

15. Fuller SJ, Sivarajah K, Sugden PH. ErbB receptors, their ligands, and the consequences of

their activation and inhibition in the myocardium. *J Mol Cell Cardiol*. 2008;44:831-854.

16. Bersell K, Arab S, Haring B, Kuhn B. Neuregulin1/ErbB4 signaling induces cardiomyocyte

proliferation and repair of heart injury. *Cell*. 2009;138:257-270.

17. Hedhli N, Huang Q, Kalinowski A, Palmeri M, Hu X, Russell RR, Russell KS.

Endothelium-derived neuregulin protects the heart against ischemic injury. *Circulation*.

2011;123:2254-2262.

18. Polizzotti BD, Ganapathy B, Walsh S, Choudhury S, Ammanamanchi N, Bennett DG, dos

Remedios CG, Haubner BJ, Penninger JM, Kuhn B. Neuregulin stimulation of

cardiomyocyte regeneration in mice and human myocardium reveals a therapeutic window.

Sci Transl Med. 2015;7:281ra45.

19. Lemmens K, Doggen K, De Keulenaer GW. Role of neuregulin-1/ErbB signaling in

cardiovascular physiology and disease: implications for therapy of heart failure.

Circulation. 2007;116:954-960.

20. Kirabo A, Ryzhov S, Gupte M, Sengsayadeth S, Gumina RJ, Sawyer DB, Galindo CL.

Neuregulin-1beta induces proliferation, survival and paracrine signaling in normal human

cardiac ventricular fibroblasts. *J Mol Cell Cardiol*. 2017;105:59-69.

21. Vermeulen Z, Hervent AS, Dugaucquier L, et al. Inhibitory actions of the NRG-1/ErbB4

pathway in macrophages during tissue fibrosis in the heart, skin, and lung. *Am J Physiol*

Heart Circ Physiol. 2017;313:H934-H945.

22. Pascual-Gil S, Abizanda G, Iglesias E, Garbayo E, Prosper F, Blanco-Prieto MJ. NRG1

PLGA MP locally induce macrophage polarisation toward a regenerative phenotype in the

heart after acute myocardial infarction. *J Drug Target*. 2019;27:573-581.

23. Krizhanovsky V, Yon M, Dickins RA, Hearn S, Simon J, Miething C, Yee H, Zender L,

Lowe SW. Senescence of activated stellate cells limits liver fibrosis. *Cell*. 2008;134:657-

667.

24. van Deursen JM. The role of senescent cells in ageing. *Nature*. 2014;509:439-446.

25.   Olayioye MA, Neve RM, Lane HA, Hynes NE. The ErbB signaling network: receptor

501   heterodimerization in development and cancer. *EMBO J*. 2000;19:3159-3167.

502   26.   Uray IP, Connelly JH, Thomazy V, Shipley GL, Vaughn WK, Frazier OH, Taegtmeyer H,

503   Davies PJ. Left ventricular unloading alters receptor tyrosine kinase expression in the

504   failing human heart. *J Heart Lung Transplant*. 2002;21:771-782.

505   27.   Suzuki T, Arumugam P, Sakagami T, et al. Pulmonary macrophage transplantation therapy.

506   *Nature*. 2014;514:450-454.

507   28.   van den Borne SW, Diez J, Blankesteijn WM, Verjans J, Hofstra L, Narula J. Myocardial

508   remodeling after infarction: the role of myofibroblasts. *Nat Rev Cardiol*. 2010;7:30-7.

509   29.   Shinde AV, Frangogiannis NG. Fibroblasts in myocardial infarction: a role in inflammation

510   and repair. *J Mol Cell Cardiol*. 2014;70:74-82.

511   30.   Park S, Jiang Z, Mortenson ED, et al. The therapeutic effect of anti-HER2/neu antibody

512   depends on both innate and adaptive immunity. *Cancer Cell*. 2010;18:160-170.

513   31.   ElZarrad MK, Mukhopadhyay P, Mohan N, Hao E, Dokmanovic M, Hirsch DS, Shen Y,

514   Pacher P, Wu WJ. Trastuzumab alters the expression of genes essential for cardiac function

515   and induces ultrastructural changes of cardiomyocytes in mice. *PLoS One*. 2013;8:e79543.

516   32.   Bladier C, Wolvetang EJ, Hutchinson P, de Haan JB, Kola I. Response of a primary human

517   fibroblast cell line to H2O2: senescence-like growth arrest or apoptosis? *Cell Growth Differ*.

518   1997;8:589-598.

519   33.   Sasaki M, Miyakoshi M, Sato Y, Nakanuma Y. Modulation of the microenvironment by

520   senescent biliary epithelial cells may be involved in the pathogenesis of primary biliary

521   cirrhosis. *J Hepatol*. 2010;53:318-325.

34. Childs BG, Durik M, Baker DJ, van Deursen JM. Cellular senescence in aging and age-

related disease: from mechanisms to therapy. *Nat Med*. 2015;21:1424-435.

35. Serrano M, Lin AW, McCurrach ME, Beach D, Lowe SW. Oncogenic ras provokes

premature cell senescence associated with accumulation of p53 and p16INK4a. *Cell*.

1997;88:593-602.

36. Kuramochi Y, Cote GM, Guo X, Lebrasseur NK, Cui L, Liao R, Sawyer DB. Cardiac

endothelial cells regulate reactive oxygen species-induced cardiomyocyte apoptosis through

neuregulin-1beta/erbB4 signaling. *J Biol Chem*. 2004;279:51141-51147.

37. Baliga RR, Pimental DR, Zhao YY, Simmons WW, Marchionni MA, Sawyer DB, Kelly

RA. NRG-1-induced cardiomyocyte hypertrophy. Role of PI-3-kinase, p70(S6K), and

MEK-MAPK-RSK. *Am J Physiol*. 1999;277:H2026-H2037.

38. Deng C, Zhang P, Harper JW, Elledge SJ, Leder P. Mice lacking p21CIP1/WAF1 undergo

normal development, but are defective in G1 checkpoint control. *Cell*. 1995;82:675-684.

39. Wang YA, Elson A, Leder P. Loss of p21 increases sensitivity to ionizing radiation and

delays the onset of lymphoma in atm-deficient mice. *Proc Natl Acad Sci U S A*.

1997;94:14590-14595.

40. Alcorta DA, Xiong Y, Phelps D, Hannon G, Beach D, Barrett JC. Involvement of the cyclin-

dependent kinase inhibitor p16 (INK4a) in replicative senescence of normal human

fibroblasts. *Proc Natl Acad Sci U S A*. 1996;93:13742-7.

41. Stein GH, Drullinger LF, Soulard A, Dulic V. Differential roles for cyclin-dependent kinase

inhibitors p21 and p16 in the mechanisms of senescence and differentiation in human

fibroblasts. *Mol Cell Biol*. 1999;19:2109-2117.

42. Yu JS, Cui W. Proliferation, survival and metabolism: the role of PI3K/AKT/mTOR

signalling in pluripotency and cell fate determination. *Development*. 2016;143:3050-3060.

43. Ikeda N, Asano K, Kikuchi K, et al. Emergence of immunoregulatory Ym1(+)Ly6C(hi)

monocytes during recovery phase of tissue injury. *Sci Immunol*. 2018;3.eaat0207.

44. Acosta JC, Banito A, Wuestefeld T, et al. A complex secretory program orchestrated by the

inflammasome controls paracrine senescence. *Nat Cell Biol*. 2013;15:978-990.

45. Tano N, Narita T, Kaneko M, Ikebe C, Coppen SR, Campbell NG, Shiraishi M, Shintani Y,

Suzuki K. Epicardial placement of mesenchymal stromal cell-sheets for the treatment of

ischemic cardiomyopathy; in vivo proof-of-concept study. *Mol Ther*. 2014;22:1864-1871.

46. Leicht M, Greipel N, Zimmer H. Comitogenic effect of catecholamines on rat cardiac

fibroblasts in culture. *Cardiovasc Res*. 2000;48:274-284.

47. Shintani Y, Kapoor A, Kaneko M, et al. TLR9 mediates cellular protection by modulating

energy metabolism in cardiomyocytes and neurons. *Proc Natl Acad Sci U S A*.

2013;110:5109-14.

Figure legends

Figure 1. Myocardial infarction (MI) induces apoptosis and senescence of cardiac

fibroblasts and upregulates Nrg1 expression in infarcted myocardium.

A, Double immunofluorescence staining for Thy1 and cleaved caspase 3 (CC3), performed on

post-MI days 7 and 28, revealed a significantly increased number of apoptotic cardiac

fibroblasts in the infarct area, relative to the number seen in the remote area. Scale bars: 100

μm. *n=4* samples each.

B, Staining for senescence-associated beta-galactosidase (SA-β-gal) on post-MI days 7 and 28

revealed a significantly increased accumulation of spindle-shaped senescent cells in the

infarct area relative to that in the remote area. Scale bars: 100 μm.

C, Quantitative reverse transcription-polymerase chain reaction (qRT-PCR) revealed significant

upregulation of senescence-associated genes (*p16* and *Glb1*) in the infarct area on post-MI

day 7 or 28 or both. *n=4* samples each.

D, qRT-PCR analysis confirmed significant upregulation of *Nrg1* in the infarct area on post-MI

day 7. *n=4* samples each.

E, On immunohistochemistry, the Thy1$^+$ErbB2$^+$ fibroblast/Thy1$^+$ fibroblast ratio and the

Thy1$^+$ErbB4$^+$ fibroblast/Thy1$^+$ fibroblast ratio determined on post-MI day 7 and 28 were

found to be significantly increased. Scale bars: 100 μm. *n=4* samples each. Gene expression

levels relative to those in non-MI heart (Day 0) are given. Mean±SEM values are shown

*$P<0.05$, **$P<0.01$, ***$P<0.005$ versus the remote area; $^\#P<0.05$, $^{\#\#}P<0.01$, $^{\#\#\#}P<0.005$

versus non-MI heart; 1-way ANOVA.

Figure 2. M2-like macrophages accumulate in the infarct area and express Nrg1.

A, Immunohistochemistry revealed increased accumulation of CD206$^+$ M2-like macrophages in

the infarct area, which peaked on post-MI day 7. Scale bars: 100 μm. *n=4* samples each.

Mean±SEM values are shown.

*P<0.05, **P<0.01, ***P<0.005 versus the remote area; $^#P$<0.05, $^{##}P$<0.01, $^{###}P$<0.005

versus non-MI heart; 1-way ANOVA.

B, Flow cytometric analysis confirmed the presence of CD206$^+$F4/80$^+$CD11b$^+$ M2-like

macrophages in normal (non-MI) hearts of adult C57BL/6 mice and in hearts of mice

subjected to MI. *n=6* samples each.

C, Microarray analysis showed the M2 macrophages after MI to have a different expression

profile from that of M2 macrophages in hearts not subjected to MI. The scatter plot shows

upregulation of 70 genes and downregulated of 39 genes in M2-like macrophages after MI

compared to M2-like macrophages . Among the upregulated genes is neuregulin 1 (*Nrg1*).

D, Quantitative reverse transcription-polymerase chain reaction confirmed post-MI upregulation

of *Nrg1* in CD206$^+$F4/80$^+$CD11b$^+$ cardiac M2-like macrophages after MI. *n=4* samples each.

Gene expression levels relative to those in M2-like macrophages from normal hearts are

shown. Mean±SEM values are shown.

$^#P$<0.05, $^{##}P$<0.01, $^{###}P$<0.005 versus M2-like macrophages from normal hearts, 2-tailed,

unpaired Student's *t*-test.

E, Double immunofluorescence staining revealed Nrg1 expression on the surface of CD206$^+$

M2-like macrophages. Scale bars: 100 μm. *n=4* samples each.

Figure 3. Bone marrow-derived macrophages (BMDMs) attenuate H_2O_2-induced apoptosis and senescence of cardiac fibroblasts via Nrg1 secretion.

A, Representative phase-contrast microscopy images. Treatment with a hydrogen peroxide (H_2O_2) solution changed the spindle-shaped appearance of cardiac fibroblasts to a significantly enlarged, flattened morphology. Addition of bone marrow-derived macrophages (BMDMs) altered the senescent morphology to a normal, spindle-shaped morphology. After addition of anti-ErbB antibody (Ab), fibroblasts displayed the same gross morphology as that of senescent fibroblasts treated with H_2O_2. Likewise, addition of recombinant neuregulin 1 (Nrg1) changed the gross morphology to a spindle shape. Scale bars: 100 μm. $n=4$ samples each.

B. Staining for SA-β-gal showed that senescence was exacerbated in H_2O_2-treated fibroblasts but not in fibroblasts cocultured with BMDMs. This suppression of senescence was attenuated by coculture with the anti-ErbB Ab. Nrg1 suppressed fibroblast senescence. Scale bars: 100 μm. $n=4$ samples each.

C. Apoptosis of cardiac fibroblasts (ratio of cleaved caspase 3^+DAPI^+ fibroblasts to $DAPI^+$ fibroblasts) was increased in coculture with H_2O_2 but decreased in coculture with BMDMs. This decrease in apoptosis was eliminated by the anti-ErbB Ab. Nrg1 suppressed apoptosis. Nuclei were counterstained with 4′,6-diamidino-2-phenylindole (DAPI). $n=4$ samples each. Scale bars: 100 μm.

D, Proliferation of cardiac fibroblasts (ratio of Ki-67$^+$ and DAPI$^+$ fibroblasts to DAPI$^+$ fibroblasts) was decreased under H_2O_2 treatment but increased when cells were cocultured with BMDMs. This enhanced proliferation was suppressed by the anti-ErbB Ab. Nrg1

accelerated proliferation. Nuclei were counterstained with DAPI. Scale bars: 50 μm. $n=4$

samples each. Mean±SEM values are shown.

[#]$P<0.05$ versus normal cardiac fibroblasts (control), *$P<0.05$ versus H_2O_2, [‡]$P<0.05$ versus

H_2O_2+BMDMs, [†]$P<0.05$ versus H_2O_2+BMDMs+Ab, [§]$P<0.05$ versus H_2O_2+Nrg1; 1-way

ANOVA.

Figure 4. Bone marrow-derived macrophages (BMDMs) promote fibroblast activation and

collagen synthesis.

A–C. Representative images of cardiac fibroblasts stained immunocytochemically for (A)

vimentin and αSMA, (B) vimentin and collagen I, and (C) vimentin and collagen III. The

staining was performed 48 hours after the start of culture.

A, Activation of cardiac fibroblasts (ratio of vimentin[+] and αSMA[+] myofibroblasts to vimentin[+]

fibroblasts) was equal under treatment with hydrogen peroxide (H_2O_2) but markedly

increased when cells were cocultured with bone marrow-derived macrophages (BMDMs).

This increase in activation was accelerated by the anti-ErbB antibody (Ab). Addition of

recombinant neuregulin 1 (Nrg1) did not affect activation. Scale bars: 100 μm. $n=4$ samples

each.

B. Collagen I synthesis was equal under treatment with H_2O_2 but significantly increased when

cells were cocultured with BMDMs. This increase in production was enhanced by treatment

with anti-ErbB Ab. Addition of Nrg1 did not affect collagen I synthesis. Scale bars: 100 μm.

$n=4$ samples each.

C. Collagen III synthesis was similar to collagen I synthesis. Scale bars: 100 μm. $n=4$ samples

each. Mean±SEM values are shown.

[#]$P<0.05$ versus normal cardiac fibroblasts (control), *$P<0.05$ versus H_2O_2, [‡]$P<0.05$ versus

H_2O_2+BMDMs, [†]$P<0.05$ versus H_2O_2+BMDMs+Ab, [§]$P<0.05$ versus H_2O_2+Nrg1; 1-way

ANOVA.

Figure 5. *PI3K/Akt signaling pathway is associated with BMDM-attenuated apoptosis and*

senescence of cardiac fibroblasts through Nrg1.

A. Representative bands of PI3K, pPI3K, Akt, pAkt, and β-actin in cardiac fibroblasts treated (or

not treated) with hydrogen peroxide (H_2O_2), in those cocultured with bone marrow-derived

macrophages (BMDMs), and those treated (or not treated) with anti-ErbB antibody

(Ab)/recombinant neuregulin 1 (Nrg1). Treatment and coculture lasted 48 hours. Bar graph

shows quantification of relative pPI3k/PI3k and pAkt/Akt. H_2O_2 alone did not affect activation

of PI3K/AKT signaling in fibroblasts. Addition of BMDMs significantly activated the signaling

pathway and, conversely, addition of the anti-ErbB Ab impaired it. Nrg1 re-stimulated the

signaling pathway. *n*=4 samples each.

B. Quantitative reverse transcription-polymerase chain reaction analysis of cardiac fibroblasts

having been treated (or not) with H_2O_2, co-cultured with BMDMs, and treated (or not) with

anti-ErbB Ab/Nrg1. Treatment and coculture lasted 48 hours. Expression levels of

senescence-associated genes (*p21, p16*, and *Glb1*) were increased in cells treated with H_2O_2

but decreased with the addition of BMDMs. Addition of the anti-ErbB Ab exacerbated

senescence and, conversely, addition of Nrg1 suppressed it. *n*=4 samples each.

C. Expression of cell cycle-associated genes (*Cdk4*, *Cdk6*, *Cdk2*, and *Ki-67*) were suppressed in

cells treated with H_2O_2 but restored in cells cocultured with BMDMs. Addition of anti-ErbB

Ab decreased such expression and, conversely, addition of Nrg1 restored such expression.

n=4 samples each.

D, Expression of the p53 suppressor gene (*MDM2*) was significantly increased in cells

cocultured with BMDMs. *n*=4 samples each.

E, Expression of cell survival-associated gene (*mTOR*) was suppressed in cells treated with H_2O_2

but restored in cells cocultured with BMDMs. Addition of anti-ErbB Ab re-suppressed the

expression and, conversely, addition of Nrg1 restored the *mTOR* expression.

F, Expression of senescence-associated secretory phenotype-associated gene (*IL-6*) was

significantly increased in cells treated with H_2O_2 and in those cocultured with BMDMs, with

ErbB Ab having been added. *n*=4 samples each.

G, Schematic representation and overview of the Nrg1/PI3K/AKT pathway. H_2O_2 treatment

contributes to the development of cellular damage, which leads to senescence, cell cycle

arrest. Nrg1 binding to coreceptor ErbB2/ErbB4 leads to activation of PI3K/AKT and

inactivation of p53 and p21, which results in anti-senescence, cell survival and proliferation.

Arrowheads indicate stimulation, whereas hammerheads represent inhibition. Gene

expression levels relative to the normal cardiac fibroblasts (control) are given. Mean±SEM

values are shown.

[#]$P<0.05$ versus control, *$P<0.05$ versus H_2O_2, [‡]$P<0.05$ versus H_2O_2+BMDMs, [†]$P<0.05$

versus H_2O_2+BMDMs+Ab, [§]$P<0.05$ versus H_2O_2+Nrg1; 1-way ANOVA.

Figure 6. In vivo inhibition of Nrg1 signaling promotes apoptosis and senescence of cardiac

fibroblasts.

A, Quantitative reverse transcription-polymerase chain reaction (qRT-PCR) analysis showed

post-myocardial infarction (MI) upregulation of senescence-associated genes (*p21*, *p16*, and

Glb1) in the infarct area after intraperitoneal injection of the mice with trastuzumab in

comparison to expression of these genes in vehicle injected post-MI heart. *n=4* samples each.

B, Double immunofluorescence staining of Thy1 and cleaved caspase 3 (CC3) showed the ratio

of apoptotic cardiac fibroblasts in the infarct area to be increased on post-MI days 7, 14, and

28 in mice injected intraperitoneally with trastuzumab compared with that in mice injected

with vehicle only. Arrow shows Thy1$^+$CC3$^+$ cells. Scale bars: 100 μm. *n=4* samples each.

C, SA-β-gal staining revealed increased senescence of cardiac cells in the infarct area on post-MI

days 7 and 14 after intraperitoneal trastuzumab injection compared with senescence after

vehicle injection. Scale bars: 100 μm. *n=4* samples each. Gene expression levels relative to

levels in non-MI heart (day 0) are given. Mean±SEM values are shown.

*$P<0.05$, **$P<0.01$, ***$P<0.005$ versus other sample(s); $^\#P<0.05$, $^{\#\#}P<0.01$, $^{\#\#\#}P<0.005$

versus non-MI heart; 1-way ANOVA.

Figure 7. In vivo inhibition of Nrg1 signaling activates cardiac fibroblasts and exacerbates fibrosis.

A, Masson trichrome staining demonstrated that deposition of collagen fibrils was increased in

the infarct area with a time lapse. Intraperitoneal trastuzumab injection significantly increased

collagen fibrils in the infarct area. Scale bars: 100 μm. *n=4* samples each.

B, Double immunofluorescence staining of Thy1 and αSMA showed increased accumulation and

activation of cardiac fibroblasts in the infarct area following trastuzumab injection versus

vehicle injection. Scale bars: 100 μm. *n=4* samples each.

C, Quantitative reverse transcription-polymerase chain reaction (qRT-PCR) analysis showed

post-MI upregulation of fibrosis-associated genes (*αSMA*, *Col1a1*, and *Col3a1*) in mice after

intraperitoneal trastuzumab injection compared with expression in mice after vehicle

injection. *n=4* samples each. Gene expression levels relative to levels in non-MI heart (day 0)

are given. Mean±SEM values are shown.

*P<0.05, **P<0.01, ***P<0.005 versus other sample(s); [#]P<0.05, [##]P<0.01, [###]P<0.005

versus the non-MI heart; 1-way ANOVA.

Figure 8. *In vivo inhibition of Nrg1 signaling exacerbates myocardial inflammation and*

promotes accumulation of M2-like macrophages.

A, Quantitative reverse transcription-polymerase chain reaction analysis (qRT-PCR) showed

post-myocardial infarction (MI) upregulation of senescence-associated secretory phenotype-

associated genes (*CCL3*, *IL-6*, and *TNF*) in mice given trastuzumab by intraperitoneal

injection compared with expression in vehicle-injected mice. *n=4* samples each.

B, Immunohistochemistry showed intraperitoneal trastuzumab injection significantly accelerated

accumulation of CD206[+] M2-like macrophages in the infarct area with a post-MI peak on day

7. Scale bars: 100 μm. *n=4* samples each. Gene expression levels relative to levels in non-MI

heart (day 0) are given. Mean±SEM values are shown.

*P<0.05, **P<0.01, ***P<0.005 versus other sample(s); [#]P<0.05 [##]P<0.01, [###]P<0.005

versus non-MI heart; 1-way ANOVA.

Table I. Primers used in PCR

	Forward	Reverse
Arg1	5'-CAAGACAGGGCTCCTTTCAG-3'	5'-AAGCAAGCCAAGGTTAAAGC-3'
CCL3	5'-CATGAAGGTCTCCACCACTG-3'	5'-CTCCATATGGCGCTGAGAA-3'
CD11b	5'-CTGAGAAATGACGGTGAGGA-3'	5'-CAGCAGGCTTTACAAACCAA-3'
CD11c	5'-GGTCCTACTGTGCACCACAC-3'	5'-GACACTCCTGCTGTGCAGTT-3'
CD206	5'-TGATTACGAGCAGTGGAAGC-3'	5'-GTTCACCGTAAGCCCAATTT-3'
p21 (Cdkn1a)	5'-GATCCACAGCGATATCCAGACA-3'	5'-AGAGACAACGGCACACTTTG-3'
p16 (Cdkn2a)	5'-ATAGACTAGCCAGGGCAGCG-3'	5'-TTGCCCATCATCATCACCTGT-3'
Col1a1	5'-TGAGCCAGCAGATTGAGAAC-3'	5'-CCAGTACTCTCCGCTCTTCC-3'
Col3a1	5'-AGTCTGGAGTCGGAGGAATG-3'	5'-AGGATGTCCAGAGGAACCAG-3'
ErbB2	5'-CTGAATACCATGCAGATGGG-3'	5'-TCACACCATAGCTCCACACA-3'
ErbB4	5'-GAGGAAAGCCCTATGATGGA-3'	5'-TCCAACATTTGACCATGACC-3'
F4/80	5'-CAACCTGCCACAACACTCTC-3'	5'-CCACATCTTCACAGGATTCG-3'
Fizz1	5'-AGGAACTTCTTGCCAATCCA-3'	5'-ACAAGCACACCCAGTAGCAG-3'
Glb1	5'-CGCTACATCTCGGGAAGCAT-3'	5'-GGGCACGTACGTCTGGATTG-3'
Gapdh	5'-CCCCTGGCCAAGGTCATCCA-3'	5'-CGGAAGGCCATGCCAGTGAG-3'
IL-1α	5'-CCATGATCTGGAAGAGACCA-3'	5'-GACAAACTTCTGCCTGACGA-3'
IL-6	5'-AGTCCGGAGAGGAGACTTCA-3'	5'-TTGCCATTGCACAACTCTTT-3'
Ki-67	5'-TATCTGGGCCACCTACCTTC-3'	5'-GCTGTTTCCAGTCCGCTTAC-3'
MHC-II	5'-ATTGCGAAAGCTGCAGAAC-3'	5'-TAGCAGCCAGTCATCCTTTG-3'
Nrg1	5'-GAATTTATGGAAGCGGAGGA-3'	5'-CAGTAGGCCACCACACACAT-3'
Spp1	5'-GAGGAAACCAGCCAAGGTAA-3'	5'-TAGTCCCTCAGAATTCAGCCA-3'
Pdgfα	5'-GAGGAGGAGACAGATGTGAGG-3'	5'-ATTGGCAATGAAGCACCATA-3'
Tgfß	5'-CAACTTCTGTCTGGGACCCT-3'	5'-CGGGTTGTGTTGGTTGTAGA-3'
Tnf	5'-TCGTAGCAAACCACCAAGTG-3'	5'-TTGTCTTTGAGATCCATGCC-3'
p53(Trp53)	5'-GTTCCGAGAGCTGAATGAGG-3'	5'-TTATGGCGGGAGGTAGACTG-3'
Vegfr2	5'-TGTGGCTTCCTGATGGCAGAA-3'	5'-AGAAACCAGTAGACATAGTTT-3'
Vegfα	5'-GTACCTCCACCATGCCAAGT-3'	5'-GCATTCACATCTGCTGTGCT-3'

739 **Table II. Antibodies used for immunocytochemistry**

Primary antibodies	Host	Dilution	Company Catalogue No.
AlexaFluor488-conjugated anti-CD206	Rat	1:100	BioLegend 141709
anti-ErbB2	Rabbit	1:100	Abcam Ab214275
anti-ErbB4	Hamster	1:100	AbD Serotec MCA1369
anti-CD90 (Thy1)	Rat	1:100	eBioscience 14-0901
anti-αSMA	Rabbit	1:100	Abcam ab5694
anti-Ki67	Rat	1:100	eBioscience 14-5698
anti-Cleaved caspase-3	Rabbit	1:200	Cell Signaling 9661
anti-Vimentin	Chicken	1:100	Abcam ab24525
Secondary antibodies			
AlexaFluor 488-conjugated antibody	Goat	1:300	Invitrogen A-11006
AlexaFluor 488-conjugated antibody	Donkey	1:300	Invitrogen A-21208
AlexaFluor 488-conjugated antibody	Goat	1:300	Invitrogen A-11039
AlexaFluor 594-conjugated antibody	Goat	1:300	Invitrogen A-11007
AlexaFluor 594-conjugated antibody	Goat	1:300	Invitrogen A-11012

741 **Table III. Antibodies used for immunoblottin**g

Primary antibodies	Host	Dilution	Company Catalogue No.
Anti-PI3K p85 alpha (phospho Y607)	Rabbit	1:1000	Abcam ab182651
Anti-phospho-Akt (Ser473)	Rabbit	1:2000	Cell Signaling 4060
HRP-conjugated anti-PI3K	Rabbit	1:5000	Abcam ab200773
HRP-conjugated anti-Akt	Rabbit	1:1000	Cell Signaling 8596
HRP-conjugated anti-beta Actin	Mouse	1:50000	Abcam ab49900
Secondary antibody			
HRP-conjugated anti-rabbit IgG	Goat	1:1000	Cell Signaling 7074S

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 7.

Figure 8.

A

B