Reparative macrophages regulate fibrosis in the infarcted myocardium by

attenuating apoptosis and senescence of fibroblasts

Manabu Shiraishi[1,3], Atsushi Yamaguchi[1], Kenta Yashiro[2,3] and Ken Suzuki[3]

[1]Department of Cardiovascular Surgery, Saitama Medical Center, Jichi Medical University, Saitama,

Japan.

[2]Division of Anatomy and Developmental Biology, Department of Anatomy, Kyoto Prefectural

University of Medicine, Kyoto, Japan.

[3]William Harvey Research Institute, Barts and The London School of Medicine and Dentistry, Queen

Mary University of London, London, United Kingdom.

Address correspondence to:

Manabu Shiraishi, Department of Cardiovascular Surgery, Saitama Medical Center, Jichi Medical

University, Saitama, Japan.

Phone: +81-48-647-2111; e-mail: manabu@omiya.jichi.ac.jp

Conflict of interest statement:

The authors have declared that no conflict of interest exists.

This PDF file includes:

Main text

Figures 1 to 9

Abstract

Appropriate fibrotic tissue formation after myocardial infarction (MI) is crucial for maintaining the structure of the heart. Reparative (or M2-like) macrophages play a vital role in fibrosis by activating cardiac fibroblasts after MI. This study aimed to investigate the molecular and cellular mechanisms by which post-MI fibrosis is formed by focusing on the role of the M2-like macrophage subset, and to examine how formation of fibrosis can be controlled. We found that cardiac fibroblasts in the infarcted mouse heart showed apoptosis and senescence, both of which are associated with the fibrotic process. Moreover, we discovered that part of the molecular mechanism underlying fibrotic tissue formation in the infarcted myocardium was attenuation of apoptosis and senescence of fibroblasts by neuregulin 1 (Nrg1)/epidermal growth factor receptor signaling. We analyzed $CD206^{+}F4/80^{+}CD11b^{+}$ M2-like macrophages in intact and infarcted hearts and identified increased Nrg1 expression in M2-like macrophages in the infarcted heart. *In vitro* and *in vivo* studies showed that selective Nrg1 receptor inhibition exacerbated senescence of cardiac fibroblasts, which resulted in excessive progression of fibrosis. These results highlight previously unidentified anti-apoptotic and anti-senescence effects of Nrg1 on cardiac fibroblasts after MI and represent the potential basis for a viable therapeutic strategy for MI.

Introduction

Myocardial infarction (MI) is one of the leading causes of mortality and disability. Even survivors of acute MI frequently develop heart failure owing to adverse ventricular remodeling (1, 2). Because the human heart has an insufficient regenerative ability, connective tissue formation is essential for maintaining integrity and rigidity of the heart. However, the mechanism by which cardiac fibrosis is regulated post-MI is not fully understood.

MI causes permanent loss of hundreds of millions of cardiomyocytes (3). Studies have also shown that even non-cardiomyocytes, including fibroblasts, disappear by apoptosis in large quantities in the infarcted area, and that senescence-associated defects occur in cardiac repair after MI (4, 5). Apoptosis plays an important role in the disappearance of infiltrated immune cells and interstitial cardiac cells after MI (5). Determining the effects of apoptosis and senescence on cell function during the tissue repair process after MI is important. Cellular senescence and apoptosis are processes of growth arrest in response to cellular stress and damage, and they limit proliferation of mammalian cells (6, 7). Senescent cells show a complex phenotype, characterized by cell cycle arrest mediated by the p16 and p53/p21 pathways, and a unique secretory phenotype, known as the senescence-associated secretory phenotype (SASP) (8). Cell cycle arrest plays a central role in the senescent phenotype of adult cardiomyocytes, and induction of cell cycle reentry of adult cardiomyocytes promotes cardiac repair post-MI (9). Furthermore, anti-apoptotic substances and Fas receptor competitive inhibitors are able to inhibit cardiomyocyte apoptosis, resulting in a decrease in the infarct size and an improvement in cardiac function after MI (10).

Previous studies have shown that even non-cardiomyocytes, including fibroblasts, undergo apoptosis in the infarcted area (4, 5). Therefore, cellular senescence and/or apoptosis in cardiac fibroblasts, not limited to cardiomyocytes, is likely to be involved in the tissue repair process after MI. To some extent, senescence of fibroblasts restricts fibrosis in itself, but the long-term presence of this condition is deleterious in the tissue repair process (11). Proper fibrosis after MI is crucial for maintaining the structure of the heart, but excessive fibrosis eventually leads to heart failure. Therefore, adjusting the balance between a pro-fibrotic versus an anti-fibrotic environment is

important for a successful regenerative outcome. To understand the tissue repair process of fibrosis,

clarifying intercellular communication between senescent and apoptotic fibroblasts and surrounding

cells is necessary.

Recent studies have shown that macrophages are essential factors in regeneration of the neonatal

mouse heart (12). We have previously shown that reparative (or M2-like) macrophages play a pivotal

role in fibrotic tissue formation post-MI through promotion of proliferation and activation of cardiac

fibroblasts (13). However, the molecular mechanism used by M2-like macrophages in the

anti-senescence and anti-fibrotic environment after MI remains unknown. We hypothesize that

M2-like macrophages play a vital role in attenuating apoptosis and senescence, which cause excessive

fibrosis, of cardiac fibroblasts.

Neuregulin 1 (Nrg1) is one of the neuregulin genes (*Nrg1–Nrg4*) that belong to the epidermal

growth factor family (14, 15), and it plays essential roles in protection and proliferation of

cardiomyocytes in response to injury (16-20). Although the association between Nrg1 and protection

of cardiomyocytes has been studied for several decades (16-20), the role of Nrg1 in protecting cardiac

fibroblasts and in post-MI regeneration has not yet been established. This study aimed to investigate

the molecular and cellular mechanism by which post-MI fibrosis is formed with a focus on the role of

the M2-like macrophage subset, and to examine how formation of fibrosis can be controlled. The

findings of this study are expected to help clarify the pathophysiology of fibrosis after MI and to form

the basis for development of new therapeutic methods focused on senescence and apoptosis of cardiac

fibroblasts.

Results

Cardiac fibroblasts undergo apoptosis and senescence after MI

First, we investigated the occurrence of cellular senescence and apoptosis in the process of post-MI

fibrosis in a mouse model of MI based on coronary artery ligation. In this model, obvious fibrotic

tissue formation and increased myocardial expression of fibrosis-associated genes (i.e., alpha-smooth

muscle actin [*αSMA*]*, Col1a1,* and *Col3a1*) were observed in the infarct area as early as day 7 post-MI

(Supplemental Figure 1A, B). This change was associated with an increase in Thy1$^+$ fibroblasts and

Thy1$^+$αSMA$^+$ myofibroblasts in the infarct area (Supplemental Figure 1C, D) with a peak at day 7

post-MI. Approximately, 40% and 15% of Thy1$^+$ fibroblasts were positive for cleaved caspase-3 in

the infarct area on days 7 and 28 after MI, respectively, which suggested robust occurrence of

apoptosis in cardiac fibroblasts. Cleaved caspase-3-positive cardiac fibroblasts were rarely found in

the non-infarcted remote area (Figure 1A). Simultaneously, senescence-associated beta galactosidase

(SA-β-Gal)-positive cells were found in the same infarct area. These cells showed a spindle shape

with many cytoplasmic processes, which suggested that they were fibroblasts (Figure 1B).

Additionally, myocardial expression of senescence-associated genes (i.e., *SA-β-Gal, p16, p53,* and

p21) (1, 6, 7, 21, 22) was upregulated in the infarct area compared with the non-infarcted remote area

(Figure 1C). Increased apoptosis and senescence in fibroblasts (and other types of cardiac cells)

exacerbated myocardial inflammation post-MI (Supplemental Figure 1E). Taken together, these

results suggest that apoptosis and senescence occur in cardiac fibroblasts during the process of fibrotic

tissue formation in the post-MI heart.

Nrg1 is upregulated in the infarcted myocardium, while cardiac fibroblasts express Nrg1 receptors

Quantitative reverse transcription-polymerase chain reaction (qRT-PCR) data showed that *Nrg1* was

upregulated in the infarct area of the mouse heart with a peak on day 7 post-MI (Figure 2A).

Immunohistochemistry (IHC) showed that most of the Thy1$^+$ cardiac fibroblasts that had accumulated

in infarcted and remote areas expressed epidermal growth factor receptor (ErbB) 2 and ErbB4, which

are co-receptors of Nrg1 (20, 23, 24). However, fibroblasts in the intact heart (no MI) did not express

these receptors (Figure 2B). These results suggest that Nrg1 is involved in changes in cardiac

fibroblasts after MI.

M2-like macrophages accumulate in the infarct area and express Nrg1

M2-like macrophages play an important role in fibrotic tissue formation post-MI (13). IHC showed

that CD206$^+$ M2-like macrophages had accumulated in the infarct area with a peak at day 7 post-MI

(Figure 3A). Interestingly, this change in M2-like macrophages post-MI corresponded to a change in

occurrence of cardiac fibroblasts post-MI (Supplemental Figure 1C). We found that CD206[+] cells

were also positive for F4/80 in the intact and MI heart (Figure 3B). Microarray analysis showed that

CD206[+]F4/80[+]CD11b[+] M2-like macrophages that were collected from hearts on day 7 after MI had a

different molecular signature than that in those from intact hearts (Supplemental Figure 2A). A range

of anti-inflammatory and reparative genes were upregulated in M2-like macrophages from MI hearts

compared with intact hearts (Supplemental Figure 2B; full data are available in the Gene Expression

Omnibus [GEO] database; GSE69879). Importantly, gene ontology analysis showed that M2-like

macrophages that were collected from hearts on day 7 after MI were associated with regulation of

apoptosis and cell death (Supplemental Figure 2C). In particular, increased *Nrg1* expression levels in

CD206[+]F4/80[+]CD11b[+] cardiac M2-like macrophages were observed after MI (Figure 3C), which was

further confirmed by qRT-PCR and IHC (Figure 3D, E). These data suggest that cardiac M2-like

macrophages may be one of the major sources of upregulation of Nrg1 post-MI.

Bone marrow-derived macrophages attenuate H_2O_2-induced apoptosis and senescence of cardiac

fibroblasts via Nrg1 secretion

We then investigated the role of macrophages in regulating senescence and apoptosis in *in vitro*

co-culture of cardiac fibroblasts with bone marrow-derived macrophages (BMDMs) using a Boyden

Chamber system (13, 25) (Supplemental Figure 3A). H_2O_2 was used to induce apoptosis and

senescence in fibroblasts (Supplemental Figure 3B, C). BMDMs cultured in the presence of H_2O_2

showed an M2-like macrophage phenotype (Supplemental Figure 4A, B). Expression of *Nrg1* in

BMDMs and *ErbB2* and *ErbB4* in isolated cardiac fibroblasts was observed (Supplemental Figure

5A–C). This gene expression was upregulated in response to H_2O_2. Phase-contrast microscopy

showed that fibroblasts treated with H_2O_2 had an enlarged, flattened, senescent morphology, which

became a spindle-shaped healthy form after co-culture with BMDMs. Senescent fibroblasts treated

with anti-ErbB4 antibody (ErbB Ab), which is a competitive inhibitor of Nrg1, displayed the same

gross morphology as senescent fibroblasts treated with H_2O_2. Recombinant Nrg1 similarly changed

the gross morphology of senescent fibroblasts treated with H_2O_2 to a spindle-shaped healthy form

again (Figure 4A). SA-β-Gal staining showed that approximately 20% of fibroblasts treated with

H_2O_2 became positive for SA-β-Gal. This change was significantly attenuated by co-culture with

BMDMs, while addition of ErbB Ab eliminated this anti-senescence effect of BMDMs. Furthermore,

administration of recombinant Nrg1 suppressed H_2O_2-induced senescence of fibroblasts (Figure 4B).

The ratio of cleaved caspase 3^+ apoptotic cells in cardiac fibroblasts was also significantly increased

with H_2O_2 stimulation (Figure 4C). Co-culture with BMDMs markedly attenuated this apoptotic

change in fibroblasts. This anti-apoptotic effect of BMDMs was attenuated by addition of ErbB Ab,

while administration of recombinant Nrg1 suppressed H_2O_2-induced apoptosis of fibroblasts.

Immunolabeling for Ki67 and a scratch assay showed that proliferation and migratory properties in

cardiac fibroblasts were significantly attenuated with H_2O_2 stimulation (Figure 4D and Supplemental

Figure 6). Co-culture with BMDMs markedly improved these cellular activities in fibroblasts. These

effects of BMDMs were attenuated by addition of ErbB Ab, while administration of recombinant

Nrg1 suppressed H_2O_2-induced functional deterioration of fibroblasts. H_2O_2 stimulation reduced

proliferation and migration of cardiac fibroblasts, which were rescued by co-culture with BMDMs.

This effect of BMDMs was eliminated by addition of ErbB Ab. Recombinant Nrg1 administration

showed a strong ability to increase proliferation of fibroblasts. These results collectively suggest that

M2-like macrophages reduce apoptosis and senescence of fibroblasts through secretion of Nrg1.

BMDMs promote activation of fibroblasts and collagen synthesis

Immunocytological staining showed that, although independent H_2O_2 treatment did not affect

transformation of cardiac fibroblasts into αSMA$^+$ myofibroblasts, co-culture with BMDMs and H_2O_2

stimulation induced activation of fibroblasts. Furthermore, addition of ErbB Ab enhanced this

BMDM-induced activation of fibroblasts. Moreover, addition of recombinant Nrg1 did not affect

transformation of cardiac fibroblasts (Figure 5A). Changes in synthesis of types I and III collagen in

fibroblasts in response to H_2O_2, BMDMs, and Nrg1 were associated with that of αSMA expression

(Figure 5B, C). These results suggest that BMDMs, which have an M2-like phenotype (Supplemental

Figure 4A, B), induce activation of fibroblasts to myofibroblasts, and this activation is accelerated

with progression of senescence of fibroblasts. Osteopontin (*Spp1*) is a key mediator of M2-like

macrophage-induced cardiac fibroblast activation (13). *Spp1* expression was increased in BMDMs in

response to H_2O_2 (Supplemental Figure 7A). In contrast, other profibrotic factors, including *Tgfb1* and

Pdgfa (26, 27), were not upregulated in M2-like macrophages (Supplemental Figure 7B).

The phosphatidylinositol 3-kinase/protein kinase B signaling pathway is associated with

BMDM-attenuated apoptosis and senescence of cardiac fibroblasts through Nrg1

We then investigated a potential mechanism underlying attenuation of apoptosis of cardiac fibroblasts

and senescence by BMDMs and/or Nrg1. The phosphatidylinositol 3-kinase (PI3K)/protein kinase B

(Akt) signaling pathway is a downstream part of the ErbB pathway. Western blot analysis showed that

co-culture with BMDMs increased activation of PI3K/Akt in fibroblasts, while this was attenuated by

addition of ErbB Ab. Recombinant Nrg1 administration also activated the PI3K/Akt pathway in

fibroblasts (Figure 6A). To determine the possible relationship between senescence-associated

p16/p53/p21 and the Nrg1/ErbB/PI3K/Akt signaling pathways, we examined mRNA expression

levels of senescence-associated genes (*p53*, *p21*, *p16, and SA-β-Gal)* (Figure 6B), cell

cycle-associated genes (cyclin-dependent kinase 4 [*Cdk4*], cyclin-dependent kinase 6 [*Cdk6*],

cyclin-dependent kinase 2 [*Cdk2*], and *Ki-67*) (Figure 6C), a p53 suppressor gene (murine double

minute 2 [*MDM2*]) (Figure 6D), a cell survival-associated gene (mechanistic target of rapamycin

[*mTOR*]) (Figure 6E), and an SASP-associated gene (interleukin-6 [*IL-6*]) (Figure 6F). Expression of

senescence-associated genes (*p16* and *p21*) and an SASP-associated gene (*IL-6*) was significantly

higher, and expression of cell cycle-associated genes (*Cdk4, Cdk6, Cdk2,* and *Ki-67*) and a cell

survival-associated gene (*mTOR*) was markedly lower in fibroblasts treated with H_2O_2 compared with

controls. Importantly, these changes in gene expression, including progression of senescence and

inflammation, and suppression of the cell cycle, recovered to those of controls by co-culture with

BMDMs. Addition of ErbB Ab resulted in gene expression similar to that with H_2O_2 treatment.

Moreover, expression of senescence-associated genes (*p53*, *p16,* and *p21)* was significantly

suppressed, and expression of cell cycle-associated genes (*Cdk4, Cdk6, Cdk2,* and *Ki-67*) was

enhanced by recombinant Nrg1 in fibroblasts treated with H_2O_2. For other related genes, BMDMs

enhanced expression of the p53 suppressor gene *MDM2*. These results suggest that the Nrg1/ErbB

system operates downstream of PI3K/Akt signaling activation and exerts a suppressive effect on cell

cycle arrest, senescence, and apoptosis. At the same time, this signaling activity is likely to increase

cellular proliferation and survival (Figure 6G).
212

In vivo inhibition of Nrg1 signaling exacerbates fibrosis

We used trastuzumab to clarify the role of Nrg1 in suppressing senescence and apoptosis of cardiac

fibroblasts post-MI *in vivo*. Trastuzumab is an anti-human epidermal growth factor receptor type 2

(HER2) monoclonal antibody that binds to the extracellular juxtamembrane domain of HER2.

Trastuzumab is an effective treatment for HER2/neu[+] tumors in animals and humans (28, 29). We

hypothesized that trastuzumab administration eliminates the anti-apoptotic and anti-senescence effects

of Nrg1, and therefore, trastuzumab increases the number of senescent cardiac fibroblasts. We also

hypothesized that, in the *in vitro* co-culture model, M2-like macrophages activate senescent

fibroblasts, resulting in escalation in fibrotic scar formation in the infarcted myocardium. We found

that intraperitoneal trastuzumab injections did not affect mRNA expression in the intact heart

(Supplemental Figure 8A–D).

Mice underwent surgery to induce MI and received intraperitoneal injections of either trastuzumab

or vehicle only (Supplemental Figure 9A). Trastuzumab did not affect post-MI mortality or body

weight (Supplemental Figure 9B, C). Gene expression profiles in the infarcted myocardium suggested

that apoptosis and senescence were augmented by trastuzumab administration in the infarct area

(Figure 7A–C). These changes in gene expression corresponded to an increase in apoptotic fibroblasts

and senescent cardiac cells in the infarct area (Figure 7D, E). Additionally, trastuzumab

administration increased fibrotic tissue formation in the infarct area (Figure 8A). In fact, the

occurrence of Thy-1[+] fibroblasts in this area was increased (Figure 8B). The number of αSMA[+]Thy1[+]

myofibroblasts was also increased in association with upregulation of *αSMA, Col1a1,* and *Col3a*1 in

the infarcted myocardium (Figure 8B, C). Increased inflammation (Figure 9A) and increased

occurrence of M2-like macrophages in this area (Figure 9B) may explain these findings. Trastuzumab

administration might have increased apoptosis and senescence in fibroblasts (and other types of

cardiac cells), which exacerbated myocardial inflammation post-MI. This led to accumulation of

M2-like macrophages. Such augmented inflammatory signals or an increased number of M2-like

macrophages could accelerate proliferation of cardiac fibroblasts, while increased M2-like

macrophages could promote activation of fibroblasts to myofibroblasts. Subsequently, trastuzumab

administration might have exacerbated fibrotic tissue formation. This concept is supported by

additional findings in the non-infarcted remote area. Trastuzumab administration induced apoptosis-

and senescence-associated expression in the remote area (Supplemental Figure 10A–E) in which such

cell damage hardly occurred post-MI. This corresponded to increased inflammation (Supplemental

Figure 11A). These changes resulted in augmented accumulation of M2-like macrophages and cardiac

fibroblasts in the remote myocardium (Supplemental Figure 11B, 12A). Activation of fibroblasts to

myofibroblasts and fibrosis-associated gene expression were also prompted (Supplemental Figure

12A, B). Consequently, fibrosis occurred even in the remote area (Supplemental Figure 13A). These

results suggest that trastuzumab augments apoptosis and senescence of cardiac fibroblasts post-MI,

resulting in excessive fibrosis, even in the non-infarcted remote area.

Discussion

Despite the enormous clinical importance of cellular and molecular processes underlying the

formation of post-MI fibrosis, they are not well understood. In this study, we described the qualitative

changes in cardiac fibroblasts during formation of fibrosis post-MI. After MI, senescence and

apoptosis of cardiac fibroblasts occurred. However, Nrg1 suppressed senescence and apoptosis of

fibroblasts by activating the Nrg1/PI3K/Akt signaling pathway and indirectly decreased excessive

fibrosis. These results support our hypothesis that M2-like macrophages play a vital role in

suppressing senescence and apoptosis of fibroblasts and regulating collagen synthesis (Figure 9).

After induction of MI, senescence-associated gene expression (i.e., *SA-β-Gal*, *p53, p21,* and *p16*) was

significantly increased in the infarct area. Cardiac fibroblasts that were accumulated in the infarct area

expressed *SA-β-Gal* and cleaved caspase 3 at a certain ratio. One of the reparative genes, *Nrg1,* was

upregulated in the infarct area. Cardiac fibroblasts showed ErbB2/ErbB4, which are co-receptors of

Nrg1 (20, 23, 24), in the damaged myocardium. To precisely determine the role of cardiac M2-like

macrophages in the activity of apoptotic and senescent fibroblasts, we analyzed gene expression

profiling of these macrophages. Interestingly, we found more than four-fold upregulation of *Nrg1*

expression in M2-like macrophages in the post-MI heart compared with the normal heart. Our

findings were strengthened by an additional *in vitro* experiment in which H_2O_2-induced senescent

cardiac fibroblasts were co-cultured with BMDMs. Senescent cells attract macrophages in

pathological conditions (30). This model allowed us to determine the precise interaction between

M2-like macrophages and senescent cardiac fibroblasts. IHC showed that after the processes of

senescence and apoptosis of fibroblasts were improved by co-culture with BMDMs, addition of ErbB

Ab exacerbated them again. Simultaneously, western blot and qRT-PCR analysis showed that

PI3K/Akt signaling pathways were activated by co-culture with BMDMs and suppressed by adding

ErbB Ab. Furthermore, adding recombinant Nrg1 resulted in the same activation of PI3K/Akt

signaling pathways as co-culture with BMDMs. Additionally, co-culture with BMDMs promoted

H_2O_2-induced senescent fibroblasts to transform fibroblasts into αSMA^+ myofibroblasts, which

resulted in increased collagen synthesis. This activation and collagen synthesis were further

augmented by adding ErbB Ab. These results corresponded to *in vivo* observations with trastuzumab.

Intraperitoneal trastuzumab injections further increased senescence and secretion of SASP. This

exacerbation of inflammation induced by trastuzumab caused increased accumulation of M2-like

macrophages in the infarct area. IHC staining and qRT-PCR analysis showed that accumulation of

M2-like macrophages promoted proliferation and activation of fibroblasts, which resulted in

augmentation of fibrosis even in the non-infarcted remote area.

Nrg1 is a cytokine that belongs to a family of proteins structurally related to epidermal growth

factor and plays essential roles in protection and proliferation of cardiomyocytes in response to injury

(16-20). Nrg1 is synthesized in endothelial cells near cardiomyocytes (20). We observed that *Nrg1*

was expressed in M2-like macrophages and that it had a specific function for rescuing

post-MI-induced senescent and apoptotic cardiac fibroblasts. Cardiac fibroblasts showed

ErbB2/ErbB4 in the damaged myocardium on days 7 and 28 post-MI. Nrg1 binds to the ErbB4

receptor. After heterodimerization with phosphorylated ErbB2, signaling pathways that are activated

downstream of ErbB2/ErbB4 signals link to the Ras-mitogen-activated protein kinase pathway and

PI3K/Akt pathways (20). The hypertrophic response to Nrg1 is mainly Ras dependent, whereas the

anti-apoptotic and cell proliferation effects are likely to be Akt dependent (31-33). Therefore, Nrg1

could attenuate the senescence-associated genes *p53*, *p21*, and *p16* through the PI3K/Akt pathways. In

our study, we observed a transient increase in *p53*, *p21*, and *p16* expression in infarcted myocardium

and cultured fibroblasts. Zhu *et al*. reported that MI promoted accumulation of senescent cardiac

fibroblasts in the heart and *p53* expression (1). Increased p53 activity in response to diverse

pathological stresses, such as MI, induces apoptosis (34, 35). Stress-induced *p53* expression increases

p21 expression in response to DNA damage and produces a reversible proliferative arrest that

provides time for DNA repair and facilitates survival of cells (36, 37). Previous studies have shown

that p21 binds to and inhibits CDK2-mediated inactivation of retinoblastoma, subsequently preventing

entry into the S phase of the cell cycle (7, 38). Furthermore, p21 is important for initiating senescence

in some settings, but its expression does not persist in senescent cells (39, 40). Simultaneously,

increased *p16* expression is found in infarct tissue. Irreversible proliferative arrest can be produced by

p16 inhibiting two cycle-dependent kinases, CDK4 and CDK6 (7, 41). Therefore, a change in

expression of these genes (i.e., *p53*, *p21*, and *p16*) in cardiac cells *in vivo* and cardiac fibroblasts *in*

vitro suggests that Nrg1 is a crucial factor controlling reversible and irreversible senescence of cardiac

cells after MI.

Previous reports have shown that trastuzumab efficiently stops or slows growth of ErbB2[+] cells

in vitro and inhibits the ability of cells to repair damaged DNA (28, 29). Our study showed that

trastuzumab injections further increased senescence and apoptosis in infarcted myocardium. As

inflammation worsens, M2-like macrophages from the bone marrow accumulate at the site of

damaged tissue (42). In our study, exacerbation of inflammation induced by trastuzumab caused

increased accumulation of M2-like macrophages and promoted proliferation and activation of

fibroblasts. Although we analyzed mRNA gene expression in tissue sections of the heart and not in

single cells, increased expression levels of senescence-associated genes were considered to reflect

senescence of fibroblasts to a certain extent. These results are consistent with our previous report,

which showed that osteopontin-mediated transformation of fibroblasts into myofibroblasts resulted in

progression of fibrosis (13). Interestingly, senescence and SASP-associated gene expression peaked

slightly later in the remote area than in the infarct area. One possible pathological mechanism of the

non-infarcted remote area is assumed to be indirect damage via SASP rather than direct cytotoxicity

due to ischemia. Senescent cells autonomously induce senescence-like gene expression in their

surrounding non-senescent cells through SASP (43). Increased SASP in the infarct area might not

simply reflect senescence and apoptosis in cells in the infarct area, but also changes in non-senescent

cells in the remote area. All of these data suggest that trastuzumab-induced exacerbation of fibrosis is

mainly mediated by progression of senescence and apoptosis in cardiac fibroblasts.

Our data provide new biological insights into the molecular mechanisms by which M2-like

macrophages regulate post-MI tissue repair by affecting senescence, apoptosis, and proliferation of

fibroblasts. This repair allows senescent cardiac fibroblasts to be maintained to a certain degree.

Although the process is not perfect, it allows them to survive. Incomplete rescue of fibroblasts from

senescence, mediated by M2-like macrophages, might lead to undesired fibrosis after diastolic

dysfunction. Although Nrg1 has been repeatedly reported to play important roles in injured

cardiomyocytes (16-20), its role in cardiac fibroblasts in this context has not been well studied. In our

study, the underlying molecular mechanisms of Nrg1 were investigated in experiments with

H_2O_2-induced senescent cardiac fibroblasts, and these showed activation of the p53/p21 and p16

pathways followed by cell cycle arrest. We present new *in vitro* evidence suggesting that M2-like

macrophages play a vital role for attenuating senescent and apoptotic fibroblasts through the

Nrg1/PI3K/Akt signaling pathways. This contributes to a variety of processes that are critical in

mediating many aspects of cellular function, including cell growth and survival (44). However,

further mechanistic studies are required to understand the signaling pathways downstream of these

factors and investigations in cell type-specific conditional *ErbB2/ErbB4*-knockout mice and to further

confirm the direct role of Nrg1 in cardiac fibroblasts. This knowledge could answer questions about

which cardiac cells are directly and indirectly affected, which cellular responses occur after MI, and

which signaling mechanisms are involved. Better understanding of the healing process of infarcts and

subsequent remodeling may result in new treatment options.

Although this study focused on Nrg1-induced anti-apoptosis and anti-senescence of cardiac

fibroblasts, M2-like macrophages are likely to mediate supplementary benefits in cardiac repair

post-MI. These benefits could include reduced inflammation, activation of fibroblasts, and

neovascular formation, as shown in our previous study (13). To develop potential therapies mediated

by M2-like macrophages, future studies will also need to determine how the gene that encodes Nrg1

is switched on by MI and to identify the other molecules that regulate apoptosis, senescence, and

proliferation of cardiac fibroblasts. There might be an alternative signal(s) to Nrg1 by which cardiac

M2-like macrophages induce proliferation of fibroblasts. This signal might be relatively negative

because we found that ErbB Ab totally depleted the ability of proliferation of fibroblasts from

BMDMs (Figure 4D).

In conclusion, our data provide evidence that Nrg1 critically regulates senescence and apoptosis of

cardiac fibroblasts in the infarcted adult murine heart and *in vitro* in a co-culture model. This process

could play a vital role in repair of the infarcted fragile ventricular wall and diastolic dysfunction due

to excessive fibrosis. Therefore, this tissue repair mechanism controls the degree of rigidity and

contraction of the infarcted heart, thus determining the prognosis post-MI. Targeted activation of

M2-like macrophages might enhance this endogenous repair mechanism in senescent cardiac

fibroblasts, suggesting that this approach could be a new therapeutic treatment of MI.

Methods

Animals. Eight- to 10-week-old mice were used in the experiments. C57BL/6 mice were purchased

from Tokyo Laboratory Animals Science Co., Ltd. The mice were maintained in a specific

pathogen-free room in our animal facility on a 12-hour light/dark cycle with free access to food and

water. The in vitro studies and in vivo procedures and assessments were performed in a blinded

manner.

In vivo treatments. Mice were treated with three i.p injections of 100 μg of Trastuzumab (Bio X Cell;

catalog BE0277) on the fourth, fifth, and sixth days after induction of myocardial infarction (MI).

Samples were collected on the seventh, 12th, and 28th days after induction of MI.

Preparation of bone marrow-derived macrophages (BMDMs). Mouse BMDMs were prepared from

the femurs and tibiae of 8-week-old wild type (WT) mice, as previously described (13). Bone marrow

mononuclear cells were collected by centrifugation on Ficoll-Paque (GE Healthcare) and were

cultivated overnight in a CO_2 incubator in Dulbecco's Modified Eagle Medium (DMEM;

Sigma-Aldrich) containing 10% fetal bovine serum (FBS), 50 U/mL penicillin, 50 μg/mL

streptomycin, and 10 ng/mL granulocyte-macrophage colony stimulating factor (GM-CSF) (R&D

Systems; 415-ML). The non-attached or weakly attached cells were collected and transferred to new

dishes and cultivated for an additional 5 days, then the cells were prepared for coculture experiments.

Coculture of cardiac fibroblasts and BMDMs in a Boyden chamber. Cardiac fibroblasts (2×10^4)

were plated on a 0.1% gelatin-coated 6-well dish (Thermo Scientific) and cultured for 48 hours in

DMEM containing 10% FBS, 50 U/mL penicillin, and 50 μg/ mL streptomycin. Cardiac fibroblasts

were damaged by 1 hour of coculture with 100 μM hydrogen peroxide (Sigma), as previously

described (45), with some modifications. Then, the damaged fibroblasts were maintained in DMEM

containing 10% FBS, 50 U/mL penicillin, and 50 μg/mL streptomycin. 2×10^4 BMDMs were seeded

onto polycarbonate membrane inserts (pore size, 0.4 μm; Thermo Scientific), which were set into the

wells containing fibroblasts. ErbB4 antibody (10 μg/mL; Thermo Scientific; catalog MA5-13016)

was added to the culture medium at the beginning of the coculture.

Induction of myocardial infarction (MI). MI was induced in mice by ligating the left coronary artery

under 2.0% isoflurane anesthesia and mechanical ventilation, as previously described (46). The

successful generation of MI was confirmed by changes in the color and motion of the left ventricular

walls. The survival rate and body weight change after MI were monitored daily.

Isolation of heart cells. Mouse heart cells were isolated as previously described (47). Immediately

after cervical dislocation, the aorta was clamped, and cooled Hank's balanced salt solution (HBSS;

Sigma-Aldrich) was injected into the left ventricular cavity. The isolated hearts were cut into 1-mm

pieces, digested with 0.05% collagenase II (Sigma-Aldrich) at 37 °C for 15 minutes and filtered with

a 40-µm cell strainer (BD Falcon). The remnant heart tissues were again digested with a fresh

digestion solution and filtered similarly; this cycle was repeated 5 times. The suspension obtained at

each cycle was combined and subjected to flow cytometric analyses or fluorescence-activated cell

sorting (FACS) after erythrocytes were depleted with Red Cell Lysis Buffer (BioLegend) according to

the manufacturer's protocol.

Flow cytometry and FACS. The isolated cells were resuspended in FACS buffer (HBSS plus 2 mM

EDTA plus 0.5% BSA) and preincubated with an anti-mouse CD16/CD32 Ab (rat, 1:100 dilution;

eBioscience; catalog 14-0161) to block the Fc receptor. The dead cells and debris were excluded by

forward scatter/side scatter (FSC/SSC) and DAPI staining (1:1,000 dilution; Sigma-Aldrich). To

determine their phenotypes, the cells were stained with the following Abs for 3 hours at 4 °C:

APC-conjugated anti-CD11b Ab (rat, 1:100 dilution; eBioscience; catalog 17-0112); phycoerythrin-

(PE-)conjugated anti-F4/80 Ab (rat, 1:20 dilution; eBioscience; catalog 12-4801); and Alexa Fluor

488-conjugated anti-CD206 Ab (rat, 1:50 dilution; BioLegend; catalog 141709). Cell sorting was

performed with a FACSAria II (BD Biosciences).

Isolation of cardiac fibroblasts. Cardiac fibroblasts were isolated from Wistar rats (Charles River

Laboratories), as previously described (48), with some modifications. The isolated rat heart was cut

into 1-mm pieces, and the fragments were plated evenly in 10-cm 0.1% gelatin-coated dishes so that

they did not touch each other. Each fragment was covered by a droplet of DMEM medium containing

10% FBS, 50 U/ mL penicillin, and 50 µg/mL streptomycin and incubated for 24 hours at 37 °C in a

CO_2 incubator. Then, sufficient medium was added to cover the entire bottom of the culture dish, and

the culture was incubated for an additional 24 hours. After this period, sufficient medium was added

to completely cover the heart fragments, and the culture was continued for an additional 5 days. When

fibroblast outgrowth was observed, the cells were collected by trypsinization. The fibroblasts were

maintained in DMEM containing 10% FBS, 50 U/mL penicillin, and 50 µg/mL streptomycin in 0.1%

gelatin-coated culture flasks. The cells were used at passage 3 or 4.

RNA extraction and real-time polymerase chain reaction (PCR). Total RNA was extracted from

cells or the heart tissue with the Gene Jet PCR purification Kit (Thermo Scientific) and quantified

with a Nano-Drop 8000 spectrophotometer (Thermo Scientific). cDNA was synthesized with the

high-capacity cDNA Reverse Transcription Kit (Applied Biosystems) from 25 ng and 150 ng of total

RNAs from M2-macrophages and the heart tissues, respectively. Real-time PCR was performed by a

QuantStudio (Applied Biosystems) with a SYBR PremixExTaqII (Takara Bio) in the following

conditions: 95 °C for 30 seconds followed by 40 cycles at 95 °C for 5 seconds and 60 °C for 30

seconds. Gene expression levels were normalized by *Gapdh*. The primers used are shown in the

Supplemental Table 1.

Microarray analysis. Total RNA of $CD206^+F4/80^+CD11b^+$ M2-like macrophages were isolated by

FACS (see above) from the heart and peritoneal cavity, amplified with the RNA Amplification

System (NuGEN) and subjected to an Illumina's beads array platform with a Mouse WG-6 v2.0

Expression BeadChip (Illumina). Two independent biological replicates were prepared for each group

category. The median per chip normalization was performed in each array. We analyzed only genes

whose signal intensity was above the value of 90 in one of the biological duplicates. A 2-fold change

cut-off was used to identify differential lay expressed genes.

Immunohistochemistry. Immediately after cervical dislocation, the aorta of the mouse was clamped

and ice-cold PBS was injected into the left ventricular cavity. The mouse heart was then perfused with

ice-cold 4% paraformaldehyde in PBS. The heart was removed, cut and embedded in optimal cutting

temperature (OCT) compound (VWR International) and frozen in isopentane chilled in liquid nitrogen.

The frozen tissue sections (8-μm thick) were prepared, and the non-specific antibody-binding sites

were pre-blocked with the blocking buffer (PBS plus 5% of goat serum). Then, the primary antibodies

were applied overnight at 4 °C. After rinsing 3 times for 15 minutes in PBS, the sections were next

incubated with appropriate fluorophore-conjugated secondary antibodies and

4',6-diamidino-2-phenylindole (DAPI) (Sigma-Aldrich, D9542) in the blocking buffer for 1 hour at

room temperature. Stained sections were mounted with a DAKO Fluorescence Mounting Medium

(Agilent, S302380-2). The primary and secondary antibodies used in this study are shown in the

Supplemental Table 2.

Picrosirius red staining. The 8-μm thick frozen sections were incubated in 1.5% phosphomolybdic

acid for 60 minutes, then in 0.1% Picrosirius Red for 15 minutes, and finally in 0.5% acetic acid

solution for 3 minutes. After dehydration by adding increasing concentrations of ethanol to xylene,

the sections were mounted with the DPX mounting medium (VWR International). The infarct area

was defined as the area in which more than 90% of cardiomyocytes were lost.

Masson trichrome staining. The frozen tissue sections (8 μm thick) were prepared as above and

stained with the Trichrome Stain Kit (Scy Tek Laboratories) according to the manufacturer's

instructions.

Immunocytochemistry. The cells were fixed by 4% of paraformaldehyde in PBS for 5 minutes at

room temperature. Apart from the cells to be used for surface antigen staining, cells were incubated in

PBS containing 0.1% of Triton x100 for 5 minutes at room temperature. Non-specific antibody

binding sites were pre-blocked with PBS containing 5% of goat serum for 30 minutes at the room

temperature. Then, primary antibodies were applied to the cells for 1 hour at room temperature. The

primary antibodies used were as follows: anti-vimentin (1:100, Abcam, ab24525), anti-cleaved

caspase-3 (1:100, Cell Signaling, 9661), anti-Ki-67 (1:100, eBioscience, 14-5698), and anti-αSMA

(1:00, Abcam, ab5694). After rinsing, cells were incubated with the fluorophore-conjugated

secondary antibodies (1:300, AlexaFluor 488-conjugated or AlexaFluor 594-conjugated polyclonal,

Invitrogen) and DAPI in blocking buffer for 1 hour at room temperature.

Imaging and analysis. All digital images were acquired with an All-in-One microscope (BZ-8000;

KEYENCE). All image analyses were performed by importing images as TIFF files into National

Institute of Health image-analysis software (ImageJ software). For quantification of fibrin formation,

the fibrin area was quantified as the percentage of the fibrin positive area on the whole screen.

Images were acquired at 5 independent regions and analyzed. A color threshold function of ImageJ

was applied to measure macrophages, fibroblasts, fibrin clot area, and cell size of cultured fibroblasts.

Immunoblotting. For western blot analysis, tissue was frozen and crushed; then, it was lysed by

T-PER Tissue Protein Extraction Regent (Thermo Scientific, 78510), according to the

manufacturer's protocol. Protein concentration was measured with a Nano-Drop 8000

spectrophotometer (Thermo Scientific), and 50-μg protein lysates were separated by PAGE,

transferred onto a polyvinylidene difluoride membrane and then identified by immunoblotting by

using the primary and second antibodies shown in the Supplemental Table 3. The protein bands

were visualized with a Super Signal West Pico Substrate (Thermo Scientific, 34077), according to

the manufacturer's instructions.

Statistics. All statistical tests were performed with GraphPad Prism software, version 8 (GraphPad

Software). Data represent the mean ± SEM. For comparisons between multiple groups,

repeated-measures analysis of variance (ANOVA) or 1- or 2-way ANOVA was performed, followed

by Bonferroni's post hoc test. Two groups were compared with the 2-tailed, unpaired Student's *t* test.

Cardiac rupture rates were compared with the χ^2 test, and survival curves were compared with the

log-rank test.

Author contributions

MS and KS conceived and designed the study. MS conducted most of the experiments and

acquired the data, with professional support from AY, KY, and KS. All authors participated in the

analysis and interpretation of the data. MS and KS primarily wrote and edited the manuscript, with

input from all other authors.

Acknowledgments

This project was funded by the Uehara Memorial Foundation, the SENSHIN Medical Research

Foundation, a Grant-in-Aid for Scientific Research (C), the Mochida Memorial Foundation for

Medical and Pharmaceutical Research, the Takeda Science Foundation, and the Public Trust Surgery

Research Fund. We thank Ellen Knapp, PhD, from Edanz Group

(https://en-author-services.edanz.com/ac) for editing a draft of this manuscript.

References

1.   Zhu F, Li Y, Zhang J, Piao C, Liu T, Li HH, et al. Senescent cardiac fibroblast is

510   critical for cardiac fibrosis after myocardial infarction. *PLoS One.* 2013;8(9):e74535.

511   2.   Dickstein K, Cohen-Solal A, Filippatos G, McMurray JJ, Ponikowski P,

512   Poole-Wilson PA, et al. ESC Guidelines for the diagnosis and treatment of acute and

513   chronic heart failure 2008: the Task Force for the Diagnosis and Treatment of Acute

514   and Chronic Heart Failure 2008 of the European Society of Cardiology. Developed in

515   collaboration with the Heart Failure Association of the ESC (HFA) and endorsed by

516   the European Society of Intensive Care Medicine (ESICM). *Eur Heart J.*

517   2008;29(19):2388-442.

518   3.   Gemberling M, Karra R, Dickson AL, and Poss KD. Nrg1 is an injury-induced

519   cardiomyocyte mitogen for the endogenous heart regeneration program in zebrafish.

520   *Elife.* 2015;4.

521   4.   Gould KE, Taffet GE, Michael LH, Christie RM, Konkol DL, Pocius JS, et al. Heart

522   failure and greater infarct expansion in middle-aged mice: a relevant model for

523   postinfarction failure. *Am J Physiol Heart Circ Physiol.* 2002;282(2):H615-21.

524   5.   Takemura G, Ohno M, Hayakawa Y, Misao J, Kanoh M, Ohno A, et al. Role of

525   apoptosis in the disappearance of infiltrated and proliferated interstitial cells after

526   myocardial infarction. *Circ Res.* 1998;82(11):1130-8.

527   6.   Sharpless NE, and Sherr CJ. Forging a signature of in vivo senescence. *Nat Rev*

528   *Cancer.* 2015;15(7):397-408.

529   7.   Munoz-Espin D, and Serrano M. Cellular senescence: from physiology to pathology.

530   *Nat Rev Mol Cell Biol.* 2014;15(7):482-96.

8.   Coppe JP, Desprez PY, Krtolica A, and Campisi J. The senescence-associated

secretory phenotype: the dark side of tumor suppression. *Annu Rev Pathol.*

2010;5:99-118.

9.   Alam P, Haile B, Arif M, Pandey R, Rokvic M, Nieman M, et al. Inhibition of

Senescence-Associated Genes Rb1 and Meis2 in Adult Cardiomyocytes Results in

Cell Cycle Reentry and Cardiac Repair Post-Myocardial Infarction. *J Am Heart Assoc.*

2019;8(15):e012089.

10. Hayakawa K, Takemura G, Kanoh M, Li Y, Koda M, Kawase Y, et al. Inhibition of

granulation tissue cell apoptosis during the subacute stage of myocardial infarction

improves cardiac remodeling and dysfunction at the chronic stage. *Circulation.*

2003;108(1):104-9.

11. Childs BG, Li H, and van Deursen JM. Senescent cells: a therapeutic target for

cardiovascular disease. *J Clin Invest.* 2018;128(4):1217-28.

12. Aurora AB, Porrello ER, Tan W, Mahmoud AI, Hill JA, Bassel-Duby R, et al.

Macrophages are required for neonatal heart regeneration. *J Clin Invest.*

2014;124(3):1382-92.

13. Shiraishi M, Shintani Y, Shintani Y, Ishida H, Saba R, Yamaguchi A, et al.

Alternatively activated macrophages determine repair of the infarcted adult murine

heart. *J Clin Invest.* 2016;126(6):2151-66.

14. Meyer D, Yamaai T, Garratt A, Riethmacher-Sonnenberg E, Kane D, Theill LE, et al.

Isoform-specific expression and function of neuregulin. *Development.*

1997;124(18):3575-86.

15. Fuller SJ, Sivarajah K, and Sugden PH. ErbB receptors, their ligands, and the

consequences of their activation and inhibition in the myocardium. *J Mol Cell Cardiol.*

2008;44(5):831-54.

16.   Bersell K, Arab S, Haring B, and Kuhn B. Neuregulin1/ErbB4 signaling induces

557   cardiomyocyte proliferation and repair of heart injury. *Cell.* 2009;138(2):257-70.

558   17.   Hedhli N, Huang Q, Kalinowski A, Palmeri M, Hu X, Russell RR, et al.

559   Endothelium-derived neuregulin protects the heart against ischemic injury.

560   *Circulation.* 2011;123(20):2254-62.

561   18.   Polizzotti BD, Ganapathy B, Walsh S, Choudhury S, Ammanamanchi N, Bennett DG,

562   et al. Neuregulin stimulation of cardiomyocyte regeneration in mice and human

563   myocardium reveals a therapeutic window. *Sci Transl Med.* 2015;7(281):281ra45.

564   19.   Yutzey KE. Regenerative biology: Neuregulin 1 makes heart muscle. *Nature.*

565   2015;520(7548):445-6.

566   20.   Lemmens K, Doggen K, and De Keulenaer GW. Role of neuregulin-1/ErbB signaling

567   in cardiovascular physiology and disease: implications for therapy of heart failure.

568   *Circulation.* 2007;116(8):954-60.

569   21.   Krizhanovsky V, Yon M, Dickins RA, Hearn S, Simon J, Miething C, et al.

570   Senescence of activated stellate cells limits liver fibrosis. *Cell.* 2008;134(4):657-67.

571   22.   van Deursen JM. The role of senescent cells in ageing. *Nature.*

572   2014;509(7501):439-46.

573   23.   Olayioye MA, Neve RM, Lane HA, and Hynes NE. The ErbB signaling network:

574   receptor heterodimerization in development and cancer. *EMBO J.*

575   2000;19(13):3159-67.

576   24.   Uray IP, Connelly JH, Thomazy V, Shipley GL, Vaughn WK, Frazier OH, et al. Left

577   ventricular unloading alters receptor tyrosine kinase expression in the failing human

578   heart. *J Heart Lung Transplant.* 2002;21(7):771-82.

579   25.   Suzuki T, Arumugam P, Sakagami T, Lachmann N, Chalk C, Sallese A, et al.

580   Pulmonary macrophage transplantation therapy. *Nature.* 2014;514(7523):450-4.

26.   van den Borne SW, Diez J, Blankesteijn WM, Verjans J, Hofstra L, and Narula J.

Myocardial remodeling after infarction: the role of myofibroblasts. *Nat Rev Cardiol.*

2010;7(1):30-7.

27.   Shinde AV, and Frangogiannis NG. Fibroblasts in myocardial infarction: a role in

inflammation and repair. *J Mol Cell Cardiol.* 2014;70:74-82.

28.   Park S, Jiang Z, Mortenson ED, Deng L, Radkevich-Brown O, Yang X, et al. The

therapeutic effect of anti-HER2/neu antibody depends on both innate and adaptive

immunity. *Cancer Cell.* 2010;18(2):160-70.

29.   ElZarrad MK, Mukhopadhyay P, Mohan N, Hao E, Dokmanovic M, Hirsch DS, et al.

Trastuzumab alters the expression of genes essential for cardiac function and induces

ultrastructural changes of cardiomyocytes in mice. *PLoS One.* 2013;8(11):e79543.

30.   Sasaki M, Miyakoshi M, Sato Y, and Nakanuma Y. Modulation of the

microenvironment by senescent biliary epithelial cells may be involved in the

pathogenesis of primary biliary cirrhosis. *J Hepatol.* 2010;53(2):318-25.

31.   Kuramochi Y, Cote GM, Guo X, Lebrasseur NK, Cui L, Liao R, et al. Cardiac

endothelial cells regulate reactive oxygen species-induced cardiomyocyte apoptosis

through neuregulin-1beta/erbB4 signaling. *J Biol Chem.* 2004;279(49):51141-7.

32.   Baliga RR, Pimental DR, Zhao YY, Simmons WW, Marchionni MA, Sawyer DB, et

al. NRG-1-induced cardiomyocyte hypertrophy. Role of PI-3-kinase, p70(S6K), and

MEK-MAPK-RSK. *Am J Physiol.* 1999;277(5 Pt 2):H2026-37.

33.   Gelb BD, and Tartaglia M. RAS signaling pathway mutations and hypertrophic

cardiomyopathy: getting into and out of the thick of it. *J Clin Invest.*

2011;121(3):844-7.

34.   Long X, Boluyt MO, Hipolito ML, Lundberg MS, Zheng JS, O'Neill L, et al. p53 and

the hypoxia-induced apoptosis of cultured neonatal rat cardiac myocytes. *J Clin Invest.*

1997;99(11):2635-43.

35.   Polyak K, Xia Y, Zweier JL, Kinzler KW, and Vogelstein B. A model for

p53-induced apoptosis. *Nature.* 1997;389(6648):300-5.

36.   Deng C, Zhang P, Harper JW, Elledge SJ, and Leder P. Mice lacking p21CIP1/WAF1

undergo normal development, but are defective in G1 checkpoint control. *Cell.*

1995;82(4):675-84.

37.   Wang YA, Elson A, and Leder P. Loss of p21 increases sensitivity to ionizing

radiation and delays the onset of lymphoma in atm-deficient mice. *Proc Natl Acad Sci*

U S A. 1997;94(26):14590-5.

38.   Childs BG, Durik M, Baker DJ, and van Deursen JM. Cellular senescence in aging

and age-related disease: from mechanisms to therapy. *Nat Med.* 2015;21(12):1424-35.

39.   Alcorta DA, Xiong Y, Phelps D, Hannon G, Beach D, and Barrett JC. Involvement of

the cyclin-dependent kinase inhibitor p16 (INK4a) in replicative senescence of

normal human fibroblasts. *Proc Natl Acad Sci U S A.* 1996;93(24):13742-7.

40.   Stein GH, Drullinger LF, Soulard A, and Dulic V. Differential roles for

cyclin-dependent kinase inhibitors p21 and p16 in the mechanisms of senescence and

differentiation in human fibroblasts. *Mol Cell Biol.* 1999;19(3):2109-17.

41.   Serrano M, Lin AW, McCurrach ME, Beach D, and Lowe SW. Oncogenic ras

provokes premature cell senescence associated with accumulation of p53 and

p16INK4a. *Cell.* 1997;88(5):593-602.

42.   Ikeda N, Asano K, Kikuchi K, Uchida Y, Ikegami H, Takagi R, et al. Emergence of

immunoregulatory Ym1(+)Ly6C(hi) monocytes during recovery phase of tissue

injury. *Sci Immunol.* 2018;3(28).

43.   Acosta JC, Banito A, Wuestefeld T, Georgilis A, Janich P, Morton JP, et al. A

complex secretory program orchestrated by the inflammasome controls paracrine

senescence. *Nat Cell Biol.* 2013;15(8):978-90.

44.   Yu JS, and Cui W. Proliferation, survival and metabolism: the role of

PI3K/AKT/mTOR signalling in pluripotency and cell fate determination.

Development. 2016;143(17):3050-60.

45.   Bladier C, Wolvetang EJ, Hutchinson P, de Haan JB, and Kola I. Response of a

primary human fibroblast cell line to H2O2: senescence-like growth arrest or

apoptosis? *Cell Growth Differ.* 1997;8(5):589-98.

46.   Tano N, Narita T, Kaneko M, Ikebe C, Coppen SR, Campbell NG, et al. Epicardial

placement of mesenchymal stromal cell-sheets for the treatment of ischemic

cardiomyopathy; in vivo proof-of-concept study. *Mol Ther.* 2014;22(10):1864-71.

47.   Shintani Y, Kapoor A, Kaneko M, Smolenski RT, D'Acquisto F, Coppen SR, et al.

TLR9 mediates cellular protection by modulating energy metabolism in

cardiomyocytes and neurons. *Proc Natl Acad Sci U S A.* 2013;110(13):5109-14.

48.   Leicht M, Greipel N, and Zimmer H. Comitogenic effect of catecholamines on rat

cardiac fibroblasts in culture. *Cardiovasc Res.* 2000;48(2):274-84.

Figure 1.

Figure 1. *Cardiac fibroblasts underwent apoptosis and senescence after MI.*

(A) Double immunofluorescence staining for Thy1 and cleaved caspase 3 (CC-3) demonstrated that apoptosis of cardiac fibroblasts in the infarct area was exacerbated at post-myocardial infarction (MI) days 7 and 28 compared with the remote area. Scale bars: 100 μm. $n = 4$ in each group.

(B) SA-β-Gal staining demonstrated that spindle-shaped senescent fibroblasts accumulated in the infarct area. Scale bars: 100 μm.

(C) Quantitative reverse transcription polymerase chain reaction analysis showed the post-MI upregulation of senescence-associated genes in the infarct area at post-MI day 7 compared with the no-MI heart (day 0). $n = 4$ in each group. Data represent the mean ± SEM. $*P < 0.05$, $**P < 0.01$, $***P < 0.005$ versus the remote area; [#]$P < 0.05$, [##]$P < 0.01$, [###]$P < 0.005$ versus the no-MI heart; 1-way ANOVA.

Figure 2.

Figure 2. *Nrg1 was upregulated in the infarct myocardium, while cardiac fibroblasts expressed Nrg1 receptors.*
(A) qRT-PCR analysis confirmed the post-MI upregulation of *Nrg1*. Expression levels relative to that of intact heart are presented. *n* = 4 in each group.

(B) Immunohistochemistry demonstrated that the ratio of ErbB4+ fibroblasts (percentage of Thy1+ErbB4+ myofibroblasts/percentage of Thy1+ fibroblasts) was markedly increased in the post-MI heart. Scale bars: 100 μm. *n* = 4 in each group. Data represent the mean ± SEM. *P < 0.05, **P < 0.01, ***P < 0.005 versus the remote area; #P < 0.05, ##P < 0.01, ###P < 0.005 versus the no-MI heart; 1-way ANOVA.

Figure 3.

Figure 3. *M2-like macrophages accumulated in the infarct and expressed Nrg1.*

(A) Immunohistochemistry showed that the accumulation of CD206[+] M2-like macrophages was increased in the infarct area, with a post-myocardial infarction (MI) peak on day 7. Scale bars: 100 μm. $n = 4$ in each group. Data represent the mean ± SEM. *$P < 0.05$, **$P < 0.01$, ***$P < 0.005$ versus the remote area; [#]$P < 0.05$, [##]$P < 0.01$, [###]$P < 0.005$ versus the no-MI heart; 1-way ANOVA.

(B) Flow cytometric analysis confirmed that CD206[+]F4/80[+]CD11b[+] M2-like macrophages were present in normal, no-myocardial infarction (no-MI) hearts and in day 7 post-MI hearts of adult C57BL/6 mice. $n = 6$ in each group.

(C) Microarray analysis demonstrated that M2 after myocardial infarction (M2: MI) had a different expression profile from that of M2: no MI. Scatter plot revealed that 70 genes were upregulated and 39 were downregulated in M2: MI macrophages compared with M2: no MI macrophages. The upregulated genes included neuregulin 1 (*Nrg1*).

(D) Quantitative reverse transcription polymerase chain reaction analysis confirmed the post-MI upregulation of *Nrg1* in CD206[+]F4/80[+]CD11b[+] cardiac M2-like macrophages [M2: MI]. $n = 4$ in each group. Data represent the mean ± SEM. [#]$P < 0.05$, [##]$P < 0.01$, [###]$P < 0.005$ versus M2: no MI; 2-tailed, unpaired Student's *t* test.

(E) Double immunofluorescence staining for CD206[+] M2-like macrophages and neuregulin 1 (Nrg1) demonstrated Nrg1 expression on the surface of CD206[+] M2-like macrophages . Scale bars: 100 μm. $n = 4$ in each group.

Figure 4.

Figure 4. *Bone marrow-derived macrophages attenuated H_2O_2-induced apoptosis and senescence of cardiac fibroblasts via Nrg1 secretion.*

(A) Representative images from phase-contrast microscopy. Treatment with hydrogen peroxide solution (H_2O_2) changed the spindle-shaped appearance to significantly enlarged, flattened morphology. After addition of bone marrow-derived macrophages (BMDMs), fibroblasts returned to spindle-shaped morphology. After addition of ErbB antibody (Ab), fibroblasts displayed the same gross morphology as senescent fibroblasts treated with H_2O_2. Recombinant neuregulin 1 (Nrg1) similarly changed the gross morphology to spindle-shaped. Scale bars: 100 μm. $n = 4$ in each group.

(B) SA-β-Gal staining showed that senescence of fibroblasts was exacerbated in coculture with H_2O_2 but not with H_2O_2 and BMDMs. This suppression of senescence was attenuated by coculture with ErbB Ab. Nrg1 suppressed fibroblast senescence. Scale bars: 100 μm. $n = 4$ in each group

(C) Apoptosis of cardiac fibroblasts (ratio of cleaved caspase-3$^+$DAPI$^+$ fibroblasts to DAPI$^+$ fibroblasts) was increased in the cocultures with H_2O_2 but decreased in those with BMDMs. This decreased apoptosis was eliminated by ErbB Ab. Nrg1 suppressed apoptosis. Nuclei were counterstained with 4',6-diamidino-2-phenylindole (DAPI). $n = 4$ in each group. Scale bars: 100 μm.

(D) Proliferation of cardiac fibroblasts (ratio of Ki-67$^+$ and DAPI$^+$ fibroblasts to DAPI$^+$ fibroblasts) was decreased in the cocultures with H_2O_2 but increased in those with BMDMs. This increased proliferation was eliminated by ErbB Ab. Nrg1 accelerated proliferation. Nuclei were counterstained with DAPI. Scale bars: 50 μm. $n = 4$ in each group. Data represent the mean $\pm$ SEM. $*P < 0.05$, $**P < 0.01$, $***P < 0.005$ versus each group; 1-way ANOVA.

Figure 5.

Figure 5. *BMDMs promote fibroblast activation and collagen synthesis.*

(A-C) Representative images of immunocytochemical staining for (A) vimentin and αSMA, (B) vimentin and collagen I, and (C) vimentin and collagen III.

(A) Activation of cardiac fibroblasts (ratio of vimentin$^+$ and αSMA$^+$ myofibroblasts to vimentin$^+$ fibroblasts) was equable in the cocultures with hydrogen peroxide solution (H_2O_2) but markedly increased in those with bone marrow-derived macrophages (BMDMs). This increased activation was further accelerated by ErbB antibody (Ab). Addition of recombinant neuregulin 1 (Nrg1) did not affect activation of cardiac fibroblasts. Scale bars: 100 μm. *n* = 4 in each group.

(B) Collagen I synthesis was equable in the cocultures with H_2O_2 but significantly increased in those with BMDMs. This increased production was further enhanced by ErbB Ab. Addition of Nrg1 did not affect collagen I synthesis. Scale bars: 100 μm. *n* = 4 in each group.

(C) Collagen III synthesis exhibited a similar tendency as collagen I. Scale bars: 100 μm. *n* = 4 in each group. Data represent the mean ± SEM. *P < 0.05, **P < 0.01, ***P < 0.005 versus each group; 1-way ANOVA

Figure 6.

Figure 6. *PI3K/Akt signaling pathway was associated with BMDM-attenuated apoptosis and senescence of cardiac fibroblasts through Nrg1.*

(A) Representative band of PI3K, pPI3K, Akt, pAkt, and β-actin. Bar graph showed that hydrogen peroxide solution (H_2O_2) alone did not affect activation of PI3K/AKT signaling in fibroblasts. Addition of bone marrow-derived macrophages (BMDMs) significantly activated the signaling pathway, and addition of ErbB antibody (Ab) impaired the activation. adding recombinant neuregulin 1 (Nrg1) stimulated the signaling pathway again. $n = 4$ in each group.

(B) Quantitative reverse transcription polymerase chain reaction analysis of cardiac fibroblasts with or without H_2O_2 / BMDMs / ErbB Ab. The expression levels of senescence-associated genes (*p53, p21, p16*, and *SA-β-Gal*) were increased in the cocultures with H_2O_2 with a time lapse but decreased in the cocultures with BMDMs. Adding ErbB Ab increased these expressions, and adding Nrg1 suppressed them. $n = 4$ in each group.

(C) The expression levels of cell cycle-associated genes (*Cdk4, Cdk6, Cdk2*, and *Ki-67*) were suppressed in the cocultures with H_2O_2 but recovered in those with BMDMs. Adding ErbB Ab decreased these expressions, and adding Nrg1 recovered these expressions. $n = 4$ in each group.

(D) p53 suppressor gene (*MDM2*) was significantly increased in coculture with BMDMs. $n = 4$ in each group.

(E) The expression levels of cell survival-associated gene (*mTOR*) were suppressed in coculture with H_2O_2 but recovered in coculture with BMDMs. Adding ErbB Ab suppressed the expression again, and adding Nrg1 increased it again.

(F) Senescence-associated secretory phenotype-associated gene (*IL-6*) was significantly increased in coculture with BMDMs and after adding ErbB Ab. $n = 4$ in each group. Data represent the mean ± SEM. *$P < 0.05$, **$P < 0.01$, ***$P < 0.005$ versus each group; 1-way ANOVA.

(G) Schematic representation and overview of the Nrg1/PI3K/AKT pathway. Ischemia contributes to the development of cellular damage, leading to senescence, cell cycle arrest, and apoptosis. Nrg1 binding to coreceptor ErbB2/ErbB4 leads to activation of PI3K/AKT and inactivation of p53 and p21. Arrowheads indicate stimulation, whereas hammerheads represent inhibition.

Figure 7.

Figure 7. *In vivo inhibition of Nrg1 signalling promotes apoptosis and senescence of cardiac fibroblasts.*

(A-C) Quantitative reverse transcription polymerase chain reaction analysis showed the post-myocardial infarction (MI) upregulation of (A) apoptosis-associated gene (*Casp3*), (B-C) senescence-associated genes (*SA-β-Gal, p53, p21*, and *p16* in mice after intraperitoneal trastuzumab injection compared with controls. *n* = 4 in each group.

(D) Double immunofluorescence staining for Thy1 and cleaved caspase 3 (CC-3) demonstrated that the ratio of apoptotic cardiac fibroblasts in the infarct area was increased in mice after intraperitoneal trastuzumab injection compared with controls at post-myocardial infarction (MI) days 7, 14 and 28. Arrow shows Thy1$^+$CC-3$^+$ cells. Scale bars: 100 μm. *n* = 4 in each group.

(E) SA-β-Gal staining demonstrated that senescence of cardiac cells in the infarct area was exacerbated in mice after intraperitoneal trastuzumab injection compared with controls at post-myocardial infarction (MI) days 7 and 14. Scale bars: 20 μm. *n* = 4 in each group. Data represent the mean ± SEM. *$P < 0.05$, **$P < 0.01$, ***$P < 0.005$ versus each group; #$P < 0.05$, ##$P < 0.01$, ###$P < 0.005$ versus the no-MI heart; 1-way ANOVA.

Figure 8.

Figure 8. *In vivo inhibition of Nrg1 signalling activates cardiac fibroblasts and exacerbates fibrosis.*

(A) Masson trichrome staining demonstrated that the deposition of collagen fibril was increased in the post-MI infarct area with a time lapse. Intraperitoneal trastuzumab injection significantly increased collagen fibril in the infarct area. Scale bars: 100 μm. $n = 4$ in each group.

(B) Double immunofluorescence staining for Thy1 and αSMA showed increased accumulation and activation of cardiac fibroblasts in the infarct area in the trastuzumab group compared with the control group. Scale bars: 100 μm. $n = 4$ in each group.

(C) Quantitative reverse transcription polymerase chain reaction (qRT-PCR) analysis showed the post-MI upregulation of fibrosis-associated genes (*αSMA, Col1a1,* and *Col3a1*) in mice after intraperitoneal trastuzumab injection mice compared with controls. $n = 4$ in each group. Data represent the mean $\pm$ SEM. $*P < 0.05$, $**P < 0.01$, $***P < 0.005$ versus each group; $^{\#}P < 0.05$, $^{\#\#}P < 0.01$, $^{\#\#\#}P < 0.005$ versus the no-MI heart; 1-way ANOVA.

Figure 9.

Figure 9. *In vivo inhibition of Nrg1 signalling exacerbates myocardial inflammation and promotes accumulation of M2-like macrophages.*

(A) Quantitative reverse transcription polymerase chain reaction analysis showed the post-myocardial infarction (MI) upregulation of senescence-associated secretory phenotype-associated gene (CCl3, *IL-6* and *TNF*) in mice after intraperitoneal trastuzumab injection compared with controls. *n* = 4 in each group.

(B) Immunohistochemistry showed increased accumulation of CD206⁺ M2-like macrophages in the infarct area, with a post-myocardial infarction (MI) peak on day 7. Intraperitoneal trastuzumab injection significantly accelerated accumulation of CD206⁺ M2-like macrophages in the infarct area. Scale bars: 100 μm. *n* = 4 in each group. Data represent the mean ± SEM. *P < 0.05, **P < 0.01, ***P < 0.005 versus each group; $^{\#}P$ < 0.05 $^{\#\#}P$ < 0.01, $^{\#\#\#}P$ < 0.005 versus the no-MI heart; 1-wayANOVA.

Figure 10.

A

Figure 10. Schematic illustration of interrelationship between M2-like macrophages and cardiac fibroblasts.

Ischemic injury contributes to the development of cellular senescence and apoptosis. Senescent cells, including cardiac fibroblasts, show the senescence-associated secretory phenotype (SASP), resulting in activation of M2-like macrophages. M2-like macrophages suppress the senescence and apoptosis of fibroblasts and simultaneously accelerate proliferation. Osteopontin-mediated induction of fibroblasts into myofibroblasts promotes fibrosis (11).