Nuclear retention of Leptin/Upd2 regulates organismal resilience to nutrient extremes

Michelle E. Poling[1, *], Camille E. Sullivan[1, *], Aditi Madan[1, *], Kevin P. Kelly[1, *], Ava E. Brent[1, *], Julien Dubrulle[2] Prashant Raghavan, and Akhila Rajan[1, #].

*equal contribution, # corresponding, and lead author: akhila@fredhutch.org

[1]Basic Sciences Division, Fred Hutch, Seattle, WA, USA.

[2]Cellular Imaging Core, Shared Resources, Fred Hutch, Seattle, WA, USA.

Abstract

Adipokines released from the adipocytes function as a systemic adipometer; they impinge on neural circuits to signal nutrient status. On starvation, adipokines must be retained to signal energy deficit; else, it significantly reduces starvation survival. But how fat cells retain adipokines is unclear. Here, we demonstrate that Atg8, a cell-intrinsic autophagy factor, regulates the starvation-induced acute retention of the Leptin *Drosophila* ortholog Upd2. We show that on starvation, as a direct consequence of Atg8's lipidation, Upd2 accumulates in the nucleus. We illustrate that Upd2's nuclear retention is critical to fat mobilization and increased starvation resilience. In conclusion, we propose that, during starvation, Atg8's role is not just limited to autophagy but is critical for withholding adipokines in the nucleus to promote starvation resilience.

INTRODUCTION

Faced with constant fluctuations in the food supply, organisms rely on accurate nutrient sensing to maintain a steady internal milieu (Leopold and Perrimon, 2007). Regulation of nutrient-sensing can be broadly divided into two categories. One includes cell-intrinsic players such as kinases like AMPK, TOR (Efeyan et al., 2015), and molecular players of the nutrient-scavenging autophagy pathway (Singh and Cuervo, 2011). Other factors that orchestrate systemic cell-extrinsic responses, including the three key metabolic hormones, Insulin, Glucagon, and Leptin (Flier, 2019). For robust energy homeostasis, cell-intrinsic and extrinsic mechanisms must tightly coordinate their roles; but the precise mechanisms underlying such coordination remain to be discovered.

The evolutionary of both cell-intrinsic nutrient-sensing (Neufeld and Baehrecke, 2008; Zirin and Perrimon, 2010) and inter-organ energy homeostasis (Droujinine and Perrimon, 2016; Kim et al., 2021; Rajan and Perrimon, 2013) between flies and mammals are now well-appreciated. In particular, this conservation includes both lipid sensing and its dysregulation, including the development of insulin resistance and 'diabetic' phenotypes (Baker and Thummel, 2007; Gillette et al., 2021; Kim *et al.*, 2021; King-Jones and Thummel, 2005; Musselman and Kuhnlein, 2018; Teleman et al., 2012; Tennessen et al., 2014; van Dam et al., 2020). Hence, discoveries in the *Drosophila* model regarding how endocrine and cell-intrinsic signals interact will broadly be relevant. In this study, using this *Drosophila* system, we investigated how a cell-intrinsic starvation sensing pathway acutely regulates one endocrine factor produced by the adipose tissue.

Adipocytes release hormones to relay fat store status systemically. Leptin in mammals (Ahima et al., 1996; Farooqi and O'Rahilly, 2009) and its functional ortholog in fruit-flies, Unpaired2 (Upd2) (Rajan and Perrimon, 2012), are primary adipokines that are released in proportion to fat stores (Hickey et al., 1996; Kolaczynski et al., 1996a; Kolaczynski et al., 1996b). Both in flies and mammals, Leptin/Upd2 impinges on brain circuits that control energy expenditure, appetite, and overall metabolism (Brent and Rajan, 2020; Ertekin et al., 2020; Friedman and Halaas, 1998; Rajan and Perrimon, 2012). Adipokines convey a permissive signal in surplus nutrient states, indicating that energy

resources can be devoted to costly activities like immunity, sleep, and reproduction. During periods of scarcity, their circulation is reduced, signaling an energy deficit, and enabling organisms to conserve energy.

Starvation-induced reduction in Leptin/Upd2 levels is crucial for an organism's neuroendocrine response to reduced energy reserves. Specifically, Leptin injections during fasting dysregulate neuroendocrine physiology and decrease survival capacity in mice (Ahima *et al.*, 1996). Based on this study (Ahima *et al.*, 1996), Flier and colleagues have proposed that "*the primary physiologic role of Leptin is to provide a signal of energy deficit to the CNS*" (Flier, 2019; Flier and Maratos-Flier, 2017). Similarly, flies with reduced levels of Upd2 display increased starvation survival (Rajan et al., 2017; Rajan and Perrimon, 2012), suggesting that Upd2's retention is critical to enable starvation resilience. But how starvation induces adipokine retention remains to be fully characterized.

In our previous work, we identified that chronic nutrient deprivation (> 5 days) results in a significant reduction in steady-state Upd2 mRNA levels (Rajan and Perrimon, 2012). We also showed that prolonged starvation (> 3 days) dismantles the secretion machinery utilized by Upd2 (Rajan *et al.*, 2017). However, these retention mechanisms operate over a temporal span of days and are not in play within an acute time window (4-8 hours). Therefore, we wondered what other acute mechanisms are utilized by fat cells to retain adipokines.

Autophagy is an acute cell-intrinsic response to nutrient deprivation(He and Klionsky, 2009; Kaushik et al., 2010; Singh et al., 2009a). In adipocytes, autophagy controls lipid metabolism by a process known as lipophagy (Singh *et al.*, 2009a; Singh et al., 2009b) Autophagy-related protein-8 (Atg8) is a central player in the autophagy pathway. Atg8 (LC3/GABARAP in mammals) is a ubiquitin-like protein family member (Kirisako et al., 1999; Shpilka et al., 2011). Atg8 is widely distributed in the nucleus in fed cells (Huang et al., 2015). It translocates to the cytoplasm during starvation, where it is conjugated to the lipid moiety phosphatidylethanolamine (PE)(Ichimura et al., 2000). Lipidated Atg8 regulates the fusion of autophagosomes to lysosomes, resulting in cargo degradation in the cytosol (Nakatogawa et al., 2007). In addition to its well-studied role in autophagy, cytosolic Atg8 regulates other non-degradative cellular activities (Galluzzi and Green, 2019), including but

not limited to phagocytosis (Cunha et al., 2018), viral replication (Reggiori et al., 2010), extracellular vesicle secretion(Leidal et al., 2020), and endocytosis(Heckmann et al., 2019).

In this study, we reveal a novel unconventional role for the nuclear pool of Atg8 in regulating nutrient-state dependent adipokine localization. In fed cells, Upd2 requires nuclear Atg8 for cytosolic localization. On starvation, Atg8 translocates to the cytosol and is lipidated; this depletes Atg8's nuclear availability. As a direct consequence of Atg8's lipidation, we show that Upd2 is withheld in the nucleus. We illustrate that Upd2's nuclear accumulation promotes starvation resilience but increases sensitivity to high-sugar diets. In sum, we uncover an unexpected link between Atg8 and adipokine signaling and illustrate how their interplay controls response to nutrient extremes.

RESULTS

ACUTE STARVATION TRIGGERS UPD2 NUCLEAR ACCUMULATION: To observe endogenous Upd2 localization in *Drosophila* adult adipocytes, we generated CRISPR-engineered tagged knock-ins into endogenous Upd2 genomic locus (See methods). The Upd2-GFP and Upd2-HA genomic knock-ins were able to rescue *upd2*-deletion (*upd2Δ*) fly fat storage defects (Figure S1A), indicating that endogenous genomic tagging (either with HA and GFP) preserved Upd2 function. In place of a robust antibody for immunostaining the Upd2 protein, HA or GFP staining of Upd2 endogenous tagged knock-ins allowed us to follow Upd2's endogenous localization. We observed that within 4 hours of starvation, Upd2-GFP levels increased overall, but especially in the nucleus (Figure 1Ab, and 1A'). This increase in the Upd2 nuclear signal was also observed in the Upd2-HA tag genomic knock-in (Figure S1B), providing a second line of evidence that endogenous Upd2 levels in the adult fly fat nuclei are higher on starvation.

Using another cell system, we sought to corroborate the increased starvation-induced increase in Upd2's nuclear localization that we observed in adult fly fat cells. *Drosophila* S2R+ cells are derived from embryonic stages and have macrophage-like properties (Schneider, 1972). *Drosophila* S2R+ cells respond robustly to amino-acid (AA) deprivation by inducing nutrient deprivation signals (Zacharogianni et al., 2014;

Zacharogianni et al., 2011). Given this, we transiently transfected cells with Upd2-WT::GFP and assayed Upd2-WT::GFP localization in cells cultured in complete media (fed) of in amino-acid (AA) depleted media (starved). Akin to the increase in Upd2 nuclear accumulation in fly fat on acute starvation, we observed a significant increase in Upd2 nuclear accumulation (Figure 1Bb) within 4-8 hours of AA deprivation. Then, we asked whether Upd2's nuclear accumulation was reversible within a short-time frame of refeeding. We noted that when AA-acid starved cells were cultured in complete media within 4-6 hours of culturing cells, Upd2::GFP localizes in punctate cytosolic structures (Figure 1Bc, Bd; See 1B's for quantification). Collectively, studies from both S2R+ cells and fly fat suggested a strong direct correlation between starvation and Upd2 nuclear accumulation.

We wanted to determine if Upd2's starvation-induced nuclear accumulation correlated with its release into extracellular space. Detection of endogenous Upd2 in the adult fly hemolymph, either by western or quantitative ELISA, is a long-standing technical hurdle. We and others have utilized the S2R+cell culture system to quantify Upd2's release into the media reliably (Hombria et al., 2005; Rajan *et al.*, 2017; Wright et al., 2011). Moreover, genome-wide functional screens for protein secretion in S2R+ cells have identified genes relevant to secretion biology *in vivo* (Bard et al., 2006; Wendler et al., 2010). Hence, given that findings from the *Drosophila* S2R+ system regarding secretion in general, and Upd2 secretion in particular, are recapitulated *in vivo*, we utilized this system to investigate the relationship between Upd2 nuclear accumulation and its secretory state.

We measured Upd2's secretion in media of S2R+ cells cultured in complete versus AA-depleted media using a quantitative ELISA for Upd2::GFP (See methods). AA deprivation, within 6 hours, reduced Upd2 secretion by 76%. Notably, Upd2 was restored to normal levels within 5 hours (Figure 1C) when cells were cultured in complete media. We noted the negative correlation between Upd2's nuclear accumulation (4-hours of starvation) and its secretory status. This led us to hypothesize that acute nuclear retention impedes Upd2 secretion. Accordingly, we observed that the knockdown of *Embargoed* (*Emb*) (Fasken et al., 2000), the *Drosophila* CRM1 nuclear export factor, causes increased Upd2 nuclear retention (Figure S1C) and impedes Upd2 secretion (Figure S1D). We cannot

exclude the possibility that a blunt disruption to nuclear export is likely to impact secretion globally. Nonetheless, we noted that Upd2's nuclear accumulation inversely correlated with its secretory potential. Altogether, we concluded that acute starvation reduces Upd2's extracellular release and increases Upd2 nuclear accumulation in diverse contexts of *Drosophila* S2R+ cells and adult fly fat.

UPD2 REQUIRES ATG8 FOR NUCLEAR EXIT: Intriguingly, although Upd2 required CRM1/Emb-based nuclear export (Figure S1D), we were unable to identify a canonical nuclear export signal (NES) on Upd2, suggesting that Upd2's nuclear export occurs indirectly via other protein adapters (See Discussion). To identify clues to what Upd2's protein partners are likely to be, we examined protein motifs in Upd2's protein sequence. We noted that Upd2 had several Atg8-interaction motifs (AIM) (Noda et al., 2010). Given Atg8's well-characterized role in nutrient-sensing, we wondered whether Atg8 might interact with Upd2 to control its nutrient-dependent nucleocytoplasmic localization.

We generated transgenic flies with two-point mutations to Upd2's putative AIM-like sequence (Figure 2A). Multiple AIM-like sequences are found in Upd2, but only one is the canonical 'WXXL' AIM(Noda *et al.*, 2010); hence we mutated that. We specifically expressed Upd2-AIM- (WXXL→AXXA) in fly fat tissue in an *upd2*-deletion background (*upd2Δ; Lpp-Gal4> UAS-Upd2-AIM-::mCherry*) and examined Upd2's localization in the abdominal fat pads of 7 day-old adult male flies, fed *ad libitum* in relation to control transgene (*upd2Δ; Lpp-Gal4> UAS-Upd2-WT::mCherry*). Upd2-AIM- expressing flies displayed a higher nuclear accumulation (2Ab) of Upd2 than controls (2Aa). It is to be noted that the Upd2-WT and Upd2-AIM- transgenes are expressed at similar levels (data not shown). Furthermore, corroborating the role of Atg8 in regulating Upd2's localization, fat tissue-specific, acute temporal knockdown of Atg8 (*upd2GFP-knockin; ppl-Gal4, tubGal80^{ts}> Atg8-RNAi*) resulted in a significant increase in the endogenous Upd2's mean fly fat nuclear intensity (Figure 2B). Thus, disruption of Upd2's putative AIM or knockdown of Atg8 increases Upd2's nuclear accumulation and phenocopies acute starvation.

Next, we wanted to test whether Upd2's intracellular localization in *Drosophila* S2R+ cells, like what we observe for fly fat, is regulated by Atg8. In Drosophila S2R+ cells, we

observed that Upd2-AIM- expression was significantly nuclear (Figure 2Cb, C') and recapitulated the Upd2-AIM- localization in fly fat (Figure 2Ab). Co-immunoprecipitation (Co-IP) experiments, performed in *Drosophila* S2R+ cells, revealed that Upd2-WT could be complex with Atg8, but point-mutations to Upd2's putative AIM (WXXL→ AXXA) impair its ability to complex with Atg8 (Figure S2A). Though these experiments are performed in supraphysiological concentrations, the co-IP complex observations indicate that Upd2 and Atg8 are likely to be complex and that Upd2's AIM sequence is required for this interaction. Furthermore, corroborating Atg8's requirement for Upd2's cytosolic localization, Atg8 knockdown in *Drosophila* S2R+ cells, using two independent dsRNAs, resulted in a significant increase in Upd2-WT::GFP nuclear signal (Figure S2B). These observations lead us the hypothesize that Atg8 regulates Upd2's cytosolic localization during a fed state.

Atg8 knockdown (Figure S2B), or point-mutations to Upd2's AIM (Figure 2C), cause increased Upd2 nuclear accumulation in S2R+ cells. Prior observations suggested that Upd2 nuclear accumulation inversely correlates with its secretory state (Figure 1C, S1D). When we performed quantitative ELISA assays for Upd2 secretion during an Atg8-knockdown (KD) or in Upd2-AIM- state, we observed a significant impairment of Upd2 secretion (Figure S2C and S2D). A previous study identified that Upd2 undergoes GRASP-mediated unconventional secretion(Rajan *et al.*, 2017). In keeping with a role for Atg8 in mediating unconventional GRASP-mediated secretion (Dupont et al., 2011; Duran et al., 2010; Leidal *et al.*, 2020; Zhang et al., 2015), we found using Co-IP experiments in S2R+ cells that Upd2-AIM- ability to interact with GRASP is impaired. In sum, these experiments suggested that Upd2, via its AIM sequence, interacts with Atg8, and this interaction regulates both its cytosolic localization and subsequent extracellular release in fed cells.

Finally, we sought to test whether reconstituting Upd2-AIM- interaction with Atg8 would be sufficient to localize Upd2-AIM- to the cytosol. To test this, we took advantage of the genetically encoded protein binder tag V_HH that acts as a nanobody recognizing GFP (vhhGFP4) (Saerens et al., 2005). Functional studies performed by fusing V_HH to different proteins and then asking how it affects the localization of a GFP-tagged protein have been used to dissect molecular players controlling protein localization (Aguilar et al., 2019; Ressurreicao et al., 2018; Rothbauer et al., 2008). Akin to the experimental design in the

studies mentioned above, which utilized nanobody fusions to probe protein localization, we tested whether reconstituting the interaction between Upd2-AIM-::GFP and V_HH tagged Atg8 is sufficient to restore the cytosolic localization of Upd2-AIM- in fed cells (See Schematic in Figure 2D).

To this end, in *Drosophila* S2R+ cells, we co-expressed with Upd2-AIM-::GFP either the control V_HH::Myc (control) or Atg8-WT was fused to V_HH::Myc (V_HH::Myc-Atg8-WT). As would be expected of a GFP-nanobody, in that it would bind to GFP-tagged proteins, we observed that V_HH::Myc showed increased nuclear accumulation, overlapping with Upd2-AIM-::GFP (See arrow; Figure 2Da). However, co-expression of the control nanobody did not result in localization change of Upd2-AIM-::GFP. Strikingly, in the presence of V_HH-tagged Atg8-WT (V_HH::Myc-Atg8-WT), we observed that Upd2-AIM- GFP is now cytosolic and co-localized with Atg8-WT in punctate structures (See Arrow; Figure 2Db), and reflecting this, Upd2-AIM- nuclear accumulation was significantly reduced (See quantification in Figure 3B'). Hence, reconstituting the physical interaction between Atg8 and Upd2-AIM- is sufficient to localize Upd2-AIM- to the cytosol. This pivotal result demonstrated that Upd2 requires Atg8 for its nuclear exit and cytosolic localization.

ATG8'S LIPIDATION CAUSES STARVATION-INDUCED UPD2 NUCLEAR RETENTION: We were puzzled, however, that despite the presence of endogenous Atg8, on starvation Upd2 nuclear retention phenocopies Atg8's knockdown and Upd2-AIM-. We reasoned that, on acute starvation, Atg8's recruitment to cytosolic vesicles might preclude Atg8 from participating in Upd2's nuclear exit. Consistent with this surmise, pharmacological activation of autophagy, by treatment with autophagy activator Torin (Thoreen et al., 2009), results in a dose-dependent increase in Upd2 nuclear accumulation (Figure S3); this indicated that increased autophagy increases Upd2 nuclear retention. Torin treatment increases Atg8's lipidation (Stolz et al., 2017). Since lipidated Atg8 associates with cytosolic membrane structures (Kirisako et al., 2000; Nakatogawa *et al.*, 2007), this prevents Atg8 from freely diffusing between the nucleus and cytosol (Kraft et al., 2016).

We posited that Atg8's lipidation is likely to reduce its nuclear availability; consequently, Upd2's nuclear exit is impeded, resulting in Upd2's nuclear accumulation.

To test this hypothesis, we generated lipidation-defective Atg8 by mutating to the C-terminal lipid acceptor moiety glycine of Atg8 (Atg8-G→A) (Ichimura *et al.*, 2000). We performed tests on Atg8 G→A to validate that lipidation was defective by using accepted assays in the field to test Atg8 lipidation (Klionsky et al., 2012). This included: i) doublet on western blots of starvation lysates (Figure S4A); ii) recruitment to cytosolic vesicular structures on starvation (Figure S4B). From this, we inferred that Atg8 G→A mutation rendered its lipidation defective (hereafter, it is referred to as Atg8-PE-for simplicity). We fused the lipidation defective Atg8-PE- to an N-terminal V_HH tag (V_HH::Myc-Atg8-PE-; see methods). In subsequent experiments, we utilized V_HH::Myc-Atg8-PE- to test the role of Atg8 lipidation in Upd2's starvation-induced nuclear accumulation.

We hypothesized that lipidation of Atg8 reduces its capacity to participate in Upd2 nuclear exit. Hence, we predicted that reconstituting Upd2-WT::GFP's interaction with lipidation-defective Atg8 should 'rescue' its nuclear accumulation defect (Figure 3A- see schematic). Consistent with this prediction, we found that, when co-transfected with lipidation defective Atg8-PE- (V_HH::Myc-Atg8-PE-), Upd2 is cytosolic even on starvation (Figure 3Af, 3A'). Furthermore, in support of our hypothesis, lipidation competent Atg8-WT (V_HH::Myc-Atg8-WT) does not reduce starvation-induced Upd2 nuclear accumulation (Figure 3Ad; A'). Studies have shown that Atg8/LC3 lipidation increased bulk preventing Atg8/LC3 from shuttling between the nucleus and cytosol (Kraft *et al.*, 2016). Accordingly, we observe that Atg8-WT is localized to cytosolic structures on starvation (Figure S4Bb), but Atg8-PE- does not (Figure S4Bd). Collectively, these experiments (Figure 3A) suggest that, on starvation, Atg8's lipidation reduces its nuclear availability, in turn resulting in increased Upd2 nuclear accumulation.

This observation also suggested that Atg8's lipidation is not required for Upd2's nuclear exit in fed cells. Accordingly, V_HH nanobody-based reconstitution of Upd2-AIM-::GFP with Atg8-PE- (Figure 3Bd), similar to Atg8-WT (Figure 3Bc), is sufficient to re-localize Upd2-AIM-::GFP to the cytosol (Figure 3B, B'). Taken together, these results (Figure 3) strongly suggest that Atg8's lipidation, on starvation, recruits it away from the nucleus to the cytosol. As a direct consequence, Upd2 is withheld in the nucleus.

ATG8'S LIPIDATION IN *DROSOPHILA* ADULT FAT REGULATES OUTCOME ON STARVATION CONTINGENT ON UPD2: The nuclear to cytosolic translocation of Atg8 on starvation is a key step of regulation in the lipidation of Atg8 (Huang *et al.*, 2015). Therefore, we predicted that consistent with mammalian data, our results in *Drosophila* S22R+ cells (Figure S4B), Atg8's nuclear amount *in vivo* in adult fly fat cells will be depleted on starvation. Flies expressing GFP tagged Atg8 in fly fat (*Lpp-Gal4> UAS-GFP::Atg8-WT*) were subjected to an acute starvation diet (4-6 hours 0% sucrose agar) or *ad libitum* fed 30% high sugar diet (HSD) for 14 days. We assessed the total nuclear signal of GFP::Atg8-WT in 3D volume between the flies fed *ad libitum* normal lab food versus starvation and HSD (Figure 4A; Also see Study Design section). As predicted, acute starvation significantly reduced the total nuclear signal of GFP::Atg8-WT, in conjunction with an increase in cytosolic punctate GFP::Atg8-WT (Figure 4Ac, 4A'). Conversely, a surplus diet (30% high-sugar) significantly increased the total nuclear signal of GFP::Atg8-WT (Figure 4Ab, 4A'). In sum, in *Drosophila* adult fly fat Atg8, nuclear signal increases on surplus diet, whereas Atg8 is less nuclear on starvation.

Next, we wondered what happens to nuclear levels of lipidation defective Atg8 in adult fly fat cells on starvation. Whereas the GFP::Atg8-WT (Figure 4Ba) was observed in cytosolic puncta on starvation, Atg8-PE- in the adult fly fat continued to be nuclear on starvation (Figure 4Bb). Specifically, we noted that the GFP::Atg8-PE-nuclear signal was the same between fed and starved states (Figure 4B'; p=0.98). Hence, when Atg8 is lipidation defective, Atg8 is nuclear on both fed and starved states.

Since Atg8-PE- displays a high level of nuclear localization in adult fly fat irrespective of the nutrient state (figure 4B), we predicted that Atg8-PE- would uncouple Upd2's exogenous expression nutrient-responsive nuclear accumulation in fly fat, as it does in S2R+ cells (Figure 3A). To test this prediction, when we acutely expressed GFP::Atg8-WT or GFP::Atg8-PE- in adult fly fat (*ppl-gal4>TubGal80ts>UAS-Atg8-WT or UAS-Atg8-PE-*), and then assessed what happens to endogenous Upd2's nuclear localization on starvation [*upd2-HA (genomic knock-in); ppl-Gal4tubGal80ts>UAS-Atg8-WT or Atg8-PE-*]. When Atg8-WT is acutely expressed, endogenous Upd2's nuclear levels increase starvation within 4 hours (Figure 4C; p=1.0e-09). However, when we express Atg8-PE- acutely, the amount of nuclear Upd2 decreases on starvation (Figure 4C; p=0.003). Hence, the presence of

exogenous lipidation defective Atg8 uncouples Upd2's nuclear levels from the systemic nutrient state *in vivo*.

Atg8-PE- expression prevented Upd2 nuclear accumulation (Figure 4C), and we wondered whether this impacted organismal starvation resilience. For conducting whole animal physiology experiments, we could not recover any viable progeny when we

expressed Atg8-PE- in an Atg8 mutant background (Figure S5A). Since the expression of the Atg8-WT and Atg8-PE- transgenes were comparable on starvation (Figure S5B), we took the approach of over-expressing Atg8-WT or Atg8-PE- in fly fat and assessing systemic effects

(See Study Design section below). First, we tested the ability of flies to mobilize fat stores and noted that flies expressing Atg8-WT (*Lpp-Gal4> UAS-Atg8-WT*) were able to break down their fat reserves on starvation (Figure 4Da; 50% mobilization; p=0.04). But flies with an

exogenous expression of lipidation-defective Atg8-PE- in fly fat (*Lpp-Gal4> UAS-Atg8-PE-*) were unable to mobilize their TAG stores on starvation (Figure 4Da); this correlated with the reduced starvation resilience of fat specific expression of Atg8-PE- (Figure 4Ea).

Next, we went one step further and predicted that the inability of flies with fat-specific Atg8-PE- over-expression to mobilize TAG on starvation is at least in part due to Upd2's cytosolic localization. If this prediction were to hold up, we should observe that the

negative effects of Atg8-PE- over-expression are mitigated in the absence of Upd2. Consequently, we performed the TAG mobilization and starvation survival experiments in an *upd2-deletion* background (*upd2Δ; Lpp-Gal4> UAS-Atg8-PE-*). In line with our prediction,

we found that exogenous expression of Atg8-PE- in fly fat of *upd2Δ* flies led to TAG mobilization (Figure 4Db) and starvation resilience (Figure 4Eb) at levels comparable to Atg8-WT transgene expression. In sum, these results support a model that Atg8's lipidation,

in addition to its critical role in autophagy, is required for Upd2 retention; this, in turn, determines how an organism survives prolonged starvation.

UPD2'S NUCLEAR RETENTION IS CRITICAL FOR STARVATION-INDUCED FAT MOBILIZATION AND HUNGER-DRIVEN FEEDING: To further assess what happens when the inverse correlation between nutrient state and adipokine nuclear accumulation is genetically perturbed. We subjected

flies expressing control transgene, Upd2-WT or Upd2-AIM- specifically in fly fat cells, to a

high sugar diet (HSD) regime (See Methods and Study Design sections). We found that over-expression of Upd2-AIM- renders flies sensitive to an HSD regime (Figure 5A), suggesting holding Upd2 in the nucleus during a surplus diet where Upd2 is less nuclear (Figure S6A) reduces HSD survival. Conversely, we predicted that genetically 'holding' Upd2 in the nucleus during starvation, a state where endogenous Upd2's nuclear accumulation increases (Figure 1A), will allow flies to survive starvation. Consistent with our prediction, the over-expression of Upd2-AIM- in fly fat cells (*upd2Δ; Lpp-Gal4> UAS-Upd2-AIM-*) on starvation provided resilience; Upd2-AIM- flies survived significantly longer than control or Upd2-WT overexpression (Figure 5B). Overall, this suggested that the inverse correlation between adipokine nuclear accumulation and nutrient state is important.

We wondered whether retention of Upd2 within the cell, but not specifically in the nucleus, would be sufficient to have the same systemic effects on starvation resilience (Figure 5B). To tease apart this nuclear versus cytosolic retention, we examined a state in which Upd2 was withheld in the cytosol but not released. To achieve this, we ectopically localized Upd2-AIM- to the cytosol by appending a constitutive nuclear export signal (NES) to Upd2-AIM- (See Methods). Expressing cytosolic Upd2-AIM- in the fly fat (*upd2Δ; Lpp-Gal4>Upd2-NES+-AIM-*), as predicted, was sufficient to re-localize Upd2-AIM- to the cytosol (Figure S6B; See arrows), whereas Upd2-AIM- expressing flies displayed no cytosolic localization (Figure S6B & Figure 2). Consistent with the role of Atg8 in mediating Upd2's secretion, we found that, despite being localized to the cytosol (Figure S6D), Upd2-NES+-AIM- was not secreted (Figure S6E). Significantly, however, we noted that fat-specific expression of Upd2-NES+-AIM- (*upd2Δ; Lpp-Gal4>Upd2-NES+-AIM-*) did much worse on starvation than control flies (Figure S6C). These results suggested that it is *not just retention within the cell, but specifically Upd2's retention in the nucleus* critical for exerting the systemic effects on starvation resilience.

We had previously shown that overexpression of Upd2-WT transgene in fly fat renders it incapable of mobilizing TAG reserves on starvation (Figure 2G in Rajan and Perrimon, 2012). Hence, we wanted to assay TAG mobilization in flies over-expressing Upd2-WT versus Upd2-AIM- specifically in fly fat (*upd2Δ; Lpp-Gal4> UAS-Upd2-WT/ AIM-*). We found that while Upd2-WT flies were unable to mobilize TAG on starvation (Figure 5C;

Upd2-WT Fed vs. Stv p=0.6), Upd2-AIM- flies could mobilize TAG stores (Figure 5C; Upd2-AIM- Fed vs. Stv; p=0.01). Notably, the increased TAG mobilization of fat-specific Upd2-AIM-expressing flies concurred with increased starvation resilience of Upd2-AIM- expressing flies (Figure 5B).

Next, we assayed whether Upd2-AIM- displayed different feeding behavior post-starvation than Upd2-WT over-expression flies. Expressing Upd2-WT transgene, specifically in fly fat (*upd2Δ; Lpp-Gal4> UAS-Upd2-WT*), displayed no significant increase in hunger-driven feeding, as measured by the FLIC assay (Figure 5D, also see methods); this is consistent with Upd2's role as a satiety signal. Conversely, Upd2-AIM- fat-specific over-expression flies (*upd2Δ; Lpp-Gal4> UAS-Upd2-AIM-*) displayed a significant increase in hunger-driven feeding (Figure 5D; Upd2-AIM- Fed vs. Starved; p=0.002). These findings suggest that nuclear accumulation of Upd2 on starvation exerts systemic effects; it promotes starvation resilience, fat-mobilization, and hunger-driven feeding.

In conclusion, our results suggest that we have revealed a specific intersection node between cell-intrinsic and extrinsic mechanisms in regulating an organism's response to nutrient extremes. We demonstrate that Atg8 regulates nutrient state-dependent localization of the *Drosophila* Leptin, Upd2 in both fed and starved cells (Summary of findings in Figure 6). Upd2's nuclear accumulation on starvation, as a direct result of Atg8's lipidation, is important for organismal adaptation to nutrient deprivation. Thus, we propose that Atg8's role in the starvation response is not just limited to autophagy but also is required for adipokine retention. This process is critical for an organism to sense and adapt to starvation.

DISCUSSION:

A NOVEL NON-CANONICAL ROLE FOR ATG8 IN NUTRIENT-STATE DEPENDENT PROTEIN LOCALIZATION:

Upd2 is a 47 kDa protein. Given the structural restraint of nuclear pore complexes (NPC) to only permit proteins <40kDA to pass through (Davis, 1995), Upd2 is likely to require active nucleocytoplasmic (NC) transport mechanisms that enable transit through the NPC (Kaffman and O'Shea, 1999). Consistent with the role for classic nuclear export being required, we find that Embargoed/CRM1 is required for Upd2's nuclear exit (Figure S1C).

But, using the NETNES mapper tool (la Cour et al., 2004) did not identify a canonical NES in Upd2. Hence, we surmised that other adapter proteins interacting with Upd2 might promote its nuclear export. Unexpectedly, we found that mutations to Upd2's AIM or KD of Atg8 phenocopies knockdown of nuclear export protein Emb (Figure 2, S2). This suggested to us that Atg8 is required for Upd2's cytosolic localization in fed cells. Consistent with this nanobody-based reconstitution of the physical interaction between Upd2-AIM-, that is only localized to the nucleus, and Atg8 is sufficient to re-localize Upd2 to the cytosol (Figure 2D, 3B, 3B').

Our study does not yet resolve how Upd2 localizes to the nucleus in the first place. But, based on the observations that Upd2 is nuclear when Atg8 is knockdown and when its AIM is disrupted, we conclude that Atg8 does not regulate Upd2's nuclear entry. Analysis of Upd2's sequence using a bioinformatic tool- classic NLS (cNLS) mapper (Kosugi et al., 2008; Kosugi et al., 2009a; Kosugi et al., 2009b), predicted that Upd2 contains an NLS at residues 141-150 (GRVIKRKHLE) with a score of 10.5. According to the cNLS mapper, a GFP reporter protein fused NLS, with a score of 10, is exclusively localized to the nucleus. Hence, we surmise that given the strong NLS, Upd2's default state is nuclear. Then, when Upd2' licensed' by cofactors like Atg8, it would Upd2 transits to the cytosol.

We considered the alternate hypothesis that Atg8 cytosolically tethers Upd2 to prevent nuclear entry. But if that were the case, we would expect that Atg8, which is highly cytosolic on starvation, would tether Upd2 to the cytosol. Given that Atg8 nuclear depletion, within 4 hours of acute starvation (Figure 4A), coincides with increased Upd2 nuclear accumulation (Figure 1A), we favor the hypothesis that the nuclear pool of Atg8 in fed cells facilitate Upd2 nuclear exit.

Further work will be needed to resolve all the events leading to Upd2's Atg8-mediated nuclear exit. We surmise that Atg8, a ubiquitin-like protein (Shpilka et al., 2011), might operate akin to other ubiquitin and ubiquitin-like proteins that have been shown to function in cellular localization (Jura et al., 2006; Nakagawa and Nakayama, 2015; Xu et al., 2010). Specifically, SUMO (small ubiquitin-like modifier) that positions its target protein (RanGAP1) to the NPC to increase the efficiency of its translocation (Mahajan et al., 1997). Similarly, Atg8 might position adipokine Upd2 to enable efficient export via the classical

nuclear export pathway or may serve to direct Upd2 to other nuclear export mechanisms such as nuclear budding (Speese et al., 2012).

How widespread this role for Atg8 in nucleocytoplasmic protein localization remains resolved. But it is quite plausible that other proteins, such as ribophagy receptor NIUFIP1 (Wyant et al., 2018) with 4 AIM sequences, and localizes to cytosol on starvation, might utilize Atg8 for context-dependent NC shuttling. Furthermore, insulin retention on acute starvation is a critical response in flies and mammals (Geminard et al., 2009). It will be important to explore whether Atg8 mediated retention, like Leptin/Upd2 in fat, also applies insulin in pancreatic beta cells. In sum, Atg8's controls context-dependent nucleocytoplasmic localization could be another layer of regulation in controlling protein localization during starvation and merits further investigation.

Atg8 controls the two step-process for Upd2 release in fed cells – 1: nuclear exit and 2: GRASP-mediated extracellular- release (Figure S2D, S2E). This study shows that lipidation-defective Atg8 (VHH-Myc::Atg8-PE-) can localize Upd2-AIM- to the cytosol (Figure 3Bd), but we do not know whether the lipidation defective Atg8 can restore Upd2-AIM-'s secretion defect. Debnath and colleagues have comprehensively delineated how Atg8/LC3 lipidation is critical to Atg8's role in extracellular vesicle secretion, a process that they term LDELS (Leidal *et al.*, 2020). Based on their elegant study (Leidal *et al.*, 2020), we surmise that though Upd2 can transit to the cytosol in the presence of lipidation-defective Atg8, but that it is not secreted. Further studies will be needed to resolve whether Upd2 undergoes an LDELS secretion process in *Drosophila* fat cells.

WHAT IMPLICATIONS DO DROSOPHILA UPD2 STUDIES HOLD FOR MAMMALIAN LEPTIN BIOLOGY?

Leptin does have two putative AIM-like sequences. But unlike Upd2 (47kDa), Leptin (15 kDa) is unlikely to require facilitated nuclear export. Even if Atg8-mediated nucleocytoplasmic shuttling does not apply to mammalian Leptin, it is possible that, like Upd2, Leptin utilizes a secretory pathway mediated by Atg8. This is not a far-fetched possibility given that human Leptin, just like Upd2, requires unconventional secretion (Rajan *et al.*, 2017). Hyperleptinemia, i.e., high levels of circulating Leptin, has been identified as a primary cause of 'leptin resistance'- a state wherein Leptin cannot effectively signal satiety state to

the brain (Knight et al., 2010). This resulted in dysfunctional feeding behavior and decreased energy expenditure. Thus, our report that Atg8's lipidation retains *Drosophila* Leptin/Upd2 represents a novel avenue to develop interventions to improve Leptin sensitivity.

In conclusion, the study on how *Drosophila* Leptin- Upd2 – is acutely retained in fly fat cells on starvation has led to numerous unanticipated insights, raises many questions that should stimulate work both in invertebrate and mammalian systems.

STUDY DESIGN- LIMITATIONS AND CONSIDERATIONS:

Atg8: In *Drosophila,* clonal analysis of larval fat tissues is the standard for studying how autophagy (Atg) core pathway genes affect cell-intrinsic processes (Takats et al., 2019). While clonal analysis addresses is an elegant genetic tool to study cell-intrinsic functions of Atg genes, it is not compatible with experimental design to study whole animal physiology. Therefore, for the physiology assays, in which we assess how Atg8's lipidation in fat affects TAG mobilization and starvation resilience (Figure 4D, 4E), we have compared the effect of Atg8-WT transgenic over-expression with the Atg8-PE- in fly fat. The reason for this design is two-fold. The first is to overcome the hurdle of studying a specific function of Atg8, its lipidation, which is critical to animal survival. We note that lipidation-defective Atg8 flies do not survive (Figure S5A). Hence the only viable strategy to study the effect of defective Atg8 lipidation is to utilize fat tissue-specific expression of Atg8-PE- and compare its effects to Atg8-WT. It is important to note that we compare the effects of two transgenes -Atg8-PE- and Atg8-WT- head-to-head. These transgenes differ in a single amino acid (the penultimate glycine of Atg8) and are expressed at comparable levels even on starvation (Figure S5B).

Furthermore, these results are to be viewed in conjunction with the V_HH-nanobody based reconstitution of interactions between Atg8-PE- and Upd2, where Atg8-PE- increases Upd'2 cytosolic exit, even on starvation, despite the presence of endogenous Atg8 (Figure 3A). Secondly, our goal is to uncover the systemic effects of only altering Atg8's lipidation status in the fat. Since Atg8 has roles in every tissue, the most feasible design is to use the tissue-specific expression of lipidation-defective Atg8 and assess how it impacts whole organism physiology in relation to Atg8-WT. Hence, within the limitations of the system, we have

utilized internally controlled experimental approaches to assess the effect of Atg8's lipidation status on endogenous Upd2 localization and whole animal physiology.

3   For experiments to study the effect of nutrition on protein-localization (Figure 4A and B), we utilized the UAS/Gal4 system for specifically expressing Atg8-WT and Atg8-PE-transgenes in the fly fat. This design of driving protein expression downstream of fat-specific

6   Gal4 allows us to uncouple transcription and translation-dependent effects of nutrient state on Atg8 mRNA and protein levels. Hence, we quantify the nuclear GFP signal in fat cells in 3D-volume, representing Atg8 levels. Furthermore, we note that on starvation, both

9   Atg8-WT and Atg8-PE- transgenes show a similar fold change in gene expression (Figure S5B). Nonetheless, Atg8-WT is cytosolic on starvation (Figure 4A), but Atg8-PE- is nuclear (figure 4B). This lends credence to our claim that we follow the nutrient state effect on

12   protein localization rather than gene expression.

Upd2: Upd2 is the functional fly ortholog of human Leptin (Rajan and Perrimon, 2012) and plays a fat-specific role in regulating the systemic nutrient state. But, as shown by multiple

15   independent groups, Upd2 exerts systemic effects by signaling from other organs and, specifically, coupling gut metabolic state to the olfactory system(Cai et al., 2021) or acting in hemocytes to regulate immunity (Chakrabarti et al., 2016). We have utilized fat-tissue-

18   specific drivers to distinguish Upd2's adipokine-specific function from its other organ-specific roles. *In lieu* of the ability to perform longitudinal diet-related experiments (10-40 days) with a knockdown of Atg8 in fat, which is lethal over seven days, we utilized Upd2-

21   AIM- over-expression in fly fat. Upd2-AIM- phenocopies Atg8-KD (Figure 2A, 2B). Using this design, we have assessed the systemic effects of various Upd2 transgenes (*Upd2-WT, Upd2-NES+-AIM-and Upd2-AIM-*). Given that these transgenes are all inserted in the same

24   genome site (attP40) and have similar expression levels, it allows for interpretation of the specific mutation to Upd2's Atg8 binding domain means. One valid reservation is that what we reveal are gain-of-function effects of Upd2-AIM-. Even if that were to be the case, it is

27   striking that we don't see these gain-of-function effects on starvation resilience when we retain Upd2-AIM- in the cytosol (*Upd2-NES+-AIM-)* (Figure S6A, S6B)*;* this suggests that retention of Upd2 in the nucleus on starvation is biologically meaningful. Hence, within the

limitations of the system, we have utilized internally controlled experimental approaches to assess the effect of Upd2's nuclear localization on whole animal physiology.

ACKNOWLEDGMENTS: We are grateful to the late Suzanne Eaton for the Lpp-Gal4 flies. We thank Laura Holderbaum for assistance with cell culture and immunoprecipitation experiments. Zachary Goldberg for technical assistance with setting up the FLIC assay. We thank Alyssa D. Dawson and Dolores Covarrubias at the Fred Hutch Genomic Shared resources facility for assistance with RNAseq. We are grateful to Feinan Wu at the Fred Hutch Bioinformatics core for RNAseq analysis. We thank David Strutt for sharing sequences for designing the V_HH tagged constructs. We thank Susan Parkhurst for her advice on the NES sequences and critical comments on the manuscript. Funding: This work is possible due to grants awarded to AR from NIGMS (GM124593) and New Development funds from Fred Hutch. KPK was supported by the NIH Chromosome Metabolism and Cancer Training Grant (T32CA009657) and is currently supported as an NSF Post-Doctoral Research Fellow (NSF Award #2109398). Genomic reagents from the DGRC, which is funded by NIH grant 2P40OD010949, were used in this study. Stocks obtained from the Bloomington Drosophila Stock Center (NIH P40OD018537) and Transgenic RNAi Resource project (NIGMS R01 GM084947 and NIGMS P41 GM132087) were used in this study. Author contributions: Conceptualization: AR, PR; Investigation: MEP, CES, AEB, KPK, AM, AR; Formal analysis, Data Curation: MEP, CES, AEB, KPK, AM, AR; Visualization: MEP, CES, AEB, KPK, AM, AR, JD; Supervision: AR; Writing- original draft: AR; Writing- review and editing: PR, KPK, AM. Funding acquisition: AR. Competing interests: Authors declare no competing interests.

RESOURCE AVAILABILITY

Lead Contact

Requests for further information, reagents, and resources should be directed to and fulfilled by the Lead Contact, Akhila Rajan (akhila@fredhutch.org).

Materials Availability

Drosophila strains generated in this study are available from the corresponding author, Akhila Rajan (akhila@fredhutch.org).

Data and Code Availability

The datasets generated in this study are available from the corresponding author, Akhila Rajan (akhila@fredhutch.org).

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Experimental Animals: *Drosophila melanogaster*

Only males were used in experiments at 5-10 days post-eclosion. Flies were cultured in a humidified incubator at 25°C with a 12h light-12h dark cycle and were fed a standard lab diet, containing per liter: 15 g yeast, 8.6 g soy flour, 63 g corn flour, 5g agar, 5g malt, 74 mL corn syrup. For acute starvation, adult male flies, seven days old, were subjected to 4-hour starvation on 0% sucrose in agar at 29°C. For survival curves, flies were subjected to 1% sucrose in an agar diet. For RNAi experiments, flies were raised at 18°C until seven days post-eclosion, after which they were shifted to 29°C for seven days.

The following fly strains used in this study were from our previous work (Rajan *et al.*, 2017; Rajan and Perrimon, 2012) and obtained from other investigators or the Bloomington *Drosophila* stock center (BDSC): upd2Δ3-62 (upd2Δ; (Hombria *et al.*, 2005)), *Lpp-Gal4* (Brankatschk and Eaton, 2010), Atg8-Trojan-Gal4 ((Lee et al., 2018); BDSC 77836). The control UAS strain used for over-expression experiments is UAS-Luciferase (Rajan and Perrimon, 2012). RNAi lines from the TRiP facility at Harvard Medical School (http://www.flyrnai.org/TRiP-HOME.html) include: Luciferase-RNAi (JF01801), Atg8-RNAi #1 (JF02895, BDSC 28989), Atg8-RNAi #2 (HMS01328, BDSC 34340).

The following UAS lines were generated for this study: *UAS-Upd2::mCherry, UAS-Upd2-AIM-::mCherry, UAS-GFP::Atg8; UAS-GFP::Atg8-PE-*. All transgenes of the same gene were inserted at the same att site to control expression levels—either attp40 or attP2. For RNAi experiments and temporal over-expression in Figure 2B and Figure 4C, flies were generated with the following genotype: *Upd2-crGFP; Ppl-Gal4, TubGal80ts* was crossed to Atg8-RNAi, Luc-RNAi for 2B; For 4B and 4C and *UAS-GFP::Atg8-WT* and UAS-GFP::Atg8-PE-. The

parents of this cross (F0 generation) were maintained at 18c for both the strains, F1s were allowed to eclose at 18c. A week after eclosion, the F1s were moved to 29c for 5-7 days before dissection and staining.

The ability of UAS-GFP::Atg8 and UAS-GFP::Atg8-PE- to rescue the loss of Atg8 function was tested by introducing Atg8 variant transgenes into the Atg8-Trojan-Gal4 line (Lee *et al.*, 2018) in which insertion of the Trojan-Gal4 into the Atg8 locus generates a lethal allele. We confirmed the presence of Gal4 by crossing to UAS-GFP.

Cell lines

Drosophila S2R+ cells were used for all cell culture-related experiments. This cell line was chosen because previous studies have validated their applicability to study autophagy 7-9 and protein secretion. The cells were maintained in Schneider's medium (GIBCO), 10% heat-inactivated FBS (SIGMA), and 5% Pen-Strep (GIBCO) at 25°C. For starvation, cells were incubated in Schneider's Insect Medium without Amino Acids (United States Biological, Cat# S0100-01.10) for the indicated amount of time.

METHOD DETAILS:

Cloning and Transgenic Flies

All cloning was done using the Gateway® Technology. For Atg8, entry cDNA was PCR amplified from Atg8 cDNA (clone LD05816 DGRC-Gold collection) and cloned into pENTR-D/TOPO using BP reaction (Gateway® BP Clonase II enzyme mix, Cat#11789-020, Invitrogen). For *Drosophila* Upd2 variants, pENTR-Upd2 (Rajan *et al.*, 2017) made for previous work in our lab was used. For site-directed mutagenesis of putative AIM sites, pENTR-Upd2 was mutagenized to convert tryptophan and leucine encoding codons to alanine. For the addition of a canonical NES+, a sequence corresponding to amino acids LQELLELLRL (Verboon et al., 2015) was inserted after the start codon in either pENTR-Upd2-WT or pENTR-Upd2-AIM-. For site-directed mutagenesis of the Atg8 lipidation site, pENTR-Atg8 was mutagenized to convert its penultimate Glycine to Alanine. All mutagenesis was done using the Q5® Site-Directed Mutagenesis Kit from NEB (Cat # E0554S). The sequence of oligonucleotides used for this mutagenesis reaction is available

on request. The entry vectors were then moved using LR clonase reaction (Gateway® LR Clonase® II Enzyme mix, Cat#11791-020, Invitrogen) into destination vectors compatible with fly transformation, or cell culture with the appropriate C-terminal tags for Upd2 and N-terminal tags for Atg8. Transgenic flies were generated by using the microinjection service provided by Bestgene Inc.

For generating the V_HH tagged constructs, the sequence for V_HH tag and appropriate linkers, utilized in a prior study by Strutt and colleagues (Ressurreicao *et al.*, 2018), was appended to either Myc (control: V_HH-Myc), Atg8-WT (V_HH-Myc::Atg8-WT), or Atg8-PE-(V_HH- Myc::Atg8-PE-). Double-stranded sequences for the three constructs (available on request) were synthesized using Codex DNA Inc. Synthetic DNA platform and cloned into PDONR221 using BP clonase (Gateway™ BP Clonase™ II Enzyme mix; Catalog number: 11789020, Invitrogen). Then, to generate expression clones, the PDONR vectors were cloned into Expression competent S2 cell culture vectors from the Murphy Collection available from DGRC using LR clonase.

Generation of C-terminal tagged Upd2 CRISPR lines:

GFP or HA-tagged Upd2 CRISPR lines were developed in collaboration with WellGenetics Inc. using modified methods of Kondo and Ueda (Kondo and Ueda, 2013). In brief, the following gRNA sequence was cloned into U6 promoter plasmids for HA lines: TCCAATGAGTCTTGAGCCCT[CGG]/GCCGAGGGCTCAAGACTCAT[TGG]. Cassette 3xHA RFP containing 3xHA and 3xP3 RFP and two homology arms were cloned into pUC57 Kan as donor template for repair. upd2/CG5988 targeting gRNAs and hs-Cas9 were supplied in DNA plasmids, together with donor plasmid for microinjection into embryos of control strain w1118. F1 flies carrying selection markers of 3xP3 RFP were further validated by genomic PCR and sequencing. CRISPR generates a break in upd2/CG5988 and is replaced by cassette 3xHA RFP. 3XP3 RFP, which facilitates the genetic screening, was flipped out by Cre recombinase. RE loxP RE (46 bp) remains after excision between Stop codon and 3'UTR. For GFP lines, gRNA sequence TCCAATGAGTCTTGAGCCCT[CGG] was used.

Tissue culture:

Before transfection, cells were passaged to 60-80% confluency. Cells were cultured in 96 well plates for transfections related to ELISA experiments and

coimmunoprecipitations in 6-well plates. For ELISAs, cells were transfected with 20ng/well of pAc-upd2::GFP 3 or indicated Upd2 variants, 10ng/well pACRenilla::Luciferase, and 150ng of dsRNA/well for knockdown experiments. For coimmunoprecipitations, cells were transfected with 200ng/well of the indicated plasmid.

For imaging experiments, cells were seeded on poly-D-lysine coated 8-well chambered culture slides for fixed imaging (MatTek CCS-8). The T7 flasks, which were 100% confluent, were used for seeing cells at 1:10 dilution at 400µl per well of the 8-well chamber dish or 1:20 dilution at 200µl per dish for FluoroDish. Transfections were done with plasmids indicated. For 8-chamber slides, 100ng/well of each plasmid was transfected. Transfections were done using the Effectene kit (Cat# 301427, QIAGEN) per the manufacturer's instructions.

dsRNA production and cell treatments:

Amplicons for dsRNAs were designed using the SnapDragon dsRNA design tool (https://www.flyrnai.org/snapdragon) and in vitro transcribed (IVT) using MEGAscript® T7 Transcription Kit (Cat# AMB1334-5, ThermoFisher). IVT reactions were carried out as per the protocol provided by the DRSC (available at: http://www.flyrnai.org/DRSC-PRS.html). The sequence of amplicons used in this study can be found in the table below. LacZ dsRNA was used as controls. All dsRNA knockdown experiments were carried out using two independent dsRNAs per gene. We found that this produced a knockdown efficiency of >85% (based on qPCR analysis) in S2R+ cells. S2R+ transfected with dsRNAs were incubated for four days to allow for gene knockdown. On the 4th day, the media was changed, and the ELISA assay was carried out on the 5th day. Note that the data is represented as percent change in Upd2/Leptin secretion normalized to transfection efficiency with 0% change indicating baseline secretion level. See the ELISA assay procedure below.

Primer sequences used for dsRNA production:

Target	Primer name	Sequence 5' - 3.'
LacZ	LacZ F1	TTTCCAGTTCCGTTATCCG

	LacZ R1	CAAAAATCCATTTCGCTGGT
Atg8a	Atg8a F1	GTACAGGGAGCCCATGGTA
	Atg8a R1	CGAGAAGGCTCCCAAGG
Atg8a	Atg8a F2	GCCCTGCGTATCAGATCAAT
	Atg8a R2	CGATGAGAATGTTTACGGCA
Emb	Emb F1	AGAGGTACATGCAACTGCCC
	Emb R1	GTTGGAACGAGACACCCACT
Emb	Emb F2	ACTAAGGCCAAGCACCTGAA
	Emb R2	TGAACCATCGTGTCCTTGAA

Treatment of cells with drugs:

For drug treatment experiments, the media was replaced with media containing the drug on day 3 after transfection with upd2::GFP. 4 hours later; the conditioned media was used for ELISA or imaging. Drugs used in this study include Torin1 (Cat# ab218606, Abcam). Stocks solutions of the drugs were made in DMSO as per the manufacturer's instructions and used at a working concentration indicated in the Figure S3 legends. DMSO treated replicates were used as controls. Note that for ELISAs, the data is represented as percent change in Upd2 secretion normalized to transfection efficiency, with 0% change indicating baseline secretion level.

ELISA assays:

GFP sandwich ELISA assay was used for detecting Upd2::GFP. On day 1, 96 well medium bind polystyrene plates (Cat#CLS3368-100EA, Sigma) were incubated overnight at 4C with coating antibody (Cat# ACT-CM-GFPTRAP, Allele Biotechnology) diluted in 0.01M pH8.0 bicarbonate buffer at a concentration of 1µg/ml. On day 2, plates were washed briefly with PBS, blocked for 30 minutes with 1% BSA block in PBS, coated with conditioned media, and incubated overnight at 4C. Recombinant GFP protein (Cat# MB-0752, Vector labs), diluted in S2R+ cell growth media (64ng/µl to 4 ng/µl), was used in every ELISA plate as a positive control to ensure linearity of GFP readings. On day 3, the plates were washed with

PBS+0.05% Tween-20 (PBS-T), blocked with 1% BSA in PBS for 30 minutes at RT. GFP detection antibody (Cat# 600-401-215, Rockland) was added to the diluted 1:1000 in 0.1% BSA in PBS-T. Plates were washed with PBS-T and incubated with secondary HRP conjugated anti Rabbit secondary antibodies (Cat# ab136636, Abcam) diluted at 1:5000 in 1% BSA block. Plates were washed in PBS-T with a final wash in PBS. For detection, each well was incubated 100µl 1-step Ultra-TMB ELISA substrate (Cat# 34028, Pierce), which was previously equilibrated to RT, for approximately 5-15 minutes until detectable blue colorimetric reaction occurred. The reaction was stopped with 2N sulphuric acid, and absorbance was measured at 450nm. The TMB readings were normalized to transfection efficiency as measured from Renilla Luciferase assays (see below).

Renilla Luciferase assay:

On day 2 of the ELISA assay, after the conditioned medium was transferred for use in ELISA assays, cells were re-suspended in 50µl of PBS and incubated with 50µl/well Renilla-Glo® Luciferase reagent (Cat# E2710, Promega) for 10 minutes and read using a multiwell luminometer.

Quantification for ELISA assays:

For ELISA assays, the ELISA signal readings are normalized to transfection efficiency; the data is represented as percent fold change from control used as the baseline. Specifically, the ratio of TMB readings to Renilla Luciferase readings is calculated. This ratio from the control is used as a baseline, and the data is represented as the percent fold change of experimental conditions with respect to the control. A 2-tailed t-test quantified statistical significance on 6 biological replicates per condition. Error bars represent %SD (Standard Deviation). For TAG analysis, statistical significance was quantified by a 2-tailed t-test on 3-6 biological replicates per condition. Error bars indicate %SD (Standard Deviation). For qPCR analysis, normalized gene expression levels were analyzed using one-way ANOVA. Error bars represent the standard error of the mean (SEM).

Immunoprecipitation and Western blots:

For Immunoprecipitation (IP) from S2R+ cells, protein for each condition was prepared by lysing 1 well of a 6-well dish, 4 days after transfection. 2mg/ml was used per IP experiments performed with camelid antibodies GFP-Trap Magnetic Agarose (Cat# gtma-

20, Chromotek); RFP-Trap Magnetic Agarose (Cat# rtma-20, Chromotek); Myc-Trap magnetic agarose (Cat# ytma-20; ChromoTek); as per the manufacturer's protocol.

Western blots were performed detailed in our prior publication 1. Antibodies used include Chicken anti-GFP (Cat#ab13970, Abcam) 1:2000 in TBS-0.05% tween-20 (TBS-T), Rabbit- anti Upd2 [custom antibody AR4398 generated at YenZym Inc., to Upd2 peptide 318-334: RRPRRNSAERRHLAAIHC; antibody was verified by its recognition of GFP tagged Upd2 pull down in western blots] 1:2000 in TBS-T 0.05%; Rabbit- anti-GRASP (Cat# ab 30315, Abcam) 1:10000 in 1% BSA TBS-T 0.1%; Rabbit-anti RFP (Cat# 600-401-379, Rockland) 1:2500 in0.1% TBS-T . The secondary antibodies used were HRP conjugated Goat antibodies directed to appropriate species. The western blot was developed using Amersham ECL kit Start or Prime (Cat# RPN3243 or # RPN2232, GE life sciences).

Tissue preparation for fixed immunostaining:

For fat body preparation, incisions using dissection scissors (Cat# 500086, World Precision Instruments Inc) were made to release the ventral abdomen from the rest of the fly body. Flies used for dissection were adult males, 7-10 days old. Dissections were done in Ringer's medium (1.8 mM $CaCl_2$, 2mM Kcl, 128mM NaCl, 4mM $MgCl_2$. 35mM sucrose, 5mM HEPES). The tissue was fixed in 4% formaldehyde for 20 minutes and rinsed with PBS. For immunohistochemistry, the fixed fat tissue was permeabilized in PBS+ 1.0% Triton-X-100 for 3X washes 5 minutes, subsequently washed with PBS+0.3% Triton-X-100 (Fat wash). Blocked for 30 minutes at room temperature (RT) with gentle agitation in Fat wash+ 5% Normal donkey serum (Block). Primary antibodies, diluted in block, included Rabbit-anti-RFP (1:500, Rockland, #600-401-379); Chicken-anti-GFP (1:2000, Abcam, #ab13970); Mouse-anti-HA (1:200; Abcam, #ab18181); mouse-anti-Lamin (1:100; ADL67.10 DSHB); Rabbit-anti-GABARAP (Atg8) (1:500; Abcam, # ab109364); and incubated overnight at 4C. Washed 3X-5X for 15 mins each the following day in the fat wash at (RT) incubated with appropriate secondary antibody (from Jackson Immunoresearch) at a concentration of 1:500 in the block for 2-4 hours at RT. Washed 3X-5X for 5-15 minutes in the fat wash, mounted in SlowFade Gold antifade reagent with DAPI (Cat# S36938, Invitrogen).

Images were captured with Zeiss LSM 800 confocal system and analyzed with Zeiss ZenLite or ImageJ.

Cell staining:

For immunohistochemistry, cells were fixed for 20 minutes in 4% formaldehyde, washed in PBS for 5 quick changes, permeabilized in PBS+ 0.1% Triton-X-100 for 3X washes 5 minutes, subsequently washed with PBS+0.1% Triton-X-100 (Cell wash). Blocked for 30 minutes at room temperature (RT) with gentle agitation in Cell wash+ 5% Normal donkey serum (Block). Primary antibodies – Rabbit-anti-RFP (1:500, Rockland, #600-401-379); Chicken-anti-GFP (1:2000, Abcam, #ab13970); mouse-anti-Lamin (1:100; ADL67.10 DSHB) – were diluted in block incubated overnight at 4oC. Washed 3X-5X for 15 mins each was the following day in the fat wash at (RT) incubated with appropriate secondary antibody at a concentration of 1:500 (from Jackson Immunoresearch) in the block for 2-4 hours at RT. Washed 3X-5X for 5-15 minutes in cell wash, mounted with SlowFade Gold antifade reagent with DAPI (Cat# S36938, Invitrogen). Images were captured with Zeiss LSM 800 or Zeiss Elyra 7 with Lattice SIM confocal systems and analyzed with Zeiss ZenLite or ImageJ 14.

Hunger-driven feeding:

For hunger-driven feeding analysis, age-matched W1118 flies were given a normal diet or HSD for 5, 7, 14, 21, 24, and 28 days after an initial 7 days of development on a normal diet. All other experiments were performed for 7-day or 14-day durations. Sixteen hours before feeding behavior assessment, half of the flies from each treatment were moved to starvation media. Individual flies were placed in a single well of fly liquid-food interaction counter (FLIC) and supplied with a 1% sucrose liquid diet. Detailed methods for how FLIC operates can be found in Ro et al., 2014 (Ro et al., 2014). Fly feeding was measured for the first three hours in the FLIC, and all FLICs were performed at 10 am local time. For each FLIC, half of the wells (n=6/FLIC) contained the fed group, and the other half contained the starved group of flies for direct comparison. 12-30 flies were measured for analysis of feeding and normalized to the control fed group as a percentage. A signal above 40 (a.u.) was considered a feeding event. Analysis of feeding events was performed using R.

Triglyceride Measurements:

TAG assays were carried out as previously described in (Rajan et al., 2017). In brief: Flies were homogenized in PBST (PBS + 0.1% Triton X-100) using 1mm zirconium beads (Cat#ZROB10, Next Advance) in a Bullet Blender® Tissue homogenizer (Model BBX24, Next

Advance). Samples were heated to 70°C for 10 minutes, then centrifuged at 14,000 rpm (in a refrigerated tabletop centrifuge). 10.0 µl of the supernatant was applied to determine the level of TAG in the sample, using the following reagents obtained from Sigma: Free glycerol (cat # F6428-40ML), Triglyceride reagent (cat# T2449-10ML), and Glycerol standard (cat# G7793-5ML). Three adult males were employed per biological replicate. Note: For adult TAG assays, the most consistent results, with the lowest standard deviations, were obtained with 10-day old males. TAG readings from whole fly lysate (n=4 replicates of 3 flies each) were normalized to the number of flies per experiment. The normalized ratio from the control served as a baseline, and data are represented as fold change of experimental genotypes with respect to the control.

Survival Assays:

Survival curves were done using flies harvested in a 24-hour time frame and aged for 7 days. Ten males per vial were flipped onto 1% sucrose agar starvation food or a 30% High-sugar diet (HSD). The number of dead flies was recorded each day until every fly had died. Flies were kept in a 25°C incubator with a 12-hour light-dark cycle for the entirety of the experiment. Survival analysis was performed using the Survival Curve module of GraphPad Prism. A Mantel-cox test was the test used to determine statistical significance. Greater than 90 flies were used per genotype per curve. Data consolidated from 3 independent experiments is shown.

Morphometrics and image analysis:

Excel or GraphPad Prism 7 software was used for data quantification and generation of graphs. Quantification of nuclear localization in S2R+ was performed as follows:

Image analysis pipelines were built in MATLAB R2020b, and the associated scripts are available on request to the lead author. Three types of analysis were performed in 3D to quantify 1- the nucleus-to-whole cell intensity ratio of Upd2 in S2 cells, 2- the vesicle-to-whole cell intensity of Atg8 in S2 cells, and 3- the mean nuclear pixel intensity of Upd2 or Atg8 in fat cells.

To generate the 3D nuclear masks, Lamin or DAPI stacks were first maximally projected along the z-axis. After global thresholding, basic morphological operations, and watershed transform, the locations of the nuclear centroids in the x-y plane were used to

scan the z-stacks and reconstruct the nuclear volume. The whole-cell volume was approximated from the nuclear volume by dilation.

Atg8 vesicles were segmented within each cell volume by performing morphological filtering and image opening with defined structural elements. The accuracy of the segmentation was assessed by manual inspection of random cells.

Once these cellular compartments had been defined, they were used as masks to extract intensity values of the signal of interest and to compute either the mean pixel intensity in the nucleus, the ratio of the integral intensities between the nucleus and the whole cell, or between the combined vesicles and the whole cell.

Two-sided Wilcoxon rank-sum tests were used to assess the statistical significance of pairwise comparisons between experimental conditions.

We determined that our data-points were normally distributed, based on two measures: i) A GraphPad outlier test did not identify any outliers in our data; and ii) the majority of our data points for a particular condition were relatively similar to one other, with only a small standard error of mean or standard deviation.

References

3   Aguilar, G., Matsuda, S., Vigano, M.A., and Affolter, M. (2019). Using Nanobodies to Study Protein Function in Developing Organisms. Antibodies (Basel) *8*. 10.3390/antib8010016.

6   Ahima, R.S., Prabakaran, D., Mantzoros, C., Qu, D., Lowell, B., Maratos-Flier, E., and Flier, J.S. (1996). Role of leptin in the neuroendocrine response to fasting. Nature *382*, 250-252. 10.1038/382250a0.

9   Baker, K.D., and Thummel, C.S. (2007). Diabetic larvae and obese flies-emerging studies of metabolism in Drosophila. Cell Metab *6*, 257-266.

12 Bard, F., Casano, L., Mallabiabarrena, A., Wallace, E., Saito, K., Kitayama, H., Guizzunti, G., Hu, Y., Wendler, F., Dasgupta, R., et al. (2006). Functional genomics reveals genes involved in protein secretion and Golgi organization. Nature *439*, 604-607. 10.1038/nature04377.

Brankatschk, M., and Eaton, S. (2010). Lipoprotein particles cross the blood-brain barrier in Drosophila. J Neurosci *30*, 10441-10447. 10.1523/JNEUROSCI.5943-09.2010.

15 Brent, A.E., and Rajan, A. (2020). Insulin and Leptin/Upd2 Exert Opposing Influences on Synapse Number in Fat-Sensing Neurons. Cell Metab *32*, 786-800 e787. 10.1016/j.cmet.2020.08.017.

Cai, X.T., Li, H., Borch Jensen, M., Maksoud, E., Borneo, J., Liang, Y., Quake, S.R., Luo, L., Haghighi, P., and Jasper, H. (2021). Gut cytokines modulate olfaction through metabolic reprogramming of glia. Nature *596*, 97-102. 10.1038/s41586-021-03756-0.

Chakrabarti, S., Dudzic, J.P., Li, X., Collas, E.J., Boquete, J.P., and Lemaitre, B. (2016). Remote Control of Intestinal Stem Cell Activity by Haemocytes in Drosophila. PLoS Genet *12*, e1006089. 10.1371/journal.pgen.1006089.

Cunha, L.D., Yang, M., Carter, R., Guy, C., Harris, L., Crawford, J.C., Quarato, G., Boada-Romero, E., Kalkavan, H., Johnson, M.D.L., et al. (2018). LC3-Associated Phagocytosis in Myeloid Cells Promotes Tumor Immune Tolerance. Cell *175*, 429-441 e416. 10.1016/j.cell.2018.08.061.

Davis, L.I. (1995). The nuclear pore complex. Annu Rev Biochem *64*, 865-896. 10.1146/annurev.bi.64.070195.004245.

Droujinine, I.A., and Perrimon, N. (2016). Interorgan Communication Pathways in Physiology: Focus on Drosophila. Annu Rev Genet *50*, 539-570. 10.1146/annurev-genet-121415-122024.

Dupont, N., Jiang, S., Pilli, M., Ornatowski, W., Bhattacharya, D., and Deretic, V. (2011). Autophagy-based unconventional secretory pathway for extracellular delivery of IL-1beta. Embo J *30*, 4701-4711. 10.1038/emboj.2011.398.

Duran, J.M., Anjard, C., Stefan, C., Loomis, W.F., and Malhotra, V. (2010). Unconventional secretion of Acb1 is mediated by autophagosomes. J Cell Biol *188*, 527-536. 10.1083/jcb.200911154.

Efeyan, A., Comb, W.C., and Sabatini, D.M. (2015). Nutrient-sensing mechanisms and pathways. Nature *517*, 302-310. 10.1038/nature14190.

Ertekin, D., Kirszenblat, L., Faville, R., and van Swinderen, B. (2020). Down-regulation of a cytokine secreted from peripheral fat bodies improves visual attention while reducing sleep in Drosophila. PLoS biology *18*, e3000548. 10.1371/journal.pbio.3000548.

Farooqi, I.S., and O'Rahilly, S. (2009). Leptin: a pivotal regulator of human energy homeostasis. The American journal of clinical nutrition *89*, 980S-984S.

Fasken, M.B., Saunders, R., Rosenberg, M., and Brighty, D.W. (2000). A leptomycin B-sensitive homologue of human CRM1 promotes nuclear export of nuclear export sequence-containing proteins in Drosophila cells. J Biol Chem *275*, 1878-1886. 10.1074/jbc.275.3.1878.

Flier, J.S. (2019). Starvation in the Midst of Plenty: Reflections on the History and Biology of Insulin and Leptin. Endocr Rev *40*, 1-16. 10.1210/er.2018-00179.

Flier, J.S., and Maratos-Flier, E. (2017). Leptin's Physiologic Role: Does the Emperor of Energy Balance Have No Clothes? Cell Metab *26*, 24-26. 10.1016/j.cmet.2017.05.013.

Friedman, J.M., and Halaas, J.L. (1998). Leptin and the regulation of body weight in mammals. Nature *395*, 763-770. 10.1038/27376.

Galluzzi, L., and Green, D.R. (2019). Autophagy-Independent Functions of the Autophagy Machinery. Cell *177*, 1682-1699. 10.1016/j.cell.2019.05.026.

Geminard, C., Rulifson, E.J., and Leopold, P. (2009). Remote control of insulin secretion by fat cells in Drosophila. Cell Metab *10*, 199-207.

Gillette, C.M., Tennessen, J.M., and Reis, T. (2021). Balancing energy expenditure and storage with growth and biosynthesis during Drosophila development. Dev Biol *475*, 234-244. 10.1016/j.ydbio.2021.01.019.

He, C., and Klionsky, D.J. (2009). Regulation mechanisms and signaling pathways of autophagy. Annu Rev Genet *43*, 67-93. 10.1146/annurev-genet-102808-114910.

Heckmann, B.L., Teubner, B.J.W., Tummers, B., Boada-Romero, E., Harris, L., Yang, M., Guy, C.S., Zakharenko, S.S., and Green, D.R. (2019). LC3-Associated Endocytosis Facilitates beta-Amyloid Clearance and Mitigates Neurodegeneration in Murine Alzheimer's Disease. Cell *178*, 536-551 e514. 10.1016/j.cell.2019.05.056.

Hickey, M.S., Considine, R.V., Israel, R.G., Mahar, T.L., McCammon, M.R., Tyndall, G.L., Houmard, J.A., and Caro, J.F. (1996). Leptin is related to body fat content in male distance runners. Am J Physiol *271*, E938-940.

Hombria, J.C., Brown, S., Hader, S., and Zeidler, M.P. (2005). Characterisation of Upd2, a Drosophila JAK/STAT pathway ligand. Dev Biol *288*, 420-433.

Huang, R., Xu, Y., Wan, W., Shou, X., Qian, J., You, Z., Liu, B., Chang, C., Zhou, T., Lippincott-Schwartz, J., and Liu, W. (2015). Deacetylation of nuclear LC3 drives autophagy initiation under starvation. Mol Cell *57*, 456-466. 10.1016/j.molcel.2014.12.013.

Ichimura, Y., Kirisako, T., Takao, T., Satomi, Y., Shimonishi, Y., Ishihara, N., Mizushima, N., Tanida, I., Kominami, E., Ohsumi, M., et al. (2000). A ubiquitin-like system mediates protein lipidation. Nature *408*, 488-492. 10.1038/35044114.

Jura, N., Scotto-Lavino, E., Sobczyk, A., and Bar-Sagi, D. (2006). Differential modification of Ras proteins by ubiquitination. Molecular Cell *21*, 679-687. 10.1016/j.molcel.2006.02.011.

     Kaffman, A., and O'Shea, E.K. (1999). Regulation of nuclear localization: a key to a door. Annu Rev Cell Dev Biol *15*, 291-339. 10.1146/annurev.cellbio.15.1.291.

27   Kaushik, S., Singh, R., and Cuervo, A.M. (2010). Autophagic pathways and metabolic stress. Diabetes Obes Metab *12 Suppl 2*, 4-14. 10.1111/j.1463-1326.2010.01263.x.

30   Kim, S.K., Tsao, D.D., Suh, G.S.B., and Miguel-Aliaga, I. (2021). Discovering signaling mechanisms governing metabolism and metabolic diseases with Drosophila. Cell Metab *33*, 1279-1292. 10.1016/j.cmet.2021.05.018.

33   King-Jones, K., and Thummel, C.S. (2005). Nuclear receptors--a perspective from Drosophila. Nat Rev Genet *6*, 311-323.

36   Kirisako, T., Baba, M., Ishihara, N., Miyazawa, K., Ohsumi, M., Yoshimori, T., Noda, T., and Ohsumi, Y. (1999). Formation process of autophagosome is traced with Apg8/Aut7p in yeast. J Cell Biol *147*, 435-446. 10.1083/jcb.147.2.435.

Kirisako, T., Ichimura, Y., Okada, H., Kabeya, Y., Mizushima, N., Yoshimori, T., Ohsumi, M., Takao, T., Noda, T., and Ohsumi, Y. (2000). The reversible modification regulates the membrane-binding state of Apg8/Aut7 essential for autophagy and the cytoplasm to vacuole targeting pathway. J Cell Biol *151*, 263-276.

Klionsky, D.J., Abdalla, F.C., Abeliovich, H., Abraham, R.T., Acevedo-Arozena, A., Adeli, K., Agholme, L., Agnello, M., Agostinis, P., Aguirre-Ghiso, J.A., et al. (2012). Guidelines for the use and interpretation of assays for monitoring autophagy. Autophagy *8*, 445-544.

Knight, Z.A., Hannan, K.S., Greenberg, M.L., and Friedman, J.M. (2010). Hyperleptinemia is required for the development of leptin resistance. PloS one *5*, e11376. 10.1371/journal.pone.0011376.

Kolaczynski, J.W., Considine, R.V., Ohannesian, J., Marco, C., Opentanova, I., Nyce, M.R., Myint, M., and Caro, J.F. (1996a). Responses of leptin to short-term fasting and refeeding in humans: a link with ketogenesis but not ketones themselves. Diabetes *45*, 1511-1515.

Kolaczynski, J.W., Ohannesian, J.P., Considine, R.V., Marco, C.C., and Caro, J.F. (1996b). Response of leptin to short-term and prolonged overfeeding in humans. J Clin Endocrinol Metab *81*, 4162-4165. 10.1210/jcem.81.11.8923877.

Kondo, S., and Ueda, R. (2013). Highly improved gene targeting by germline-specific Cas9 expression in Drosophila. Genetics *195*, 715-721. 10.1534/genetics.113.156737.

Kosugi, S., Hasebe, M., Entani, T., Takayama, S., Tomita, M., and Yanagawa, H. (2008). Design of peptide inhibitors for the importin alpha/beta nuclear import pathway by activity-based profiling. Chem Biol *15*, 940-949. 10.1016/j.chembiol.2008.07.019.

Kosugi, S., Hasebe, M., Matsumura, N., Takashima, H., Miyamoto-Sato, E., Tomita, M., and Yanagawa, H. (2009a). Six classes of nuclear localization signals specific to different binding grooves of importin alpha. J Biol Chem *284*, 478-485. 10.1074/jbc.M807017200.

Kosugi, S., Hasebe, M., Tomita, M., and Yanagawa, H. (2009b). Systematic identification of cell cycle-dependent yeast nucleocytoplasmic shuttling proteins by prediction of composite motifs. Proc Natl Acad Sci U S A *106*, 10171-10176. 10.1073/pnas.0900604106.

Kraft, L.J., Manral, P., Dowler, J., and Kenworthy, A.K. (2016). Nuclear LC3 Associates with Slowly Diffusing Complexes that Survey the Nucleolus. Traffic *17*, 369-399. 10.1111/tra.12372.

la Cour, T., Kiemer, L., Molgaard, A., Gupta, R., Skriver, K., and Brunak, S. (2004). Analysis and prediction of leucine-rich nuclear export signals. Protein Eng Des Sel *17*, 527-536. 10.1093/protein/gzh062.

Lee, P.T., Zirin, J., Kanca, O., Lin, W.W., Schulze, K.L., Li-Kroeger, D., Tao, R., Devereaux, C., Hu, Y., Chung, V., et al. (2018). A gene-specific T2A-GAL4 library for Drosophila. Elife *7*. 10.7554/eLife.35574.

Leidal, A.M., Huang, H.H., Marsh, T., Solvik, T., Zhang, D., Ye, J., Kai, F., Goldsmith, J., Liu, J.Y., Huang, Y.H., et al. (2020). The LC3-conjugation machinery specifies the loading of RNA-binding proteins into extracellular vesicles. Nat Cell Biol *22*, 187-199. 10.1038/s41556-019-0450-y.

Leopold, P., and Perrimon, N. (2007). Drosophila and the genetics of the internal milieu. Nature *450*, 186-188.

3   Mahajan, R., Delphin, C., Guan, T., Gerace, L., and Melchior, F. (1997). A small ubiquitin-related polypeptide involved in targeting RanGAP1 to nuclear pore complex protein RanBP2. Cell *88*, 97-107. 10.1016/s0092-8674(00)81862-0.

6   Musselman, L.P., and Kuhnlein, R.P. (2018). Drosophila as a model to study obesity and metabolic disease. J Exp Biol *221*. 10.1242/jeb.163881.

Nakagawa, T., and Nakayama, K. (2015). Protein monoubiquitylation: targets and diverse functions.
9   Genes Cells *20*, 543-562. 10.1111/gtc.12250.

Nakatogawa, H., Ichimura, Y., and Ohsumi, Y. (2007). Atg8, a ubiquitin-like protein required for autophagosome formation, mediates membrane tethering and hemifusion. Cell *130*, 165-178.
12   10.1016/j.cell.2007.05.021.

Neufeld, T.P., and Baehrecke, E.H. (2008). Eating on the fly: function and regulation of autophagy during cell growth, survival and death in Drosophila. Autophagy *4*, 557-562. 10.4161/auto.5782.

15   Noda, N.N., Ohsumi, Y., and Inagaki, F. (2010). Atg8-family interacting motif crucial for selective autophagy. FEBS letters *584*, 1379-1385. 10.1016/j.febslet.2010.01.018.

Rajan, A., Housden, B.E., Wirtz-Peitz, F., Holderbaum, L., and Perrimon, N. (2017). A Mechanism
18   Coupling Systemic Energy Sensing to Adipokine Secretion. Dev Cell *43*, 83-98 e86. 10.1016/j.devcel.2017.09.007.

Rajan, A., and Perrimon, N. (2012). *Drosophila* Cytokine Unpaired 2 Regulates Physiological
21   Homeostasis by Remotely Controlling Insulin Secretion. Cell *151*, 123-137. 10.1016/j.cell.2012.08.019.

Rajan, A., and Perrimon, N. (2013). Of flies and men: insights on organismal metabolism from fruit
24   flies. BMC Biol *11*, 38. 10.1186/1741-7007-11-38.

Reggiori, F., Monastyrska, I., Verheije, M.H., Cali, T., Ulasli, M., Bianchi, S., Bernasconi, R., de Haan, C.A., and Molinari, M. (2010). Coronaviruses Hijack the LC3-I-positive EDEMosomes, ER-derived
27   vesicles exporting short-lived ERAD regulators, for replication. Cell Host Microbe *7*, 500-508. 10.1016/j.chom.2010.05.013.

Ressurreicao, M., Warrington, S., and Strutt, D. (2018). Rapid Disruption of Dishevelled Activity
30   Uncovers an Intercellular Role in Maintenance of Prickle in Core Planar Polarity Protein Complexes. Cell Rep *25*, 1415-1424 e1416. 10.1016/j.celrep.2018.10.039.

Ro, J., Harvanek, Z.M., and Pletcher, S.D. (2014). FLIC: high-throughput, continuous analysis of
33   feeding behaviors in Drosophila. PLoS One *9*, e101107. 10.1371/journal.pone.0101107.

Rothbauer, U., Zolghadr, K., Muyldermans, S., Schepers, A., Cardoso, M.C., and Leonhardt, H. (2008). A versatile nanotrap for biochemical and functional studies with fluorescent fusion proteins.
36   Mol Cell Proteomics *7*, 282-289. 10.1074/mcp.M700342-MCP200.

Saerens, D., Pellis, M., Loris, R., Pardon, E., Dumoulin, M., Matagne, A., Wyns, L., Muyldermans, S., and Conrath, K. (2005). Identification of a universal VHH framework to graft non-canonical antigen-binding loops of camel single-domain antibodies. Journal of molecular biology *352*, 597-607. 10.1016/j.jmb.2005.07.038.

Schneider, I. (1972). Cell lines derived from late embryonic stages of Drosophila melanogaster. J Embryol Exp Morphol *27*, 353-365.

Shpilka, T., Weidberg, H., Pietrokovski, S., and Elazar, Z. (2011). Atg8: an autophagy-related ubiquitin-like protein family. Genome Biol *12*, 226. 10.1186/gb-2011-12-7-226.

Singh, R., and Cuervo, A.M. (2011). Autophagy in the cellular energetic balance. Cell Metab *13*, 495-504. 10.1016/j.cmet.2011.04.004.

Singh, R., Kaushik, S., Wang, Y., Xiang, Y., Novak, I., Komatsu, M., Tanaka, K., Cuervo, A.M., and Czaja, M.J. (2009a). Autophagy regulates lipid metabolism. Nature *458*, 1131-1135. 10.1038/nature07976.

Singh, R., Xiang, Y., Wang, Y., Baikati, K., Cuervo, A.M., Luu, Y.K., Tang, Y., Pessin, J.E., Schwartz, G.J., and Czaja, M.J. (2009b). Autophagy regulates adipose mass and differentiation in mice. J Clin Invest *119*, 3329-3339. 10.1172/JCI39228.

Speese, S.D., Ashley, J., Jokhi, V., Nunnari, J., Barria, R., Li, Y., Ataman, B., Koon, A., Chang, Y.T., Li, Q., et al. (2012). Nuclear envelope budding enables large ribonucleoprotein particle export during synaptic Wnt signaling. Cell *149*, 832-846. 10.1016/j.cell.2012.03.032.

Stolz, A., Putyrski, M., Kutle, I., Huber, J., Wang, C., Major, V., Sidhu, S.S., Youle, R.J., Rogov, V.V., Dotsch, V., et al. (2017). Fluorescence-based ATG8 sensors monitor localization and function of LC3/GABARAP proteins. EMBO J *36*, 549-564. 10.15252/embj.201695063.

Takats, S., Toth, S., Szenci, G., and Juhasz, G. (2019). Investigating Non-selective Autophagy in Drosophila. Methods Mol Biol *1880*, 589-600. 10.1007/978-1-4939-8873-0_38.

Teleman, A.A., Ratzenbock, I., and Oldham, S. (2012). Drosophila: a model for understanding obesity and diabetic complications. Exp Clin Endocrinol Diabetes *120*, 184-185. 10.1055/s-0032-1304566.

Tennessen, J.M., Barry, W.E., Cox, J., and Thummel, C.S. (2014). Methods for studying metabolism in Drosophila. Methods *68*, 105-115. 10.1016/j.ymeth.2014.02.034.

Thoreen, C.C., Kang, S.A., Chang, J.W., Liu, Q., Zhang, J., Gao, Y., Reichling, L.J., Sim, T., Sabatini, D.M., and Gray, N.S. (2009). An ATP-competitive mammalian target of rapamycin inhibitor reveals rapamycin-resistant functions of mTORC1. J Biol Chem *284*, 8023-8032. 10.1074/jbc.M900301200.

van Dam, E., van Leeuwen, L.A.G., Dos Santos, E., James, J., Best, L., Lennicke, C., Vincent, A.J., Marinos, G., Foley, A., Buricova, M., et al. (2020). Sugar-Induced Obesity and Insulin Resistance Are Uncoupled from Shortened Survival in Drosophila. Cell Metab *31*, 710-725 e717. 10.1016/j.cmet.2020.02.016.

Verboon, J.M., Rincon-Arano, H., Werwie, T.R., Delrow, J.J., Scalzo, D., Nandakumar, V., Groudine, M., and Parkhurst, S.M. (2015). Wash interacts with lamin and affects global nuclear organization. Curr Biol *25*, 804-810. 10.1016/j.cub.2015.01.052.

Wendler, F., Gillingham, A.K., Sinka, R., Rosa-Ferreira, C., Gordon, D.E., Franch-Marro, X., Peden, A.A., Vincent, J.P., and Munro, S. (2010). A genome-wide RNA interference screen identifies two novel components of the metazoan secretory pathway. EMBO J *29*, 304-314. 10.1038/emboj.2009.350.

Wright, V.M., Vogt, K.L., Smythe, E., and Zeidler, M.P. (2011). Differential activities of the Drosophila JAK/STAT pathway ligands Upd, Upd2 and Upd3. Cell Signal *23*, 920-927, 2011. S0898-6568(11)00021-0 [pii]

10.1016/j.cellsig.2011.01.020.

Wyant, G.A., Abu-Remaileh, M., Frenkel, E.M., Laqtom, N.N., Dharamdasani, V., Lewis, C.A., Chan, S.H., Heinze, I., Ori, A., and Sabatini, D.M. (2018). NUFIP1 is a ribosome receptor for starvation-induced ribophagy. Science *360*, 751-758. 10.1126/science.aar2663.

Xu, L., Lubkov, V., Taylor, L.J., and Bar-Sagi, D. (2010). Feedback regulation of Ras signaling by Rabex-5-mediated ubiquitination. Curr Biol *20*, 1372-1377. 10.1016/j.cub.2010.06.051.

Zacharogianni, M., Aguilera-Gomez, A., Veenendaal, T., Smout, J., and Rabouille, C. (2014). A stress assembly that confers cell viability by preserving ERES components during amino-acid starvation. Elife *3*. 10.7554/eLife.04132.

Zacharogianni, M., Kondylis, V., Tang, Y., Farhan, H., Xanthakis, D., Fuchs, F., Boutros, M., and Rabouille, C. (2011). ERK7 is a negative regulator of protein secretion in response to amino-acid starvation by modulating Sec16 membrane association. EMBO J *30*, 3684-3700. 10.1038/emboj.2011.253.

Zhang, M., Kenny, S., Ge, L., Xu, K., and Schekman, R. (2015). Translocation of interleukin-1beta into a vesicle intermediate in autophagy-mediated secretion. Elife *4*. 10.7554/eLife.11205.

Zirin, J., and Perrimon, N. (2010). Drosophila as a model system to study autophagy. Semin Immunopathol *32*, 363-372. 10.1007/s00281-010-0223-y.

Figure 1: Acute starvation triggers Upd2 nuclear accumulation

Figure 1. Acute Starvation triggers Upd2 nuclear accumulation.

(A) <u>Effect of starvation on Upd2 nuclear accumulation in adult fly fat tissue:</u> Confocal micrographs of fixed and immunostained adult fly fat from flies CRISPR-engineered (see methods) with GFP tag in the endogenous Upd2 locus (that rescues the upd2Δ mutant- see Figure S1). Fat is stained with GFP antibody (in a, b) in well-fed flies (a), or flies starved for 4 hours. Upd2 shows increased nuclear accumulation on starvation [see look up table (LUT)]. Scale bar is 1μM. (A') Upd2-GFP endogenous nuclear accumulation is quantified using 3D-volumetric segmentation-based calculations (See Methods). Each dot represents a fat cell nucleus, 75-100 fat nuclei were counted per condition per genotype. Statistical significance is quantified by the two-sided Wilcoxon test. Also see Figure S1 for similar experiment performed on an HA-tag knock into the Upd2 genomic locus (Upd2-HA).

(B) <u>Effect of starvation and re-feeding on Upd2's nuclear accumulation in *Drosophila* S2R+ cells:</u> Confocal micrographs of single optical-slices of *Drosophila* S2R+ cells transiently transfected with Upd2-WT::GFP (green; anti-GFP) and co-stained with Lamin (red). Scale bar is 5um. In B', the ratio of Upd2::GFP nuclear/whole cell intensity is plotted. Each dot represents a cell, 50-100 cells were counted per time point. Statistical significance is quantified by the two-sided Wilcoxon test.

(C) <u>Effect of starvation and re-feeding on Upd2's release into media:</u> Normalized percent fold change in secreted GFP signal detected by GFP sandwich ELISA assay performed on conditioned media of S2R+ cells transiently transfected with Upd2-WT::GFP. Cells were incubated either with complete media (control) or Amino-Acid (AA) free media for the indicated times. The refeeding % fold change is indicated with respect to starvation. Statistical significance is quantified by unpaired two-tailed t-test on 6 biological replicates per condition.

Figure S1: companion to Main Figure 1

A *Upd2::GFP and Upd2::HA CRISPR mediated genomic locus tag knock-in*

Rescue of Upd2-deletion TAG phenotype by Upd2 tag knock-in flies

Aa)

Legend:
- ■ Upd2Δ
- ▨ Upd2-crGFP
- ▢ Upd2-crGFP/Upd2Δ

Normalized TAG level (fold change)

p = 0.025
p = 0.01

Ab)

Legend:
- ■ Upd2Δ
- ▨ Upd2-crHA
- ▢ Upd2-crHA/Upd2Δ

Normalized TAG level (fold change)

p = 0.003
p = 0.005

B *Upd2-HA genomic locus tag knock in*

Upd2::HA nuclear accumulation levels in relation to nutrient state

p=0.03848

Mean Upd2 nuclear mean pixel intensity

FED   STV

C *Transient transfection of Upd2::GFP and dsRNA in S2R+ cells*

Upd2-WT::GFP localization when Embargoed/CRM1 nuclear export pathway is disrupted

anti-Lamin anti-GFP

Control dsRNA — **a** — 5 µm — *LacZ-KD*

Emb dsRNA — **b** — *Emb-KD*

C'

Upd2::GFP Nuclear/Cell intensity ratio

p=0.01
p=0.0009

Control dsRNA   *Embargoed dsRNA #1*   *Embargoed dsRNA #2*

D *Upd2::GFP transfected in S2R+ cells*

Upd2 secretion in S2R+ cells during Emb/CRM1 knock-down

% change in secretion of Upd2::GFP

Legend:
- ▢ *Control dsRNA*
- ▨ *Emb dsRNA #1*
- ■ *Emb dsRNA #2*

0%   **-33%** **-49%**

p=0.008
p=0.001

Figure S1: Companion to Main Figure 1.

(A) Rescue of upd2Δ triglyceride (TAG) storage reduction phenotype by endogenous CRISPR tag knock-in of HA [Upd2-HA] (S1B) and GFP [Upd2-GFP] (1A) in the Upd2 locus.

(B) Upd2-HA endogenous nuclear accumulation is quantified using 3D-volumetric segmentation-based calculations (See Methods). Compare with similar results observed for a GFP-tagged Upd2 in Main Figure 1A'.

(C) Confocal micrographs of single optical-slices of *Drosophila* S2R+ cells transiently co-transfected with Upd2-WT::GFP (green) stained with Lamin (red). In fed state, cells are co-transfected with (a) LacZ-dsRNA (control), or (b) CRM1/Embragoed (Emb)-KD (Emb dsRNA). Emb-KD increases Upd2-WT::GFP nuclear accumulation and shows an enrichment of Upd2-WT::GFP at the interface of nuclear lamina and cytoplasm (see blue arrows). Scale bar is 5μm. Two independent *Emb-dsRNAs* (#1 and #2) were used.

For B, C' statistical significance is quantified by the two-sided Wilcoxon test. Each dot represents a fat cell nucleus (B) or cell (C'). 50-100 replicates were quantified per genotype per condition.

(D) Normalized percent fold change in secreted GFP signal detected by GFP sandwich ELISA assay performed on conditioned media of S2R+ cells transiently transfected with Upd2-WT::GFP dsRNA for control (LacZ) or two independent *Emb dsRNAs*. ELISA for Upd2-WT::GFP was performed after 5 days of knockdown, and the amount of Upd2-WT::GFP in a 24-hour period (Day 5) was assessed. Statistical significance is quantified by unpaired two-tailed t-test on 6 biological replicates per condition.

Figure 2: *Upd2 requires Atg8 for nuclear exit in fed state*

A Upd2's AIM sequence: yfqylnnv**W**kt**L**hrvlrrprr

upd2Δ; Lpp-Gal4>UAS-Upd2-X::mCherry

Upd2-WT::mCherry vs. Upd2-AIM-::mCherry fat-specific overexpression

anti-Lamin anti-mCherry (LUT)

Upd2-WT

a *Upd2-WT* Upd2 (LUT) Upd2 (LUT)

Merge Fed 5um High / Low

Upd2-AIM-

b *Upd2-AIM-*

Merge Fed

B *Upd2::GFP knock in; ppl-GAL4tubGal80ᵗˢ> UAS- RNAi*

Nuclear levels of endogenous Upd2::GFP protein in fat-specific temporal adult fly knock-down of Atg8

p=0.008

Mean Upd2 Nuclear pixel intensity

3.0 ×10²

2.5

2.0

1.5

1.0

0.5

Luciferase-RNAi   *Atg8-RNAi*

C *Transient transfection of Upd2-X::GFP in Drosophila S2R+ cells*

Upd2-WT::GFP vs. Upd2-AIM-::GFP in well-fed S2R+ cells

anti-Lamin   anti-GFP

Upd2-WT

a anti-GFP anti-Lamin anti-GFP *Upd2-WT::GFP* Merge 5 µm

Upd2-AIM-

b anti-GFP anti-Lamin anti-GFP *Upd2-AIM-::GFP* Merge

C'

p=1.11e-06

Upd2::GFP Nuclear/Cell intensity ratio

1 0.9 0.8 0.7 0.6 0.5 0.4 0.3 0.2 0.1 0

Upd2-WT   *Upd2-AIM-*

D *Transient transfection of Upd2-AIM-::GFP with V_HH-Atg8-WT in Drosophila S2R+ cells*

Upd2-AIM- localization on nanobody-directed interaction with Atg8-WT in S2R+ cells

Da) Upd2-AIM- GFP + V_HH (GFP-nanobody)

Db) Upd2-AIM- GFP + V_HH (GFP-nanobody) Atg8-WT

Da Merge *Upd2-AIM-::GFP V_HH-Myc*

Upd2-AIM-::GFP *V_HH-Myc*

Db Merge *Upd2-AIM-::GFP V_HH-Myc::Atg8-WT* 2 µm

Upd2-AIM-::GFP *V_HH-Myc::Atg8-WT* 2 µm

anti-Lamin   **anti-GFP**   **anti-Myc**

Figure 2. Atg8 controls Upd2 nucleocytoplasmic localization.

(A) <u>Effect of point mutations to Upd2's AIM (Upd2-AIM-) on its nuclear accumulation in adult fly fat:</u> Upd2-AIM- includes two point mutations to Upd2's putative AIM (WXXL→ AXXA) – see sequence above the figure. Confocal micrographs of single optical sections of fixed and immunostained adult fly fat from transgenic flies expressing mCherry tagged Upd2-cDNAs for (a) Upd2-WT (UAS-Upd2-WT) or (b) Upd2's AIM mutated (UAS-Upd2-AIM-) under the control of a fat-cell specific promoter (Lpp-Gal4) in a upd2-deletion background (upd2Δ; Lpp-Gal4).

(B) <u>Effect of Atg8-KD on Upd2 nuclear accumulation in adult fly fat tissue:</u> Endogenous Upd2 nuclear signal in fly fat (upd2::GFP-endogenous knock-in) of flies expressing control (Luciferase-grey bar) or Atg8-RNAi (red bar) in fat cells during adult stages (ppl-Gal4tubGal80ts).

(C) <u>Effect of point mutations to Upd2's AIM (Upd2-AIM-::GFP) on its nuclear accumulation in *Drosophila* S2R+ cells:</u> Confocal micrographs (a, b) of single optical-slices of *Drosophila* S2R+ cells transiently co-transfected with (a) Upd2-WT::GFP or (b) Upd2-AIM-::GFP (green- anti-GFP) stained with Lamin (red) in a fed state. Scale bars are 5um for a, b. In C', the ratio of Upd2::GFP nuclear/whole cell intensity is plotted. See methods of details on quantification.

For (B) each dot on the graph is a fat cell nucleus, in (C') each dot is a cell. Statistical significance for B and C' are quantified by the two-sided Wilcoxon Rank sum tests.

(D) <u>Effect of nanobody-directed interaction between Atg8-WT and Upd2-AIM-::GFP on Upd2's localization in *Drosophila* S2R+ cells:</u> Top panel shows schematic of the experimental design. V_HH tag, a small 15kDa antibody to GFP, is expressed by itself (a) or fused to the N-terminal of Atg8-WT and co-transfected with Upd2-AIM-::GFP. Bottom panel shows a single XY-slice of confocal image in *Drosophila* S2R+ cells cultured in complete media. Cells were fixed and immunostained with antibodies to detect V_HH-Myc / V_HH-Myc::Atg8-WT (anti-Myc-red) and Upd2-AIM-::GFP (anti-GFP). In a, note that Upd2-AIM-::GFP displays nuclear accumulation and so does V_HH-Myc, consistent with its role as an antibody to GFP (see arrows). In b, punctate overlap between V_HH Atg8 and GFP are observed- see arrow and inset. Note Upd2-AIM- is present in the cytosolic puncta, instead of nuclear accumulation (compare with Figure 2Cb). Also see Figure 3B' for quantification of the nuclear/cytosolic localization of Upd2-AIM- in the presence of V_HH-Myc::Atg8-WT compared to control (V_HH-Myc empty).

Figure S2: *companion to Main Figure 2*

A *Transient transfection of Upd2-X::GFP and Myc::Atg8-WT in S2R+ cells*

Upd2-WT::GFP vs. Upd2-AIM-::GFP's ability to co-immunoprecipitate with Atg8

GFP-IP — Anti-GFP, Anti-Upd2
Myc-IP — Anti-Myc, Anti-GFP, Anti-Upd2
INPUT — Anti-Upd2, Anti-Myc

B *Transient transfection of Upd2::GFP and dsRNA in S2R+ cells*

Upd2-WT::GFP localization in Atg8 knock-down cells

anti-Lamin anti-GFP

Control dsRNA — *LacZ-KD*
Atg8 dsRNA — *Atg8-KD*
5 μm

B'

Upd2::GFP Nuclear/Cell intensity ratio

p=0.017
p=0.002

Control dsRNA   Atg8 dsRNA #1   Atg8 dsRNA #2

C *Upd2::GFP transfected in S2R+ cells*

Upd2 secretion in S2R+ cells during Atg8 knock-down

% change in secretion of Upd2::GFP

☐ Control dsRNA
■ Atg8 dsRNA

-85%

p=0.048

D *Upd2::GFP transfected in S2R+ cells*

Upd2-WT vs. Upd2-AIM- secretion in S2R+ cells

Upd2 AIM sequence:yfqylnnvWktLhrvlrrprr

% change in secretion of Upd2::GFP

☐ Upd2-WT::GFP
☐ Upd2-AIM-::GFP

-96%

p=4.29E-05

E *Transient transfection of Upd2-X::GFP and GRASP::RFP in S2R+ cells*

Upd2-WT::GFP vs. Upd2-AIM-::GFP's ability to co-immunoprecipitate with GRASP

Upd2-WT::GFP   Upd2-AIM-::GFP
1   2

GFP-IP — Anti-Upd2
RFP-IP — Anti-GRASP, Anti-Upd2
INPUT — Anti-Tubulin

GRASP::RFP

Figure S2: Companion to Main Figure 2.

(A) GFP- and Myc-IPs were prepared from S2R+ cells transiently transfected with the indicated cDNAs. GFP, Myc IPs and 2% of the input were analyzed by immunoblotting for the indicated proteins. Note Upd2-AIM- always runs slightly higher than Upd2 (see asterisk) and is always more abundant in the lysate as the Upd2-AIM- is not secreted (see D below), but Upd2-AIM- it is not readily detectable in the Myc-IPs (see red asterisk), despite being more abundant in input. Atg8-Myc is always more abundant in Upd2-AIM- input than in Upd2 -WT input, despite loading same amount of protein in input.

(B) Confocal micrographs of single optical-slices of *Drosophila* S2R+ cells transiently co-transfected with Upd2-WT::GFP (green) stained with Lamin (red). (a) In fed state, in cells co-transfected with LacZ-dsRNA (control) Upd2-WT::GFP is localized to cytosolic puncta and nuclear localization is not obvious. (b) In fed state, Atg8-KD (Atg8 dsRNA) significant increase in Upd2-WT::GFP nuclear accumulation is observed. Scale bar is 5um and right most panel shows DIC image merge. In B', the ratio of Upd2::GFP nuclear/whole cell intensity is plotted. Each dot represents a cell, 20-40 cells were counted per dsRNA condition. Two independent Atg8-dsRNAs were used. Statistical significance is quantified by two-sided Wilcoxon sum rank tests.

(C) Normalized percent fold change in secreted GFP signal detected by GFP sandwich ELISA assay performed on conditioned media of *Drosophila* S2R+ cells transiently transfected with Upd2-WT::GFP dsRNA for control (LacZ) or Atg8 dsRNA. ELISA for Upd2-WT::GFP was performed after 5 days of knockdown, and the amount of Upd2-WT::GFP in a 24-hour period (Day 5) was assessed. Statistical significance is quantified by unpaired two-tailed t-test on 6 biological replicates per condition.

(D) Normalized percent fold change in secreted GFP signal detected by GFP sandwich ELISA assay performed on conditioned media of S2R+ cells transiently transfected with Upd2-WT::GFP or Upd2-AIM-::GFP. Mutated residues in Upd2's AIM is shown in red. Statistical significance is quantified by unpaired two-tailed t-test on 6 biological replicates per condition.

(E) GFP- and RFP-IPs were prepared from S2R+ cells transiently co-transfected with GRASP-RFP and Upd2-WT::GFP or Upd2-AIM-::GFP. GFP, RFP IPs and 2% of the input were analyzed by immunoblotting for the indicated proteins. Note Upd2-AIM- always runs slightly higher than Upd2 and is always more abundant in the lysate as the Upd2-AIM- is not secreted, but it is not readily detectable in the RFP-IPs, despite loading more input.

Figure 3: Atg8's lipidation status regulates starvation-induced Upd2 nuclear retention

A

Transient transfection of Upd2-WT-::GFP with V_HH-Atg8-WT/PE- in Drosophila S2R+ cells

Upd2-WT::GFP's localization on starvation in the nanobody-directed interaction with Atg8-WT or Atg8-PE-

B

Transient transfection of Upd2-AIM-::GFP with V_HH-Atg8-WT/PE- in Drosophila S2R+ cells

Upd2-AIM-::GFP localization on nanobody-directed interaction with Atg8-WT or Atg8-PE-

Figure 3. Atg8 lipidation status regulates starvation-induced nuclear retention.

In (A) and (B), schematics illustrate experimental design. V_HH tag, a small 15kDa antibody to GFP, is the control and V_HH tag fused to the N-terminal of Atg8-WT or Atg8-G$\rightarrow$A mutation (Atg8-PE-) Also see companion Figure S3 and S4.

(A) <u>Effect of nanobody-directed interaction between Upd2-WT::GFP and Atg8-WT or Atg8-PE- in *Drosophila* S2R+ cells during an acute amino-acid starvation:</u> Side panels show schematic of the experimental design. Upd2's starvation induced nuclear retention is assessed. Aa-Af show a single XY-slice of confocal image in *Drosophila* S2R+ cells cultured in complete media (a, c, e) or amino-acid free media (b, d, f). Cells were fixed and immunostained with anti-Lamin (blue) and anti-GFP (green) to detect Upd2-WT::GFP. Scale bar represents 1μM.

(B) <u>Effect of nanobody-directed interaction between Upd2-AIM-::GFP and Atg8-WT or lipidation defective Atg8-PE- in *Drosophila* S2R+ cells:</u> Top panels show schematic of the experimental design. Ba-Bd show a single XY-slice of confocal image in *Drosophila* S2R+ cells cultured in complete media. Cells were fixed and immunostained with anti-Lamin (blue) and anti-GFP (green) to detect Upd2-WT::GFP (a) or Upd2-AIM-::GFP (b-c). The effect of co-transfection with of Upd2-WT with V_HH tag (a) is used as baseline and compared to Upd2-AIM- localization in the presence of 'empty' V_HH tag (b) or V_HH tag fused to Atg8-WT (c) or Atg8-PE- (d) is assessed. Scale bar represent 5μM.

A' and B' graphs show 3D-volumetric quantification of nuclear to whole cell intensity of GFP signal. Statistical significance is calculated using two-sided Wilcoxon sum rank tests. Each circle represents a cell.

Figure S3: *companion to Main Figure 3*

Upd2-WT::GFP localization in S2R+ cells treated with autophagy activator Torin

A

anti-Lamin anti-GFP

DMSO

Torin

50nM

250nM

750nM

5 µm

DMSO

Torin 50nm

Torin 250nm

Torin 750nm

A'

Upd2::GFP Nuclear/Cell intensity ratio

p=1.94e-19

p=2.94e-16

p=1.64e-10

DMSO   50nM Torin   250nM Torin   750nM Torin

Supplemental Figure 3: Companion to Main Figure 3.

(A) Confocal micrographs of single optical-slices of *Drosophila* S2R+ cells transiently transfected with Upd2-WT::GFP (green) stained with Lamin (red) and treated with DMSO (a) or indicated concentrations of autophagy activator Torin1 (b, c, d). Scale bar is 5μm and right most panel shows DIC image merge. In A', the ratio of Upd2::GFP nuclear/whole cell intensity is plotted. Each dot represents a cell, 50-100 cells were counted per condition. Statistical significance is quantified by unpaired two-tailed t-test on 6 biological replicates per condition.

Figure S4: companion to Main Figure 3

A *Transient transfection of mCherry::Atg8-WT/ G->A in Drosophila S2R+ cells*

Figure S4: Companion to Main Figure 3.

(A) Western blot on lysates prepared from cells transfected with mCherry::Atg8-WT or mCherry::Atg8-G→ A. Cells were either maintained in complete media (Fed-CM) or AA-acid starved for 2 hours (2hr) and 4 hours (4hr). The cell lysates were probed with anti-RFP antibody. Atg8 (15kDa Atg8+ 27kDa mCherry) is recognized at ~42kDa. On 4 hr AA starvation, only in Atg8-WT, a lower doublet band is observed (see arrow), but not in the mCherry::Atg8 G→A, suggesting defective lipidation. Loading is shown as a total protein stain for the blot.

(B) Confocal micrographs of single optical-slices of *Drosophila* S2R+ cells transiently transfected with (a, b) mCherry::Atg8-WT (red) or (c, d) mCherry::Atg8-PE- (red) stained with Lamin (blue) in either well-fed state (a, c) or AA-starved (b, d). Scale bar is 2um and right most panel shows DIC image merge. Atg8 vesicles were segmented within each cell volume by performing morphological filtering (See Methods). Representative images of thresholding analyses corresponding to confocal images are shown in right most panel. Red represents the vesicular segmentation. Portion of Atg8 signal trapped in the vesicles, defined as the ratio between Atg8 vesicle intensity and whole cell is quantified. Each dot represents one cell. P-value calculated using the two-sided Wilcoxon rank sum tests. Note the significant increase in Atg8-WT vesicular/whole cell ratio during starvation is not observed in Atg8-PE-. Atg8-PE- vesicle amounts phenocopy Atg8-WT in the fed state.

Figure 4: *Effects of lipidation-defective Atg8 expression in fly fat in relation to Upd2*

A Atg8-WT nuclear signal in adult fly fat in correlation to nutrient state

Lpp-Gal4>UAS-GFP::Atg8-WT
anti-GFP anti-Lamin

Normal Diet (a)
High-Sugar Diet (b)
Starved 4-6 hours (c)

A'
GFP::Atg8-WT nuclear mean pixel intensity
p=7.5e-05
p=0.008
p=0.0008
Fed | High-Sugar | Stv

B Atg8-PE- nuclear signal in adult fly fat at 4-hours post-starvation

GFP-tagged Atg8 transgene, anti-Atg8 (antibody), Lamin

a) Lpp-Gal4> UAS-GFP::Atg8-WT
GFP:Atg8-WT

b) Lpp-Gal4> UAS-GFP::Atg8-PE-
GFP:Atg8-PE-

B'
GFP::Atg8 nuclear mean pixel intensity
p=2.43e-27
p=0.94/NS
p=1.7e-20
Stv Atg8-WT | Fed Atg8-PE- | Stv Atg8-PE-

C Upd2 endogenous nuclear signal in flies with fat-specific expression of Atg8-WT/PE-

upd2-HA (genomic knock-in); ppl-Gal4tubGal80ts> UAS-X

Upd2 Nuclear mean intensity

Upd2 nuclear signal on starvation ↑
p=1.0e-09****

Upd2 nuclear signal on starvation ↓
p=0.003

Fed | Stv (Atg8-WT) | Fed | Stv (Atg8-PE-)

D Triacylglycerol (TAG) mobilization of flies with fat -specific expression of Atg8-WT/PE-

a) Lpp-Gal4> UAS-X
upd2+ (endogenous levels)
TAG level (fold change)
p=0.04 *
p=0.3 ns
Fed | Stv (Atg8-WT) | Fed | Stv (Atg8-PE-)

b) upd2-del; Lpp-Gal4> UAS-X
upd2-deletion
TAG level (fold change)
p=0.02 *
p=0.03 *
Fed | Stv (Atg8-WT) | Fed | Stv (Atg8-PE-)

E Starvation survival of flies with fat -specific expression of Atg8-WT/PE-

a) Lpp-Gal4>UAS-X
upd2+ (endogenous levels)
Probability of Survival
— UAS-Atg8-WT
— UAS-Atg8-PE-
— UAS-Control
Days on starvation diet
p-values:
Atg8-WT vs control= ns (0.1608)
Control vs Atg8-PE-< p=0.0001***
Atg8-WT vs Atg8-PE-< p=0.0001***

b) upd2-del; Lpp-Gal4>UAS-X
upd2-deletion
Probability of Survival
— UAS-Atg8-WT
— UAS-Atg8-PE-
— UAS-Control
Days on starvation diet
p-values:
Atg8-WT vs control= ns (0.2307)
Control vs Atg8-PE-=ns (0.6329)
Atg8-WT vs Atg8-PE-= ns (0.6579)

Figure 4. Effects of lipidation-defective Atg8 expression in fly fat in relation to Upd2.

(A) <u>Atg8-WT::GFP nuclear levels in fly fat correlate with nutrient state:</u> Confocal micrographs of XY-slices from fixed and immunostained adult fly fat from flies expressing GFP::Atg8-WT transgene stained with anti-GFP (green) to detect Atg8. Atg8's nuclear localization in three different nutrient states- normal lab food (a), 14- day high sugar diet (b) and acute starvation (c) are shown.

(B) <u>Lipidation defective Atg8 (Atg8-PE-) remains nuclear in *Drosophila* adult fly fat on acute starvation:</u> Confocal micrographs of single optical sections of fixed and immunostained [anti-GFP(green); anti-Atg8 (red); anti-Lamin (blue)] adult fly fat from transgenic flies- starved for 4 hours- expressing GFP-tagged Atg8-cDNAs for (Da) Atg8-WT (*UAS-GFP::Atg8-WT*) or (b) lipidation-defective Atg8 (*UAS-GFP::Atg8-PE-*) under the control of a fat-cell specific promoter (*Lpp-Gal4*) in an Atg8-WT background. Scale bar represents 2µM.

In A' and B' the amount of GFP::Atg8 in the nucleus in quantified in using 3D-volumteric image segmentation (see methods). Each circle represents a nucleus and statistical significance is calculated using two-sided Wilcoxon sum rank tests.

(C) <u>Effect of fat-specific expression of Atg8-WT/PE- on Upd2 nuclear accumulation in adult fly fat tissue:</u> Upd2-GFP endogenous nuclear accumulation is quantified using 3D-volumetric segmentation-based calculations in flies with acute fat-specific expression of Atg8-WT or Atg8-PE- in fed versus 4 hours starvation. Each dot represents a fat cell nucleus, 75-100 fat nuclei were counted per condition per genotype. Statistical significance is quantified by the two-sided Wilcoxon test.

For D and E, Flies expressing cDNAs (UAS-X) for GFP::Atg8-WT, GFP::Atg8-PE- and control (*UAS-Luciferase*), specifically in fat tissue (*Lpp-Gal4>UAS-X*), with endogenous levels of Upd2 (a) or upd2-deletion (*upd2Δ; Lpp-Gal4>UAS-X*).

(D) <u>Effect of fat-specific expression of Atg8-PE- on triglyceride mobilization during starvation in the presence and absence of endogenous Upd2:</u> TAG levels at fed state were held at 1, and fold change in TAG at starvation was calculated. Error bars represent standard deviation of 9 biological replicates. Statistical significance is calculated using the two-tailed student's T test.

(E) <u>Effect of fat-specific expression of Atg8-PE- on starvation survival in the presence and absence of endogenous Upd2:</u> Probability of survival of transgenic flies with fat-specific overexpression of Atg8-PE- in a WT (a) and upd2 deletion background (b) are shown. Statistical significance is calculated using the Log-rank (Mantel-Cox) test. Per genotype >100 flies were used per experiment. Data shown is then consolidation of three independent experiments.

Figure S5: companion to Main Figure 4

Atg8-Trojan-Gal4 .. ± ♀ X w1118 .. UAS-Atg8-WT or UAS-Atg8-PE- ♂
FM7a   '' +   Y '' UAS-Atg8-WT   UAS-Atg8-PE-

		Transgene on chromosome III			
		UAS-Atg8-WT		UAS-Atg8-PE-	
		# males	% males	# males	% males
X chromosome genotype	Atg8-Trojan-Gal4	36	46.8	0	0
	FM7a	41	53.2	36	100

B | *Atg8 mRNA levels in fat-specific Atg8-WT/PE- expression compared to wild-type controls*

Figure S5: Companion to Main Figure 4.

(C) Assessing the rescue of an Atg8 knock out allele- *Atg8-Trojan-Gal4* (see methods), by the UAS-Atg8 transgenes. In the text box is outlined the crossing scheme depicting the Atg8-Trojan-Gal4 allele rescue with wild-type Atg8 transgene (UAS-GFP::Atg8-WT) or the lipidation-defective Atg8 transgene (UAS-GFP::Atg8-PE-). Table shows the number of progenies counted from the cross and the % of transgenic rescue. While the expected percentage of flies with UAS-GFP-Atg8-WT, as expected the UAS-Atg8-PE- (lipidation-defective version) does not rescue the Atg8 knockout allele.

(D) Atg8 mRNA levels assessed by RNAseq in the indicated genotypes. Each bar represents difference in fold change between starved and fed state of the indicated genotype. Error bars represent the standard deviation of log2 fold change among biological replicates (n=3). Significance was calculated using t-test and followed by *post-hoc* analyses of q-values to account for false discovery rate (See Methods). Values of q<0.05 were considered significant. Atg8 transgenes (red and violet bars) are upregulated on starvation and expressed at higher levels than control (grey). The difference between Atg8-WT and Atg8-PE- expression is not significant.

Figure 5: *Upd2's nuclear retention promotes starvation resilience and hunger-driven feeding*

Figure 5. Upd2's nuclear retention promotes starvation resilience and hunger-driven feeding.

(A) and (B) Effect of fat-specific expression of Upd2-AIM- on high sugar diet and starvation diets: Flies expressing cDNAs (UAS-X) for Upd2-WT, Upd2-AIM- and control (UAS-Luciferase) specifically in fly fat (*upd2Δ Lpp-Gal4>UAS-X*) are subjected to (A) 30% high-sugar diet (HSD) and (B) 1% sucrose agarose diet. Per genotype >100 flies were used per experiment. Data shown is a consolidation of three independent experiments. Statistical differences in starvation survival were quantified using the Log-rank (Mantel-Cox) test. See also companion figure S6B for data on Upd2-NES$^+$AIM$^-$ starvation survival.

(C) Effect of fat-specific expression of Upd2-AIM- on triglyceride mobilization during starvation: Flies expressing cDNAs (UAS-X) for Upd2-WT, Upd2-AIM- and control (UAS-Luciferase) specifically in fat of flies in a Upd2 deletion background (*upd2Δ; Lpp-Gal4>UAS-X*. TAG levels at fed state were held at 1, and fold change in TAG at starvation was calculated. Error bars represent standard deviation of 9 biological replicates. Statistical significance is calculated using the two-tailed student's T test. Statistical significance is calculated using the two-tailed student's T-test.

(D) Effect of fat-specific expression of Upd2-AIM- on hunger-driven response: Graph depicts boxplots of lick events of 7d old adult flies overexpressing cDNAs (UAS-X) for Upd2-WT, Upd2-AIM- and control (UAS-Luciferase) specifically in fly fat (*upd2Δ; Lpp-Gal4>UAS-X*) assessed for feeding behavior using FLIC (Fly liquid interaction counter- see methods). Flies experience either uninterrupted access to food (Fed *ad libitum*) or a 16 hour fast on 1% agarose (Stv). Number of licks was determined through contact using a fly liquid interaction counter (FLIC) on fed and stv flies in parallel for the first 3 hours after the 16-hour starvation period of the Stv group flies. Since flies were monitored in parallel, data was normalized as a percentage to the median fed lick values for each respective genotype, with median fed values being considered 100% lick events. Black dots represent individual fly licks (n=12). Significant differences between fed and starved groups were calculated using a t-test. Note only the UAS-Upd2AIM show a significant increase in licks events in the starved group compared to the fed control (p=0.002).

Figure S6: *companion to Main Figure 5*

Upd2::GFP genomic locus tag knock-in

A — Upd2 nuclear signal in adult fly fat on high-Sugar Diet (HSD)

p=3.09e-12

Upd2 nuclear mean pixel intensity

Normal | HSD

B — upd2Δ; Lpp-Gal4>UAS-Upd2-X::mCherry

Upd2-WT vs. Upd2-NES+AIM- fat-specific overexpression

Upd2-X::mcherry (anti-mcherry), anti-Lamin

UAS-Upd2-WT | UAS-Upd2-AIM- | UAS-Upd2-NES+AIM-

a — Upd2-WT | b — Upd2-AIM- | c — Upd2-NES+AIM-

a' anti-mCherry (LUT) | b' anti-mCherry (LUT) | c' anti-mCherry (LUT)

High / Low

C — Starvation survival in flies with fat-specific over-expression of Upd2-NES+AIM-

upd2Δ; Lpp-Gal4>UAS-X

Probability of Survival

— UAS-Luciferase
— UAS-Upd2NES/AIM

p-value <0.0001 (****)

Day

D — Transient transfection of Upd2-X::GFP in Drosophila S2R+ cells

Upd2-AIM-::GFP vs. Upd2-NES+AIM-::GFP localization

Upd2::GFP (anti-GFP), DAPI

Upd2-WT::GFP | Upd2-AIM-

Fed
a — Fed — Upd2-WT | c — Fed — Upd2-AIM-
Stv, 8hr
b — Stv — Upd2-WT | d — Stv — Upd2-AIM-

5 µm

Upd2-NES+AIM-

Fed
e — Fed — NES+AIM-
Stv, 8hr
f — Stv — NES+AIM-

D'

Upd2::GFP Nuclear/Cell intensity ratio

p=1.39e-09
p=5.27e-10

AIM-FED | NES+AIM-FED | NES+AIM-STV

E — Upd2::GFP transfected in S2R+ cells

Upd2-WT vs. Upd2-NES+AIM- secretion in S2R+ cells

% change in secretion of Upd2::GFP

□ Upd2-WT
□ Upd2-NES+AIM-

p=0.0018

0%
-60%

Upd2-WT | Upd2-NES+AIM-

Figure S6: Companion to Main Figure 5.

(A) Confocal micrographs of single optical sections of fixed and immunostained adult fly fat from transgenic flies expressing mCherry tagged Upd2-cDNAs for (a, a') Upd2-WT (*UAS-Upd2-WT*); (b, b') Upd2's AIM mutated (*UAS-Upd2-AIM-*); (c,c') Upd2-AIM- engineered with a N-terminal canonical nuclear export signal (NES) (*UAS-Upd2-NES+-AIM-*) under the control of a fat-cell specific promoter (Lpp-Gal4) in a upd2-deletion background (*upd2Δ; Lpp-Gal4*). In Upd2-WT in fed state is detectable in very low levels in the nucleus (see look up table (LUT)) and yellow arrows in a Upd2-AIM- shows significantly nuclear accumulation even in a fed state (b'). Whereas addition of NES+ to Upd2-AIM (c,c') causes Upd2 to be localized in a diffuse pattern in the cytosol and localize to periphery of nuclear lamina see arrows.

(B) Flies expressing cDNAs (UAS-X) for *Upd2-NES+AIM-* and control (*UAS-Luciferase*) specifically in fly fat in upd2 deletion background (*upd2Δ Lpp-Gal4>UAS-X*). Transgenic flies with fat-specific overexpression of AIM- that has a canonical NES (Upd2-NES+-AIM-) results in starvation-sensitivity in comparison to controls as assessed by the Log-rank (Mantel-Cox) test. See also Main Figure 5B on characterization of starvation sensitivity of *UAS-Upd2- AIM-*.

(C) Confocal micrographs of single optical-slices of *Drosophila* S2R+ cells transiently co-transfected with Upd2::GFP variants (a, b) WT; (c, d) (e, f) NES+AIM- Upd2::GFP (green) stained with DAPI (blue) and anti-GFP (green) in a fed state (a ,c,) or AA-starved for 8 hours (b, d). Scale bars are 5um right most panel shows DIC image merge. Note that addition of NES+ renders Upd2-AIM- cytosolic in both fed and starved states. Note also that unlike Upd2::WT which are localized to punctate in fed states(Ca), the addition of NES+ to Upd2-AIM- display a diffuse cytosolic localization (Ce). C' Quantification of nuclear to cytosolic ratio of Upd2-GFP based on 3D-volumetric methods, to assess the difference between Upd2-AIM- in fed state versus Upd2-NES+-AIM-. Upd2-NES+AIM- is significantly less nuclear than Upd2-AIM- in both fed and starved states.

(D) Normalized percent fold change in secreted GFP signal detected by GFP sandwich ELISA assay performed on conditioned media of S2R+ cells transiently transfected with Upd2::GFP variants: WT; NES+AIM- the amount of Upd2-WT::GFP in a 24-hour period was assessed. Statistical significance is quantified by unpaired two-tailed t-test on 6 biological replicates per condition. Note that despite being localized to the cytosol Upd2-NES+-AIM- is not secreted, consistent with the requirement for Atg8 during secretion.

Figure 6

Upd2's localization is regulated by Atg8 during different nutrient states

Figure 6. Summary of findings: Upd2's intracellular localization is contingent on nutrient state and Atg8.

Each illustration summarizes the key findings of the study. The upper panel is the key. Top three panels are fed state and lower three panels are in acute starvation (stv). Each panel illustrates a single cell. Notations: N- nucleus, C- cytosol, E- extracellular space.

- *Top left:* In a normal baseline fed state, the ortholog of human Leptin in flies, Upd2, enters the nucleus. Subsequently, Upd2's Atg8-interaction motif (AIM) mediates interactions with Atg8 that enables Upd2 nuclear exit to the cytosol and then Upd2 is released to the extracellular space.

- *Bottom left:* During starvation, Atg8 transits to the cytosol and is lipidated. Depletion of the nuclear pool of Atg8 on starvation increases Upd2 nuclear accumulation and causes cells to retain the adipokine on starvation. Thus, Atg8's lipidation state serves as a switch to license adipokine Upd2's cytosolic or nuclear localization.

- *Top middle and right:* When Atg8's interaction with Upd2 is disrupted (Upd2-AIM-) and when Atg8 levels are reduced by RNAi-mediated knock-down, Upd2 accumulates in the nucleus even in fed cells. This provides evidence for Atg8's role in facilitating Upd2's cytosolic localization in fed cells. We note that constitutive expression of Upd2-AIM- in fly fat renders flies sensitive to a HSD. This suggests that maintaining Upd2 in the nucleus, during a nutrient state where it is usually cytosolic, has a negative impact on resilience to a surplus diet.

- *Bottom middle:* Upd2-AIM- remains in the nucleus on starvation, while Atg8 translocates to the cytosol. Upd2-AIM- flies are starvation resilient and upregulate a systemic innate immune gene expression signature.

- *Bottom right:* Lipidation-defective Atg8-PE- continues to be nuclear on starvation. Presence of Atg8-PE- allows Upd2 to be cytosolic even on starvation and renders flies sensitive to starvation.