1 **Cluster-specific gene markers enhance *Shigella* and Enteroinvasive *Escherichia coli in***
2 ***silico* serotyping**
3

4 Xiaomei Zhang[1], Michael Payne[1], Thanh Nguyen[1], Sandeep Kaur[1], Ruiting Lan[1]*
5
6 [1]School of Biotechnology and Biomolecular Sciences, University of New South Wales, Sydney,
7 New South Wales, Australia
8
9
10
*Corresponding Author
Email: r.lan@unsw.edu.au
Phone: 61-2-9385 2095
Fax: 61-2-9385 1483

Keywords: Phylogenetic clusters, cluster-specific gene markers, *Shigella*/EIEC serotyping
Running title: *Shigella* EIEC Cluster Enhanced Serotyping
Repositories: Raw sequence data are available from NCBI under the BioProject number
PRJNA692536.

20
21
22

Abstract

Shigella and enteroinvasive *Escherichia coli* **(**EIEC) cause human bacillary dysentery with similar invasion mechanisms and share similar physiological, biochemical and genetic characteristics. The ability to differentiate *Shigella* and EIEC from each other is important for clinical diagnostic and epidemiologic investigations. The existing genetic signatures may not discriminate between *Shigella* and EIEC. However, phylogenetically, *Shigella* and EIEC strains are composed of multiple clusters and are different forms of *E. coli*. In this study, we identified 10 *Shigella* clusters, 7 EIEC clusters and 53 sporadic types of EIEC by examining over 17,000 publicly available *Shigella*/EIEC genomes. We compared *Shigella* and EIEC accessory genomes to identify the cluster-specific gene markers or marker sets for the 17 clusters and 53 sporadic types. The gene markers showed 99.63% accuracy and more than 97.02% specificity. In addition, we developed a freely available *in silico* serotyping pipeline named *Shigella* EIEC Cluster Enhanced Serotype Finder (ShigEiFinder) by incorporating the cluster-specific gene markers and established *Shigella*/EIEC serotype specific O antigen genes and modification genes into typing. ShigEiFinder can process either paired end Illumina sequencing reads or assembled genomes and almost perfectly differentiated *Shigella* from EIEC with 99.70% and 99.81% cluster assignment accuracy for the assembled genomes and mapped reads respectively. ShigEiFinder was able to serotype over 59 *Shigella* serotypes and 22 EIEC serotypes and provided a high specificity with 99.40% for assembled genomes and 99.38% for mapped reads for serotyping. The cluster markers and our new serotyping tool, ShigEiFinder (https://github.com/LanLab/ShigEiFinder), will be useful for epidemiologic and diagnostic investigations.

Impact statement

The differentiation of *Shigella* strains from enteroinvasive *E. coli* (EIEC) is important for clinical diagnosis and public health epidemiologic investigations. The similarities between *Shigella* and EIEC strains make this differentiation very difficult as both share common ancestries within *E. coli*. However, *Shigella* and EIEC are phylogenetically separated into multiple clusters, making high resolution separation using cluster specific genomic markers possible. In this study, we identified 17 *Shigella* or EIEC clusters including five that were newly identified through examination of over 17,000 publicly available *Shigella* and EIEC genomes. We further identified an individual or a set of cluster-specific gene markers for each cluster using comparative genomic analysis. These markers can then be used to classify isolates into clusters and were used to develop an *in silico* pipeline, ShigEiFinder

(https://github.com/LanLab/ShigEiFinder) for accurate differentiation, cluster typing and serotyping of *Shigella* and EIEC from Illumina sequencing reads or assembled genomes. This study will have broad application from understanding the evolution of *Shigella*/EIEC to diagnosis and epidemiology.

Data summary

Sequencing data have been deposited at the National Center for Biotechnology Information under BioProject number PRJNA692536.

Introduction

Shigella is one of the most common etiologic agents of foodborne infections worldwide and can cause diarrhea with a very low infectious dose (1, 2). The infections can vary from mild diarrhea to severe bloody diarrhea referred to as bacillary dysentery. The estimated cases of *Shigella* infections are 190 million with at least 210,000 deaths annually, predominantly in children younger than 5 years old in developing countries (3-7). *Shigella* infections also have a significantly impact on public health in developed countries although most cases are travel-associated (8).

The *Shigella* genus consists of four species, *Shigella sonnei*, *Shigella flexneri*, *Shigella boydii* and *Shigella dysenteriae* (9). Serological testing further classifies *Shigella* species into more than 55 serotypes through the agglutination reaction of antisera to *Shigella* serotype specific O-antigens (10, 11). Up to 89.6% *Shigella* infections were caused by *S. flexneri* (65.9%) and *S. sonnei* (23.7%) globally (12, 13). The predominant serotype reported in *Shigella* infections has been *S. flexneri* serotype 2a while *S. dysenteriae* serotype 1 has caused the most severe disease (11, 14). Note that for brevity, in all references to *Shigella* serotypes below, *S. sonnei*, *S. flexneri*, *S. boydii* and *S. dysenteriae* are abbreviated as SS, SF, SB and SD respectively and a serotype is designated with abbreviated "species" name plus the serotype number e.g. *S. dysenteriae* serotype 1 is abbreviated as SD1.

Enteroinvasive *Escherichia coli* (EIEC) is a pathovar of *E. coli* that causes diarrhoea with less severe symptoms to *Shigella* infections in humans worldwide, particularly in developing countries (8, 13, 15-18). EIEC infections in developed countries are mainly imported (19). EIEC has more than 18 specific E. coli O-serotypes (19, 20). Although the incidence of EIEC

is low (17), EIEC serotypes have been associated with outbreaks and sporadic cases of infections (20-22). In contrast to *Shigella*, EIEC infections are not notifiable in many countries (23, 24).

Shigella and EIEC have always been considered very closely related and share several characteristics (25-28). *Shigella* and EIEC are both non-motile and lack the ability of ferment lactose (24). Some of EIEC O antigens are identical or similar to *Shigella* O antigens (O112ac, O124, O136, O143, O152 and O164) (26, 29-31). Furthermore, *Shigella* and EIEC both carry the virulence plasmid pINV, which encodes virulence genes required for invasion (32, 33) and contain *ipaH* (invasion plasmid antigen H) genes with the exception of some SB13 isolates (10, 23, 24, 34, 35). *Shigella* and EIEC have arisen from *E. coli* in multiple independent events and should be regarded as a single pathovar of E. coli (25, 26, 28, 36-38). Previous phylogenetic studies suggested that *Shigella* isolates were divided into 3 clusters (C1, C2 and C3) with 5 outliers (SS, SB13, SD1, SD8 and SD10) (25, 38) whereas EIEC isolates were grouped into four clusters (C4, C5, C6 and C7) (26). The seven *Shigella*/EIEC clusters and 5 outliers of *Shigella* are within the broader non-enteroinvasive *E. coli* species except for SB13 which is closer to *Escherichia albertii* (39, 40). WGS-based phylogenomic studies have also defined multiple alternative clusters of *Shigella* and EIEC (23, 28, 41).

The traditional biochemical test for motility and lysine decarboxylase (LDC) activity (42) and molecular test for the presence of *ipaH* gene have been used to differentiate *Shigella* and EIEC from non-enteroinvasive E. coli (24, 43-45). Agglutination with *Shigella*/EIEC associated antiserum further classify *Shigella* or EIEC to serotype level. However, cross-reactivity, strains not producing O antigens, and newly emerged *Shigella* serotypes may all prevent accurate serotyping (10, 46). Serotyping by antigenic agglutination is being replaced by molecular serotyping (47, 48), which can be achieved through examination of the sequences of O antigen biosynthesis and modification genes (8, 24, 49-52).

Recently, PCR-based molecular detection methods targeting the gene *lacY* were developed to distinguish *Shigella* from EIEC (53, 54). However, the ability of the primers described in these methods to accurately differentiate between *Shigella* and EIEC was later questioned (23, 28). With the uptake of whole-genome sequencing technology, several studies have identified phylogenetic clade specific markers, species specific markers and EIEC lineage-specific genes for discrimination between *Shigella* and EIEC and between *Shigella* species (23, 27, 28, 41, 55,

56). More recently, genetic markers *lacY*, *cadA*, *Ss_methylase* were used for identification of *Shigella* and EIEC (10). However, these markers failed to discriminate between *Shigella* and EIEC when a larger genetic diversity is considered (23, 28, 55). A Kmer-based approach can identify *Shigella* isolates to the species level but misidentification was also observed (56).

In this study, we aimed to i), identify phylogenetic clusters of *Shigella* and EIEC through large scale examination of publicly available genomes; ii), identify cluster-specific gene markers using comparative genomic analysis of *Shigella* and EIEC accessory genomes for differentiation of *Shigella* and EIEC; iii), develop a pipeline for *Shigella* and EIEC *in silico* serotyping based on the cluster-specific gene markers combined with *Shigella* and EIEC serotype-specific O antigen and H antigen genes. We demonstrate that these cluster-specific gene markers enhance *in silico* serotyping using genomic data. We also developed an automated pipeline for cluster typing and serotyping of *Shigella*/EIEC from WGS data.

Materials and Methods

Identification of *Shigella*/EIEC isolates from NCBI database

 E. coli/Shigella isolates from the NCBI SRA (National Center for Biotechnology Information Sequence Read Archive) as May of 2019 were queried. Raw reads were retrieved from ENA (European Nucleotide Archive). The *ipaH* gene (GenBank accession number M32063.1) was used to screen E. coli/*Shigella* reads using Salmon v0.13.0 (57). Taxonomic classification for *E. coli/Shigella* was confirmed by Kraken v1.1.1 (58). Molecular serotype prediction of *ipaH* negative *Shigella* isolates was performed by ShigaTyper v1.0.6 (10). Isolates that were *ipaH* positive and isolates with designation of SB13 by ShigaTyper were selected as *Shigella*/EIEC database.

The sequence types (STs) and ribosomal STs (rSTs) of *ipaH* gene negative E. coli (non-enteroinvasive E. coli) isolates were examined. STs and rSTs for these isolates were obtained from the E. coli/*Shigella* database in the Enterobase (59) as of May 2019. For STs and rSTs with only one isolate, the isolates were selected. For STs and rSTs with more than one isolates, one representative isolate for each ST and rST were randomly selected. In total, 12,743 ipaH negative E. coli isolates representing 3,800 STs and 11,463 rSTs were selected as non-enteroinvasive E. coli control database.

Genome sequencing

159 Whole-genome sequencing (WGS) of 31 EIEC strains used in a previous study (26) was
160 performed by Illumina NextSeq (Illumina, Scoresby, VIC, Australia). DNA libraries were
161 constructed using Nextera XT Sample preparation kit (Illumina Inc., San Diego, CA, USA) and
162 sequenced using the NextSeq sequencer (Illumina Inc.). FASTQ sequences of the strains
163 sequenced in this study were deposited in the NCBI under the BioProject (PRJNA692536).
164

165 **Genome assembly and data processing**
166 Raw reads were *de novo* assembled using SPADES v3.14.0 assembler with default settings
167 [http://bioinf.spbau.ru/spades] (60). The metrics of assembled genomes were obtained with
168 QUAST v5.0.0 (61). Three standard deviations (SD) from the mean for contig number, largest
169 contig, total length, GC, N50 and genes were used as quality filter for assembled genomes.
170

171 The STs for isolates in *Shigella*/EIEC database was checked by using mlst
172 (https://github.com/tseemann/mlst) with the *E. coli* scheme from PubMLST (62). rSTs were
173 extracted from the E. coli/*Shigella* rMLST database in Enterobase (59) as of May 2019.
174 Serotype prediction for isolates in *Shigella*/EIEC was performed by ShigaTyper v1.0.6 (10).
175 Serotyping of E. coli O and H antigens were predicted by using SerotypeFinder v2.0.1 (63).
176

Selection of isolates for *Shigella*/EIEC identification dataset
The selection of isolates for the identification dataset was based on the representative isolates
for each ST, rST and serotype of *Shigella* and EIEC in the *Shigella*/EIEC database. For STs
and rSTs with only one isolate, the isolate was selected. For STs and rSTs with more than one
isolates, one representative isolate for each ST, rST was randomly selected. A representative
experimentally confirmed isolate of each serotype of *Shigella* and EIEC was also randomly
selected. 72 ECOR strains downloaded from Enterobase (59) and 18 *E. albertii* strains were
used as controls for the identification dataset. The details of the identification dataset are listed
in Table S1. The remaining isolates in *Shigella*/EIEC database were referred as validation
dataset (Table S2).
187

188 The identification dataset was used to characterise the phylogenetic relationships of *Shigella*
189 and EIEC. The identification dataset was also used to identify cluster-specific genes. The
190 validation dataset was used to evaluate the performance of cluster-specific gene markers using
191 the *in-silico* serotyping pipeline.
192

Phylogeny of *Shigella* and EIEC based on WGS

Three phylogenetic trees including identification tree, confirmation tree and validation tree
were constructed by Quicktree v1.3 (64) with default parameters to identify and confirm the
phylogenetic clustering of *Shigella* and EIEC isolates. The phylogenetic trees were visualised
by Grapetree and ITOL v5 (65, 66).

The identification phylogenetic tree was generated based on isolates in the identification
dataset for the characterisation of clusters of *Shigella* and EIEC isolates (Fig. 1). A subset of
485 isolates known to represent each identified cluster from the identification dataset were then
selected. The subset of 485 isolates from the identification dataset and 1,872 non-
enteroinvasive E. coli isolates from non-enteroinvasive E. coli control dataset (2,357 isolates
total) were used to construct a confirmation tree. This tree was used for confirmation of the
phylogenetic relationships between identified *Shigella*/EIEC clusters in the identification
dataset and non-enteroinvasive E. coli isolates. The validation tree was generated based on
1,159 representative isolates from the validation dataset that were selected in the same way as
the identification dataset and a subset of 485 isolates from the identification dataset to assign
validation dataset isolates to clusters.

Investigation of *Shigella* virulence plasmid pINV

The presence of *Shigella* virulence plasmid pINV in isolates were investigated by using BWA-
MEM v0.7.17 (Burrows-Wheeler Aligner) (67) to align isolate raw reads onto the reference
sequence of pINV (68) (NC_024996.1). Mapped reads were sorted and indexed using
Samtools v1.9 (69). The individual gene coverage from mapping was obtained using Bedtools
coverage v2.27.1 (70).

Identification of the cluster-specific gene markers

Cluster-specific gene markers were identified from *Shigella/*EIEC accessory genomes. The
genomes from the identification dataset were annotated using PROKKA v1.13.3 (71). Pan- and
core-genomes were analysed by roary v3.12.0 (72) using an 80% sequence identity threshold.
The genes specific to each cluster were identified from the accessory genes with an in-house
python script. In this study, the number of genomes from a given cluster containing all specific
genes for that cluster was termed true positives (TP), the number of genomes from the same
cluster lacking any of those same genes was termed false negatives (FN). The number of

226 genomes from other clusters containing all of those same genes was termed false positives
227 (FP).
228
229 The sensitivity (True positive rate, TPR) of each cluster-specific gene marker was defined as
230 TP/(TP+FN). The specificity (True negative rate, TNR) was defined as TN/(TN+FP).
231

Validation of the cluster-specific gene markers

233 The ability of cluster-specific gene markers to assign *Shigella*/EIEC isolates was examined by
234 using BLASTN to search against the validation dataset (Table S2) and non-enteroinvasive E.
235 coli control database for the presence of any of the cluster-specific gene marker or a set of
236 cluster-specific gene markers. The BLASTN thresholds were defined as 80% sequence identity
237 and 50% gene length coverage.
238

Development of ShigEiFinder, an automated pipeline for molecular serotyping of
***Shigella*/EIEC**

ShigEiFinder was developed using paired end illumina genome sequencing reads or assembled
genomes identify cluster-specific gene markers combined with *Shigella*/EIEC serotype specific
O antigen genes (wzx and wzy) and modification genes (Fig. 2, Data S1). We used the same
signature O and H sequences from ShigaTyper and SerotypeFinder (Data S2) (10, 63). These
include *Shigella* serotype-specifc wzx/wzy genes and modification genes from ShigaTyper and
E. coli O antigen and *fliC* (H antigen) genes from SerotypeFinder. *ipaH* gene and 38 virulence
genes used in analysis of virulence of 59 sporadic EIEC isolates were also included in the
typing reference sequences database. Seven House Keeping (HK) genes -*recA*, *purA*, *mdh*, *icd*,
gyrB, *fumC* and *adk* downloaded from NCBI were used for contamination checking.
250

Raw reads were aligned to the typing reference sequences by using BWA-MEM v0.7.17 (67).
The mapping length percentage and the mean mapping depth for all genes were calculated
using Samtools coverage v1.10 (69). To determine whether the genes were present or absent,
50% of mapping length for all cluster-specific genes, virulence genes and O antigen genes and
10% for *ipaH* gene were used as cutoff value. The ratio of mean mapping depth to the mean
mapping depth of the 7 HK genes was used to determine a contamination threshold with ratios
less than 1% for *ipaH* gene and less than 10% for other genes assigned as contamination.
Reads coverage mapped to particular regions of genes were checked by using samtools
mpileup v1.10.

Assembled genomes were searched against the typing reference sequences using BLASTN v2.9.0 (73) with 80% sequence identity and 50% gene length coverage for all genes with exception of *ipaH* gene which was defined as 10% gene length coverage.

ShigEiFinder was tested with the identification dataset and validated with the *Shigella*/EIEC validation dataset and non-enteroinvasive E. coli control database. The specificity defined as (1 - the number of non-enteroinvasive E. coli isolates being detected / the total number of non-enteroinvasive E. coli isolates) * 100.

Results
Screening sequenced genomes for *Shigella*/EIEC isolates

We first screened available *E. coli* and *Shigella* genomes based on the presence of *ipaH* gene. We examined 122,361 isolates with the species annotation of E. coli (104,256) or *Shigella* (18,105) with paired end illumina sequencing reads available in NCBI SRA database. Of 122,361 isolates, 17,989 isolates were positive to the *ipaH* gene including 455 out of 104,256 E. coli isolates and 17,434 out of 18,105 *Shigella* isolates. The 17,989 *ipaH* positive *E. coli* and *Shigella* genomes and 571 *ipaH* negative "*Shigella*" genomes were checked for taxonomic classification and genome assembly quality. 17,320 *ipaH* positive *E. coli* and *Shigella* genomes and 246 *ipaH* negative "*Shigella*" genomes passed quality filters. Among 246 *ipaH* negative "*Shigella*" genomes, 11 isolates belonged to SB13 by using ShigaTyper (10) while the remaining 235 isolates were classified with taxonomic identifier of E. coli by Kraken v1.1.1 (58) and were removed from analysis. A total of 17,331 genomes including 17,320 *ipaH* positives and 11 SB13 genomes were selected to form the *Shigella*/EIEC database, which contained 429 genomes with species identifier of *E. coli* and 16,902 genomes with species identifier of *Shigella.*

Isolates in *Shigella*/EIEC database were typed using MLST, ShigaTyper and SerotypeFinder. MLST and rMLST divided the 17,331 *Shigella*/EIEC isolates into 252 STs (73 isolates untypeable by MLST) and 1,128 rSTs (3,513 isolates untypeable by rMLST). Of 16,902 genomes with species identifier of *Shigella*, 8,313 isolates and 8,189 isolates were typed as *Shigella* and EIEC respectively by ShigaTyper while 400 isolates were untypeable. ShigaTyper

typed the majority of the 8,313 isolates as SF (66.82%) including 25.43% SF2a isolates,

followed by SS (19.69%), SB (7.22%) and SD (6.27%).

SerotypeFinder typed 293 of the 429 *E. coli* genomes into 71 *E. coli* O/H antigen types.

Among these 293 isolates with typable O/H antigen types, 190 isolates belonged to 22 known

EIEC serotypes (O28ac:H-, O28ac:H7, O29:H4, O112ac:H26, O121:H30, O124:H30,

O124:H24, O124:H7, O132:H7, O132:H21, O135:H30, O136:H7, O143:H26, O144:H25,

O152:H-, O152:H30, O164:H-, O164:H30, O167:H26, O173:H7 and 2 newly emerged EIEC

serotypes O96:H19 and O8:H19) (20-22). The remaining 136 of 429 genomes were O antigen

untypable and typed to 15 H antigen types only by SerotypeFinder, of which H16 was the

predominant type.

Identification of *Shigella* and EIEC clusters

Shigella and EIEC are known to have been derived from *E. coli* independently. To identify

previously defined clusters (25, 26) and any new clusters from the 17,331 *Shigella*/EIEC

genomes, we selected representative genomes to perform phylogenetic analysis as it was

impractical to construct a tree with all genomes. The selection was based on ST, rST and

serotype of the 17,331 *Shigella*/EIEC genomes. One isolate was selected to represent each ST,

rST and serotype for a total of 1,830 isolates. The selection included 252 STs, 1,128 rSTs, 59

Shigella serotypes (21 SB serotypes, 20 SF serotypes, 17 SD serotypes and SS), 22 EIEC

known serotypes and 31 other or partial antigen types. A further 31 in-house sequenced EIEC

isolates, 18 EIEC isolates used in a previous typing study (41), 72 ECOR strains and 18 *E.*

albertii strains were also included to form the identification dataset of 1,969 isolates. Details

are listed in Table S1. A phylogenetic tree was constructed based on the identification dataset

to identify the clusters (Fig. 1).

All known clusters were identified (Fig. 1) including 3 *Shigella* clusters (C1, C2, C3) and 5

outliers (SD1, SD8, SD10, SB13 and SS) as defined by Pupo et al (25) and 4 EIEC clusters

(C4, C5, C6 and C7) defined by Lan et al. (26). Each of these clusters was supported by a

bootstrap value of 80% or greater (Fig. S1). 1,789 isolates of the 1,879 *Shigella*/EIEC isolates

(1,830 isolates from the *Shigella*/EIEC database, 31 in-house sequenced EIEC isolates and 18

EIEC isolates from Hazen *et al.*) fell within these clusters.

Of the remaining 90 *Shigella*/EIEC unclustered isolates, 31 belonged to 5 *Shigella*/EIEC serotypes including 5 SB13 isolates, 8 SB12 isolates, 2 EIEC O135:H30 isolates, 12 EIEC serotype O96:H19 isolates and 4 EIEC O8:H19 isolates, while 59 isolates were sporadic EIEC isolates which are described in detail in the separate section below. The 5 SB13 isolates were grouped into one lineage within *E. coli* and close to known *Shigella*/EIEC clusters rather than the established SB13 cluster outside *E. coli* which was within the *E. albertii* lineage. The former was previously named as atypical SB13 while the latter was previously named as typical SB13 (39). The 8 SB12 isolates formed one single cluster close to SD1 and atypical SB13 clusters. Two EIEC O135:H30 isolates were grouped as a separate cluster close to C5. Twelve isolates belonging to EIEC serotype O96:H19 and 4 isolates typed as O8:H19 were clustered into two separate clusters, both of which were more closely related to SD8 than other *Shigella*/EIEC clusters. Therefore, atypical SB13 and SB12 were defined as new clusters of *Shigella* while EIEC O96:H19, EIEC O8:H19 and EIEC O135:H30 were defined as C8, C9 and C10 respectively. In total there were 10 *Shigella* clusters and 7 EIEC clusters (Table 1).

Analysis of the 59 sporadic EIEC isolates

To determine the phylogenetic relationships of the above defined clusters and the remaining 59 sporadic EIEC isolates within the larger non-enteroinvasive *E. coli* population a confirmation tree was generated using 485 isolates representing the known clusters and 1,872 representative non-*Shigella*/EIEC isolates (Fig. S2). The 59 sporadic EIEC isolates including 2 EIEC isolates M2330 (O152:H51) and M2339 (O124:H7) sequenced in this study and 57 isolates were interspersed among non-*Shigella*/EIEC isolates and did not form large clusters. Groups of these isolates that were not previously identified were named as sporadic EIEC lineage followed by their serotype. For example, M2339 (O124:H7) grouped together with one other EIEC isolate with the same O and H antigens O124:H7 and were named 'sporadic EIEC lineage O124:H7'. There were 53 sporadic EIEC lineages including 5 lineages with 2 or more isolates and 48 lineages with only one isolate. The STs, rSTs and antigen types of these 59 isolates were listed in the Table S1.

Some of the sporadic EIEC isolates fell into STs containing *ipaH* negative isolates. We therefore examined the presence of the pINV virulence plasmid in the sporadic EIEC isolates. We selected 38 genes that are essential for virulence including 35 genes (12 *mxi* genes, 9 *spa* genes, 5 *ipaA-J* genes, 6 *ipgA-F* genes as well as *acp*, *virB*, *icsB*) in the conserved entry region encoding the Mxi-Spa-Ipa type III secretion system and its effectors and 3 regulator genes

(*virF*, *virA* and *icsA*/*virG*) (24, 33, 68) and determined the presence of pINV in the 59 sporadic

EIEC isolates by mapping the sequence reads onto a pINV reference sequence (68). Reads

from 18 non-*Shigella/EIEC* isolates that shared the same ST as one of 58 sporadic isolates

were positive for these genes.

The number of essential virulence genes with mapped reads in the 59 sporadic EIEC isolates

were analysed (Fig. S3). Those isolates containing more than 25 of the 38 essential virulence

genes were defined as virulence plasmid positive. While isolates containing between 13 and 25

were defined as intermediate and less than 13 were defined as virulence plasmid negative.

The 2 newly sequenced sporadic EIEC isolates (M2330 and M2339) were positive for the

virulence plasmid and of the other 57 sporadic EIEC isolates, 39 isolates were positive, 9

isolates were negative and 9 isolates were intermediate (Table S1). The results were compared

with those non-*Shigella/*EIEC isolates belonging to the same ST. The virulence plasmid was

absent in all non-*Shigella/*EIEC isolates while all sporadic EIEC isolates in these STs were

either positive or intermediate. Therefore, this analysis confirmed the sporadic isolates

belonged to EIEC and the STs contained both EIEC and non- EIEC isolates.

Identification of cluster-specific gene markers

In this study, cluster-specific gene markers were either a single gene present in all isolates of a

cluster and absent in all other isolates or a set of genes (two or more) that as a combination

were only found in one cluster. For the marker sets, a subset of cluster-specific gene markers

for a given cluster could be found in other clusters but the entire set was only found in the

target cluster.

Comparative genomic analysis on 1,969 accessory genomes from the identification dataset was

used to identify cluster-specific gene markers or marker sets. Multiple candidate cluster-

specific gene markers or marker sets of markers for each of 17 *Shigella*/EIEC clusters and 53

sporadic EIEC lineages were identified through screening the accessory genes from 1,969

genomes. These gene markers or marker sets were 100% sensitive to clusters but with varying

specificity. The cluster-specific gene markers or marker sets with the lowest FP rates were then

selected from candidate cluster-specific gene markers by BLASTN searches against genomes

in the identification dataset using 80% sequence identity and 50% gene length threshold.

Five single cluster-specific gene markers (C7, C10, SB12, SB13 and atypical SB13) and 12

sets of cluster-specific gene markers (C1, C2, C3, C4, C5, C6, C8, C9, SS, SD1, SD8 and

SD10) were selected for *Shigella*/EIEC cluster typing. The sensitivity and specificity for each

cluster-specific gene marker or a set of cluster-specific gene markers for the identification

dataset were listed in Table 2. The cluster-specific gene markers or marker sets were all 100%

sensitive and 100% specific with the exception of C1 (99.94% specificity), C3 (99.91%

specificity) and SS (99.8% specificity). A single specific gene for each of 53 sporadic EIEC

lineages were also selected with the exception of one lineage which has a set of 2 genes. These

genes were all 100% sensitive and specific for a given sporadic EIEC lineage.

All cluster-specific gene markers, 37 in total (5 single, 32 genes in 12 sets) and 54 sporadic

EIEC lineages specific gene markers were located on chromosome but one of C4 gene markers

and 5 sporadic EIEC lineages specific genes were located on plasmid by NCBI BLAST

searching. None of the cluster-specific gene markers were contiguous in the genomes. The

location of these cluster-specific gene markers was determined by BLASTN against

representative complete genomes of *Shigella*/EIEC containing gene features downloaded from

NCBI GenBank. In those cluster or sporadic lineages with no representative complete genome

specific gene markers were named using their cluster or sporadic EIEC lineage followed by the

cluster or lineage number. For example, C7 specific gene marker was named "C7 specific

gene".

The functional characterization of these specific gene markers were identified from RAST

annotation (74). For 37 cluster-specific gene markers, 22 had known functions and 15 encoded

hypothetical proteins with unknown functions, while 11 sporadic EIEC lineages specific gene

markers were identified with known functions and 43 were hypothetical proteins with

unknown functions. The location and functions of specific gene markers are listed in Table S3.

Validation of cluster-specific gene markers

The ability of cluster-specific gene markers to correctly assign *Shigella*/EIEC isolates was

evaluate with 15,501 *Shigella*/EIEC isolates in the validation dataset, 12,743 isolates from non-

enteroinvasive *E. coli* control database.

Using cluster-specific gene markers, 15,443 of the 15,501 (99.63%) *Shigella*/EIEC isolates

were assigned to clusters which included 15,337 *Shigella* isolates, 102 EIEC isolates, 4

sporadic EIEC isolates, and 38 (0.24%) isolates with more than one clusters. Twenty of the 15,501 (0.13%) *Shigella*/EIEC isolates were not assigned to any of identified clusters.

To confirm the assignment of cluster-specific gene markers, we constructed a "validation" phylogenetic tree (Fig. S4) using 1,159 representative isolates from the validation dataset and a subset of 485 isolates from each cluster from the identification dataset. Isolates that grouped with known cluster isolates (from identification dataset) with strong bootstrap support were assigned to that cluster. All 1,159 isolates were grouped into known clusters on the validation phylogenetic tree. The cluster-specific gene markers assignments were entirely consistent with cluster assignments by phylogenetic tree.

We tested cluster-specific gene markers with the 12,743 non-enteroinvasive *E. coli* isolates. The *Shigella*/EIEC cluster-specific gene markers were highly specific with specificity varying from 98.8% to 100% for cluster-specific genes and 97.02% to 100% for sporadic EIEC specific genes. Details are listed in Table S4.

Development of an automated pipeline for molecular serotyping of *Shigella*/EIEC
Above results showed that cluster-specific gene markers were sensitive and specific and can distinguish *Shigella* and EIEC isolates. We therefore used these genes combined with established *Shigella*/EIEC serotype specific O antigen and H antigen genes to develop an automated pipeline for *in silico* serotyping of *Shigella*/EIEC (Fig. 2).

The pipeline is named *Shigella* EIEC Cluster Enhanced Serotype Finder (ShigEiFinder). ShigEiFinder can process either paired end Illumina sequencing reads or assembled genomes (installable package: https://github.com/LanLab/ShigEiFinder, online too: https://mgtdb.unsw.edu.au/ShigEiFinder/). ShigEiFinder classifies isolates into Non-*Shigella*/EIEC, *Shigella* or EIEC clusters based on the presence of *ipaH* gene, number of virulence genes, cluster specific genes. The "Not *Shigella*/EIEC" assignment was determined by the absence of the *ipaH* gene, virulence genes (<26) and absence of cluster-specific gene markers. The "*Shigella* or EIEC clusters" assignments were made based on the presence of *ipaH* gene, and/or more than 25 virulence genes together with the presence of any of cluster-specific gene markers or marker set, whereas the presence of *ipaH* gene and/or more than 25 virulence genes with absence of any of cluster-specific gene markers were assigned as "*Shigella*/EIEC unclustered".

Shigella and EIEC isolates were differentiated and serotypes were assigned after cluster

assignment. ShigEiFinder predicts a serotype through examining the presence of any of

established *Shigella* serotype specific O antigen and modification genes and E. coli O and H

antigen genes that differentiate the serotypes as ShigaTyper and SerotypeFinder (10, 63). A

"novel serotype" is assigned if no match to known serotypes.

Two pairs of *Shigella* serotypes, SB1/SB20 and SB6/SB10, are known to be difficult to

differentiate as they share identical O antigen genes (10, 46, 75). ShigaTyper used a heparinase

gene for the differentiation of SB20 from SB1 and *wbaM* gene for the separation of SB6 from

SB10. We found that fragments of the heparinase and *wbaM* genes may be present in other

serotypes and cannot accurately differentiate SB1/SB20 and SB6/SB10. We found a SB20

specific gene which encoded hypothetical proteins with unknown functions and located on a

plasmid by comparative genomic analysis of all isolates in C1 accessory genome. The SB20

specific gene can reliably differentiate SB20 from SB1and also one SNP each in *wzx* and *wzy*

genes that can differentiate SB6 from SB10. We used these differences (Data S1) in

ShigEiFinder for the prediction of these serotypes.

The accuracy and specificity of ShigEiFinder in cluster typing

The accuracy of ShigEiFinder was tested with 1,969 isolates (1,969 assembled genomes and

1,951 Illumina reads [note no reads available for 18 EIEC isolates from NCBI) from the

identification dataset and 15,501 isolates from the validation dataset. The results are listed in

Table 3.

ShigEiFinder was able to assign 99.54% and 99.28% of the isolates in the identification dataset

to clusters for assembled genomes and read mapping respectively. The accuracy was 99.70%

and 99.81% for assembled genomes and read mapping respectively when applied to the

validation dataset. Discrepancies were observed between assembled genomes and read

mapping (Table 3). There were more isolates assigned to "*Shigella*/EIEC unclustered" in read

mapping, in contrast there were more isolates assigned to multiple clusters in genome

assemblies. The specificity of ShigEiFinder was 99.40% for assembled genomes and 99.38%

for read mapping when evaluated with 12,743 non-*Shigella*/EIEC *E. coli* isolates. An

additional 2 isolates were detected as sporadic EIEC lineages by read mapping.

Comparison of ShigEiFinder and ShigaTyper

To demonstrate ShigEiFinder for differentiation of *Shigella* from EIEC and enhancement of cluster based serotyping, the comparison of read mapping results between ShigEiFinder and the existing *in silico Shigella* identification pipeline ShigaTyper (10) was performed with 488 isolates used in ShigaTyper and 15,501 isolates from *Shigella*/EIEC validation dataset used in the present study.

The 488 isolates used in ShigaTyper consisted of 23 other species, 45 *E. coli* isolates and 420 *Shigella* isolates. ShigEiFinder identified 23 other species isolates and 453 out of 465 *E. coli* and *Shigella* isolates, in agreement with ShigaTyper assignment. ShigEiFinder also assigned the remaining 3 EIEC isolates and 9 (either multiple *wzx* or no *wzx* genes found) isolates untypeable by ShigaTyper to *Shigella*/EIEC clusters.

ShigEiFinder assigned 15,471 of 15,501 *Shigella*/EIEC isolates to *Shigella* or EIEC clusters and then to a serotype. The accuracy of ShigEiFinder to correctly assign isolates to *Shigella* or EIEC clusters was 99.81% (15,471/15,501). By contrast, ShigaTyper assigned 7,277 isolates (46.95%) to *Shigella*, 7.976 isolates (51.45%) to EIEC, 177 (1.14%) isolates to multiple *wzx* genes and failed to type 71 (0.46%) isolates.

The predicted serotype of 7,277 (46.96%) *Shigella* isolates by ShigaTyper agreed with the results of ShigEiFinder. For 8,224 isolates typed as EIEC or untypable by ShigaTyper, 99.73% (8,202/8,224) of the isolates were assigned to *Shigella* or EIEC clusters by ShigEiFinder (Table 4). Of these isolates, the majority belonged to SS, SD1 and SF which were erroneously predicted as EIEC by ShigaTyper.

Discussion

Shigella and EIEC cause human bacillary dysentery with similar invasion mechanisms, however the pathogenicity of these 2 groups varies (8, 43). The prevalence of each of the four *Shigella* "species" also varies (11-13). Differentiation of *Shigella* and EIEC from each other is important for epidemiologic and diagnostic investigations. However, their similar physiological, biochemical and genetic characteristics make this differentiation difficult.

Determining phylogenetic clusters for better separation of *Shigella* isolates from EIEC

From a phylogenetic perspective, *Shigella* and EIEC strains consisted of multiple phylogenetic

lineages derived from commensal *E. coli*, which do not reflect the nomenclature of *Shigella*

and EIEC (23, 25, 26, 28, 38, 41). In the present study, we identified all phylogenetic clusters

of *Shigella* and EIEC through large scale examination of publicly available genomes.

Phylogenetic results demonstrated that *Shigella* isolates had at least 10 clusters while EIEC

isolates had at least 7 clusters. The 10 *Shigella* clusters included the 7 previously defined

lineages including 3 major clusters (C1, C2 and C3) and 5 outliers (SD1, SD8, SD10, SB13

and SS) (25) and 2 newly identified clusters (SB12 and SB13-atypical). The 7 EIEC clusters

consisted of 4 previously defined EIEC clusters (C4, C5, C6 and C7) (26) and 3 newly

identified EIEC clusters (C8 EIEC O96:H19, C9 EIEC O8:H19 and C10 EIEC O135:H30).

Our WGS-based phylogeny provided high resolution for assigning *Shigella* and EIEC isolates

to clusters. Several serotypes that are currently increasing in frequency (SB19, SB20, SD14,

SD15, SD provisional serotype 96-626) (76-79) were assigned to clusters and five new

clusters/outliers were identified. SB13 isolates in this study formed two known lineages. One

lineage was located outside of *Shigella*/EIEC clusters and represented the outlier SB13 which

is in fact belonging to the newly defined species *E. albertii* (25, 26, 38, 39). The second lineage

was with *E. coli*, and was defined as atypical SB13 previously (39). The newly identified

Shigella outlier SB12 was previously grouped into C3 based on housekeeping gene trees (25,

38) but was seen as outliers in two other studies (28, 56).

Newly identified clusters C8 (EIEC O96:H19) and C9 (EIEC O8:H19) represented the

emergence of novel EIEC serotypes. A recent study revealed that EIEC serotype O96:H19 (C8)

could be the result of a recent acquisition of the invasion plasmid by commensal E. coli (80).

The EIEC serotype O8:H19 (C9) had not been reported previously.

Apart from the 17 major and outlier clusters of *Shigella* and EIEC, the presence of 53 sporadic

EIEC lineages indicated greater genetic diversity than has been observed previously. Isolates

belonging to these sporadic EIEC groups were more closely related to non-enteroinvasive E.

coli isolates than to major *Shigella*/EIEC lineages. However, 41 of these isolates, representing

38 sporadic EIEC lineages, carried pINV. *Shigella* and EIEC both carry the *Shigella* virulence

plasmid pINV which is vital for virulence and distinguishes *Shigella*/EIEC from other *E. coli*

(24, 33, 68). Therefore, these isolates may represent recently formed EIEC lineages through

acquisition of the pINV. The remaining 18 isolates contained the *ipaH* gene but may or may

not carry pINV. It is possible that these strains carried very low copy number of pINV or the

pINV plasmid was lost during culture.

Highly sensitive and specific cluster-specific gene markers for differentiation of *Shigella*

and EIEC isolates

Several studies have identified phylogenetic related genomic markers for discrimination of

Shigella and EIEC and between *Shigella* species (23, 27, 28, 41, 55, 56). However, these

phylogenetic analyses were performed only with a small number of genomes (23, 28, 55). In

addition, non-invasive E. coli isolates were included in some of the phylogenetic clusters

identified (28) which led to non-invasive E. coli isolates being identified by the markers.

We identified cluster-specific gene markers for each respective cluster which were only

composed of *Shigella* or EIEC isolates. Sets of cluster-specific gene markers were identified

for those clusters where no single suitable marker is present. The combination of genes

enhances the specificity of cluster-specific gene markers as demonstrated by the 100%

sensitivity and very high specificity in this analysis (Table 2). Genes specific to each of the 53

sporadic EIEC lineages were also identified and they were sensitive and specific, although it

should be noted that these values are based on very small sample sizes.

The cluster-specific gene markers or marker sets can be used to differentiate *Shigella*/EIEC

from non-enteroinvasive *E. coli* independent of *ipaH* gene. The *ipaH* gene as a molecular

target has been used to differentiate *Shigella* and EIEC from non-enteroinvasive E. coli (24,

43-45). In our study, the cluster-specific gene markers were specific to *Shigella*/EIEC with

98.8% to 100% specificity when evaluated on non-enteroinvasive *E. coli* control database,

providing confidence that the cluster-specific genes or sets are robust markers for the

identification of *Shigella*/EIEC. 53 sporadic EIEC lineage specific gene markers also have

very high specificity (97.02% to 100%) against non-enteroinvasive *E. coli* control database.

The cluster-specific gene markers or marker sets are able to assign *Shigella*/EIEC isolates to

correct clusters in 99.63% of cases and can clearly distinguish *Shigella* isolates from EIEC

when applied to the validation dataset. While ShigaTyper assigned 46.95% isolates to *Shigella*

and 51.45% isolates to EIEC in the same dataset we tested, leading to a large proportion of

isolates incorrectly assigned. The majority of the isolates predicted as EIEC by ShigaTyper

were SS or SD1 as they belonged to SS and SD1 specific STs and were positive to a set of SS

or SD1 specific gene markers and grouped into SS or SD1 cluster on our phylogenetic tree. The genes used in ShigaTyper were SS specific marker Ss_methylase gene (81, 82) together with SS O antigen wzx gene. However, SS specific marker Ss_methylase gene was found in other *Shigella* serotypes and EIEC (10) and SS O antigen wzx gene were located on a plasmid which is frequently lost (83). Similarly, the SD1 O antigen genes used in ShigaTyper were plasmid-borne which may also lead to inconsistent detection (84, 85). A previous study identified 6 loci to distinguish EIEC from *Shigella* (23). We searched the 6 loci against our *Shigella*/EIEC database and found that some *Shigella* isolates were misidentified as EIEC isolates, such as SD8 isolates incorrectly identified as EIEC subtype 13. Our cluster-specific genes can differentiate SD8 isolates from EIEC with 100% accuracy. Therefore, the cluster-specific gene markers marker sets described here provided nearly perfect differentiation of *Shigella* from EIEC.

The cluster-specific gene markers or marker sets are able to differentiate SS and SF (with exception of SF6) from SB and SD. SF and SS are the major cause of *Shigella* infections, accounting for up to 89.6% annual cases (11-13). Differentiation of SS and SF isolates from SB and SD is also beneficial for diagnosis and surveillance. A recent study identified "species" specific markers for the detection of each of the four *Shigella* "species" and validated with only one isolate per species (55). A molecular algorithm based on *Shigella* O antigen genes can detect 85% of SF isolates (52). In contrast, a set of SF specific genes in our study can correctly identify SF isolates with 99.62% accuracy.

The cluster-specific gene markers or marker sets can also assign *Shigella*/EIEC isolates to serotype level if the cluster has single serotype such as SD1, SD8, SD10, SB13, SB12, EIEC O144:H25 (C7), EIEC O96:H19 (C8), EIEC O8:H19 (C9) and EIEC O135:H30 (C10). The remaining EIEC, SF, SB and SD serotypes were distributed over the major clusters C4-6, C3, C1 and C2 respectively. Cluster-specific gene markers or marker sets combined with serotype associated O antigen and modification genes can further identify these isolates to serotype level if the cluster has multiple serotypes. Once an isolate is assigned to a cluster, only serotype associated O antigen and modification genes found in that cluster need be examined. This allows the elimination of ambiguous or incorrect serotype assignments that may otherwise occur, increasing the overall accuracy of the method.

Cluster-specific gene marker based ShigEiFinder can accurately type *Shigella* and EIEC

To facilitate the use of cluster-specific gene markers or marker sets for typing, we developed

an automated pipeline, ShigEiFinder, for *in silico* molecular serotyping of *Shigella*/EIEC.

ShigEiFinder provided *Shigella*/ EIEC differentiation as well as serotype prediction by yielding

"presence or absence" of cluster-specific gene markers or marker sets combined with

Shigella/EIEC O antigen genes and modification genes in a query isolate (either reads or

assembled genomes). We showed 99.70% and 99.81% accuracy to assign isolates to the correct

clusters from 15,501 *Shigella*/EIEC isolates in validation dataset for the assembled genomes

and reads mapping respectively. In contrast, the existing *in silico Shigella* serotyping pipeline

ShigaTyper had 46.95% accuracy for reads mapping when tested with the same validation

dataset, with 51.45% of isolates in validation dataset being predicted as EIEC by ShigaTyper.

The genetic determinants used in ShigaTyper for differentiation of *Shigella* from EIEC and

identification of SS were *lacY*, *cadA*, *Ss_methylase,* SS and SD1 O antigen *wzx* genes (10). As

discussed above some of these genes were found to be non-specific in this study. Compared

with ShigaTyper, the cluster-specific gene markers used in ShigEiFinder for identification of

Shigella and EIEC provided higher discriminatory power than ShigaTyper. ShigEiFinder also

provided a high specificity with 99.40% for assembled genomes and 99.38% for reads

mapping.

ShigEiFinder can differentiate *Shigella* isolates from EIEC and distinguish SS and SF (with

exception of SF6) isolates from SB and SD accurately. It also can identify SD1 isolates

directly. ShigEiFinder was able to serotype over 59 *Shigella* serotypes and 22 EIEC serotypes.

Therefore, ShigEiFinder will be useful for clinical, epidemiological and diagnostic

investigations and the cluster-specific gene markers identified could be adapted for

metagenomics or culture independent typing.

Conclusion

This study analysed over 17,000 publicly available *Shigella*/EIEC genomes and identified 10

clusters of *Shigella*, 7 clusters of EIEC and 53 sporadic types of EIEC. Cluster-specific gene

markers or marker sets for the 17 major clusters and 53 sporadic types were identified and

found to be valuable for *in silico* typing. We additionally developed ShigEiFinder, a freely

available *in silico* serotyping pipeline incorporating the cluster-specific gene markers to

facilitate serotyping of *Shigella*/EIEC isolates using genome sequences with very high

specificity and sensitivity.

Authors and contributors

Conceptualization: R.L, M.P.; Investigation: X.Z., M.P., T.N., S.K.; Methodology: M.P., R.L.

Writing – original draft: X.Z.; Writing – review and editing: M.P., R.L.

Conflicts of interest

The authors declare that there are no conflicts of interest.

Funding information

This work was funded in part by a National Health and Medical Research Council project grant (grant number 1129713) and an Australian Research Council Discovery Grant (DP170101917).

Acknowledgements

The authors thank Duncan Smith and Robin Heron from UNSW Research Technology Services for computing assistance.

Data bibliography

Zhang X, Payne M, Nguyen T, Kaur S, Lan R. All the sequencing data generated within this study, NCBI BioProject number (PRJNA692536).

Abbreviations

SS, *Shigella sonnei*; SF, *Shigella flexneri*; SB, *Shigella boydii*; SD, *Shigella dysenteriae*; EIEC, Enteroinvasive *Escherichia coli*; NCBI SRA, National Center for Biotechnology Information Sequence Read Archive; ST, sequence type; rST, ribosomal ST; MLST, Multilocus sequence typing; rMLST, Ribosomal MLST; ECOR, *Escherichia coli* reference collection; WGS, whole-genome sequencing; TP, true positive; FN, false negative; FP, false positive; HK, House Keeping.

References

1. DuPont HL, Levine MM, Hornick RB, Formal SB. Inoculum size in shigellosis and implications for expected mode of transmission. *The Journal of infectious diseases*. 1989;159(6):1126-8.
2. Troeger C, Forouzanfar M, Rao PC, Khalil I, Brown A, Reiner Jr RC, et al. Estimates of global, regional, and national morbidity, mortality, and aetiologies of diarrhoeal diseases: a systematic

analysis for the Global Burden of Disease Study 2015. *The Lancet Infectious Diseases*. 2017;17(9):909-
48.

3.   World Health Organization. Guidelines for the control of shigellosis, including epidemics due
to *Shigella dysenteriae* type 1. 2005.

4.   Kirk MD, Pires SM, Black RE, Caipo M, Crump JA, Devleesschauwer B, et al. World Health
Organization estimates of the global and regional disease burden of 22 foodborne bacterial, protozoal,
and viral diseases, 2010: a data synthesis. *PLoS medicine*. 2015;12(12):e1001921.

5.   Brengi SP, Sun Q, Bolaños H, Duarte F, Jenkins C, Pichel M, et al. PCR-based method for
Shigella flexneri serotyping: international multicenter validation. *Journal of clinical microbiology*.
2019;57(4):e01592-18.

6.   Kotloff KL, Nataro JP, Blackwelder WC, Nasrin D, Farag TH, Panchalingam S, et al. Burden and
aetiology of diarrhoeal disease in infants and young children in developing countries (the Global
Enteric Multicenter Study, GEMS): a prospective, case-control study. *The Lancet* 2013;382(9888):209-
22.

7.   Khalil IA, Troeger C, Blacker BF, Rao PC, Brown A, Atherly DE, et al. Morbidity and mortality
due to *shigella* and enterotoxigenic *Escherichia coli* diarrhoea: the Global Burden of Disease Study
1990–2016. *The Lancet Infectious Diseases*. 2018;18(11):1229-40.

8.   van den Beld MJ, Warmelink E, Friedrich AW, Reubsaet FA, Schipper M, de Boer RF, et al.
Incidence, clinical implications and impact on public health of infections with *Shigella spp.* and entero-
invasive *Escherichia coli* (EIEC): results of a multicenter cross-sectional study in the Netherlands during
2016–2017. *BMC Infectious Diseases*. 2019;19(1):1037.

9.   Edwards PR, Ewing WH. Identification of enterobacteriaceae. Identification of
Enterobacteriaceae. 1972(Third edition).

10.   Wu Y, Lau HK, Lee T, Lau DK, Payne J. *In silico* serotyping based on Whole-Genome sequencing
improves the accuracy of *shigella* identification. *Appl Environ Microbiol*. 2019;85(7):e00165-19.

11.   The HC, Thanh DP, Holt KE, Thomson NR, Baker S. The genomic signatures of *Shigella*
evolution, adaptation and geographical spread. *Nature Reviews Microbiology*. 2016;14(4):235.

12.   Livio S, Strockbine NA, Panchalingam S, Tennant SM, Barry EM, Marohn ME, et al. *Shigella*
isolates from the global enteric multicenter study inform vaccine development. *Clinical Infectious*
Diseases. 2014;59(7):933-41.

13.   OzFoodNet Working Group. Monitoring the incidence and causes of diseases potentially
transmitted by food in Australia: Annual report of the OzFoodNet network, 2011. *Communicable*
diseases intelligence quarterly report. 2015;39(2):E236.

14.   Connor TR, Barker CR, Baker KS, Weill F-X, Talukder KA, Smith AM, et al. Species-wide whole
genome sequencing reveals historical global spread and recent local persistence in *Shigella flexneri*.
Elife. 2015;4:e07335.

15.   Taylor D, Echeverria P, Sethabutr O, Pitarangsi C, Leksomboon U, Blacklow N, et al. Clinical and
microbiologic features of *Shigella* and enteroinvasive *Escherichia coli* infections detected by DNA
hybridization. *Journal of clinical microbiology*. 1988;26(7):1362-6.

16.   Levine MM. Escherichia coli that cause diarrhea: enterotoxigenic, enteropathogenic,
enteroinvasive, enterohemorrhagic, and enteroadherent. *Journal of infectious Diseases*.
1987;155(3):377-89.

17.   Tai AY, Easton M, Encena J, Rotty J, Valcanis M, Howden BP, et al. A review of the public health
management of shigellosis in Australia in the era of culture‐independent diagnostic testing. *Australian*
New Zealand journal of public health. 2016;40(6):588-91.

18.   Gomes TA, Elias WP, Scaletsky IC, Guth BE, Rodrigues JF, Piazza RM, et al. Diarrheagenic
Escherichia coli. *brazilian journal of microbiology* 2016;47:3-30.

19.   Pasqua M, Michelacci V, Di Martino ML, Tozzoli R, Grossi M, Colonna B, et al. The Intriguing
Evolutionary Journey of Enteroinvasive *E. coli* (EIEC) toward Pathogenicity. *Frontiers in microbiology*.
2017;8:2390-.

20.   Herzig CT, Fleischauer AT, Lackey B, Lee N, Lawson T, Moore ZS, et al. Notes from the Field:
Enteroinvasive *Escherichia coli* Outbreak Associated with a Potluck Party—North Carolina, June–July
2018. *Morbidity and Mortality Weekly Report*. 2019;68(7):183.
21.   Pettengill EA, Hoffmann M, Binet R, Roberts RJ, Payne J, Allard M, et al. Complete genome
sequence of enteroinvasive *Escherichia coli* O96: H19 associated with a severe foodborne outbreak.
Genome Announcements. 2015;3(4):e00883-15.
22.   Escher M, Scavia G, Morabito S, Tozzoli R, Maugliani A, Cantoni S, et al. A severe foodborne
outbreak of diarrhoea linked to a canteen in Italy caused by enteroinvasive *Escherichia coli*, an
uncommon agent. *Epidemiology and infecttion*. 2014;142(12):2559-66.
23.   Dhakal R, Wang Q, Lan R, Howard P, Sintchenko V. Novel multiplex PCR assay for identification
and subtyping of enteroinvasive *Escherichia coli* and differentiation from *Shigella* based on target
genes selected by comparative genomics. *Journal of medical microbiology*. 2018;67(9):1257-64.
24.   Van den Beld M, Reubsaet F. Differentiation between *Shigella*, enteroinvasive *Escherichia coli*
(EIEC) and noninvasive *Escherichia coli*. *European Journal of Clinical Microbiology & Infectious*
Diseases. 2012;31(6):899-904.
25.   Pupo GM, Lan R, Reeves PR. Multiple independent origins of *Shigella* clones of *Escherichia coli*
and convergent evolution of many of their characteristics. *Proceedings of the National Academy of*
Sciences 2000;97(19):10567-72.
26.   Lan R, Alles MC, Donohoe K, Martinez MB, Reeves PR. Molecular evolutionary relationships of
enteroinvasive *Escherichia coli* and *Shigella spp*. *Infection and immunity*. 2004;72(9):5080-8.
27.   Sahl JW, Morris CR, Emberger J, Fraser CM, Ochieng JB, Juma J, et al. Defining the
phylogenomics of *Shigella* species: a pathway to diagnostics. *Journal of clinical microbiology*.
2015;53(3):951-60.
28.   Pettengill EA, Pettengill JB, Binet R. Phylogenetic analyses of *Shigella* and enteroinvasive
Escherichia coli for the identification of molecular epidemiological markers: whole-genome
comparative analysis does not support distinct genera designation. *Frontiers in microbiology*.
2016;6:1573.
29.   Cheasty T, Rowe B. Antigenic relationships between the enteroinvasive *Escherichia coli* O
antigens O28ac, O112ac, O124, O136, O143, O144, O152, and O164 and *Shigella* O antigens. *Journal of*
clinical microbiology. 1983;17(4):681-4.
30.   Landersjö C, Weintraub A, Widmalm G. Structure determination of the O-antigen
polysaccharide from the enteroinvasive *Escherichia coli* (EIEC) O143 by component analysis and NMR
spectroscopy. *Carbohydrate research*. 1996;291:209-16.
31.   Linnerborg M, Weintraub A, Widmalm G. Structural studies of the O‑antigen polysaccharide
from the enteroinvasive *Escherichia coli* O164 cross‑reacting with *Shigella* dysenteriae type 3.
European journal of biochemistry. 1999;266(2):460-6.
32.   Sansonetti P, d'Hauteville H, Écobichon Ct, Pourcel C, editors. Molecular comparison of
virulence plasmids in *Shigella* and enteroinvasive *Escherichia coli*. InAnnales de l'Institut
Pasteur/Microbiologie; 1983 May 1 (Vol. 134, No. 3, pp. 295-318). Elsevier Masson.
33.   Lan R, Lumb B, Ryan D, Reeves PR. Molecular Evolution of Large Virulence Plasmid in *Shigella*
Clones and Enteroinvasive *Escherichia coli*. Infection and immunity. 2001;69(10):6303-9.
34.   Venkatesan MM, Buysse JM, Kopecko DJ. Use of *Shigella flexneri ipaC* and *ipaH* gene
sequences for the general identification of *Shigella spp*. and enteroinvasive *Escherichia coli*. *Journal of*
Clinical Microbiology. 1989;27(12):2687-91.
35.   Hale TL. Genetic basis of virulence in *Shigella* species. *Microbiology and Molecular Biology*
Reviews. 1991;55(2):206-24.
36.   Jin Q, Yuan Z, Xu J, Wang Y, Shen Y, Lu W, et al. Genome sequence of *Shigella flexneri* 2a:
insights into pathogenicity through comparison with genomes of *Escherichia coli* K12 and O157.
Nucleic acids research. 2002;30(20):4432-41.
37.   Yang F, Yang J, Zhang X, Chen L, Jiang Y, Yan Y, et al. Genome dynamics and diversity of
Shigella species, the etiologic agents of bacillary dysentery. *Nucleic acids research*. 2005;33(19):6445-
58.

38. Yang J, Nie H, Chen L, Zhang X, Yang F, Xu X, et al. Revisiting the molecular evolutionary history
of *Shigella spp*. *Journal of molecular evolution*. 2007;64(1):71-9.
39. Hyma KE, Lacher DW, Nelson AM, Bumbaugh AC, Janda JM, Strockbine NA, et al. Evolutionary
genetics of a new pathogenic *Escherichia* species: *Escherichia albertii* and related *Shigella boydii*
strains. *Journal of bacteriology*. 2005;187(2):619-28.
40. Walters LL, Raterman EL, Grys TE, Welch RA. Atypical *Shigella boydii* 13 encodes virulence
factors seen in attaching and effacing *Escherichia coli*. *FEMS Microbiol Lett*. 2012;328(1):20-5.
41. Hazen TH, Leonard SR, Lampel KA, Lacher DW, Maurelli AT, Rasko DA. Investigating the
relatedness of enteroinvasive *Escherichia coli* to other *E. coli* and *Shigella* isolates by using
comparative genomics. *Infection and immunity*. 2016;84(8):2362-71.
42. Silva RM, Toledo M, Trabulsi LR. Biochemical and cultural characteristics of invasive
Escherichia coli. *Journal of Clinical Microbiology*. 1980;11(5):441-4.
43. van den Beld MJ, Friedrich AW, van Zanten E, Reubsaet FA, Kooistra-Smid MA, Rossen JW.
Multicenter evaluation of molecular and culture-dependent diagnostics for *Shigella* species and
Entero-invasive *Escherichia coli* in the Netherlands. *Journal of microbiological methods*. 2016;131:10-5.
44. Van Lint P, De Witte E, Ursi J, Van Herendael B, Van Schaeren J. A screening algorithm for
diagnosing bacterial gastroenteritis by real-time PCR in combination with guided culture. *Diagnostic
microbiology*. 2016;85(2):255-9.
45. De Boer RF, Ott A, Kesztyüs B, Kooistra-Smid AM. Improved detection of five major
gastrointestinal pathogens by use of a molecular screening approach. *Journal of clinical microbiology*.
2010;48(11):4140-6.
46. Liu B, Knirel YA, Feng L, Perepelov AV, Senchenkova SyN, Wang Q, et al. Structure and genetics
of *Shigella* O antigens. *FEMS microbiology reviews*. 2008;32(4):627-53.
47. Wattiau P, Boland C, Bertrand S. Methodologies for *Salmonella enterica ssp enterica*
subtyping: gold standards and alternatives. *Applied and environmental microbiology*. 2011:AEM.
05527-11.
48. Cai H, Lu L, Muckle C, Prescott J, Chen S. Development of a novel protein microarray method
for serotyping *Salmonella enterica* strains. *Journal of clinical microbiology*. 2005;43(7):3427-30.
49. van der Ploeg CA, Rogé AD, Bordagorría XL, de Urquiza MT, Castillo ABC, Bruno SB. Design of
Two Multiplex PCR Assays for Serotyping *Shigella flexneri*. *Foodborne pathogens and disease*.
2018;15(1):33-8.
50. Sun Q, Lan R, Wang Y, Zhao A, Zhang S, Wang J, et al. Development of a multiplex PCR assay
targeting O-antigen modification genes for molecular serotyping of *Shigella flexneri*. *Journal of clinical
microbiology*. 2011;49(11):3766-70.
51. Li Y, Cao B, Liu B, Liu D, Gao Q, Peng X, et al. Molecular detection of all 34 distinct O-antigen
forms of *Shigella*. *Journal of Medical Microbiolgy*. 2009;58(1):69-81.
52. van den Beld MJ, de Boer RF, Reubsaet FA, Rossen JW, Zhou K, Kuiling S, et al. Evaluation of a
Culture-Dependent Algorithm and a Molecular Algorithm for Identification of *Shigella spp*., *Escherichia
coli*, and Enteroinvasive *E. coli*. *Journal of clinical microbiology*. 2018;56(10):e00510-18.
53. Pavlovic M, Luze A, Konrad R, Berger A, Sing A, Busch U, et al. Development of a duplex real‐
time PCR for differentiation between *E. coli* and *Shigella spp*. *Journal of applied microbiology*.
2011;110(5):1245-51.
54. Løbersli I, Wester AL, Kristiansen Å, Brandal LT. Molecular differentiation of *Shigella spp*. from
enteroinvasive *E. coli*. *European Journal of Microbiology*. 2016;6(3):197-205.
55. Kim H-J, Ryu J-O, Song J-Y, Kim H-Y. Multiplex polymerase chain reaction for identification of
shigellae and four *Shigella* species using novel genetic markers screened by comparative genomics.
Foodborne pathogens. 2017;14(7):400-6.
56. Chattaway MA, Schaefer U, Tewolde R, Dallman TJ, Jenkins C. Identification of *Escherichia coli*
and *Shigella* species from whole-genome sequences. *Journal of clinical microbiology*. 2017;55(2):616-
23.
57. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-aware
quantification of transcript expression. *Nature methods*. 2017 Apr;14(4):417-9.

58.   Wood DE, Salzberg SL. Kraken: ultrafast metagenomic sequence classification using exact
alignments. *Genome biology*. 2014;15(3):R46.
59.   Alikhan N-F, Zhou Z, Sergeant MJ, Achtman M. A genomic overview of the population
structure of *Salmonella*. *PLoS genetics*. 2018;14(4):e1007261.
60.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new
genome assembly algorithm and its applications to single-cell sequencing. *Journal of computational
biology*. 2012;19(5):455-77.
61.   Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for genome
assemblies. *Bioinformatics*. 2013;29(8):1072-5.
62.   Jolley KA, Maiden MCJ. BIGSdb: Scalable analysis of bacterial genome variation at the
population level. *BMC Bioinformatics*. 2010;11(1):595.
63.   Joensen KG, Tetzschner AM, Iguchi A, Aarestrup FM, Scheutz F. Rapid and easy *in silico*
serotyping of *Escherichia coli* isolates by use of whole-genome sequencing data. *Journal of clinical
microbiology*. 2015;53(8):2410-26.
64.   Hu D, Liu B, Wang L, Reeves PR. Living Trees: High-Quality Reproducible and Reusable
Construction of Bacterial Phylogenetic Trees. *Molecular Biology and Evolution*. 2019;37(2):563-75.
65.   Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new developments.
Nucleic acids research. 2019;47(W1):W256-W9.
66.   Zhou Z, Alikhan N-F, Sergeant MJ, Luhmann N, Vaz C, Francisco AP, et al. GrapeTree:
visualization of core genomic relationships among 100,000 bacterial pathogens. *Genome research*.
2018;28(9):1395-404.
67.   Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv
preprint arXiv*. 2013.
68.   Buchrieser C, Glaser P, Rusniok C, Nedjari H, d'Hauteville H, Kunst F, et al. The virulence
plasmid pWR100 and the repertoire of proteins secreted by the type III secretion apparatus of *Shigella
flexneri*. *Molecular microbiology*. 2000;38(4):760-71.
69.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map
format and SAMtools. *Bioinformatics* 2009;25(16):2078-9.
70.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics. 2010;26(6):841-2.
71.   Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics*. 2014;30(14):2068-
9.
72.   Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MT, et al. Roary: rapid large-scale
prokaryote pan genome analysis. *Bioinformatics*. 2015;31(22):3691-3.
73.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:
architecture and applications. *BMC bioinformatics*. 2009;10(1):421.
74.   Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, et al. The RAST Server: rapid
annotations using subsystems technology. *BMC genomics*. 2008;9(1):75.
75.   Sof'ya NS, Feng L, Yang J, Shashkov AS, Cheng J, Liu D, et al. Structural and genetic
characterization of the *Shigella boydi*i type 10 and type 6 O antigens. *Journal of bacteriology*.
2005;187(7):2551-4.
76.   Ansaruzzaman M, Kibriya A, Rahman A, Neogi P, Faruque A, Rowe B, et al. Detection of
provisional serovars of *Shigella* dysenteriae and designation as *S*. dysenteriae serotypes 14 and 15.
Journal of clinical microbiology. 1995;33(5):1423-5.
77.   Balows A. Manual of clinical microbiology 8th edition: PR Murray, EJ Baron, JH Jorgenson, MA
Pfaller, and RH Yolken, eds., ASM Press, 2003, 2113 pages, 2 vol, 2003+ subject & author indices, ISBN:
1-555810255-4, US $189.95. *Diagnostic Microbiology*. 2003;47(4):625.
78.   Woodward DL, Clark CG, Caldeira RA, Ahmed R, Soule G, Bryden L, et al. Identification and
characterization of *Shigella boydii* 20 serovar nov., a new and emerging *Shigella* serotype. *Journal of
Medical Microbiolgy*. 2005;54(8):741-8.

79.   Kim J, Lindsey RL, Garcia-Toledo L, Loparev VN, Rowe LA, Batra D, et al. High-quality whole-
genome sequences for 59 historical *Shigella* strains generated with PacBio sequencing. *Genome*
Announcements. 2018;6(15).
80.   Michelacci V, Prosseda G, Maugliani A, Tozzoli R, Sanchez S, Herrera-León S, et al.
Characterization of an emergent clone of enteroinvasive *Escherichia coli* circulating in Europe. *Clinical*
Microbiology. 2016;22(3):287. e11-. e19.
81.   Liu J, Platts-Mills JA, Juma J, Kabir F, Nkeze J, Okoi C, et al. Use of quantitative molecular
diagnostic methods to identify causes of diarrhoea in children: a reanalysis of the GEMS case-control
study. *The Lancet*. 2016 Sep 24;388(10051):1291-301.
82.   Cho MS, Ahn T-Y, Joh K, Kwon O-S, Jheong W-H, Park DS. A novel marker for the species-
specific detection and quantitation of *Shigella sonnei* by targeting a methylase gene. *Journal of*
microbiolgy and biotechnology. 2012;22(8):1113-7.
83.   Sansonetti PJ, Kopecko DJ, Formal SB. *Shigella sonnei* plasmids: evidence that a large plasmid
is necessary for virulence. *Infection and immunity*. 1981;34(1):75-83.
84.   Feng L, Perepelov AV, Zhao G, Shevelev SD, Wang Q, Sof'ya NS, et al. Structural and genetic
evidence that the *Escherichia coli* O148 O antigen is the precursor of the *Shigella dysenteriae* type 1 O
antigen and identification of a glucosyltransferase gene. *Microbiology*. 2007;153(1):139-47.
85.   Göhmann S, Manning P, Alpert C-A, Walker M, Timmis K. Lipopolysaccharide O-antigen
biosynthesis in *Shigella* dysenteriae serotype 1: analysis of the plasmid-carried rfp determinant.
Microbial pathogenesis. 1994;16(1):53-64.
925
926

 Table 1: The summary of identified *Shigella*/EIEC clusters and outliers in identification dataset

Clusters (no of serotypes)[#]	No of isolates	No. STs	No. rSTs	Serotypes
C1 (25)	288	36	166	SB1-4, SB6, SB8, SB10, SB14, SB18, SB11[b], SB19-20[b]; SD3-7, SD9, SD11-13, SD14-15[*], SD-96-265[*]; SF6
C2 (9)	101	19	56	SB5, SB7, SB9, SB11, SB15, SB16, SB17; SD2, SD-E670-74[b]; SD2
C3 (20)	744	81	437	SF1a, SF1b, SF1c (7a), SF2a, SF2b, SF3a, SF3b, SF4a, SF4av, SF4b, SF4bv, SF5a, SF5b, SF7b, SFX, SFXv (4c), SFY, SFYv, SF novel serotype; SB-E1621-54*
C4 (9)	51	6	21	O28ac:H-, O28ac:H7, O136:H7, O164:H-, O164:H7, O29:H4, O173:H7, O124:H7, O132:H7[*]
C5 (6)	62	4	15	O121:H30, O124:H30, O164:H30, O132:H21, O152:H30, O152:H-
C6 (3)	20	2	6	O143:H26, O167:H26, O112ac:H26[b]
C7	10	1	3	O144:H25
C8[a]	12	2	1	O96:H19
C9[a]	4	1	2	O8:H19
C10 [#]	2	1	1	O135:H30
CSS	427	39	294	
CSD1	70	8	56	SD1
CSD8	7	3	3	SD8
CSD10	2	2	1	SD10
CSB12[a]	8	2	6	SB12
CSB13	7	3	3	SB13

Clusters (no of serotypes)[#]	No of isolates	No. STs	No. rSTs	Serotypes
CSB13-atypical[a]	5	3	3	SB13
Sporadic EIEC lineages[a] (53)	59	49	53	53 antigen types

929   [#]Numbers in parentheses are the number of serotypes within that cluster.

930   [a]: Clusters identified as new clusters in this study.

931   [b]: Serotypes were inconsistent with previous analyses.

934   **Table 2: The sensitivity and specificity of cluster-specific genes**

Clusters	Cluster-specific genes (Single/sets)[b]	Identification dataset (1969 isolates)		
		No of isolates	Sensitivity	Specificity
C1	Set of 4 genes	288	100	99.94[a]
C2	Set of 3 genes	101	100	100
C3	Set of 3 genes	744	100	99.59[a]
C4	Set of 2 genes	51	100	100
C5	Set of 3 genes	62	100	100
C6	Set of 2 genes	20	100	100
C7	Single gene	10	100	100
C8	Set of 2 genes	12	100	100
C9	Set of 2 genes	4	100	100
C10	Single gene	2	100	100
CSS	Set of 5 genes	427	100	99.87[a]
CSD1	Set of 2 genes	70	100	100
CSD8	Single gene	7	100	100
CSD10	Single gene	2	100	100
CSB12	Single gene	8	100	100
CSB13	Single gene	7	100	100
CSB13-atypical	Single gene	5	100	100
53 Sporadic EIEC lineages	Single gene / lineage	59	100	100

936   [a]:The specificity of cluster-specific gene set less than 100% was due to at least one FP found in

937   that set.

938   [b]: The sequences of these genes were listed in Data S1.

Table 3: The accuracy of ShigEiFinder with identification dataset and validation dataset

ShigEiFinder assignments	Identification Dataset (n=1,969)[a]		Validation dataset (n=15,501)	
	Genomes	**Reads mapping**	**Genomes**	**Reads mapping**
Shigella/EIEC clusters	1871	1848	15,455	15,471
Multiple *Shigella*/EIEC clusters	9	6	33	7
Shigella/EIEC unclustered	0	8	13	23
Not *Shigella*/EIEC	89	89	0	0
Accuracy[b]	99.54%	99.28%	99.70%	99.81%

[a]: Identification dataset has 90 non-*Shigella*/EIEC strains including 72 ECOR strains and 18 *E.albertii* strains. 1,969 assembled genomes and 1,951 reads (reads not available for 18 EIEC isolates downloaded from NCBI) in identification dataset. One of *E.albertii* strain was assigned as SB13 which was grouped into SB13 cluster on the phylogenetic tree.

[b]: The accuracy was defined as the number of *Shigella*/EIEC isolates being correctly assigned to cluster over the total number of tested.

Table 4: Discrepant assignment of 8,224 isolates by ShigEiFinder and Shigatyper

ShigEiFinder Assignment	ShigaTyper assignment			Total
	EIEC	**Multiple wzx**	**Non-prediction**	
SS	7,465	12	7	7,484
SF	117	61	10	188
C1 and C2 (SB/SD)	17	99	51	167
SB12	0	2	0	2
SD1	244	1	1	246
SD8	1	0	0	1
SD10	0	0	2	2
EIEC	97	0	0	97
Sporadic EIEC lineages	15	0	0	15
Multiple clusters	5	2	0	7
Shigella/EIEC unclustered	15	0	0	15
Total	7,976	177	71	8,224

Figure legends:

Figure 1: *Shigella*/EIEC cluster Identification phylogenetic tree

Representative isolates from the identification dataset were used to construct the phylogenetic

tree by Quicktree v1.3 (64) to identify *Shigella* and EIEC clusters and visualised by

Grapetree's interactive mode. The dendrogram tree shows the phylogenetic relationships of

1879 *Shigella* and EIEC isolates represented in the identification dataset. Branch lengths are

log scale for clarity. The tree scales indicated the 0.2 substitutions per locus. *Shigella* and EIEC

clusters are coloured. Numbers in square brackets indicate the number of isolates of each

identified cluster. CSP is sporadic EIEC lineages.

Figure 2: *in silico* serotyping pipeline workflow

Schematic of *in silico* serotyping *Shigella* and EIEC by cluster-specific genes combined with

the *ipaH* gene and O antigen and modification genes and H antigen genes, implemented in

ShigEiFinder. Both assembled genomes and raw reads are accepted as data input.

Supplementary Material

Figure S1: **Identification phylogenetic tree**

An identification phylogenetic tree constructed by Quicktree v1.3 (64) and visualised by ITOL

v5 shows the phylogenetic relationships of 1879 *Shigella* and EIEC isolates in identification

dataset. The tree scales indicated the 0.01 substitutions per locus. *Shigella* and EIEC clusters

are colored. The internal branches are colored to represent the bootstrap values. Green color

indicates the maximum bootstrap value (1). The red color shows the minimum bootstrap value

(0). Each of cluster is well supported by bootstrap value. CSP is sporadic lineages.

Figure S2-A: **Confirmation phylogenetic tree**

A confirmation phylogenetic tree was constructed by Quicktree v1.3 (64) based on 2375

isolates and visualised by Grapetree's interactive mode. The tree shows the phylogenetic

relationships between identified *Shigella*/EIEC clusters in identification dataset and non-

enteroinvasive *E.coli* isolates. Branch lengths are log scale for clarity. The tree scales indicated

the 0.1 substitutions per locus. Known *Shigella* and EIEC clusters from identification dataset

are colored. Numbers in square brackets indicate the number of isolates of each identified

cluster. CSP is sporadic lineages.

Figure S2-B: **Confirmation phylogenetic tree**

A confirmation phylogenetic tree constructed by Quicktree v1.3 (64) and visualised by ITOL

v5 shows the phylogenetic relationships between identified *Shigella*/EIEC clusters in

identification dataset and non-enteroinvasive *E.coli* isolates. The tree scales indicated the 0.01

substitutions per locus. *Shigella* and EIEC clusters are colored. The internal branches are

colored to represent the bootstrap values. Green color indicates the maximum bootstrap value

(1). The red color shows the minimum bootstrap value (0). Each of cluster is well supported by

bootstrap value. CSP is sporadic lineages.

Figure S3: **Distribution of mapped 38 virulence genes in 59 sporadic isolates**

The presence of *Shigella* virulence plasmid pINV in 59 sporadic isolates in identification

dataset was determined by the mapped 38 virulence genes. Detailed genes were described in

Results "**Investigation of *Shigella* virulence plasmid pINV in 59 sporadic isolates**". Three

categories were defined based on the number of virulence genes mapped to isolate. Virulence

plasmid positive: > 25 genes mapped to isolate; Intermediate: 13 to 25 genes mapped to isolate;

Virulence plasmid negative: less than 13 genes mapped to isolate.

Figure S4 (A): **Validation phylogenetic tree**

A validation tree was generated by Quicktree v1.3 (64) and visualised by Grapetree's

interactive mode to assign representative isolates in validation dataset to clusters. Branch

lengths are log scale for clarity. The tree scales indicated the 0.2 substitutions per locus.

Known *Shigella* and EIEC clusters from identification dataset are colored. Numbers in square

brackets indicate the number of isolates of each identified cluster. Isolates in validation dataset

are colored white. The isolates are assigned to clusters if they grouped into known cluster

isolates. CSP is sporadic lineages.

Figure S4 (B): **Validation phylogenetic tree**

A validation phylogenetic tree was constructed by Quicktree v1.3 (64) and visualised by ITOL

v5 to assign representative isolates in validation dataset to clusters. The tree scales indicated

the 0.01 substitutions per locus. *Shigella* and EIEC clusters are colored. The internal branches

are colored to represent the bootstrap values. Green color indicates the maximum bootstrap

value (1). The red color shows the minimum bootstrap value (0). Each of cluster is well

supported by bootstrap value. Isolates that grouped with known cluster isolates (from

identification dataset) with strong bootstrap support are categorised into that cluster. CSP is

sporadic lineages.

Table S1: 1,969 isolates used in identification dataset

Table S2: 15,501 isolates used in validation dataset

Table S3: The location and function of cluster-specific genes

1063 **Table S4**: The results of cluster-specific gene markers tested with 12,743 non-enteroinvasive

1064 *E.coli* isolates

1065

1066 **Data S1**: Algorithms incorporated into the ShigEiFinder

1067 **Data S2**: Genetic signature O and H genes from ShigaTyper and SerotypeFinder

1068

1069 **Data Availability Statement**

1070

1071 Custom python scripts used in this study are available from the authors on request.

1072

0.2