7-Dehydrocholesterol-derived oxysterols cause neurogenic defects in Smith-Lemli-Opitz syndrome

Hideaki Tomita [a], Kelly M. Hines [a,b], Josi M. Herron [a], Amy Li [a], David W. Baggett [a,c], Libin Xu [a,*]

[a] Department of Medicinal Chemistry, University of Washington, Seattle, WA, 98115, USA

[b] Current address: Department of Chemistry, University of Georgia, Athens, GA, 30602, USA

[c] Current address: Department of Structural Biology, St. Jude Children's Research Hospital, Memphis, TN, 38105, USA

*, Correspondence: libinxu@uw.edu

Abstract

Defective 3β-hydroxysterol-Δ^7-reductase (DHCR7) in the developmental disorder, Smith-Lemli-Opitz syndrome (SLOS), results in deficiency in cholesterol and accumulation of its precursor, 7-dehydrocholesterol (7-DHC). Here, we show that loss of *DHCR7* causes accumulation of 7-DHC-derived oxysterol metabolites, premature neurogenesis, and perturbation of neuronal localization in developing murine or human cortical neural precursors, both *in vitro* and *in vivo*. We found that a major oxysterol, 3β,5α-dihydroxycholest-7-en-6-one (DHCEO), mediates these effects by initiating crosstalk between glucocorticoid receptor (GR) and neurotrophin receptor kinase TrkB. Either loss of *DHCR7* or direct exposure to DHCEO causes hyperactivation of GR and TrkB and their downstream MEK-ERK-C/EBP signaling pathway in cortical neural precursors. Moreover, direct inhibition of GR activation with an antagonist or inhibition of DHCEO accumulation with antioxidants rescues the premature neurogenesis phenotype caused by the loss of *DHCR7*. These results suggest that GR could be a new therapeutic target against the neurological defects observed in SLOS.

Keywords: Smith-Lemli-Opitz syndrome, cholesterol, 7-dehydrocholesterol, oxysterol, neurogenesis, neural progenitor, glucocorticoid receptor, TrkB, cortical development

Introduction

Brain is rich in cholesterol, contributing to 25% of total cholesterol in human body, and nearly all cholesterol in the brain is synthesized locally (Dietschy and Turley, 2004). Therefore, dysregulation of cholesterol metabolism in CNS can potentially cause significant defects in CNS development and functions. Indeed, enzymatic deficiencies in cholesterol biosynthesis cause a number of inherited diseases with severe neurodevelopmental phenotypes (Porter and Herman, 2011).

Smith-Lemli Opitz syndrome (SLOS) is an autosomal recessive, neurological and developmental disorder characterized by multiple developmental defects, such as distinctive facial features, cleft palate, microcephaly and holoprosencephaly, as well as severe intellectual impairment and behavioral problems (Porter and Herman, 2011; Thurm et al., 2016). Notably, SLOS patients display high incidence (> 50%) of autism spectrum disorders (Bukelis et al., 2007; Sikora et al., 2006; Tierney et al., 2006). SLOS is caused by mutations in the 3β-hydroxysterol-Δ^7-reductase gene (*DHCR7*), which encodes the enzyme that converts 7-dehydrocholesterol (7-DHC) to cholesterol in the final step of the cholesterol biosynthesis pathway (Fitzky et al., 1998; Tint et al., 1994; Wassif et al., 1998). Defective DHCR7 resulting from the mutations leads to deficiency in cholesterol and accumulation of 7-DHC in tissues and fluids of affected individuals (Tint et al., 1994; Tint et al., 1995). 7-DHC was found to be highly reactive toward free radical oxidation, leading to the formation of its oxidative metabolites, *i.e.*, oxysterols (Xu et al., 2009; Xu et al., 2010; Xu et al., 2013; Xu et al., 2011). 7-DHC-derived oxysterols can exert cytotoxicity in neuronal cells, induce gene expression changes, and increase formation of dendritic arborization from cortical neurons (Korade et al., 2010; Xu et al., 2012). Interestingly, increased dendrite and axon formation has also been observed in neurons isolated

from *Dhcr7*-KO mouse brain (Jiang et al., 2010). These studies suggest that 7-DHC-derived

oxysterols may be underlying the alterations in neuronal processes in SLOS.

Cholesterol, 7-DHC, and oxysterols derived from both have been found to play important

roles in modulating signaling pathways in developing tissues and organs, such as Hedgehog (Hh)

and Wnt signaling pathways (Byrne et al., 2016; Corcoran and Scott, 2006; Huang et al., 2018;

Myers et al., 2017; Porter et al., 1996; Raleigh et al., 2018). Related to SLOS, ring-B oxysterols

derived from 7-DHC oxidation inhibit Smo in the Hh pathway (Sever et al., 2016). On the other

hand, cholesterol was recently found to selectively activate canonical Wnt signaling over non-

canonical Wnt signaling (Sheng et al., 2014). However, in human induced pluripotent stem cells

(hiPSCs) derived from SLOS patient fibroblasts, accumulation of 7-DHC was found to inhibit

Wnt/β-catenin pathway, which contributes to the precocious neuronal specification in SLOS

neural progenitors (Francis et al., 2016). Depending on the position of the oxidation, oxysterols

have also been shown to bind and activate other signaling molecules and nuclear receptors,

including estrogen receptors, liver X receptors, and glucocorticoid receptor (GR), and thus, play

important roles in neurodevelopment and diseases (DuSell et al., 2008; Theofilopoulos et al.,

2013; Voisin et al., 2017). However, it is unknown how alteration of sterol composition

influences neural stem cell/progenitor behaviors during cortical development, and it remains

elusive whether neural defects of SLOS is due to deficiency in cholesterol or accumulation of 7-

DHC or its oxysterols.

Here we examine the effects of *DHCR7* mutations in developing neural precursors,

focusing on the cerebral cortex. We demonstrate that 7-DHC-derived oxysterols start to

accumulate at embryonic day 12.5 (E12.5) and continue to increase at E14.5 and E16.5, and that

loss of *DHCR7* causes decreased proliferation and self-renewal of mouse cortical neural

precursors and aberrant premature neurogenesis in both mouse and human neural progenitor cells

(NPCs). We then provide evidence that a 7-DHC-derived oxysterol, 3β,5α-dihydroxycholest-7-

en-6-one (DHCEO), activates GR and the downstream receptor tyrosine kinases (RTKs)-

mediated neurogenic signaling through TrkB, and in doing so, promotes premature NPC

differentiation and perturbs neuronal positioning. Either inhibition of GR activation with an

antagonist or inhibition of DHCEO accumulation with antioxidants rescues the premature

neurogenesis defect.

Results

Dhcr7 is Expressed in Embryonic Cortical Precursors and Neurons of Developing Murine
Cortex

To understand the role of *Dhcr7* in neural development, we studied murine developing

cortical precursors during embryonic neurogenesis. We first analyzed the expression of *Dhcr7*

mRNA during mouse cortical development. RT-PCR analysis showed expression of *Dhcr7* in

embryonic day 11 (E11) through postnatal day 0 (P0) cortex (Figure 1A). Quantitative PCR

(qPCR) also suggested the stable expression of *Dhcr7* from E11 to E18 (Figure 1B). Western

blot analysis further confirmed the presence of Dhcr7 protein in developing cortex (Figure 1C).

Immunostaining showed that Dhcr7 was broadly expressed in the developing cortex and was

detectable in precursors in the E12 ventricular and subventricular (VZ/SVZ) zones (Figure 1D).

Immunostaining of cortical sections of *Dhcr7*-KO mouse brain at E12 with this particular Dhcr7

antibody confirmed the antibody specificity against Dhcr7. To further establish the expression of

Dhcr7 in neural lineage cells, we analyzed cortical precursor cultures from E12.5, which consists

of proliferating radial glial cells that generate neurons in culture (Figure 1E). In the E12.5

cortical precursor cultures, Dhcr7+ cells were co-labeled with the precursor markers, Sox2 and

Pax6. Dhcr7 was also co-labeled with the neuronal marker, βIII-tubulin, and the neural lineage

marker, Nestin. The results suggested that Dhcr7 was consistently expressed in neural lineage

cells, which is consistent with our *in vivo* analysis.

121
122

Figure 1. Loss of *Dhcr7* alleles causes decreased proliferation and increased neurogenesis in murine cortical precursors. (A) RT-PCR for *Dhcr7* mRNA in the E11.5 to P0 cortex. *β-actin* mRNA was used as loading control. (B) qRT-PCR for *Dhcr7* mRNA in the E11.5 to E18.5 cortex. Data is expressed as fold change over E11.5 cortex. (C) Western blot of Dhcr7 in total cortical lysates from E13.5 to P0. The blot was re-probed for Erk1/2 as a loading control. (D) Images of E13.5 *Dhcr7$^{+/+}$* (*Dhcr7*-WT, left panel) and *Dhcr7$^{-/-}$* (*Dhcr7*-KO, right panel) cortical sections immunostained for Dhcr7 (red). The subventricular/ventricular zone (SVZ/VZ) is denoted. The right panels show image of boxed area. Scale bar =100 μm. (E) Images of cultured mouse cortical precursors immunostained for Dhcr7 (red) and Sox2, Pax6, βIII-tubulin and Nestin (green) and counterstained with DAPI (blue). Scale bar = 50 μm. (F-J) E12.5 cortical precursors from single *Dhcr$^{+/+}$* and *Dhcr7$^{-/-}$* embryos were cultured 3 days and analyzed. (F) Cells were immunostained for Ki67 (green) and βIII-tubulin (red) after 3 days and quantified for the proportions of Ki67+ (G), Pax6+ (H), βIII-tubulin+ (I) and CC3+ cells (J). Scale bar = 50 μm. *, $p < 0.001$; n = 3 embryos per genotype. (K-O) E13.5 cortical precursor cells from single *Dhcr7$^{+/+}$* or *Dhcr7$^{-/-}$* embryos were cultured as primary neurosphere (K) and the number and diameter of primary neurospheres were quantified 6 days later (L, M). Equal numbers of primary neurospheres were then passaged, and the number and diameter of secondary neurospheres were quantified 6 days later (N, O). Representative images of *Dhcr$^{+/+}$* and *Dhcr7$^{-/-}$* neurospheres are shown as inserts in the left corner.   *, $p < 0.001$; n = 3 embryos per genotype. Scale Bar = 100 μm.

Figure 1-Figure Supplement 1. Characterization of the pluripotency of SLOS-derived human iPSCs.

Figure 1-Figure Supplement 2. Loss of *DHCR7* alleles causes decreased proliferation and increased neurogenesis in human cortical precursors.

Loss of Dhcr7 Alleles Causes Decreased Proliferation and Increased Neurogenesis in Murine and Human Cortical Precursors

SLOS is characterized by neurodevelopmental defects, such as microcephaly. To ask if

Dhcr7 plays important roles in neural precursor (progenitor) development as seen in human

patients with SLOS, we intercrossed *Dhcr7$^{+/-}$* mice and prepared single embryo cultures from

E12.5 *Dhcr7$^{-/-}$* (knockout or KO) or *Dhcr7$^{+/+}$* embryos. Cultures were immunostained for Ki67

and βIII-tubulin 3 days after plating, which revealed that loss of *Dhcr7* caused significant

decrease in the proportion of Ki67+ and Pax6+ precursors whereas increased the proportion of

βIII-tubulin+ neurons (Figure 1F-I). Immunostaining for cleaved caspase 3 (CC3) showed that

the loss of *Dhcr7* alleles did not affect the survival of cortical precursor cells in culture (Figure

1J). These results suggested that loss of *Dhcr7* leads to premature neurogenesis and decreased

proliferation of cortical precursors.

To determine if Dhcr7 is important for the proliferation and self-renewal of cortical

precursors, we performed neurosphere assays, which assess if sphere-forming precursors can

self-renew and generate new spheres. E13.5 cortical precursors from *Dhcr7*$^{-/-}$ and *Dhcr7*$^{+/+}$

embryos were cultured in the presence of FGF2 and EGF. The number and diameter of spheres

were measured 7 days post-plating (Figure 1K-O). Significantly fewer number of and smaller

neurospheres were generated from *Dhcr7*$^{-/-}$ embryonic cortices comparing to its wild-type

littermates (Figure 1L-M). These results were consistent with the reduced proportion of Ki67+

precursors in the adherent cultures. Following formation of the primary neurospheres, these

spheres were triturated and re-plated at equal density to form the secondary neurospheres. The

results showed that there was an approximately 2-fold decrease in the number and diameter of

secondary spheres from *Dhcr7*$^{-/-}$ cortical precursors relative to *Dhcr7*$^{+/+}$ (Figure 1N-O). Thus,

the loss of *Dhcr7* alleles disrupts proliferation and self-renewal of cortical precursors.

We then asked whether *DHCR7* was also necessary for neurogenesis of human

embryonic stem cell-derived NPCs. To examine function of DHCR7, we generated human

induced pluripotent stem cells (hiPSCs) from two lines of SLOS patient fibroblasts and one line

of wild-type fibroblasts and verified their pluripotency and stemness as described in Methods

(Figure 1-Figure Supplement 1) (Okita et al., 2011; Yu et al., 2007). The SLOS hiPSCs were

then differentiated into hNPCs as described previously. Immunoreactivity of DHCR7 was

detected in almost all wild-type hNPCs in these cultures with Sox2+ and Nestin+ cells (Figure 1-

Figure Supplement 2; Panel A). To examine neural differentiation, the SLOS and wild-type

NPCs were differentiated in cultures and immunostained 4 days later for Ki67 and βIII-tubulin

for proliferating precursors and newborn neurons, respectively (Figure 1-Figure Supplement 2;

Panels B-D). Notably, the proportion of Ki67+ precursors was significantly decreased whereas

the proportion of βIII-tubulin+ neurons was increased in SLOS NPC cultures, similar to the

phenotype seen in murine cortical precursors from $Dhcr7^{-/-}$ embryos. Taken together, DHCR7 is

involved in proliferation and differentiation of human NPCs.

Cholesterol precursor 7-DHC and 7-DHC-derived Oxysterols are accumulated in Dhcr7$^{-/-}$
developing cortex and human NPCs
190
7-DHC is highly susceptible to free radical oxidation (Xu et al., 2009), which leads to

formation of numerous oxysterols in cells and tissues (Xu et al., 2013; Xu et al., 2011) (Figure

2A). Liquid chromatography-tandem mass spectrometry (LC-MS/MS) was performed on

cortices from $Dhcr7^{-/-}$ and $Dhcr7^{+/+}$ embryos during cortical development. These analyses

revealed significant accumulation of 7-DHC and reduction of cholesterol in the embryonic

cortices from $Dhcr7^{-/-}$ embryos throughout cortical development from E12.5 to E16.5 (Figure

2B). Increased levels of 7-dehydrodesmosterol (7-DHD), the precursor to desmosterol via

Dhcr7, and 8-dehydrocholesterol (8-DHC) and decreased level of desmosterol in $Dhcr7^{-/-}$

cortices are also consistent with the loss of the function of Dhcr7. Furthermore, 7-DHC

oxysterols, such as DHCEO, 4α-hydroxy-7-DHC (4α-OH-7DHC), and 4β-hydroxy-7-DHC (4β-

OH-7DHC) (Xu et al., 2013; Xu et al., 2011) showed substantial accumulation in the $Dhcr7^{-/-}$

cortices, reaching up to 4-8 ng/mg of tissue, throughout the development (Figure 2A and 2C).

Note that 4 ng/mg of tissue would translate into approximately 10 μM assuming the density of

the brain is 1 g/mL. On the other hand, cholesterol-derived oxysterols, such as 24-keto-Chol,

24,25-epoxy-Chol, 24- or 25-OH-Chol, 7-OH-Chol, and 7-keto-Chol decreased or did not

change significantly in $Dhcr7^{-/-}$ cortices relative to $Dhcr7^{+/+}$ cortices.

Furthermore, significant accumulation of 7-DHC and 7-DHC-derived oxysterols and

reduction of cholesterol were also found in both SLOS patient-derived hiPSCs and NPCs relative

to wild type (Figure 2-Figure Supplement 1). Taken together, *DHCR7* mutations led to

accumulation of 7-DHC and 7-DHC-derived oxysterols in murine and human cortical precursors.

211
Figure 2. Cholesterol precursor 7-DHC and 7-DHC-derived Oxysterols are accumulated in
***Dhcr7*^{-/-} mouse embryonic cortex.** (A) Chemical structures of 7-DHC-derived oxysterols. LC-
MS/MS analysis of (B) cholesterol and its precursors and (C) 7-DHC-derived oxysterols in
Dhcr7^{+/+} and *Dhcr7*^{-/-} embryonic cortex during development. Error bars indicate standard
deviation. *, $p < 0.05$; **, $p < 0.005$; ***, $p < 0.001$; n = 3 biological replicates per group.

Figure 2-Figure Supplement 1. Cholesterol precursor 7-DHC and 7-DHC-derived Oxysterols
are accumulated *in* SLOS-derived human iPSCs and NPCs.
219

220 ***Knockdown (KD) of Dhcr7 Causes Increased Neurogenesis and Depletion of Cycling***
221 ***Precursors in Murine and Human NPC cultures***
222
To examine a potential role of *Dhcr7*, we generated three *Dhcr7* short hairpin RNAs

(shRNAs)-EGFP reporters and transfected them into 293T cells along with murine *Dhcr7*

cDNA-expressing plasmids. We found that *Dhcr7* shRNA2 was the most effective among those

shRNAs and was chosen to examine *Dhcr7* function during neurogenesis from E12.5 cortical

precursors in culture (Figure 3A). When cortical precursors were transfected with *Dhcr7*

shRNA2-EGFP, the shRNA significantly decreased the percentage of transfected cells

expressing *Dhcr7* (Figure 3B-C). *Dhcr7* KD led to a significant decrease in proliferation of

cortical precursors as measured by Ki67 immunostaining and significant increase of βIII-

tubulin+ neurons 3 days post-transfection, but it did not affect cell survival as examined by

immunostaining for CC3 (Figure 3D-G). The significant decrease of precursor proliferation was

caused by a decrease in Pax6+ radial precursors (Figure 3H).

Dhcr7 KD also decreased self-renewal of radial precursor as demonstrated by clonal

analysis with piggybac (PB) transposon, which permanently labels precursors and their progeny.

Cortical precursors were transfected with PB transposase and PB-*Dhcr7* shRNA2-EGFP or

control shRNA-EGFP, and cultures were immunostained 3 days post-transfection for EGFP, the

precursor markers, Sox2, and βIII-tubulin (Figure 3I-L). KD of *Dhcr7* reduced EGFP+

multicellular clones (Figure 3J) whereas neuron-only (βIII-tubulin+) clones were increased

(Figure 3K). Furthermore, the number of precursors in clones containing at least one Sox2+

cells are decreased (Figure 3L).

To ensure that these changes were *Dhcr7* shRNA-dependent, we performed a rescue

experiments using human *DHCR7* cDNA that is resistant to the murine *Dhcr7* shRNA (Figure 3-

Figure Supplement 1). The murine shRNA did not affect the expression of human DHCR7

cDNA as confirmed by co-transfection of the murine shRNA and human *DHCR7* cDNA in 293T

cells and western blot analysis 2 days later (Figure 3-Figure Supplement 1; Panel A).

Furthermore, precursors were co-transfected with murine *Dhcr7* shRNA-EGFP or control

shRNA-EGFP +/- human *DHCR7* cDNA and immunostained 3 days later for Ki67 or βIII-

tubulin (Figure 3-Figure Supplement 1). The human *DHCR7* cDNA showed significant rescue

of the murine *Dhcr7* shRNA KD phenotypes, confirming the specificity of the shRNA.

251
Figure 3. Knockdown of *Dhcr7* causes increased neurogenesis and depletion of cycling
precursors in murine and human NPC cultures. (A) Western blot for Dhcr7 in 293T cells
transfected with control or individual murine *Dhcr7* shRNAs. The blot was re-probed for Erk1/2
as a loading control. (B and C) Mouse cortical precursors were transfected EGFP-*Dhcr7*
shRNAs (sh2) or EGFP-control (Con) and immunostained for EGFP and Dhcr7 (red) 2 days later

and EGFP+ cells expressing detectable Dhcr7 were quantified by fluorescence intensity (C).
Arrow and arrowhead in (B) denote EGFP+/Dhcr7- and EGFP-/Dhcr7+ cells, respectively. (D-
G) E12.5 cortical precursors were transfected with control or *Dhcr7* shRNAs and analyzed 3
days later. (D) Cultures were immunostained for EGFP (green) and Ki67 or βIII-tubulin (red;
double-labelled cells in orange are indicated with arrows) or CC3 and the proportion of total
EGFP+ cells that were also positive for Ki67 (E), βIII-tubulin (F) or CC3 (G) was quantified. *,
p < 0.001; n = 3. Scale bar = 50 μm. In all cases, error bars denote SEM. **(**H-K) E12.5
precursors were co-transfected with the PB transposase and PB-EGFP-control (Con) or PB-
EGFP-*DHCR7* shRNA (sh2). (H) Cultured cells were immunostained for EGFP (green), Sox2
(red) and βIII-tubulin (red) after 3 days and quantified for clones greater than one cell in size (I),
neuron-only clones (J), and clones with at least one Sox2+ precursors (K). Arrows in (I) top
denote EFGP+/ Sox2+ precursors. Arrow in (I) bottom denote EGFP+/βIII-tubulin+ cells. *, *p* <
0.001; n = 3. (M) Western blots of DHCR7 in 293T cells transfected with human control (Con)
or human-specific *DHCR7* shRNA (sh3) plus human *DHCR7*-expressing plasmid, analyzed after
24 hr. The blot was re-probed for Erk1/2. (N) Human cortical precursors were transfected with
EGFP-control (Con) or EGFP-*DHCR7* shRNA (sh3). Cells were immunostained 3 days later for
EGFP (green) and Ki67 (red), βIII-tubulin (red) or CC3 and the proportion of total EGFP+ cells
that were also positive for Ki67 (O), βIII-tubulin (P), or CC3 (Q) was quantified. *, *p* < 0.001; n
= 3. Scale Bar = 50 μm. Allows in (N) denote double-positive cells.
Figure 3-Figure Supplement 1. Rescue of the neurogenesis phenotype in *Dhcr7*-knockdown
mouse cortical precursors by human *DHCR7* cDNA expression vector.
278

Finally, we asked whether *Dhcr7* also plays a role in human iPSC-derived NPCs (Figure

3M-Q). To examine the function of DHCR7, we generated a human-targeted *DHCR7* shRNA-

EGFP and confirmed its efficiency by transfecting it into 293T cells along with human DHCR7

expressing vectors (Figure 3M). Human NPCs were transfected with this shRNA-EGFP and

analyzed by immunostaining 4 days later, which showed that the KD of human *DHCR7* reduced

EGFP+ and Ki67+ proliferating precursors whereas increased EGFP+ and βIII-tubulin+ newborn

neurons (Figure 3N-Q), similar to those observed in murine cortical precursors and SLOS

hNPCs.

287
288 ***7-DHC Derived Oxysterols Lead to Similar Neurogenic Defects as loss of Dhcr7 in Murine***
289 ***Cortical precursors in vitro***
290
Oxysterols have been found to influence biological processes, including proliferation,

differentiation and cell survival (DuSell et al., 2008; Theofilopoulos et al., 2013; Voisin et al.,

2017). Thus, we asked whether 7-DHC-derived oxysterols play regulatory roles in cortical

precursor biology. To assess potential effect of these 7-DHC-derived oxysterols on cortical

precursor behaviors, cortical precursors from E12.5 wild-type embryos were cultured and treated

with different concentrations of individual oxysterols for 3 days, followed by immunostaining

for Ki67, βIII-tubulin, and CC3 (Figure 4A). Notably, DHCEO-treated cortical precursor

showed significant increase in the proportion of βIII-tubulin+ newborn neurons and significant

decrease in the proportion of Ki67+ precursors (Figure 4B-C). This effect of DHCEO was

observed in a dose-dependent manner up to 3.5 µM, which is roughly the concentration of

DHCEO in the brains of P0 *Dhcr7*-KO mice (Xu et al., 2011). There was no significant change

in cell death/survival with DHCEO treatment (Figure 4D). The other oxysterols showed notable

toxic effect at a concentration of 5 µM as indicated by increased CC3+ cells (Figure 4G, 4J, and

4M). Note that the level of 7k-DHC is the lowest among the oxysterols, only reaching 0.9 ng/mg

of tissue (equivalent to 2.3 µM), so we do not expect the highest concentration examined, 5 µM,

for this oxysterol is relevant to the neurogenic phenotype. Interestingly, 4α-OH-7DHC treatment

at a concentration of 2 µM also displayed significant increases in βIII-tubulin+ neurons and

decreases in Ki67+ precursors without significant changes in cell death/survival (Figure 4H-J).

However, at 5 µM, CC3+ cells were significantly elevated at the expense of βIII-tubulin+

neurons (Figure 4J). To summarize, treatment with DHCEO at physiologically relevant

concentrations completely replicated the premature neurogenic phenotype observed in murine

and human SLOS NPCs while 4α-OH-7DHC may also contribute to the phenotype despite its

toxicity at high concentration.

Figure 4. 7-DHC Derived Oxysterols Lead to Similar Neurogenic Defects as loss of *Dhcr7* in Murine Cortical precursors *in vitro*. (A-M) E12.5 cortical precursors were cultured for 2 days in the presence of different concentrations of 7-DHC-derived oxysterols and quantified.

(A) Cell were immunostained for Ki67 (green), βIII-tubulin (red), and CC3 (red, arrow) after 3 days and the proportions of Ki67+ (B,E,H,K), βIII-tubulin+ (C,F,I,L), and CC3+ (D,G,J,M) cells were determined. Error bars indicate SEM. *, $p < 0.05$; **, $p < 0.005$; ***, $p < 0.001$. n = 3 per experiment. Scale Bar = 50 μm.

Figure 5. *Dhcr7$^{-/-}$* **mice display premature neurogenesis, altered neocortical cytoarchitecture *in vivo*, and increased activity of TrkB neurogenic signaling pathway.** (A) E18.5 cortical sections from *Dhcr7$^{+/+}$* and *Dhcr7$^{-/-}$* were immunostained for Tbr1 (red), Ctip2 (green) and counterstained with DAPI (blue). (B and C) Quantifications of absolute thickness (B) and number of positive cells (C). (D-G) Cortical sections from E15.5 *Dhcr7$^{+/+}$* and *Dhcr7$^{-/-}$* mice were immunostained for Satb2 (D, green) and Tbr1 (E, red). Quantifications of absolute thickness (F) and number of positive cells (G) for Satb2 and Tbr1. (H) E15.5 cortical sections from *Dhcr7$^{+/+}$* and *Dhcr7$^{-/-}$* embryos EdU-labeled at E12.5 were immunostained for EdU (green) and counterstained with DAPI (blue). (I) Quantification of relative location of EdU+ cells in cortical sections. (J) E15.5 cortices were isolated from *Dhcr7$^{+/+}$* and *Dhcr7$^{-/-}$* embryos and analyzed by western blot for phospho-TrkB, phospho-MEK, or phospho-C/EBPβ. Blots were re-probed with antibodies for total GR, TrkB, MEK, C/EBPβ and GAPDH as loading controls. Error bars indicate SEM. *, $p < 0.05$; **, $p < 0.005$; ***, $p < 0.001$. n = 3 per experiment. Scale Bar = 50 μm.

Loss of Dhcr7 causes premature differentiation and alter neocortical cytoarchitecture in vivo

We asked whether loss of *Dhcr7* disrupted cortical development *in vivo* as was observed

in vitro. To examine this, cortical sections from *Dhcr7$^{-/-}$* mice at E18.5 were immunostained for

the expression of well-defined cortical layer markers: the layer 2-4 marker Brn2, the layer 5

marker CTIP2, and the layer 6 marker Tbr1 (Molyneaux et al., 2007) (Figure 5A). The analyses

showed loss of *Dhcr7* led to reduction of all cortical layers, resulting in reduction of overall size

of the neocortex (Figure 5B-C). To further evaluate the effects of *Dhcr7* mutations in fate

decision of cortical precursors during corticogenesis, we immunostained cortical sections from

Dhcr7$^{-/-}$ mice at E14.5 and E15.5 for Tbr1 and Satb2, which label early-born and late-born

cortical neurons, respectively (Figure 5D-E). The analyses showed the significant increase of

Tbr1+ neurons and Satb2+ neurons in cortical section from *Dhcr7$^{-/-}$* mice at E14.5 and E15.5,

respectively (Figure 5F-G). This aberrant increased production of neurons with loss of *Dhcr7*

potentially hinders proper localization of developing neurons in the cortex. To test this

hypothesis, we injected pregnant dams with 5-ethynyl-2-deoxyuridine (Edu) at E12.5 to label

proliferating radial precursors and analyzed *Dhcr7$^{+/+}$* and *Dhcr7$^{-/-}$* littermates three days later at

E15.5 to identify the locations of Edu-labeled cells in the cortex (Figure 5H-I). Majority of Edu-

labeled cells migrated into the cortical plate in the KO mice whereas Edu-labeled cells were

scattered throughout the intermediate zone (IZ) and the cortical plates in the wild-type

littermates. These observations suggest premature increases in precursor differentiation, which

raises the possibility of activating major neurogenic signaling pathways during corticogenesis.

RTKs and their downstream targets are known to play important roles in precursor proliferation

and differentiation in developing cortex, and TrkB, one of the major RTKs expressed in cortical

precursors, regulates proliferation and differentiation into neurons by activating the MEK-ERK-

C/EBP pathway (Barnabe-Heider and Miller, 2003; Bartkowska et al., 2007; Menard et al.,

2002). Thus, we examined the activity of TrkB-MEK-ERK-C/EBP pathway by western blot with

antibodies for phosphorylated activated MEK1 and C/EBPβ at E15.5 of $Dhcr7^{-/-}$ cortices (Figure

5J). This analysis showed that relative to total MEK1 and ERK1/2 levels, their phosphorylated

forms increased in $Dhcr7^{-/-}$ cortices. This observation suggests that loss of $Dhcr7$ can

dysregulate an RTK-dependent signaling pathway that is critical for neural precursor

proliferation and differentiation.

The above observations indicate premature precursor differentiation at the expense of

depleting the proliferative precursor pool. To examine this hypothesis, pregnant dams were

injected with Edu at different time points of cortical development, and $Dhcr7^{+/+}$ and $Dhcr7^{-/-}$

littermates were analyzed 18 hours later to quantify the number of cycling progenitors that have

exited the cell cycles (Figure 6A-B). Cells that had left the cell cycle were stained as Edu+ and

Ki67- whereas cells that re-entered the cell cycle were stained as Edu+ and Ki67+. The index of

the cell cycle exit is defined by the percentage of Edu+/Ki67- cells in total Edu+ cells. The

index showed that the number of cells exiting the cell cycle in $Dhcr7^{-/-}$ cortices was significantly

increased at E13.5 and E14.5 compared with *Dhcr7$^{+/+}$* cortices, indicating that loss of *Dhcr7*

increased cell cycle exit. Increased cell cycle exit of cortical precursors can potentially arise from

changes in proliferation. To test this possibility, pregnant dams were pulsed with Edu for 2 hours

to label all cells in S-phase and the fraction of cortical precursors in S-phase, a proliferating

index from E13.5 to E15.5 in *Dhcr7$^{-/-}$* and *Dhcr7$^{+/+}$* cortices, was quantified (Figure 6C-D). The

proliferating index was determined by the percentage of Edu+ and Ki67+ cells out of total Ki67+

cells, which provides an estimate of cell cycle length because the length of S phase remains

relatively constant in mammalian cells, whereas the length of G1 phase regulates proliferation

(DiSalvo et al., 1995). The analyses showed a significant decrease in the proliferating index in

Dhcr7$^{-/-}$ cortices compared to *Dhcr7$^{+/+}$* cortices from E13.5 to E15.5 (Figure 6C-D), indicating

slower cycle progression and longer cycle length in *Dhcr7$^{-/-}$* cortices (Chenn and Walsh, 2002).

Indeed, this accelerated depletion of the progenitor pool in *Dhcr7$^{-/-}$* cortices result in decreases in

the size of VZ/SVZ compared to *Dhcr7$^{+/+}$* cortices from E13.5 to E15.5 (Figure 6E-F). Taken

together, these results suggest that loss of *Dhcr7* leads to a decrease in the overall number of

cortical precursors with increased cell cycle exit and decreased proliferation index via activation

of RTK-mediated MEK-ERK-C/EBP pathway.

Figure 6. ***Dhcr7$^{-/-}$*** **mice show accelerated cell cycle exiting and depletion of cortical precursor cells** ***in vivo.*** (A) Cortical sections from *Dhcr7$^{-/-}$* and *Dhcr7$^{+/+}$* embryos labeled by Edu at different developmental stages were immunostained 18 hours later for Edu (green) and Ki67 (Magenta). (B) Quantification of cell-cycle exit index of *Dhcr7$^{-/-}$* cortices compared with *Dhcr7$^{+/+}$* cortices. (C) Cortical sections from *Dhcr7$^{-/-}$* and *Dhcr7$^{+/+}$* embryos labeled by Edu injection at different developmental stages were immunostained 2 hours later for Edu (green) and Ki67 (Magenta). (D) Quantification of proliferation index of *Dhcr7$^{-/-}$* cortices compared with *Dhcr7$^{+/+}$* cortices. (E) Coronal cortical sections immunostained for Pax6 cortical precursor marker at different developmental stages. (F) Quantification of the relative size of the Pax6+ region shown as fractions of the whole cortical radial thickness. Error bars indicate SEM. *, $p < 0.05$; **, $p < 0.005$; ***, $p < 0.001$. n = 3 per experiment. Scale Bar = 50 μm.

Perturbation in DHCR7 Causes Gene Expression Changes in Neurogenic Pathways in Human NPCs

To ask whether *DHCR7* mutations influence the NPC biology, we aim to compare gene

expression profiles between hNPCs with *DHCR7* mutations versus hNPCs from healthy

individuals. To do so, we expanded the hNPCs derived from hiPSCs in the presence of essential

growth factors and carried out RNA sequencing (RNAseq) on these cells. Relative to wild-type

hNPCs, 2357 genes were significantly differentially expressed in SLOS hNPCs (1072

upregulated and 1285 downregulated genes with adjusted *p* value < 0.05), distinguishing the

SLOS and wild-type precursors (Figure S1A; Table S6 for RNAseq data). Gene ontology and

pathway analysis found that PI3K-Akt signaling pathway, MAPK signaling, and Ras signaling

pathways are among those with the highest *p*-values (Figure S1B-C; Table S5). Notably,

MAP2K1(also known as MEK1) was found to be upregulated in SLOS patient-derived hNPCs.

MAP2K1 is a key signaling molecule in MAPK and Ras signaling pathways, which are known to

play important roles in precursor proliferation and differentiation (Bonni et al., 1999; Menard et

al., 2002; Yang et al., 2013), consistent with the phenotypes that were observed in *Dhcr7*-KO

mice and SLOS patient-derived hNPCs *in vitro*.

7-DHC-derived oxysterol, DHCEO, activates cortical neurogenesis via activation of glucocorticoid receptor and inhibition of GR or inhibition of the formation of DHCEO rescues the neurogenic defects in SLOS NPCs

As shown in Figure 4, treatment of WT mouse NPCs with 7-DHC-derived oxysterols,

particularly DHCEO, can replicate the same aberrant premature neurogenesis observed in SLOS

NPCs. However, the mechanism of action of DHCEO remains unknown. Interestingly, 6-oxo-

cholestan-3β,5α-diol (OCDO), a cholesterol-derived oxysterol that is structurally similar to

DHCEO, is shown to bind and activate glucocorticoid receptor (GR) (Voisin et al., 2017) (Figure

7-Figure Supplement 1; Panel A). Additionally, GR activation is shown to activate TrkB

(Jeanneteau et al., 2008), which leads to further activation of RTK-mediated MEK-ERK pathway

(Barnabe-Heider and Miller, 2003; Bartkowska et al., 2007; Menard et al., 2002). TrkB and

RTK-mediated MEK-ERK pathway are necessary for neurogenesis during embryonic cortical

development. Thus, we asked whether DHCEO can activate the GR and further lead to the

activation of MEK-ERK neurogenic pathway. To investigate this, we first examined if GR was

phosphorylated and activated in the embryonic cortices from $Dhcr7^{-/-}$ and $Dhcr7^{+/+}$ embryos at

E15.5 in addition to TrkB, MEK, and C/EBPβ shown in Figure 5J. Western blots demonstrated

that phosphorylation of GR was indeed increased in $Dhcr7^{-/-}$ cortices comparing to $Dhcr7^{+/+}$

cortices (Figure 7A).

To ask if DHCEO could physically interact with GR, we performed molecular docking

simulation between DHCEO and the ligand binding domain of GR. Molecular docking is an

effective computational approach to understand protein-ligand interaction between small

molecules and receptor proteins both energetically and geometrically (Ferreira et al., 2015). As

seen in Figure 7B, DHCEO docks in the binding pocket of human GR favorably with a docking

score of -9 (a negative value suggests favorable interactions), comparable to the docking score of

OCDO at -9.5.

We further examined whether DHCEO could initiate GR and TrkB activation in wild-

type hNPCs $in\ vitro$. Cultured hNPCs were exposed to DHCEO (3.5 μM), a physiologically

relevant concentration, harvested at the different time points (3, 6, 9 and 24 hrs), and evaluated

for GR and TrkB activation by western blots (Figure 7C), which revealed that DHCEO-treated

hNPCs showed a gradual increase of phosphorylated GR starting at 3 hrs and peaking at 10 hrs

of exposure. As GR phosphorylation increased, DHCEO-treated hNPCs also showed the

phosphorylation of TrkB from 6 hrs to 10 hrs exposure, which dissipated at 24hrs exposure. As

no exogenous GR ligands and neurotrophins were added in these experiments, these results

indicated that DHCEO caused the increased phosphorylation of GR, which led to the activation

of TrkB phosphorylation in human NPCs *in vitro*.

Figure 7. DHCEO activates cortical neurogenesis via activation of glucocorticoid receptor and inhibition of the effect or the formation of DHCEO rescues the neurogenic defects in

SLOS NPCs. (A) Western blot showing increased phospho-GR in E15.5 $Dhcr7^{-/-}$ mouse brain relative to $Dhcr7^{+/+}$. (B) Image of the docked position of DHCEO (white) and OCDO (blue) in the ligand binding pocket of GR. (C) Human neural progenitors were treated with 3.5 μM DHCEO over the indicated time periods. Lysates were probed with phosphor-GR and phosphor-TrkB and re-probed with antibodies for total GR, total TrkB or total ERK as loading controls. (D) Human NPCs were transfected with EGFP-control (Con) or EGFP-$DHCR7$ (sh3) shRNA. Cells were treated with 1 μM RU38486, a selective GR antagonist one day after transfection. Three days post-transfection, cells were immunostained for EGFP and βIII-tubulin and quantified. (E) Wild type hNPCs were transfected with EGFP-control (Con) or EGFP-$DHCR7$ shRNA (sh3), and then treated with vitamin E, vitamin C, vitamin E/C, or glutathione (GSH). Three days post-transfection, cells were immunostained for EGFP and βIII-tubulin. (F,G) Quantification of EGFP and βIII-tubulin+ cells in $DHCR7$-KD wild-type hNPCs treated with various antioxidants, SAG, or LDL+SAG. (H) SLOS hNPCs were treated with cholesterol or various antioxidants and were immunostained for βIII-tubulin and DAPI. (I) Quantification of the proportion of βIII-tubulin+ cells in wild type hNPCs, and SLOS hNPCs treated with cholesterol or various antioxidants. (J) Quantification of DHCEO by LC-MS/MS in SLOS hiPSCs treated with vitamin E, vitamin E/C, and GSH. (K) Proposed mechanisms of actions of vitamin E and GSH during free radical oxidation of 7-DHC. Error bars indicate SEM. *, $p < 0.05$; **, $p < 0.005$; ***, $p < 0.001$. n = 3 per experiment. Scale Bar = 50 μm.
Figure 7-Figure Supplement 1. Antioxidants rescue the neurogenic phenotype in human and murine NPCs with $Dhcr7$ mutations.

To confirm that GR activation could cause aberrant premature neurogenesis, we cultured human cortical precursors and transfected them with $hDHCR7$ shRNA-EGFP and control shRNA-EGFP. 12 hours post-transfection, a glucocorticoid receptor antagonist, RU38486, or vehicle control was added and cultured for 3 days. Cultures were immunostained for EGFP and βIII-tubulin to evaluate the effects on neurogenesis. The analysis showed that inhibition of GR by RU38486 rescued the $DHCR7$-KD-mediated increase in neurogenesis down to the level observed in the control shRNA (Figure 7D).

Finally, we assessed whether aberrant premature neurogenesis was due to DHCEO, as it is a major oxysterol derived from free radical (non-enzymatic) oxidation of 7-DHC (Xu et al., 2011) (Figure 2). Antioxidants can prevent or slow down formation of oxysterols via the free radical mechanism and protect cells from deleterious effects of oxysterols (Yin et al., 2011). In particular, vitamin E (V_E), but not vitamin C (V_C), has been shown to effectively inhibit

formation of 7-DHC derived oxysterols in SLOS patient-derived fibroblasts (Korade et al.,

2014). Combination with V_C, which recycle oxidized V_E, enhances the antioxidant activity of V_E

against lipid peroxidation (Doba et al., 1985). Furthermore, glutathione (GSH) is an important

detoxifying molecule that is abundant (in mM) in cells and can potentially inhibit the formation

of DHCEO by reacting with its electrophilic precursor, 7-DHC $5\alpha,6\alpha$-epoxide (Porter et al.,

2020; Xu et al., 2011). To test this possibility, we treated *hDHCR7* KD human cortical

precursors with antioxidants (V_E, V_C, V_E/V_C, or GSH) and then immunostained for EGFP and

βIII-tubulin to evaluate the effects on neurogenesis (Figure 7E-F). The analysis indicated that

V_E, V_E/V_C, or GSH, but not V_C alone, effectively reversed increased neurogenesis caused by

hDHCR7 KD to the level observed in the control. Potential detoxifying effects of these

antioxidants were further tested in human SLOS patient-derived NPCs as well as cortical

precursors from *Dhcr7$^{-/-}$* mouse embryos *in vitro* (Figure 7H-I and Figure 7-Figure Supplement

1, panels B-E). Treatments with V_E, V_E/V_C and GSH were again found to effectively reduce the

increased neurogenesis in both human and murine cortical precursors with *Dhcr7* mutations

down to the level observed in WT samples. LC-MS/MS analysis confirmed that V_E, V_E/V_C, or

GSH indeed inhibited the levels of DHCEO in SLOS hiPSCs (Figure 7J). These results suggest

that free radical chain-breaking antioxidant, such as V_E, and nucleophilic antioxidant, such as

GSH, can effectively rescue the neurogenesis phenotype observed in SLOS NPCs by inhibiting

the free radical oxidation of 7-DHC at different stages (Figure 7K).

In addition to antioxidant treatment, we tested whether restoration of cholesterol could

potentially rescue aberrant premature neurogenesis observed in both human and murine SLOS

NPCs (Figure 7I, S6C and S6E). Low-density lipoprotein was added to cultures to examine the

effects of cholesterol on neurogenesis. The results revealed that addition of cholesterol did not

prevent increased neurogenesis in SLOS NPCs. Because DHCEO has been shown to inhibit

Smo in the Hh signaling pathway (Sever et al., 2016), we also tested whether supplementation of

SAG (a Smo agonist) can rescue the phenotype with or without cholesterol supplementation in

DHCR7-KD hNPCs (Figure 7G). However, none rescued the neurogenesis phenotype. Thus, the

lack of cholesterol and inhibition of Hh signaling do not contribute to the neurogenesis

phenotype observed in SLOS NPCs.

Taken together, the abnormal premature neurogenesis in cortical precursor with *Dhcr7*

mutations was caused by DHCEO activation of GR and further activation of Trk-mediated

neurogenic signaling pathway, but not by the deficiency in cholesterol, and inhibition of the

formation of DHCEO or GR rescues the neurogenic defects.

Discussion

Whether deficiency in cholesterol, accumulation of 7-DHC, or accumulation of 7-DHC

oxysterols is more important for the neurological and developmental phenotypes observed in

patients in SLOS remains unknown. Here, we characterized the neurogenic phenotype of SLOS

cortical precursors *in vitro* and *in vivo* and showed that the accumulated 7-DHC-derived

oxysterols, particularly DHCEO, disrupts normal embryonic neurogenesis by accelerating

progressive depletion of cortical precursors through neurogenesis, ultimately resulting in greatly

reduced cerebral cortex thickness with abnormal cortical layering.

The data presented here support four major conclusions. First, LC-MS/MS analyses

indicate that *DHCR7* mutations promote significant accumulation of 7-DHC-derived oxysterols

in murine and human embryonic cortical NPCs during cortical development. Second, our studies

with KD or KO of *DHCR7* indicate that *DHCR7* is necessary for normal proliferation of

embryonic NPCs and genesis of newborn neurons in culture and within the environment of the

embryonic cortex. KD or KO of *DHCR7* leads to premature differentiation of cortical precursors

to neurons and ultimately a perturbation in cortical cytoarchitecture. This occurs through the

activation of neurogenic TrkB-MEK-C/EBP pathway. Third, we show that 7-DHC-derived

oxysterols, especially DHCEO, display detrimental effects to normal neural development, and

that DHCEO can activate GR, which is responsible for abnormal cortical precursor

differentiation, since *DHCR7* KD/KO phenotype can be rescued by concurrent inhibition of

glucocorticoid receptor. Fourth, we show that premature differentiation of cortical precursors

with *DHCR7* mutations is cholesterol-independent since inhibition of 7-DHC-derived oxysterols

via antioxidant treatment can rescue *DHCR7* KD/KO phenotype whereas the cholesterol

supplementation cannot.

Based on these data, we propose a model where DHCEO binds and activates GR, and in

doing so, controls the RTK-mediated neurogenic pathway, TrkB-MEK-C/EBP, and thus fate-

decision of developing precursors and neurons. This model is consistent with previous studies

showing that an oxysterol structurally similar to DHCEO, OCDO, can bind and regulate GR and

its transcriptional activity (Voisin et al., 2017) and ligand-bound GR promoted TrkB

phosphorylation, leading to activation of its downstream effectors, MEK and ERK (Jeanneteau et

al., 2008).   Our model of DHCEO exerting its effect on neurogenesis through GR does not

contradict the previous finding by Francis et al., which suggests that Wnt/β-catenin inhibition

contributes to precocious neurogenesis (Francis et al., 2016), because activation of GR has been

shown to inhibit Wnt/β-catenin signaling in several systems (Olkku and Mahonen, 2009; Zhou et

al., 2020).

This work represents the first comprehensive characterization of the neurogenic

phenotype in SLOS mouse model and SLOS patient-derived NPCs. We demonstrate that the

aberrant neural development observed in mice and human NPCs with *DHCR7* mutations occurs

in a cholesterol-independent and 7-DHC-derived oxysterol-dependent manner. Successful

rescue of the neurogenic phenotype by inhibiting GR or inhibiting the formation of oxysterols

with antioxidants pave the wave for potential therapies toward the neurological defects observed

in SLOS patients.

Acknowledgements

This work was supported by a National Institutes of Health (NIH) grant R01HD092659 (LX) and

a grant from Smith-Lemli-Opitz/RSH Foundation. JMH was supported by the University of

Washington (UW) Environmental Pathology/ Toxicology Training Program (NIH T32

ES007032) and AL was supported by the UW Pharmacological Sciences Training Program (NIH

T32 GM007750) and the Institute of Translational Health Sciences TL1 Program (NIH TL1

TR002318). We also thank Dr. Carol Ware and Christopher Cavanaugh for technical advices on

hiPSC culture.

Author Contributions

HT conceptualized, designed, performed, and analyzed most of the experiments and co-wrote the

paper. LX conceptualized, designed experiments, analyzed data and co-wrote the paper. KMH

and AL performed and analyzed LC-MS/MS. JMH analyzed RNA-sequencing experiments. DB

performed and analyzed docking simulation.

Competing interests

The authors declare no competing interests.

Materials and Methods

Animals. Animal use was approved by the University of Washington Institutional Animal Care

and Use Committee. *Dhcr7* mice (B6.129P2(Cg)-*Dhcr7*[tm1Gst]/J) were maintained as heterozygous

and were genotyped as described previously (Fitzky et al., 2001). The primers used are the

following: *Dhcr7*-WT-F: 5'-GGATCTTCTGAGGGCAGCTT-3'; *Dhcr7*-WT-R: 5'-

TCTGAACCCTTGGCTGATC-3'; Delta-Mut: 5'-CTAGACCGCGGCTAGAGAAT-3'. For *in*

vitro transfection, C57BL/6J E12.5 pregnant mice were obtained by time-mating. Mating pairs

of wild-type C57BL/6J mice were purchased from Jackson Laboratories (Bar Harbor, ME). All

mice had free access to rodent chow and water in a 12-hour dark-light cycle room.

Plasmids. The target sequence for murine *Dhcr7* shRNA was cloned into pLKO-UBI-GFP

digested with EcoRI and PacI. pLKO-UBI-GFP was generated by digesting out hPGK promoter

from pLKO.3G and ligating ubiquitin promoter (UBI) from pCLX-UBI-VenusN with PacI and

BamHI. pLKO.3G was a gift from Christophe Benoist & Diane Mathis (Addgene plasmid #

14748; http://n2t.net/addgene:14748; RRID:Addgene_14748) and pCLX-UBI-VenusN was a gift

from Patrick Salmon (Addgene plasmid # 27247; http://n2t.net/addgene:27247;

RRID:Addgene_27247). cDNAs encoding murine (pCMV-mDHCR7-Myc/DDK) and human

DHCR7 (pCMV-hDHCR7-Myc/DDK) were purchased from Origene (MR223420 and

RC228922 for murine and human cDNA clone respectively). PB-EF1α-GreenPuro-H1MCS and

Super PiggyBac transposase expression vector from System Biosciences (Cat#PBS506A-1 and

Cat# PB210PA-1 respectively) were used for clonal analysis.

Cortical precursor cell cultures. Murine cortical precursor cells were cultured as described

previously (Barnabe-Heider and Miller, 2003; Tomita et al., 2020). Briefly, mouse cortical

precursor cells from cortices were dissected from E12.5 wild-type or *Dhcr7*-KO mouse embryos

in ice-cold HBSS (Invitrogen) and transferred into cortical precursor medium. The cortical

precursor medium consisted of Neurobasal medium (Invitrogen) with 500 μM L-glutamine

(Invitrogen), 2% B27 supplement (Invitrogen), 1% penicillin-streptomycin (Invitrogen) and 40

ng/ml FGF2 (BD Biosciences). The dissected tissue was mechanically triturated by a fire-

polished glass pipette and plated onto 24-well plates coated with 2% laminin (BD Biosciences)

and 1% poly-D-lysine (Sigma Aldrich). Plating density of the cortical precursors was 150,000

cells/well in 24-well plates for single embryo cultures and plasmid transfections.

Human induced pluripotent stem cells (hiPSCs). Human fibroblast cell lines (GM03044 and

GM05788) isolated from patients with SLOS were obtained from Coriell Institute and

reprogrammed as described elsewhere (Okita et al., 2011; Yu et al., 2007). Briefly, 1×10^6 cells

of human fibroblast cells at early passages were trypsinized (0.25% Trypsin/0.5mM EDTA,

Gibco) and electroporated with indicated episomal plasmids using Amaxa Basic Nucleofector

Kit for primary mammalian fibroblasts, program A-24 (Lonza). 1.0μg pSIN4-EF2-O2S and

pSIN4-EF2-N2L were used for each electroporation. The electroporated human fibroblasts were

seeded onto a gelatin-coated tissue culture dish and cultured with fibroblast medium (DMEM

with 10% FBS, 2mM L-glutamine, 1% non-essential amino acids, 1% sodium pyruvate, and 1%

penicillin and streptomycin) for 7 days and re-seeded onto feeder layer cells (irradiated SNL 76/7

feeder cells; a gift from Dr. Allan Bradley, Sanger Institute, UK) with hiPSC medium

(DMEM/F12 medium with 20% knockout serum replacer, 1% sodium pyruvate, 1% non-

essential amino acids, 0.007% 2-mercaptoethanol,1% penicillin and streptomycin, 0.1mM

sodium butyrate, 50nM suberoylanilide hydroxamic acid, and 4ng/ml of bFGF) in 100mm tissue

culture dishes and continued to be cultured until hiPSC colonies were visible (28 days from

electroporation). The hiPSC colonies were then picked and further cultured for expansion.

Maintenance and Differentiation of hiPSC. Undifferentiated hiPSCs were cultured and

maintained in mTeSR Plus medium (StemCell Technologies) on Matrigel (BD Biosciences)-

coated plates prior to the generation of human neural progenitor cells (hNPCs). hNPCs was

generated using the STEMdiff SMADi Neural Induction kit (StemCell Technologies) according

to the manufacturer's protocol. Briefly, neuralized embryoid bodies (EBs) were generated by

culturing small aggregates of hESCs in ultra-low attachment plates (Corning) in STEMdiff

SMADi Neural induction medium. EBs were replated and cultured in poly-D-lysine/Laminin-

coated plates with the neural induction medium until EBs formed neural rosette formations.

Neural rosettes were collected and replated in poly-D-lysine/Laminin-coated plates for hNPCs

outgrowth in neural induction medium until hNPCs were ready for the first passage. hNPCs

were maintained in STEMdiff Neural Progenitor medium. For neurogenesis of hNPCs, hNPCs

were cultured in Neurobasal A medium supplemented with B27 minus vitamin A, 1% penicillin-

streptomycin and glutamax (all from Invitrogen) in poly-D-lysine/Laminin-coated plates

(Konopka et al., 2012; Usui et al., 2017). For quantification, an average of 700 cells was

counted per condition in 5-6 random fields per independent experiment.

In vitro differentiation of iPSCs. For in-vitro differentiation of human iPSCs to the three germ

layers, STEMdiff Trilineage Differentiation Kit (StemCell Technologies) was used as described

by the manufacturer. Briefly, cells were harvested by Gentle Cell Dissociation Reagent

(StemCell Technologies) and plated onto Matrigel-coated 24 well plates with mTeSR1 medium

(StemCell Technologies). Cell density and viability were determined using trypan blue

exclusion. Cells were seeded at a clonal density of 200,000 cell/cm^2 for ectoderm and endoderm

differentiation and 50,000 cell/cm^2 for mesoderm differentiation. 24 hours after plating, cells

were switched to the appropriate STEMdiff Trilineage medium for ectoderm, mesoderm and

endoderm differentiation. Cells were cultured in the lineage specific medium for 5 days

(mesoderm and endoderm lineages) and 7 days (ectoderm lineage) and were harvested and/or

fixed for analyses of lineage-specific markers for the three germ layers (Figure 1-Figure

Supplement 1).

Transfection and quantification. For plasmid transfection of mouse cortical precursors,

Lipofectamine LTX and Plus Reagent (Invitrogen) were used as described by the manufacturer.

Briefly, 1 μg of DNA and 1 μl of Lipofectamine LTX and Plus Regent in 100 μl of Opti-MEM

(both from Invitrogen) were mixed, incubated for 20 min, and added to precursors three hours

after plating. For clonal analysis, 1.5 μg DNA, at 1:3 ratio of Super PiggyBac transposase

expression vector to shRNA or control plasmids, were incubated as described above and added

to cortical precursors three hours after plating. For plasmid transfection of human cortical

precursors, Lipofectamine Stem Transfection Reagent (Invitrogen) was used as described by the

manufacturer. Briefly, 1μg of DNA and 1μl of Lipofectamine Stem Transfection Reagent were

prepared separately in 25μl of Opti-MEM medium (Invitrogen), mixed, incubated for 30 min and

added to precursors one day after plating. The target sequences for murine and human *Dhcr7*

shRNA were 5'- GGAAGGTGCTTCTTGTTTA-3' and 5'- GGAAGTGGTTTGACTTCAA-3'

respectively. The target sequence for the control shRNA was 5'-TCCCAACTGTCACGTTCTC-

3'. For quantification, immunostaining and image acquisition were performed, and > 100 cells

per condition per experiment were counted and analyzed, and experiments were performed with

3 embryos per plasmid transfected and analyzed individually.

Neurosphere cultures. E13.5 cortices from *Dhcr7$^{+/+}$* or *Dhcr7$^{-/-}$* embryos were dissected and

mechanically dissociated into a single cell suspension by fire-polished glass pipette as previously

described (Capecchi and Pozner, 2015; Tomita et al., 2020). Cell density and viability were

determined using trypan blue exclusion. Cells were seeded in triplicate at a clonal density of 10

cells/μl in 6 well (2 ml/well) ultra-low attachment culture plates (Coster) in serum-free medium

supplemented with 20 ng/ml EFG (Sigma), 20 ng/ml FGF2 (Sigma), 2% B27 supplement

(Invitrogen) and 2 μg/ml heparin (Sigma). Neurospheres were cultured for 6 days at 37 ºC. To

evaluate self-renewal potential, neurospheres were mechanically dissociated into single cell

suspensions by fire-polished glass pipette, passed through a 45μm nylon screen cell strainer, and

cultured at a clonal density of 2 cells/μl for an additional 6 days.

Immunocytochemistry and histological analysis. For morphometric analysis, immunostaining of

tissue sections was performed as described (Capecchi and Pozner, 2015). Briefly, brain sections

were permeabilized and blocked in PBST (1X PBS, 0.5% (v/v) Triton X-100) containing 10%

NGS for 1 hour. Brain slices were incubated with primary antibodies in PBST with 5% NGS at 4

ºC overnight. The sections were incubated with secondary antibodies in PBST with 5% NGS for

1-2 hours at room temperature. Sections were counterstained with DAPI (Thermo Fisher

Scientific). Slides were mounted in Fluoromount-G anti-fade reagent (Southern Biotech).

Digital image acquisition was performed with EVO-FL Imagining System (Thermo Fisher

Scientific). For quantification of precursor and neuron numbers, we analyzed sections at the

medial-lateral level, counting all marker-positive cells in a 200 μm wide strip of the cortex,

extending from the meninges to the ventricle. In all cases, we analyzed at least 3 similar cortical

sections/embryo or pup from 3 different embryos or pup per genotype (for a total of at least 9

sections per genotype).

Antibodies. The primary antibodies used for immunostaining were chicken anti-GFP (1:1000;

Abcam), rabbit anti-Dhcr7 (1:100; Thermo Fisher Scientific), rabbit anti-Sox2 (1:200; Cell

Signaling Technology), rabbit anti-Pax6 (1:1000; Covance), mouse anti-Ki67 (1:200; BD

Biosciences), mouse anti-βIII-tubulin (1:1000; Covance), rabbit anti-βIII-tubulin (1:1000;

Covance), rabbit anti-Tbr1 (1:500; Abcam), mouse anti-Satb2 (1:400; Abcam), rat anti-Ctip2

(1:500; Abcam), and rabbit anti-cleaved caspase 3 (1:400; Cell Signaling Technology), mouse

anti-Nestin (1:1000; StemCell Technologies), rabbit anti-Brachyury (1:200; R&D Systems), goat

anti-NCAM (1:200; R&D Systems), goat anti-Sox17 (1:200; R&D Systems), rabbit anti-Foxa2

(1:1000; Abcam). The secondary antibodies used for immunostaining were Rhodamine

(TRITC)-conjugated goat anti-mouse and anti-rabbit IgG (1:500; Jackson ImmunoResearch

Laboratories) and Alexa Fluor 488-conjugated goat anti-mouse, anti-rat and anti-rabbit IgG

(1:800; Jackson ImmunoResearch Laboratories), Alexa Flour 488-conjugated donkey anti-goat

IgG (1:800; Jackson ImmunoResearch Laboratories), Rhodamine (TRITC)-conjugated donkey

anti-goat (1:200; Jaclson ImmunoResearch Laboratories). The primary antibodies used for

immunoblotting were rabbit anti-Dhcr7 (1:1000; Abcam), rabbit anti-GAPDH (1:5000; Santa

Cruz Biotechnology), rabbit anti Erk1/2 (1:5000; Santa Cruz Biotechnology), rabbit anti-TrkB

(1:1000; Cell Signaling Technology), rabbit anti-phospho-TrkB (1:1000; Cell Signaling

Technology), rabbit anti-glucocorticoid receptor (1:1000; Cell Signaling Technology), rabbit

anti-phospho-glucocorticoid receptor (1:1000; Cell Signaling Technology), rabbit anti-MEK

(1:1000; Cell Signaling Technology), rabbit anti-phospho-MEK (1:1000; Cell Signaling), rabbit

anti-phospho-cebpβ (1:1000; Cell signaling Technology) and rabbit anti-cebpβ (1:1000; Cell

Signaling Technology), rabbit anti-Pax6 (1:1000; Covance), mouse anti-Nestin (1:1000;

StemCell Technologies), rabbit anti-Brachyury (1:200; R&D Systems), rabbit anti-Tbr2 (1:500;

Abcam), goat anti-Sox17 (1:200; R&D Systems), rabbit anti-Foxa2 (1:1000; Abcam). The

secondary antibodies used for immunoblotting were HRP-conjugated goat anti-mouse IgG

(1:5000; Jackson ImmunoResearch Laboratories) and anti-rabbit IgG (1:10,000; Jackson

ImmunoResearch Laboratories), HRP-conjugated donkey anti-goat IgG (1:1000; Jackson

ImmunoResearch Laboratories).

RT-PCR. Total RNA was isolated with Trizol and cDNA was prepared using the SuperScript III

Reverse Transcriptase kit (Invitrogen) according to the manufacture's protocols. Primer

sequences are the following: *Dhcr7*-F: 5'-TATGAGGTGAATGGGCTGCA-3'; *Dhcr7*-R: 5'-

GGTTAATGAGGGTCCAGGCT -3'; *β-actin*-F: 5'-GATGACGATATCGCTGCGCTG-3'; *β-*

actin-R: 5'- GTACGACCAGAGGCATACAGG-3'. All PCR products were single bands with

predicted molecular weights and confirmed by DNA sequencing.

Quantitative PCR. Total RNA was extracted with Tri-Reagent (Sigma) treated with DNAse I

(Fermentas, Thermo Scientific, Waltham, MA, USA) and cDNA was synthesized from 1μg of

RNA using the SuperScript IV Reverse Transcriptase Kit (Invitrogen) according to the

manufacturer's protocols. Quantitative PCR was performed using Taqman Fast Advance Mater

Mix (Thermo Fisher Scientific) and Taqman probes targeted against either *Dhcr7*

(Mm01164321_m1) or *β-Actin* (Mm00607939_s1). *β-Actin* mRNA was used as an endogenous

control for all reactions, and all reactions were performed in triplicate. Quantitative PCR was

performed and analyzed using StepOne Plus Real-Time PCR system (ThermoFisher Scientific).

Western Blotting. Embryonic cortices or neurosphere cultures were lysed in RIPA buffer (50

mM Tri pH8, 150mM NaCl, 1% NP-40, 0.1% SDS, 1mM EDTA) containing 1mM PMSF

(phenylmethanesulfonyl fluoride), 1mM sodium vanadate, 20mM sodium fluoride 10 μg/ml

aprotinin and 10 μg/ml leupeptin. 10-20 μg of protein lysate was electrophoresed, and western

blots were performed as described previously (Barnabe-Heider and Miller, 2003).

RNA sequencing and data analysis. Raw RNA sequencing reads in FASTQ format were

mapped to the human genome using HISAT (https://ccb.jhu.edu/software/hisat/; Last accessed

January 22, 2021), and format conversions were performed using Samtools. Cufflinks

(http://cole-trapnell-lab.github.io/cufflinks/; Last accessed January 22, 2021) was used to

estimate relative abundances of transcripts from each RNA sample. Cuffdiff, a module of

Cufflinks, was then used to determine differentially expressed genes (DEGs) between WT and

KO hNPCs. DEGs met the following criteria: adjusted p value < .05 (corresponding to the

allowed false discovery rate of 5%) and fold-change > 1.5 between genotypes. A two-way

hierarchical clustering dendrogram (complete-linkage method, Euclidean distance scale) of

DEGs was used to visualize biological variability among samples, generated by R software using

the "pheatmap" package (https://cran.r-project.org/web/packages/pheatmap/). To elucidate

biological functions of DEGs, the Core Analysis feature of Ingenuity Pathway Analysis (IPA®,

Qiagen) was used to identify significantly enriched Diseases and Biological Functions related to

the nervous system. Network interactions among DEGs involved in the Biological Function

"development of the central nervous system" were assessed using STRING (Search Tool for the

Retrieval of Interacting Genes/Proteins) analysis, set at the highest confidence interaction score

and only connected nodes displayed (https://string-db.org/cgi/input.pl?sessionId=xCahIfrzvltC;

Last accessed January 22, 2021). Enriched KEGG (Kyoto Encyclopedia of Genes and Genomes)

pathways were identified among DEGs in the STRING network. Finally, a Venn diagram was

generated to demonstrate the overlap between genes dysregulated in KO hNPCs and genes in the

SFARI database, a collection of genes implicated in autism susceptibility (https://gene.sfari.org/;

Last accessed January 22, 2021). Raw data of the RNA sequencing has been deposited at Dryad

(https://doi.org/10.5061/dryad.zw3r2287f; available upon publication). The list of differentially

expressed genes can be found in Supplementary File 2 (Excel).

Oxysterol and Sterol Analysis. Cell pellets were resuspended in 300 μL of 1X PBS and lysed by

sonication in an ice bath for 30 min. Protein determination was performed using the BioRad DC

protein mass assay (BioRad, Hercules, CA). Internal standard mixtures for sterols and oxysterols

analysis were added to each sample (see Tables S1-4 for a list of standards and their

concentrations used). Lipid extraction was performed using the Folch method as described

previously (Fliesler et al., 2018; Folch et al., 1957; Herron et al., 2018). The dried lipid extract

was reconstituted with 200 μL of methylene chloride and stored at -80 ºC until analysis. Prior to

analysis, 50 μL of extract was transferred to into glass LC vials, dried under argon, and

reconstituted with 50 μL of 90% methanol in water with 0.1% formic acid. For tissue samples,

sterol and oxysterol internal standard mixtures were added to whole tissues, which were

subsequently homogenized in 4 mL Folch solution with 1 mL 0.9% NaCl. The dried lipid extract

was reconstituted with 0.5 mL (for tissues <50 mg), 1.0 mL (tissues >50 <100 mg) or 1.5 mL

(>100 mg) of methylene chloride. Prior to analysis, 50 μL of lipid extract (30 μL for tissues

>100 mg) was transferred into glass LC vials, dried under argon, and reconstituted with 50 μL

90% methanol in water with 0.1% formic acid. Determination of oxysterol and sterol

concentrations in cells and tissues was performed by ultra-performance liquid chromatography

(UPLC) tandem mass spectrometry (MS/MS) on a SCIEX 6500 triple quadrupole mass

spectrometer (for oxysterols) or a SCIEX 4000 QTRAP (for sterols) mass spectrometer with

atmospheric pressure chemical ionization (APCI) coupled to a Waters Acquity UPLC system, as

described previously (Fliesler et al., 2018; Herron et al., 2018). Briefly, sterols and oxysterols

were separated by reversed phase chromatography on a C18 column (1.7 mm, 2.1 x 100 mm,

Phenomenex Kinetex) using a 15 min isocratic gradient of 90% methanol with 0.1% formic acid

at a flow of 0.4 mL/min. Selective reaction monitoring (SRM) was used to monitor the

dehydration of the sterol and oxysterol $[M+H]^+$ ions to generate $[M+H-H_2O]^+$ ions (See Table

S1-4 for a list of retention times and MS/MS transitions for the standards used). The APCI

parameters were as follows: nebulizer current, 3 mA; temperature, 350 ºC; curtain gas, 20 psi;

ion source gas, 55 psi. The MS conditions for SRM analysis were as follows: declustering

potential, 80 V; entrance potential, 10 V; collision energy, 25 V; collision cell exit potential, 20

V. Data analysis was performed with Analyst (v. 1.6.2) Quantitation Wizard. Analyte

concentrations in cells and tissues were determined relative to the internal standard levels and the

relative response factor (RRF) of each analyte calculated from a mixture of sterol and oxysterol

standards and internal standards. Concentrations were normalized to cell protein weight or tissue

weight.

Docking simulation. Docking simulation was performed with Autodock Vina 1.1.2. The model

ligands and 3D models were built and generated by Openbabel 2.3.2. Crystal structure of

glucocorticoid receptor and docking of cortisol to glucocorticoid receptor were from crystal

structure 4p6w available from Protein Data Bank (http://www.rcsb.org/structure/4p6w).

Statistical analysis. In general, three biological replicates were used for each experiment based

on the strong phenotype observed in the *Dhcr7*-KO SLOS model and our previous publication

showing the large differences in oxysterols and sterol levels in WT and KO brains at birth (Xu et

al., 2011). For culturing of mouse cortical precursors or neurospheres from single embryos, one

litter is considered a biological replicate. For hNPC culture, each separate preparation of hNPC

from hiPSC is considered a biological replicate. For *in vivo* study, one litter is considered a

biological replicate. Statistical analyses were performed using two-tailed Student's *t*-test

assuming unequal variance when comparing two different groups unless otherwise indicated in

the text. For immunostaining of cell culture or tissue sections, at least three technical replicates

per biological replicate were performed. To analyze the multi-group neuroanatomical studies, we

used one-way ANOVA unless otherwise indicated in the text. Significant interactions or main

effects were further analyzed using Newman-Keuls post-hoc tests. All statistical tests were

performed using Microsoft Excel or Prism 8 (GraphPad). In all cases, error bars indicate standard

error of the mean.

Figure Supplements

Figure 1-Figure Supplement 1. Characterization of the pluripotency of SLOS-derived
human iPSCs. Related to Figure 1. Pluripotency of SLOS patient derived human iPSC lines,
3044 and 5788 were characterized. (A) Representative images of established SLOS patient
derived human iPSC lines (3044 and 5788) and human iPSC line from healthy individual
(emhf2). (B) RT-PCR analysis of hES cell marker genes in human iPSCs derived from SLOS
patients (3044 and 5788) and healthy individual (emhf2). Primers used for Oct3/4, Sox2, Klf4
and c-Myc specifically amplified endogenous transcripts but not transgenic transcripts. (C)
Immunostaining confirming in-vitro differentiation of SLOS patient derived human iPSCs (3044
and 5788) into all three germ layers. (D) RT-PCR analyses and Western blot analyses of
differentiation markers for the three germ layers.

855
856

Figure 1-Figure Supplement 2. Loss of *DHCR7* alleles causes decreased proliferation and
increased neurogenesis in human cortical precursors. (A) Images of cultured human cortical
precursors immunostained for Dhcr7 (red) and Sox2 or Nestin (green). (B-D) Human SLOS-
patient-derived (SLOS 3044 and SLOS 5788) and unaffected-patient-derived cortical precursors
were cultured for 3 days and analyzed. (B) Cells were immunostained for Ki67 (green) and βIII-
tubulin (red) after 3 days and quantified the proportions of Ki67+ (C) and βIII-tubulin+ cells (D).
Scale Bar = 50 μm. *, $p < 0.001$; n = 3 biological replicates per genotype.

Figure 2-Figure Supplement 1. Cholesterol precursor 7-DHC and 7-DHC-derived
Oxysterols are accumulated *in* SLOS-derived human iPSCs and NPCs. Related to Figure 2.
SLOS patient derived hiPSCs and NPCs (3044 and 5788) along hiPSCs and NPC from healthy
individual (emhf2) were cultured and used for LC-MS/MS analysis sterols and oxysterols.
Quantitation of cholesterol, its precursors and accumulated 7-DHC derived oxysterols in SLOS
patient derived hiPSCs and hiPSCs from healthy individuals (A) and in SLOS patient derived
NPCs and NPCs from healthy individuals (B).

Figure 3-Figure Supplement 1. Rescue of the neurogenesis phenotype in *Dhcr7*-knockdown
mouse cortical precursors by human *DHCR7* cDNA expression vector. Related to Figure 3.
Potential off-target effects of murine *Dhcr7* shRNAs were determined. Human *DHCR7* cDNA
expressing vectors transfected murine NPCs with either murine Dhcr7 shRNA vector or control
shRNA vectors. The transfected cells were cultured for 3 days, immunostained and analyzed for
proliferation marker, Ki67 or neuronal marker, βIII tubulin. (A) Cultures were immunostained
for EGFP-mDhcr7shRNA and proliferation marker Ki67, the proportion of total EGFP+ cells
that were also positive for Ki67 was quantified. (B) Cultures were immunostained for EGFP-
mDhcr7shRNA and neuronal marker, βIII tubulin, the proportion of total EGFP+ cells that were
also positive for βIII tubulin was quantified.

Figure 7-Figure Supplement 1. Antioxidants rescue the neurogenic phenotype in human
and murine NPCs with *Dhcr7* mutations. Related to Figure 7. (A) Structures of DHCEO and
known ligands of GR, OCDO and cortisol. (B, D) Another SLOS hNPC line (SLOS 5788) and
mNPCs from E12.5 *Dhcr7*$^{-/-}$ and *Dhcr7*$^{+/+}$ embryonic cortices were cultured and treated with

cholesterol, vitamin E, vitamin C, vitamin E/C or glutathione (GSH). The cells were
immunostained for βIII-tubulin and DAPI three days after plated. (C) Quantification of the
proportion of βIII-tubulin positive cells in wild type hNPCs and SLOS hNPCs treated with
cholesterol or antioxidants. (D) Quantification of the proportion of βIII-tubulin positive cells in
Dhcr7$^{-/-}$ and *Dhcr7*$^{+/+}$ NPCs treated with cholesterol or antioxidants. Error bars indicate SEM. *,
p< 0.001. n = 3 per experiment. Scale Bar = 50 μm.
894

914   **Supplementary File 1**
915
916   **Table S1. Retention times and MS/MS transitions for oxysterol internal standards. Relate to**
917   **Figure 2 and S3.**
918
919   **Table S2. Retention times and MS/MS transitions for all oxysterol standards. Relate to**
920   **Figure 2 and S3.**
921
922   **Table S3. Retention times and MS/MS transitions for sterol internal standards. Relate to**
923   **Figure 2 and S3.**
924
925   **Table S4. Retention times and MS/MS transitions for sterol standards. Relate to Figure 2**
926   **and S3.**
927
Table S5. Ingenuity Pathway Analysis (IPA®) reveals "development of the central nervous
system" as one of the top 10 enriched Diseases and Biological Functions related to the
nervous system. Related to Figure S1. DEGs were further analyzed with (IPA®) to identify the
most enriched biological functions related to the nervous system in SLOS mutant NPCs. The
table below shows the top ten enriched terms corresponding to Diseases/Bio-functions along
with the *p*-value and overlapping number of genes in the dataset.
934
Figure S1. Perturbation in *DHCR7* causes gene expression changes in neurogenic
pathways in human NPCs. (A) Hierarchical clustering heatmap of differentially expressed
genes show distinct expression pattern changes in transcript abundance for SLOS mutant NPCs
as compared to wild type. Red color represents increase in abundance, blue color represents
relative decrease, and white color represents no change. (B) Enriched KEGG pathways identified
among DEGs from SLOS mutant NPCs and involved in the Biological Function "development
of the central nervous system", identified by Ingenuity Pathway Analysis (see Supporting
Information). Color of bar corresponds to DEGs in the STRING network that are in enriched
pathways. (C) String analysis of DEGs from SLOS mutant NPCs and involved in the Biological
Function "Development of the central nervous system" identified by Ingenuity Pathway
Analysis. Parameters of high confidence have been applied and only connected nodes are
displayed. Arrows indicate whether the gene was up- or down-regulated. Color of DEG
corresponds to enriched KEGG pathway. *n* =3 biological replicates per genotype; DEGs met the
criteria of fold-change > 1.5 between genotypes and adjusted *p*-value < 0.05.
949
950

951   **Supplementary File 2**
952
953   **Table S6 (Excel file). List of differentially expressed genes in SLOS hNPCs relative to wild-**
954   **type hNPCs. Relate to Figure S1.**
955

References

Barnabe-Heider, F., and Miller, F.D. (2003). Endogenously produced neurotrophins regulate survival and differentiation of cortical progenitors via distinct signaling pathways. J Neurosci *23*, 5149-5160.

Bartkowska, K., Paquin, A., Gauthier, A.S., Kaplan, D.R., and Miller, F.D. (2007). Trk signaling regulates neural precursor cell proliferation and differentiation during cortical development. Development *134*, 4369-4380. doi: 10.1242/dev.008227.

Bonni, A., Brunet, A., West, A.E., Datta, S.R., Takasu, M.A., and Greenberg, M.E. (1999). Cell survival promoted by the Ras-MAPK signaling pathway by transcription-dependent and - independent mechanisms. Science *286*, 1358-1362. doi: 10.1126/science.286.5443.1358.

Bukelis, I., Porter, F.D., Zimmerman, A.W., and Tierney, E. (2007). Smith-Lemli-Opitz syndrome and autism spectrum disorder. Am J Psychiatry *164*, 1655-1661. doi: 164/11/1655 [pii] 10.1176/appi.ajp.2007.07020315.

Byrne, E.F., Sircar, R., Miller, P.S., Hedger, G., Luchetti, G., Nachtergaele, S., Tully, M.D., Mydock-McGrane, L., Covey, D.F., Rambo, R.P.*, et al.* (2016). Structural basis of Smoothened regulation by its extracellular domains. Nature *535*, 517-522. doi: 10.1038/nature18934.

Capecchi, M.R., and Pozner, A. (2015). ASPM regulates symmetric stem cell division by tuning Cyclin E ubiquitination. Nat Commun *6*, 8763. doi: 10.1038/ncomms9763.

Chenn, A., and Walsh, C.A. (2002). Regulation of cerebral cortical size by control of cell cycle exit in neural precursors. Science *297*, 365-369. doi: 10.1126/science.1074192.

Corcoran, R.B., and Scott, M.P. (2006). Oxysterols stimulate Sonic hedgehog signal transduction and proliferation of medulloblastoma cells. Proc Natl Acad Sci U S A *103*, 8408-8413. doi: 10.1073/pnas.0602852103.

Dietschy, J.M., and Turley, S.D. (2004). Cholesterol metabolism in the central nervous system during early development and in the mature animal. J Lipid Res *45*, 1375-1397. doi: 10.1194/jlr.R400004-JLR200 R400004-JLR200 [pii].

DiSalvo, C.V., Zhang, D., and Jacobberger, J.W. (1995). Regulation of NIH-3T3 cell G1 phase transit by serum during exponential growth. Cell Prolif *28*, 511-524. doi: 10.1111/j.1365-2184.1995.tb00089.x.

Doba, T., Burton, G.W., and Ingold, K.U. (1985). Antioxidant and co-antioxidant activity of vitamin C. The effect of vitamin C, either alone or in the presence of vitamin E or a water-soluble vitamin E analogue, upon the peroxidation of aqueous multilamellar phospholipid

liposomes. Biochimica et Biophysica Acta (BBA) - Lipids and Lipid Metabolism *835*, 298-303.
doi: https://doi.org/10.1016/0005-2760(85)90285-1.

DuSell, C.D., Umetani, M., Shaul, P.W., Mangelsdorf, D.J., and McDonnell, D.P. (2008). 27-
Hydroxycholesterol Is an Endogenous Selective Estrogen Receptor Modulator. Molecular
Endocrinology *22*, 65-77. doi: 10.1210/me.2007-0383.

Ferreira, L.G., Dos Santos, R.N., Oliva, G., and Andricopulo, A.D. (2015). Molecular docking
and structure-based drug design strategies. Molecules *20*, 13384-13421. doi:
10.3390/molecules200713384.

Fitzky, B.U., Moebius, F.F., Asaoka, H., Waage-Baudet, H., Xu, L., Xu, G., Maeda, N.,
Kluckman, K., Hiller, S., Yu, H.*, et al.* (2001). 7-Dehydrocholesterol-dependent proteolysis of
HMG-CoA reductase suppresses sterol biosynthesis in a mouse model of Smith-Lemli-
Opitz/RSH syndrome. J Clin Invest *108*, 905-915. doi: 10.1172/jci12103.

Fitzky, B.U., Witsch-Baumgartner, M., Erdel, M., Lee, J.N., Paik, Y.K., Glossmann, H.,
Utermann, G., and Moebius, F.F. (1998). Mutations in the Delta 7-sterol reductase gene in
patients with the Smith-Lemli-Opitz syndrome. Proc Natl Acad Sci U S A *95*, 8181-8186.

Fliesler, S.J., Peachey, N.S., Herron, J., Hines, K.M., Weinstock, N.I., Ramachandra Rao, S., and
Xu, L. (2018). Prevention of Retinal Degeneration in a Rat Model of Smith-Lemli-Opitz
Syndrome. Sci Rep *8*, 1286. doi: 10.1038/s41598-018-19592-8.

Folch, J., Lees, M., and Sloane Stanley, G.H. (1957). A simple method for the isolation and
purification of total lipides from animal tissues. J Biol Chem *226*, 497-509.

Francis, K.R., Ton, A.N., Xin, Y., O'Halloran, P.E., Wassif, C.A., Malik, N., Williams, I.M.,
Cluzeau, C.V., Trivedi, N.S., Pavan, W.J.*, et al.* (2016). Modeling Smith-Lemli-Opitz syndrome
with induced pluripotent stem cells reveals a causal role for Wnt/beta-catenin defects in neuronal
cholesterol synthesis phenotypes. Nat Med *22*, 388-396. doi: 10.1038/nm.4067.

Herron, J., Hines, K.M., and Xu, L. (2018). Assessment of Altered Cholesterol Homeostasis by
Xenobiotics Using Ultra-High Performance Liquid Chromatography-Tandem Mass
Spectrometry. Curr Protoc Toxicol *78*, e65. doi: 10.1002/cptx.65.

Huang, P., Zheng, S., Wierbowski, B.M., Kim, Y., Nedelcu, D., Aravena, L., Liu, J., Kruse,
A.C., and Salic, A. (2018). Structural Basis of Smoothened Activation in Hedgehog Signaling.
Cell *174*, 312-324. doi: 10.1016/j.cell.2018.04.029.

Jeanneteau, F., Garabedian, M.J., and Chao, M.V. (2008). Activation of Trk neurotrophin
receptors by glucocorticoids provides a neuroprotective effect. Proc Natl Acad Sci U S A *105*,
4862-4867. doi: 10.1073/pnas.0709102105.

Jiang, X.S., Wassif, C.A., Backlund, P.S., Song, L., Holtzclaw, L.A., Li, Z., Yergey, A.L., and
Porter, F.D. (2010). Activation of Rho GTPases in Smith-Lemli-Opitz syndrome:
pathophysiological and clinical implications. Hum Mol Genet *19*, 1347-1357. doi:
10.1093/hmg/ddq011.

Konopka, G., Wexler, E., Rosen, E., Mukamel, Z., Osborn, G.E., Chen, L., Lu, D., Gao, F., Gao,
K., Lowe, J.K.*, et al.* (2012). Modeling the functional genomics of autism using human neurons.
Molecular Psychiatry *17*, 202-214. doi: 10.1038/mp.2011.60.

Korade, Z., Xu, L., Harrison, F.E., Ahsen, R., Hart, S.E., Folkes, O.M., Mirnics, K., and Porter,
N.A. (2014). Antioxidant Supplementation Ameliorates Molecular Deficits in Smith-Lemli-
Opitz Syndrome. Biol Psychiatry *75*, 215-222. doi: S0006-3223(13)00591-X [pii]
10.1016/j.biopsych.2013.06.013.

Korade, Z., Xu, L., Shelton, R., and Porter, N.A. (2010). Biological activities of 7-
dehydrocholesterol-derived oxysterols: implications for Smith-Lemli-Opitz syndrome. J Lipid
Res *51*, 3259-3269. doi: jlr.M009365 [pii] 10.1194/jlr.M009365.

Menard, C., Hein, P., Paquin, A., Savelson, A., Yang, X.M., Lederfein, D., Barnabe-Heider, F.,
Mir, A.A., Sterneck, E., Peterson, A.C.*, et al.* (2002). An essential role for a MEK-C/EBP
pathway during growth factor-regulated cortical neurogenesis. Neuron *36*, 597-610. doi:
10.1016/s0896-6273(02)01026-7.

Molyneaux, B.J., Arlotta, P., Menezes, J.R., and Macklis, J.D. (2007). Neuronal subtype
specification in the cerebral cortex. Nat Rev Neurosci *8*, 427-437. doi: 10.1038/nrn2151.

Myers, B.R., Neahring, L., Zhang, Y., Roberts, K.J., and Beachy, P.A. (2017). Rapid, direct
activity assays for Smoothened reveal Hedgehog pathway regulation by membrane cholesterol
and extracellular sodium. Proc Natl Acad Sci U S A *114*, E11141-E11150. doi:
10.1073/pnas.1717891115.

Okita, K., Matsumura, Y., Sato, Y., Okada, A., Morizane, A., Okamoto, S., Hong, H.,
Nakagawa, M., Tanabe, K., Tezuka, K.*, et al.* (2011). A more efficient method to generate
integration-free human iPS cells. Nat Methods *8*, 409-412. doi: 10.1038/nmeth.1591.

Olkku, A., and Mahonen, A. (2009). Calreticulin mediated glucocorticoid receptor export is
involved in β-catenin translocation and Wnt signalling inhibition in human osteoblastic cells.
Bone *44*, 555-565. doi: https://doi.org/10.1016/j.bone.2008.11.013.

Porter, F.D., and Herman, G.E. (2011). Malformation syndromes caused by disorders of
cholesterol synthesis. J Lipid Res *52*, 6-34. doi: jlr.R009548 [pii] 10.1194/jlr.R009548.

Porter, J.A., Young, K.E., and Beachy, P.A. (1996). Cholesterol modification of hedgehog
signaling proteins in animal development. Science *274*, 255-259.

Porter, N.A., Xu, L., and Pratt, D.A. (2020). Reactive Sterol Electrophiles: Mechanisms of
Formation and Reactions with Proteins and Amino Acid Nucleophiles. Chemistry *2*, 390-417.
doi: 10.3390/chemistry2020025.

Raleigh, D.R., Sever, N., Choksi, P.K., Sigg, M.A., Hines, K.M., Thompson, B.M., Elnatan, D.,
Jaishankar, P., Bisignano, P., Garcia-Gonzalo, F.R.*, et al.* (2018). Cilia-Associated Oxysterols
Activate Smoothened. Mol Cell *72*, 316-327 e315. doi: 10.1016/j.molcel.2018.08.034.

Sever, N., Mann, R.K., Xu, L., Snell, W.J., Hernandez-Lara, C.I., Porter, N.A., and Beachy, P.A.
(2016). Endogenous B-ring oxysterols inhibit the Hedgehog component Smoothened in a manner
distinct from cyclopamine or side-chain oxysterols. Proc Natl Acad Sci U S A *113*, 5904-5909.
doi: 10.1073/pnas.1604984113.

Sheng, R., Kim, H., Lee, H., Xin, Y., Chen, Y., Tian, W., Cui, Y., Choi, J.C., Doh, J., Han, J.K.*,*
et al. (2014). Cholesterol selectively activates canonical Wnt signalling over non-canonical Wnt
signalling. Nat Commun *5*, 4393. doi: 10.1038/ncomms5393.

Sikora, D., Pettit-Kekel, K., Penfield, J., Merkens, L., and Steiner, R. (2006). The near universal
presence of autism spectrum disorders in children with Smith-Lemli-Opitz syndrome. Am J Med
Genet *140A*, 1511-1518. doi: 10.1002/(issn)1552-4833.

Theofilopoulos, S., Wang, Y., Kitambi, S.S., Sacchetti, P., Sousa, K.M., Bodin, K., Kirk, J.,
Salto, C., Gustafsson, M., Toledo, E.M.*, et al.* (2013). Brain endogenous liver X receptor ligands
selectively promote midbrain neurogenesis. Nat Chem Biol *9*, 126-133. doi:
10.1038/nchembio.1156.

Thurm, A., Tierney, E., Farmer, C., Albert, P., Joseph, L., Swedo, S., Bianconi, S., Bukelis, I.,
Wheeler, C., Sarphare, G.*, et al.* (2016). Development, behavior, and biomarker characterization
of Smith-Lemli-Opitz syndrome: an update. J Neurodev Disord *8*, 12. doi: 10.1186/s11689-016-
9145-x.

Tierney, E., Bukelis, I., Thompson, R.E., Ahmed, K., Aneja, A., Kratz, L., and Kelley, R.I.
(2006). Abnormalities of cholesterol metabolism in autism spectrum disorders. Am J Med Genet
B *141B*, 666-668. doi: 10.1002/ajmg.b.30368.

Tint, G.S., Irons, M., Elias, E.R., Batta, A.K., Frieden, R., Chen, T.S., and Salen, G. (1994).
Defective cholesterol biosynthesis associated with the Smith-Lemli-Opitz syndrome. N Engl J
Med *330*, 107-113.

Tint, G.S., Seller, M., Hughes-Benzie, R., Batta, A.K., Shefer, S., Genest, D., Irons, M., Elias,
E., and Salen, G. (1995). Markedly increased tissue concentrations of 7-dehydrocholesterol
combined with low levels of cholesterol are characteristic of the Smith-Lemli-Opitz syndrome. J
Lipid Res *36*, 89-95.

Tomita, H., Cornejo, F., Aranda-Pino, B., Woodard, C.L., Rioseco, C.C., Neel, B.G., Alvarez,
A.R., Kaplan, D.R., Miller, F.D., and Cancino, G.I. (2020). The Protein Tyrosine Phosphatase
Receptor Delta Regulates Developmental Neurogenesis. Cell Rep *30*, 215-228 e215. doi:
10.1016/j.celrep.2019.11.033.

Usui, N., Araujo, D.J., Kulkarni, A., Co, M., Ellegood, J., Harper, M., Toriumi, K., Lerch, J.P.,
and Konopka, G. (2017). Foxp1 regulation of neonatal vocalizations via cortical development.
Genes Dev *31*, 2039-2055. doi: 10.1101/gad.305037.117.

Voisin, M., de Medina, P., Mallinger, A., Dalenc, F., Huc-Claustre, E., Leignadier, J., Serhan,
N., Soules, R., Ségala, G., Mougel, A*., et al.* (2017). Identification of a tumor-promoter
cholesterol metabolite in human breast cancers acting through the glucocorticoid receptor.
Proceedings of the National Academy of Sciences *114*, E9346. doi: 10.1073/pnas.1707965114.

Wassif, C.A., Maslen, C., Kachilele-Linjewile, S., Lin, D., Linck, L.M., Connor, W.E., Steiner,
R.D., and Porter, F.D. (1998). Mutations in the human sterol Delta(7)-reductase gene at 11q12-
13 cause Smith-Lemli-Opitz syndrome. American Journal of Human Genetics *63*, 55-62.

Xu, L., Davis, T.A., and Porter, N.A. (2009). Rate Constants for Peroxidation of Polyunsaturated
Fatty Acids and Sterols in Solution and in Liposomes. J Am Chem Soc *131*, 13037-13044.

Xu, L., Korade, Z., and Porter, N.A. (2010). Oxysterols from free radical chain oxidation of 7-
dehydrocholesterol: product and mechanistic studies. J Am Chem Soc *132*, 2222-2232. doi:
10.1021/ja9080265.

Xu, L., Korade, Z., Rosado, D.A., Jr., Mirnics, K., and Porter, N.A. (2013). Metabolism of
oxysterols derived from nonenzymatic oxidation of 7-dehydrocholesterol in cells. J Lipid Res *54*,
1135-1143. doi: 10.1194/jlr.M035733
jlr.M035733 [pii].

Xu, L., Korade, Z., Rosado, D.A., Liu, W., Lamberson, C.R., and Porter, N.A. (2011). An
oxysterol biomarker for 7-dehydrocholesterol oxidation in cell/mouse models for Smith-Lemli-
Opitz syndrome. J Lipid Res *52*, 1222-1233. doi: jlr.M014498 [pii] 10.1194/jlr.M014498.

Xu, L., Mirnics, K., Bowman, A.B., Liu, W., Da, J., Porter, N.A., and Korade, Z. (2012).
DHCEO accumulation is a critical mediator of pathophysiology in a Smith-Lemli-Opitz
syndrome model. Neurobiol Dis *45*, 923-929. doi: S0969-9961(11)00386-X [pii]
10.1016/j.nbd.2011.12.011.

Yang, S.H., Sharrocks, A.D., and Whitmarsh, A.J. (2013). MAP kinase signalling cascades and
transcriptional regulation. Gene *513*, 1-13. doi: 10.1016/j.gene.2012.10.033.

Yin, H., Xu, L., and Porter, N.A. (2011). Free radical lipid peroxidation: mechanisms and
analysis. Chem Rev *111*, 5944-5972. doi: 10.1021/cr200084z.

Yu, J., Vodyanik, M.A., Smuga-Otto, K., Antosiewicz-Bourget, J., Frane, J.L., Tian, S., Nie, J.,
Jonsdottir, G.A., Ruotti, V., Stewart, R.*, et al.* (2007). Induced pluripotent stem cell lines derived
from human somatic cells. Science *318*, 1917-1920. doi: 10.1126/science.1151526.

Zhou, H., Mehta, S., Srivastava, S.P., Grabinska, K., Zhang, X., Wong, C., Hedayat, A., Perrotta,
P., Fernandez-Hernando, C., Sessa, W.C.*, et al.* (2020). Endothelial cell-glucocorticoid receptor
interactions and regulation of Wnt signaling. JCI Insight *5*. doi: 10.1172/jci.insight.131384.
1131

1132